Intron Retention Controls Localization of lncRNAs *PURPL* and *MALAT1* to Promote Cell Proliferation and Migration

Ioannis Grammatikakis[1], Chosita Norkaew[1], You Jin Song[2], Amit K. Behera[3], Erica C. Pehrsson[4,5], Corrine Corrina R. Hartford[1], Shreya Kordale[3], Rishabh Prasanth[2], Yongmei Zhao[6], Biraj Shrethsa[6], Xiao Ling Li[1], Ravi Kumar[1], Ragini Singh[1], Tayvia Brownmiller[7], Xinyu Wen[4,5], Natasha J. Caplen[7], Pablo Perez-Pinera[8], Kannanganattu V. Prasanth[2*], Thomas Gonatopoulos-Pournatzis[3*], Ashish Lal[1*]

[1]Regulatory RNAs and Cancer Section, Genetics Branch, Center for Cancer Research (CCR), National Cancer Institute (NCI), National Institutes of Health (NIH), Bethesda, Maryland, USA

[2]Department of Cell and Developmental Biology, Cancer Center at Illinois, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA

[3]Functional Transcriptomics Section, RNA Biology Laboratory, NCI, NIH, Frederick, Maryland, USA

[4]OMICS Technology Facility, Bioinformatics, Genetics Branch, CCR, NCI, NIH, Bethesda, Maryland, USA

[5]Advanced Biomedical Computational Science, Frederick National Laboratory for Cancer Research, Frederick, Maryland, USA

[6]Sequencing Facility Bioinformatics Group, Bioinformatics and Computational Science Directorate, Frederick National Laboratory for Cancer Research, NCI, NIH, Frederick, Maryland, USA

[7]Functional Genetics Section, Genetics Branch, CCR, NCI, NIH, Bethesda, Maryland, USA

[8]Department of Bioengineering, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA

24 *Corresponding authors: ashish.lal@nih.gov; kumarp@illinois.edu;

25 thomas.gonatopoulos@nih.gov

26 **Classification:** Biological Sciences

27 **Keywords:** U2AF2, lncRNA, *PURPL*, *MALAT1*, intron retention, splicing, CRISPR/Cas9, nuclear

28 speckles, proliferation, cell migration

Abstract

Intron retention (IR) is a form of alternative splicing in which introns that are normally removed are retained in mature transcripts. Despite emerging evidence of widespread IR in protein-coding genes and lncRNAs, the mechanisms and functional consequences underlying this process remain poorly understood. Here, we performed a genome-wide screen, to dissect the mechanisms governing IR in the lncRNA *PURPL*. Unexpectedly, the top hit from the screen was the essential splicing activator U2AF2, which promotes IR in *PURPL* through direct binding to a weak polypyrimidine tract. Retention of this intron drives nuclear localization of *PURPL* and enhances cell proliferation, revealing a functional role for IR. Transcriptome-wide analysis showed that while U2AF2 promotes splicing of most transcripts, consistent with its canonical role, it also promotes IR in a distinct subset of RNAs. This subset includes the nuclear speckle localized lncRNA *MALAT1*, whose speckle localization is impaired upon U2AF2 depletion. Using *MALAT1* knockout cells reconstituted with wild-type or intron-deleted *MALAT1* variants, we identified a single intron that is essential for *MALAT1* nuclear speckle localization. Deletion of this intron from endogenous *MALAT1* disrupted speckle localization and reduced cell migration, phenocopying the loss of *MALAT1*. Together, these findings uncover a previously unrecognized role for U2AF2 in promoting intron retention and establish IR as a key mechanism regulating lncRNA localization and function.

Significance statement

Intron retention is an understudied form of alternative splicing that can profoundly influence protein isoform diversity, RNA stability, localization, and function. We investigated the mechanisms and consequences of intron retention in the lncRNAs *PURPL* and *MALAT1*. Using a genome wide CRISPR screen and transcriptome wide analyses, we discovered that the splicing factor U2AF2, classically known for promoting intron removal, unexpectedly enhances intron retention by binding to a weak polypyrimidine tract in the retained intron of these lncRNAs. Intron retention, in turn, dictates their localization and modulates cell proliferation and migration, revealing a previously unrecognized regulatory layer in lncRNA biology.

Introduction

Splicing refers to the process of intron removal and exon "stitching" as part of RNA processing that occurs in the nucleus and is catalyzed by a large nucleoprotein complex called the spliceosome (1-3). Splicing is a highly regulated process where *cis*- and *trans*-acting factors coordinate proper and efficient spliceosomal recruitment (4). Several consensus sequences in the RNA are crucial for splicing: the 5′ and 3′ splice sites (ss), which demarcate the exon-intron boundaries of the intron to be removed, typically conform to the consensus sequence of GU-AG. The polypyrimidine (Py)-tract is a stretch of pyrimidines close to the 3′ ss and the branch point that lies upstream of the Py-tract (1). The splicing factor U2AF1 (U2-auxiliary factor 1) binds to the 3′ ss and U2AF2 binds to the Py-tract. The U2AF heterodimer, along with SF1 (Splicing factor 1), is recruited to the branch point to form the pre-spliceosomal E complex. This complex serves as a critical intermediate for the subsequent assembly of the spliceosome and splicing catalysis (5-9).

The vast majority of human genes encode pre-mRNAs that contain introns that are spliced out and most nascent transcripts undergo alternative splicing, which leads to the combinatorial joining of exons. Alternative splicing is a major contributor to protein isoform diversity, enabling the generation of multiple transcript variants from a single gene and frequently shaping measurable cellular phenotypes (10-13). There are several types of alternative splicing such as exon skipping, intron retention, mutually exclusive exons, and alternative 5′ and 3′ ss (14). One of the least studied forms of alternative splicing is intron retention (IR). Lately, it has drawn more attention since it has an important role in post-transcriptional gene expression regulation, and it is believed that more than half of human multi-exonic genes are affected by IR (15). IR frequently leads to nuclear detention of transcripts and eventually degradation which may serve as a defense mechanism of the cell against deleterious products. The extra sequence added by the retained intron may change

the open reading frame (ORF) of protein-coding genes and lead to nonsense-mediated decay

(NMD) (16-18) or initiate the use of an upstream ORF (19, 20). In addition, they can contain

binding sites for microRNAs (miRNAs) or RNA-binding proteins (RBPs), or they may form

structural changes that impact the function of the transcript (21). Interestingly, IR has been

associated with several diseases such as cancer where tumor cells show higher levels of IR

compared to normal tissues (22). Studies on IR have shown that retained introns have been

associated with weaker (non-consensus) splice sites, higher GC content, various lengths of introns,

and higher intronic sequence conservation (15, 23). However, the molecular mechanisms of IR

regulation remain largely elusive.

Long non-coding RNAs (lncRNAs) are RNA molecules >200 nucleotides (nt) long that

have no obvious coding potential. Similar to protein-coding genes, lncRNAs are transcribed by

RNA Polymerase II, are 5′-capped, spliced and polyadenylated (24). However, several

characteristics distinguish lncRNAs from protein-coding genes, including higher tissue-specific

expression, lower abundance and stability, predominant nuclear localization, and having fewer but

longer exons (25-29). One of the most distinguished features of lncRNAs is that they are less

efficiently spliced compared to mRNAs (27, 30-32). Interestingly, lncRNAs contain introns with

much slower intron-excision kinetics compared to protein-coding genes (33). However, the

regulatory mechanisms governing lncRNA splicing remain poorly understood.

lncRNAs have been shown to participate in splicing regulation, with *MALAT1* as the most

prominent example. *MALAT1* is a highly abundant nuclear lncRNA which has oncogenic activity

and is one of the first lncRNAs associated with cancer (34-36). It also has the ability to regulate

alternative splicing because it resides in the nuclear speckles and can interact with several SR

(Serine/Arginine rich) proteins to regulate their phosphorylation state and distribution between

transcription sites and speckles (35, 37-39). Nuclear speckles are sub-nuclear compartments that

act as hubs in coordinating transcription, RNA processing and mRNA export (40). Several RBPs

reside in nuclear speckles such as the protein SON which is a key component and marker of nuclear

speckles (41). However, *MALAT1* itself is not essential for the structural integrity of nuclear

speckles (42, 43). Nevertheless, it remains unclear whether *MALAT1* undergoes alternative

splicing and how this process impacts its localization to nuclear speckles.

Another lncRNA that we previously characterized and termed *PURPL* is transcriptionally

induced by p53 (44). *PURPL* plays a growth-promoting role in colorectal and liver cancer cells by

distinct mechanisms including suppressing p53 protein levels. Interestingly, transcriptomic

analyses using both long-read and short-read sequencing revealed multiple *PURPL* isoforms,

several of which retain intron 2 (45). However, the molecular mechanism governing *PURPL* intron

2 splicing has yet to be elucidated.

Here, we show that the intron 2-retaining *PURPL* isoform, which localizes to the nucleus,

has a shorter half-life than the spliced isoform and promotes cell proliferation. To identify

regulatory factors controlling the splicing of this functionally relevant intron, we utilized our

recently developed CRISPR-based genome-wide screening strategy using a pooled guide RNA

library with a *PURPL* intron 2 minigene reporter containing its flanking exons. Unexpectedly, this

approach revealed a role of the splicing activator U2AF2in promoting *PURPL* IR, contrary to its

canonical function in facilitating intron removal. U2AF2 was further found to promote a broader

intron retention program that includes two introns within *MALAT1*. Notably, depletion of U2AF2

resulted in *MALAT1* exclusion from nuclear speckles. *MALAT1* knockout cells exogenously

expressing *MALAT1* truncation mutants lacking specific introns identified one intron essential for

the nuclear speckle association of the *MALAT1* transcript. Breast cancer cells where this intron is

deleted showed reduced cell migration demonstrating a functional role of the intronic sequences.

These findings provide mechanistic insights into the splicing regulation of *PURPL* and *MALAT1*,

uncovering a previously unrecognized non-canonical function of U2AF2 in promoting

functionally relevant intron retention events.

Results

A CRISPR-based screen identifies U2AF2 as an activator of *PURPL* intron 2 retention

We previously identified *PURPL* as a p53-regulated lncRNA that promotes the growth of

colorectal and liver cancer cells (44, 45). In those studies, we reported that *PURPL* undergoes

alternative splicing, with the ~1.9 kb intron 2 either retained or spliced in the mature transcript

(45). To identify regulators of *PURPL* intron 2 retention, we applied our recently developed

screening platform, CRISPR-based identification of Regulators of Alternative Splicing with

Phenotypic Sequencing (CRASP-seq) (46). This approach utilizes a genome-wide knockout

lentiviral library targeting ~18,500 protein-coding genes using a dual hybrid guide (hg)RNA

system that combines *Streptococcus pyogenes* (*Sp*)Cas9 and optimized *Acidaminococcus sp.*

(op)Cas12a nucleases. (13, 47-49). The library also includes ~2,400 pairs and ~500 curated genetic

interaction pairs, allowing combinatorial perturbations from a single construct. Each gene or gene

pair is targeted by four independent hgRNAs alongside ~500 intergenic and non-targeting controls,

totaling ~ 91,000 hgRNAs. The lentiviral library incorporates a doxycycline-inducible *PURPL* IR

reporter comprised of the full native intron 2 sequences (~1.9 kb) flanked by the adjacent exons 2

(161 bp) and 3 (163 bp) (Figure 1A).

We performed the CRASP-screen in two cell lines: HAP1 and RPE1. Following puromycin

selection for 48 h and doxycycline induction for an additional 24 h, RNA was extracted, and poly-

A RNA was purified using oligo(dT) beads to exclude transcripts still undergoing processing.

cDNA was synthesized using customized primers containing unique molecular identifiers (UMIs), facilitating the removal of PCR duplicates during downstream analysis. The synthesized cDNA was then utilized to construct Illumina sequencing libraries for paired-end sequencing. This sequencing approach enables simultaneous quantification of transcripts that either retain or splice out an intron of interest from one read, while the corresponding hgRNA sequences were captured in the other read to pinpoint specific genetic perturbations. By directly linking *PURPL* splicing outcomes with genetic modifications introduced by each hgRNA in the library, this platform allowed the genome-wide identification of *PURPL* splicing regulators (Figure 1B).

We employed an analytical pipeline that prioritized candidate genes based on multiple hgRNAs showing significant deviations from intergenic controls, with consistent identification across two technical replicates. This analysis identified 62 genes that influence *PURPL* intron 2 splicing, the majority of which act as suppressors of intron 2 splicing (Figure 1C and Table S1). Gene ontology analysis of biological processes revealed a strong enrichment for terms related to mRNA processing and RNA splicing. Additionally, analysis of cellular components indicated that the identified hits are predominantly associated with Spliceosomal Complex, including the U2-type Spliceosomal Complex and the Catalytic step 2 Spliceosome (Figure 1D). These findings validate the effectiveness of our screening approach.

Consistent with the endogenous *PURPL* transcript isoforms, the basal levels of IR of the reporter were very high, with the majority of amplified transcripts corresponding to intron-containing isoforms, as expected based on the intensity of the Illumina libraries before sequencing (Figure S1A). Thus, the detection of genes that suppress IR might be below the detection levels of this screen. Therefore, we focused on the genes that promote intron retention. By calculating the change in Percent Intron Retention ($\Delta\text{PIR} = \text{PIR}_{\text{guide}} - \text{PIR}_{\text{intergenic}}$) across the two cell lines, we

identified 5 unique genes as regulators of *PURPL* intron 2 retention, all of which act to suppress

intron 2 splicing (Figure 1E and Table S1). This highly significant overlap between the two cell

lines (Figure 1E; odds ratio = 204.4, p=2.81xe^{-10}; Fisher's exact test) indicated a shared regulatory

mechanism governing *PURPL* IR, although cell-type-specific effects may also contribute.

Surprisingly, the gene with the strongest impact based on the average ΔPIR metric and

identified as a hit in both cell lines, was U2AF2 (Figure 1F). This was followed by Small Nuclear

Ribonucleoprotein U1 Subunit 70 (SNRNP70), a core component of the U1 snRNP, and

Elongation Factor Tu GTP Binding Domain, which is associated with U5 snRNP (Figure 1F). Our

finding that U2AF2 promotes IR of *PURPL* intron 2 was unexpected because U2AF2 is a

component of the spliceosome machinery, and its canonical function is to promote intron removal

(5, 9).

U2AF2 binds to the *PURPL* intron 2 transcript and promotes intron retention

U2AF2, as a heterodimer with U2AF1, binds to the Py-tract and the 3′ ss, respectively, to promote

splicing (5, 8). To elucidate its role in *PURPL* splicing, we investigated whether U2AF2 binds to

intron 2. Enhanced Crosslinking and Immunoprecipitation sequencing (eCLIP-seq) (50), from the

ENCODE project (51, 52) from HepG2 cells revealed multiple U2AF2 and U2AF1 binding sites

not only proximal to the 3′ ss of *PURPL* intron 2 but also broadly distributed across intron 2 (Figure

2A). Compared to eCLIP data for other RBPs, U2AF1 and U2AF2 showed the highest density of

binding sites and the most prominent peaks. Specific binding of U2AF2 to intron 2 was confirmed

by performing RNA immunoprecipitation (RNA-IP) using a U2AF2-specific antibody and primer

pairs designed to detect *PURPL* transcripts. U2AF2 was found to interact with transcripts

containing *PURPL* intron 2, but not with the spliced *PURPL* transcripts (Figure 2B and S1B).

To functionally validate the role of U2AF2 in *PURPL* intron 2 retention, we performed

RT-PCR assays from HAP1 and RPE1 cells transduced with *PURPL* minigene reporter constructs

and control or *U2AF2*-targeting sgRNAs. In both cell lines, we observed a striking decrease in

intron 2 retention upon *U2AF2* depletion as compared to control cells expressing sgRNAs targeting

intergenic regions (Figure 2C). As an alternative approach, we knocked down *U2AF2* using two

independent siRNAs and conducted semi-quantitative RT-PCR assays from SKHEP1, HepG2, and

normal human diploid fibroblasts W138 cells, using a primer triplet (Figures 2D-2F; Figures S2A

and S2B). In all tested cell lines, U2AF2 depletion increased *PURPL* intron 2 splicing, from ~20%

to ~40-60%, more than doubling the abundance of the spliced isoform. Given that U2AF2

functions as a heterodimer with U2AF1, we also knocked down *U2AF1* in SKHEP1 cells and

observed an effect similar to *U2AF2* knockdown on *PURPL* intron 2 retention regulation (Figure

S2C). We expected that U2AF1 would also come as one of the top hits in the screen but *U2AF1*

sgRNAs were not included in the library due to high off-target scores of the guides. These results

demonstrate that the U2AF complex promotes *PURPL* intron 2 retention and that both U2AF1 and

U2AF2 promote intron retention in *PURPL*.

The intron-retained *PURPL* transcript is unstable and localizes to the nucleus

We previously observed an enrichment of the intron 2-retaining *PURPL* transcript in the nucleus

in liver cancer cells (45). To validate these findings, we conducted single molecule RNA-FISH in

HCT116 cells using probes hybridizing to *PURPL* intron 2 (Figure 3A). Because *PURPL* is

upregulated in response to DNA damage (44), we treated the cells with hydroxyurea for 24 h to

induce *PURPL* expression. RNA-FISH revealed that the *PURPL* intron 2 containing transcript was

exclusively localized to the nucleus. To assay for subnuclear compartmentalization, we also

performed RNA-FISH to detect *MALAT1*, which resides in nuclear speckles (38). Intron 2-retained

PURPL transcripts did not colocalize with *MALAT1* indicating that *PURPL* is excluded from nuclear speckles (Figure 3A). We further validated this result by conducting nuclear/cytoplasmic fractionation assays followed by RT-qPCR and found that *PURPL* intron 2-containing transcripts were almost exclusively nuclear (Figure S3A).

Intron retention can affect transcript stability, with longer RNA species often being more vulnerable to degradation. To test if *PURPL* intron 2 effects transcript turnover, we conducted RNA stability assays using the transcription inhibitor Actinomycin D (ActD) and observed that the intron 2-retained *PURPL* transcript was less stable and had a significantly lower half-life than the *PURPL* transcript in which intron 2 was spliced out (Figure 3B). To determine the functional impact of *PURPL* intron 2 retention, we used CRISPRi to generate *PURPL*-depleted SKHEP1 cells by targeting 3 independent sgRNAs to the p53-response element of the *PURPL* promoter that we had used in our previous study (45). We observed >80% reduction in the levels of endogenous *PURPL* in these experiments (Figure 3C). We next stably transduced these *PURPL*-depleted cells with a lentivirus to express exogenous *PURPL* transcript specifically containing intron 2 under the control of a doxycycline (doxy)-inducible promoter and observed ~2-3-fold induction of the exogenous *PURPL* intron 2 containing transcript (Figure 3D). Because we and others have previously shown that *PURPL* is overexpressed in some cancers and *PURPL* promotes proliferation (44, 45, 53, 54), we next performed cell proliferation assays from the doxy-inducible cells. Our results showed that cells expressing intron 2-contaning *PURPL* proliferate faster (~2-fold) as compared to the no-doxy cells after 6 days of doxy treatment (Figure 2E). Induction of intron 2-contaning *PURPL* transcripts did not have any effect on live cell percentage which remained at ~100% (Figure S3B). Importantly, as a negative control, there was no effect of doxy treatment on the proliferation of parental SKHEP1 cells (Figure S3C). To confirm that

proliferation increase induced by exogenous *PURPL* transcript containing intron 2 was not due to

the spliced isoform, we performed semi-quantitative RT-PCR. As shown in Figure 3F, doxy

treatment resulted exclusively in the detection of the intron-retained *PURPL* transcript. These data

suggest that intron 2 retention modulates *PURPL* function by sequestering the transcript in the

nucleus and reducing its stability, while simultaneously promoting cell proliferation, highlighting

a critical regulatory role for intron 2.

A weak polypyrimidine tract is critical for *PURPL* intron 2 retention

U2AF2 binds to the Py-tract to initiate spliceosomal assembly leading to splicing and intron

removal (5). Suboptimal splicing efficiency may be the result of multiple factors, including splice

site strength and the Py-tract sequence composition. The Py-tract is a stretch of 15-20 pyrimidines

(mostly Us) immediately upstream of the 3′ ss. We analyzed the Py-tract of *PURPL* intron 2 and

noticed that it deviates from the consensus, containing numerous purines (Figure S4A). To

investigate whether U2AF2 regulation on intron 2 retention relies on the Py-tract sequence, we

used the *PURPL* intron 2 minigene reporter system (Figure 1A). We mutated the Py-tract,

replacing it either with a strong Py-tract from *PURPL* intron 1 or with a stretch of Us, simulating

a strong Py-tract. *U2AF2* was knocked down in SKHEP1 cells stably expressing the mutated Py-

tract constructs and RT-PCR was conducted to monitor intron 2 splicing. Interestingly, the basal

levels of intron retention were very different with different mutants; ~90% in intron 2 WT (wild-

type), ~30% in *PURPL* intron 1 and ~20% in U-rich construct (Figure S4B and S4C). This data

demonstrates that the weak Py-tract is responsible for the inefficient intron 2 removal in *PURPL*

(Figure S4C). *U2AF2* knockdown resulted in modest increase in splicing of the mutant constructs

as well. Because the minigene reporter contained the mutant Py-tract along with the remaining

endogenous sequences from *PURPL* intron 2, the modest increase in splicing observed for the

mutants after *U2AF2* knockdown may reflect U2AF2 inhibitory binding at additional sites within

PURPL intron 2 beyond the Py-tract (Figure 2A). These data indicate that the endogenous intron

2 Py-tract in *PURPL* is a non-canonical, weak Py-tract that controls the high intron retention

observed in *PURPL*.

303   U2AF2 directly regulates multiple IR events including *MALAT1*

To determine if U2AF2 promotes IR in additional genes, we next performed total RNA-seq and

PacBio Iso-seq from SKHEP1 cells transfected with siCTRL or siU2AF2. Consistent with our

data, *U2AF2* knockdown resulted in a marked increase in the splicing efficiency of *PURPL* intron

2 (Figures S5A and S5B). Additionally, *U2AF2* depletion increased overall *PURPL* expression,

with increased read coverage across all exons (Figures S5A and S5B; Tables S3 and S4). To

identify IR events regulated by U2AF2 at a transcriptome-wide level, we utilized IRFinder, an

algorithm specifically designed to calculate changes in IR (55). This analysis revealed ~1000

statistically significant IR events regulated by U2AF2. For majority of these events (732), there

was an increase in the IR ratio (Intronic abundance/(Intronic abundance + exonic abundance)) after

U2AF2 depletion, indicating that U2AF2 predominantly promotes splicing, as expected. However,

in 269 events, *U2AF2* knockdown resulted in decreased IR, indicating that U2AF2 promotes IR in

a subset of genes (Figure 4A and Table S2). We next analyzed ENCODE U2AF2 eCLIP-seq data

from HepG2 cells to determine if U2AF2 directly binds to the regulated transcripts. U2AF2

binding was detected in 96 out of the 269 IR events supporting a direct role for U2AF2 in

promoting IR (Figure S5C and Table S2).

To validate these findings, we selected three of the identified introns from *MDM1*,

RETREG2, and *TMEM41B* (Figure S6A). We observed multiple U2AF2 binding peaks within the

regulated introns from *MDM1* (Figure S6B). We next assessed their splicing in SKHEP1 cells

following *U2AF2* knockdown using two independent siRNAs. In all three cases, *U2AF2* depletion

significantly enhanced intron removal (Figures S6C-S6E). IRFinder validated that *PURPL* intron

2 was one of the events showing a significant decrease (~2-fold) in IR ratio upon *U2AF2* depletion

(Figure 4B).

Among the lncRNAs in which IR is promoted by U2AF2, was the lncRNA *MALAT1*.

IRFinder identified two introns in *MALAT1* that have weak Py-tract sequences similar to *PURPL*

intron 2 (Figure S7A-S7C). For both introns, *U2AF2* knockdown resulted in a significant decrease

(~2-fold) in the IR ratio compared to siCTRL (Figures 4A and 4B). This result was confirmed by

RT-PCR from HCT116 and SKHEP1 cells upon *U2AF2* knockdown (Figures 4C-4F).

Intron 2 of *MALAT1* drives its localization to nuclear speckles

MALAT1 regulates splicing by interacting with splicing factors in nuclear speckles.(38) To

determine the impact of IR on *MALAT1* localization to nuclear speckles, we performed RNA-FISH

for *MALAT1* upon *U2AF2* depletion in SKHEP1 and HCT116 cells. Remarkably, *U2AF2*

knockdown resulted in a significant decrease in the ratio of *MALAT1* signal intensity in speckles

relative to nucleoplasm (Figures 5A and 5B; Figures S8A and S8B). U2AF2 depletion did not

affect total *MALAT1* levels as observed in our RNA-seq (Table S3). This reduction in *MALAT1*

levels in nuclear speckles did not compromise the structural integrity of nuclear speckles as

indicated by immunostaining for the RNA-binding protein SON, a well-established nuclear

speckle marker (41) (Figures 5A and S8A) and is consistent with previous reports showing that

MALAT1 is not essential for the integrity of nuclear speckles (42, 43). These data suggest that

U2AF2 plays an important role in promoting *MALAT1* localization to nuclear speckles.

To determine whether it is the intron retention in the transcript *per se* which directs

MALAT1 localization or if U2AF2 indirectly regulates its localization, we assessed the nuclear

speckle localization of wild-type (WT) and intron-deleted *MALAT1* transcripts. To do this, we used CRISPR/Cas9 to generate two *MALAT1* knockout (KO) clones in HCT116 by inserting ~100 bp downstream of the transcription start site, a puromycin resistance gene in one allele and a hygromycin resistance gene in the other allele. In both KO clones, >95% reduction in *MALAT1* RNA levels was observed as compared to a wild-type clone (Figure S9A).

Next, we transduced the *MALAT1* KO clones with an empty vector or doxycycline-inducible lentiviral constructs that exogenously express full-length *MALAT1*-WT in which both introns are intact, intron 1 deleted *MALAT1* (del-1), intron 2 deleted *MALAT1* (del-2) or both introns were deleted (del-1+2). After 48 h of doxy-treatment to induce *MALAT1* expression, which was confirmed with RT-qPCR (Figure S9B), RNA-FISH combined with immunofluorescence for the nuclear speckle marker SON, revealed distinct localization patterns. As expected, exogenously expressed *MALAT1*-WT predominantly (~80%) localized to nuclear speckles (Figures 6A and S10A). Similarly, *MALAT1*-del-1 also predominantly (~80%) localized to nuclear speckles. In contrast, *MALAT1* lacking intron 2 (del-2) or both introns (del-1+2) exhibited a marked reduction (<15%) in nuclear speckle localization and instead displayed diffused localization in the nucleus (Figures 6A and S10A). Importantly, this diffused localization upon intron 2 deletion was observed in both *MALAT1* knockout clones and was further confirmed through quantitative analyses (Figures 6B and S10B).

Deletion of *MALAT1* Intron 2 using CRISPR/Cas9 impairs cell migration

MALAT1 has been shown to promote cell migration and metastasis (34, 56) and its expression has been correlated with high tumor progression and metastasis in various tumors (57). We therefore tested whether the intronic sequences that regulate *MALAT1* nuclear speckle localization are related to the ability of *MALAT1* to promote cell migration. To investigate this, we used the

metastatic breast cancer cell line MDA-MB-231 and first performed RNA-FISH for *MALAT1* after

knocking down *U2AF2*. Similar to SKHEP1 and HCT116 cells, *U2AF2* knockdown resulted in

decreased localization of *MALAT1* to speckles in MDA-MB-231 cells (Figures 7A and 7B). We

next utilized CRISPR/Cas9 to generate KO clones in MDA-MB-231 cells where either the whole

MALAT1 locus or intron 2 of *MALAT1* were deleted. *MALAT1* depletion in the *MALAT1* KO clone

was confirmed with RT-qPCR (Figure S11A). RNA-FISH in 2 separate clones showed that intron

2 deletion reduced *MALAT1* localization to nuclear speckles compared to the WT clone (Figures

7C-7E). In the intron-2 deletion clones, the transcripts containing intron 2 were, as expected,

undetected with primers specifically amplifying intron 2. Interestingly, there was a modest

increase in total *MALAT1* expression suggesting a compensatory mechanism for the loss of intron

2 (Figure S11B). Functionally, when we tested the migration potential of these cells in transwell

cell migration assays, we found significantly reduced cell migration in the intron 2-deleted clones,

phenocopying the complete loss of *MALAT1* locus (Figures 7F and S11C). These data suggest that

the retention of intron 2 is essential for *MALAT1* localization to nuclear speckles and it plays a key

role in promoting cell migration. Together, our data uncovers a U2AF2-driven intron retention

program in lncRNAs that regulates cell proliferation and migration.

Discussion

The goal of this study was to examine the causes and consequences of IR in lncRNAs. Intron

retention is a regulated process that shapes gene expression networks through mechanisms such as

NMD and influences alternative protein isoform generation during differentiation, development,

and disease (17, 58). We previously identified a p53-induced lncRNA, *PURPL*, which exhibits

growth-promoting functions, and that intron 2 of *PURPL* is retained in multiple cell lines (45). To

accomplish our goal, we employed an innovative RNA-seq coupled genome-wide CRISPR screen

on a reporter containing *PURPL* intron 2, and unexpectedly identified U2AF2 as a major inhibitor

of intron 2 removal in *PURPL*. U2AF2 forms a heterodimer with U2AF1, binding to the Py-tract

and the 3′ ss, respectively, to promote splicing by recruiting core spliceosomal components (8, 9,

59). Therefore, given the established role of U2AF2 as a critical positive regulator of splicing

catalysis, we were surprised that U2AF2 inhibits the removal of *PURPL* intron 2. While U2AF1

was not identified as a hit in our screen, follow-up experiments revealed that U2AF1 also

suppresses *PURPL* intron 2 removal, suggesting a previously unrecognized role for the U2AF

complex as a negative regulator *PURPL* IR.

Beyond IR in *PURPL*, we sought to determine whether U2AF2 promotes IR in other

transcripts. Using IR Finder (55), we identified other genes where U2AF2 promoted IR. One of

the genes that drew our attention was the lncRNA *MALAT1* which showed two intron retention

events regulated and bound by U2AF2. These introns contain weak splice sites suggesting that this

may be the reason that they are regulated, and they are not constitutively spliced. Because *MALAT1*

interacts with splicing factors within nuclear speckles (38), we examined whether U2AF2 affects

MALAT1 localization to speckles. Indeed, our results establish *MALAT1* intron 2 as a critical

determinant of its nuclear speckle localization. Future studies involving mapping of the intronic

sequences and targeted mutagenesis of the splice sites within *MALAT1* intron 2 could help

distinguish whether speckle localization of *MALAT1* is inhibited by the splicing process itself or

depends on a discrete element within the intron.

For *PURPL*, previous work established its context-dependent oncogenic activity (44, 45),

but those studies depleted all transcripts from the *PURPL* locus despite the presence of multiple

alternatively spliced isoforms. In the current study, overexpression of the intron 2-retaining

PURPL isoform in CRISPRi-mediated *PURPL*-depleted cells increased proliferation,

demonstrating isoform-specific, growth-promoting function. Our findings reveal that intron

retention in both *MALAT1* and *PURPL* is a regulatory mechanism with direct consequences on

lncRNA localization and cancer-related phenotypes.

The surprising finding of the non-canonical function of U2AF2 promoted us to determine

the underlying mechanism. We first asked whether U2AF2 influences transcription initiation or

elongation at the *PURPL* locus, given the tight coupling between transcription and splicing (60,

61). Transcription kinetics can modulate IR (15, 62) and lncRNAs often lack proximal RNA-pol

II phosphorylation over 5′ ss (33). We therefore assayed for PolII-pSer5 (transcription initiation)

and PolII-pSer2 (transcription elongation) occupancy after U2AF2 knockdown using CUT&RUN-

Seq but there was no significant difference around intron 2 or elsewhere across the *PURPL* locus

(data not shown). We also examined the chromatin mark H3K36me3 which has been linked to IR

(63) and again observed no effect after U2AF2 depletion (data not shown).

U2AF2 functions by binding to the Py-tract, and the Py-tract of *PURPL* intron 2 and

MALAT1 introns 1 and 2 deviate from the consensus, containing a mix of pyrimidines and purines.

These atypical sequences likely contribute to splicing inhibition as revealed by our data from the

mini-gene reporter for *PURPL*. Given that U2AF1 and U2AF2 bind to multiple intronic sites, a

plausible model is that their docking obstructs subsequent spliceosomal assembly. lncRNAs

generally undergo less efficient splicing and consequently, they have higher levels of IR compared

to protein coding genes (31, 33). A study showed that lncRNA splicing strength correlates with 5′

ss strength and lncRNA Py-tract has higher thymidine content compared to protein coding genes

(64). The reduced evolutionary pressure to maintain efficient splicing as they are not used as

templates for protein synthesis, likely contributes to this pattern. This is also supported by their

predominant nuclear localization (26, 27).

Our data shows that *PURPL* transcripts can be either nuclear or cytoplasmic, and U2AF2

regulation of IR offers a mechanism of nuclear detention. Similar to this notion, the lncRNA *TUG1*

remains in the nucleus due to intron retention (65), and the mouse lncRNA *pCharme* undergoes

an IR event which renders it chromatin-bound through its association with MATR3 and PTBP1

(66). Another aspect of the effects of IR lies in the turnover of the transcripts where additional

sequences added by the introns may cause lower transcript stability. In support of this, nuclear

lncRNAs have been shown to be less stable than cytoplasmic ones (28), and *PURPL* also follows

this rule. Notably, the human lncRNA *hFAST* is cytoplasmic and promotes pluripotency through

WNT signaling, but the mouse homolog does not because of its nuclear localization (67).

Introns have been proposed to act as docking elements that tether lncRNAs to chromatin

(27, 66), and a recent study showed that nuclear speckles are associated with distinct classes of

retained introns (68). This raises the question of what makes intron 2 important for speckle

localization of *MALAT1*. One possibility is that this region recruits specific RBPs or adopts a

structural configuration required for speckle association. Our data further indicates that intron 2

retention contributes to *MALAT1's* role in cell migration, potentially because speckle localization

is necessary to sustain this phenotype. Future studies are warranted to dig deeper into the

underlying mechanism and to determine how intron retention shapes the function of *PURPL* and

MALAT1.

Collectively, our findings reveal an expected function of U2AF2 in promoting intron

retention that governs the subcellular localization and function of the lncRNAs *PURPL* and

MALAT1. As intron retention continues to emerge as a regulated form of alternative splicing, the

evidence presented here expands the framework for the mechanisms and biological consequences

of intron retention.

Materials and Methods

Cell culture and treatment

HAP1, HCT116, HEK293T, HepG2, RPE1, SKHEP1, U2OS, WI38, and MDA-MB-231 cells were purchased from ATCC and maintained in Dulbecco's Modified Eagle Medium (DMEM) (Gibco) supplemented with 10% (v/v) fetal bovine serum (Gibco) and 1% (v/v) PenStrep (Gibco) in a 5% CO_2 atmosphere at 37°C. All cell lines were routinely confirmed to be free of mycoplasma using Venor GeM Mycoplasma Detection Kit (Millipore Sigma-Aldrich).

CRASP-Seq *PURPL* screen

The *PURPL* minigene reporter, composed of exons 2 and 3 with the full native intronic sequence, was cloned into the CRASP-Seq genome-wide library as described previously (46). Reporter fragments were synthesized by TWIST Biosciences and assembled into the Eco32I-digested CRASP-Seq knockout library using NEBuilder (49). Library coverage (>250-fold) was maintained through parallel cloning reactions and electroporation into Endura competent cells followed by plasmid amplification and extraction.

Lentivirus was produced following established protocols (49). The pooled library was used to infect HAP1 and RPE1 cells expressing *Sp*Cas9 and opCas12a nucleases (49). at a multiplicity of infection (MOI) of ~0.2. After puromycin selection (2 µg/mL for 48 hours), cells were pooled, induced with doxycycline to activate reporter expression and harvested 24 hours later for RNA analysis.

Total RNA was extracted, poly(A)+ RNA isolated, and cDNA synthesized using a U6-annealing primer. Spliced and intron-retaining reporter transcripts were amplified using exon- or intron-specific primers incorporating Illumina barcodes. Libraries were sequenced on the NovaSeq 6000 platform with a 25% PhiX spike-in and a 300-cycle kit. The sequencing strategy consisted

of the following configuration: Read 1 (210 cycles), Index Read 1 (8 cycles), Index Read 2 (8

cycles) and Read 2 (85 cycles). Each sample yielded over 500 million total reads.

CRASP-seq data analysis

CRASP-Seq data were processed using a custom pipeline (46). Guides, cell barcodes, and UMIs

were retrieved using STAR, and alignment were managed with SAMTools. Reads containing

complete hybrid guide, barcodes, UMI, and splicing outcomes were deduplicated prior to analysis.

Splicing outcomes were quantified as percent intron retention (PIR): $PIR = [IE / (IE + EE)] \times 100$,

where IE represents intron retention reads, and EE represents exon-exon junction reads.

Differential PIR (ΔPIR) was calculated as: $\Delta PIR = PIR_{guide} - PIR_{intergenic}$

The ΔPIR for each gene and for each guide separately is shown in Table S1. Guides with

fewer than 10 reads were excluded. A ΔPIR threshold corresponding to 10% FDR (based on

intergenic controls) defined significant hits. At the gene level, hits required $FPKM \geq 0.1$, at least

two qualifying guides, and $\geq 50\%$ of guides exceeding the ΔPIR threshold in both technical

replicates. Genes identified through combinatorial targeting that met these criteria were included

as hits (Table S1). The heatmap and Venn diagram in Figure 1C and 1E display single-gene hits

as well as collapsed gene hits resulting from combinatorial targeting.

eCLIP-seq data and functional enrichment analysis

eCLIP-seq data for U2AF2, U2AF1, PTBP1, PRPF8, and SRSF1 were obtained from the

ENCODE database (encodeproject.org) with the following identifiers: ENCFF989JBA (CTRL),

ENCFF566CFJ (CTRL), ENCFF536AFD (U2AF2), ENCFF913WRH (U2AF2), ENCFF159SPZ

(U2AF2), ENCFF368XEI (U2AF2), ENCFF542NOA (U2AF1), ENCFF651VRQ (U2AF1),

ENCFF363UDO (PTBP1), ENCFF044RKN (PRPF8), and ENCFF327NVE (SRSF1). For

visualization of RNA-seq data, the Integrated Genome Viewer (IGV) version 17 was used. The

functional enrichment analysis was performed using g:Profiler (69) and the results are shown in

the 'gProfiler_Results' tab in Table S1.

siRNA transfections

For siRNA transfections, we used AllStars Negative Control siRNA (QIAGEN). U2AF2 siRNAs

were purchased from Dharmacon (J-012380-18 and J-012380-19). U2AF1 siRNA was also

purchased from Dharmacon (J-012325-10). Cells were reverse transfected with siRNA at a final

concentration of 20 nM using Lipofectamine RNAiMAX (Invitrogen) in Opti-MEM I Reduced

Serum Medium (Gibco) according to the manufacturer's protocol. SKHEP1 cells were transfected

for 72 hr before RNA harvesting. U2OS, HepG2 and WI38 cells were transfected for two rounds

of 48 h + 72 h or 72 h + 72 h. For RNA FISH, SKHEP1, HCT116, and MDA-MB-231 cells were

transfected for 72 hr. SKHEP1 expressing the *PURPL* Py-tract mutants were transfected for two

rounds of 48 h and 24 h before harvesting, the cells were treated with 1 µg/mL doxycycline to

induce the transgene expression. For RNA-seq, the cells were transfected for 72 h before harvested.

For RNA stability assay, the cells were reseeded in 12-well plates 48 h after siRNA transfection

and 24 h later they were treated with Act D for the indicated time points at 5 µg/mL before RNA

extraction and subsequent assays. For RNA-IP, the cells were transfected for 72 h before harvested

in 10 cm-plates.

RNA extraction, RT-qPCR, semi-quantitative RT-PCR, and splicing gels

Total RNA was extracted using TRIzol (Invitrogen) according to the manufacturer's protocol.

cDNA was then made using iScript™ Reverse Transcription Supermix (Bio-Rad). For real-time

qPCR, all reactions were carried out on StepOnePlus real-time PCR System (Applied Biosystems)

using FastStart SYBR Green Master Mix (Millipore Sigma). *GAPDH* mRNA was used to

normalize expression except for RNA-IPs in Figure 2B where *18S* was used for normalization, and

the relative expression/enrichment of RNAs was calculated using the $2^{-\Delta\Delta Ct}$ method. RT-qPCR

primer sequences for each gene are indicated in Table S3. For splicing gels, cDNA was used to

amplify the region flanking the retained intron with primers in the flanking exons with the addition

of one primer inside the intron as indicated in Figure S1B. In the case of *MALAT1*, primers in the

flanking exons were utilized due to the short length of introns. For the assay of the effect of U2AF2

on the Py-tract mutants of *PURPL*, primers on the transgene were used instead of the endogenous

PURPL exon 2 and 3. The sequences of the primers can be found in Table S5.

RNA-IPs

For RNA-IP, two 10 cm-plates of SKHEP1 cells were lysed with RIPA buffer (Thermo Scientific)

for 10 min on ice before spin down for 10 min at 4°C. The lysate was incubated with 1 μg of anti-

U2AF2 (CST) or rabbit IgG (CST) in 500 μL total volume overnight at 4°C. The next day, 50 μL

of Protein A/G Dynabeads (Pierce) were added to the mix and incubated for 2 h at 4°C. Beads

were then washed 4 times with RIPA buffer and RNA was isolated using TRIzol. Equal volumes

of extracted RNA were used for cDNA synthesis and RT-qPCR as described above.

Immunoblotting

For immunoblotting, total cell lysate was prepared using RIPA buffer (Thermo Scientific)

containing protease inhibitor cocktail (Roche). Protein concentration was determined using the

BCA protein quantitation kit (Thermo Scientific). Cell lysate was loaded onto SDS-PAGE gel and

transferred to a PVDF membrane (Thermo Scientific) using a semi-dry apparatus (Bio-Rad). The

membrane was blocked in 5% skim milk (Millipore) in TBS containing 0.05% Tween 20. The

following antibodies were used: anti-U2AF2 (1:1000, rabbit) (CST, 70471S), anti-GAPDH

(1:10,000, rabbit) (CST, 2118L) and secondary anti-rabbit (1:5000) (CST, 7074S).

Nuclear cytoplasmic fractionation

For nuclear cytoplasmic fractionation, SKHEP1 cells were harvested after trypsinization at 80-

90% confluence. Cells were pelleted and resuspended in 500 µL RSB buffer (10 mM Tris-HCl,

100 mM NaCl, and 2.5 mM MgCl2) containing 40 µg/mL digitonin (Invitrogen), and 100 U of

RNase Out. After incubation on ice for 10 min, cell viability was tested with trypan blue to make

sure that the cells were permeabilized. The cytoplasmic fraction was harvested as supernatant after

centrifugation at 1,500 × g for 5 min at 4°C. The nuclear pellet was resuspended in 100 µL RSB

buffer. 100 µL of the cytoplasmic fractionation were transferred to a fresh tube and 900 µL of

TRIzol was added as well as to the nuclear extracts for RNA extraction and subsequent cDNA

synthesis and RT-qPCR.

Cell proliferation assay

The cell proliferation assay was conducted as previously described (70). To measure proliferation,

parental SKHEP1 cells and *PURPL*-CRISPRi cells overexpressing *PURPL* containing intron 2

were seeded at $1x10^{\wedge 5}$ cells per well in 6-well plates in the presence of 1 µg/mL doxycycline. After

3 days, the cell numbers and viability were measured. The cells were then passaged at $1x10^5$ and

measured again at 6 days, and cumulative cell numbers were plotted.

RNA-seq

The TruSeq Stranded mRNA libraries were pooled and sequenced on one NovaSeq6000 S1

flowcell using 2x101 cycles for pair-end run. The Real Time Analysis software (RTA v3.4.4) was

used for processing raw data files, the Illumina bcl2fastq v2.20 was used to demultiplex and

convert binary base calls and quality scores to fastq format. The sequencing reads were trimmed

of adapters and low-quality bases using Cutadapt (v1.18). The trimmed reads were mapped to

human reference genome (hg38) and Gencode annotation GENCODE_v30 GTF using STAR

aligner (version 2.7.0f) with two-pass alignment option. RSEM (v1.3.1) was used for gene and

transcript quantification based on GENCODE annotation file. Palantir Foundry was used for analysis of Differential Expression of Genes inside the secure NIH Integrated Data Analysis Platform (NIDAP). The DEG Analysis results are shown in Table S3.

Iso-Seq

The libraries were prepared using PacBio standard Iso-seq library prep using SMRTbell prep kit 3.0. The libraries were sequenced on PacBio Sequel II platform using PacBio v2.0 chemistry. PacBio SMRTlink raw subreads were converted into HiFi circular consensus sequences (CCS) with minimum 3 passes. PacBio IsoSeq v3 pipeline was used to demultiplex the barcodes and remove primers. Additional refine steps included trimming poly-A tails and removing concatemers to generate Full Length Non-Concatemer (FLNC) reads. The FLNC reads were used to map to human reference (hg38) using Minimap2 software to generate alignment bam files. Hierarchical clustering and merging steps were performed to obtain consensus isoforms and the full-length (FL) consensus sequences. The high-quality FL transcripts were mapped to the reference genome by using the minimap2 software. Isoform classification and quality control were done using SQANTI3 software and Illumina short-read gene expression data was combined with PacBio Iso-seq for transcript quantification. The Iso-Seq Analysis results are shown in Table S4. The resulting bam files from the Iso-seq analysis were used for plotting and visualization with the R package Gviz (https://bioconductor.org/packages/release/bioc/html/Gviz.html).

IRFinder

IRFinder (55) was used to identify intron retention events that changed significantly on U2AF2 knockdown. A docker image was downloaded from Docker Hub (docker://cloxd/irfinder:2.0.1). The reference was built using pre-downloaded resources (IRFinder BuildRefProcess), using Ensembl release 92 and the 100 bp mappability file provided by IRFinder. The Build-BED-refs.sh

598   script was adjusted so that transcript biotype was not restricted to "processed_transcript" or

599   "protein_coding". Intron retention was quantified using IRFinder in FastQ mode with adapter

600   trimming. Differential intron retention events between siU2AF2 vs. siCTRL were identified with

601   IRFinder Diff using DESeq2 with the default IR ratio ≥ 0.05 in at least one sample and no warning

602   filters, which profiled 16,859 unique introns.

Construct generation

For Py-tract mutants, gene fragments were purchased from Twist Biosciences and cloned into the

pLCHKO-puro vector containing the WT sequences of *PURPL* (Figure 1A), between sites for

NseI and AfeI. The *PURPL* transcript containing intron 2 was cloned into the pCW57 vector

(Hygro) (Addgene #80922) in SalI and MluI sites. *MALAT1* transcripts were cloned into the

pCW57 vector (BSD) (Addgene #80921) in SalI and MluI sites. The *PURPL* insert was generated

by Twist Biosciences and the *MALAT1* sequence was amplified using Phusion DNA polymerase

(NEB). For deletion mutants, we used the Q5® Site-Directed Mutagenesis Kit (NEB). The primers

used for cloning are listed in Table S5.

Generation of *MALAT1* KO cell lines

For HCT116 *MALAT1* knockout, we used integration by non-homologous end joining, which was

accomplished by introducing a simultaneous double-strand break in genomic DNA and in the

targeting vectors (pCMV-Puro and pCMV-Hygro) (71, 72). Plasmids encoding spCas9 and

sgRNAs were obtained from Addgene (Plasmids #41815 and #47108). Oligonucleotides for

construction of sgRNAs were obtained from Integrated DNA Technologies, hybridized,

phosphorylated and cloned into the sgRNA plasmid or targeting vector using BbsI sites (73).

Target sequences for sgRNAs are provided in Table S5. These vectors were transfected into

HCT116 using Lipofectamine 2000 following recommendations from the manufacturer. Three

days after transfection, the cells were selected with Puromycin (0.5 µg/mL) and Hygromycin (100

µg/mL) to generate clonal populations. Genomic DNA from each clone was isolated using DNEasy

Blood and Tissue Kit (Qiagen). PCRs to detect integration of the targeting vector at the target site

were performed using KAPA2G Robust PCR kits (Kapa Biosystems) according to the

manufacturer's instructions. A typical reaction contained 20-100 ng of genomic DNA in Buffer A

(5 µL), Enhancer (5 µL), dNTPs (0.5 µL), primers forward (PINCR Det FP, 1.25 µL) and reverse

(Targeting vector Det RP, 1.25 µL) and KAPA2G Robust DNA Polymerase (0.5 U). The DNA

sequences of the primers for each target are provided in Table S5. PCR products were visualized

in 2% agarose gels and images were captured using a ChemiDoc-It2 (UVP).

To generate MDA-MB-231 whole-locus and intron 2 *MALAT1* knockout clones, we used

the CRIPSR/Cas9 system by transfecting the cells with gRNAs and purified Cas9. 2 µL of each

30 µM gRNA was mixed with 1 µL (~5 µg) of purified Cas9 and 100 µL of 4D-Nucleofector® X

SE buffer SE from the Cell Line Kit (Lonza) supplemented with 18.5 µL of SE supplement per

100 µL total buffer volume. The mixture was incubated at room temperature for 10 min and

1×10^6 cells were resuspended in nucleofection cuvettes. Nucleofections were carried out using

the 4D-Nucleofector® System (Lonza). The cells were transferred to a 6-well plate, and they were

single-cell sorted 48 h later. The clones were grown for 3-4 weeks before genomic DNA extraction

and genotyping with PCR using the primers in Table S5.

***PURPL* and *MALAT1* overexpression and transduction**

PURPL intron 2 constructs were transduced in previously generated *PURPL*-CRISPRi SKHEP1

cells (45). *MALAT1*-WT and deletion mutants were transduced in the two HCT116-*MALAT1* KO

clones. For lentivirus generation, HEK293T cells were seeded at 2×10^5 cells/well onto 6-well

plates, and 1,200 ng of each construct was transfected with lentiviral package vectors using Lipofectamine 2000 (Life Technologies, Invitrogen) according to the manufacturer's protocol. Virus-containing media were harvested at 48 and 72 h post-transfection and added to cells for 24 h. Transduced cells were selected with 500 µg/mL hygromycin for 2 weeks (SKHEP1) or 10 µg/mL Blasticidin (HCT116).

RNA-FISH, IF and Quantification of FISH

The *PURPL* intron probe set was custom-designed using the Stellaris Probe Designer, tagged with Quasar 570 dye and purchased from Stellaris. The *MALAT1* smFISH probe set with Quasar 670 dye was purchased from Stellaris (Cat# VSMF-2211-5). For *PURPL* intron smRNA-FISH, SKHEP1 cells were seeded on coverslips and cultured for 24 hours at 37°C with 5% CO_2 before inducing DNA damage by incubating the cells with 2 mM hydroxyurea for 24 hours (or regular growth media for control). For *MALAT1* smRNA-FISH, HCT116 and MDA-MB-231 *MALAT1* KO cells were seeded on coverslips and cultured for 24 hours at 37°C with 5% CO_2. *MALAT1* WT, *MALAT1* del-1, *MALAT1* del-2, *MALAT1* del1+2, or empty vector expression was induced by incubating the cells with 1 µg/mL doxycycline for 48 hours. The cells that were transfected with siRNA for U2AF2 were seeded on coverslips at the time of transfection. Cells were fixed with 4% PFA for 15 min at room temperature, permeabilized with 0.5% Triton X-100, and washed with washing buffer (10% formamide, 2XSSC) for 5 min. Probe was added to hybridization buffer (10% formamide in Stellaris hybridization buffer Cat# SMF-HB1-10) at a final concentration of 125 nM. Hybridization was done in a humidified chamber in the dark overnight at 37°C. After hybridization, the coverslips were washed twice with wash buffer for 30 min at 37°C and then post-fixed with 4% PFA for 10 min at room temperature. The coverslips were blocked with 5% normal goat serum and then incubated with SON antibody (Sigma Cat# HPA023535). Primary

and secondary antibodies were diluted in 1% normal goat serum. Antibody incubations were done

at room temperature for 1 h and washed with PBS. DNA was counterstained by DAPI. The

coverslips were then washed with 4XSSC for 5 min at room temperature and mounted in

VectaShield Antifade Mounting Medium (Vector Laboratories). Images were taken using

DeltaVision microscope (GE Healthcare) equipped with 60 X/1.42 NA oil immersion objective

(Olympus) and CoolSNAP-HQ2 camera. To quantify the speckle to nucleoplasm signal intensity

ratio of *MALAT1* in Figures 5B, 7B, 7D, and S8B ImageJ was used to create nucleus masks based

on DAPI and speckles mask based on SON. The signal intensity of *MALAT1* in the speckles mask

was divided by the *MALAT1* signal intensity in the nucleus-minus-speckles area. For the

quantification of Figures 6B, 7E and S10B, the cells with *MALAT1* FISH signal showing

prominent speckle patterns similar to SON were classified as "Speckle" pattern, and the cells with

a diffused *MALAT1* FISH signal were classified as "Diffused". Only *MALAT1*-positive cells were

counted for quantification.

Transwell cell migration assays

For the transwell assay, WT and KO MDA-MB-231 clones were seeded at 10×10^5 cells per well

in 12-well plates. 24 h later, the media were removed and replaced with media containing 1% FBS.

After 5 h, the cells were trypsinized and seeded in the top chamber of the transwell plates (Corning,

Catalog#354578) at 25,000 cells per well at a volume of 100 μL. The lower chamber of the

transwell is filled with 700 μL of media containing 10% FBS. The cells were incubated for 24 h

and the transwell were removed from the plates. The non-migrated cells were scraped with a cotton

swab, and the migrated cells were fixed with ice-cold methanol for 5 min and stained with 0.05%

crystal violet blue. The migrated cells were counted under a light microscope.

Statistics

Statistics were performed using the Student's t-test.

Data availability

691 The Iso-Seq data files have been deposited at GEO with accession number GSE288684 and
692 reviewer token: ktcnasayrbgnjev (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi). The RNA-
693 Seq data files have been deposited at GEO with accession number GSE288685 and reviewer token:
694 qvwhuawixxqjnml (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi).

Acknowledgments

This research was supported by the Intramural Research Program (A.L.) of the National Cancer
Institute (NCI), Center for Cancer Research (CCR) (project numbers ZIA BC011646, ZIA
BC012019), NIH. KVP laboratory is supported by grants from NIH (R01-GM132458), Cancer
center at Illinois seed grant, ARPA-H, and National Science Foundation (NSF) center for
quantitative cell biology. YJS is supported by NIH T32EB019944. We thank the CCR Genomics
Core, CCR, NCI, Bethesda, MD for its valuable assistance with Sanger and Nanopore sequencing.
We also thank the Sequencing Facility, CCR, NCI, Frederick, MD for performing the RNA
sequencing assays. The contributions of the NIH author(s) were made as part of their official duties
as NIH federal employees, are in compliance with agency policy requirements, and are considered
Works of the United States Government. However, the findings and conclusions presented in this
paper are those of the author(s) and do not necessarily reflect the views of the NIH or the U.S.
Department of Health and Human Services.

Author Contributions

709 IG, CN, YJS, CCH, SK, RP, XLL, RK, RS, and PPZ conducted experiments. IG, AKB, EP, YZ,
710 BS, TB, XW, TGP, KP, and AL did data analysis. IG and AL wrote the manuscript. NC, KP, and
711 TGP edited the manuscript.

Declaration of interests

The authors declare they have no competing interest.

References

1.   M. E. Wilkinson, C. Charenton, K. Nagai, RNA Splicing by the Spliceosome. *Annu Rev Biochem* **89**, 359-388 (2020).

2.   L. E. Marasco, A. R. Kornblihtt, The physiology of alternative splicing. *Nat Rev Mol Cell Biol* **24**, 242-254 (2023).

3.   A. G. Matera, Z. Wang, A day in the life of the spliceosome. *Nat Rev Mol Cell Biol* **15**, 108-121 (2014).

4.   D. L. Black, Mechanisms of alternative pre-messenger RNA splicing. *Annu Rev Biochem* **72**, 291-336 (2003).

5.   B. Ruskin, P. D. Zamore, M. R. Green, A factor, U2AF, is required for U2 snRNP binding and splicing complex assembly. *Cell* **52**, 207-219 (1988).

6.   J. A. Berglund, K. Chua, N. Abovich, R. Reed, M. Rosbash, The splicing factor BBP interacts specifically with the pre-mRNA branchpoint sequence UACUAAC. *Cell* **89**, 781-787 (1997).

7.   J. A. Berglund, N. Abovich, M. Rosbash, A cooperative interaction between U2AF65 and mBBP/SF1 facilitates branchpoint region recognition. *Genes Dev* **12**, 858-867 (1998).

8.   S. Wu, C. M. Romfo, T. W. Nilsen, M. R. Green, Functional recognition of the 3' splice site AG by the splicing factor U2AF35. *Nature* **402**, 832-835 (1999).

9.   D. A. Zorio, T. Blumenthal, Both subunits of U2AF recognize the 3' splice site in Caenorhabditis elegans. *Nature* **402**, 835-838 (1999).

734   10.   E. T. Wang *et al.*, Alternative isoform regulation in human tissue transcriptomes. *Nature*

735   **456**, 470-476 (2008).

736   11.   T. W. Nilsen, B. R. Graveley, Expansion of the eukaryotic proteome by alternative

737   splicing. *Nature* **463**, 457-463 (2010).

738   12.   Q. Pan, O. Shai, L. J. Lee, B. J. Frey, B. J. Blencowe, Deep surveying of alternative

739   splicing complexity in the human transcriptome by high-throughput sequencing. *Nat*

740   *Genet* **40**, 1413-1415 (2008).

741   13.   R. A. Gier *et al.*, High-performance CRISPR-Cas12a genome editing for combinatorial

742   genetic screening. *Nat Commun* **11**, 3455 (2020).

743   14.   H. Keren, G. Lev-Maor, G. Ast, Alternative splicing and evolution: diversification, exon

744   definition and function. *Nat Rev Genet* **11**, 345-355 (2010).

745   15.   U. Braunschweig *et al.*, Widespread intron retention in mammals functionally tunes

746   transcriptomes. *Genome Res* **24**, 1774-1786 (2014).

747   16.   J. J. Wong *et al.*, Orchestrated intron retention regulates normal granulocyte

748   differentiation. *Cell* **154**, 583-595 (2013).

749   17.   G. Monteuuis, J. J. L. Wong, C. G. Bailey, U. Schmitz, J. E. J. Rasko, The changing

750   paradigm of intron retention: regulation, ramifications and recipes. *Nucleic Acids Res* **47**,

751   11497-11513 (2019).

752   18.   T. Eom *et al.*, NOVA-dependent regulation of cryptic NMD exons controls synaptic

753   protein levels after seizure. *Elife* **2**, e00178 (2013).

754   19.   M. A. Hossain *et al.*, Posttranscriptional Regulation of Gcr1 Expression and Activity Is

755   Crucial for Metabolic Adjustment in Response to Glucose Availability. *Mol Cell* **62**, 346-

756   358 (2016).

20.   S. Cha, C. P. Hong, H. A. Kang, J. S. Hahn, Differential activation mechanisms of two

isoforms of Gcr1 transcription factor generated from spliced and un-spliced transcripts in

Saccharomyces cerevisiae. *Nucleic Acids Res* **49**, 745-759 (2021).

21.   U. Schmitz *et al.*, Intron retention enhances gene regulatory complexity in vertebrates.

Genome Biol **18**, 216 (2017).

22.   H. Dvinge, R. K. Bradley, Widespread intron retention diversifies most cancer

transcriptomes. *Genome Med* **7**, 45 (2015).

23.   P. L. Boutz, A. Bhutkar, P. A. Sharp, Detained introns are a novel, widespread class of

post-transcriptionally spliced introns. *Genes Dev* **29**, 63-80 (2015).

24.   J. J. Quinn, H. Y. Chang, Unique features of long non-coding RNA biogenesis and

function. *Nat Rev Genet* **17**, 47-62 (2016).

25.   I. Ulitsky, D. P. Bartel, lincRNAs: genomics, evolution, and mechanisms. *Cell* **154**, 26-

46 (2013).

26.   C. J. Guo, G. Xu, L. L. Chen, Mechanisms of Long Noncoding RNA Nuclear Retention.

Trends Biochem Sci **45**, 947-960 (2020).

27.   B. Zuckerman, I. Ulitsky, Predictive models of subcellular localization of long RNAs.

RNA **25**, 557-572 (2019).

28.   M. B. Clark *et al.*, Genome-wide analysis of long noncoding RNA stability. *Genome Res*

22, 885-898 (2012).

29.   M. N. Cabili *et al.*, Localization and abundance analysis of human lncRNAs at single-cell

and single-molecule resolution. *Genome Biol* **16**, 20 (2015).

30.   H. Tilgner *et al.*, Deep sequencing of subcellular RNA fractions shows splicing to be

predominantly co-transcriptional in the human genome but inefficient for lncRNAs.

Genome Res **22**, 1616-1625 (2012).

31.   M. Mele *et al.*, Chromatin environment, transcriptional regulation, and splicing

distinguish lincRNAs and mRNAs. *Genome Res* **27**, 27-37 (2017).

32.   M. Kumar, I. Grammatikakis, Causes, Consequences and Challenges of Intron Retention

in lncRNAs. *Mol Cell Biol* **45**, 576-584 (2025).

33.   N. Mukherjee *et al.*, Integrative classification of human coding and noncoding genes

through RNA metabolism profiles. *Nat Struct Mol Biol* **24**, 86-96 (2017).

34.   T. Gutschner *et al.*, The noncoding RNA MALAT1 is a critical regulator of the

metastasis phenotype of lung cancer cells. *Cancer Res* **73**, 1180-1189 (2013).

35.   J. N. Hutchinson *et al.*, A screen for nuclear transcripts identifies two linked noncoding

RNAs associated with SC35 splicing domains. *BMC Genomics* **8**, 39 (2007).

36.   P. Ji *et al.*, MALAT-1, a novel noncoding RNA, and thymosin beta4 predict metastasis

and survival in early-stage non-small cell lung cancer. *Oncogene* **22**, 8031-8041 (2003).

37.   D. Bernard *et al.*, A long nuclear-retained non-coding RNA regulates synaptogenesis by

modulating gene expression. *EMBO J* **29**, 3082-3093 (2010).

38.   V. Tripathi *et al.*, The nuclear-retained noncoding RNA MALAT1 regulates alternative

splicing by modulating SR splicing factor phosphorylation. *Mol Cell* **39**, 925-938 (2010).

39.   M. K. Shinn *et al.*, Nuclear speckle proteins form intrinsic and MALAT1-dependent

microphases. *Cell* **189**, 832-852 e824 (2026).

40.   D. L. Spector, A. I. Lamond, Nuclear speckles. *Cold Spring Harb Perspect Biol* **3** (2011).

41. A. Sharma, H. Takata, K. Shibahara, A. Bubulya, P. A. Bubulya, Son is essential for

nuclear speckle organization and cell cycle progression. *Mol Biol Cell* **21**, 650-663

(2010).

42. S. Nakagawa *et al.*, Malat1 is not an essential component of nuclear speckles in mice.

RNA **18**, 1487-1499 (2012).

43. B. Zhang *et al.*, The lncRNA Malat1 is dispensable for mouse development but its

transcription plays a cis-regulatory role in the adult. *Cell Rep* **2**, 111-123 (2012).

44. X. L. Li *et al.*, Long Noncoding RNA PURPL Suppresses Basal p53 Levels and

Promotes Tumorigenicity in Colorectal Cancer. *Cell Rep* **20**, 2408-2423 (2017).

45. C. C. R. Hartford *et al.*, Context-Dependent Function of Long Noncoding RNA PURPL

in Transcriptome Regulation during p53 Activation. *Mol Cell Biol* **42**, e0028922 (2022).

46. A. K. Behera *et al.*, RNA-coupled CRISPR screens reveal ZNF207 as a regulator of

LMNA aberrant splicing in progeria. *Mol Cell* **86**, 41-59 e15 (2026).

47. M. Aregger, K. Xing, T. Gonatopoulos-Pournatzis, Application of CHyMErA Cas9-

Cas12a combinatorial genome-editing platform for genetic interaction mapping and gene

fragment deletion screening. *Nat Protoc* **16**, 4722-4765 (2021).

48. T. Gonatopoulos-Pournatzis *et al.*, Genetic interaction mapping and exon-resolution

functional genomics with a hybrid Cas9-Cas12a platform. *Nat Biotechnol* **38**, 638-648

(2020).

49. M. S. Xiao *et al.*, Genome-scale exon perturbation screens uncover exons critical for cell

fitness. *Mol Cell* **84**, 2553-2572 e2519 (2024).

50. E. L. Van Nostrand *et al.*, Robust transcriptome-wide discovery of RNA-binding protein

binding sites with enhanced CLIP (eCLIP). *Nat Methods* **13**, 508-514 (2016).

51.   E. P. Consortium, An integrated encyclopedia of DNA elements in the human genome.

824   *Nature* **489**, 57-74 (2012).

825   52.   Y. Luo *et al.*, New developments on the Encyclopedia of DNA Elements (ENCODE)

826   data portal. *Nucleic Acids Res* **48**, D882-D889 (2020).

827   53.   T. Berhane *et al.*, Knockdown of the long noncoding RNA PURPL induces apoptosis and

828   sensitizes liver cancer cells to doxorubicin. *Sci Rep* **12**, 19502 (2022).

829   54.   S. Han *et al.*, PURPL represses autophagic cell death to promote cutaneous melanoma by

830   modulating ULK1 phosphorylation. *Cell Death Dis* **12**, 1070 (2021).

831   55.   R. Middleton *et al.*, IRFinder: assessing the impact of intron retention on mammalian

832   gene expression. *Genome Biol* **18**, 51 (2017).

833   56.   E. Martinez-Terroba *et al.*, Overexpression of Malat1 drives metastasis through

834   inflammatory reprogramming of the tumor microenvironment. *Sci Immunol* **9**, eadh5462

835   (2024).

836   57.   G. Arun, D. Aggarwal, D. L. Spector, MALAT1 Long Non-Coding RNA: Functional

837   Implications. *Noncoding RNA* **6** (2020).

838   58.   A. G. Jacob, C. W. J. Smith, Intron retention as a component of regulated gene

839   expression programs. *Hum Genet* **136**, 1043-1057 (2017).

840   59.   C. Shao *et al.*, Mechanisms for U2AF to define 3' splice sites and regulate alternative

841   splicing in the human genome. *Nat Struct Mol Biol* **21**, 997-1005 (2014).

842   60.   S. Naftelberg, I. E. Schor, G. Ast, A. R. Kornblihtt, Regulation of alternative splicing

843   through coupling with transcription and chromatin structure. *Annu Rev Biochem* **84**, 165-

844   198 (2015).

61.   L. Herzel, D. S. M. Ottoz, T. Alpert, K. M. Neugebauer, Splicing and transcription touch

base: co-transcriptional spliceosome assembly and function. *Nat Rev Mol Cell Biol* **18**,

637-650 (2017).

62.   J. J. Wong *et al.*, Intron retention is regulated by altered MeCP2-mediated splicing factor

recruitment. *Nat Commun* **8**, 15134 (2017).

63.   R. Guo *et al.*, BS69/ZMYND11 reads and connects histone H3.3 lysine 36

trimethylation-decorated chromatin to regulated pre-mRNA processing. *Mol Cell* **56**,

298-310 (2014).

64.   Z. Krchnakova *et al.*, Splicing of long non-coding RNAs primarily depends on

polypyrimidine tract and 5' splice-site sequences due to weak interactions with SR

proteins. *Nucleic Acids Res* **47**, 911-928 (2019).

65.   G. Dumbovic *et al.*, Nuclear compartmentalization of TERT mRNA and TUG1 lncRNA

is driven by intron retention. *Nat Commun* **12**, 3308 (2021).

66.   F. Desideri *et al.*, Intronic Determinants Coordinate Charme lncRNA Nuclear Activity

through the Interaction with MATR3 and PTBP1. *Cell Rep* **33**, 108548 (2020).

67.   C. J. Guo *et al.*, Distinct Processing of lncRNAs Contributes to Non-conserved Functions

in Stem Cells. *Cell* **181**, 621-636 e622 (2020).

68.   A. R. Barutcu *et al.*, Systematic mapping of nuclear domain-associated transcripts reveals

speckles and lamina as hubs of functionally distinct retained introns. *Mol Cell* **82**, 1035-

1052 e1039 (2022).

69.   L. Kolberg *et al.*, g:Profiler-interoperable web service for functional enrichment analysis

and gene identifier mapping (2023 update). *Nucleic Acids Res* **51**, W207-W212 (2023).

70. J. S. Lee *et al.*, An ultraconserved snoRNA-like element in long noncoding RNA

CRNDE promotes ribosome biogenesis and cell proliferation. *Mol Cell* **85**, 1543-1560

e1510 (2025).

71. A. Brown *et al.*, Multiplexed and tunable transcriptional activation by promoter insertion

using nuclease-assisted vector integration. *Nucleic Acids Res* **47**, e67 (2019).

72. A. Brown, W. S. Woods, P. Perez-Pinera, Multiplexed Targeted Genome Engineering

Using a Universal Nuclease-Assisted Vector Integration System. *ACS Synth Biol* **5**, 582-

588 (2016).

73. A. Brown, W. S. Woods, P. Perez-Pinera, Targeted Gene Activation Using RNA-Guided

Nucleases. *Methods Mol Biol* **1468**, 235-250 (2017).

Figure legends

Figure 1. CRISPR-based screen identifies U2AF2 as a promoter of intron retention. (A)

Schematic of the *PURPL* IR reporter used in the screen to identify IR regulators. The pLCHKO

vector contains the puromycin resistance gene used for selection. The transgene also contains the

genomic sequences of exon 2 (161 bp), intron 2 (1,943 bp), and exon 3 (163 bp) of the *PURPL*

gene under a Doxycycline-inducible promoter. A hybrid guide (hg)RNA for guiding *Sp*Cas9 and

*Lb*Cas12a is expressed under a U6 promoter. Top arrow: The hgRNA is cleaved by Cas12a to

produce 2 guide RNAs for dual targeting of each protein coding gene, paralog targeting, and

genetic interaction mapping. Bottom arrow: Upon intron splicing or intron retention, two different

reads are generated. **(B)** Two different cell lines, stably expressing Cas9 and Cas12a, are used:

HAP1, and RPE1. The CRASP-seq pipeline until Illumina paired-end sequencing of CRISPR

libraries is depicted. **(C)** Heatmap showing single genes only (after removing gene pairs), whether

identified individually or as part of a gene pair. The hits are ordered by the average ΔPIR values

across HAP1 and RPE1, from lowest to highest. **(D)** Left: Gene ontology enrichment analysis of

biological processes regulated by the hits of the screen. The most significant pathways involve

mRNA processing and RNA splicing. Right: Gene ontology enrichment analysis of cellular

components regulated by the hits of the screen. These components include mainly Spliceosomal

Complex, small nuclear ribonucleoprotein complex and the Catalytic step 2 Spliceosome. **(E)**

Venn diagram showing that there are 5 common hits which cause a reduction in the percentage of

intron retention (ΔPIR) of the splicing reporter in the two cell lines. **(F)** Table showing the top 5

hits of the screen with a ΔPIR for each hit in each cell line and their full names.

Figure 2. U2AF2 binds to intron 2-containing *PURPL* transcripts and promotes intron 2

retention. (A) IGV snapshot of eCLIP-seq data showing binding sites and enrichment of U2AF2

and U2AF1 on the *PURPL* transcripts around intron 2. eCLIP-seq data for PTBP1, PRPF8, and

SRSF1 are indicated. The annotated locus by RefSeq is also indicated. eCLIP data-seq was

downloaded from encodeproject.org. **(B)** RT-qPCR after RNA-IPs using a U2AF2 antibody with

primer pairs specifically detecting *PURPL* transcripts as indicated in Figure S1B. U2AF2 binds to

transcripts containing intron 2 but not the ones with intron 1, intron 3, or spliced exons 2 and 3.

Samples were normalized to IgG-IP. 18S was used as a loading control. **(C)** Validation of hgRNAs

using the *PURPL* minigene in HAP1 and RPE1 cells. The gels show RT-PCR products for *PURPL*

transcripts with specific primers for the minigene. The hgRNAs used were targeting intergenic

region, or U2AF2. The schematics next to the gel indicate the expected products of the intron-

retained and spliced isoforms. **(D)** Western blot for U2AF2 showing successful knockdown of

U2AF2 protein in SKHEP1 cells. GAPDH was used as a loading control (lower panel). **(E)**

Schematic of the PCR primer triplet used to detect intron 2 retention (red) or splicing (purple). The

length for each PCR product is indicated. **(F)** Gel with RT-PCR products for *PURPL* upon

knockdown of U2AF2 with 2 different siRNAs in SKHEP1 cells. The schematics next to the gel

indicate the expected products of the intron-retained and spliced isoforms. Between the two

expected PCR products, we observed an extra band corresponding to the inclusion of an alternative

exon inside *PURPL* intron 2 as observed in RefSeq, the inclusion of which is not affected by

U2AF2. **(G)** Graph with quantitation of the gel bands in **(F)**. Error bars represent standard

deviations from 3 **(G)** experiments. ***p<0.001.

Figure 3. Expression of Intron 2 Retaining *PURPL* leads to high proliferation. (A) RNA-FISH

images for *PURPL* with intron 2 retention and *MALAT1* in HCT116 cells without treatment or

after 24 h of 2 mM of Hydorxyurea (HU) to induce *PURPL* expression. Scale bar is 10 μm. **(B)**

RNA stability assays were performed for *PURPL* transcripts by measuring their levels by RT-

qPCR following 0, 2, and 4 h of ActD treatment in SKHEP1 cells. *MYC* was used as a positive

control for unstable RNA. **(C)** RT-qPCR showing *PURPL* depletion in in SKHEP1 cells using 3

different gRNAs (sg2, sg4, sg5) compared to a Non-Targeting Control gRNA. **(D)** RT-qPCR for

intron 2-containing PURPL transcript after 48 h of 1 μg/mL doxycycline treatment in comparison

to no treatment in SKHEP1 *PURPL*-CRISPRi populations using the 3 gRNAs as in **(C)**. **(E)**

Proliferation assay showing the effect of overexpression of intron 2-containing *PURPL* transcript

in the proliferation of SKHEP1 cells where the endogenous *PURPL* is knocked down with

CRISPRi. Error bars represent standard deviation from 3 populations with different gRNAs. The

cells were treated with 1 μg/mL doxycycline to induce intron 2-retained *PURPL* expression and

cell proliferation was monitored at 3 and 6 days. **(F)** Gel with RT-PCR products for transcripts

using primers as indicated in Fig. 2E upon doxycycline treatment of SKHEP1 CRIPSRi cells. The

three repeats represent 3 different clones of cells. The last lane is an RT- control. The schematics

next to the gel indicate the expected products of the intron-retained and spliced isoforms. NS

indicates a non-specific band. Error bars in **(B)**, **(C)**, and **(D)** represent standard deviations from 2

experiments. *p<0.05, **p<0.01, ***p<0.001.

Figure 4. U2AF2 directly promotes Intron Retention of *MALAT1*. (A) U2AF2 was knocked

down in SKHEP1 cells and 72 h later, RNA was extracted and RNA-seq was performed. *Left*:

Number of decreased (blue) and increased (red) IR events at various p-values after U2AF2

knockdown as analyzed with the IR Finder algorithm. The purple arrow indicates the *PURPL* IR

event and green arrows indicate *MALAT1* IR events *Right*: Pie chart of the numbers of increased

and decreased IR events upon U2AF2 knockdown. **(B)** Floating bar plots showing the IR ratio of

PURPL intron 2 and the IR ratio of intron 1 (middle) and intron 2 (right) of *MALAT1* in siCTRL

and siU2AF2#1 samples as analyzed with the IRFinder algorithm. **(C)** and **(E)** RT-PCR for

MALAT1 using a primer pair flanking the regulated intron 1 **(C)** or intron 2 **(E)** upon knockdown

of U2AF2 with 2 different siRNAs in HCT116 and SKHEP1 cells. The schematics next to the gel

indicate the expected products of the intron-retained and spliced isoforms. **(D)** and **(F)** Bar graph

with quantitation of the gel bands from **(C)** and **(E)** in SKHEP1 cells. Error bars represent standard

deviations from 2 independent experiments. *p<0.05, **p<0.01, ***p<0.001.

Figure 5. U2AF2 promotes localization of *MALAT1* to nuclear speckles. (A) RNA-FISH

images for *MALAT1* and Immunofluorescence images for SON is shown upon transfection of

SKHEP1 cells with siCTRL or siU2AF2 with two different siRNAs. *MALAT1* is enriched in

nuclear speckles in the siCTRL but not upon U2AF2 knockdown. **(B)** Quantitation of the speckle

to nuclear plasma *MALAT1* signal ratio in the three replicates in panel **(A)**. ****p<0.0001.

Figure 6. Intron 2 of *MALAT1* dictates its localization to nuclear speckles. (A) RNA-FISH

images for *MALAT1* and Immunofluorescence images for SON in a HCT116 *MALAT1*-KO clone

(clone 1) where Empty Vector (EV) or constructs that exogenously express *MALAT1* transcripts

were re-introduced using a doxy-inducible lentivirus. Expressed *MALAT1* transcripts were full-

length (WT) or intron 1 deleted (del-1) or intron 2 deleted (del-2) or both introns deleted

simultaneously (del-1+2). The cells were first treated with 1 μg/mL doxycycline for 48 h to induce

MALAT1 expression. *MALAT1* without intron 2 does not enrich in speckles as it is shown with

SON protein. Scale bar is 10 μm. **(B)** Graph showing quantification of the percentage of cells

where *MALAT1* transcripts are enriched in speckles or being diffused in HCT116 cells clone 1.

Error bars represent standard deviations from 2 independent experiments.

Figure 7. Intron 2 regulates the migration capacity of *MALAT1* in breast cancer cells. (A)

RNA-FISH images for *MALAT1* and Immunofluorescence images for SON are shown upon

transfection of MDA-MB-231 cells with siCTRL or siU2AF2. *MALAT1* is enriched in nuclear

speckles in the siCTRL but not upon U2AF2 knockdown. **(B)** Quantitation of the speckle to nuclear

plasma *MALAT1* signal ratio in the three replicates in panel **(A)**. **(C)** RNA-FISH images for

MALAT1 and Immunofluorescence images for SON in MDA-MB-231 clones either WT, or whole

locus *MALAT1* deletion, or intron 2 deletion. When intron 2 is deleted, *MALAT1* does not enrich

in speckles as it is shown with SON protein. Scale bar is 10 μm. **(D)** Quantitation of the speckle to

nuclear plasma *MALAT1* signal ratio in the three replicates in panel **(C)**. **(E)** Graph showing

quantification of the percentage of *MALAT1* transcripts enriched in speckles or being diffused in

MDA-MB-231 cells in panel **(C)**, N=2. **(F)** Bar graph showing the number of cells that have

migrated in transwell migration assays conducted in MDA-MB-231 WT, *MALAT1* KO and

MALAT1 – intron 2 deletion clones. Intron 2 deletion leads to decreased migration potential. The

number of cells are the sum of cells from 5 different fields, N=4. *p<0.05, **p<0.01, ***p<0.001,

****p<0.0001.

Figure 1

Figure 3

Figure 4

A

SKHEP1

	SON	MALAT1	Merge + DAPI
siCTRL			
siU2AF2 #1			
siU2AF2 #2			

B

Speckle to nucleoplasm MALAT1 signal intensity ratio

siCTRL   siU2AF2#1 ****   siU2AF2#2 ****

Figure 7

A MDA-MB-231

B

C

D

E

F