Non-canonical function of the splicing activator U2AF2 in promoting intron retention in the lncRNAs *PURPL* and *MALAT1*

Ioannis Grammatikakis[1], Chosita Norkaew[1], You Jin Song[2], Amit K. Behera[3], Erica C. Pehrsson[4,5], Corrine Corrina R. Hartford[1], Shreya Kordale[3], Rishabh Prasanth[6], Yongmei Zhao[7], Biraj Shrethsa[7], Xiao Ling Li[1], Ravi Kumar[1], Ragini Singh[1], Tayvia Brownmiller[8], Xinyu Wen[4,5], Natasha J. Caplen[8], Pablo Perez-Pinera[9], Kannanganattu V. Prasanth[2*], Thomas Gonatopoulos-Pournatzis[3*], Ashish Lal[1*]

[1]Regulatory RNAs and Cancer Section, Genetics Branch, Center for Cancer Research (CCR), National Cancer Institute (NCI), National Institutes of Health (NIH), Bethesda, Maryland, USA

[2]Department of Cell and Developmental Biology, Cancer Center at Illinois, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA

[3]Functional Transcriptomics Section, RNA Biology Laboratory, NCI, NIH, Frederick, Maryland, USA

[4]OMICS Technology Facility, Bioinformatics, Genetics Branch, CCR, NCI, NIH, Bethesda, Maryland, USA

[5]Advanced Biomedical Computational Science, Frederick National Laboratory for Cancer Research, Frederick, Maryland, USA

[6]University Laboratory High School, Urbana, Illinois, USA.

[7]Sequencing Facility Bioinformatics Group, Bioinformatics and Computational Science Directorate, Frederick National Laboratory for Cancer Research, NCI, NIH, Frederick, Maryland, USA

[8]Functional Genetics Section, Genetics Branch, CCR, NCI, NIH, Bethesda, Maryland, USA

[9]Department of Bioengineering, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA

*Corresponding authors:** ashish.lal@nih.gov;kumarp@illinois.edu;

thomas.gonatopoulos@nih.gov

Keywords: U2AF2, lncRNA, *PURPL*, *MALAT1*, intron retention, splicing, CRISPR/Cas9, nuclear speckles, non-canonical

Summary

Intron retention (IR) is a form of alternative splicing in which an intron that is typically spliced out is retained in the mature RNA. Despite emerging evidence of widespread IR in protein-coding genes and lncRNAs, the underlying molecular mechanisms remain unclear. Here, we developed a genome-wide screen termed CRASP-seq, to investigate the mechanisms underlying IR in the lncRNA *PURPL.* Unexpectedly, the top hit was the essential splicing activator U2AF2 that promoted IR in *PURPL* by directly binding to a weak polypyrimidine tract. U2AF2 promoted IR in additional transcripts, including the nuclear speckle-localized *MALAT1* whose localization to nuclear speckles was disrupted upon U2AF2 depletion. Importantly, retention of a specific endogenous *MALAT1* intron was critical for its nuclear speckle localization and promoting cell migration. These findings uncover a non-canonical function of U2AF2 in promoting IR and reveal how IR contributes to the subcellular localization and functions of *PURPL* and *MALAT1*.

Introduction

Splicing refers to the process of intron removal and exon "stitching" as part of RNA processing that occurs in the nucleus and is catalyzed by a large nucleoprotein complex called the spliceosome.[1-3] Splicing is a highly regulated process where *cis-* and *trans*-acting factors coordinate proper and efficient spliceosomal recruitment.[4] Several consensus sequences in the RNA are crucial for splicing: the 5′ and 3′ splice sites (ss), which demarcate the exon-intron boundaries of the intron to be removed, typically conform to the consensus sequence of GU-AG. The polypyrimidine (Py)-tract is a stretch of pyrimidines close to the 3′ ss and the branch point that lies upstream of the Py-tract.[1] The splicing factor U2AF1 (U2-auxiliary factor 1) binds to the 3′ ss and U2AF2 binds to the Py-tract. The U2AF heterodimer, along with SF1 (Splicing factor 1), is recruited to the branch point to form the pre-spliceosomal E complex. This complex serves as a critical intermediate for the subsequent assembly of the spliceosome and splicing catalysis.[5-9]

The vast majority of human genes encode pre-mRNAs that contain introns that are spliced out and most nascent transcripts undergo alternative splicing, which leads to the combinatorial joining of exons. Alternative splicing is a major contributor to protein isoform diversity where multiple transcript variants can be produced from a single gene.[10-12] There are several types of alternative splicing such as exon skipping, intron retention, mutually exclusive exons, and alternative 5′ and 3′ ss.[13] One of the least studied forms of alternative splicing is intron retention (IR). Lately, it has drawn more attention since it has an important role in post-transcriptional gene expression regulation, and it is believed that more than half of human multi-exonic genes are affected by IR.[14] IR frequently leads to nuclear detention of the transcripts and

eventually degradation which may serve as a defense mechanism of the cell against deleterious

products. The extra sequence added by the retained intron may change the open reading frame

(ORF) of protein-coding genes and lead to nonsense-mediated decay (NMD)[15-17] or initiate the

use of an upstream ORF.[18,19] In addition, they can contain binding sites for microRNAs

(miRNAs) or RNA-binding proteins (RBPs), or they may form structural changes that impact the

function of the transcript.[20] Interestingly, IR has been associated with several diseases such as

cancer where tumor cells show higher levels of IR compared to normal tissues.[21] Studies on IR

have shown that retained introns have been associated with weaker (non-consensus) splice sites,

higher GC content, various lengths of introns, and higher intronic sequence conservation.[14,22]

However, the molecular mechanisms of IR regulation remain largely elusive.

Long non-coding RNAs (lncRNAs) are RNA molecules >200 nucleotides (nt) long that

have no obvious coding potential. Similar to protein-coding genes, lncRNAs are transcribed by

RNA Polymerase II, are 5′-capped, spliced and polyadenylated.[23] However, several

characteristics distinguish lncRNAs from protein-coding genes, including higher tissue-specific

expression, lower abundance and stability, predominant nuclear localization, and having fewer

but longer exons.[24-28] One of the most distinguished features of lncRNAs is that they are less

efficiently spliced compared to mRNAs.[26,29,30] Interestingly, lncRNAs contain introns with much

slower intron-excision kinetics compared to protein-coding genes.[31] However, the regulatory

mechanisms governing lncRNA splicing remain poorly understood.

lncRNAs have been shown to participate in splicing regulation, with *MALAT1* as the

most prominent example. *MALAT1* is a highly abundant nuclear lncRNA which has oncogenic

activity and is one of the first lncRNAs associated with cancer.[32-34] It also has the ability to

regulate alternative splicing because it resides in the nuclear speckles and can interact with

several SR (Serine/Arginine rich) proteins to regulate their phosphorylation state and distribution between transcription sites and speckles.[33,35-37] Nuclear speckles are sub-nuclear compartments that act as hubs in coordinating transcription, RNA processing and mRNA export.[38] Several RBPs reside in nuclear speckles such as the protein SON which is a key component and marker of nuclear speckles.[39] However, *MALAT1* itself is not essential for the structural integrity of nuclear speckles.[40,41] Nevertheless, it remains unclear whether *MALAT1* undergoes alternative splicing and how this process impacts its localization to nuclear speckles.

Another lncRNA that we previously characterized and termed *PURPL* is transcriptionally induced by p53.[42] *PURPL* plays a growth-promoting role in colorectal and liver cancer cells by distinct mechanisms including suppressing p53 protein levels. Interestingly, through transcriptomic analyses with both long-read and short-read sequencing, we identified multiple *PURPL* isoforms, several of which retain intron 2.[43] However, the molecular mechanism governing *PURPL* intron 2 splicing has yet to be elucidated.

Here, we developed a novel CRISPR-based screening approach using a genome-wide guide RNA library and a minigene reporter containing the sequence of *PURPL* intron 2 and its flanking exons. This approach unexpectedly discovered a direct role of the splicing activator U2AF2 as a key factor promoting *PURPL* IR, contrary to its canonical function in facilitating intron removal. U2AF2 was similarly found to promote an intron retention program that includes two *MALAT1* introns. Interestingly, depletion of U2AF2 resulted in *MALAT1* exclusion from nuclear speckles. *MALAT1* knockout cells exogenously expressing *MALAT1* truncation mutants lacking specific introns identified one intron essential for the nuclear speckle association of the *MALAT1* transcript. Breast cancer cells where this intron is deleted showed reduced cell migration implying a functional role of the intronic sequences. These findings provide

mechanistic insights into the splicing regulation of *PURPL* and *MALAT1*, uncovering a

previously unrecognized non-canonical function of U2AF2 in promoting intron retention.

Results

A CRISPR-based screen identifies U2AF2 as an activator of *PURPL* intron 2 retention

Previously, we identified *PURPL* as a p53-regulated lncRNA that promotes the growth of

colorectal and liver cancer cells.[42,43] In those studies, we reported that *PURPL* undergoes

alternative splicing, and interestingly intron 2 of *PURPL* which is ~1.9 kb long, was either

retained or spliced in the mature transcript.[43] To identify proteins that regulate *PURPL* intron 2

retention, in the present study we used a CRISPR-based screening methodology that we recently

developed, termed CRISPR-based identification of Regulators of Alternative Splicing with

Phenotypic Sequencing (CRASP).[44] This approach utilizes a genome-wide knockout lentiviral

library targeting ~18,500 protein-coding genes using a dual hybrid guide (hg)RNA system. This

system combines *Streptococcus pyogenes* (*Sp*)Cas9 and optimized *Acidaminococcus sp.*

(op)Cas12a nucleases, where the hgRNA is cleaved by the intrinsic RNase activity of Cas12a to

produce two guide RNAs for dual targeting in a given cell.[45-48] The library also includes hgRNAs

targeting paralogs (~2,400 pairs), and a curated set of genetic interaction pairs (~500), enabling

combinatorial gene targeting from a single construct. Each gene and gene pair are targeted by

four independent hgRNAs. Additionally, the library includes ~500 intergenic and non-targeting

controls, totaling around 91,000 hgRNAs. Importantly, the lentiviral library incorporates a

PURPL IR reporter expressed under a doxycycline-inducible Tet Response Element (TRE). The

PURPL minigene reporter is comprised of the full native intron 2 sequences (~1.9 kb) flanked by

the adjacent exons 2 (161 bp) and 3 (163 bp) (Figure 1A).

We performed the CRASP-screen in two cell lines: HAP1 and RPE1. Following

puromycin selection for 48 h and doxycycline induction for an additional 24 h, RNA was

extracted, and poly-A RNA was purified using oligo(dT) beads to exclude transcripts still

undergoing processing. cDNA was synthesized using customized primers containing unique

molecular identifiers (UMIs), facilitating the removal of PCR duplicates during downstream

analysis. The synthesized cDNA was then utilized to construct Illumina sequencing libraries for

paired-end sequencing. This sequencing approach enabled simultaneous quantification of

transcripts that either retain or splice out the intron of interest from one read, while the

corresponding hgRNA sequences were captured in the other read to pinpoint specific genetic

perturbations. By directly linking *PURPL* splicing outcomes with genetic modifications

introduced by each hgRNA in the library, this platform allowed the genome-wide exploration of

PURPL splicing regulators (Figure 1B).

We employed an analytical pipeline that prioritized candidate genes based on multiple

hgRNAs showing significant deviations from intergenic controls, with consistent identification

across two technical replicates. This analysis identified a total of 62 genes that influence *PURPL*

intron 2 splicing, the majority of which act as suppressors of intron 2 splicing (Figure 1C and

Table S1). Gene ontology enrichment analysis of biological processes revealed a strong

enrichment for terms related to mRNA processing and RNA splicing. Additionally, analysis of

cellular components indicated that the identified hits are predominantly associated with

Spliceosomal Complex, including the U2-type Spliceosomal Complex and the Catalytic step 2

Spliceosome (Figure 1D). These findings validate the effectiveness of our screening approach.

Consistent with the endogenous *PURPL* transcript isoforms, the basal levels of IR of the

reporter were very high, with the majority of amplified transcripts corresponding to intron-

containing isoforms, as expected based on the intensity of the Illumina libraries before

sequencing (Figure S1A). Thus, the detection of genes that suppress IR might be below the

detection levels of this screen. Therefore, we focused on the genes that promote intron retention.

By calculating the change in Percent Intron Retention (ΔPIR = PIR_{guide} − $PIR_{intergenic}$) across the

two cell lines, we identified 5 unique genes as regulators of *PURPL* intron 2 retention, all of

which act to suppress intron 2 splicing (Figure 1E and Table S1). The small number of shared

hits between the two cell lines primarily reflects the stringent hit-calling criteria (odds ratio =

204.4, p=2.81xe^{-10}) and cell-type-specific editing constraints.

Surprisingly, the gene with the strongest impact based on the average ΔPIR metric and

identified as a hit in both cell lines, was U2AF2 (Figure 1F). This was followed by Small

Nuclear Ribonucleoprotein U1 Subunit 70 (SNRNP70), a core component of the U1 snRNP, and

Elongation Factor Tu GTP Binding Domain, which is associated with U5 snRNP (Figure 1F).

Our finding that U2AF2 was promoting IR of *PURPL* intron 2 was unexpected because U2AF2

is a component of the spliceosome machinery, and its canonical function is to promote intron

removal.[5,9] RT-PCR assays confirmed our findings, showing that HAP1 and RPE1 cells

transduced with *PURPL* minigene reporter constructs exhibit a striking increase in intron 2

splicing when U2AF2-targeting hgRNAs are expressed, relative to cells expressing hgRNAs

targeting intergenic regions, which served as controls. When the control hgRNAs are used,

PURPL intron 2 is mostly retained (Figure S1B). Our results suggest that depletion of U2AF2

results in decreased *PURPL* intron 2 retention, contradictory to its canonical function.

U2AF2 binds to *PURPL* intron 2 transcripts and promotes intron retention

U2AF2, as a heterodimer with U2AF1, binds to the Py-tract and the 3′ ss, respectively, to

promote splicing.[5,8] To elucidate its role in *PURPL* splicing, we investigated whether U2AF2

binds to intron 2. Enhanced Crosslinking and Immunoprecipitation sequencing (eCLIP-seq),[49]

from the ENCODE project[50,51] from HepG2 cells revealed multiple U2AF2 and U2AF1 binding

sites not only proximal to the 3′ ss of *PURPL* intron 2 but also broadly distributed across intron 2

(Figure 2A). Compared to eCLIP data for other RBPs, U2AF1 and U2AF2 showed the highest

density of binding sites and the most prominent peaks. Specific binding of U2AF2 to intron 2

was confirmed by performing RNA immunoprecipitation (RNA-IP) using a U2AF2-specific

antibody and primer pairs designed to detect *PURPL* transcripts (Figure S1C). U2AF2 was found

to interact with transcripts containing *PURPL* intron 2, but not with the spliced *PURPL*

transcripts (Figure 2B).

To functionally validate the role of U2AF2 in *PURPL* intron 2 retention, we performed RT-

qPCR to quantify endogenous *PURPL* transcript isoforms upon U2AF2 knockdown using two

independent siRNAs (Figure S1D). This analysis revealed an increase in the spliced isoform and

decrease in *PURPL* intron 2 retention in SKHEP1 cells (Figure S1E), as well as other cell lines

such as HepG2 and U2OS cells (Figures S2A and S2C). To further validate the role of U2AF2

on *PURPL* intron 2 splicing, we also conducted semi-quantitative RT-PCR assays from

SKHEP1, HepG2, and W138 cells using a primer triplet (Figure 2C, S2B and S2D). In all cell

lines tested, U2AF2 depletion led to an increase in *PURPL* intron 2 splicing, consistent with

findings from the minigene reporter assay and orthogonal CRISPR-mediated U2AF2 knockdown

experiments.

It should be noted that although U2AF2 knockdown did not completely abolish intron

retention, the effect of U2AF2 knockdown was reproducible and increased splicing of *PURPL*

intron 2 from ~20% to ~40% across multiple cell lines, representing a highly significant change

that effectively doubled the abundance of the spliced *PURPL* isoform. There are several

plausible explanations for the partial nature of this effect. First, because U2AF2 is an essential

gene, we were unable to perform multiple rounds of siRNA transfection, which likely limited the

extent of downstream effects upon U2AF2 knockdown that could be achieved without

compromising cell viability. Consistent with this, as shown in Figure S1B, when a sgRNA was

used to target U2AF2 we observed a strong reduction in *PURPL* intron 2 retention and increased

splicing in both HAP1 and RPE1 cells. Second, *PURPL* intron retention may be regulated by

additional RNA-binding proteins in addition to U2AF2 as revealed by our CRISPR-screen

(Figure 1F). In such a combinatorial regulatory context, depletion of U2AF2 alone would be

expected to produce only a partial reduction in intron retention. Together, these observations

support a significant and biologically meaningful role for U2AF2 in promoting *PURPL* intron 2

retention despite the incomplete penetrance of the knockdown.

Given that U2AF2 functions as a heterodimer with U2AF1, we also knocked down

U2AF1 in SKHEP1 cells and observed a similar effect of U2AF2 knockdown on *PURPL* intron

2 retention regulation (Figures S2E and S2F). We would expect that U2AF1 would also come as

one of the top hits in the screen but U2AF1 gRNAs were not included in the library due to high

off-target scores of the guides. These results demonstrate that the U2AF complex promotes

PURPL intron 2 retention and that both U2AF1 and U2AF2 can promote intron retention.

The intron-retained *PURPL* transcript is unstable and localizes to the nucleus

We previously observed an enrichment of the intron 2-retained *PURPL* transcript in the

nucleus.[43] To confirm this result, we conducted single molecule RNA-FISH in HCT116 cells

using probes hybridizing to *PURPL* intron 2 (Figure 2D). Because *PURPL* is a known target of

p53 and is upregulated upon DNA damage,[42] we treated the cells with the DNA damaging agent

hydroxyurea for 24 hr to induce *PURPL* expression. RNA-FISH confirmed the exclusive nuclear

localization of the *PURPL* intron 2 containing transcript. To assay for subnuclear

compartmentalization, we also performed RNA-FISH to detect *MALAT1*, which resides in

nuclear speckles.[36] Intron 2-retained *PURPL* transcripts did not colocalize with *MALAT1*

indicating that it does not localize to nuclear speckles (Figure 2D). We further validated this

result by conducting nuclear/cytoplasmic fractionation assays followed by RT-qPCR showing

that intron 2-containing transcripts are almost exclusively nuclear (Figure S3A).

Intron retention can affect transcript stability, with longer RNA species often being more

vulnerable to degradation. To test if *PURPL* intron 2 effects transcript turnover, we conducted

RNA stability assays using the transcription inhibitor Actinomycin D (ActD) and observed that

the intron 2-retained *PURPL* transcript is less stable and has a significantly lower half-life than

the *PURPL* transcript in which intron 2 is spliced out (Figure S3B). To identify the functional

impact of *PURPL* intron 2, we first used CRISPRi to generate *PURPL*-depleted SKHEP1 cells

by targeting 3 independent sgRNAs to the p53-response element of the *PURPL* promoter.[43] We

observed >80% reduction in the levels of endogenous *PURPL* in these experiments (Figure

S3C). We next stably transduced these *PURPL*-depleted cells with a lentivirus to express

exogenous *PURPL* transcript specifically containing intron 2 under the control of a doxycycline

(doxy)-inducible promoter and observed ~2-3-fold induction of the exogenous *PURPL* intron 2

containing transcript (Figure 2E). Because we and others have previously shown that *PURPL* is

overexpressed in some cancers and *PURPL* is a pro-proliferative lncRNA,[42,43,52,53] we next

performed cell proliferation assays from the doxy-inducible cells. Our results show that the cells

expressing the intron 2-contaning *PURPL* proliferate faster (~2-fold) as compared to the no-doxy

cells, after 6 days of doxy treatment (Figure 2F). To make sure that the observed increase in

proliferation upon expression of exogenous *PURPL* transcript containing intron 2 was not due to

the spliced *PURPL* isoform, we performed semi-quantitative RT-PCR, and as shown in Figure

S3D, upon doxy treatment, only the intron retained *PURPL* transcript was detected. Importantly,

as a negative control, there was no effect of doxy treatment on the proliferation of parental

SKHEP1 cells (Figure S3E). Induction of intron 2-contaning *PURPL* transcripts did not have any

effect in live cell percentage which remained at ~100% (Figure S3F). Taken together, these

findings suggest that intron 2 retention affects *PURPL* by sequestering the transcript in the

nucleus, reducing its stability and this transcript promotes cell proliferation.

The weak polypyrimidine-tract in *PURPL* intron 2 is important for IR regulation by

U2AF2

U2AF2 binds to the Py-tract to initiate spliceosomal assembly leading to splicing and intron

removal.[5] Suboptimal splicing efficiency may be the result of multiple factors, including splice

site strength and the Py-tract sequence composition. The Py-tract is a stretch of 15-20

pyrimidines (mostly Us) immediately upstream of the 3′ ss. We analyzed the Py-tract of *PURPL*

intron 2 and noticed that it deviates from the consensus, containing numerous purines (Figure

S4A). To investigate whether U2AF2 regulation on intron 2 retention relies on the Py-tract

sequence, we used the *PURPL* intron 2 minigene reporter system (Figure 1A). We mutated the

Py-tract, replacing it either with a strong Py-tract from *PURPL* intron 1 or with a stretch of Us,

simulating a strong Py-tract. U2AF2 was knocked down in SKHEP1 cells stably expressing the

mutated Py-tract constructs and RT-PCR was conducted to monitor intron 2 splicing.

Interestingly, the basal levels of intron retention were very different with different mutants;

~90% in intron 2 WT, ~30% in *PURPL* intron 1 and ~20% in the All-U construct (Figure S4B

and S4C). This shows that the Py-tract is indeed, at least partially, responsible for the inefficient

intron 2 removal in *PURPL* (Figure S4C). Surprisingly, when the WT sequence was replaced

with the strong Py-tract from intron 1 or with a stretch of Us, U2AF2 knockdown continued to

promote splicing. Because the minigene reporter contained the mutant Py-tract along with the

remaining endogenous sequences from *PURPL* intron 2, the modest increase in splicing

observed for the mutants after *U2AF2* knockdown may reflect U2AF2 inhibitory binding at

additional sites within *PURPL* intron 2 beyond the Py-tract (Figure 2A). It may be that these

regions function as a weak Py-tracts, which are sufficient for U2AF2 binding but not strong

enough to efficiently recruit the spliceosome or promote intron removal. These results indicate

that the endogenous intron 2 element in *PURPL* is a non-canonical, low-affinity Py-tract, and

that its weakness may contribute to the high intron retention observed in *PURPL*.

U2AF2 directly regulates multiple IR events including *MALAT1*

To determine if U2AF2 promotes IR in additional genes, we next performed total RNA-seq and

PacBio Iso-seq from SKHEP1 cells transfected with siCTRL or siU2AF2. Consistent with our

data, U2AF2 knockdown resulted in a marked increase in the splicing efficiency of *PURPL*

intron 2 (Figures S5A and S5B). Additionally, we observed that U2AF2 depletion increased

overall *PURPL* expression, with more reads mapping to all exons (Figures S5A and S5B and

Tables S3 and S4).

To identify IR events regulated by U2AF2 at a transcriptome-wide level, we utilized

IRFinder, an algorithm specifically designed to calculate changes in IR.[54] This analysis revealed

~1000 statistically significant IR events regulated by U2AF2. For the majority of these events

(732), there was an increase in the IR ratio (Intronic abundance/(Intronic abundance + exonic

abundance)) after U2AF2 depletion showing that U2AF2 predominantly promotes splicing, as

expected. However, in 269 events, U2AF2 knockdown resulted in decreased IR, indicating that U2AF2 promotes IR in a subset of genes (Figure 3A and Table S2). We next analyzed ENCODE U2AF2 eCLIP-seq data from HepG2 cells to determine if U2AF2 directly binds to the regulated transcripts. We found that in 96 out of the 269 events, U2AF2 was directly bound to the RNA, implying a direct intron retention regulation by U2AF2 (Figure S5C and Table S2).

To validate these findings, we selected three of the identified introns from *MDM1*, *RETREG2*, and *TMEM41B* (Figure S6A). We observed multiple U2AF2 binding peaks within the regulated introns from *MDM1* (Figure S6B). We next assessed their splicing in SKHEP1 cells following U2AF2 knockdown using two independent siRNAs. In all three cases, U2AF2 depletion significantly enhanced intron removal (Figures S6C, S6D, and S6E). IRFinder validated that *PURPL* intron 2 is one of the events showing a significant decrease in IR ratio upon U2AF2 depletion (Figure 3B).

Among the genes in which IR is promoted by U2AF2, there were only a few lncRNAs other than *PURPL*. One of the lncRNAs was *MALAT1*. IRFinder identified two introns in *MALAT1* that have weak Py-tract sequences similar to *PURPL* intron 2 (Figure S7). For both these introns in *MALAT1*, U2AF2 knockdown resulted in a significant decrease in the IR ratio compared to siCTRL (Figures 3A and 3B). This result was confirmed by RT-PCR from HCT116 and SKHEP1 cells upon U2AF2 knockdown (Figures 3C-F). These findings suggest that U2AF2 directly promotes IR in multiple transcripts through binding to the regulated introns. Similar to *PURPL*, depletion of U2AF2 does not result in complete loss of IR likely reflecting the fact that *U2AF2* is an essential gene which limited our ability to perform extended knockdown experiments without compromising cell viability. Additionally, depletion of U2AF2 alone may produce only partial effects when other factors contribute to the regulation of these introns.

SON depletion rescues the effect of U2AF2 knockdown in *PURPL* intron 2 retention

To investigate the mechanism by which U2AF2 inhibits the removal of introns in *PURPL* and *MALAT1*, we considered the possibility of U2AF2 preventing the recruitment of other splicing factors which function towards removal of introns when weak splice sites are present. SON is a large RS-like protein mainly located in nuclear speckles where splicing factors reside, and it has been shown to have the ability of removing constitutive introns with weak splice sites.[55] Based on this, we knocked down SON with 2 different siRNAs in SKHEP1 cells either alone or with simultaneous U2AF2 knockdown (Figures 4A-D). When SON alone was depleted, the effect on intron retention was minimal. However, SON depletion rescued the effect of U2AF2 depletion (Figures 4A-E). This indicates that U2AF2 and SON compete for *PURPL* intron 2 and *MALAT1* intron removal. We next asked the question whether U2AF2 presence affects the binding of SON in *PURPL* and *MALAT1* precursor RNAs. To test this, we conducted RNA-IP using a SON antibody and followed by RT-qPCR using primers specifically detecting *PURPL* and *MALAT1* transcripts (Figure 4F). SON showed higher binding to *MALAT1* and *PURPL* transcripts after U2AF2 depletion compared to negative control. These data suggest that U2AF2 inhibits *PURPL* intron 2 and *MALAT1* splicing partly due to the inhibition of recruitment of SON.

Intron 2 of *MALAT1* drives its localization to nuclear speckles

MALAT1 plays a role in regulating splicing by interacting with splicing factors in the nuclear speckles.[36] To determine the impact of IR on *MALAT1* localization to nuclear speckles we performed RNA-FISH for *MALAT1* upon U2AF2 depletion with two different siRNAs in SKHEP1 and HCT116 cells (Figures 5A and S8A). Remarkably, U2AF2 knockdown resulted in a significant decrease in the ratio of *MALAT1* signal intensity in speckles relative to nucleoplasm (Figures 5A-B and S8A-B). U2AF2 depletion did not affect total *MALAT1* levels as observed in

the RNA-seq results (Table S3). This reduction does not compromise the structural integrity of

nuclear speckles as indicated by immunostaining for the RNA-binding protein SON, a well-

established nuclear speckle marker[39] (Figures 5A and S8A) and is consistent with previous

reports showing that *MALAT1* is not essential for the integrity of nuclear speckles.[40,41] Together,

these results suggest that U2AF2 plays a role in promoting *MALAT1* localization to nuclear

speckles.

To determine whether it is the intron retention in the transcript *per se* which directs

MALAT1 localization or if U2AF2 indirectly regulates its localization, we assessed the nuclear

speckle localization of wild-type (WT) and intron-deleted *MALAT1* transcripts. To do this, we

used CRISPR/Cas9 to generate two *MALAT1* knockout (KO) clones in HCT116 by inserting

~100 bp downstream of the transcription start site, a puromycin resistance gene in one allele and

a hygromycin resistance gene in the other allele (Figure S9A). In both KO clones, there was

>95% reduction in *MALAT1* RNA levels as compared to a wild-type clone (Figure S9A).

Next, we transduced the *MALAT1* KO clones with an empty vector or doxycycline-

inducible lentiviral constructs that exogenously express the full-length *MALAT1*-WT in which

both introns are intact, intron 1 deleted *MALAT1* (del-1), intron 2 deleted *MALAT1* (del-2) or

both introns were deleted (del-1+2). After 48 hours of doxycycline treatment to induce *MALAT1*

expression, which was confirmed with RT-qPCR (Figure S9B), RNA-FISH combined with

immunofluorescence for the nuclear speckle marker SON, revealed distinct localization patterns.

As expected, exogenously expressed *MALAT1*-WT was predominantly (~80%) localized to

nuclear speckles (Figures 6A and S10A). Similarly, *MALAT1*-del-1 was also predominantly

(~80%) localized to nuclear speckles. Remarkably, *MALAT1* lacking intron 2 (del-2) or both

introns (del-1+2) no longer localized (<15%) to nuclear speckles and showed diffused

localization in the nucleus (Figures 6A and S10A). Importantly, this diffused localization upon

intron 2 deletion was observed in both *MALAT1* knockout clones and was further confirmed

through quantitative analyses (Figures 6B and S10B).

Intron 2 of *MALAT1* promotes cell migration

MALAT1 has been shown to promote cell migration and metastasis[32,56] and its expression has

been correlated with high tumor progression and metastasis in various tumors.[57] We therefore

tested whether the intronic sequences that regulate *MALAT1* nuclear speckle localization are

related to the ability of *MALAT1* to promote cell migration. To investigate this, we used the

metastatic breast cancer cell line MDA-MB-231 and first performed RNA-FISH for *MALAT1*

after knocking down U2AF2 with 2 different siRNAs. Similar to SKHEP1 and HCT116 cells,

U2AF2 knockdown resulted in decreased localization of *MALAT1* to speckles in MDA-MB-231

cells (Figures 7A and B). We next utilized CRISPR/Cas9 to generate KO clones in MDA-MB-

231 cells where either the whole *MALAT1* locus or intron 2 of *MALAT1* were deleted. *MALAT1*

depletion in the *MALAT1* KO clone was confirmed with RT-qPCR (Figure S11A). RNA-FISH in

2 separate clones showed that intron 2 deletion reduced *MALAT1* localization to nuclear speckles

compared to the WT clone (Figures 7C-E). In the intron-2 deletion clones, the transcripts

containing intron 2 were, as expected, undetected with primers specifically amplifying intron 2.

Interestingly, there was a modest but significant increase in total *MALAT1* expression suggesting

a compensatory mechanism for the loss of intron 2 and loss of nuclear speckle localization

(Figure S11B). We then tested the migration potential of these cells in transwell cell migration

assays and found reduced cell migration in the intron 2-deleted clones, phenocopying the

complete loss of *MALAT1* (Figures 7F and S11C). These data suggest that the retention of intron

2 is essential for *MALAT1* localization to nuclear speckles and it plays a key role in promoting

cell migration.

Discussion

Intron retention is a regulated process that shapes gene expression networks through mechanisms

such as NMD and influences alternative protein isoform generation during differentiation,

development, and disease.[16,58] We previously identified a p53-induced lncRNA, *PURPL*, which

exhibits growth-promoting functions, and that intron 2 of *PURPL* is retained in multiple cell

lines.[43] In this study, using an innovative RNA-coupled genome-wide CRISPR screen, we

unexpectedly identified U2AF2 as a major inhibitor of intron 2 removal in *PURPL*. U2AF2

forms a heterodimer with U2AF1, binding to the Py-tract and the 3′ ss, respectively, to promote

splicing by recruiting core spliceosomal components.[8,9,59] Therefore, given the established role of

U2AF2 as a critical positive regulator of splicing catalysis, we were surprised that U2AF2

inhibits the removal of *PURPL* intron 2. While U2AF1 was not identified as a hit in our screen,

follow-up experiments revealed that U2AF1 also suppresses *PURPL* intron 2 removal,

suggesting a previously unrecognized role for the U2AF complex as a negative regulator *PURPL*

IR.

Beyond IR in *PURPL*, we sought to determine whether U2AF2 promotes IR in other

transcripts. Using IR Finder,[54] we identified other genes where U2AF2 promoted IR. One of the

genes that drew our attention was the lncRNA *MALAT1* which showed two intron retention

events regulated and bound by U2AF2. These introns contain weak splice sites suggesting that

this may be the reason that they are regulated, and they are not constitutively spliced. Because

MALAT1 interacts with splicing factors within nuclear speckles[36], we examined whether U2AF2

affects *MALAT1* localization to speckles. To determine whether this effect is directly linked to

the retention of one or two of the introns or whether it is a secondary effect, we engineered cell

lines expressing distinct *MALAT1* transcript isoforms. We observed that splicing is directly

involved in the process and that intron 2 sequences are responsible for the inclusion of *MALAT1*

in the nuclear speckles. We next tested the functional consequences of intron 2 retention in a

breast cancer cell line by knocking out *MALAT1* intron 2 using CRISPR/Cas9. Clones in which

intron 2 of *MALAT1* was deleted showed reduced cell migration, phenocopying *MALAT1*-null

cells. It should be noted that although *MALAT1* transcripts lacking intron 2 whether expressed

from constructs or endogenous intron 2 knockout, display predominantly diffused nucleoplasmic

localization, this observation alone cannot distinguish whether the effect arises from a specific

speckle-localization element embedded within intron 2 or from intron removal by the splicing

process *per se*. Future studies involving mapping of the intronic sequences and targeted

mutagenesis of the splice sites within *MALAT1* intron 2 could help distinguish whether speckle

localization of *MALAT1* is inhibited by the splicing process itself or depends on a discrete

element within the intron.

For *PURPL*, previous work established its context-dependent oncogenic activity[42,43], but

those studies depleted all transcripts from the *PURPL* locus despite the presence of multiple

alternatively spliced isoforms. In the current study, overexpression of the intron 2-retaining

PURPL isoform in CRISPRi-mediated *PURPL*-depleted cells increased proliferation,

demonstrating isoform-specific oncogenic function. Together, these findings reveal that intron

retention in both *MALAT1* and *PURPL* is a regulatory mechanism with direct consequences on

lncRNA localization and cancer-related phenotypes.

The surprising finding of the non-canonical function of U2AF2 promoted us to determine

the underlying mechanism. We first asked whether U2AF2 influences transcription initiation or

elongation at the *PURPL* locus, given the tight coupling between transcription and splicing.[60,61]

Transcription kinetics can modulate IR[14,62] and lncRNAs often lack proximal RNA-pol II

phosphorylation over 5′ ss.[31] We therefore assayed for PolII-pSer5 (transcription initiation) and

PolII-pSer2 (transcription elongation) occupancy after U2AF2 knockdown using CUT&RUN-

Seq but there was no significant difference around intron 2 or elsewhere across the *PURPL* locus

(data not shown). We also examined the chromatin mark H3K36me3 which has been linked to

IR[63] and again observed no effect after U2AF2 depletion (data not shown).

U2AF2 functions by binding to the Py-tract, and the Py-tract of *PURPL* intron 2 and

MALAT1 introns 1 and 2 deviate from the consensus, containing a mix of pyrimidines and

purines. These atypical sequences likely contribute to splicing inhibition as revealed by our data

from the mini-gene reporter for *PURPL*. Given that U2AF1 and U2AF2 bind to multiple intronic

sites, a plausible model is that their docking obstructs subsequent spliceosomal assembly.

lncRNAs generally undergo less efficient splicing and consequently, they have higher levels of

IR compared to protein coding genes.[30,31] A study showed that lncRNA splicing strength

correlates with 5′ ss strength and lncRNA Py-tract has higher thymidine content compared to

protein coding genes.[64] The reduced evolutionary pressure to maintain efficient splicing as they

are not used as templates for protein synthesis, likely contributes to this pattern. This is also

supported by their predominant nuclear localization.[25,26]

Our data shows that *PURPL* transcripts can be either nuclear or cytoplasmic, and U2AF2

regulation of IR offers a mechanism of nuclear detention. Similar to this notion, the lncRNA

TUG1 remains in the nucleus due to intron retention[65], and the mouse lncRNA *pCharme*

undergoes an IR event which renders it chromatin-bound through its association with MATR3

and PTBP1.[66] Another aspect of the effects of IR lies in the turnover of the transcripts where

additional sequences added by the introns may cause lower transcript stability. In support of this,

nuclear lncRNAs have been shown to be less stable than cytoplasmic ones[27], and *PURPL* also

follows this rule. Notably, the human lncRNA *hFAST* is cytoplasmic and promotes pluripotency

through WNT signaling, but the mouse homolog does not because of its nuclear localization.[67]

If U2AF2 is not required for intron removal in a subset of transcripts including those

from the *PURPL* and *MALAT1* locus, how would the splicing occur? To address this question,

we surveyed the literature for RNA-binding proteins known to facilitate splicing of transcripts

with weak splice sites. This led us to SON, an RNA-binding protein reported to promote the

splicing of transcripts that harbor suboptimal splice signals[68] SON depletion leads to inefficient

intron removal since it serves as a co-factor to promote splicing of weak splice sites, particularly

in genes encoding for cell-cycle regulating proteins.[69] Interestingly, SON knockdown increased

intron-retention of RNAs interacting with nuclear speckles.[55] We therefore investigated whether

U2AF2 interacts with SON protein in *PURPL* and *MALAT1* intron removal regulation. Although

SON depletion did not have any effect on IR, co-depletion of SON and U2AF2 restored *PURPL*

and *MALAT1* intron retention. Notably, SON binding to *PURPL* and *MALAT1* increased in the

absence of U2AF2 suggesting that U2AF2 inhibits the recruitment of SON for intron removal.

Whether this functional interaction is accompanied by physical interaction of the two splicing

regulators or mutual exclusive binding to *PURPL* remains to be tested. These observations point

to a potential regulatory interplay between SON and U2AF2 in intron removal in *PURPL* and

MALAT1, that will require more targeted mechanistic studies in future studies.

Introns have been proposed to act as docking elements that tether lncRNAs to chromatin,

[26,66] and a recent study showed that nuclear speckles are associated with distinct classes of

retained introns.[70] This raises the question of what makes intron 2 important for speckle

localization of *MALAT1*. One possibility is that this region recruits specific RBPs or adopts a
structural configuration required for speckle association. Our data further indicates that intron 2
retention contributes to *MALAT1's* role in cell migration, potentially because speckle localization
is necessary to sustain this phenotype. Future studies are warranted to dig deeper into the
underlying mechanism and to determine how intron retention shapes the function of *PURPL* and
MALAT1.

Collectively, these findings reveal a non-canonical function of U2AF2 in promoting an
intron retention program that governs the subcellular localization and function of the lncRNAs.
As intron retention continues to emerge as a regulated form of alternative splicing, the evidence
presented here expands the framework for the mechanisms and biological consequences of intron
retention.

Limitations of the study

Limitations of the study stem from the fact that the screen was conducted with an artificial
system where only the sequences from *PURPL* intron 2 and its flanking exons were present in
the construct. The system cannot capture regulatory elements outside the cloned regions or
account for factors that influence splicing such as transcription and chromatin marks. In addition,
although the library contained thousands of gRNAs, it is not exhaustive and may have missed
additional regulators of *PURPL* intron 2 retention, such as U2AF1. Consistent with this, our data
showed that U2AF2 knockdown did not completely abolish *PURPL* or *MALAT1* intron retention.
Thus, intron retention in *PURPL* and *MALAT1* may be regulated by additional RNA-binding
proteins in addition to U2AF2. In such a combinatorial regulatory context, depletion of U2AF2
alone would be expected to produce only a partial reduction in intron retention. Together, these

observations support a significant and biologically meaningful role for U2AF2 in promoting

intron retention in *PURPL* and *MALAT1* despite the incomplete penetrance of the knockdown.

Materials and Methods

Cell culture and treatment

HAP1, HCT116, HEK293T, HepG2, RPE1, SKHEP1, U2OS, WI38, and MDA-MB-231 cells

were purchased from ATCC and maintained in Dulbecco's Modified Eagle Medium (DMEM)

(Gibco) supplemented with 10% (v/v) fetal bovine serum (Gibco) and 1% (v/v) PenStrep (Gibco)

in a 5% CO_2 atmosphere at 37°C. All cell lines were routinely confirmed to be free of

mycoplasma using Venor GeM Mycoplasma Detection Kit (Millipore Sigma-Aldrich).

CRASP-Seq *PURPL* screen

The *PURPL* minigene reporter, composed of exons 2 and 3 with the full native intronic sequence

between them, was integrated into the CRASP-Seq genome-wide library. The *PURPL* intron 2

minigene reporter fragments were synthesized by TWIST Biosciences and cloned using

NEBuilder into the Eco32I-digested CRASP-Seq knockout library vector as described

previously.[48] To ensure a minimum 250-fold library coverage, 16 Eco32I-digested vector

reactions were prepared following the manufacturer's protocol, which included incubation at

50°C for 60 minutes. The reaction mixture was then precipitated with sodium acetate and

ethanol, purified, and transformed into Endura competent cells (LGC Biosearch Technologies)

by electroporation (1 mm cuvette, 25 µF, 200 Ω, 1,600 V). Colonies were plated on LB agar

plates with 100 µg/mL carbenicillin. After overnight incubation at 30°C, bacterial colonies were

pooled, and plasmids were extracted using an Endonuclease Free Plasmid Maxi Kit (QIAGEN).

Lentivirus was produced following established protocols.[48] HEK293T cells (8 million per

15-cm plate) were transfected with a mixture containing 8 µg of the CRASP-Seq lentiviral

library vector, 6.5 µg psPAX2 (Addgene #12260), 4 µg pMD2.G (Addgene #12260), 48 µL X-

tremeGENE 9 DNA Transfection Reagent (Sigma-Aldrich), and 1.4 mL Opti-MEM (Gibco).

After 24 hours, the medium was replaced with serum-free, high-BSA DMEM. Virus-containing

medium was collected 48 hours later, centrifuged at 475 rcf for 5 minutes at 4°C, aliquoted, and

stored at −80°C.

The lentiviral library was used to infect HAP1 and RPE1 cells expressing spCas9 and

opCas12a nucleases.[48] at a multiplicity of infection (MOI) of ~0.2. Infected cells were selected

with 2 µg/mL puromycin for 48 hours, pooled, replated, and induced with doxycycline to

activate reporter expression. Twenty-four hours post-induction, cells were harvested in aliquots

of 10 million for downstream analyses.

Total RNA was extracted using the Qiagen RNeasy Plus Mini Kit, and polyadenylated

mRNA was isolated with oligo(dT) Dynabeads (ThermoFisher Scientific). First-strand cDNA

synthesis was performed with the Maxima H Minus First Strand cDNA Synthesis Kit and a

custom primer annealing to the U6 promoter (ThermoFisher Scientific). PCR amplification

targeted spliced and intron-containing transcripts using custom primers specific to exon 3 or

intron 2, respectively, incorporating Illumina barcodes. PCR reactions were carried out with

NEBNext Ultra II Q5 Master Mix (NEB), and products were resolved on a 1% agarose gel

stained with SYBR Safe (ThermoFisher Scientific). Target bands were extracted and purified

using a ThermoFisher gel extraction kit. Validated libraries were quantified using the Qubit

dsDNA HS Assay (ThermoFisher Scientific) and the Agilent TapeStation system. Libraries were

sequenced on the NovaSeq 6000 platform with a 25% PhiX spike-in and a 300-cycle kit. The

sequencing strategy consisted of the following configuration: Read 1 (210 cycles), Index Read 1

(8 cycles), Index Read 2 (8 cycles) and Read 2 (85 cycles). Each sample yielded over 500

million total reads, ensuring robust data coverage.

Data Analysis

A custom bioinformatics pipeline has been developed to analyze CRASP-Seq data. Cas9 and

Cas12a guides, cell barcodes (CBCs), and unique molecular identifiers (UMIs) were retrieved

using STAR (Dobin et al., 2013). SAMTools (Li et al., 2009) was used to manage sequencing

alignment files. The aligned and annotated reads with all the necessary information including

hybrid guide information, CBC, UMI, and splicing outcomes were deduplicated to carry out

splicing regulation analysis only for complete and unique reads. Splicing reporter reads (R2)

were analyzed for splicing outcomes by computing percent intron retention (PIR) and differential

PIR (ΔPIR) as follows: PIR = [IE / (IE + EE)] $\times$ 100, where IE represents intron retention read

counts, and EE represents exon-exon junction read counts.

- $\Delta PIR = PIR_{guide} - PIR_{intergenic}$

The ΔPIR for each gene and for each guide separately is shown in Table S1. To determine screen

hits, a multi-step filtering process was employed to ensure precision and reliability in the results.

Initially, guides with fewer than 10 reads were excluded from the analysis to eliminate low-

confidence data. Next, a stringent ΔPIR threshold was established based on the intergenic guide

distribution, with a False Discovery Rate (FDR) of 10%, to define statistically significant hits.

This ΔPIR threshold was then applied at several levels of analysis. At the gene level, only genes

meeting the threshold and with FPKM $\geq$ 0.1 were considered hits. For a gene to qualify, at least

50% of its associated guides had to pass the guide-level ΔPIR threshold. Additionally, a gene

required a minimum of two guides to meet the criteria, which excluded instances where a single

guide accounted for the signal. To classify genes as hits, they were required to meet the

aforementioned threshold in both technical replicates, which were generated based on CBC-

defined single-cell lineage groupings. All genes identified through combinatorial gene targeting

that satisfied the previously described criteria were considered hits and they are shown in the

'HITS' tab of Table S1. The heatmap and Venn diagram in Figure 1C and 1E display single-gene

hits as well as collapsed gene hits resulting from combinatorial targeting.

eCLIP-seq data and functional enrichment analysis

eCLIP-seq data for U2AF2, U2AF1, PTBP1, PRPF8, and SRSF1 were obtained from the

ENCODE database (encodeproject.org) with the following identifiers: ENCFF989JBA (CTRL),

ENCFF566CFJ (CTRL), ENCFF536AFD (U2AF2), ENCFF913WRH (U2AF2), ENCFF159SPZ

(U2AF2), ENCFF368XEI (U2AF2), ENCFF542NOA (U2AF1), ENCFF651VRQ (U2AF1),

ENCFF363UDO (PTBP1), ENCFF044RKN (PRPF8), and ENCFF327NVE (SRSF1). For

visualization of RNA-seq data, the Integrated Genome Viewer (IGV) version 17 was used. The

functional enrichment analysis was performed using g:Profiler[71] and the results are shown in the

'gProfiler_Results' tab in Table S1.

siRNA transfections

For siRNA transfections, we used AllStars Negative Control siRNA (QIAGEN). U2AF2 siRNAs

were purchased from Dharmacon (J-012380-18 and J-012380-19). U2AF1 siRNA was also

purchased from Dharmacon (J-012325-10), as well as SON siRNAs (J-012983-05 and J-012983-

06). Cells were reverse transfected with siRNA at a final concentration of 20 nM using

Lipofectamine RNAiMAX (Invitrogen) in Opti-MEM I Reduced Serum Medium (Gibco)

according to the manufacturer's protocol. SKHEP1 cells were transfected for 72 hr before RNA

harvesting. U2OS, HepG2 and WI38 cells were transfected for two rounds of 48 h + 72 h or 72 h

+ 72 h. For RNA FISH, SKHEP1, HCT116, and MDA-MB-231 cells were transfected for 72 hr.

SKHEP1 expressing the *PURPL* Py-tract mutants were transfected for two rounds of 48 h and 24

h before harvesting, the cells were treated with 1μg/ml doxycycline to induce the transgene

expression. For RNA-seq, the cells were transfected for 72 hr before harvested. For RNA

stability assay, the cells were reseeded in 12-well plates 48 hr after siRNA transfection and 24 hr

later they were treated with Act D for the indicated time points at 5ug/ml before RNA extraction

and subsequent assays. For RNA-IP, the cells were transfected for 72 hr before harvested in 10

cm-plates.

RNA extraction, RT-qPCR, RT-PCR, and splicing gels

Total RNA was extracted using TRIzol (Invitrogen) according to the manufacturer's protocol.

cDNA was then made using iScript™ Reverse Transcription Supermix (Bio-Rad). For real-time

qPCR, all reactions were carried out on StepOnePlus real-time PCR System (Applied

Biosystems) using FastStart SYBR Green Master Mix (Millipore Sigma). *GAPDH* mRNA was

used to normalize expression except for RNA-IPs in Figure 2B where *18S* was used for

normalization, and the relative expression/enrichment of RNAs was calculated using the $2^{-\Delta\Delta Ct}$

method. RT-qPCR primer sequences for each gene are indicated in Table S3. For splicing gels,

cDNA was used to amplify the region flanking the retained intron with primers in the flanking

exons with the addition of one primer inside the intron as indicated in Figure S1C. In the case of

MALAT1, primers in the flanking exons were utilized due to the short length of introns. For assay

of the effect of U2AF2 on the Py-tract mutants of *PURPL*, primers on the transgene were used

instead of the endogenous *PURPL* exon 2 and 3. The sequences of the primers can be found in

Table S5.

RNA-IPs

For RNA-IP, two 10 cm-plates of SKHEP1 cells were lysed with RIPA buffer (Thermo Scientific) for 10 min on ice before spin down for 10 min at 4°C. The lysate was incubated with 1 µg of anti-U2AF2 (CST) or rabbit IgG (CST) in 500 µl total volume overnight at 4°C. The next day, 50 µl of Protein A/G Dynabeads (Pierce) were added to the mix and incubated for 2 hr at 4 $^{\circ}$C. Beads were then washed 4 times with RIPA buffer and RNA was isolated using TRIzol. Equal volumes of extracted RNA were used for cDNA synthesis and RT-qPCR as described above. For siU2AF2-transfected cells, one 10 cm-plate was used, and all the lysate was incubated with SON antibody and not IgG.

Immunoblotting

For immunoblotting, total cell lysate was prepared using RIPA buffer (Thermo Scientific) containing protease inhibitor cocktail (Roche). Protein concentration was determined using the BCA protein quantitation kit (Thermo Scientific). Cell lysate was loaded onto SDS-PAGE gel and transferred to a PVDF membrane (Thermo Scientific) using a semi-dry apparatus (Bio-Rad). The membrane was blocked in 5% skim milk (Millipore) in TBS containing 0.05% Tween 20. The following antibodies were used: anti-U2AF2 (1:1000, rabbit) (CST, 70471S), anti-GAPDH (1:10,000, rabbit) (CST, 2118L), anti-SON (1:500, rabbit) (Abcam, 121759), and secondary anti-rabbit (1:5000) (CST, 7074S).

Nuclear cytoplasmic fractionation

For nuclear cytoplasmic fractionation, SKHEP1 cells were harvested after trypsinization at 80-90% confluence. Cells were pelleted and resuspended in 500 µL RSB buffer (10 mM Tris-HCl, 100 mM NaCl, and 2.5 mM MgCl2) containing 40 µg/mL digitonin (Invitrogen), and

100 U of RNase Out. After incubation on ice for 10□min, cell viability was tested with trypan

blue to make sure that the cells were permeabilized. The cytoplasmic fraction was harvested as

supernatant after centrifugation at 1,500□×□g for 5□min at 4°C. The nuclear pellet was

resuspended in 100ul RSB buffer. 100ul of the cytoplasmic fractionation were transferred to a

fresh tube and 900ul of Trizol was added as well as to the nuclear extracts for RNA extraction

and subsequent cDNA synthesis and RT-qPCR.

Cell proliferation assay

The cell proliferation assay was conducted as previously described.[72] To measure proliferation,

parental SKHEP1 cells and *PURPL* CRISPRi cells overexpressing *PURPL* containing intron 2

were seeded at 1 x10^5 cells per well in 6-well plates in the presence of 1μg/ml doxycycline.

After 3 days, the cell numbers and viability were measured. The cells were then passaged at

1x10^5 and measured again at 6 days and cumulative cell numbers were plotted.

RNA-seq

The TruSeq Stranded mRNA libraries were pooled and sequenced on one NovaSeq6000 S1

flowcell using 2x101 cycles for pair-end run. The Real Time Analysis software (RTA v3.4.4)

was used for processing raw data files, the Illumina bcl2fastq v2.20 was used to demultiplex and

convert binary base calls and quality scores to fastq format. The sequencing reads were trimmed

of adapters and low-quality bases using Cutadapt (v1.18). The trimmed reads were mapped to

human reference genome (hg38) and Gencode annotation GENCODE_v30 GTF using STAR

aligner (version 2.7.0f) with two-pass alignment option. RSEM (v1.3.1) was used for gene and

transcript quantification based on GENCODE annotation file. Palantir Foundry was used for

analysis of Differential Expression of Genes inside the secure NIH Integrated Data Analysis

Platform (NIDAP). The DEG Analysis results are shown in Table S3.

Iso-Seq

The libraries were prepared using PacBio standard Iso-seq library prep using SMRTbell prep kit

3.0. The libraries were sequenced on PacBio Sequel II platform using PacBio v2.0 chemistry.

PacBio SMRTlink raw subreads were converted into HiFi circular consensus sequences (CCS)

with minimum 3 passes. PacBio IsoSeq v3 pipeline was used to demultiplex the barcodes and

remove primers. Additional refine steps included trimming poly-A tails and removing

concatemers to generate Full Length Non-Concatemer (FLNC) reads. The FLNC reads were

used to map to human reference (hg38) using Minimap2 software to generate alignment bam

files. Hierarchical clustering and merging steps were performed to obtain consensus isoforms

and the full-length (FL) consensus sequences. The high-quality FL transcripts were mapped to

the reference genome by using the minimap2 software. Isoform classification and quality control

were done using SQANTI3 software and Illumina short-read gene expression data was combined

with PacBio Iso-seq for transcript quantification. The Iso-Seq Analysis results are shown in

Table S4. The resulting bam files from the Iso-seq analysis were used for plotting and

visualization   with   the   R   package   Gviz

(https://bioconductor.org/packages/release/bioc/html/Gviz.html).

IRFinder

IRFinder[54] was used to identify intron retention events that changed significantly on U2AF2

knockdown. A docker image was downloaded from Docker Hub (docker://cloxd/irfinder:2.0.1).

The reference was built using pre-downloaded resources (IRFinder BuildRefProcess), using

Ensembl release 92 and the 100bp mappability file provided by IRFinder. The Build-BED-

refs.sh script was adjusted so that transcript biotype was not restricted to "processed_transcript"

or "protein_coding". Intron retention was quantified using IRFinder in FastQ mode with adapter

trimming. Differential intron retention events between siU2AF2 vs. siCTRL were identified with

IRFinder Diff using DESeq2 with the default IR ratio ≥ 0.05 in at least one sample and no

warning filters, which profiled 16,859 unique introns.

Construct generation

For Py-tract mutants, gene fragments were purchased from Twist Biosciences and cloned into the

pLCHKO-puro vector containing the WT sequences of *PURPL* (Figure 1A), between sites for

NseI and AfeI. The *PURPL* transcript containing intron 2 was cloned into the pCW57 vector

(Hygro) (Addgene #80922) in SalI and MluI sites. *MALAT1* transcripts were cloned into the

pCW57 vector (BSD) (Addgene #80921) in SalI and MluI sites. The *PURPL* insert was

generated by Twist Biosciences and the *MALAT1* sequence was amplified using Phusion DNA

polymerase (NEB). For deletion mutants, we used the Q5® Site-Directed Mutagenesis Kit

(NEB). The primers used for cloning are listed in Table S5.

Generation of *MALAT1* KO cell lines

For HCT116 *MALAT1* knockout, we used integration by non-homologous end joining, which

was accomplished by introducing a simultaneous double-strand break in genomic DNA and in

the targeting vectors (pCMV-Puro and pCMV-Hygro).[73,74] Plasmids encoding spCas9 and

sgRNAs were obtained from Addgene (Plasmids #41815 and #47108). Oligonucleotides for

construction of sgRNAs were obtained from Integrated DNA Technologies, hybridized,

phosphorylated and cloned into the sgRNA plasmid or targeting vector using BbsI sites.[75] Target

sequences for sgRNAs are provided in Table S5. These vectors were transfected into HCT116

using Lipofectamine 2000 following recommendations from the manufacturer. Three days after

transfection, the cells were selected with Puromycin (0.5 µg/ml) and Hygromycin (100 µg/ml) to

generate clonal populations. Genomic DNA from each clone was isolated using DNEasy Blood

and Tissue Kit (Qiagen). PCRs to detect integration of the targeting vector at the target site were

performed using KAPA2G Robust PCR kits (Kapa Biosystems) according to the manufacturer's

instructions. A typical reaction contained 20–100 ng of genomic DNA in Buffer A (5 µl),

Enhancer (5 µl), dNTPs (0.5 µl), primers forward (PINCR Det FP, 1.25 µl) and reverse

(Targeting vector Det RP, 1.25 µl) and KAPA2G Robust DNA Polymerase (0.5 U). The DNA

sequences of the primers for each target are provided in Table S5. PCR products were visualized

in 2% agarose gels and images were captured using a ChemiDoc-It2 (UVP).

To generate MDA-MB-231 whole-locus and intron 2 *MALAT1* knockout clones, we used

the CRIPSR/Cas9 system by transfecting the cells with gRNAs and purified Cas9. 2 µl of each

30 µM gRNA was mixed with 1 µl (~5 µg) of purified Cas9 and 100 µl of 4D-Nucleofector® X

SE buffer SE from the Cell Line Kit (Lonza) supplemented with 18.5 µl of SE supplement per

100 µl total buffer volume. The mixture was incubated at room temperature for 10 min and

1×10^6 cells were resuspended in nucleofection cuvettes. Nucleofections were carried out using

the 4D-Nucleofector® System (Lonza). The cells were transferred to a 6-well plate, and they

were single-cell sorted 48 h later. The clones were grown for 3-4 weeks before genomic DNA

extraction and genotyping with PCR using the primers in Table S5.

PURPL and MALAT1 overexpression and transduction

PURPL intron 2 constructs were transduced in previously generated *PURPL* CRISPRi SKHEP1

cells.[43] *MALAT1*-WT and deletion mutants were transduced in the two HCT116-*MALAT1* KO

clones. For lentivirus generation, HEK293T cells were seeded at 2×10^5 cells/well onto 6-well

plates, and 1,200 ng of each construct was transfected with lentiviral package vectors using

Lipofectamine 2000 (Life Technologies, Invitrogen) according to the manufacturer's protocol.

Virus-containing media were harvested at 48 and 72 h post-transfection and added to cells for 24

h. Transduced cells were selected with 500 µg/mL hygromycin for 2 weeks (SKHEP1) or 10

µg/ml Blasticidin (HCT116).

RNA FISH, IF and Quantification of FISH

The *PURPL* intron probe set was custom-designed using the Stellaris Probe Designer, tagged

with Quasar 570 dye and purchased from Stellaris. The *MALAT1* smFISH probe set with Quasar

670 dye was purchased from Stellaris (Cat# VSMF-2211-5). For *PURPL* intron smRNA-FISH,

SKHEP1 cells were seeded on coverslips and cultured for 24 hours at 37°C with 5% CO_2 before

inducing DNA damage by incubating the cells with 2 mM hydroxyurea for 24 hours (or regular

growth media for control). For *MALAT1* smRNA-FISH, HCT116 and MDA-MB-231 *MALAT1*

KO cells were seeded on coverslips and cultured for 24 hours at 37°C with 5% CO_2. *MALAT1*

WT, *MALAT1* del-1, *MALAT1* del-2, *MALAT1* del1+2, or empty vector expression was induced

by incubating the cells with 1 µg/mL doxycycline for 48 hours. The cells that were transfected

with siRNA for U2AF2 were seeded on coverslips at the time of transfection.Cells were fixed

with 4% PFA for 15 min at room temperature, permeabilized with 0.5% Triton X-100, and

washed with washing buffer (10% formamide, 2XSSC) for 5 min. Probe was added to

hybridization buffer (10% formamide in Stellaris hybridization buffer Cat# SMF-HB1-10) at a

final concentration of 125 nM. Hybridization was done in a humidified chamber in the dark

overnight at 37°C. After hybridization, the coverslips were washed twice with wash buffer for 30

min at 37°C and then post-fixed with 4% PFA for 10 min at room temperature. The coverslips

were blocked with 5% normal goat serum and then incubated with SON antibody (Sigma Cat#

HPA023535). Primary and secondary antibodies were diluted in 1% normal goat serum.

Antibody incubations were done at room temperature for 1 hour and washed with PBS. DNA

was counterstained by DAPI. The coverslips were then washed with 4XSSC for 5 min at room

temperature and mounted in VectaShield Antifade Mounting Medium (Vector Laboratories).

Images were taken using DeltaVision microscope (GE Healthcare) equipped with 60 X/1.42 NA

oil immersion objective (Olympus) and CoolSNAP-HQ2 camera. To quantify the speckle to

nucleoplasm signal intensity ratio of *MALAT1* in Figures 5B, 7B, 7D, and S8B ImageJ was used

to create nucleus masks based on DAPI and speckles mask based on SON. The signal intensity

of *MALAT1* in the speckles mask was divided by the *MALAT1* signal intensity in the nucleus-

minus-speckles area. For the quantification of Figures 6B, 7E and S10B, the cells with *MALAT1*

FISH signal showing prominent speckle patterns similar to SON were classified as "Speckle"

pattern, and the cells with a diffused *MALAT1* FISH signal were classified as "Diffused". Only

MALAT1-positive cells were counted for quantification.

Transwell cell migration assays

For the transwell assay, WT and KO MDA-MB-231 clones were seeded at 10 x 10^5 cells per

well in 12-well plates. 24 hr later, the media were removed and replaced with media containing

1% FBS. After 5 hr, the cells were trypsinized and seeded in the top chamber of the transwell

plates (Corning, #354578) at 25,000 cells per well at a volume of 100 μl. The lower chamber of

the transwell is filled with 700 μl of media containing 10% FBS. The cells were incubated for 24

hr and the transwell were removed from the plates. The non-migrated cells were scraped with a

cotton swab and the migrated cells were fixed with ice-cold methanol for 5 min and stained with

0.05% crystal violet blue. The migrated cells were counted under a light microscope.

Statistics

Statistics were performed using the Student's t-test.

Data availability

The Iso-Seq data files have been deposited at GEO with accession number GSE288684 and the RNA-Seq data files have been deposited at GEO with accession number GSE288685.

Acknowledgments

This research was supported by the Intramural Research Program (A.L.) of the National Cancer Institute (NCI), Center for Cancer Research (CCR) (project number ZIA BC011646), NIH. KVP laboratory is supported by grants from NIH (R01-GM132458), Cancer center at Illinois seed grant, ARPA-H, and National Science Foundation (NSF) center for quantitative cell biology. YJS is supported by NIH T32EB019944. We thank the CCR Genomics Core, CCR, NCI, Bethesda, MD for its valuable assistance with Sanger and Nanopore sequencing. We also thank the Sequencing Facility, CCR, NCI, Frederick, MD for performing the RNA sequencing assays. The contributions of the NIH author(s) were made as part of their official duties as NIH federal employees, are in compliance with agency policy requirements, and are considered Works of the United States Government. However, the findings and conclusions presented in this paper are those of the author(s) and do not necessarily reflect the views of the NIH or the U.S. Department of Health and Human Services.

Author Contributions

IG, CN, YJS, AB, CCH, SK, RP, XLL, RK, RS, and PPZ conducted experiments. IG, EP, YZ, BS, TB, XW, TGP, KP, and AL did data analysis. IG and AL wrote the manuscript. NC, KP, and TGP edited the manuscript.

Declaration of interests

The authors declare they have no competing interest.

Figure legends

Figure 1. CRISPR-based screen identifies U2AF2 as a promoter of intron retention. (A)

Schematic of the *PURPL* IR reporter used in the screen to identify IR regulators. The pLCHKO

vector contains the puromycin resistance gene used for selection. The transgene also contains the

genomic sequences of exon 2 (161 bp), intron 2 (1,943 bp), and exon 3 (163 bp) of the *PURPL*

gene under a Doxycycline-inducible promoter. A hybrid guide (hg)RNA for guiding SpCas9 and

LbCas12a is expressed under a U6 promoter. A construct with a hgRNA targeting an intergenic

region was used as a negative control. Top arrow: The hgRNA is cleaved by Cas12a to produce 2

guide RNAs for dual targeting of each protein coding gene, paralog targeting, and genetic

interaction mapping. Bottom arrow: Upon intron splicing or intron retention, two different reads

are generated. **(B)** Two different cell lines, stably expressing Cas9 and Cas12a, are used: HAP1,

and RPE1. The pipeline before RNA-seq is depicted. **(C)** Heatmap showing single genes only

(after removing gene pairs), whether identified individually or as part of a gene pair. The hits are

ordered by the average ΔPIR values across HAP1 and RPE1, from lowest to highest. **(D)** Left:

Gene ontology enrichment analysis of biological processes regulated by the hits of the screen.

The most significant pathways involve mRNA processing and RNA splicing. Right: Gene

ontology enrichment analysis of cellular components regulated by the hits of the screen. These

components include mainly Spliceosomal Complex, small nuclear ribonucleoprotein complex

and the Catalytic step 2 Spliceosome. **(E)** Venn diagram showing that there are 5 common hits

which cause a reduction in the percentage of intron retention (ΔPIR) of the splicing reporter in

the two cell lines. **(F)** Table showing the top 5 hits of the screen with a ΔPIR for each hit in each

cell line and their full names.

Figure 2. U2AF2 binds to intron 2-containing *PURPL* transcripts and promotes intron 2

retention. (A) IGV snapshot of eCLIP-seq data showing binding sites and enrichment of U2AF2

and U2AF1 on the *PURPL* transcripts around intron 2. eCLIP-seq data for PTBP1, PRPF8, and

SRSF1 are indicated. The annotated locus by RefSeq is also indicated. eCLIP data-seq was

downloaded from encodeproject.org. **(B)** RT-qPCR after RNA-IPs using a U2AF2 antibody with

primer pairs specifically detecting *PURPL* transcripts as indicated in Figure S1C. U2AF2 binds

to transcripts containing intron 2 but not the ones with intron 1, intron 3, or spliced exons 2 and

3. Samples were normalized to IgG-IP. 18S was used as a loading control. **(C)** Top panel: Gel

with RT-PCR products for *PURPL* upon knockdown of U2AF2 with 2 different siRNAs in

SKHEP1 cells. The schematics next to the gel indicate the expected products of the intron-

retained and spliced isoforms. Between the two expected PCR products, we observed an extra

band corresponding to the inclusion of an alternative exon inside *PURPL* intron 2 as observed in

RefSeq, the inclusion of which is not affected by U2AF2. Quantitation of the gel bands is shown

in the graphs on the right. Bottom panel: Schematic of the PCR primer triplet used to detect

intron 2 retention (red) or splicing (purple). The length for each PCR product is indicated. **(D)**

RNA-FISH images for *PURPL* with intron 2 retention and *MALAT1* in HCT116 cells without

treatment or after 24 hr of 2 mM of Hydorxyurea (HU) to induce *PURPL* expression. Scale bar is

10μm. **(E)** RT-qPCR for intron 2-containing *PURPL* transcript after 48 hr of 1μg/ml doxycycline

treatment in comparison to no treatment in SKHEP1 *PURPL*-CRISPRi populations using 3

different gRNAs. **(F)** Proliferation assay showing the effect of overexpression of intron 2-

containing *PURPL* transcript in the proliferation of SKHEP1 cells where the endogenous *PURPL*

is knocked down with CRISPRi. The graph depicts the average of 3 populations with different

gRNAs. The cells were treated with 1ug/ml doxycycline to induce intron 2-retained *PURPL*

expression and cell proliferation was monitored at 3 and 6 days. Error bars represent standard

deviations from 2 **(E,)** and 3 **(C)** experiments. **p<0.01, ***p<0.001.

Figure 3. U2AF2 directly promotes Intron Retention of *MALAT1*. (A) U2AF2 was knocked

down in SKHEP1 cells and 72 hr later, RNA was extracted and RNA-seq was performed. *Left*:

Number of decreased (blue) and increased (red) IR events at various p-values after U2AF2

knockdown as analyzed with the IR Finder algorithm. The purple arrow indicates the *PURPL* IR

event and green arrows indicate MALAT IR events *Right*: Pie chart of the numbers of increased

and decreased IR events upon U2AF2 knockdown. **(B)** Floating bar plot showing the IR ratio of

PURPL intron 2 and the IR ratio of intron 1 (middle) and intron 2 (right) of *MALAT1*. siCTRL

and siU2AF2#1 samples as analyzed with the IRFinder algorithm. **(C)** and **(E)** RT-PCR for

MALAT1 using a primer pair flanking the regulated intron 1 **(C)** or intron 2 **(E)** upon knockdown

of U2AF2 with 2 different siRNAs in HCT116 and SKHEP1 cells. The schematics next to the

gel indicate the expected products of the intron-retained and spliced isoforms. **(D)** and **(F)** Bar

graph with quantitation of the gel bands from **(C)** and **(E)** in SKHEP1 cells. Error bars represent

standard deviations from 2 independent experiments. *p<0.05, **p<0.01, ***p<0.001.

Figure 4. SON depletion rescues the effect of U2AF2 knockdown in *PURPL* and *MALAT1*

intron 2 retention. (A) and **(C)** RT-PCR for *PURPL* **(A)** and *MALAT1* **(C)** transcripts upon

knockdown of U2AF2, knockdown of SON or simultaneous knockdown indicating a rescue

effect. The schematics next to the gel indicate the expected products of the intron-retained and

spliced isoforms. **(B)** and **(D)** Quantitation of the gel bands from **(A)** and **(C)**. Error bars

represent the SD of 2 experiments. **(E)** Western blot for SON (upper panel) and U2AF2 (middle

panel) showing successful knockdown of SON and U2AF2. GAPDH was used as a loading

control (lower panel). **(F)** RT-qPCR after RNA-IP using a SON antibody with primer pairs

specifically detecting *MALAT1* or *PURPL* transcripts upon U2AF2 knockdown as indicated in

Figure S1C. Transcript -specific primers for *MALAT1* are shown in Table S5. SON binds to

spliced transcripts more strongly upon U2AF2 depletion. Samples were normalized to input.

GAPDH was used as a loading control. The graph shows a representative of 2 replicates.

Figure 5. U2AF2 promotes localization of *MALAT1* to nuclear speckles. (A) RNA-FISH

images for *MALAT1* and Immunofluorescence images for SON is shown upon transfection of

SKHEP1 cells with siCTRL or siU2AF2. *MALAT1* is enriched in nuclear speckles in the siCTRL

but not upon U2AF2 knockdown. **(B)** Quantitation of the speckle to nuclear plasma *MALAT1*

signal ratio in the three replicates in panel **(A)**. ****p<0.0001.

Figure 6. Intron 2 of *MALAT1* dictates its localization to nuclear speckles. (A) RNA-FISH

images for *MALAT1* and Immunofluorescence images for SON in a HCT116 *MALAT1*-KO clone

(clone 1) where Empty Vector (EV) or constructs that exogenously express *MALAT1* transcripts

were re-introduced using a doxy-inducible lentivirus. Expressed *MALAT1* transcripts were full-

length (WT) or intron 1 deleted (del-1) or intron 2 deleted (del-2) or both introns deleted

simultaneously (del-1+2). The cells were first treated with 1ug/ml doxycycline for 48 hr to

induce *MALAT1* expression. *MALAT1* without intron 2 does not enrich in speckles as it is shown

with SON protein. Scale bar is 10μm. **(B)** Graph showing quantification of the percentage of

cells where *MALAT1* transcripts are enriched in speckles or being diffused in HCT116 cells

clone 1. Error bars represent standard deviations from 2 independent experiments.

Figure 7. Intron 2 regulates the migration capacity of *MALAT1* in breast cancer cells. (A)

RNA-FISH images for *MALAT1* and Immunofluorescence images for SON are shown upon

transfection of MDA-MB-231 cells with siCTRL or siU2AF2. *MALAT1* is enriched in nuclear

speckles in the siCTRL but not upon U2AF2 knockdown. **(B)** Quantitation of the speckle to

nuclear plasma *MALAT1* signal ratio in the three replicates in panel **(A)**. **(C)** RNA-FISH images

for *MALAT1* and Immunofluorescence images for SON in MDA-MB-231 clones either WT, or

whole locus *MALAT1* deletion, or intron 2 deletion. When intron 2 is deleted, *MALAT1* does not

enrich in speckles as it is shown with SON protein. Scale bar is 10μm. **(D)** Quantitation of the

speckle to nuclear plasma *MALAT1* signal ratio in the three replicates in panel **(C)**. **(E)** Graph

showing quantification of the percentage of *MALAT1* transcripts enriched in speckles or being

diffused in MDA-MB-231 cells in panel **(C)**, N=2. **(F)** Bar graph showing the number of cells

that have migrated in transwell migration assays conducted in MDA-MB-231 WT, *MALAT1* KO

and *MALAT1* – intron 2 deletion clones. Intron 2 deletion leads to decreased migration potential.

The number of cells are the sum of cells from 5 different fields, N=4. *p<0.05, **p<0.01,

***p<0.001, ****p<0.0001.

References

1.   Wilkinson, M.E., Charenton, C., and Nagai, K. (2020). RNA Splicing by the

Spliceosome. Annu Rev Biochem *89*, 359-388. 10.1146/annurev-biochem-091719-

064225.

2.   Marasco, L.E., and Kornblihtt, A.R. (2023). The physiology of alternative splicing. Nat

Rev Mol Cell Biol *24*, 242-254. 10.1038/s41580-022-00545-z.

3.   Matera, A.G., and Wang, Z. (2014). A day in the life of the spliceosome. Nat Rev Mol

Cell Biol *15*, 108-121. 10.1038/nrm3742.

4.   Black, D.L. (2003). Mechanisms of alternative pre-messenger RNA splicing. Annu Rev

Biochem *72*, 291-336. 10.1146/annurev.biochem.72.121801.161720.

5.   Ruskin, B., Zamore, P.D., and Green, M.R. (1988). A factor, U2AF, is required for U2

snRNP binding and splicing complex assembly. Cell *52*, 207-219. 10.1016/0092-

8674(88)90509-0.

6.   Berglund, J.A., Chua, K., Abovich, N., Reed, R., and Rosbash, M. (1997). The splicing

factor BBP interacts specifically with the pre-mRNA branchpoint sequence UACUAAC.

Cell *89*, 781-787. 10.1016/s0092-8674(00)80261-5.

7.   Berglund, J.A., Abovich, N., and Rosbash, M. (1998). A cooperative interaction between

U2AF65 and mBBP/SF1 facilitates branchpoint region recognition. Genes Dev *12*, 858-

867. 10.1101/gad.12.6.858.

8.   Wu, S., Romfo, C.M., Nilsen, T.W., and Green, M.R. (1999). Functional recognition of

the 3' splice site AG by the splicing factor U2AF35. Nature *402*, 832-835.

10.1038/45590.

9.   Zorio, D.A., and Blumenthal, T. (1999). Both subunits of U2AF recognize the 3' splice

site in Caenorhabditis elegans. Nature *402*, 835-838. 10.1038/45597.

10.   Wang, E.T., Sandberg, R., Luo, S., Khrebtukova, I., Zhang, L., Mayr, C., Kingsmore,

S.F., Schroth, G.P., and Burge, C.B. (2008). Alternative isoform regulation in human

tissue transcriptomes. Nature *456*, 470-476. 10.1038/nature07509.

11.   Nilsen, T.W., and Graveley, B.R. (2010). Expansion of the eukaryotic proteome by

alternative splicing. Nature *463*, 457-463. 10.1038/nature08909.

12.   Pan, Q., Shai, O., Lee, L.J., Frey, B.J., and Blencowe, B.J. (2008). Deep surveying of

alternative splicing complexity in the human transcriptome by high-throughput

sequencing. Nat Genet *40*, 1413-1415. 10.1038/ng.259.

13.   Keren, H., Lev-Maor, G., and Ast, G. (2010). Alternative splicing and evolution:

953   diversification, exon definition and function. Nat Rev Genet *11*, 345-355.

954   10.1038/nrg2776.

955   14.   Braunschweig, U., Barbosa-Morais, N.L., Pan, Q., Nachman, E.N., Alipanahi, B.,

956   Gonatopoulos-Pournatzis, T., Frey, B., Irimia, M., and Blencowe, B.J. (2014).

957   Widespread intron retention in mammals functionally tunes transcriptomes. Genome Res

958   *24*, 1774-1786. 10.1101/gr.177790.114.

959   15.   Wong, J.J., Ritchie, W., Ebner, O.A., Selbach, M., Wong, J.W., Huang, Y., Gao, D.,

960   Pinello, N., Gonzalez, M., Baidya, K., et al. (2013). Orchestrated intron retention

961   regulates normal granulocyte differentiation. Cell *154*, 583-595.

962   10.1016/j.cell.2013.06.052.

963   16.   Monteuuis, G., Wong, J.J.L., Bailey, C.G., Schmitz, U., and Rasko, J.E.J. (2019). The

964   changing paradigm of intron retention: regulation, ramifications and recipes. Nucleic

965   Acids Res *47*, 11497-11513. 10.1093/nar/gkz1068.

966   17.   Eom, T., Zhang, C., Wang, H., Lay, K., Fak, J., Noebels, J.L., and Darnell, R.B. (2013).

967   NOVA-dependent regulation of cryptic NMD exons controls synaptic protein levels after

968   seizure. Elife *2*, e00178. 10.7554/eLife.00178.

969   18.   Hossain, M.A., Claggett, J.M., Edwards, S.R., Shi, A., Pennebaker, S.L., Cheng, M.Y.,

970   Hasty, J., and Johnson, T.L. (2016). Posttranscriptional Regulation of Gcr1 Expression

971   and Activity Is Crucial for Metabolic Adjustment in Response to Glucose Availability.

972   Mol Cell *62*, 346-358. 10.1016/j.molcel.2016.04.012.

973   19.   Cha, S., Hong, C.P., Kang, H.A., and Hahn, J.S. (2021). Differential activation

974   mechanisms of two isoforms of Gcr1 transcription factor generated from spliced and un-

spliced transcripts in Saccharomyces cerevisiae. Nucleic Acids Res *49*, 745-759.

10.1093/nar/gkaa1221.

20. Schmitz, U., Pinello, N., Jia, F., Alasmari, S., Ritchie, W., Keightley, M.C., Shini, S.,

Lieschke, G.J., Wong, J.J., and Rasko, J.E.J. (2017). Intron retention enhances gene

regulatory complexity in vertebrates. Genome Biol *18*, 216. 10.1186/s13059-017-1339-3.

21. Dvinge, H., and Bradley, R.K. (2015). Widespread intron retention diversifies most

cancer transcriptomes. Genome Med *7*, 45. 10.1186/s13073-015-0168-9.

22. Boutz, P.L., Bhutkar, A., and Sharp, P.A. (2015). Detained introns are a novel,

widespread class of post-transcriptionally spliced introns. Genes Dev *29*, 63-80.

10.1101/gad.247361.114.

23. Quinn, J.J., and Chang, H.Y. (2016). Unique features of long non-coding RNA

biogenesis and function. Nat Rev Genet *17*, 47-62. 10.1038/nrg.2015.10.

24. Ulitsky, I., and Bartel, D.P. (2013). lincRNAs: genomics, evolution, and mechanisms.

Cell *154*, 26-46. 10.1016/j.cell.2013.06.020.

25. Guo, C.J., Xu, G., and Chen, L.L. (2020). Mechanisms of Long Noncoding RNA Nuclear

Retention. Trends Biochem Sci *45*, 947-960. 10.1016/j.tibs.2020.07.001.

26. Zuckerman, B., and Ulitsky, I. (2019). Predictive models of subcellular localization of

long RNAs. RNA *25*, 557-572. 10.1261/rna.068288.118.

27. Clark, M.B., Johnston, R.L., Inostroza-Ponta, M., Fox, A.H., Fortini, E., Moscato, P.,

Dinger, M.E., and Mattick, J.S. (2012). Genome-wide analysis of long noncoding RNA

stability. Genome Res *22*, 885-898. 10.1101/gr.131037.111.

28. Cabili, M.N., Dunagin, M.C., McClanahan, P.D., Biaesch, A., Padovan-Merhar, O.,

Regev, A., Rinn, J.L., and Raj, A. (2015). Localization and abundance analysis of human

lncRNAs at single-cell and single-molecule resolution. Genome Biol *16*, 20.

10.1186/s13059-015-0586-4.

29.   Tilgner, H., Knowles, D.G., Johnson, R., Davis, C.A., Chakrabortty, S., Djebali, S.,

Curado, J., Snyder, M., Gingeras, T.R., and Guigo, R. (2012). Deep sequencing of

subcellular RNA fractions shows splicing to be predominantly co-transcriptional in the

human genome but inefficient for lncRNAs. Genome Res *22*, 1616-1625.

10.1101/gr.134445.111.

30.   Mele, M., Mattioli, K., Mallard, W., Shechner, D.M., Gerhardinger, C., and Rinn, J.L.

(2017). Chromatin environment, transcriptional regulation, and splicing distinguish

lincRNAs and mRNAs. Genome Res *27*, 27-37. 10.1101/gr.214205.116.

31.   Mukherjee, N., Calviello, L., Hirsekorn, A., de Pretis, S., Pelizzola, M., and Ohler, U.

(2017). Integrative classification of human coding and noncoding genes through RNA

metabolism profiles. Nat Struct Mol Biol *24*, 86-96. 10.1038/nsmb.3325.

32.   Gutschner, T., Hammerle, M., Eissmann, M., Hsu, J., Kim, Y., Hung, G., Revenko, A.,

Arun, G., Stentrup, M., Gross, M., et al. (2013). The noncoding RNA MALAT1 is a

critical regulator of the metastasis phenotype of lung cancer cells. Cancer Res *73*, 1180-

1189. 10.1158/0008-5472.CAN-12-2850.

33.   Hutchinson, J.N., Ensminger, A.W., Clemson, C.M., Lynch, C.R., Lawrence, J.B., and

Chess, A. (2007). A screen for nuclear transcripts identifies two linked noncoding RNAs

associated with SC35 splicing domains. BMC Genomics *8*, 39. 10.1186/1471-2164-8-39.

34.   Ji, P., Diederichs, S., Wang, W., Boing, S., Metzger, R., Schneider, P.M., Tidow, N.,

Brandt, B., Buerger, H., Bulk, E., et al. (2003). MALAT-1, a novel noncoding RNA, and

thymosin beta4 predict metastasis and survival in early-stage non-small cell lung cancer.

Oncogene *22*, 8031-8041. 10.1038/sj.onc.1206928.

35.   Bernard, D., Prasanth, K.V., Tripathi, V., Colasse, S., Nakamura, T., Xuan, Z., Zhang,

M.Q., Sedel, F., Jourdren, L., Coulpier, F., et al. (2010). A long nuclear-retained non-

coding RNA regulates synaptogenesis by modulating gene expression. EMBO J *29*,

3082-3093. 10.1038/emboj.2010.199.

36.   Tripathi, V., Ellis, J.D., Shen, Z., Song, D.Y., Pan, Q., Watt, A.T., Freier, S.M., Bennett,

C.F., Sharma, A., Bubulya, P.A., et al. (2010). The nuclear-retained noncoding RNA

MALAT1 regulates alternative splicing by modulating SR splicing factor

phosphorylation. Mol Cell *39*, 925-938. 10.1016/j.molcel.2010.08.011.

37.   Shinn, M.K., Tomares, D.T., Liu, V., Pant, A., Qiu, Y., Vitalis, A., Song, Y.J., Ayala, Y.,

Ruff, K.M., Strout, G.W., et al. (2026). Nuclear speckle proteins form intrinsic and

MALAT1-dependent microphases. Cell *189*, 832-852 e824. 10.1016/j.cell.2025.11.026.

38.   Spector, D.L., and Lamond, A.I. (2011). Nuclear speckles. Cold Spring Harb Perspect

Biol *3*. 10.1101/cshperspect.a000646.

39.   Sharma, A., Takata, H., Shibahara, K., Bubulya, A., and Bubulya, P.A. (2010). Son is

essential for nuclear speckle organization and cell cycle progression. Mol Biol Cell *21*,

650-663. 10.1091/mbc.e09-02-0126.

40.   Nakagawa, S., Ip, J.Y., Shioi, G., Tripathi, V., Zong, X., Hirose, T., and Prasanth, K.V.

(2012). Malat1 is not an essential component of nuclear speckles in mice. RNA *18*, 1487-

1499. 10.1261/rna.033217.112.

41.   Zhang, B., Arun, G., Mao, Y.S., Lazar, Z., Hung, G., Bhattacharjee, G., Xiao, X., Booth,

C.J., Wu, J., Zhang, C., and Spector, D.L. (2012). The lncRNA Malat1 is dispensable for

mouse development but its transcription plays a cis-regulatory role in the adult. Cell Rep

2, 111-123. 10.1016/j.celrep.2012.06.003.

42.　Li, X.L., Subramanian, M., Jones, M.F., Chaudhary, R., Singh, D.K., Zong, X., Gryder,

B., Sindri, S., Mo, M., Schetter, A., et al. (2017). Long Noncoding RNA PURPL

Suppresses Basal p53 Levels and Promotes Tumorigenicity in Colorectal Cancer. Cell

Rep *20*, 2408-2423. 10.1016/j.celrep.2017.08.041.

43.　Hartford, C.C.R., Shrestha, R.L., Pongor, L., Zhao, Y., Chen, X., Fromont, C.,

Chaudhary, R., Li, X.L., Pasterczyk, K.R., Kumar, R., et al. (2022). Context-Dependent

Function of Long Noncoding RNA PURPL in Transcriptome Regulation during p53

Activation. Mol Cell Biol *42*, e0028922. 10.1128/mcb.00289-22.

44.　Behera, A.K., Kim, J.J., Kordale, S., Pekovic, F., Damodaran, A.P., Kumari, B., Vidak,

S., Dickson, E., Xiao, M.S., Duncan, G., et al. (2025). RNA-coupled CRISPR screens

reveal ZNF207 as a regulator of LMNA aberrant splicing in progeria. Mol Cell.

10.1016/j.molcel.2025.12.003.

45.　Aregger, M., Xing, K., and Gonatopoulos-Pournatzis, T. (2021). Application of

CHyMErA Cas9-Cas12a combinatorial genome-editing platform for genetic interaction

mapping and gene fragment deletion screening. Nat Protoc *16*, 4722-4765.

10.1038/s41596-021-00595-1.

46.　Gier, R.A., Budinich, K.A., Evitt, N.H., Cao, Z., Freilich, E.S., Chen, Q., Qi, J., Lan, Y.,

Kohli, R.M., and Shi, J. (2020). High-performance CRISPR-Cas12a genome editing for

combinatorial genetic screening. Nat Commun *11*, 3455. 10.1038/s41467-020-17209-1.

47.　Gonatopoulos-Pournatzis, T., Aregger, M., Brown, K.R., Farhangmehr, S.,

Braunschweig, U., Ward, H.N., Ha, K.C.H., Weiss, A., Billmann, M., Durbic, T., et al.

(2020). Genetic interaction mapping and exon-resolution functional genomics with a

hybrid Cas9-Cas12a platform. Nat Biotechnol *38*, 638-648. 10.1038/s41587-020-0437-z.

48.   Xiao, M.S., Damodaran, A.P., Kumari, B., Dickson, E., Xing, K., On, T.A., Parab, N.,

King, H.E., Perez, A.R., Guiblet, W.M., et al. (2024). Genome-scale exon perturbation

screens uncover exons critical for cell fitness. Mol Cell *84*, 2553-2572 e2519.

10.1016/j.molcel.2024.05.024.

49.   Van Nostrand, E.L., Pratt, G.A., Shishkin, A.A., Gelboin-Burkhart, C., Fang, M.Y.,

Sundararaman, B., Blue, S.M., Nguyen, T.B., Surka, C., Elkins, K., et al. (2016). Robust

transcriptome-wide discovery of RNA-binding protein binding sites with enhanced CLIP

(eCLIP). Nat Methods *13*, 508-514. 10.1038/nmeth.3810.

50.   Consortium, E.P. (2012). An integrated encyclopedia of DNA elements in the human

genome. Nature *489*, 57-74. 10.1038/nature11247.

51.   Luo, Y., Hitz, B.C., Gabdank, I., Hilton, J.A., Kagda, M.S., Lam, B., Myers, Z., Sud, P.,

Jou, J., Lin, K., et al. (2020). New developments on the Encyclopedia of DNA Elements

(ENCODE) data portal. Nucleic Acids Res *48*, D882-D889. 10.1093/nar/gkz1062.

52.   Berhane, T., Holm, A., Karstensen, K.T., Petri, A., Ilieva, M.S., Krarup, H., Vyberg, M.,

Lovendorf, M.B., and Kauppinen, S. (2022). Knockdown of the long noncoding RNA

PURPL induces apoptosis and sensitizes liver cancer cells to doxorubicin. Sci Rep *12*,

19502. 10.1038/s41598-022-23802-9.

53.   Han, S., Li, X., Wang, K., Zhu, D., Meng, B., Liu, J., Liang, X., Jin, Y., Liu, X., Wen, Q.,

and Zhou, L. (2021). PURPL represses autophagic cell death to promote cutaneous

melanoma by modulating ULK1 phosphorylation. Cell Death Dis *12*, 1070.

10.1038/s41419-021-04362-8.

54.   Middleton, R., Gao, D., Thomas, A., Singh, B., Au, A., Wong, J.J., Bomane, A., Cosson,

B., Eyras, E., Rasko, J.E., and Ritchie, W. (2017). IRFinder: assessing the impact of

intron retention on mammalian gene expression. Genome Biol *18*, 51. 10.1186/s13059-

017-1184-4.

55.   Wu, J., Xiao, Y., Liu, Y., Wen, L., Jin, C., Liu, S., Paul, S., He, C., Regev, O., and Fei, J.

(2024). Dynamics of RNA localization to nuclear speckles are connected to splicing

efficiency. Sci Adv *10*, eadp7727. 10.1126/sciadv.adp7727.

56.   Martinez-Terroba, E., Plasek-Hegde, L.M., Chiotakakos, I., Li, V., de Miguel, F.J.,

Robles-Oteiza, C., Tyagi, A., Politi, K., Zamudio, J.R., and Dimitrova, N. (2024).

Overexpression of Malat1 drives metastasis through inflammatory reprogramming of the

tumor microenvironment. Sci Immunol *9*, eadh5462. 10.1126/sciimmunol.adh5462.

57.   Arun, G., Aggarwal, D., and Spector, D.L. (2020). MALAT1 Long Non-Coding RNA:

Functional Implications. Noncoding RNA *6*. 10.3390/ncrna6020022.

58.   Jacob, A.G., and Smith, C.W.J. (2017). Intron retention as a component of regulated gene

expression programs. Hum Genet *136*, 1043-1057. 10.1007/s00439-017-1791-x.

59.   Shao, C., Yang, B., Wu, T., Huang, J., Tang, P., Zhou, Y., Zhou, J., Qiu, J., Jiang, L., Li,

H., et al. (2014). Mechanisms for U2AF to define 3' splice sites and regulate alternative

splicing in the human genome. Nat Struct Mol Biol *21*, 997-1005. 10.1038/nsmb.2906.

60.   Naftelberg, S., Schor, I.E., Ast, G., and Kornblihtt, A.R. (2015). Regulation of alternative

splicing through coupling with transcription and chromatin structure. Annu Rev Biochem

84, 165-198. 10.1146/annurev-biochem-060614-034242.

61.   Herzel, L., Ottoz, D.S.M., Alpert, T., and Neugebauer, K.M. (2017). Splicing and

1111   transcription touch base: co-transcriptional spliceosome assembly and function. Nat Rev

1112   Mol Cell Biol *18*, 637-650. 10.1038/nrm.2017.63.

1113   62.   Wong, J.J., Gao, D., Nguyen, T.V., Kwok, C.T., van Geldermalsen, M., Middleton, R.,

1114   Pinello, N., Thoeng, A., Nagarajah, R., Holst, J., et al. (2017). Intron retention is

1115   regulated by altered MeCP2-mediated splicing factor recruitment. Nat Commun *8*,

1116   15134. 10.1038/ncomms15134.

1117   63.   Guo, R., Zheng, L., Park, J.W., Lv, R., Chen, H., Jiao, F., Xu, W., Mu, S., Wen, H., Qiu,

1118   J., et al. (2014). BS69/ZMYND11 reads and connects histone H3.3 lysine 36

1119   trimethylation-decorated chromatin to regulated pre-mRNA processing. Mol Cell *56*,

1120   298-310. 10.1016/j.molcel.2014.08.022.

1121   64.   Krchnakova, Z., Thakur, P.K., Krausova, M., Bieberstein, N., Haberman, N., Muller-

1122   McNicoll, M., and Stanek, D. (2019). Splicing of long non-coding RNAs primarily

1123   depends on polypyrimidine tract and 5' splice-site sequences due to weak interactions

1124   with SR proteins. Nucleic Acids Res *47*, 911-928. 10.1093/nar/gky1147.

1125   65.   Dumbovic, G., Braunschweig, U., Langner, H.K., Smallegan, M., Biayna, J., Hass, E.P.,

1126   Jastrzebska, K., Blencowe, B., Cech, T.R., Caruthers, M.H., and Rinn, J.L. (2021).

1127   Nuclear compartmentalization of TERT mRNA and TUG1 lncRNA is driven by intron

1128   retention. Nat Commun *12*, 3308. 10.1038/s41467-021-23221-w.

1129   66.   Desideri, F., Cipriano, A., Petrezselyova, S., Buonaiuto, G., Santini, T., Kasparek, P.,

1130   Prochazka, J., Janson, G., Paiardini, A., Calicchio, A., et al. (2020). Intronic

1131   Determinants Coordinate Charme lncRNA Nuclear Activity through the Interaction with

1132   MATR3 and PTBP1. Cell Rep *33*, 108548. 10.1016/j.celrep.2020.108548.

67.   Guo, C.J., Ma, X.K., Xing, Y.H., Zheng, C.C., Xu, Y.F., Shan, L., Zhang, J., Wang, S.,

Wang, Y., Carmichael, G.G., et al. (2020). Distinct Processing of lncRNAs Contributes

to Non-conserved Functions in Stem Cells. Cell *181*, 621-636 e622.

10.1016/j.cell.2020.03.006.

68.   Sharma, A., Markey, M., Torres-Munoz, K., Varia, S., Kadakia, M., Bubulya, A., and

Bubulya, P.A. (2011). Son maintains accurate splicing for a subset of human pre-

mRNAs. J Cell Sci *124*, 4286-4298. 10.1242/jcs.092239.

69.   Ahn, E.Y., DeKelver, R.C., Lo, M.C., Nguyen, T.A., Matsuura, S., Boyapati, A., Pandit,

S., Fu, X.D., and Zhang, D.E. (2011). SON controls cell-cycle progression by

coordinated regulation of RNA splicing. Mol Cell *42*, 185-198.

10.1016/j.molcel.2011.03.014.

70.   Barutcu, A.R., Wu, M., Braunschweig, U., Dyakov, B.J.A., Luo, Z., Turner, K.M.,

Durbic, T., Lin, Z.Y., Weatheritt, R.J., Maass, P.G., et al. (2022). Systematic mapping of

nuclear domain-associated transcripts reveals speckles and lamina as hubs of functionally

distinct retained introns. Mol Cell *82*, 1035-1052 e1039. 10.1016/j.molcel.2021.12.010.

71.   Kolberg, L., Raudvere, U., Kuzmin, I., Adler, P., Vilo, J., and Peterson, H. (2023).

g:Profiler-interoperable web service for functional enrichment analysis and gene

identifier mapping (2023 update). Nucleic Acids Res *51*, W207-W212.

10.1093/nar/gkad347.

72.   Lee, J.S., Dan, T., Zhang, H., Cheng, Y., Rehfeld, F., Brugarolas, J., and Mendell, J.T.

(2025). An ultraconserved snoRNA-like element in long noncoding RNA CRNDE

promotes ribosome biogenesis and cell proliferation. Mol Cell *85*, 1543-1560 e1510.

10.1016/j.molcel.2025.03.006.

73.   Brown, A., Winter, J., Gapinske, M., Tague, N., Woods, W.S., and Perez-Pinera, P.

1157   (2019). Multiplexed and tunable transcriptional activation by promoter insertion using

1158   nuclease-assisted vector integration. Nucleic Acids Res *47*, e67. 10.1093/nar/gkz210.

1159   74.   Brown, A., Woods, W.S., and Perez-Pinera, P. (2016). Multiplexed Targeted Genome

1160   Engineering Using a Universal Nuclease-Assisted Vector Integration System. ACS Synth

1161   Biol *5*, 582-588. 10.1021/acssynbio.6b00056.

1162   75.   Brown, A., Woods, W.S., and Perez-Pinera, P. (2017). Targeted Gene Activation Using

1163   RNA-Guided Nucleases. Methods Mol Biol *1468*, 235-250. 10.1007/978-1-4939-4035-

1164   6_16.

1165

A

*Sp*Cas9   *Lb*Cas12a

combinatorial genome editing

Cas9 guide | Cas12a guide

| Cas9 spacer | tracrRNA | direct repeat | Cas12a spacer |

hybrid guide (hg)RNA

- Dual targeting of each protein coding gene
- Paralogue targeting
- GI mapping

5′ LTR | BGH | U6 → Cas9 guide | Cas12a guide | **PURPL Splicing reporter**

| Cas9 spacer | tracrRNA | direct repeat | Cas12a spacer | CBC | Exon 3 — Exon 2 | FG | TRE ← rtTA DOX

hybrid guide (hg)RNA

3′ LTR | PGK | Puro | 2A | rtTA

P7 | i7 | Read_2 | IR | CBC | Cas12a spacer | Cas9 spacer | UMI | Read_1 | i5 | P5

P7 | i7 | Read_2 | Exon | CBC | Cas12a spacer | Cas9 spacer | UMI | Read_1 | i5 | P5

B   **Figure 1**

HAP1   RPE1

1. Total RNA extraction

2. oligo-dT RNA purification

3. cDNA synthesis (UMI incorporation)

4. PCR amplification (Dual indexes incorporation)

5. RNA-sequencing and analysis

C

U2AF2, SDE2, H2AC14, SRSF2, PRPF8, ZNF878, SRSF1, MNT, SSU72, SCAF1, NECTIN2, MEPCE, PDCD11, CTSK, GRN, CPSF1, HEXIM1, INTS2, INTS9, SNRNP70, EFTUD2, NCBP2, URM1, DMPK, BAMBI, DNAJC17, SF3A2, COPG1, DDX23, GTF2A1, SART3, CPSF2, UHMK1, CCNL2, SYMPK, SF3A1, PYM1, SLC16A9, EIF4A3, SF3B3, MAGOH, AP3S2, PRELID3A, FIP1L1, SF3B4, AP3S1, TRA2A, CCNL1, PSMA7, DHX8, SS18L1, PRELID3B, SF3A3, SLC16A14, ARCN1, BUD23, COPA, SS18, SF3B6, LARS1, KPNB1, EIF2S1

RPE1 / HAP1

ΔPIR   −20 −10 0 10 20

D

Gene Ontology Biological Process
- mRNA processing
- RNA splicing
- mRNA splicing, via spliceosome
- protein-RNA complex organization
- spliceosomal complex assembly
- U2-type prespliceosome assembly
- ribonucleoprotein complex biogenesis
- regulation of RNA splicing
- protein-RNA complex assembly
- RNA 3'-end processing

−log10 (FDR)

Gene Ontology Cellular Component
- spliceosomal complex
- small nuclear ribonucleoprotein complex
- catalytic step 2 spliceosome
- U2-type spliceosomal complex
- spliceosomal snRNP complex
- nuclear speckles
- precatalytic spliceosome
- U2 snRNP
- mRNA cleavage factor complex

−log10 (FDR)

E

HAP1 **45**   **5**   RPE1 **12**

F

HIT	ΔPIR HAP1	ΔPIR RPE1	Name
U2AF2	-28.42	-23.98	U2 Small Nuclear RNA Auxiliary Factor 2
SNRNP70	-16.05	-3.69	Small Nuclear Ribonucleoprotein U1 Subunit 70
EFTUD2	-14.64	-12.84	Elongation Factor Tu GTP Binding Domain Containing 2
DDX23	-9.49	-6.28	DEAD-Box Helicase 23
SART3	-2.01	-14.01	Spliceosome Associated Factor 3, U4/U6 Recycling Protein

Figure 2

Figure 3

Figure 4

A

SKHEP1

B

Figure 7

A MDA-MB-231

SON | MALAT1 | Merge + DAPI

siCTRL

siU2AF2 #1

siU2AF2 #2

B

C WT1 | MALAT1 KO | INT2 KO-1 | INT2 KO-2

SON

MALAT1

Merge+DAPI

D

E

F