A stepwise route to polyploidy in yeast

Cintia Gómez-Muñoz[1,2], Nina Vittorelli[1,2,3], Maxime Gaudin[4], Nicolas Agier[1,2], Stéphane Delmas[1,2], Yann Corbeau[1,2], Marco Cosentino Lagomarsino[5,6], Gianni Liti[7], Bertrand Llorente[4] and Gilles Fischer[1,2]*

[1] Sorbonne Université, CNRS, Computational, Quantitative and Synthetic Biology, CQSB, F-75005 Paris, France.

[2] Sorbonne Université, CNRS, Inserm, Institut de Biologie Paris-Seine, IBPS, F-75005 Paris, France.

[3] Centre Interdisciplinaire de Recherche en Biologie, Collège de France, CNRS, INSERM, Paris, France.

[4] CRCM, Aix Marseille Université, CNRS, INSERM, Institut Paoli Calmettes, Marseille, France.

[5] IFOM-ETS, AIRC institute of molecular oncology, Milan, Italy.

[6] Dipartimento di Fisica, Università degli Studi di Milano, Milan, Italy and INFN sezione di Milano, Milan, Italy.

[7] IRCAN, CNRS, INSERM, Côte d'Azur University, Nice, France.

*Corresponding author

Email address: gilles.fischer@sorbonne-universite.fr

Abstract

Polyploidy is a ubiquitous biological phenomenon that promotes genetic innovation and adaptation[1]. In the budding yeast *Saccharomyces cerevisiae*, polyploid strains are widespread in natural, industrial, and clinical niches, yet their origin is unclear[2–4]. Here, we identify a novel mechanism of stepwise polyploidization that involves a Sporulate-Endoreplicate-Mate (*SEM*) sequence. We show that spore endoreplication during germination doubles the genome while preserving mating competence, enabling incremental increases in ploidy through mating with a non-endoreplicated spore from the same ascus. We experimentally demonstrate the transition from diploidy to triploidy and from triploidy to tetraploidy. Stepwise polyploidization thus positions triploids as central intermediates in ploidy evolution rather than evolutionary dead-ends. We further show that spores from intact asci can spontaneously undergo one or two successive *SEM* cycles and generate novel triploid and tetraploid strains without genetic manipulation. Natural polyploids harbor specific genomic features that are consistent with the *SEM* mechanism, including the prevalence of triploidy, extensive aneuploidy, pervasive heterozygosity, and a strong association with heterothallism[4–11]. Together, our findings establish stepwise polyploidization through iterative *SEM* cycles as the predominant natural route to polyploidy in *S. cerevisiae*.

Main

Polyploidy, defined as the presence of more than two complete sets of chromosomes, is widespread across the tree of life, with both ancient and recent polyploidization events documented in all major lineages[1]. In plants and fungi, polyploidy promotes genomic and phenotypic diversification, facilitates adaptation to stress, and contributes to ecological success and domestication[3,4,12,13]. Although less common in animals, polyploidy has played an important evolutionary role, notably through two ancient rounds of whole genome duplication (WGD) at the base of vertebrate evolution[14], and remains prevalent in fish, amphibians, and some invertebrates[15–17]. Polyploidy also arises at the tissue level in multicellular organisms[1] and is a hallmark of cancer, where it fuels genome instability, tumor evolution, and therapy resistance[18–20]. Overall, polyploidy is a powerful driver of genetic innovation and is increasingly recognized as a multifaceted force shaping evolution, adaptation, and disease[1,21].

Autopolyploidization, the increase in chromosome sets within a single species, can arise through diverse mitotic or meiotic mechanisms, including somatic cell fusion, WGD through endoreplication or cytokinesis failure, polyspermy and meiotic restitution leading to unreduced gametes[1,21,22]. Somatic genome doubling produces exclusively even ploidy states, whereas the fusion of reduced and unreduced gametes can generate triploids and other odd ploidy levels. Although triploidy has often been viewed as an evolutionary dead end due to impaired fertility, triploids can potentially serve as a bridge in the transition from diploidy to tetraploidy[21,22].

In the budding yeast *Saccharomyces cerevisiae*, between 8% and 20% of natural isolates are autopolyploid and triploids are at least as frequent as tetraploids in natural populations[6–9], a pattern that remains difficult to reconcile with the canonical yeast life-cycle[2]. Despite early reports of triploid and tetraploid isolates more than seventy years ago[23] and laboratory demonstrations that triploids can be generated through mating-based routes[24], the natural mechanism underlying triploidization in yeast remains unknown. A prevailing model posits that *S. cerevisiae* undergoes saltational transitions from diploidy to tetraploidy through WGD (Fig. 1A), a view supported by the ancestral WGD event approximately 100 million years ago[25,26] and

observations of spontaneous autotetraploidization during vegetative growth[27–29]. In

this framework, triploids are viewed as transient intermediates arising from

chromosome loss during the reversion of tetraploids to diploids[31–33]. However, this

model does not account for the high frequency and apparent stability of triploids in

natural populations[33]. An alternative possibility would be stepwise polyploidization

through successive gains of chromosome sets, in which triploids arise directly from

diploids through mating with haploids, rather than as derivatives of tetraploids (Fig.

1A). Although at first sight such a stepwise route appears unlikely because natural

diploids do not mate and haploids exist only transiently between spore germination

and mating, we show that this double challenge is circumvented by a sequence of

events that we term Sporulate–Endoreplicate–Mate (*SEM*) (Fig. 1B). We

demonstrate that polyploidization through iterative *SEM* sequences is the

predominant natural route to polyploidy in *S. cerevisiae*, enabling stepwise

transitions from diploidy to triploidy and from triploidy to tetraploidy.

A

B

Figure 1

Figure 1: Saltational and stepwise routes to polyploidy in *S. cerevisiae*.
A, Schematic comparison between saltational and stepwise polyploidization, where x denotes ploidy level. In the saltational model, diploid cells ($2x$) undergo Whole Genome Duplication (WGD) to directly generate tetraploids ($4x$), whereas triploids ($3x$) arise secondarily through chromosome loss from tetraploids. In stepwise polyploidization, ploidy increases incrementally through successive Sporulate–Endoreplicate–Mate (*SEM*) cycles. **B**, Schematic representation of the *SEM* sequence, where $2n$ and n denote vegetative and gametic chromosome number, respectively. Sporulation of a $2n$ vegetative cell produces four $1n$ spores. Upon germination, one spore undergoes post-meiotic endoreplication, doubling the genome while preserving mating competence. Mating between the endoreplicated spore and a compatible spore from the same ascus generates a cell with increased ploidy.

The Sporulate-Endoreplicate-Mate sequence

In this study, we follow the terminology of Ramsey & Schemske[22], where $2n$ refers to the vegetative chromosome number and n to the gametic chromosome number, independently of ploidy. Ploidy levels are indicated by x, with diploids, triploids, and tetraploids corresponding to $2x$, $3x$, and $4x$, respectively.

The *SEM* sequence describes a succession of events that begins with *S*, the sporulation of a $2n$ vegetative cell which produces four $1n$ spores enclosed within an ascus (Fig 1B). This is followed by *E*, an endoreplication event that occurs upon spore germination. Here, endoreplication is used as a generic term encompassing both endocycling and endomitosis[34,35], reflecting the fact that the exact molecular mechanism underlying genome doubling is currently unknown. Endoreplication of a germinating spore generates a cell with a duplicated genome and carrying a single mating type that is thus capable of mating, while the ascus environment concomitantly supplies a compatible non-duplicatd partner. The final event is *M*, the mating between the endoreplicated spore and a compatible spore from the same ascus, resulting in a cell carrying a $+1x$ ploidy increment (Fig. 1B). Each iteration of the *SEM* sequence increases ploidy by one chromosome set (Fig. 1A).

Below, we provide experimental evidence for each step of *SEM* sequence. Additionally, we were able to experimentally recapitulate the full process of stepwise ploidy increases $2x \rightarrow 3x \rightarrow 4x$ through two iterations of the *SEM* sequence. Importantly, the stepwise polyploidization model predicts the frequent emergence of both triploids and tetraploids and suggests that increasing ploidy above $4x$ is rare. Finally, we verified that the genomic characteristics of most natural polyploids are

120   fully compatible with their stepwise origin and that the saltational route accounts

121   for a minority of cases.

Stepwise diploid-to-triploid transition

Iteration one of the *SEM* sequence starts with *S1* the sporulation of a diploid parental

cell and is directly followed by *E1* the endoreplication of one of the germinating

spores (Fig. 2A). Previous studies showed that endoreplication occurs frequently in

vegetative cells, with approximately 17% of haploids spontaneously diploidizing

after 100 generations in the absence of external stress, and can even reach 100% in

the presence of EtOH or KCl[36]. Endoreplicated strains are fully homozygous across

the entire genome, hence carry a single mating type enabling them to mate.

To detect endoreplication in natural populations, we screened a collection of 742

natural diploids using a combination of mating assays (Extended Methods) to

identify endoreplicated strains capable of mating but persisted without finishing the

SEM sequence. As expected, most strains exhibited no or extremely low levels of

mating activity. Nevertheless, a subset of strains showed clear and reproducible

mating signals (Table S1, Fig. 2B). We further characterized these strains by PCR

genotyping the *MAT* locus. We identified 26 single mating type diploids capable of

mating. About half of these were heterozygous at the genome level, suggesting they

arose through *MAT* locus hemizygotization or homozygotization (Fig. 2B). These

strains could contribute to polyploidization upon outcrossing, independently from

the *SEM* sequence. The other eleven strains were fully homozygous, indicating that

they most likely resulted from endoreplication (Fig. 2B). They belong to eight

different clades suggesting that endoreplication is pervasive across *S. cerevisiae*

populations (Table S1).

This initial screen provides clear evidence that spontaneous endoreplication occurs

at measurable frequencies in natural yeast populations. To determine whether

endoreplication can occur concomitantly with spore germination, as proposed for

the *E1* step of the *SEM* sequence (Fig. 2A), we screened 23 homothallic and 9

heterothallic diploid strains for their capacity to undergo spontaneous spore

endoreplication. We sporulated all 32 strains, measured the ploidy by flow

cytometry of 684 microdissected spores from their meiotic progeny, and determined

their mating genotype (Table S2, Table S3). Homothallic diploids produce haploid
spores that self-mate through mating-type switching, generating *MATa*/*MATα*
diploid monosporic isolates. As expected, all 23 homothallic diploids produced
exclusively *MATa*/*MATα* diploid monosporic isolates, and none showed evidence
of endoreplication, which would have resulted in tetraploid monosporic isolates
(Fig. 2C). In contrast, heterothallic diploids produce haploid spores that cannot
switch mating-type and therefore develop into single-mating-type haploid
monosporic isolates. Consistent with this expectation, all heterothallic strains
produced single–mating-type haploid monosporic isolates (Fig. 2C). Notably, five
of the nine heterothallic strains also produced ~2*x* single-mating-type monosporic
isolates, suggesting that endoreplication occurred after spore germination.

We next tested whether these putatively endoreplicated isolates were able to
generate triploid strains by crossing them with compatible haploid testers
(corresponding to the *M1* step in Fig. 2A). The ~2*x* monosporic isolates derived from
the two heterothallic strains AKR and ALH, both belonging to the Asian
fermentation clade, produced triploid progeny upon mating, confirming that these
isolates were endoreplicated (Supplementary Fig. 1).

Together, these results demonstrate that (i) a subset of natural isolates likely
originates from spore endoreplication, (ii) endoreplication can occur upon spore
germination, and (iii) endoreplicated isolates are capable of generating triploid
strains. These findings further reinforce the strong association between
heterothallism and natural polyploidy[5], as only heterothallic diploids generated
triploid isolates through the *SEM* pathway.

Figure 2

Figure 2: Endoreplication during spore germination enables diploid-to-triploid transition
A, Schematic of the first iteration of the *SEM* sequence, in which sporulation (*S1*), endoreplication (*E1*), and mating (*M1*) collectively enable the transition from diploid to triploid. **B**, Identification of natural diploid strains capable of mating. Mating scores include mass mating assays (medium shading), observation of zygote formation (dark shading) and creeping phenotypes (light shading), revealing a subset of diploid isolates exhibiting reproducible mating activity, as *a*-mater (blue) or *α*-mater (red). *MAT* locus hemizygotization corresponds to chromosome III monosomy. Genome-wide homo and heterozygosity are symbolized by open and stripped circles, respectively. Fully homozygous single-mating-type diploids are inferred to result from haploid endoreplication. **C**, Ploidy and mating-type composition of monosporic isolates derived from homothallic and heterothallic diploid strains. Homothallic strains produce exclusively *MATa*/*MATα* diploids (purple), whereas heterothallic strains generate haploid *MATa* (blue) or *MATα* (pink). Black crosses indicate parental strain ploidy. The shaded area indicates the ploidy range expected from endoreplication. Several heterothallic diploids produce monosporic isolates consistent with endoreplication. The homothallic CAS strain produces monosporic isolates showing increased ploidy compared to the parental, although lower than expected for endoreplication.

Stepwise triploid-to-tetraploid transition

The second iteration of the *SEM* sequence allows stepwise acquisition of tetraploidy and begins with *S2*, the sporulation of a triploid parental cell (Fig. 3A). Among 46 triploids from the 1,011 collection, 38 (87.5%) were able to sporulate, with efficiency ranging from 3% to 97% (median 44%)[37]. We microdissected 180 spores from the meiotic progeny of each of three representative euploid triploid strains and confirmed that spore viability was extremely low (5.0–7.8%; Supplementary Fig. 2, Table S3). This could be due to chromosome missegregation or mutational load. To disentangle their respective contribution, we constructed a fully homozygous triploid strain (AKN-3x) and measured its gamete viability. This artificial triploid showed 56.7% gamete viability (Supplementary Fig. 2), versus 96% in the original AKN diploid strain (Table S3), indicating that segregation defects alone reduce fertility by approximately 2-fold in triploids, as previously reported[38]. The even lower viability of natural triploids (5.0–7.8%) highlights the additional contribution of mutational load in their genomes, estimated to comprise several heterozygous high-impact variants in essential genes[39].

Because triploid cells carry an odd number of homologs for each chromosome, segregation at meiosis I is necessarily unbalanced, and triploid meiosis is therefore expected to produce spores with variable chromosomal contents ranging from 1x to 3x (expected mean 1.5x, Fig. 3A). Genome sequencing of 32 monosporic isolates from

three natural euploid triploids showed that all spore genomes were aneuploid, *i.e.* none were perfectly haploid, diploid or triploid (Fig. 3B; Table S2).

We identified genomic signatures of *E2*, the endoreplication step of the *SEM* sequence (Fig. 3A), in six of the 32 sequenced monosporic isolates (Supplementary Fig. 3). First, these isolates contain multiple chromosomes present in four copies (Fig. 3B). Because regular meiotic segregation of parental triploid genomes can generate at most three copies of a chromosome per spore, tetrasomes are most parsimoniously explained by endoreplication of originally disomic chromosomes. Second, these six isolates are completely devoid of monosomes, in contrast to their widespread presence in non-endoreplicated spores, suggesting that all originally monosomic chromosomes were converted into disomes by endoreplication (Fig. 3B). Third, allele balance ratios (*ABRs*) at heterozygous sites show ~50% heterozygosity for tetrasomes and complete homozygosity for disomes in all six isolates, as expected under endoreplication, although three isolates also exhibit additional *ABR* complexity affecting a small number of chromosomes (Fig. 3C, Supplementary Fig. 3). Additionally, the six isolates exhibited ploidy levels ranging from $2.2x$ to $3.0x$ (mean $2.6x$), significantly higher than the $1.1x$ to $2.4x$ range (mean $1.7x$) observed in the other isolates (Fig. 3D). Together, these findings indicate that endoreplication occurs at appreciable frequency in triploid progeny (6 of 32, 19%), supporting the endoreplicate step of the second iteration of the *SEM* sequence (Fig. 3A).

We assessed the mating ability of the six endoreplicated monosporic isolates, along with 26 non-endoreplicated isolates, by genotyping their mating types and evaluating mating proficiency using mass mating assays. Five of the six isolates behaved as *MATα* strains, while the remaining one was a non-mater (Table S2). To validate the *M2* step of the *SEM* sequence (Fig. 3A), we crossed the five mating-competent endoreplicated isolates with a haploid tester strain and measured the ploidy of the resulting progeny. All crosses resulted in increased ploidy, with increments ranging from $+0.7x$ to $+2.1x$, demonstrating the stepwise transition from triploidy to tetraploidy (Supplementary Fig. 4).

Together, these results show that triploid sporulation is frequent but inefficient, producing few viable spores. Nonetheless, these viable progenies are highly

242   aneuploid, often undergo endoreplication, and can act as mating-competent
243   partners, enabling stepwise increases in ploidy toward tetraploidy.

Figure 3

Figure 3: Stepwise triploid-to-tetraploid transition through the second iteration of the SEM sequence.
A, Schematic of the second *SEM* iteration (*S2–E2–M2*) initiated by sporulation of a triploid parental strain. Triploid meiosis produces highly aneuploid spores with variable chromosome copy numbers that average to 1.5*x*. The mating genotype of the parental triploid strain is *MATa/α/α*. **B**, Chromosome copy number profiles of 32 monosporic isolates derived from triploid meiosis, showing extensive aneuploidy. Six isolates (bolded names) display tetrasomic chromosomes as well as complete absence of monosomic chromosomes, indicative of post-meiotic endoreplication. **C**, Genome-wide (left) and chromosome-level (right) copy number (top) and allele balance ratio (*ABR*; bottom) profiles of the T317c monosporic isolate. Chromosome copy numbers correspond to the sequencing depth normalized by the genome ploidy. Centromeres are marked with black circles and the rounded chromosome copy number is indicated in colored circles. In *ABR* profiles, only alternative alleles are plotted. **D**, Inferred ploidy distributions of monosporic isolates, highlighting increased ploidy in endoreplicated progeny (the 6 isolates in bold in panel B are highlighted). The shaded area indicates the approximate ploidy range expected from endoreplication in monosporic isolates.

Spontaneous generation of novel polyploids through stepwise polyploidization

A key assumption of the *SEM* model is that endoreplication and mating occur sequentially and rapidly after spore germination within the same ascus. To directly test this prediction, we designed an experimental system in which diploid strains were sporulated and intact asci, *i.e.* without spore dissection, were individually micromanipulated and plated onto rich medium. This setup allowed spore germination, potential endoreplication, and intra-ascus mating to occur spontaneously under laboratory conditions, after which ascus viability was assessed and the ploidy of the resulting colonies was determined (Fig. 4A, Table S4).

We micromanipulated asci from 12 diploid strains naturally defective for mating type switching due to loss of function mutations in the *HO* gene. We focused on heterothallic strains because most natural polyploids also lack a functional *HO* gene[5], and because our results showed that only heterothallic spores endoreplicated (Fig. 2C). These strains span a broad genetic diversity, having been isolated from both natural and anthropologic niches across multiple geographic regions (Table S2). In total, we micromanipulated 1,191 intact asci (50 - 150 asci per strain). Ascus viability varied widely among strains (Fig. 4B). We recovered both haploid and diploid monoascus isolates. Haploid monoascus isolates corresponded to unmated monosporic progeny, whereas diploid isolates most likely resulted from intra-ascus mating between compatible spores (Fig. 4B). Notably, five of the twelve tested

diploid strains produced triploid monoascus isolates at frequencies ranging from 1.2% to 9.1% (Fig. 4B). In total, nine triploid isolates were recovered, subcloned, and confirmed as triploid by both flow cytometry and whole-genome sequencing (Table S4 and S5). Chromosome copy number and *ABR* profiles displayed values characteristic of triploidy (Supplementary Fig. 6). Together, these results demonstrate that sporulation of a heterothallic diploid strain can spontaneously generate novel triploid progeny through at least one complete iteration of the *SEM* sequence (Fig. 2A).

We next tested whether these newly formed triploids could undergo a second iteration of the *SEM* sequence and spontaneously generate tetraploid progeny. A randomly selected novel triploid isolate from each of two genetic backgrounds (ALH and CNT) was resporulated. In both cases, ascus viability was reduced relative to the diploid parent, consistent with impaired triploid meiosis (Fig. 4C). The resulting monoascus isolates displayed a range of ploidies, including haploid and diploid isolates, indicating that ploidy often regressed through meiosis as expected. However, other isolates were triploid, suggesting that ploidy was restored through sporulation followed by mating within the same ascus (Fig. 4C). Importantly, the triploid isolates generated in this process carry novel allelic combinations relative to their parental genotypes and potentially altered aneuploidy landscapes resulting from meiotic resegregation of trivalents. Thus, even without further increase in ploidy, these isolates enable exploration of novel genotypic space. Notably, among the ascus progeny of ALH-3x, we identified a tetraploid isolate (ALH-4x; Fig. 4C). This isolate was subcloned, and its ploidy was validated by both flow cytometry and whole-genome sequencing (Fig. 4D and E). Comparison of chromosome *ABR* profiles between ALH-3x and ALH-4x revealed the expected +1x increase in genome content, providing direct evidence that ploidy can further increase through a second iteration of the SEM sequence (Fig. 3A).

Together, these results demonstrate that the entire ploidy transition, from diploid to tetraploid through intermediate triploid states, can be fully recapitulated under laboratory conditions and harnessed to generate polyploids with novel genetic compositions.

Figure 4

A barrier to increase ploidy beyond tetraploidy

We explored whether further stepwise ploidy increase beyond tetraploidy was

achievable through the *SEM* sequence. We found that 25 of 38 (65.8%) tetraploids

efficiently sporulate as 25 of 38 (65.8%) were able to sporulate, with a median

efficiency of 57%[37]. Using two representative strains, we confirmed the low spore

viability of tetraploids (6.7% to 12.2%, Supplementary Fig. 2). We constructed an

artificial homozygous tetraploid (AKN-4x) to estimate gamete viability in the

absence of deleterious recessive mutations. This strain showed 77.2% spore viability,

significantly higher than the 56.7% in the homozygous triploid AKN-3x

(Supplementary Fig. 2), showing that chromosome segregation is less affected in

tetraploid than in triploids, as previously reported[41]. Consistently, genome

sequencing of 28 monosporic isolates obtained in the meiotic progeny of two natural

euploid tetraploids revealed very few monosmic chromosomes as well as a much

lower global incidence of aneuploidies than in triploid progenies (Supplementary

Fig. 5A). Eleven isolates were euploid diploids while the remaining showed either

monosomies (in seven cases) and/or trisomies (in 14 cases). Thus, the low spore

viability of tetraploids probably reflects the major contribution of recessive

mutational load exposed at homo- or hemizygous state in the gametes. No sign of

spore endoreplication was detected in the progeny of tetraploids, suggesting that

further ploidy increase through the *SEM* sequence is limited, or at least less frequent than in the progeny of triploids (Supplementary Fig. 5B). Monosporic isolates displayed various mating-type combinations associated with complex mating phenotypes (Supplementary Fig. 5B and 5C, Table S3), including bisexual mating[42] and dioecy[43].

Finally, genome sequencing of 16 monosporic isolates obtained in the meiotic progeny of one natural pentaploid revealed consistent segregation of disomes and trisomes across all isolates (a single case of monosomic chromosome I was observed, Supplementary Fig. 5A). No sign of spore endoreplication was detected in the progeny of the pentaploid strain.

In conclusion, we found no evidence supporting stepwise ploidy increase beyond tetraploidy, suggesting that $4x$ might represent an upper limit for this process. This is consistent with the rarity of natural pentaploids, which encompass only about 0.2% of strains in the largest available collection[9], and with the absence of hexaploid or higher-ploidy isolates.

Stepwise polyploidization as the main route to natural polyploidy

The stepwise and saltational routes to polyploidization rely on distinct mechanisms, namely the *SEM* sequence and the WGD, respectively (Fig. 1), and are therefore expected to leave contrasting genomic signatures. To discriminate between these modes of origin, we analyzed the genomic constitution of the 91 natural polyploids from the 1,011 collection[7] and assessed whether their features are more consistent with a stepwise or a saltational process.

We first measured the DNA content of the 91 natural polyploids by flow cytometry and inferred ploidy without rounding to integer values. Rather than forming discrete clusters corresponding to strict $3x$, $4x$, or $5x$ ploidy levels, the data revealed an almost continuous ploidy range from $2.5x$ to $5.5x$ (Fig. 5A, Table S6), with clear discontinuities near $3.5x$ and $4.5x$ delimiting triploid, tetraploid, and pentaploid classes. Independent analysis of genome-wide *ABRs* at heterozygous sites yielded ploidy estimates largely concordant with DNA content, with only three exceptions (ASA, BEM, and CBR; Fig. 5A). This integrated reassessment led us to reclassify two previously reported tetraploids (AFT and CFS) as triploids and to resolve one

previously ambiguous strain (AEC) as a tetraploid, resulting in a final dataset

comprising 48 triploids, 38 tetraploids, and 5 pentaploids (Table S6).

Under a saltational polyploidization model, the prevalence of triploids would

require chromosome loss from tetraploids to occur more frequently than WGD from

diploids. Although rapid chromosome loss has been documented in laboratory

engineered homozygous tetraploids[30–32], it typically results in diploid rather than

triploid genomes[3]. Moreover, long term vegetative propagation of natural

tetraploids revealed no reduction in ploidy, indicating that polyploid levels are

stable in natural isolates[33]. Together, these observations argue against a dominant

saltational origin and instead support a stepwise route to polyploidization. This

conclusion is further reinforced by the strong enrichment of heterothallic strains

among natural polyploids[5], which is expected under the *SEM* model given that

endoreplication is restricted to heterothallic spores (Fig. 2C).

Additional genomic features are likewise inconsistent with a WGD-based origin. If

WGD is independent from zyogosity, a saltational model predicts that the fraction

of homozygous polyploids should be comparable to that observed among diploids

(47% in the 1,011 collection[7]). In contrast, we detected only two homozygous

triploids and no homozygous tetraploids among the 91 natural polyploids.

Furthermore, WGD of a heterozygous diploid genome would produce tetraploids

characterized by a dominant *ABR* peak near 0.5 (Supplementary Fig. 7), a signature

that is never observed in our data (Fig. 5B), providing strong evidence against WGD

as the primary mechanism of tetraploidization.

Finally, combining chromosome-specific sequencing coverage with *ABRs* revealed

extensive aneuploidy among natural polyploids, including 122 whole-chromosome

and 63 segmental aneuploidies (Table S6, Fig. 5C). Under the *SEM* model, such

aneuploidy is expected to accumulate with increasing ploidy, as tetraploids and

pentaploids originate from highly aneuploid spores produced by triploid meiosis.

In contrast, a saltational route predicts higher aneuploidy levels in triploids arising

from chromosome loss in tetraploids. Consistent with the stepwise model, we

observed a monotonic increase in the proportion of aneuploid strains with ploidy,

from 52% in triploids to 78% in tetraploids and 100% in pentaploids.

Together, these complementary genomic signatures indicate that higher ploidy
levels in *S. cerevisiae* arise predominantly through iterative cycles of the *SEM*
sequence rather than through saltational whole-genome duplication, establishing
stepwise polyploidization as the main evolutionary route to natural polyploidy.

Figure 5

Figure 5 | Genomic signatures of natural polyploids support a stepwise origin.
A, Ploidy estimates of 91 natural polyploid isolates inferred from flow cytometry, showing a near-
continuous distribution from ~2.5*x* to ~5.5*x* with discontinuities delineating triploid, tetraploid, and
pentaploid classes. **B**, Genome-wide allele balance ratio (*ABR*) distributions for the 91 natural
polyploids. Observed *ABR* patterns are inconsistent with whole-genome duplication of heterozygous
diploids and instead support stepwise ploidy increase. **C**, Chromosome copy number and *ABR*
profiles of a representative natural polyploid strain (AQT) presenting several whole-chromosome
and segmental aneuploidies.

Discussion

Here, we identify a natural route to polyploidization in yeast that we term stepwise polyploidization, mediated by iterative cycles of the *SEM* sequence. This route begins with sporulation, followed by post-meiotic endoreplication during spore germination. The sequence ends with mating between the endoreplicated spore and a compatible meiotic product within the ascus. This process contrasts sharply with the somatic WGD in diploids, which is often invoked to explain polyploid formation in yeast, as well as with the dominant mechanism described in plants, which relies on the fusion of unreduced gametes produced by meiotic restitution[22]. Notably, a conceptually related process has been proposed in plants, where a subset of 2*x* nuclei in mature pollen may arise from mitotic endoreplication after completion of meiosis rather than from meiotic restitution[44]. This conceptual parallel suggests that post-meiotic endoreplication may contribute to polyploid formation beyond fungi.

The *SEM* model provides a unified explanation for several genomic characteristics of natural yeast polyploids, including strong association with heterothallism, pervasive heterozygosity, frequent triploid state and extensive aneuploidy. In a recent work, we showed that natural polyploidy in yeast was strongly associated with repeated losses of self-fertility[5], yet the causal basis of this association remained unclear. Our results provide a mechanistic explanation: only heterothallic spores experience a prolonged single–mating-type state upon germination that permits endoreplication, whereas homothallic spores rapidly self-mate and therefore do not enter the *SEM* pathway. At the same time, ploidy increases through mating between genetically distinct meiotic products preserve heterozygosity and account for the observed allele-balance distributions, in contrast to duplication of a single diploid genome. Stepwise polyploidization further implies that triploids are not rare anomalies but expected and stable intermediates produced by a single iteration of the *SEM* sequence. Because meiosis in triploids is intrinsically unbalanced, viable progeny are highly aneuploid, providing a direct explanation for the extensive aneuploidy observed in tetraploids. Given the broad ploidy range of spores produced by triploid meiosis, pentaploids could theoretically be generated by mating between an endoreplicated ~3*x* spore and a near-diploid partner.

The endoreplication step itself emerges as the pivotal mechanism of stepwise
polyploidization. Although the precise molecular mechanisms underlying
spontaneous spore endoreplication remain to be elucidated, an early study
identified a mutation in *CDC31*, a gene required for spindle pole body duplication,
as a driver of endoreduplication[45]. To date, at least 16 genes have been implicated in
ploidy increase in *S. cerevisiae*[46]. These genes encode proteins that primarily regulate
spindle pole body duplication, chromosome segregation, cytokinesis and cell-cycle
control, thereby highlighting the central role of cell-division and genome-
partitioning pathways in the control of ploidy. Moreover, high rate of
endoreplication depends on the presence of a functional copy of *SSD1*, a gene that
also regulates aneuploidy tolerance in wild yeast[47,48]. Future work will be required
to assess the extent of natural variation in these genes across populations and to
dissect the molecular events governing genome doubling associated with spore
germination.
In conclusion, our findings demonstrate that stepwise polyploidization through
iterative cycles of the *SEM* sequence constitutes the predominant route to natural
polyploidy in *S. cerevisiae*, although alternative routes, such as those involving
mating-competent diploids arising from *MAT* hemi- or homozygotization, may also
contribute. Beyond its evolutionary significance, the *SEM* mechanism has practical
implications. Polyploid yeast strains are widely used in fermentation industries,
including brewing and baking, and are also frequently isolated from clinical
settings[4,7–11]. The identification of the natural mechanism underlying their formation,
together with our ability to experimentally recapitulate *SEM*-mediated
polyploidization under laboratory conditions without genetic modification, opens
new avenues for the rational generation of novel polyploid strains. We envision that
controlled and potentially automated implementation of the *SEM* sequence could
enable the production of large numbers of polyploids with unprecedented genetic
compositions, offering a powerful platform for strain improvement for diverse
applications in the food production and fermentation industries.

Methods

Strains

Natural parental diploid (Table S1) and polyploid (Table S5) strains analyzed in this study were obtained from the 1,011 *S. cerevisiae* collection[7] and are referred to using the standardized three-letter strain code. Artificial homozygous polyploids derived from the AKN (YPS128) background, as well as heterozygous polyploids derived from AKN and SACE_YBP (YJM1248) backgrounds (YPB012, YPB022, YPB021), were constructed as described previously[49]. Standard reference strains used for ploidy calibration were as previously described[50]. Tester strains used for mass mating assays were OS104a and OS104α (YPS128 background)[51].

Media composition and growth conditions

Standard growth medium was YPD, consisting of yeast extract (10 g L^{-1}), peptone (20 g L^{-1}), and dextrose (20 g L^{-1}), prepared with or without agar (20 g L^{-1}). Sporulation medium was potassium acetate (KAc) agar, containing potassium acetate (20 g L^{-1}) and agar (25 g L^{-1}). Selective synthetic complete media were prepared using yeast nitrogen base without ammonium sulfate (YNB; 1.7 g L^{-1}), arginine (2 g L^{-1}), glucose (20 g L^{-1}), and agar (20 g L^{-1}), supplemented with appropriate complete supplement mixtures (CSM). CSM + G418 medium contained standard CSM and was supplemented with Geneticin (G418; 200 μg mL^{-1}). CSM - Ura medium contained CSM lacking uracil (CSM - Ura). CSM - Ura + G418 medium was identical to CSM - Ura and additionally supplemented with G418 (200 μg mL^{-1}).

Unless otherwise stated, yeast cells were revived from glycerol stocks on YPD agar and grown at 30 °C until the formation of a visible patch prior to experimental assays. Qualitative sporulation and resporulation capacity were assessed by transferring cells to KAc agar and incubating them at 30 °C. After 7 days, cultures were examined by light microscopy for the presence of asci.

Flow cytometry

Cellular ploidy was estimated by propidium iodide staining followed by flow cytometry. Ploidy validations were performed as previously described[50], using cultures grown in YPD. Rapid ploidy assessments were conducted either from

overnight 1 mL YPD microcultures or by direct fixation of colonies in 70% ethanol.
Flow cytometry data were gated using FSC-A versus SSC-A and FSC-A versus FSC-
H to exclude debris and doublets, followed by FSC-A versus PI fluorescence
intensity. When required, cell aggregation was reduced by proteinase K treatment.
Ploidy values were inferred using MuPETFlow[50] (see Extended Methods).

Genotyping mating type loci

Mating genotype was determined by colony PCR as previously described[5]. In a
limited number of cases, PCR reactions were performed or repeated using *MAT*
locus primers described in[52] to confirm mating genotype assignments.

Mass mating assay

Query and tester cells were mixed on YPD agar and incubated at 30 °C for 24 h to
allow mating. Parental query cells were then streaked onto single-selection medium
to confirm marker compatibility, while cells from the mating patch were plated onto
double-selection medium to assess mating ability (see Media composition and
growth conditions).

Multi-test screening to detect diploid maters

All 742 query strains were initially tested using the mass mating assay with minor
modifications, and mating outcomes were scored semi-quantitatively, considering
as positive only the crosses that formed a confluent cellular layer (see Extended
Methods). A subset of 102 strains was further characterized by microscopic
examination of zygote formation after 4 h of co-incubation with tester strains, and
by assessment of creeping behavior as previously described[53] (see Extended
Methods). Results from the three assays were integrated into a single composite
mating score, and strains with a score greater than 0 were classified as potential
maters. Mating efficiency of candidate strains was then evaluated using a
quantitative mating assay adapted from[54]. Briefly, equal numbers of query and tester
cells (2.5×10^8 cells of each) were mixed and spotted onto a nitrocellulose filter
placed on YPD agar. After 4 h of co-incubation at 30 °C, cells were recovered from
the filter, serially diluted, and plated onto selective media to quantify colony-
forming units (CFUs).

Candidate diploid mater strains were subsequently subjected to supplementary
analyses to further characterize their mating behavior and genetic features. Mating
genotype was determined as described in the section Genotyping mating type loci,
and chromosome III aneuploidies and zygosity were retrieved from the original
sequencing data[7].

Asci dissection and spore isolation

Sporulated asci were treated with lyticase and dissected using a micromanipulator
(Singer Instruments MSM 400) as previously described[5]. Spores were plated on YPD
agar and cultured at 30 °C, and germination was monitored for up to 7 days.
Resulting colonies were picked and stored in glycerol stocks until further analyses.

Ascus micromanipulation and polyploid detection

Complete asci, irrespective of spore number, were randomly isolated and plated
onto YPD agar using a micromanipulator (Singer Instruments MSM 400) and
incubated at 30 °C. After colony formation (approximately 2 days), a rapid
preliminary ploidy assessment was performed as described in the section Flow
cytometry. Colonies suspected to be polyploid were subsequently single-colony
streaked, and ploidy was validated by flow cytometry.

Genomic DNA isolation

Genomic DNA was extracted either using the E-Z 96 Tissue DNA Kit (Omega Bio-
Tek), following the manufacturer's instructions, or using the SDS-Kac protocol as
previously described[55].

Genome sequencing and bioinformatic analysis

Genomic DNA was sequenced using Illumina paired-end sequencing (2 × 150 bp)
with an average insert size of approximately 300 bp. Read quality was assessed
using FastQC (v0.12.1)[56] and summarized with MultiQC (v1.13)[57]. Read trimming
and filtering were performed using Trimmomatic (v0.39)[58], with parallel execution
managed using GNU Parallel (v20190322)[59].

Read alignment and variant calling were performed using a custom Snakemake-
based pipeline (https://github.com/CintiaG/poly_CCN_and_VC.git). Sequencing
reads were aligned to the reference genome using BWA (v0.7.18)[60]. SAM files were
converted to BAM format and sorted using SAMtools (v1.21)[61]. Duplicate reads were

identified and marked using GATK MarkDuplicatesSpark (v4.5.0.0)[62]. Variants were

called per strain using GATK HaplotypeCaller in gVCF mode. Individual gVCF files

were merged using GATK CombineGVCFs, and joint genotyping was performed

with GATK GenotypeGVCFs to generate final VCF files.

Chromosome copy number (CCN) was estimated from sequencing depth using

BEDTools (v2.29.2)[63]. CCN and segmental aneuploidies were inferred using in-

house scripts (plot_depth.R and plot_copies.R; https://github.com/CintiaG/wgs-

aneuploidy-abr.git).

Allele balance ratio (*ABR*) was calculated by extracting read depth (DP) and allele

depth (AD) information from VCF files using BCFtools (v1.13)[64]. Text outputs were

converted to fst format using the fst package (v0.9.8, http://www.fstpackage.org).

ABR values were computed using the allele_balance_ratio.R script

(https://github.com/CintiaG/wgs-aneuploidy-abr.git), retaining variants with

ABR between 0.125 and 0.875. The number of heterozygous sites was estimated

considering only variants with up to three alleles. In selected cases, CCN estimates

were manually refined using *ABR* information, and genome-wide visualizations

were generated using the plot_allele_balance_ratio.R script

(https://github.com/CintiaG/wgs-aneuploidy-abr.git).

Data availability

The data for this study have been deposited in the European Nucleotide Archive

(ENA) at EMBL-EBI under accession number PRJEB108394.

References

1.	Morris, J. P., Baslan, T., Soltis, D. E., Soltis, P. S. & Fox, D. T. Integrating the Study of Polyploidy Across Organisms, Tissues, and Disease. *Annu. Rev. Genet.* **58**, 297–318 (2024).

2.	Fischer, G., Liti, G. & Llorente, B. The budding yeast life cycle: More complex than anticipated? *Yeast Chichester Engl.* **38**, 5–11 (2021).

3.	Selmecki, A. *et al.* Polyploidy can drive rapid adaptation in yeast. *Nature* **519**, 349–352 (2015).

4.	Albertin, W. & Marullo, P. Polyploidy in fungi: evolution after whole-genome duplication. *Proc. R. Soc. B Biol. Sci.* **279**, 2497–2509 (2012).

5.	Vittorelli, N. *et al.* Repeated losses of self-fertility shaped heterozygosity and polyploidy in yeast evolution. *Proc. Natl. Acad. Sci.* **123**, e2525679123 (2026).

6.	Ward, C. M. *et al.* Recombination, admixture and genome instability shape the genomic landscape of Saccharomyces cerevisiae derived from spontaneous grape ferments. *PLOS Genet.* **20**, e1011223 (2024).

7.	Peter, J. *et al.* Genome evolution across 1,011 Saccharomyces cerevisiae isolates. *Nature* **556**, 339 (2018).

8.	Duan, S.-F. *et al.* The origin and adaptive evolution of domesticated populations of yeast from Far East Asia. *Nat. Commun.* **9**, 2690 (2018).

9.	Loegler, V., Friedrich, A. & Schacherer, J. Overview of the Saccharomyces cerevisiae population structure through the lens of 3,034 genomes. *G3 Bethesda Md* **14**, jkae245 (2024).

10.	Ezov, T. K. *et al.* Molecular-Genetic Biodiversity in a Natural Population of the Yeast Saccharomyces cerevisiae From "Evolution Canyon": Microsatellite Polymorphism, Ploidy and Controversial Sexual Status. *Genetics* **174**, 1455–1468 (2006).

11.	Muller, L. A. H. & McCusker, J. H. Microsatellite analysis of genetic diversity among clinical and nonclinical Saccharomyces cerevisiae isolates suggests heterozygote advantage in clinical environments. *Mol. Ecol.* **18**, 2779–2786 (2009).

12.   Segraves, K. A. & Anneberg, T. J. Species interactions and plant polyploidy.
Am. J. Bot. **103**, 1326–1335 (2016).

13.   Wood, T. E. *et al.* The frequency of polyploid speciation in vascular plants.
Proc. Natl. Acad. Sci. **106**, 13875–13879 (2009).

14.   MABLE, B. K. 'Why polyploidy is rarer in animals than in plants': myths and
mechanisms. *Biol. J. Linn. Soc.* **82**, 453–466 (2004).

15.   Mable, B. K., Alexandrou, M. A. & Taylor, M. I. Genome duplication in
amphibians and fish: an extended synthesis. *J. Zool.* **284**, 151–182 (2011).

16.   David, K. T. Global gradients in the distribution of animal polyploids. *Proc.*
Natl. Acad. Sci. U. S. A. **119**, e2214070119 (2022).

17.   Mezzasalma, M., Brunelli, E., Odierna, G. & Guarino, F. M. Evolutionary and
Genomic Diversity of True Polyploidy in Tetrapods. *Anim. Open Access J. MDPI* **13**,
1033 (2023).

18.   Bielski, C. M. *et al.* Genome doubling shapes the evolution and prognosis of
advanced cancers. *Nat. Genet.* **50**, 1189–1195 (2018).

19.   Lambuta, R. A. *et al.* Whole-genome doubling drives oncogenic loss of
chromatin segregation. *Nature* **615**, 925–933 (2023).

20.   Martínez-Jiménez, F. *et al.* Pan-cancer whole-genome comparison of primary
and metastatic solid tumours. *Nature* **618**, 333–341 (2023).

21.   Otto, S. P. & Whitton, J. Polyploid incidence and evolution. *Annu. Rev. Genet.*
34, 401–437 (2000).

22.   Ramsey, J. & Schemske, D. W. PATHWAYS, MECHANISMS, AND RATES
OF POLYPLOID FORMATION IN FLOWERING PLANTS. *Annu. Rev. Ecol. Evol.*
Syst. **29**, 467–501 (1998).

23.   Pomper, S., Daniels, K. M. & McKee, D. W. Genetic Analysis of Polyploid
Yeast. *Genetics* **39**, 343–355 (1954).

24.   Gunge, N. & Nakatomi, Y. Genetic Mechanisms of Rare Matings of the Yeast
SACCHAROMYCES CEREVISIAE Heterozygous for Mating Type. *Genetics* **70**, 41–
58 (1972).

25.   Wolfe, K. H. & Shields, D. C. Molecular evidence for an ancient duplication
of the entire yeast genome. *Nature* **387**, 708–713 (1997).

26.   Marcet-Houben, M. & Gabaldón, T. Beyond the Whole-Genome Duplication:
Phylogenetic Evidence for an Ancient Interspecies Hybridization in the Baker's
Yeast Lineage. *PLOS Biol.* **13**, e1002220 (2015).

27.   Marsit, S., Hénault, M., Charron, G., Fijarczyk, A. & Landry, C. R. The neutral
rate of whole-genome duplication varies among yeast species and their hybrids. *Nat.*
Commun. **12**, 3126 (2021).

28.   Charron, G., Marsit, S., Hénault, M., Martin, H. & Landry, C. R. Spontaneous
whole-genome duplication restores fertility in interspecific hybrids. *Nat. Commun.*
10, 1–10 (2019).

29.   Tong, K. *et al.* Genome duplication in a long-term multicellularity evolution
experiment. *Nature* **639**, 691–699 (2025).

30.   Mayer, V. W. & Aguilera, A. High levels of chromosome instability in
polyploids of Saccharomyces cerevisiae. *Mutat. Res.* **231**, 177–186 (1990).

31.   Gerstein, A. C., Chun, H.-J. E., Grant, A. & Otto, S. P. Genomic convergence
toward diploidy in Saccharomyces cerevisiae. *PLoS Genet.* **2**, e145 (2006).

32.   Gerstein, A. C., McBride, R. M. & Otto, S. P. Ploidy reduction in
Saccharomyces cerevisiae. *Biol. Lett.* **4**, 91–94 (2008).

33.   Dutta, A., Dutreux, F. & Schacherer, J. Loss of Heterozygosity Spectrum
Depends on Ploidy Level in Natural Yeast Populations. *Mol. Biol. Evol.* **39**, msac214
(2022).

34.   Fox, D. T. & Duronio, R. J. Endoreplication and polyploidy: insights into
development and disease. *Development* **140**, 3–12 (2013).

35.   Edgar, B. A. & Orr-Weaver, T. L. Endoreplication Cell Cycles: More for Less.
Cell **105**, 297–306 (2001).

36.   Harari, Y., Ram, Y., Rappoport, N., Hadany, L. & Kupiec, M. Spontaneous
Changes in Ploidy Are Common in Yeast. *Curr. Biol. CB* **28**, 825-835.e4 (2018).

37.   De Chiara, M. *et al.* Domestication reprogrammed the budding yeast life
cycle. *Nat. Ecol. Evol.* **6**, 448–460 (2022).

38.   Charles, J. St., Hamilton, M. L. & Petes, T. D. Meiotic Chromosome
Segregation in Triploid Strains of Saccharomyces cerevisiae. *Genetics* **186**, 537–550
(2010).

39.   Mozzachiodi, S., Krogerus, K., Gibson, B., Nicolas, A. & Liti, G. Unlocking the

functional potential of polyploid yeasts. *Nat. Commun.* **13**, 2580 (2022).

40.   Barrere, J., Nanda, P. & Murray, A. W. Alternating selection for dispersal and

multicellularity favors regulated life cycles. *Curr. Biol. CB* **33**, 1809-1817.e3 (2023).

41.   Loidl, J. Meiotic chromosome pairing in triploid and tetraploid

Saccharomyces cerevisiae. *Genetics* **139**, 1511–1520 (1995).

42.   Haber, J. E. BISEXUAL MATING BEHAVIOR IN A DIPLOID OF

SACCHAROMYCES CEREVISIAE: EVIDENCE FOR GENETICALLY

CONTROLLED NON-RANDOM CHROMOSOME LOSS DURING VEGETATIVE

GROWTH. *Genetics* **78**, 843–858 (1974).

43.   Al Safadi, R., Weiss-Gayet, M., Briolay, J. & Aigle, M. A polyploid population

of Saccharomyces cerevisiae with separate sexes (dioecy). *FEMS Yeast Res.* **10**, 757–

768 (2010).

44.   Pan, G., Zhou, Y., Fowke, L. C. & Wang, H. An efficient method for flow

cytometric analysis of pollen and detection of 2n nuclei in Brassica napus pollen.

Plant Cell Rep. **23**, 196–202 (2004).

45.   Schild, D., Ananthaswamy, H. N. & Mortimer, R. K. An endomitotic effect of

a cell cycle mutation of Saccharomyces cerevisiae. *Genetics* **97**, 551–562 (1981).

46.   Andreychuk, Y., Zhuk, A., Tarakhovskaya, E., Inge-Vechtomov, S. &

Stepchenkova, E. Rate of spontaneous polyploidization in haploid yeast

Saccharomyces cerevisiae. *Biol. Commun.* **67**, 88–96 (2022).

47.   Tung, S., Bakerlee, C. W., Phillips, A. M., Nguyen Ba, A. N. & Desai, M. M.

The genetic basis of differential autodiploidization in evolving yeast populations.

G3 GenesGenomesGenetics **11**, jkab192 (2021).

48.   Hose, J. *et al.* The genetic basis of aneuploidy tolerance in wild yeast. *eLife* **9**,

e52063 (2020).

49.   Garrido, C. *et al.* Natural diversity of telomere length distributions across 100

Saccharomyces cerevisiae strains. *Genome Res.* gr.281132.125 (2026)

doi:10.1101/gr.281132.125.

50.   Gómez-Muñoz, C. & Fischer, G. MuPETFlow: multiple ploidy estimation tool

from flow cytometry data. *BMC Genomics* **26**, 301 (2025).

51. Cubillos, F. A., Louis, E. J. & Liti, G. Generation of a large set of genetically
tractable haploid and diploid Saccharomyces strains. *FEMS Yeast Res.* **9**, 1217–1225
(2009).

52. Huxley, C., Green, E. D. & Dunham, I. Rapid assessment of S. cerevisiae
mating type by PCR. *Trends Genet. TIG* **6**, 236 (1990).

53. Arras, S. D. M. *et al.* Creeping yeast: a simple, cheap and robust protocol for
the identification of mating type in Saccharomyces cerevisiae. *FEMS Yeast Res.* **22**,
foac017 (2022).

54. de Barros Lopes, M., Bellon, J. R., Shirley, N. J. & Ganter, P. F. Evidence for
multiple interspecific hybridization in *Saccharomyces* sensu stricto species. *FEMS*
Yeast Res. **1**, 323–331 (2002).

55. Agier, N. *et al.* A transient mutational burst occurs during yeast colony
development. *Mol. Syst. Biol.* **21**, 1214–1236 (2025).

56. Andrews, S. (2010) FastQC A Quality Control Tool for High Throughput
Sequence Data. - References - Scientific Research Publishing.
https://www.scirp.org/reference/referencespapers?referenceid=2781642.

57. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize
analysis results for multiple tools and samples in a single report. *Bioinforma. Oxf.*
Engl. **32**, 3047–3048 (2016).

58. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for
Illumina sequence data. *Bioinforma. Oxf. Engl.* **30**, 2114–2120 (2014).

59. GNU Parallel 2018. *Guide books*
https://dl.acm.org/doi/abs/10.5555/3235180 doi:10.5555/3235180.

60. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–
Wheeler transform. *Bioinformatics* **25**, 1754–1760 (2009).

61. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinforma.*
Oxf. Engl. **25**, 2078–2079 (2009).

62. Van der Auwera, G. A. *et al.* From FastQ data to high confidence variant calls:
the Genome Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinforma.* **43**,
11.10.1-11.10.33 (2013).

63.	Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for
comparing genomic features. *Bioinforma. Oxf. Engl.* **26**, 841–842 (2010).
64.	Danecek, P. & McCarthy, S. A. BCFtools/csq: haplotype-aware variant
consequences. *Bioinformatics* **33**, 2037–2039 (2017).
755