A High-fat, High-salt Diet Model of MDAKD Impairs Bioenergetic Efficiency for ATP Synthesis

Stephen T Decker, PhD[1], Za'rya T Smith[1,2], Precious C Opurum[1,3], Venisia L Paula[1,2,4,], Kennedy N Moses[1], Deborah Stuart[5], Anu S Kurian[1], Subhasmita Rout[1,3], Nirupama Ramkumar MD, MPH[5], Katsuhiko Funai, PhD[1,3]

[1]Center for Metabolic Health, University of Utah, Salt Lake City, Utah

[2]Haumana 'O Pasifika Program, University of Utah, Salt Lake City, Utah

[3]Department of Nutrition and Integrative Physiology, University of Utah, Salt Lake City, Utah

[4]Department of Kinesiology, University of Utah, Salt Lake City, Utah

[5]Division of Nephrology and Hypertension, University of Utah School of Medicine, Salt Lake City, Utah

Corresponding Author: Katsuhiko Funai, kfunai@utah.edu

15 N 2030 E

University of Utah

Salt Lake City, UT 84112

Keywords: Mitochondria, Metabolism, High-Salt, High-Fat, MDAKD

Abstract

Metabolic dysfunction-associated kidney disease (MDAKD) is closely linked to dietary

excess, but models that capture early kidney injury without obesity are limited. We fed male

C57BL/6J (6J) and C57BL/6N (6N) mice a high-fat, high-sodium (HF/HNa) or control diet for 16

weeks. HF/HNa feeding did not alter body weight, adiposity, or total food intake; however, it

increased dietary energy and sodium exposure, kidney mass, water intake, and urine volume.

GFR declined modestly in 6J mice, whereas 6N mice maintained or slightly increased GFR.

Both substrains showed increased urinary albumin, creatinine, KIM-1, and NGAL, while cystatin

C rose predominantly in 6N mice, indicating strain-dependent tubular injury. Whole-kidney

trichrome staining revealed increased fibrotic area with HF/HNa, particularly in 6N mice, without

significant changes in glomerular morphology. In isolated renal mitochondria, oxygen

consumption was preserved, but ATP production and ATP:O ratios were reduced, with

unchanged citrate synthase activity and OXPHOS protein abundance, consistent with early

mitochondrial bioenergetic uncoupling. Exploratory urinary proteomics in 6J mice identified

HF/HNa-associated changes in proteins linked to tubular stress and extracellular matrix

remodeling. These findings define an early MDAKD-like renal phenotype with strain-specific

functional responses, tubular injury, fibrosis, and impaired mitochondrial ATP efficiency.

Translational Statement

Metabolic Dysfunction-Associated Kidney Disease (MDAKD) is a leading driver of chronic

kidney disease (CKD) in the world. In addition to obesity and related comorbidities, renal

mitochondrial dysfunction is thought to be a key contributor to the development of CKD in

patients with MDAKD; however, few models recapitulate the progression of MDAKD. We couple

well-established mouse models of obesity, namely the C57Bl/6J and C57Bl/6N mouse lines,

with a high-fat, high-salt diet to induce renal mitochondrial dysfunction, leading to early stages of

MDAKD as indicated by widespread fibrosis and mild reduction in glomerular filtration rate,

though these effects were strain-dependent. We identify diet-induced mitochondrial dysfunction

as a common feature in both mouse strains, suggesting impairments in mitochondrial respiration

and oxidative ATP production are indeed a contributing factor to the development of MDAKD.

This study highlights the role of energetic impairments in the pathogenesis of MDAKD and may

guide future therapies for CKD.

Introduction

Chronic kidney disease (CKD) is a major global health burden, affecting nearly 10% of the population worldwide and accounting for more than 800 million individuals overall.[1,2] CKD is strongly associated with increased risk of cardiovascular disease and premature mortality, with the majority of patients dying from cardiovascular complications rather than progressing to kidney failure.[1,3,4] One of the dominant drivers of CKD in contemporary populations is the metabolic dysfunction-associated kidney disease (MDAKD), a pathophysiological state defined by the convergence of obesity, metabolic dysregulation, and renal dysfunction.[3-5] Patients with MDAKD exhibit accelerated loss of glomerular filtration rate (GFR), higher albuminuria, lipid deposition in the kidney, and a greater risk of progression to end-stage kidney disease compared to individuals with isolated kidney injury.[5,6]

Despite its prevalence and clinical importance, preclinical models that recapitulate the multifactorial nature of MDAKD-associated CKD remain lacking.[7] Traditional murine models of CKD rely on surgical interventions such as unilateral ureteral obstruction or 5/6 nephrectomy, which produce acute and severe injury but fail to mimic the gradual, systemic progression of human disease.[8] Genetic models replicate rare monogenic disorders but lack relevance to the common multifactorial pathogenesis of MDAKD.[9] Dietary manipulations such as high-fat feeding[10-17], which induce obesity and insulin resistance, and high-salt diets, which promote hypertension and tubular stress[10,12,18,19], have been widely used. However, in isolation, these approaches rarely lead to measurable declines in GFR or sustained CKD progression. Thus, no standardized diet-focused murine paradigm faithfully captures the integration of systemic metabolic load, tubular injury, and mitochondrial dysfunction that typifies human MDAKD-associated CKD.[3,8,20-22]

There is increasing recognition that impaired renal mitochondrial respiration and ATP synthesis are a central mediator of progressive CKD, particularly in the setting of increased tubular sodium reabsorption and metabolic overload.[16,21,23-29] The kidney is among the most

energy-demanding organs, with sodium reabsorption accounting for the majority of ATP

utilization.[22] Maladaptive increases in transport work, particularly in the proximal tubule and thick

ascending limb, can impose a metabolic burden that exceeds mitochondrial capacity, leading to

oxidative injury, fibrosis, and nephron loss.[13,16,21,22] However, few murine studies have

systematically linked dietary stressors with renal functional decline, structural injury, transporter

adaptations, and mitochondrial bioenergetics in vivo. Furthermore, although reports have

suggested that standard mouse models of obesity, such as the C57Bl/6J and C57Bl/6N strains,

are less prone to kidney disease,[30] the combination of a high-fat, high-salt diet in these strains

has not yet been utilized in a renal-specific context.

In this study, we tested whether long-term exposure (16 weeks) to a combined high-fat,

high-salt (HF/HNa) diet could reproduce key renal features of MDAKD. We evaluated systemic

and renal physiology, histopathology, injury marker expression, and mitochondrial respiration

using a substrate–uncoupler–inhibitor titration (SUIT) protocol.[31] We aimed to determine if

HF/HNa feeding recapitulates traits of human CKD, and whether this approach could serve as a

relevant model for future mechanistic and therapeutic research.

Methods

Animal Care & Husbandry

8-10-week-old male C57BL/6J mice (Strain No. 000664; n = 23) were purchased from

Jackson Laboratory, and C57BL/6NCrl (Strain code: 027; n = 28) mice were purchased from

Charles River. Figure 1A shows a diagram of the experimental approach. All mice were

maintained in our colony and randomly assigned to a standard chow diet or a 42% fat, ~8% salt

diet (HF/HNa; Envigo Custom TD.230162; Figure 1B) for 16 weeks. Mice were housed in

standard conditions, and all animals had access to *ad libitum* food and water throughout the

study duration. Body weights were measured weekly. End-of-study body composition was

determined using a Bruker Minispec NMR (Bruker, Billerica, MA). All animals were handled

according to approved University of Utah Animal Use and Care Committee (Protocol No. 1633).

Glomerular Filtration Rate (GFR)

Transcutaneous GFR was measured using the clearance of fluorescein isothiocyanate-

labeled sinistrin (FITC-sinistrin) with a commercially available device (NIC-Kidney, MediBeacon

GmbH, Mannheim, Germany), as previously described,[32] according to the manufacturer. Mice

were placed under non-lethal isoflurane-induced anesthesia. A patch of dorsal skin was shaved,

and the monitor was attached. Background fluorescence was measured for 10 min, after which

FITC-sinistrin (7.5 mg/100 g body wt) was administered retro-orbitally. FITC-sinistrin

fluorescence was monitored for the next 90 minutes. The kinetics of fluorescence

disappearance were used to determine GFR using the manufacturer's software, according to

their standard guidelines. We measured the GFR of all mice between 10:00 AM and 02:00 PM.

Urine Collection & Urinalysis

Mice were placed in an MMC100 Metabolic Cage, and urine was collected every 24 h for

48 h at the end of the study. Body weight, food, and water intake were all observed throughout

the experiment. Mice were housed with their respective chow or HF/HNa diet converted into a

gel with free access to water. Urine samples were centrifuged at 21,000 g for 15 min, then

aliquoted and stored at -80°C. Urine albumin and creatinine were assessed using an

absorbance-based Albumin Creatinine Ratio Assay Kit (ab241018; Abcam Ltd., Cambridge,

UK). Urine Cystatin C (CysC; kit KE10066), lipocalin 2 (Lcn2, or NGAL; kit KE10045), and

kidney injury marker-1 (KIM-1, or HAVCR1; kit KE10064) were assessed using commercially

available kits from Proteintech (Rosemont, IL, USA). All assays were performed according to the

manufacturer's recommendations.

Euthanasia, Organ Harvesting, and Blood Plasma Collection and Analysis

After 16 weeks of the dietary intervention, mice were fasted for 2 hours (beginning at

7:00-8:00 AM to minimize feeding artifacts and variability in metabolic activity caused by food

intake and circadian rhythm) before blood was collected, and animals were euthanized using a

lethal mix of ketamine and xylazine. Blood was collected in EDTA-coated blood collection tubes

and centrifuged at 1,000 x g for 10 minutes. The separated plasma was then aliquoted and

stored at –80°C for later analysis. Blood urea nitrogen (BUN) was analyzed using the

QuantiChrom^{TM} Urea Assay Kit (DIUR-100; BioAssay Systems, Hayward, CA, USA) according

to the manufacturer's protocol.

Following euthanasia, the kidneys were then harvested and weighed. At random, one

kidney was finely cut into thirds, snap-frozen in liquid nitrogen, and stored for later western

blotting and qPCR. The other kidney was finely sliced for histological staining and mitochondrial

function assays.

Histology

Kidney samples were fixed in 10% formaldehyde for 24–48 h and transferred to the

ARUP Research Histology Core Laboratory at the University of Utah. Kidney slices were fixed in

paraffin, sectioned at 4 µm, and stained with H&E, PAS, and Masson's trichrome to detect

morphological changes and fibrosis. Trichrome-stained kidney sections were imaged using

Zeiss AxioScan.Z1 (ZEISS, Oberkochen, Germany) slide scanner at the University of Utah

Imaging core.

Histology Analysis

Trichrome-stained kidney sections were analyzed in Fiji software[33] (version 2.7.0) using

a combination of custom macros, the Labkit plugin[34] (version 0.4.0), and a custom-made Python

script to segment images using the MONAI framework[35-37] in a Google Colab environment, and

are described in detail in the supplementary methods. Briefly, glomerular and tubular structures

were identified to generate mask image files for all slides. The generated masks were used to

crop individual glomeruli and tubular structures for analysis. Features were analyzed following

postprocessing feature cutoffs and the colour deconvolution2[38-40] (version 2.1) plugin

incorporated into a custom macro in Fiji.

Kidney Mitochondrial Isolation

Mitochondria isolation was performed as previously described.[41] Briefly, kidney sections

were minced in ice-cold mitochondrial isolation medium (MIM) buffer (300 mM sucrose, 10 mM

HEPES, 1 mM EGTA; pH 7.4) with 1 mg/mL bovine serum albumin (BSA)and carefully

homogenized with a Teflon-glass pestle. The homogenate was centrifuged at 800 g for 10 min at

4°C. The resulting supernatant was transferred to another tube and centrifuged at 1,300 g for 10

min. The resulting supernatant was transferred to another tube and centrifuged at 10,000 g for

10 min at 4°C. The pellet was then resuspended in 100-300 µL of MIM buffer. Protein

concentrations were assessed with Pierce BCA Protein Assay Kits (No. 23225, Thermo Fisher

Scientific, Waltham, MA, USA).

High-Resolution Respirometry and Fluorometry

Mitochondrial O_2 consumption (JO_2) was evaluated using the Oroboros Oxygraph O2K

(Oroboros Instruments, Innsbruck, Austria), as previously described.[41] Isolated mitochondria

were introduced to the oxygraphy chambers containing 2 mL of Buffer Z [105 mM MES

potassium salt, 30 mM potassium chloride (KCl), 10 mM monopotassium phosphate (KH2PO4),

5 mM magnesium chloride (MgCl2), and 0.5 mg/mL BSA]. Respiration was assessed by adding

the following substrates: 0.5 mM malate (M), 5 mM pyruvate (P), 5 mM glutamate (G), 2 mM

ADP, 10 mM succinate (S), and 1.5 µM carbonyl cyanide-p-trifluoromethoxyphenylhydrazone

(FCCP).

ATP production ($JATP$) was fluorometrically quantified with a Horiba Fluoromax 4

(Horiba Scientific, Piscataway, NJ, USA) by enzymatically linking it to NADPH synthesis, as

previously described.[41] ATP generation was assessed with 0.5 mM malate, 5 mM pyruvate, 5

mM glutamate, 10 mM succinate, and successive 2, 20, and 200 µM ADP additions.

Western Blot Analyses

Whole kidney tissue samples were homogenized in cold RIPA lysis buffer (No. 89901,

Thermo Fisher Scientific) containing protease inhibitor (No. 78446, Thermo Fisher Scientific),

incubated at 4°C for 10 min, and centrifuged at 12,000 g for 15 min at 4°C, and the supernatant

was collected into a fresh tube. The BCA Protein Assay Kit (No. 23225, Thermo Fisher

Scientific) was used to measure the protein concentration in the supernatant. Western blotting

was carried out as previously described[41] with slight modifications. In summary, for

electrophoresis, appropriate volumes of protein were combined with 4✕ Laemmli sample buffer

supplemented with 10% β-mercaptoethanol (#No. 1610710, Bio-Rad, Hercules, CA) and loaded

onto a Bio-Rad 4%–20% gradient SDS-polyacrylamide gel. After that, the proteins were

transferred from the gel onto membranes made of polyvinylidene fluoride. The membranes were

blocked for 1 h at room temperature using 5% low-fat skim milk in Tris-buffered saline

containing 0.1% Tween 20 (TBS-T), and then, the kidney samples were incubated overnight

with specific primary antibodies (Table 1) to probe for protein abundance. Following multiple

TBS-T washes, the membranes were incubated for 1 h at room temperature with appropriate

secondary antibodies diluted 1:5,000 in 5% BSA. After multiple washes in TBS-T and one final

rinse in TBS. Membranes were exposed with Western Lightning Plus-ECL (PerkinElmer) and

imaged using a ChemiDoc Imaging System (Bio-Rad, Hercules, CA). The images were

quantified using Image Lab software (Bio-Rad).

Real-Time Quantitative Polymerase Chain Reaction (qPCR)

For qPCR investigations, whole mouse kidney tissues were lysed in 1 mL of TRIzol

(Thermo Fisher Scientific), and RNA was extracted using conventional methods. The iScript

cDNA Synthesis Kit (Bio-Rad, Hercules, CA) was used to reverse-transcribe total RNA. To

assess the expression of kidney damage biomarkers (Table 1). Prevalidated primer sequences

were taken from either the primer bank or previously published papers. All mRNA levels were

standardized to RPL32 with SYBR Green reagents (Thermo Fisher Scientific), and 384-well

plates were loaded and sent to the University of Utah Genomics Core laboratories for

processing.

Mass Spectrometry-based Proteomics

Mass spectrometry–based proteomics was performed as described in detail in the

supplementary methods. Briefly, urinary proteins (~200 µL) were acetone-precipitated,

processed using S-TRAP columns with reduction, alkylation, and trypsin digestion, and peptides

were eluted, desalted, and resuspended in formic acid; a pooled QC sample and a fractionated

pooled library were generated. Peptides (200 ng) were analyzed by nano–LC-MS/MS on a

nanoElute 2 coupled to a Bruker timsTOF Pro2 using a 45-min reversed-phase gradient, with

PASEF-DDA for library generation and dia-PASEF for sample analysis. A custom spectral library

was built in FragPipe v22.0 against the UniProt mouse reference proteome, and protein

quantification was performed in DIA-NN v2.0 with standard variable modifications, one missed

cleavage allowed, and a 1% false discovery rate at the peptide and protein levels.

Statistical Analysis

All statistical analyses were performed in R (version 4.5.1). Data are reported as mean ±

SEM unless otherwise noted. Group comparisons were analyzed using n-way ANOVA with diet

(chow vs. HF/HNa) and strain (C57Bl/6J vs. C57Bl/6N) as fixed factors. Normality and variance

assumptions were evaluated, and nonparametric aligned rank transform[42,43] (ART) procedures

were applied when appropriate. Post hoc multiple comparisons were performed using the

Holm–Šidák method. For histological segmentation and glomerular morphometry, linear mixed-

effects models (LMM) were employed with individual mice as a random factor to account for

nested sampling. For urine proteomics analysis, data were processed as described above, and

significance was corrected for the False Discovery Rate (FDR). Analyses were conducted using

the R packages *tidyverse, ggprism, ggbeeswarm, rstatix, ARTool, ggpubr, multcomp, reshape2,*

ggcompare, mutoss, emmeans, and *svglite*. Statistical significance was defined as $p < 0.05$.

Results

HF/HNa feeding alters fluid balance and intake without changing body weight or adiposity

Over the 16-week intervention (Figures 1A & 1B), both C57Bl/6J (chow n = 13, HF/HNa n = 10) and C57Bl/6N mice (chow n = 15, HF/HNa n = 13) gained weight progressively, but endpoint body weight and liver mass did not differ by diet or strain (Figures 1C & 1D). However, kidney mass was increased in the HF/HNa diet for both strains (Figure 1E), indicating that the combined high-fat, high-salt diet did not induce obesity or gross changes in adiposity compared with chow controls (likely due to no differences in food overall intake, shown in Supplementary Figure S1), but did induce renal hypertrophy. As expected, mice on the HF/HNa diet had a greater energy, fat, and salt intake compared to the chow groups (all $p < 0.001$; Figures 1F-H), consistent with a sustained shift in osmotic balance. In parallel, diet increased water intake ($p < 0.01$; Figure 1I) and urine output ($p < 0.001$; Figure 1J). Body composition was not influenced by diet in either strain (Supplementary Figure S1).

Renal Function

GFR, assessed by transcutaneous FITC–sinistrin clearance, differed by diet and substrain (Figures 2A–B). In C57Bl/6J mice, HF/HNa feeding significantly reduced GFR compared with chow controls, whereas in C57Bl/6N mice, HF/HNa-fed animals showed a small, non-significant increase in GFR relative to chow.

HF/HNa feeding increased urinary albumin and creatinine excretion and decreased plasma BUN in both substrains (Figures 2C–E). Urinary albumin–creatinine ratio (ACR) showed a main effect of diet and was significantly higher in HF/HNa-fed C57Bl/6J mice, with more variable responses in C57Bl/6N mice (Figure 2F). BUN:creatinine ratios did not differ significantly between diets in either substrain (Figure 2G). Urinary cystatin C excretion was significantly higher in HF/HNa-fed C57Bl/6N mice but not in C57Bl/6J mice (Figure 2H). Urinary

244   KIM-1 and LCN2 concentrations were also higher in HF/HNa-fed animals, with a significant

245   main effect of diet (p = 0.005 and p = 0.001, respectively; Figures 2I–J).

247   *Histopathological Analysis*

248   Structural analysis of glomeruli revealed significant remodeling with HF/HNa feeding.

Representative Masson's trichrome staining showed increased collagen deposition in HF/HNa

kidneys compared with chow (Figure 3A; representative images of H&E and PAS stains are

shown in Supplemental Figure 3). Whole-slide kidney fibrosis stain area was increased in the

HF/HNa-fed animals (main effect p < 0.001; Figure 3B). However, 6N showed a much greater

increase in fibrotic area relative to their chow counterparts (Chow vs HFHNa: p < 0.001) than

the 6J mice (Chow vs HFHNa: p = ns).

255   A custom Python pipeline was used to segment (representative image shown in

Supplementary Figure 3A) and generate binary masks of individual glomeruli, which were

subsequently filtered by size and circularity to exclude artifacts. A schematic of the glomerular

compartments, highlighting the features subjected to quantification, is shown in Figure 3C.

Quantitative analysis following color deconvolution revealed multiple abnormalities in glomerular

structure. A HF/HNa diet did not change total corpuscle area (Figure 3D), nor did it change tuft

or Bowman's Space areas (Figures 3E, 3F, 3H, and 3I), but it did significantly decrease

glomerular fibrosis area (p < 0.05; Figure 3G and 3J). We did not observe any significant

changes in the fibrotic areas of the tubules or in the basement membrane, nor did we see

substantial changes in Col1α1 or Col3α1 mRNA (Supplementary Figure 3). Taken together with

the functional data, these findings suggest that the HF/HNa-associated decline in GFR

observed in 6J mice likely reflects widespread tubular impairments and/or altered

hemodynamics rather than established glomerulosclerosis or tubulointerstitial fibrosis at this

stage.

Transporter and Inflammatory Protein Expression

Western blot analysis revealed evidence of strain-dependent responses of renal

transporters and podocyte proteins to HF/HNa feeding (representative blots shown in Figure

4A). In 6N mice, AQP2 abundance was increased (p = 0.051; Figure 4B), whereas 6J mice

showed no change compared with chow controls. NKCC2 abundance was unchanged in both

strains (Figure 4C). However, SGLT2 abundance trended toward a decrease in 6N mice on

HF/HNa (p = 0.062; Figure 4D), but not in 6J mice, suggesting adaptation to sodium

reabsorptive stress is a feature unique to 6N mice. Further, the mechanism underlying these

adaptations is not the result of changes in ER stress, as the ER stress markers BiP, CHOP, and

PERK were unchanged (Figures 4E-4G). Podocin abundance was unchanged in both strains

(Supplemental Figure 4B), suggesting minimal podocyte injury in these models. Likewise,

Na/K$^+$-ATPase and TNF-α abundance were also unaffected by diet (Supplemental Figure 4C

and 4D).

At the mRNA level, 6J, but not 6N, mice fed a HF/HNa diet exhibited significantly

increased mRNA abundance of KIM-1 (p < 0.001; Figure 4H). Surprisingly, diet did not induce

any changes in Lcn2, NFκB, or any canonical markers of inflammation (Figures 4I-M). Together,

these findings suggest that 6J mice experience a greater burden of osmotic imbalance under

HF/HNa feeding, which results in tubular injury that is not dependent on ER stress or

inflammatory pathways. Alternatively, the adaptive responses in 6N kidneys may underlie their

preserved or slightly elevated GFR despite similar exposure to HF/HNa feeding.

Mitochondrial Respiration and ATP Production

Kidney cells (especially proximal tubules) are highly energetic and rely heavily on

mitochondrial oxidative phosphorylation to maintain the reabsorptive load placed on the kidneys,

and thus, global osmotic balance. High-resolution respirometry of isolated kidney mitochondria

showed diet-linked differences in mitochondrial function (Figure 5A–D). While the 6N mice

exhibited constantly higher JO_2 compared to 6J mice, HF/HNa feeding did not alter

mitochondrial respiration under any of the substrate conditions. However, the HF/HNa diet did

significantly impair JATP and, thus, lowered the ATP:O ratio, indicating an uncoupling effect and

possible bioenergetic impairment of HF/HNa diet on kidney mitochondria. Furthermore, it is

unlikely that these changes were due to adaptations in mitochondrial content or ETC protein

abundance, as the abundance of citrate synthase (Figures 5E & 5F) was not different in any of

the groups. Furthermore, PGC1-a mRNA abundance (Supplementary Figure 5A), and ETC

complexes I, II, and IV, and ATP synthase protein abundance (Supplementary Figures 5B-E)

were not significantly different between strains or diets.

Urine Proteomics

Last, preliminary analysis of urine proteins in 6J mice (chosen due to the diet-induced

decline in GFR) revealed several urinary markers of interest that were increased with HF/HNa

feeding (Figure 6A). Consistent with our urinalysis markers, albumin, cystatin C, and Lcn2 levels

were increased in the urine of HF/HNa-fed mice, but the increase was not statistically

significant, likely due to sample size limitations (Figure 6B). Interestingly, other kidney injury

markers, such as Villin-1 (a structural protein found in tubular microvilli), were also increased in

urine. Markers of proteostasis, such as the subunit c of the V0 domain of vacuolar H^+-ATPase

(Atp6v0c), sinal peptide peptidase-2A (Sppl2a), and programmed cell death 6 (Pdcd6), were

increased in HF/HNa-fed mice urine. Likewise, several transport proteins, such as succinate

dehydrogenase A (Sdha), aldolase B (Aldob), and the organic ion transporter (Slc22a19) were

also increased in the urine. Collectively, these support several novel markers of interest to track

the severity of renal failure in mice.

Discussion

In this study (summarised in Figure 6C), we demonstrate that 16 weeks of HF/HNa feeding in C57Bl/6 mice recapitulates several key features of MDAKDassociated CKD. Despite no effect on body weight or adiposity, HF/HNa-fed animals exhibited altered feeding behavior, increased water intake, and higher urinary output, consistent with prior reports that sodium-rich diets impose systemic metabolic stress independent of obesity.[12,18,19] Functionally, 6J mice developed a mild but significant reduction in GFR, while 6N mice maintained or slightly increased GFR, reflecting strain-dependent susceptibility to renal injury. Both strains, however, displayed biochemical evidence of tubular stress with elevated U-cystatin C, U-KIM-1, and U-Lcn2, aligning with human CKD, where tubular injury markers often rise before overt albuminuria.[6,8,14,17,20] At the molecular level, transporter expression patterns diverged between strains: 6N mice exhibited trends toward adaptive regulation of AQP2 and SGLT2, while 6J mice showed no changes in protein abundance or fibrotic gene expression, consistent with prior evidence that sodium transport and renal stress differ by genetic background.[30] Finally, mitochondrial analyses revealed that HF/HNa feeding impaired ATP production and coupling efficiency without altering respiratory chain protein abundance, indicating functional bioenergetic uncoupling, in line with accumulating evidence that mitochondrial dysfunction is a central mediator of CKD progression.[16,21,23,25,28,41] Further, we did not find any evidence of activation of pro-inflammatory or ER stress pathways, suggesting that mitochondrial bioenergetic deficiency is a significant contributor to early progression of renal dysfunction, regardless of genetic background. Together, these results establish HF/HNa feeding as a multifactorial model of CKD and highlight strain-specific adaptations that may underlie resistance versus susceptibility to diet-induced kidney dysfunction.

Genetic Background Influences Renal Pathological Progression

A key novel finding of this study is the divergence between C57Bl/6J and C57Bl/6N mice

in their renal response to HF/HNa feeding. While 6J mice exhibited a modest but significant

decline in GFR and structural remodeling of glomeruli, 6N mice maintained or even slightly

increased GFR despite equivalent dietary exposure. This contrast underscores the profound

influence of genetic background on CKD susceptibility, consistent with prior work showing that

allelic differences between C57 substrains shape cardiovascular, renal, and metabolic

phenotypes.[30] In all mice, lack of diet-induced glomerular morphological changes suggests that

other factors, such as early hemodynamic impairment, rather than glomerulosclerosis, drive the

initial loss of filtration capacity. Such a dissociation between functional decline and fibrotic

remodeling closely mirrors early-stage human CKD, where reductions in GFR often precede

substantial structural injury and where fibrosis becomes detectable only at later stages of

progression.[1,3,13]

The 6N substrain, in contrast, exhibited relative preservation of filtration despite

biochemical evidence of tubular stress, highlighting potential protective adaptations at the

glomerular and vascular level. This observation is particularly important given that murine

models often fail to develop measurable declines in GFR in response to dietary stress alone,

limiting their translational relevance.[7,10,12-14,16,19,20,41] Our results therefore establish the 6J strain

as uniquely susceptible to hemodynamic perturbation under combined high-fat and high-salt

feeding, while the 6N strain provides a valuable comparator in which protective adaptations can

be studied. One possibility is that 6N mice exhibit more favorable autoregulatory control of renal

blood flow, buffering glomeruli against hypertensive and osmotic stress. Another is that

differences in sodium transporter regulation reduce the metabolic burden imposed on

glomerular hemodynamics in 6N mice. By contrasting these two substrains, our study highlights

the utility of genetic background as a tool to dissect mechanisms of susceptibility versus

resilience, paralleling the heterogeneity observed in human CKD populations where some

individuals remain resistant to renal decline despite comparable metabolic or hypertensive

stressors.

Tubular Stress is a Common Feature of HF/HNa Diet Feeding Regardless of Strain

Another important novel aspect of our study is the demonstration that tubular stress and

transporter regulation diverge between strains, potentially explaining the preserved filtration

capacity in 6N mice. Both substrains exhibited biochemical evidence of tubular injury, as shown

by elevations in urinary cystatin C, KIM-1, and NGAL. These markers are increasingly

recognized as early indicators of CKD progression in humans, often rising before overt

albuminuria or reductions in eGFR are detectable.[6,13,44] However, the molecular responses at

the level of tubular transporters were strikingly strain-dependent. In 6N mice, AQP2 expression

trended upward while SGLT2 trended downward, changes that would be expected to reduce

osmotic stress by limiting glucose reabsorption while maintaining water handling. By contrast,

6J mice did not exhibit these adaptations.

These findings support a model in which protective adaptations in sodium and glucose

transport in 6N mice reduce the energy burden of reabsorption and buffer glomeruli against

hemodynamic overload. The absence of significant changes in ER stress markers (BiP, CHOP,

PERK) and fibrotic and inflammatory transcripts (Col1a1, Col3a1, NF-κB, IL-6, CCL2, SGK1) in

both strains suggests that these adaptations occur independently of canonical unfolded protein

responses or pro-inflammatory pathways. Taken together, these results align with the

hypothesis that tubular dysfunction and transport burden are central drivers of CKD progression

and suggest that the balance between adaptive and maladaptive transporter regulation

determines whether GFR is preserved or declines under chronic dietary stress.

Mitochondrial Energetics as a Driver of Renal Stress

A central novel finding of this study is that HF/HNa feeding impaired renal mitochondrial

energetics in both C57Bl/6 substrains, providing a unifying mechanism linking tubular transport

burden to kidney injury. High-resolution respirometry demonstrated that while oxygen

consumption rates (JO$_2$) were not significantly altered by diet, ATP production (JATP) and the

ATP:O ratio were reduced, consistent with bioenergetic uncoupling. This functional impairment

occurred without significant changes in mitochondrial content, as assessed by citrate synthase

abundance, or in respiratory chain complex protein levels, indicating that the observed deficits

reflect bioenergetic dysfunction rather than reduced mitochondrial mass. Such uncoupling has

been described as an early event in CKD pathogenesis, where increased sodium transport

demand drives ATP turnover beyond the capacity of mitochondria to maintain efficient oxidative

phosphorylation [16,21,23-28,41,45]

The implications of these findings are twofold. First, they suggest that mitochondrial

inefficiency may be a primary causative mechanism of HF/HNa-induced renal injury,

independent of overt ER stress, fibrosis, or inflammation. Even modest reductions in ATP yield

per oxygen consumed can compromise tubular reabsorptive function and contribute to the

release of tubular injury markers, all of which were observed in this study. Second, the similar

bioenergetic impairment in both strains suggests that mitochondrial uncoupling represents a

shared pathogenic mechanism, while strain-specific differences in transporter regulation and

glomerular hemodynamics determine whether GFR is preserved (6N) or declines (6J). These

results align with recent reviews proposing mitochondrial dysfunction as a central mechanism of

CKD progression[16,21,23,25] and highlight mitochondrial energetics as a therapeutic target in diet-

induced kidney disease.

Conclusion & Future Directions

In conclusion, 16 weeks of HF/HNa feeding in C57Bl/6 mice reproduces several key

features of MDAKD-associated CKD, including tubular injury, glomerular remodeling, and

mitochondrial uncoupling, while revealing substrain-dependent susceptibility to filtration loss.

These findings establish HF/HNa feeding as a tractable, non-surgical model of early diet-

induced kidney disease and highlight the central role of mitochondrial energetics in driving renal

stress. Future studies should expand on these findings by applying complementary approaches

to dissect bioenergetic regulation at higher resolution,[46,47] including measurements of

mitochondrial membrane potential, reactive oxygen species production, and respiratory control

in isolated nephron segments.[24,48] In addition, comprehensive lipidomic profiling of mitochondrial

phospholipids, particularly cardiolipin and phosphatidylethanolamine species, will provide critical

insights into how changes in the mitochondrial membrane environment contribute to uncoupling

and impaired ATP synthesis.[49-53] Such studies will clarify the mechanistic links between tubular

transport burden, mitochondrial dysfunction, and nephron loss, and may identify new

therapeutic targets to preserve bioenergetic efficiency in CKD.

Disclosures

The authors have no conflicts of interest to disclose at this time.

Data Sharing

Data are available upon request. Scripts, training images, and output data are publicly available and accessible on GitHub at https://github.com/stdecker.

Acknowledgments

We gratefully acknowledge the support of the University of Utah Health Sciences Core Facilities. Specifically, we thank the Cell Imaging Core Facility for support with microscopy, ARUP histological services, and the Genomics Core Facility for quantitative PCR services. Their expertise and resources were instrumental in the completion of this study. Schematics in this manuscript were generated using BioRender.

Funding

This work was supported by the following grants from the National Institutes of Health: NIDDK R01-DK107397, NIDDK R01-DK127979, NIGMS R01-GM144613, and NIA R01-AG074535 to Katsuhiko Funai; NIDDK R01-DK133271 to Nirupama Ramkumar; and Ruth L. Kirschstein National Research Service Award 5T32DK091317 to Stephen Decker. Additional support for S.D. was provided by the National Kidney Foundation of Utah and Idaho.

Author Contributions

Supplementary material: Supplementary material is available online at www.kidney-

international.org.

References

1. Faucon AL. CKD Beyond the Clinic: A Holistic Projection of its Global Burden.
Kidney Int Rep. Aug 2025;10(8):2524-2526. doi:10.1016/j.ekir.2025.06.031

2. Hu J, Ke R, Teixeira W, et al. Global, Regional, and National Burden of CKD due
to Glomerulonephritis from 1990 to 2019: A Systematic Analysis from the Global Burden
of Disease Study 2019. *Clin J Am Soc Nephrol*. Jan 1 2023;18(1):60-71.
doi:10.2215/CJN.0000000000000017

3. Hall RK, Sloane R, Pieper C, et al. Competing Risks of Fracture and Death in
Older Adults with Chronic Kidney Disease. *J Am Geriatr Soc*. Mar 2018;66(3):532-538.
doi:10.1111/jgs.15256

4. Go AS, Chertow GM, Fan D, McCulloch CE, Hsu CY. Chronic kidney disease and
the risks of death, cardiovascular events, and hospitalization. *N Engl J Med*. Sep 23
2004;351(13):1296-305. doi:10.1056/NEJMoa041031

5. Bansal A, Chonchol M. Metabolic dysfunction-associated kidney disease:
pathogenesis and clinical manifestations. *Kidney Int*. Aug 2025;108(2):194-200.
doi:10.1016/j.kint.2025.01.044

6. Uduman J. Epidemiology of Cardiorenal Syndrome. *Adv Chronic Kidney Dis*. Sep
2018;25(5):391-399. doi:10.1053/j.ackd.2018.08.009

7. Liang J, Liu Y. Animal Models of Kidney Disease: Challenges and Perspectives.
Kidney360. Oct 1 2023;4(10):1479-1493. doi:10.34067/KID.0000000000000227

8. Eddy AA, Neilson EG. Chronic kidney disease progression. *J Am Soc Nephrol*.
Nov 2006;17(11):2964-6. doi:10.1681/ASN.2006070704

9.   Molinari E, Sayer JA. Disease Modeling To Understand the Pathomechanisms of

Human Genetic Kidney Disorders. *Clin J Am Soc Nephrol*. Jun 8 2020;15(6):855-872.

doi:10.2215/CJN.08890719

10.   Carvalho L, Shimari M, Boeder AM, et al. A novel model of cardiovascular-

kidney-metabolic syndrome combining unilateral nephrectomy and high-salt-sugar-fat

diet in mice. *Lab Anim (NY)*. Nov 2024;53(11):336-346. doi:10.1038/s41684-024-01457-

5

11.   Opurum PC, Decker ST, Stuart D, et al. Six months of physical inactivity is

insufficient to cause chronic kidney disease in C57BL/6J mice. *bioRxiv*. Aug 30

2024;doi:10.1101/2024.08.29.610415

12.   Costa MV, Fernandes-Santos C, Faria Tda S, Aguila MB, Mandarim-de-Lacerda

CA. Diets rich in saturated fat and/or salt differentially modulate atrial natriuretic peptide

and renin expression in C57BL/6 mice. *Eur J Nutr*. Feb 2012;51(1):89-96.

doi:10.1007/s00394-011-0196-1

13.   Hall JE, do Carmo JM, da Silva AA, Wang Z, Hall ME. Obesity, kidney

dysfunction and hypertension: mechanistic links. *Nat Rev Nephrol*. Jun 2019;15(6):367-

385. doi:10.1038/s41581-019-0145-4

14.   Deji N, Kume S, Araki S, et al. Structural and functional changes in the kidneys of

high-fat diet-induced obese mice. *Am J Physiol Renal Physiol*. Jan 2009;296(1):F118-

26. doi:10.1152/ajprenal.00110.2008

15.   Jiang T, Wang Z, Proctor G, et al. Diet-induced obesity in C57BL/6J mice causes

increased renal lipid accumulation and glomerulosclerosis via a sterol regulatory

element-binding protein-1c-dependent pathway. *J Biol Chem.* Sep 16
2005;280(37):32317-25. doi:10.1074/jbc.M500801200

16.   Sun Y, Ge X, Li X, et al. High-fat diet promotes renal injury by inducing oxidative
stress and mitochondrial dysfunction. *Cell Death Dis.* Oct 24 2020;11(10):914.
doi:10.1038/s41419-020-03122-4

17.   Dissard R, Klein J, Caubet C, et al. Long term metabolic syndrome induced by a
high fat high fructose diet leads to minimal renal injury in C57BL/6 mice. *PLoS One.*
2013;8(10):e76703. doi:10.1371/journal.pone.0076703

18.   Bernhardt A, Krause A, Reichardt C, et al. Excessive sodium chloride ingestion
promotes inflammation and kidney fibrosis in aging mice. *Am J Physiol Cell Physiol.* Aug
1 2023;325(2):C456-C470. doi:10.1152/ajpcell.00230.2023

19.   Escano CS, Armando I, Wang X, et al. Renal dopaminergic defect in C57Bl/6J
mice. *Am J Physiol Regul Integr Comp Physiol.* Dec 2009;297(6):R1660-9.
doi:10.1152/ajpregu.00147.2009

20.   Vasquez F, Tiscornia C, Lorca-Ponce E, Aicardi V, Vasquez S. Cardiorenal
Syndrome: Molecular Pathways Linking Cardiovascular Dysfunction and Chronic Kidney
Disease Progression. *Int J Mol Sci.* Aug 1 2025;26(15)doi:10.3390/ijms26157440

21.   Chen J, Zhou Q, Su L, Ni L. Mitochondrial dysfunction: the hidden catalyst in
chronic kidney disease progression. *Ren Fail.* Dec 2025;47(1):2506812.
doi:10.1080/0886022X.2025.2506812

22.   Chevalier RL. Bioenergetics: the evolutionary basis of progressive kidney
disease. *Physiol Rev.* Oct 1 2023;103(4):2451-2506. doi:10.1152/physrev.00029.2022

23.   Verissimo T, de Seigneux S. New evidence of the impact of mitochondria on

kidney health and disease. *Nat Rev Nephrol.* Feb 2024;20(2):81-82.

doi:10.1038/s41581-023-00803-z

24.   McCrimmon A, Domondon M, Sultanova RF, Ilatovskaya DV, Stadler K.

Comprehensive assessment of mitochondrial respiratory function in freshly isolated

nephron segments. *Am J Physiol Renal Physiol.* May 1 2020;318(5):F1237-F1245.

doi:10.1152/ajprenal.00031.2020

25.   Bhargava P, Schnellmann RG. Mitochondrial energetics in the kidney. *Nature*

Reviews Nephrology. Oct 2017;13(10):629-646. doi:10.1038/nrneph.2017.107

26.   Hall AM, Rhodes GJ, Sandoval RM, Corridon PR, Molitoris BA. In vivo

multiphoton imaging of mitochondrial structure and function during acute kidney injury.

Kidney Int. Jan 2013;83(1):72-83. doi:10.1038/ki.2012.328

27.   Hall AM, Unwin RJ, Parker N, Duchen MR. Multiphoton imaging reveals

differences in mitochondrial function between nephron segments. *J Am Soc Nephrol.*

Jun 2009;20(6):1293-302. doi:10.1681/ASN.2008070759

28.   Hall AM, Unwin RJ. The not so 'mighty chondrion': emergence of renal diseases

due to mitochondrial dysfunction. *Nephron Physiol.* 2007;105(1):p1-10.

doi:10.1159/000096860

29.   Rosca MG, Mustata TG, Kinter MT, et al. Glycation of mitochondrial proteins from

diabetic rat kidney is associated with excess superoxide formation. *American Journal of*

Physiology-Renal Physiology. 2005 2005;doi:10.1152/ajprenal.00415.2004

30.   Bufi R, Korstanje R. The impact of genetic background on mouse models of

kidney disease. *Kidney Int.* Jul 2022;102(1):38-44. doi:10.1016/j.kint.2022.03.020

31. Doerrier C, Garcia-Souza LF, Krumschnabel G, Wohlfarter Y, Mészáros AT,

Gnaiger E. High-resolution fluorespirometry and oxphos protocols for human cells,

permeabilized fibers from small biopsies of muscle, and isolated mitochondria. *Methods*

in Molecular Biology. 2018.

32. Schock-Kusch D, Xie Q, Shulhevich Y, et al. Transcutaneous assessment of

renal function in conscious rats with a device for measuring FITC-sinistrin

disappearance curves. *Kidney Int*. Jun 2011;79(11):1254-8. doi:10.1038/ki.2011.31

33. Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for

biological-image analysis. *Nat Methods*. Jun 28 2012;9(7):676-82.

doi:10.1038/nmeth.2019

34. Arzt M, Deschamps J, Schmied C, et al. LABKIT: Labeling and Segmentation

Toolkit for Big Image Data. Methods. *Frontiers in Computer Science*. 2022-February-10

2022;Volume 4 - 2022doi:10.3389/fcomp.2022.777728

35. Spiess E, Muller D, Dinser M, et al. Automatic Segmentation of Histopathological

Glioblastoma Whole-Slide Images Utilizing MONAI. *Stud Health Technol Inform*. May 15

2025;327:88-92. doi:10.3233/SHTI250279

36. Tejashwini PS, Thriveni J, Venugopal KR. Enhanced abdominal multi-organ

segmentation with 3D UNet and UNet + + deep neural networks utilizing the MONAI

framework. *Abdom Radiol (NY)*. Jun 30 2025;doi:10.1007/s00261-025-05041-4

37. Diaz-Pinto A, Alle S, Nath V, et al. MONAI Label: A framework for AI-assisted

interactive labeling of 3D medical images. *Med Image Anal*. Jul 2024;95:103207.

doi:10.1016/j.media.2024.103207

38.   Luchian A, Ressel L. Combined colour deconvolution and artificial intelligence
approach for region-selective immunohistochemical labelling quantification: The
example of alpha smooth muscle actin in mouse kidney. *J Biophotonics*. Jan
2024;17(1):e202300244. doi:10.1002/jbio.202300244

39.   Landini G, Martinelli G, Piccinini F. Colour deconvolution: stain unmixing in
histological   imaging.   *Bioinformatics*.   Jun   16   2021;37(10):1485-1487.
doi:10.1093/bioinformatics/btaa847

40.   Ruifrok AC, Johnston DA. Quantification of histochemical staining by color
deconvolution. *Anal Quant Cytol Histol*. Aug 2001;23(4):291-9.

41.   Opurum PC, Decker ST, Stuart D, et al. Combined sedentarism and high-fat diet
induce early signs of kidney injury in C57BL/6J mice. *Am J Physiol Renal Physiol*. Jun 1
2025;328(6):F850-F860. doi:10.1152/ajprenal.00259.2024

42.   Elkin LA, Allen PG, Kay M, Higgins JJ, Wobbrock JO. An Aligned Rank Transform
Procedure for Multifactor Contrast Tests; An Aligned Rank Transform Procedure for
Multifactor Contrast Tests. 2021 2021;15(21)doi:10.1145/3472749.3474784

43.   Wobbrock JO, Findlater L, Gergle D, Higgins JJ. The Aligned Rank Transform for
nonparametric factorial analyses using only ANOVA procedures. In: *Conference on*
Human Factors in Computing Systems - Proceedings. ACM Press; 2011:143-146.

44.   Hall AM, Unwin RJ. The not□so□simple proximal tubule. *Am J Physiol Renal*
Physiol. 2007;

45.   Rosca MG, Mustata TG, Kinter MT, et al. Glycation of mitochondrial proteins from
diabetic rat kidney is associated with excess superoxide formation. 2005.

46.   Nelson MAM, Fisher-Wellman KH. Mitochondrial Diagnostics: A Discovery-Based

Biochemical Platform for Phenotyping Human Peripheral Blood Cell Mitochondria.

Methods Mol Biol. 2021;2277:371-389. doi:10.1007/978-1-0716-1270-5_23

47.   Fisher-Wellman KH, Davidson MT, Narowski TM, Lin CT, Koves TR, Muoio DM.

Mitochondrial Diagnostics: A Multiplexed Assay Platform for Comprehensive

Assessment of Mitochondrial Energy Fluxes. *Cell Rep*. Sep 25 2018;24(13):3593-3606

e10. doi:10.1016/j.celrep.2018.08.091

48.   Decker ST, Opurum PC, Choi RH, et al. A Platform for Mitochondrial Profiling in

Enriched Kidney Segments Under Thermodynamic Control. *bioRxiv*.

2025:2025.05.05.652276. doi:10.1101/2025.05.05.652276

49.   Decker ST, Funai K. Mitochondrial membrane lipids in the regulation of

bioenergetic flux. *Cell Metab*. Sep 3 2024;36(9):1963-1978.

doi:10.1016/j.cmet.2024.07.024

50.   Pennington ER, Funai K, Brown DA, Shaikh SR. The role of cardiolipin

concentration and acyl chain composition on mitochondrial inner membrane molecular

organization and function. *Biochim Biophys Acta Mol Cell Biol Lipids*. Jul

2019;1864(7):1039-1052. doi:10.1016/j.bbalip.2019.03.012

51.   Heden TD, Johnson JM, Ferrara PJ, et al. Mitochondrial PE potentiates

respiratory enzymes to amplify skeletal muscle aerobic capacity. *Science Advances*.

2019/09/11/ 2019;5(9):eaax8352. doi:10.1126/sciadv.aax8352

52.   Verkerke ARP, Ferrara PJ, Lin C-T, et al. Phospholipid methylation regulates

muscle metabolic rate through Ca2+ transport efficiency. *Nature Metabolism*. 2019/09//

2019;1(9):876-885. doi:10.1038/s42255-019-0111-2

53.   Heden TD, Neufer PD, Funai K. Looking Beyond Structure: Membrane

Phospholipids of Skeletal Muscle Mitochondria. *Trends in Endocrinology & Metabolism*.

2016/08/01/ 2016;27(8):553-562. doi:10.1016/j.tem.2016.05.007

54.   Hu C, Lakshmipathi J, Stuart D, Kohan DE. Profiling renal sodium transporters in

mice with nephron Ift88 disruption: Association with sex, cysts, and blood pressure.

Physiol Rep. Mar 2022;10(5):e15206. doi:10.14814/phy2.15206

Western Blot		
Target	Dilution	Source
Aquaporin 2 (AQP2)	1:1000	Santa Cruz (sc-47710)
Sodium-Potassium-Calcium Transporter (NKCC2)	1:1000	Kohan Lab[54]
Sodium-glucose cotransporter (SGLT2)	1:1000	Santa Cruz (sc-393350
Podocin	1:1000	Invitrogen (PA5-79757)
Sodium-potassium ATPase (Na/K$^+$-ATPase)	1:5000	Cell Signaling (#30105)
Tumor Necrosis Factor alpha (TNF-α)	1:1000	Cell Signaling (#3707)
GAPDH	1:1000	
Actin	1:1000	Sigma-Aldrich (A2066)
Citrate Synthase (CS)	1:1000	Abcam (ab9660)
OXPHOS Cocktail	1:1000	Abcam (STN-19467)
PERK	1:5000	Cell Signaling (#3192)
BiP	1:1000	Cell Signaling (#3177)
CHOP	1:1000	Santa Cruz (sc-793)
qPCR		
Target	Forward Sequence	Reverse Sequence
KIM-1	5'-CTGGAATGGCACTGTGACATCC-3'	5'-GCAGATGCCAACATAGAAGCCC-3'
Lcn2	5'-CACTGACTACGACCAGTTTGCC-3'	5'-GATGTTGTTATCCTTGAGGCCC-3'
PGC1-α	5'-TGAGGACCGCTAGCAAGTTT-3'	5'-TGAAGTGGTGTAGCGACCAA-3'
Col1a1	5'-GACCAAAAGGTGATGCTGGACAG-3'	5'-CAAGACCTCGTGCTCCAGTTAG-3'
Col3a1	5'-CCTCAGGGTATTGCTGGACAAC-3'	5'-CAGAAGGACCTTGTTTGCCAGG-3'
NFκB	5'-GCTGCCAAAGAAGGACACGACA-3'	5'-GGCAGGCTATTGCTCATCACAG-3'
IL-6	5'-TACCACTTCACAAGTCGGAGGC-3'	5--CTGCAAGTGCATCATCGTTGTTC-3'
SGK1	5'-CTCATTCCAGACCGCTGACAAAC-3'	5'-CCAAGGCACTGGCTATTTCAGC-3'
CCL2	5'-GCTACAAGAGGATCACCAGCAG-3'	5'-GTCTGGACCCATTCCTTCTTGG-3'

Figure 1. Experimental setup (A) and composition of the diets (B) of the present study. Body

weight (C) and liver mass (D) were not different between groups at the time of termination;

however, kidney mass (E) was significantly greater in high-fat, high-salt (HFHNa) fed mice

compared to their respective controls. Free-living, per-cage energy intake (F) was greater in the

HFHNa-fed mice than in the chow-fed mice, as was fat (G), salt (H), and water (I) intake.

Individual urine output (J) collected during a 24-hour period in metabolic cages was also

increased in the HFHNa-fed mice.

Figure 2. Glomerular filtration rate (GFR; A & B) was mildly (~20%) decreased in the 6J high-fat,

high-salt-fed (HFHNa) compared with chow-fed mice, but no differences between diet groups

were observed in 6N mice. However, 24-hour urine albumin (C), creatinine (D), and albumin

were significantly increased in the HFHNa-fed mice, while plasma blood urea nitrogen (BUN; F)

was significantly decreased. Albumin-to-creatinine ratio (ACR; F) was likewise significantly

increased in the 6J strain, but there were no significant differences in the BUN-to-creatinine ratio

(G). In agreement with other markers of impaired kidney function, urine cystatin C (H), kidney

injury marker 1 (KIM-1; I), and lipocalin 2 (LCN2; J) were increased by HFHNa feeding.

Figure 3. A representative image of Masson's trichrome stain is shown in panel A. A high-fat,

high-salt diet (HFHNa) significantly increased fibrosis area in the 6N, but not 6J, mice.

Glomerular-specific analysis of the glomerular corpuscle, tuft, Bowman's space, and fibrosis

areas was then assessed (C). There was no significant difference in corpuscle area (D), tuft

area (E & H), or Bowman's space area (F & I). However, HFHNa diet decreased the absolute

(G) but not the relative (J) fibrosis stain area.

Figure 4. Representative images of western blots are shown in panel A. A high-fat, high-salt diet

(HFHNa) resulted in a significantly increased protein abundance of aquaporin 2 (AQP2; B), but

had no effect on sodium-potassium cotransporter 2 (NKCC2; C). On the other hand, HFHNa

feeding resulted in a significant decrease in the protein abundance of the sodium-glucose

cotransporter (SGLT2; D). No changes were observed in the protein abundance of the ER

stress markers BIP (E), CHOP (F), or PERK (G). mRNA abundance of kidney injury marker

(KIM-1; H) was increased in only the 6J HFHNa-fed mice. Diet did not significantly alter the

mRNA abundance of lipocalin 2 (LCN2; I), NFκB (J), interleukin-6 (IL-6; K), C-C motif

chemokine ligand 2 (CCL2; L), or serum and glucocorticoid-induced protein kinase 1 (SGK1; M).

Figure 5. Mitochondrial respiration (JO_2) in isolated mitochondria was not impacted by diet

following a maximal (A) or submaximal (B) protocol. However, ATP production (JATP; C) was

significantly impaired in the mice fed a high-fat, high-salt (HFHNa) diet. Consequentially,

mitochondrial ATP-to-O ratio (P:O Ratio; C) was decreased in the HFHNa-fed mice. These

differences occurred despite no changes in mitochondrial citrate synthase (CS) protein

abundance in whole-tissue lysate (F) or isolated mitochondria (G) preparations.

Figure 6. Urine proteomic analysis in a subset (n = 3 per group) of 6J mice (A) revealed that

high-fat, high-salt (HFHNa) feeding significantly increased the presence of several urinary

proteins. Several of these excreted proteins included those related to kidney injury, proteostasis,

cell stability, and membrane transport (B). Collectively (C), our data support the hypothesis that

a HFHNa diet induces impairments in mitochondrial ATP production, which drives decreases in

glomerular filtration rate (GFR) and altered solute transport, leading to metabolic dysfunction-

associated kidney disease (MDAKD).

Figure 2

Figure 3

Figure 4

A

B

Kidney Injury
- Alb — p = 0.058
- Cst3 — p = 0.142
- Lcn2 — p = 0.180
- Alpl — p = 0.057
- Vil1 — p = 0.054

Proteostasis
- Gpc1 — **p = 0.033**
- Atp6v0c — p = 0.055
- Sppl2a — **p = 0.031**
- Ppic — **p = 0.043**
- Aoah — p = 0.307
- Atraid — p = 0.084
- Pdcd6 — **p = 0.025**
- Creg1 — p = 0.059

Transport Stability
- Cadm4 — **p = 0.024**
- Rhoc — **p = 0.043**
- S100a11 — **p = 0.026**
- Sdha — **p = 0.046**
- Aldob — p = 0.050
- Aldh3b2 — p = 0.070
- Slc22a19 — **p = 0.038**

WT   HF/HNa

C

Figure 6