Growth and molecular responses of potato to lunar regolith simulants

David Handy, Anika Loeffler, Medora Knudson, Sydney Campbell, Pankaj Jaiswal, Jeffrey C. Anderson, Aymeric Goyer*

Oregon State University, United States, Department of Botany and Plant Pathology, Corvallis

* Corresponding author:

Aymeric Goyer

Aymeric.goyer@oregonstate.edu

Abstract

Background

On-site food production will be required to achieve NASA's goal of a sustainable Lunar habitat. Toward this end, the use of fine, soil-like material on the Lunar surface, known as regolith, has been proposed as a plant growth substrate. However, how this substrate may affect plant growth is not well understood. Lunar regolith is devoid of the organic materials that make soils on earth fertile for plant growth, and has been weathered by solar winds, cosmic rays, and micrometeorite impacts. Additionally, regolith at certain lunar sites may contain heavy metals. These metal ions may leech, thus posing challenges with accumulation in plant material. To address and verify the efficacy of regolith-based crop production, we used lunar regolith simulants (LRS). We investigated the effects of LRS on potato (*Solanum tuberosum* cv 'Modoc') plant and tuber development, gene expression, and nutrition profiles.

Results

Growth in LRS negatively impacted the potato plant size and tuber yield. While the degree of impact differed between simulants, all plants grown in LRS were statistically significantly shorter in height than plants grown in control soil. Further experiments with the lunar mare simulant 1E (LMS-1E) show that these effects can be ameliorated through the addition of vermicompost, an organic component, with a 70:30 v/v ratio of LMS-1E to compost being virtually indistinguishable from controls. Changes in gene expression profiles also differed between simulants, with genes related to

photosynthesis, biotic and abiotic stress responses, signaling, and terpenes and

flavonoids metabolism being commonly altered. Despite these observed differences in

transcription, broad changes in metabolite profiles were not observed.

Conclusions

LRS are clearly stressful on plants. However, amendment of the substrate with

composted materials appears to be a viable strategy to alleviate stress. Given these

observations, regolith-based agriculture may not be viable for very early food production

when organic matter content is low. However, this would improve over time with

continual incorporation of organic matter to regolith. As such, we believe regolith-based

agriculture is a viable long-term strategy.

Keywords

Potato, lunar regolith, omics, space biology, heavy metals

Background

It is the goal of NASA's Artemis missions to establish a permanent human

presence on the Moon in the 2030s (https://www.nasa.gov/feature/artemis/). However,

achieving a sustainable and affordable presence on the lunar surface will require on-site

food production to reduce resupply costs and increase mission success probabilities [1].

Regolith-based agriculture has been proposed as an *in-situ* method of supporting crops

for food production. However, surface lunar regolith as a viable substrate needs to be

verified to ensure adequate food production and food safety. As genuine lunar regolith is

not available for large-scale plant studies, lunar regolith simulants (LRS) are commonly

used for experimentation [2–4].

Lunar regolith falls into two categories: lunar mare and lunar highlands. The

primary component in lunar mare regolith is basalt formed by lava flows during the

volcanically active phase, while the geologically older, crater-filled lunar highlands

primarily consist of ferroan anorthosite [5]. The most unique feature of lunar regolith

compared to terrestrial soils and LRS is the presence of agglutinates, aggregate

particles formed by micrometeorite impacts and comprised of nanophase iron, mineral

fragments, and glass. While LRS often appropriately contains volcanic glass,

agglutinates are more difficult to manufacture and are often excluded from LRS. While

lacking nitrogen, lunar regolith and its simulants contain micronutrients such as

phosphorous, potassium, magnesium, iron, manganese, and calcium [6,7]. While these

nutrients may appear replete, it should be noted that the bioavailability of some, such as

phosphorous, is dependent on pH and the presence of other elements.

Potato (*Solanum tuberosum*) is an ideal candidate for space crop production

because it is high-yielding, clonally propagated, rich in calories and phytonutritients, and

has a high satiety index. Potato is also a good model organism for plants that produce

underground tubers. It has a rich history of available genetic resources in the form of

germplasm accessions, sequenced genome and public access to many large-scale

omics datasets for undertaking comparative studies. Early experiments involving lunar

samples returned from Apollo-era missions exposed potato seedlings, among other

plants, to suspensions of lunar regolith with no reported effects on plant development

[8]. Most recently, *Arabidopsis thaliana* grown in lunar regolith returned from three

Apollo sites had stunted roots, and slowed aerial growth compared to a LRS, and each

regolith induced unique overall gene expression changes [9]. However, plants showed

abbreviated growth and arrest in the vegetative stages, activated stress response

pathways and negatively impacted the telomerase activity and shortened telomeres[10].

Negative effects on potato tuber yield has been reported when grown in Martian regolith

simulant [11], but the effects of LRS on potato growth and yield has not yet been

ascertained.

Here, we present findings from two greenhouse experiments performed on the

potato cultivar 'Modoc'. In the first experiment, we tested potato growth on LRS

amended with different percentages of organic matter. This enabled us to find a ratio

LRS/compost that supports sufficient plant growth for analyzing growth phenotype traits,

physiology, and metabolism but is also suboptimal to investigate the stress response to

LRS. Data was used to design our second experiment. In the second experiment, we

assessed the effects of a variety of LRS types mixed with compost on growth,

physiology, leaf and tuber metabolite profile, and leaf gene expression. To our

knowledge, this is the first report of the response of potato to LRS with fully-grown

plants.

Methods

Chemicals

97   Chlorogenic acid, HPLC-grade acetonitrile, HPLC-grade methanol, and meta-

98   phosphoric acid were from VWR (Radnor, PA, USA). L-Phenylalanine, L-tyrosine,

99   ascorbic acid, solanine, chaconine, methoxamine, pyridine, and ribitol were from

100   Millipore-Sigma (St. Louis, MO, USA). N-methyl-N-(trimethylsilyl) trifluoroacetamide with

101   chlorotrimethylsilane was purchased from CovaChem (Loves Park, IL, USA).

Potato and growth substrate materials

Mini-tubers of the potato cultivar Modoc were obtained from CSS Farms (Colorado City,

CO, USA). Lunar regolith simulants LHS-1E and LMS-1E were purchased from the

Exolith Lab (Orland, FL, USA), and JSC-1A, OPRH4W30, and NUW-LHT-5M were

obtained from NASA. The Adkins-series soil was obtained from the top 30-cm layer of a

non-cultivated, Rye grass field from Umatilla County, Oregon (45.817318° N -

119.294922° W). Vermicompost was sourced from Earthworm Castings (Green Bay, WI,

USA).

Growth conditions

Experiment 1: Determination of optimal LRS/compost ratio

Vermicompost was selected as the green compost to mimic the use of inedible

plant matter as a soil amendment in a bioregenerative life support system. Mixtures

were prepared by combining 5% non-autoclaved vermicompost with 10% or 25%

autoclaved vermicompost in 85% or 70% LMS-1E (volume ratios), respectively. The

consistent amount of non-autoclaved vermicompost between batches enabled us to

obtain mixtures with the same starting microbiome. Adkins-series soil was used as a

control of optimal potato growth and was mixed with vermicompost in the same ratios as

LRSs. Mini-tubers were planted on 2 August 2024 in 8.89 x 7.62 cm square pots lined

with landscaping fabric (Greenhouse Megastore, Danville, IL, USA), containing

approximately 300 cm^3 of soil mixture. Each treatment was done with four biological

replicates with one plant per pot. Pots were placed in a fixed, randomized configuration

under a PlantEye600 scanner (Phenospex, Heerlen, The Netherlands) for phenotyping,

with a border row of plants around the experimental plants to reduce edge-effects.

Plants were grown using natural sunlight in a greenhouse set to 22℃ day and 18℃

night. Plants were watered with 40 mL of reverse osmosis filtered water every 2-3 days,

increasing frequency as plants matured.

Plants were scanned using the PlantEye600 scanner every 12 hours for the first

six weeks after emergence until plant foliage from different plants overlapped (Table S1

& S2). The scanner recorded digital biomass, greenness, height, hue, leaf angle, leaf

area, normalized difference vegetation index (NDVI), normalized pigments chlorophyll

ratio index (NPCI), and plant senescence reflectance index (PSRI). At nine weeks, a

MultispeQ (PhotosynQ Inc., East Lansing, MI, USA) was used to analyze plant leaves

for pigment content, and photosynthetic parameters such as quantum yield,

electrochromic shift, non-photochemical quenching, etc. (Table S3 &S4). Plants were

harvested on 15 October 2024, 74 days after planting (DAP). Plants were removed from

the substrate and the roots and tubers rinsed with tap water. The substrate was sieved

to remove any remaining roots, stolons, and Osmocote pellets, then collected for

chemical and microbial analysis (see below). For additional phenotyping

measurements, total mass of above- and below-ground tissues was first measured,

then plant parts were separated to measure shoot (leaves and stems) mass, root mass,

and tuber mass. Tuber and total masses were used to calculate harvest index as a

percentage of edible biomass (100 x edible biomass/total biomass). Leaves, roots, and

tubers were flash frozen in liquid nitrogen, lyophilized, and ground to powder with mortar

and pestle for further analysis.

Experiment 2: Effect of various lunar regolith simulant types

Adkins series soil and LRSs LHS-1E, LMS-1E, JSC-1A, OPRH4W30, and NUW-

LHT-5M were mixed with vermicompost (5% v/v). Mini-tubers were planted once sprouts

were approximately 2-cm long. Five tubers were planted in each soil-type (one tuber per

pot) in 8.89 x 7.62 cm square pots lined with landscaping fabric and arranged in a

randomized format. Plants were scanned with a Phenospex PlantEye600 scanner every

12 hours for the initial 35 days of growth (5 May through 9 June) until plants outgrew the

spacing required for scanning. Leaf samples were collected from the upper canopy at

39 DAP and flash frozen for RNA-seq and metabolite analysis. We did not collect leaves

from plants grown in NUW-LHT-5M because plants were stunted and did not have

leaves. Plants were harvested at 59 DAP and tubers were weighed, flash frozen in

liquid nitrogen, lyophilized, and ground to powder with mortar and pestle for later

metabolite analysis.

Soil and tissue analyses

Electroconductivity (EC), pH, moisture, microbial respiration, organic matter

content and mineral content, including heavy metals, of soil samples and mineral

content of leaf and tuber samples were measured by the Oregon State University Soil

Health Lab. For Experiment 1, soil and freeze-dried tissue samples of the same

treatment were pooled by measuring and combining equivalent portions of three

biological replicates. For experiment 2, four biological replicates were analyzed

independently. Additionally, a sample was reserved from the pre-growth batch of each

LRS for analysis. EC and pH of soil were determined by suspending the substrate in

deionized water in a 1:1 ratio and measured using a Hanna HI5522 instrument (Hanna

Instruments, Woonsocket, RI, USA) with either a temperature and pH probe, or an EC

probe as appropriate. Microbial respiration was determined by carbon dioxide burst test.

Wet substrate was incubated in a vessel with an airtight seal at 23°C for up to 96 h.

Carbon dioxide was measured using a Picarro isotopic CO_2 analyzer (Picarro, Inc.,

Santa Clara, CA, USA) at 0 h, 24 h, and 96 h. For Experiment 1, minerals were

analyzed as follows. Five hundred mg of substrate or 250 mg of plant tissue were

acidified using 10 mL nitric acid, then analyzed via microwave digestion using an Anton

Paar microwave (Ashland, VA, USA), followed by elemental determination via

Inductively Coupled Plasma-Optical Emission Spectrometry (ICP-OES) using an Agilent

5110 instrument (Agilent Technologies, Inc., Santa Clara, CA, USA). For Experiment 2,

soil minerals were extracted by adding 20 mL of Mehlich-3 extractant (20 g L^{-1} NH_4NO_3,

5.556 mg mL^{-1} NH_4F, 0.372 mg mL^{-1} EDTA, 11.5 mL L^{-1} glacial acetic acid, and 0.82 mL

L^{-1} HNO_3) to 2 g of sample, shaking for 5 minutes, filtered, then analyzed via ICP-OES.

Plant tissue was analyzed using the same method as experiment 1. Note that heavy

metal analysis in experiment 1 characterized total elemental content and did not

account for bioavailable versus non-bioavailable forms of these metals. Therefore,

analysis of these elements in plant tissues will likely provide more meaningful

information than those made on soil mixtures. The Mehlich-3 extraction was used in

experiment 2 because this method focuses on bioavailable forms of these metals and

nutrients.

Untargeted metabolite analysis using gas chromatography-mass spectrometry (GC-MS)

Tissue extracts were analyzed by GC-MS as previously described [12]. Briefly, 50

mg of dried tissue was extracted using a water:methanol:chloroform (1:2.5:1) solution with

ribitol (40 µg mL^{-1}) as an internal standard. Extracts were centrifuged to remove cellular

debris, then phase separated to isolate aqueous compounds. Fifty µL aliquots were

frozen and lyophilized then stored at -80 °C. Samples were derivatized by resuspension

of lyophilized material into 20 µL methoxamine HCl (30 mg mL^{-1}) dissolved in pyridine and

incubating the sample at 37 °C for 90 min with shaking. Then, 40 µL of N-methyl-N-

(trimethylsilyl) trifluoroacetamide with 1% chlorotrimethylsilane was added to each sample

followed by a 30-min incubation at 37 °C with shaking. One µL of each sample was

injected with a 4:1 split for tuber samples and a 10:1 split for leaf samples on an Agilent

7890B GC equipped with an Agilent 5977B MSD (Agilent Technologies). The sample

injection order was randomized, and the reagent blank was injected after every three

samples to control for potential carryover. Resulting spectra were analyzed using AMDIS

[13] and the Agilent Fiehn 2013 GC/MS Metabolomics RTL Library [14] to identify

components.

Targeted metabolite analysis using High-Performance Liquid Chromatography (HPLC)

For HPLC analysis, 100 mg of ground, freeze-dried tuber tissue was extracted in 1 mL of a 50% methanol, 2.5% metaphosphoric acid, 1 mM EDTA extraction solution. Samples were shaken for 10 minutes at room temperature in the dark, then centrifuged at 14,000 x g for 10 minutes. Supernatant was moved to a new tube and re-centrifuged to remove debris. The supernatant was transferred to amber HPLC vials for injection. Metabolites were analyzed using standardized methods as previously published [15]. Standards were prepared in two combination sets: phenolic compounds (ascorbic acid, chlorogenic acid, tyrosine, tryptophan, and phenylalanine) and glycoalkaloids (solanine and chaconine). Phenolic compound standards were prepared with a seven-step 2x serial dilution ranging from 400 to 6.125 mg L^{-1}. Serial dilutions of glycoalkaloid standards were prepared at 400, 300, 200, 100, 50, 25, 12.5 and 6.125 mg L^{-1}. Each standard was injected in triplicate to generate standard curves used to calculate the compound concentrations in each sample. Compounds were analyzed using an UltiMate 3000 UHPLC system (Thermo Fisher Scientific, Waltham, MA, USA) with an LPG-3400SD quaternary analytical pump, solvent degasser, WPS-3000TSL autosampler, and TCC-3000 column compartment. Compounds were separated using an Acclaim Polar Advantage II C18 column (4.6 x 150 mm, 5 µm) (Dionex, Sunnyvale, CA, USA) using previously published methods [15,16] . Compounds were detected by UV absorbance using a diode array detector DAD 3000 (Thermo Fisher Scientific).

RNA sequencing

Total RNAs were extracted using a PureLink RNA mini kit (Invitrogen, Waltham, MA, USA), followed by treatment with DNase I (DNA-*free* DNA removal kit (Invitrogen)) to remove traces of genomic DNA. RNA integrity was verified on agarose gel and by analysis on an Agilent Bioanalyzer 2100 (Agilent Technologies, Inc). RNAs were then sent to Novogene for sequencing on an Illumina Novaseq X Plus Series (150-bp, paired-ends) as previously described [17]. Raw .fastq files were run through cutadapt to remove adapter sequences. The DM v6.1 doubled monoploid potato genome and Gene Ontology (GO) term annotations were downloaded from spuddb.uga.edu [18]. Reads were aligned to the genome using HISAT2 [19], sorted using SAMtools [20], and mapped using STRINGTIE [21]. Gene counts were then analyzed in R using the DESeq2 package to generate lists of differentially expressed genes (DEGs), principal component analysis (PCA) plots, and heatmaps [22]. Replicate A of the OPRH4W30 treatment was removed as an outlier based on PCA clustering. Gene set enrichment analysis (GSEA) was performed using the ClusterProfiler package in R [23,24] and used to generate dotplots.

Statistical analysis

Statistical analyses of GC-MS data were performed using MetaboAnalyst 6.0 (https://www.metaboanalyst.ca/) using default processing settings and data filtering and using the reference feature ribitol for normalization. Features with missing values across replicates were excluded, and a 5% low-variance filter was applied based on the interquantile range. Principle component analyses were generated based on these

251 settings. For other data including harvest metrics, LCMS measurements, PhenospeX

252 data, and PhotsynQ measurements, data was analyzed via ANOVA in R using the 'dplyr'

253 and 'multcompview' packages. Figures were generated using the 'ggplot2',

254 'RColorBrewer' and 'ggpubr' packages.

Results

Experiment 1: Determination of suboptimal LRS/compost

ratio

In this experiment, we tested potato growth on lunar mare simulant LMS-1E

amended with varying amounts of vermicompost. Our goal was to find a ratio

LRS/compost that supports sufficient plant growth for analysis of growth, physiology,

and metabolism but is also suboptimal to investigate the stress response to LRS. In

early stages of development, no statistically significant differences in growth and

physiology phenotypes recorded by the PlantEye600 scanner were observed (Table S1

& S2). This was likely due to plants emerging from the soil asynchronously, creating

large variations within treatment groups at early time points. However, plant growth had

leveled out by harvest, and statistically significant differences were observed for multiple

growth-related phenotypes (Figure 1). Plants grown in 100% regolith simulant were

severely impacted, with reduced total, shoot, root and tuber biomass, as well as harvest

index, compared to control plants grown in Adkins soil (Figure 1 & Figure **2**, Table S5).

Most notably, the roots of three out of four plants grown in 100% regolith were too small

to be weighed on the available balance (less than 0.01 g). Plants grown in 70% regolith

simulant were statistically larger than those in 100% regolith in all mass metrics. For

some phenotypes,70% LMS grown plants were similar to plants grown in Adkins soil,

and even outperformed Adkins soils in tuber mass (Figure 1F). The impact on total tuber

mass was also notable, as this translates to the edible biomass of the plant. Plants

grown in 85% regolith had a biomass between that of plants grown in 100% regolith

simulant and Adkins soil, with no statistically significant difference to either treatment.

The variable ratios of compost amendment in the Adkins series control soil had no

significant effect on any plant phenotypes. While some photosynthesis related

phenotypes measured using PhotosynQ were found to have statistically significant

differences, clear patterns across related parameters were not observed (Table S3 &

S4).

Figure 1. Harvest phenotypes of potato plant variety Modoc grown in variable ratios of LMS-1E and compost. A) Total biomass, B) Shoot biomass, C) Tuber number, D) Total tuber mass, E) Harvest index, and F) Average tuber mass. Shared letters indicate statistically similar groupings as determined by one-way ANOVA and Tukey's tests.

Figure 2. Post-harvest images of potato plants grown in LMS-1E (left) and Adkins soil (right), in

100%, 85%, or 70% regolith to compost volume ratios.

Next, we analyzed soil samples for various metrics to find clues that could

explain the phenotypic differences observed. The results presented hereafter are based

on analysis of pooled biological replicates, so statistical analysis was not possible.

However, several trends are worth noting. Analysis of pre-growth LMS-1E substrate

revealed differences in soil characteristics with compost amendment (Table S6). The pH

of 100% LMS-1E was 7.9. However, this value decreased as vermicompost content

increased, which is expected as pH of vermicompost is between 6 and 7. Increased

compost ratio also correlated with an increase in electroconductivity (EC), microbial

respiration, and detectable heavy metals. Note that some of these metals most likely

come from the Osmocote fertilizer, *i.e.*, Cu, Ni, Zn, as these elements are not present in

LMS-1E [4]. Post-growth analysis showed a consistent decrease in pH and increase in

EC as compared to pre-growth analyses (Table S6). A drop in pH is expected as plants

take up essential ions like K^+ and Ca^{2++} and compensate by releasing protons into the

soil. Post-growth concentrations of heavy metals Al, Cr, Cu, Fe, Ni and Zn were higher

than pre-growth. In addition, post-growth microbial respiration decreased compared to

pre-growth analysis for 85% and 70% regolith simulants, while 100% regolith had

slightly increased levels. It should be noted that microbial respiration differed between

85% and 70% regolith treatment, despite containing the same amount of non-sterilized

compost. This is likely due to the method of assessment, which involves incubating the

soil for 24-96 h, during which the microbes reproduce at different rates based on the

quantity of compost in the substrate.

To further understand what factors might have led to underdeveloped roots and

overall decreased plant growth in 100% regolith simulant, we quantified heavy metal

contents in leaves and tubers (Table S7). The most striking observation was the large

accumulation of copper in leaves compared to all other treatments, and to a lesser

extent in tubers, of plants grown in 100% regolith simulant compared to plants grown in

Adkins soil and 70% and 85% regolith simulant, suggesting that copper concentrations

available to plants were at toxic levels compared to Adkins.

Experiment 2: Effect of various lunar regolith simulant types

The growth of potato plants varies depending on LRS type

In Experiment 1, we found a relatively linear relationship between added compost and improvement of plant growth. This information was used to inform the design of the second experiment. Stunting caused by 100% regolith simulant limits analyses due to lack of sample, while 70% and 85% simulant may mask stress responses caused by the regolith. As such, we used a 95:5 v/v ratio of regolith to compost in this second experiment. In addition, we planted actively sprouting mini-tubers (approximately 2-cm long sprouts) to decrease variability in plant development due to asynchronous emergence of shoots from tubers. We selected three lunar highland simulants (LHS-1E, OPRH4W30, and NUW-LHT-5M) and two lunar mare simulants (JSC-1A and LMS-1E). Plants grown in lunar highland simulant NUW-LHT-5M were the most severely affected, with statistically significant reduction in plant size and tuber biomass (Figures 3 & 4). This can be partly explained by the severely delayed emergence of plants observed in this LRS (approximately 24 DAP compared to 4-8 DAP for other types of LRS). Additionally, only three out of five plants grown on NUW-LHT-5M grew large enough to be detected by the Phenospex scanner and produced tubers (Tables S8 & S9). Statistically significant reduction in tuber mass was also observed in all other LRS types compared to Adkins soil (Figure 4), while plant size was statistically reduced only in lunar highland simulant OPRH4W30 compared to Adkins when analyzing digital canopy biomass (Figure 3). Colorimetric indices did not have any statistical differences between treatments (Table S8).

Figure 3. (A) Digital above-ground biomass across the first 34 days after planting (DAP) of
potato plants grown in Adkins soil (ADK) and five regolith simulants; JSC-1A (JSC), LHS-1E
(LHS), LMS-1E (LMS), NUW-LHT-5M (NUW), and OPRH4W30 (OPR). Data was collected via
the Phenospex PlantEye600 scanner. (B) Boxplot of final above-ground digital biomass at 34
DAP. All substrates were amended with 5% compost by volume. Shared letters indicate
statistically significant groupings determined by ANOVA followed by Tukey's post-hoc. N=5
except NUW-LHT-5M where N=3.

Figure 4. Harvested tuber mass of plants. Tubers were harvested at 59 DAP from plants grown in Adkins soil (ADK) and five regolith simulants; JSC-1A (JSC), LHS-1E (LHS), LMS-1E (LMS), NUW-LHT-5M (NUW), and OPRH4W30 (OPR). Shared letters a, b and c indicate statistically significant groupings determined by ANOVA followed by Tukey's post-hoc. N=5 except NUW-LHT-5M where N=3.

The potato transcriptome response differs between LRSs

To better understand the molecular changes that led to the observed phenotypes, we analyzed the transcriptome of leaves harvested from plants grown on lunar mare simulants JSC1-A and LMS-1E, lunar highland simulants OPRH4W30 and LHS1-E, and the Adkins terrestrial soil. A total of 972,140,076 raw reads were obtained (Table S10). Differential gene expression analysis between Adkins and each LRS revealed varying numbers of differentially expressed genes (DEGs) (p-adjusted value < 0.05) (Figure 5, Table S11). Comparison between lunar highland simulant OPRH4W30 and Adkins produced the lowest number of DEGs at 124, while 958, 1397, and 360 genes were differentially expressed when comparing lunar mare simulants JSC1-A and LMS1-E and lunar highland simulant LHS1-E, respectively, with Adkins. The majority of DEGs in

response to lunar mare simulants JSC-1A and LMS-1E were unique to each of these

LRSs, whereas most DEGs in response to lunar highland simulant LHS-1E were

common with those found in response to lunar mare simulant LMS-1E. Gene set

enrichment (GSE) analysis of lunar mare simulants JSC1-A and LMS-1E and lunar

highland simulants LHS1-E and OPRH4W30 revealed enrichment of 6, 20, 6, and 7

gene ontology terms , respectively (p-adj < 0.05) (Figure 6, Table S12). Many affected

genes are related to various stress response and signaling pathways. Notably, all four

LRSs affected genes related to 'kinase activity' (GO:0016301). Four pathways were

enriched in LMS-1E, LHS-1E, and OPRH4W30: 'extracellular region' (GO:0005576),

'protein modification process' (GO:0036211), 'protein metabolic process' (GO:0019538),

and 'enzyme regulator activity' (GO:0030234). The pathways 'cell wall' (GO:0005618)

was enriched in both LMS-1E and OPRH4W30. LHS-1E, JSC-1A, and OPRH4W30

shared one enriched process: 'signal transduction' (GO:0007165). Thirteen pathways

were uniquely enriched by LMS-1E: 'DNA-binding transcription factor activity'

(GO:0003700), 'DNA binding' (GO:0003677), 'cytosol' (GO:0005829), 'nucleobase-

containing compound metabolic process' (GO:0006139), 'response to chemical'

(GO:0042221), 'catalytic activity' (GO:0003824), 'endoplasmic reticulum' (GO:0005783),

'nuclease activity' (GO:0004518), 'signaling receptor activity' (GO:0038023), 'thylakoid'

(GO:0009579), 'RNA binding' (GO:0003723), 'hydrolase activity' (GO:0016787) and

'membrane' (GO:0016020). Three pathways were uniquely enriched by JSC-1A:

'carbohydrate binding' (GO:0030246), 'molecular function' (GO:0003674), and

'response to external stimulus' (GO:0009605). LHS-1E and OPRH4W30 did not

uniquely enrich any pathway.

Figure 5. Venn diagram of differentially expressed genes ($p_{adj} < 0.05$) of plants grown in LHS-1E (LHS), LMS-1E (LMS), JSC-1A (JSC), and OPRH4W30 (OPR) compared with the terrestrial soil Adkins.

Figure 6. Gene Set Enrichment (GSE) analysis of differentially expressed genes between
Adkins soil treatment and LRS. LMS-1E (LMS), LHS-1E (LHS), JSC-1A (JSC), OPRH4W30
(OPR). Dotplots display the gene ratio (count of enriched genes/count of total genes in pathway)
of various GO terms based on comparison to Adkins control soil.

Metabolites and minerals profiles are unique to each LRS

We complemented transcriptome analysis with metabolite analysis in both leaf

and tuber tissues. Analysis of leaf metabolites by GC-MS revealed unique metabolic

profiles of plants grown in LRS as compared to Adkins soil (Figures 7, Table S13).

Compounds contributing to these differences include sucrose and fructose as well as

other nutrients such as citric acid (Figure 8). Analysis of tuber metabolites by GC-MS

revealed only minor changes between treatments (Figure 8, Table S14). In addition,

analysis of phenolic compounds and two major glycoalkaloids via HPLC-UV/VIS did not

reveal significant changes in 6 out of 7 compounds analyzed (Figure 9). Tyrosine

concentrations increased significantly in plants grown in lunar mare simulant JSC-1A

compared to Adkins, but did not increase significantly in lunar mare simulant LMS-1E

and lunar highland simulants LHS-1E and OPRH4W30. Heavy metal analysis of tubers

revealed increased accumulation of Cu and Zn in all LRS grown-tubers, though not

statistically significant in some cases. Increased accumulation of Al in tubers grown in

lunar highland simulants LHS-1E and OPRH4W30 was also observed (Table 2, Table

S15).

Figure 7. PCA biplot of GC-MS metabolite analysis of leaves collected at 39 DAP from potato

plants grown in Adkins soil (ADK) and four lunar regolith simulants. JSC-1A (JSC), LHS-1E

(LHS), LMS-1E (LMS), and OPRH4W30 (OPR).

Figure 8. PCA biplot of GC-MS metabolite analysis of tubers harvested at 59 DAP from potato plants grown in Adkins soil (ADK) and four lunar regolith simulants; JSC-1A (JSC), LHS-1E (LHS), LMS-1E (LMS), and OPRH4W30 (OPR).

Figure 9. Quantities of targeted compounds extracted from harvested tubers of potato plants grown in Adkins soil (ADK) and five lunar regolith simulants. JSC-1A (JSC), LHS-1E (LHS), LMS-1E (LMS), and OPRH4W30 (OPR). (A) ascorbic acid, (B) chlorogenic acid, (C) phenylalanine, (D) tryptophan, and (E) tyrosine, (F) chaconine, and (G) solanine. Shared letters indicate statistically significant groupings determined by ANOVA followed by Tukey's post-hoc.

Analysis of soils and tuber heavy metal content

Soil analysis revealed differences between LRSs and Adkins soil (Table 1 or

Table S16). All LRSs had higher moisture content, indicating reduced drainage

capability. Lunar mare simulants LMS1-E and JSC-1A and lunar highland simulant,

LHS1-E all experienced a reduction in pH from 7.5-7.72 range to ~6.5, whereas lunar

highland simulant OPRH4W30 remained ~7.5, compared to pre-growth samples (Table

S17). Electroconductivity decreased in lunar mare simulants LMS1-E and JSC-1A and

increased in lunar highland simulants LHS-1E and OPRH4W30 compared to pre-growth

samples. Elemental availability was also altered for some elements, such as Ca, K, Mg,

and Zn, post-growth. However, this is consistent with changes in pH, which determines

the solubility of many nutrients.

Discussion

The goal of this study was to assess the effects of LRS on potato growth and physiology and to characterize the underlying molecular changes. Because LRSs do not contain any nitrogen, we provided plants with this macronutrient along with other macro and micronutrients via fertilization with the slow-release fertilizer Osmocote. Growth was severely affected when potato plants were grown in 100% lunar simulant LMS-1. Symptoms included stunting and underdeveloped roots, which are typical of plants with excess copper [25,26]. Interestingly, leaf and tuber tissues of plants grown in 100% LMS-1E accumulated higher copper levels compared to Adkins control. The only notable source of copper was Osmocote, as copper is not present in LMS-1E [27]. Excess aluminum can also lead to similar phenotypes [28]. However, we did not observe any substantial differences in leaf and tuber Al concentrations between plants grown in 100% LMS-1E and Adkins. This may be due to the low solubility of aluminum at pH measured in the substrates used in this study, which were consistently above 5. Aluminum becomes an issue at pH below 5 that increases solubility [29]. The negative effect of 100% LMS-1E was alleviated by adding compost to the growth substrate. If copper toxicity is the reason for the stunting phenotype, the addition of organic matter would provide binding sites for copper [30], thereby alleviating hyperaccumulation. These types of interactions should therefore be considered when designing fertilizer management plans for growing crops in lunar regolith that had little to no biological amendments.

Potato plants grown on different lunar mare and lunar highland regolith simulants

amended with 5% compost were all negatively affected in canopy size and tuber yield

compared to those grown in Adkins. Therefore, LRS is stressful for potato plants, which

responded to this stressful environment by modifying the expression of genes involved

in photosynthesis, stress responses, signaling, and metabolism of terpenes and

flavonoids. We have several hypotheses to help explain these negative phenotypes.

First, we cannot rule out that copper and possibly aluminum toxicity was completely

alleviated by adding compost. Indeed, heavy metals analysis in tubers showed that

copper concentrations were significantly higher in lunar mare highland JSC-1A

compared to Adkins, and aluminum concentrations were higher in lunar highland

simulants LHS-1E and OPRH4W30, indicating higher uptake of these heavy metals in

LRS than in Adkins. Copper is also moderately increased in all other LRS, though not

statistically different according to ANOVA. Unfortunately, we were unable to confirm

these results in leaf tissues as plants had senesced at harvest, preventing us from

collecting leaf samples. It remains unclear how aluminum could accumulate in tubers

grown on lunar highland simulants as pH was above 5. Second, the presence of glassy

agglutinates in lunar highland simulants OPRH4W30 and NUW-LHT-5M may be

particularly problematic for plant growth [31,32]. Lunar mare simulant JSC-1A, despite

lacking in agglutinates, does contain a high glass fraction [33] and caused similar issues

as OPRH4W30. These sharp micro-particles may cause abrasions on root/stolon/tuber

epidermis. Given the findings of the transcriptomic analysis, these microabrasions are

likely being perceived by the plant as herbivory by nematodes and other soil-borne

insects (Figure 6). Third, high levels of compaction, cementation, small particle size and

higher density of lunar highland simulant NUW-LHT-5M and lunar mare simulant JSC-

1A after watering might also have negatively affected plant growth and development.

This is consistent with the higher moisture content recorded in LRSs vs the Adkins soil,

and is likely a result of high particle density and small particle size. Finally, it is likely that

all together these factors caused the stunting observed.

Growth in LRS did not significantly alter potato tuber metabolite profiles,

indicating negligible changes to nutritional content. Additionally, stress induced by the

LRS did not cause any changes in glycoalkaloid levels. These findings support that

tuber nutritional quality is maintained in LRS-grown tubers, despite lower potential

yields. Should this limitation prove true with larger-scale growth of mature plants,

agricultural modules may need to be larger than currently predicted to produce required

quantities of edible matter. Heavy metal analysis revealed mild accumulation of Zn and

Cu in tubers grown in LRS. In addition, plants grown in highland-type LRS were found to

accumulate aluminum, which is not of immediate concern for human consumption

purposes, but could contribute to long-term negative impacts on plant production.

Conclusions

From these observations, we conclude that LRS are clearly stressful on potato

plants, given the reduction in overall and edible biomass, and the changes to stress

response related gene expression. However, significant changes were not found in

metabolite profiles of leaves or tubers, indicating that nutrition was not impacted by

growth in LRS. Finally, addition of vermicompost to LMS-1E improves growth of potato.

This highlights that remediation of lunar regolith via the addition of biomatter is possible,

which bodes well for later generations of plant growth in lunar regolith as biomatter

builds up over the course of the first generations, though the lack of organic matter

presents an issue for early stages of bioregenerative life support development.

Declarations

Ethics approval and consent to participate

525   Not applicable

Consent for publication

527   Not applicable

Availability of data and materials

529   The raw sequence reads have been deposited in the Open Science Data Repository

530   under doi 10.26030/w56j-2d69.

Competing interests

532   The authors declare that they have no competing interests.

Funding

534   This research was supported by funding from NASA (Grant# 80NSSC24K0755). Anika

535   Loeffler and Sydney Campbell were supported by funding from the Oregon State

536   University URSA Engage program, and Medora Knudson was supported by funding

537 from the Oregon State University College of Agricultural Sciences Beginning

538 Researcher Program.

Authors' contributions

DH: design of the work, acquisition, analysis, and interpretation of data, wrote first draft

of the manuscript; AL: data acquisition; MK: data acquisition; SC: data acquisition; PJ:

funding acquisition, design of the work, reviewed and edited the manuscript; JCA: data

acquisition, reviewed and edited the manuscript; AG: conception, funding acquisition,

design of the work, analysis and interpretation of data, reviewed and edited the

manuscript. All authors have approved the submitted manuscript.

Acknowledgements

547 We thank Matthew Barrow at CSS Farms for providing minitubers of Modoc.

References

[1]   D.W. Ming, D.L. Henninger, Use of lunar regolith as a substrate for plant growth,

Adv. Sp. Res. 14 (1994) 435–443. https://doi.org/10.1016/0273-1177(94)90333-6.

[2]   L.G. Duri, A.G. Caporale, Y. Rouphael, S. Vingiani, M. Palladino, et al., The

Potential for Lunar and Martian Regolith Simulants to Sustain Plant Growth: A

Multidisciplinary Overview, Front. Astron. Sp. Sci. 8 (2022).

https://doi.org/10.3389/fspas.2021.747821.

[3]   L.E. Fackrell, S. Humphrey, R. Loureiro, A.G. Palmer, J. Long-Fox, Overview and

recommendations for research on plants and microbes in regolith-based

agriculture, Npj Sustain. Agric. 2 (2024) 15. https://doi.org/10.1038/s44264-024-

00013-5.

[4]   A. Madison, Z. Landsman, J. Long-Fox, A. Metke, K. Krol, et al., Lunar Dust

Simulants and Their Applications, Earth Sp. 2022 Sp. Explor. Util. Eng. Constr.

Extrem. Environ. - Sel. Pap. from 18th Bienn. Int. Conf. Eng. Sci. Constr. Oper.

Challenging Environ. (2023) 86–94. https://doi.org/10.1061/9780784484470.008.

[5]   S.R. Taylor, The Moon, Encycl. Sol. Syst. (2007) 227–250.

https://doi.org/10.1016/B978-012088589-3/50016-5.

[6]   E. Hill, M.J. Mellin, B. Deane, Y. Liu, L.A. Taylor, Apollo sample 70051 and high-

and low-Ti lunar soil simulants MLS-1A and JSC-1A: Implications for future lunar

exploration, J. Geophys. Res. Planets 112 (2007).

https://doi.org/10.1029/2006JE002767.

[7]   A. Madison, Z. Landsman, J. Long-Fox, A. Metke, K. Krol, et al., Lunar Dust

Simulants and Their Applications, in: Earth Sp. 2022, American Society of Civil

Engineers, Reston, VA, 2023: pp. 86–94.

https://doi.org/10.1061/9780784484470.008.

[8]   C.H. Walkinshaw, P.H. Johnson, Analysis of Vegetable Seedlings Grown in

Contact with Apollo 14 Lunar Surface Fines1, HortScience 6 (1971) 532–535.

https://doi.org/10.21273/HORTSCI.6.6.532.

[9]   A.-L. Paul, S.M. Elardo, R. Ferl, Plants grown in Apollo lunar regolith present

stress-associated transcriptomes that inform prospects for lunar exploration,

Commun. Biol. 5 (2022) 382. https://doi.org/10.1038/s42003-022-03334-8.

[10] B.B. Barcenilla, I. Kundel, E. Hall, N. Hilty, P. Ulianich, et al., Telomere dynamics

and oxidative stress in Arabidopsis grown in lunar regolith simulant, Front. Plant

Sci. 15 (2024). https://doi.org/10.3389/fpls.2024.1351613.

[11] A.G. Caporale, R. Paradiso, G. Liuzzi, M. Palladino, C. Amitrano, et al., Green

compost amendment improves potato plant performance on Mars regolith

simulant as substrate for cultivation in space, Plant Soil 486 (2023) 217–233.

https://doi.org/10.1007/s11104-022-05860-0.

[12] C. Bvindi, K. Howe, Y. Wang, R.T. Mullen, C.J. Rogan, et al., Potato Non-Specific

Lipid Transfer Protein StnsLTPI.33 Is Associated with the Production of Reactive

Oxygen Species, Plant Growth, and Susceptibility to Alternaria solani, Plants 12

(2023) 3129. https://doi.org/10.3390/plants12173129.

[13] T. Davies, The new automated mass spectrometry deconvolution and

identification system (AMDIS), Spectrosc. Eur 10 (1998) 24–27.

[14] T. Kind, G. Wohlgemuth, D.Y. Lee, Y. Lu, M. Palazoglu, et al., FiehnLib: Mass

Spectral and Retention Index Libraries for Metabolomics Based on Quadrupole

and Time-of-Flight Gas Chromatography/Mass Spectrometry, Anal. Chem. 81

(2009) 10038–10048. https://doi.org/10.1021/ac9019522.

[15] A. Goyer, M. Picard, H.A. Hellmann, S.L. Mooney, Effect of low-temperature

storage on the content of folate, vitamin B 6 , ascorbic acid, chlorogenic acid,

tyrosine, and phenylalanine in potatoes, J. Sci. Food Agric. 99 (2019) 4842–4848.

https://doi.org/10.1002/jsfa.9750.

[16] S. Torabian, A. Goyer, R. Qin, K. Jones, C. Flagg, et al., Effect of potassium

fertilizer, genetic makeup, and environment on the potato tuber nutrition, J. Agric.

Food Res. 19 (2025) 101645. https://doi.org/10.1016/j.jafr.2025.101645.

[17] T.W. Berrian, M.L. Fabian, C.J. Rogan, J.C. Anderson, C.R. Clarke, et al.,

Investigation of the Effectiveness and Molecular Mechanisms of Thiamin Priming

to Control Early Blight Disease in Potato, Phytopathology® 115 (2025) 234–246.

https://doi.org/10.1094/PHYTO-09-24-0277-R.

[18] J.P. Hamilton, J. Brose, C.R. Buell, SpudDB: a database for accessing potato

genomic data, Genetics 229 (2025). https://doi.org/10.1093/genetics/iyae205.

[19] D. Kim, J.M. Paggi, C. Park, C. Bennett, S.L. Salzberg, Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype, Nat. Biotechnol. 37

(2019) 907–915. https://doi.org/10.1038/s41587-019-0201-4.

[20] H. Li, B. Handsaker, A. Wysoker, T. Fennell, J. Ruan, et al., The Sequence

Alignment/Map format and SAMtools, Bioinformatics 25 (2009) 2078–2079.

https://doi.org/10.1093/bioinformatics/btp352.

[21] M. Pertea, G.M. Pertea, C.M. Antonescu, T.-C. Chang, J.T. Mendell, et al.,

StringTie enables improved reconstruction of a transcriptome from RNA-seq

reads, Nat. Biotechnol. 33 (2015) 290–295. https://doi.org/10.1038/nbt.3122.

[22] M.I. Love, W. Huber, S. Anders, Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2, Genome Biol. 15 (2014) 550.

https://doi.org/10.1186/s13059-014-0550-8.

[23] S. Xu, E. Hu, Y. Cai, Z. Xie, X. Luo, et al., Using clusterProfiler to characterize

multiomics data, Nat. Protoc. 19 (2024) 3292–3320.

https://doi.org/10.1038/s41596-024-01020-z.

[24] G. Yu, Thirteen years of clusterProfiler, Innov. 5 (2024) 100722.

https://doi.org/10.1016/j.xinn.2024.100722.

[25] A. Moore, M. Satterwhite, J. Ippolito, Soil copper thresholds for potato production,

West. Nutr. Manag. Conf. 10 (2013) 152–155.

[26] A.R. Mir, J. Pichtel, S. Hayat, Copper: uptake, toxicity and tolerance in plants and

management of Cu-contaminated soil, BioMetals 34 (2021) 737–759.

https://doi.org/10.1007/s10534-021-00306-z.

[27] J.M. Long-Fox, Z.A. Landsman, P.B. Easter, C.A. Millwater, D.T. Britt,

Geomechanical properties of lunar regolith simulants LHS-1 and LMS-1, Adv. Sp.

Res. 71 (2023) 5400–5412. https://doi.org/10.1016/j.asr.2023.02.034.

[28] R. Ofoe, R.H. Thomas, S.K. Asiedu, G. Wang-Pruski, B. Fofana, et al., Aluminum

in plant: Benefits, toxicity and tolerance mechanisms., Front. Plant Sci. 13 (2022)

1085998. https://doi.org/10.3389/fpls.2022.1085998.

[29] J.L. Domingo, ALUMINUM (ALUMINIUM) | Toxicology, in: Encycl. Food Sci. Nutr.,

Elsevier, 2003: pp. 160–166. https://doi.org/10.1016/B0-12-227055-X/00035-3.

[30] E.J.M. Temminghoff, S.E.A.T.M. Van der Zee, F.A.M. de Haan, Copper Mobility in

a Copper-Contaminated Sandy Soil as Affected by pH and Solid and Dissolved

Organic Matter, Environ. Sci. Technol. 31 (1997) 1109–1115.

https://doi.org/10.1021/es9606236.

[31]   M.C. Roth, V.G. Roux, L. Roux, MECHANICAL PROPERTIES OF LUNAR

REGOLITH SIMULANTS, 2025.

[32]   D.L. Rickman, P. Douglas Archer, R.N. Kovtun, M. Barmatz, B. Dotson, et al.,

Characterization of NUW-LHT-5M, A Lunar Highland Simulant, 2024.

http://www.sti.nasa.gov.

[33]   S. Sen, D. Butts, C.S. Ray, G.B. Thompson, R.A. Morris, et al., Production of high

fidelity lunar agglutinate simulant, Adv. Sp. Res. 47 (2011) 1912–1921.

https://doi.org/10.1016/j.asr.2011.02.005.

Table 1. Post-growth soil characteristics of soil substrates amended with 5% v/v vermicompost. Data are means ± SE (N =4). Superscript letters indicate statistically similar groupings according to ANOVA followed by Tukey's post-hoc. Analysis was performed in R.

	Moisture (%)	pH	EC (dS m^{-1})	Elemental quantities (mg kg^{-1})									Microbial respiration (µg CO$_2$-C g^{-1} dry soil day^{-1})		Organic matter (%)
				Al	Ca	Cu	Fe	K	Mg	Mn	P	Zn	24 hr	96 hr	
ADK	7.70 ± 0.90^b	6.38 ± 0.10^b	0.51 ± 0.09^a	464.24 ± 9.64^e	1578.67 ± 29.92^b	2.70 ± 0.29^{bc}	115.76 ± 5.06^e	221.24 ± 13.92^{ab}	229.49 ±6.03^c	31.88 ± 0.54^c	56.83 ± 4.97^b	2.00 ± 0.29^b	5.56 ± 0.25^a	2.26 ± 0.23^b	1.13 ± 0.03^a
JSC	11.63 ± 2.07^{ab}	6.69 ± 0.16^b	0.85 ± 0.10^a	1161.58 ± 30.19^b	1666.09 ± 45.28^b	2.70 ± 0.15^{bc}	1526.69 ± 37.23^b	268.39 ± 22.28^a	567.91 ± 15.34^a	38.59 ± 1.01^b	29.21 ± 2.02^c	4.50 ± 0.35^b	5.14 ± 0.11^a	2.45 ± 0.11^b	0.52 ± 0.03^c
LHS	12.60 ± 1.39^{ab}	6.52 ± 0.04^b	0.74 ± 0.05^a	866.32 ± 11.59^c	1567.52 ± 33.05^b	1.66 ± 0.20^c	417 ± 12.34^d	177.23 ± 12.13^{bc}	135.91 ± 5.06^d	10.42 ± 0.23^e	66.88 ± 3.25^{ab}	4.21 ± 0.30^b	6.30 ± 0.38^a	3.27 ± 0.27^{ab}	0.58 ± 0.03^c
LMS	17.06 ± 3.14^{ab}	6.46 ± 0.11^b	0.94 ± 0.14^a	664.90 ± 30.36^d	1210.03 ± 60.24^c	3.34 ± 0.24^{ab}	1066.22 ± 38.93^c	203.57 ± 20.36^{abc}	368.12 ± 20.74^b	24.32 ± 1.01^d	83.94 ± 4.55^b	4.83 ± 0.76^b	6.37 ± 0.63^a	3.49 ± 0.25^a	0.61 ± 0.03^c
OPR	19.75 ± 0.53^a	7.53 ± 0.05^a	0.95 ± 0.07^a	1337.32 ± 13.10^a	7035.16 ± 87.10^a	4.09 ± 0.13^a	1924.68 ± 26.19^a	138.20 ± 8.83^c	203.55 ± 3.27^c	43.36 ± 0.70^a	23.18 ± 1.27^c	10.05 ± 1.70^a	6.10 ± 0.43^a	2.62 ± 0.26^{ab}	0.85 ± 0.03^b

Table 2. Heavy metal content of tubers grown in various LRS with 5% v/v vermicompost, determined by ICP-OES. Letters indicate statistically similar groupings determined by one-way ANOVA and Tukey's post-hoc, performed in R. N=4. As, Cd, Co, Cr, Ni, Pb were also analyzed but found below quantifiable levels in all samples.

Soil type	Heavy metal content (mg kg^{-1} dry weight)		
	Al	Cu	Zn
ADK	55.9 ± 3.03^b	2.8 ± 0.32^b	6.6 ± 0.31^b
JSC	59.3 ± 6.42^b	6.0 ± 0.60^a	12.3 ± 1.08^a
LHS	144.6 ± 11.60^a	4.7 ± 1.08^{ab}	9.5 ± 0.63^{ab}
LMS	50.8 ± 3.74^b	5.1 ± 0.00^{ab}	11.4 ± 0.51^a
OPR	185.1 ± 26.88^a	6.0 ± 0.32^{ab}	12.0 ± 0.64^a