# Calcium signals natural transformation in *Acinetobacter baumannii*

Jason Baby Chirakadavil[1]*, Kelly Goldlust[1]*, Ludovic Poiré[1], Nicolas Gaudin[1], Alexandre Chassard[1], Adrien Camilli[1,2], Manon Bouvier[1], Maria-Halima Laaberki[1,2#@], Xavier Charpentier[1#@]

1. CIRI, Center for Integrative Research in Infectious diseases and Immunology, Inserm, U1111, Université Claude Bernard Lyon 1, CNRS, UMR5308, École Normale Supérieure de Lyon, Univ Lyon, 69007, Lyon, France
2. Université de Lyon, VetAgro Sup, 69280 Marcy l'Etoile, France.

* These authors contributed equally
# Co-senior authors
@ Correspondence to Maria-Halima Laaberki, maria-halima.laaberki@vetagro-sup.fr
@ Correspondence to Xavier Charpentier, xavier.charpentier@univ-lyon1.fr

**Running Head:** Calcium signals natural transformation

| Name | email | ORCID |
|---|---|---|
| Jason Baby Chirakadavil | jason.chirakadavil@gmail.com | 0000-0001-8596-018X |
| Kelly Goldlust | kelly.goldlust@inserm.fr | 0000-0003-4372-6840 |
| Ludovic Poiré | ludovic.poire@univ-lyon1.fr | 0009-0008-7763-389X |
| Alexandre Chassard | alexandre.chassard8@gmail.com | |
| Adrien Camilli | adrien.camilli@hotmail.fr | |
| Manon Bouvier | manon.bouvier@univ-lyon1.fr | |
| Maria-Halima Laaberki | maria-halima.laaberki@vetagro-sup.fr | 0000-0001-5796-5995 |
| Xavier Charpentier | xavier.charpentier@univ-lyon1.fr | 0000-0001-9039-8021 |

# Abstract

The acquisition of resistance to antibiotics in the opportunistic pathogen *Acinetobacter baumannii* may be linked to its capacity to undergo natural transformation. This mode of horizontal gene transfer relies on the import of extracellular DNA and its chromosomal integration by homologous recombination. Type IV pilus activity initiates the capture of extracellular DNA, which is then transported to the cytoplasm for recombination in the chromosome. While most *Acinetobacter baumannii* strains are transformable, the conditions allowing the expression of type IV pilus and other transformation genes remain largely unexplored. By investigating transformation-permissive conditions, we uncovered that calcium is a potent inducer of natural transformation. Type IV pilus genes and other transformation-specific genes (*comEA*, *dprA*) are upregulated by submillimolar concentrations of calcium ions, in a growth phase-dependent manner. In contrast, sodium chloride represses expression of *pilA*, counteracting the calcium-dependent induction, explaining the reported absence of transformation in NaCl-containing medium (such as LB). Independently of transcriptional induction, calcium ions also directly bind the type IV

pilus through the calcium-dependent adhesin PilY1. Our data support a model in which calcium strengthens the interaction of PilY1 with the minor pilin complex, increasing pilus dynamics and subsequent pilus-dependent DNA capture. Hence, calcium signals natural transformation through both transcriptional and structural activation of type IV pilus. In addition to providing new insights into the regulation of natural transformation in *A. baumannii* this work led us to establish a protocol for genetic engineering of *A. baumannii* by natural transformation.

## Importance

*Acinetobacter baumannii* is a nosocomial pathogen considered a critical research priority due to its resistance to last resort antibiotics. Understanding how *A. baumannii* evolves and acquires resistance to antibiotics is thus of prime importance. This species is capable of natural transformation, a means to acquire and spread genetic information, including antibiotic resistance genes. However, the conditions under which this process is active in this species remain elusive. We identify calcium ions as potent inducers of natural transformation and propose a model of the signaling of natural transformation by calcium ions. This opens the way for further investigations into the contribution of natural transformation to acquisition of antibiotic resistance. In addition, it provides an efficient way to genetically manipulate most *A. baumannii* strains.

# Introduction

Natural transformation is one of the main mechanisms of horizontal gene transfer driving the evolution of bacterial genomes (1). Over 80 bacterial species have been experimentally demonstrated to be capable of natural transformation (2). This ever-growing list includes *Acinetobacter baumannii*, a critical priority for research and development of new therapeutic strategies (3) due to its resistance to last resort antibiotics. Natural transformation likely plays a major role in the evolution of *A. baumannii* genomes, constituting a route of gene exchange promoting recombination events and acquisition of antibiotic resistance (4–6). Although natural transformation in *A. baumannii* appears to play a limited role in overall gene acquisition, at least 15% of antibiotic resistance genes had been acquired through this route (5). Indeed, even large genetic elements, such as AbaR resistance islands, could be acquired by natural transformation during the co-culture of *A. baumannii* isolates (7). Yet, it is the subsequent partial inhibition of natural transformation by these resistance islands that favors the prevalence of antibiotic resistance genes (8). Most *A. baumannii* are naturally transformable, but at varying levels (9–12). Such variations in natural transformation could be ascribed to genomic elements like restriction-modification systems that can limit transformation (11) and prophages inhibiting transformability (10). It remained possible that not all strains respond to the same stimuli for expression of transformation, yet little is known about the extracellular conditions that impact the expression of natural transformation in *A. baumannii*.

Natural transformation relies on the concerted expression of genes involved in DNA capture, internalization and recombination in the chromosome. DNA capture is mediated by extracellular filaments known as type IV pili (13). The Type IV pilus is a dynamic nanomachine that undergoes cycles of extension and retraction energized by at least two respective ATPase, PilB and PilT (14). Current models of Type IV pilus assembly propose that a complex of minor pilins (PilX, PilW, FimT, FimU, PilV) primes the assembly of the major pilin PilA to form a thin flexible filament (15–17). Pilus extension, capture of extracellular DNA, followed by retraction, bring DNA to the cytoplasmic space where it is bound by the ComEA receptor (13). Then DNA is converted into single-stranded DNA and translocated to the cytoplasm through the ComEC channel (18) and rapidly protected by DprA which recruits RecA for homologous recombination (19). Integration of DNA,

facilitated by the ComM helicase and YraN nuclease (20), followed by replication, results in genetic transformation (21, 22).

The conditions and genetic control allowing the concerted expression of the genes involved in natural transformation – often referred to as competence – vary between species (2). For example, *Legionella pneumophila* becomes transiently competent for natural transformation in the early stationary phase when the post-transcriptional repression of transformation genes (Type IV minor pilins, comEA, comFC) by the RocC/RocR chaperone sRNA-pair is relieved (23). In *Vibrio cholerae*, chitin induces a transformation regulon, including type IV pilus genes, under control of the transcriptional activator TfoX (24). In the cyanobacterium *Synechococcus elongatus*, natural transformation is strictly regulated by circadian clock with Type IV pilus machinery produced in the morning, and natural transformation occurring at the beginning of the dark phase (evening phase) when the other genes required for transformation (including *dprA*) are expressed (25). In *Acinetobacter baumannii*, the inducing signals are undefined. Transformation protocols typically involve growth on agar or agarose-based solid medium compatible with Type IV pilus-mediated motility (7, 9, 26–30). Although the signals remained elusive, natural transformation was found to occur at the beginning of the growth phase and linked to the expression of type IV pilus genes (28).

Here, we explored the regulation of natural transformation in *Acinetobacter baumannii* by experimental conditions. Using a selection-free luminescence-based assay for natural transformation we screened a wide range of conditions, ultimately identifying a trigger of natural transformation. Genetics and molecular modeling indicate that calcium ions induce transformation by both signaling expression of the Type IV pilus and stabilization of the pilus assembly. The identified conditions allow the natural transformation of *A. baumannii* with high efficiency in liquid medium, opening the way for further investigations into the dynamics of transformation but also facilitating genetic engineering of this opportunistic pathogen.

## Results

**Mechanical sensing is not involved in induction of natural transformation**

Natural transformation of *A. baumannii* is routinely done on medium solidified with agar or agarose (7, 9, 26–30). However, we and others have experienced inconsistent results that could be linked to the source of the agar or agarose used to solidify the medium. To formally establish this, we tested five references of agarose powders from four suppliers.

Used at the same concentration (15 g/L) all agarose powders have identical solidifying
power. We  tested natural transformation of strain AB5075 using the NanoLuc (Nluc)
transformation assay previously reported (10). Briefly, a suspension of AB5075 is spotted
on the agarose plated together with a plasmid DNA that cannot replicate in *A. baumannii*
but that carries the *nluc* gene flanked by 2 kb-long regions homologous to the
chromosome. Import and recombination of this sequence leads to expression of Nluc in
the transformants which can be detected by measuring luminescence with the Nluc
substrate furimazine. We previously established that the luminescence signal is linearly
correlated with the number of transformants over eight orders of magnitude ($R^2$=0.99).
Using this assay we found that only two of the tested agarose could generate
transformants with RFU up to $10^6$ (Fig. 1A), corresponding to transformation frequencies of
$10^{-3}$-$10^{-4}$. Three other agarose gave luminescence signal barely above the value of the
control without DNA ($2 \times 10^2$ RFU). This suggested that either the perception of a solid
surface is required to induce transformation but some agarose contain an inhibitor, or the
perception of a solid surface is dispensable for induction but some agarose contain an
inducer of transformation. We tested this latter hypothesis by extracting any water-soluble
compound from the unmelt agarose powder of the Euromedex D3 agarose that is
permissive for transformation. A D3 agarose soluble extract (ASE) in liquid tryptone
medium strongly induces transformation of strain AB5075, up to $10^7$ RLU (transformation
frequency of $10^{-2}$), a 10,000-fold increase relative to a no-agarose liquid tryptone medium
(Fig. 1B). As expected, no transformation was observed in an AB5075 *comEC* mutant.
This shows that the D3 ASE contains a water-soluble potent inducer of natural
transformation, aguing against a mechanical triggering of natural transformation.

**A soluble molecule induces a growth phase-dependent regulon of natural**
**transformation**
Natural transformation in *A. baumannii* was previously found regulated by the growth
phase, correlating with the expression of Type IV pilus genes (28). To determine how the
ASE stimulates transformation, we tested the transformability of bacteria over the growth
curve. We confirmed previous results, showing that transformation peaks at 3 hours post-
inoculation, corresponding to the exponential growth phase (Fig. 1C). Transformation
stimulated by ASE followed the same kinetic, peaking at 3 hours, with a wider
transformation window but also falling under the detection limit in the stationary phase at 7
hours post-inoculation (Fig. 1C). We then tested if increased transformation was tied to
type IV pilus induction. To do so, we compared the effect of ASE from inducing (D3) and

non-inducing agarose (Roth) on the global transcriptome using RNAseq profiling at 3 hours post-inoculation. Each of the non-inducing (Roth) (Fig. 2D) and inducing (D3) agarose (Fig. 2E) downregulate specific set of genes (blue dots). Yet, they also downregulate the expression of the same genes (green dots) including the phenylacetic acid catabolic pathway (*paaA*, *paaB*, *paaC*, *paaX*, *paaK*, *paaZ*). This clearly shows that adding agarose to culture medium is not without consequences on gene expression. Most importantly, while the Roth ASE induces only a dozen genes, the D3 ASE induces the upregulation of nearly all type IV pilus genes, several of which are organized in operons (*pilMONPQ*, *pilA*, *fimU-pilVWXY1-pilE1-pilE2*, *pilGHIL-chpA*, *pilBCD*), but also transformation-specific genes such as *yraN*, *dprA* or *comEA* (Fig. 1E and Fig. 1F). Notably, all of these genes are downregulated in the stationary phase (D3 ASE, 7 hours) relative to the exponential phase (D3 ASE, 3 hours) (Fig. 1F). Hence, a water-soluble compound present in agarose preparations induces a growth-phase regulated transformation regulon.

**Figure 1. A water-soluble molecule induces a growth phase-dependent regulon of natural transformation.**
A. Efficiencies of natural transformation of *A. baumannii* strain AB5075 in tryptone medium solidified by different agaroses. Transformation is determined by incubating with a DNA fragment containing the Nluc gene flanked by 2 kb-long sequence of homology with the chromosome on each side. Transformants are

detected by luminescence units (LU) produced when adding the substrate furimazine, and expressed relative to the optical density of the culture (RLU). The detection limit, and background luminescence is $10^2$.

B. Efficiencies of natural transformation of strain AB5075 in liquid tryptone medium or tryptone agarose soluble extract (ASE) using the powder of the Euromedex D3 agarose. The method is identical as in panel A.

C. Growth and natural transformation of strain AB5075 in tryptone medium or tryptone agarose soluble extract (ASE) using the powder of the Euromedex D3 agarose. Growth of the culture is monitored by reading the absorbance at 600 nm ($Abs_{600}$). Transformation efficiency is tested at 30 minutes interval by addition of transforming DNA for 30 minutes followed by the addition of DNAse I. The culture is then further incubated. All samples were analyzed for luminescence signal, indicating effective transformation, at the end of the kinetic.

D and E. RNAseq analysis of strain AB5075 grown for 3 hours in tryptone and tryptone using the non-transforming Roth agarose (D) or the transforming Euromedex D3 agarose (E). In both panel D and E, differentially-expressed genes ($log_2FC<-1$ or $log_2FC>1$ and P-value<0.05) are shown in blue. Genes down-regulated in both conditions are highlighted in green. Genes specifically upregulated by each condition are highlighted in deep red.

F. Variation in expression of selected genes, required for natural transformation, in Roth ASE, D3 ASE and in D3 ASE at 8 hours (stationary phase) compared to D3 ASE at 3 hours (exponential phase). In AB5075, the *comM* gene is disrupted by the insertion of an AbaR element, forming two pseudogenes (*comM'* and '*comM*).

## Screen identifies Ca and Mg as inducers and NaCl as repressor

RNAseq profiling revealed no upregulated genes involved in metabolism, suggesting that the transformation inducer is not metabolizable. Chemical preparations are often contaminated by ions and salt. We thus screened the Biolog PM-PM5 plates containing four undisclosed concentrations of ions for induction of natural transformation. This revealed strong induction by $CaCl_2$ and to a lesser extent by $MgCl_2$ (Fig. 2A). Addition of $CaCl_2$ or $MgCl_2$ to the tryptone medium strongly stimulated natural transformation, in a dose-dependent manner (Fig. 2B). $CaCl_2$ is a stronger inducer, matching the induction by the D3 ASE (Fig. 2B), yet $CaCl_2$ and $MgCl_2$ showed the same dose-response effect, with maximal induction at 0.68 mM, suggesting the same mechanism of action. Indeed, the effects of $CaCl_2$ and $MgCl_2$ are not additive (Fig. 2C). Although $CaCl_2$ is a stronger inducer than $MgCl_2$, $MgCl_2$ shows stimulation of growth like observed for D3 ASE in the stationary phase (Fig. 2D and Fig. 1C). This suggested that D3 ASE contained $Ca^{2+}$ ions stimulating transformation as well as $Mg^{2+}$ ions stimulating growth. To test this hypothesis, we evaluated the effect on growth of adding the $Ca^{2+}/Mg^{2+}$ chelator EDTA to the medium. EDTA at 0.1 mM abolished growth indicating that the tryptone medium readily contains trace amounts of $Ca^{2+}$ (less than 0.1 mM, given the 1:1 stoichiometry of EDTA:cation). Adding $CaCl_2$ at 0.68 mM did not change the growth yield (same final absorbance), but growth is expectedly inhibited at EDTA concentration of at least 0.8 mM. D3 ASE improved growth yield, identically to adding $MgCl_2$ at 0.68 mM, and for both an EDTA concentration of 0.8 mM is needed to prevent growth. Thus, given the increased growth of D3 ASE and its effect on transformation similar to that of adding $CaCl_2$, we conclude that D3 ASE

contains both $Ca^{2+}$ and $Mg^{2+}$ ions, in undetermined concentrations, but likely greater than 0.68 mM.

Screening for the effect of ions on transformation, we also observed reduced transformation with increasing amounts of NaCl (Fig. 2A). This was confirmed in tryptone medium as well as when tryptone supplemented with $CaCl_2$ (Fig. 2E). NaCl at concentration greater than 0.07% inhibits transformation and completely abolishes it at 1%, which is the typical concentration of the commonly used lysogeny broth (LB medium). This is consistent with previous results obtained with strain A118 (29). In LB medium, $CaCl_2$ was previously reported to only moderately induce transformation in strain A118 (31). In this strain, we confirmed that $CaCl_2$ has a modest effect in standard LB, and also found no effect of $MgCl_2$ (Fig. 2F). The results suggest that LB might readily contain enough $Mg^{2+}$ and a sub-optimal amount of $Ca^{2+}$ to induce transformation but that the induction is counteracted by inhibition by NaCl. Consistent with this hypothesis, LB prepared without NaCl (LB no salt) produces a 500-fold increase in transformation, with a further increase when supplemented with $CaCl_2$, while $MgCl_2$ had no additional effect (Fig. 2E).

Having identified the optimal conditions for natural transformation, we tested these on a panel of strains, including commonly used laboratory strains. Transformation medium ($CaCl_2$ 0.68 mM, no NaCl) improved transformation of most strains, including CNRAB1 and 40288, which were non-transformable in the absence of supplementation (Fig. 2G). Addition of $CaCl_2$ also further increased transformation of the highly transformable strain A118, but did not change the transformation of the M2, 29D2 and 31D1 strains which readily showed high transformation levels in the absence of $CaCl_2$ supplementation.

In conclusion, $CaCl_2$ and to a lesser extent $MgCl_2$, induce natural transformation. Both divalent cations are the likely contaminants contained in the transformation-inducing agarose. Adding $CaCl_2$ to a tryptone medium without sodium chloride allows *A. baumannii* to undergo natural transformation in liquid medium, further demonstrating that surface sensing is dispensable for this type IV pilus-mediated process.

**Figure 2. Induction of natural transformation by calcium and magnesium is antagonized by sodium chloride**

A. Natural transformation of strain AB5075 in tryptone medium supplemented with the dried content of the Biolog PM-PM5 plate.

B. Natural transformation in tryptone medium supplemented with increasing concentration of CaCl$_2$ or MgCl$_2$.

C. Natural transformation in tryptone medium supplemented with CaCl$_2$ (0.68 mM) or MgCl$_2$ (0.68 mM) or both. Bars represent the average of technical replicates.

D. Effect on growth of EDTA on tryptone medium and D3 ASE, measured at 16 hours post-inoculation by absorbance reading at 600 nm. Data are presented as average values of three independent experiments.

E. Natural transformation in tryptone medium and tryptone supplemented with CaCl$_2$ (0.68 mM) at an increasing concentration of NaCl. Bars represent the geometric average of three independent experiments.

F. Effect of CaCl$_2$ and MgCl$_2$ on natural transformation of A118 in LB and LB no salt. Bars represent average of five technical replicates.

G. Natural transformation of a panel of *A. baumannii* and *A. nosocomialis* (M2) strains in tryptone medium alone or supplemented with CaCl$_2$ (0.68 mM). Bars represent the geometric average of three independent experiments.

**Ca$^{2+}$ and Mg$^{2+}$ are equivalent in inducing expression of transformation genes, independently of T4P activity**

In order to understand the mechanism by which CaCl$_2$ and MgCl$_2$ induce transformation, we analyzed the transcriptome of AB5075 exposed to CaCl$_2$ or MgCl$_2$ in the exponential growth phase. Consistent with D3 ASE containing calcium/magnesium salts, CaCl$_2$ and MgCl$_2$ induce the upregulation of the type IV pilus genes and other natural transformation genes (*comEA*, *dprA*) (Fig. 3A). The transcriptional responses to CaCl$_2$ and MgCl$_2$ are strikingly similar, indicating that the two metal ions are sensed by a common signaling pathway. With the *pilA* gene being upregulated and a central component of type IV pilus regulation and activity, we followed its expression using a transcriptional fusion of the full *pilA* gene to *gfp* (*pilA::gfp*) in a Δ*pilA* mutant. In agreement with the RNAseq data, *pilA* is induced to the same extent by CaCl$_2$ and MgCl$_2$ and is expressed during the exponential phase and then repressed in the stationary phase (Fig. 3B). The inhibition of transformation by NaCl suggested that it counteracts the Ca$^{2+}$-induced expression of the type IV pilus. Indeed, in the presence of calcium, *pilA* expression is repressed by increasing concentration of NaCl, indicating that inhibition of natural transformation is the consequence of a transcriptional response to NaCl (Fig. 3C).

The coordinated expression of the type IV pilus and the genes required for DNA import/recombination is indicative of a mechanism that equally senses Ca$^{2+}$ and Mg$^{2+}$ ions and triggers a transcriptional response. It is not currently known how the *comEA* and *dprA* genes are regulated in *A. baumannii*, but regulation of type IV pilus in *A. baumannii* has some similarities with the better-known regulation in *P. aeruginosa*. The chemosensory systems Pil-Chp (32), could have been a candidate sensor/regulator, yet in *A. baumannii*, the absence of *chpA*, *pilG*, or *pilH* in transformation conditions (D3 agarose) did not change the *pilA* transcript levels (28). Expression of the major pilin PilA is under control of the two component PilRS system which can increase *pilA* transcription in response to low levels of periplasmic PilA caused by its mobilization in the extended pilus (33). In addition, in *P. aeruginosa* the minor pilin PilY1 was found to act as a Ca$^{2+}$-dependent regulator of bacterial surface motility, favoring cycles of pilus extension/retraction (34). Hence, we hypothesized that PilY1 could sense Ca$^{2+}$, activates pilus dynamics, which would in turn trigger *pilA* transcriptional activation through the PilRS system. To test this hypothesis, we deleted the *pilY1* gene, which, as shown for several species including *Acinetobacter baylyi* should cause complete loss of pilus assembly (34–36). As expected, the Δ*pilY1* mutant is completely defective for natural transformation (Fig. 3D). However, the loss of PilY1 did not alter the transcriptional activation of *pilA* in response to CaCl$_2$ as determined at the

populual level (Fig. 3E) and at the single cell level (Fig. 3F and 3G). Thus, transcriptional activation of *pilA* occurs independently of type IV pilus dynamics. In conclusion, $Ca^{2+}$ and $Mg^{2+}$ ions alike induce a transcriptional response mainly increasing the expression of type IV pilus and transformation genes. Interestingly, although $CaCl_2$ and $MgCl_2$ induce transcriptional activation of the same magnitude, $CaCl_2$ can increase transformation 10- to 20-times more than $MgCl_2$ (Fig. 2B and Fig. 2C), suggesting that its transformation-stimulating effect is not solely due to increased pilus expression.

**Figure 3. Calcium and magnesium induce a type IV pilus and transformation regulon.**
A. RNAseq analysis of AB5075 grown for 3 hours in tryptone medium or and tryptone medium supplemented with $CaCl_2$ or $MgCl_2$. Differentially-expressed genes (log2FC<-1 or log2FC>1 and P-value<0.05) in both $CaCl_2$ and $MgCl_2$ conditions are shown in yellow. Genes specifically differentially-regulated by $CaCl_2$ and $MgCl_2$ are highlighted in green and blue, respectively.
B. Expression of a transcriptional fusion of the *pilA* gene to *gfp* in AB5075-T Δ*pilA*. Bacterial growth ($Abs_{600}$) and change in *pilA*::*gfp* expression relative to cell density (dGFP/dt, arbitrary units) were measured every 10 min over an 8-h period. $CaCl_2$ and $MgCl_2$ were added to tryptone medium at 0.68 mM.
C. Same as B, with increasing concentrations of $CaCl_2$ in tryptone medium.
D. The PilY1 minor pilin is essential for natural transformation in strain AB5075. Transformation conditions are identical to those of Figure 2. Presented data are an average of three independent experiments.
E. Fold-change in *pilA*::*gfp* expression caused by $CaCl_2$ at 0.68 mM in AB5057-T Δ*pilA* and AB5075-T Δ*pilA* Δ*pilY1*. Expression kinetics were recorded as in panel B, and fold-change represents the ratio of the maximal dGFP/dt of the $CaCl_2$ condition relative to tryptone alone. Individual results from four independent experiments are shown.
F. Fluorescence microscopy *pilA*::*gfp* expression in AB5057-T Δ*pilA* and AB5075-T Δ*pilA* Δ*pilY1* with or without $CaCl_2$. Scale bars (bottom right corner) represent 10µm.
G. Single cell expression of *pilA*::*gfp* expression in AB5057-T Δ*pilA* in tryptone medium with (n=4680,

median level=772) or without CaCl$_2$ (n=2449, median level=187), and AB5075-T Δ*pilA* Δ*pilY1* with (n=4373, median level=885) or without CaCl$_2$ (n=2848, median level=203). Results are from two independent experiments.

## Calcium stabilizes the minor pilin complex

In *P. aeruginosa*, recent structural analyzes show that PilY1 forms a champagne cork like-structure that plugs PilQ secretin and primes the minor pilins facilitating Type IV pilus assembly and dynamics (16). Ca$^{2+}$ binding to the conserved calcium binding loop of PilY1 is proposed to inhibit PilT-mediated pilus retraction, while in the absence of calcium pilus retraction is favored (34). We hypothesize that in *A. baumannii*, Ca$^{2+}$ could also have a structural effect on the PilY1-minor pilin complex, directly influencing the dynamics of the pilus and, consequently, natural transformation. We first modeled PilY1 in complex with the minor pilins PilX, PilV, PilW, FimU, FimT, PilE1, PilE2 and PilA major protein using Alphafold3. Modeling of the PilY1-PilA-minor pilin complex produced a top rank model, with metrics indicating a good confidence for the overall topology (pTM of 0.72) and for inter-protein interface (ipTM of 0.73) (Fig. 4A). The modeled structure of PilY1 exhibits the characteristic architecture described in the crystal structure of PilY1 of *P. aeruginosa* (34), with an N-terminal globular head domain and a C-terminal beta-propeller fold (Fig. 4A). Importantly, the model shows that the beta-propeller fold makes direct contact with the minor pilin complex, in agreement with previously proposed pilus assembly models, based on genetic and biochemical evidence in *P. aeruginosa* (16, 37) and on cryo-electron tomography and protein interaction networks of the *M. xanthus* pilus (17). PilX is positioned directly below the C-terminal domain of PilY1 and appears to be the only direct contact with PilY1. The others minor pilin (PilV, PilW, FimT, FimU, PilE1, PilE2) and the major pilin PilA show no direct contact with PilY1 but organize into a compact helical bundle positioned beneath or beside PilX (Fig. 4A). These observations suggest that PilX may act as a structural bridge between the minor pilin complex and PilY1, potentially serving as a central element of type IV pilus initiation complex. To locate the calcium binding site on PilY1, we next modeled PilY1 in complex with PilX and performed a Ca$^{2+}$ docking using Alphafold3. The best model exhibits a high ranking confidence of 0.8, a pTM of 0.76, an ipTM of 0.81. Two Ca$^{2+}$-binding motifs are predicted. One is located in the N-terminal globular head domain and has no known function, and was not present in the crystal structure of the *P. aeruginosa* PilY1 (34). The other one is located in the beta-propeller and correspond to the one required for twitching motility in *P. aeruginosa* (34) (Fig. 4B). Given that the only direct interaction of PilY1 with PilX involves the beta-propeller fold containing the calcium binding site, we next tested the role of Ca$^{2+}$ binding to

the PilX-PilY1 interaction. To do so, we performed a molecular mechanics, generalized Born surface area calculation (MMGBSA) in presence or absence of $Ca^{2+}$ and estimated the binding free energy ($\Delta G\_bind$) (Table1). This approach assesses how $Ca^{2+}$ influences the stability and interaction strength between PilY1 and PilX. To ensure robustness and reproducibility, we performed two independent simulations, each on 5 structural models, including the best model presented in (Fig. 4B). In the absence of divalent cation, $\Delta G\_bind$ values ranged between -195 and -210 Kcal/mol depending on the model, with negative values indicating stable interaction between PilY1 and PilX. In the presence of $Ca^{2+}$, $\Delta G\_bind$ values ranged between -212 and -230 Kcal/mol corresponding to a $\Delta\Delta G$ values of -18,77 Kcal/mol compared to the condition without $Ca^{2+}$. This decrease of binding free energy value clearly supports that $Ca^{2+}$ enhances the stability of PilY1-PilX complex. In contrast, in the presence of a $Mg^{2+}$ ion, $\Delta G\_bind$ values ranged between -188 and -207 Kcal/mol with a $\Delta\Delta G$ values of –2,34 Kcal/mol compared to the condition without ion. This small change in binding free energy indicates that the $Mg^{2+}$ has a moderate role in stabilizing PilY1-PilX complex. To validate these results, we also performed the more computationally intensive molecular mechanics Poisson-Boltzmann surface area (MMPBSA) calculation. Consistent with the MMGBSA results, the MMPBSA analysis confirmed that $Ca^{2+}$ binding strengthens the PilY1-PilX interaction (Table S3). To further investigate the role of this calcium binding loop in natural transformation, we introduced in PilY1 the substitution D947A predicted to impair $Ca^{2+}$ coordination without altering the overall structure of PilY1 (Fig. 4C) (34, 38). In agreement with predictions, transformation efficiency is drastically reduced in the D947A substitution compared to the wild type (Fig. 4D). Taken together, the results indicate that $Ca^{2+}$ helps stabilize the PilY1-PilX complex. $Ca^{2+}$ binding, more than $Mg^{2+}$, strengthens the interaction between PilY1 and PilX, which in turn promotes the recruitment of the other minor pilins and favors type IV pilus assembly.

| Condition | replicate | ΔG_bind (best model) | Mean ΔG_bind (5 models) | ΔΔG (best model) | ΔΔG (5 models) |
|---|---|---|---|---|---|
| No ion | N1 | -212.69 | -194.52 | | |
| | N2 | -227.06 | -209.74 | | |
| | Mean | -219.88 | -202.13 | 0 | 0 |
| Calcium | N1 | -246.02 | -211.53 | | |
| | N2 | -221.18 | -230.27 | | |
| | Mean | -233.6 | -220.9 | -13.72 | -18.77 |
| Magnesium | N1 | -206.99 | -183.13 | | |
| | N2 | -237.45 | -207.32 | | |
| | Mean | -222.22 | -197.73 | -2.34 | 4.4 |

**Table 1 : Binding free energy of PilY1-PilX.** Binding free energy of PilY1-PilX ($\Delta G\_bind$) in the absence (control) or presence of $Ca^{2+}/Mg^{2+}$. Energy was calculated using MMGBSA (CHARMM36, GB8). $\Delta G\_bind$ are shown for both replicate (N1,N2) of the mean of 5 independent structural models and the best model. $\Delta\Delta G$ values is the binding free energy difference between the control and the others conditions. A decrease of $\Delta\Delta G$ reflects a higher complex stability.

**Figure 4 : Calcium stabilizes the PilY1-minor pilin complex.**
A. Alphafold multimer modeling of PilY1, Type IV minor pilins and major pilin complex. Structure confidence metrics best model ipTM = 0.69, pTM = 0.71, ranking confidence = 0.73. Cartoon representation PilY1 (green), PilX (yellow), PilA (orange), PilV (blue), PilW (hotpink), FimU (lime), FimT (wheat).
B. Close up on the AlphaFold model of PilY1 and PilX with calcium docking and highlighting the calcium-binding loop (red). Structure confidence metrics ipTM = 0.80, pTM = 0.76, ranking confidence = 0.81. PilY1 (green), PilX (yellow).
C. Calcium binding loop modeling with relative distance in angstroms.
D. Natural transformation of AB5075-T encoding the wild-type PilY1 (WT) or the PilY1 D947A substitution in the presence of $CaCl_2$. Presented data are an average of three independent experiments.

# Discussion

We here report the conditions that are permissive to natural transformation in *A. baumannii* in liquid medium. Up until now, natural transformation of *A. baumannii* required the use of solid medium, and DNA uptake seemed to correlate with type IV pilus-dependent motility (26–28, 30). It could be hypothesized that surface sensing was the trigger for expression of the Type IV genes, and possibly also for other transformation genes that are needed to

take up and recombine DNA. This hypothesis is supported by work in *P. aeruginosa* and *Caulobacter crescentus* showing that the surface-dependent resistance to pilus retraction is a trigger for intracellular signaling, further inducing Type IV pilus genes, virulence or chemotaxis genes (14, 39, 40). Here we found that transformation can occur in liquid medium if supplemented with submillimolar concentrations of $CaCl_2$ or $MgCl_2$. Under these conditions, the bacteria are not attached to a surface and cycles of extension-retraction would occur in planktonic cells, allowing the capture of extracellular DNA, likely through the binding to the minor pilin PilT (41). Activation of Type IV pilus in liquid medium is also supported by the $CaCl_2$-dependent and PilA-dependent formation of large planktonic clumps of cells (Fig. S1). Activation of Type IV pilus, and processes that depend on it, can thus be achieved in the absence of surface sensing by supplementing the medium with calcium ions. Interestingly, our earlier findings suggested that $CaCl_2$ and $MgCl_2$ had minimal impact on transformation (9). These experiments used transformation conditions based on agarose D3, which we show here readily contains calcium and magnesium ions at levels sufficient for optimal transformation efficiency, hence explaining the lack of effect of additional calcium or magnesium ions. Another work also reported that $CaCl_2$ can induce transformation in the *A. baumannii* strain A118 in LB medium but with only a 4-fold increase in transformation frequencies, a much weaker effect than the >1,000-fold increase reported here. We did confirm here the weak induction of $CaCl_2$ in LB medium, and show it to be due to the presence of NaCl inhibiting transformation (Fig. 2E). Sodium chloride was indeed previously reported to inhibit transformation, causing a 100-fold decrease in transformation frequencies in A118 (29) and most other tested strains (12). Our results indicate that this is due, at least in part, to sodium chloride causing the transcriptional repression of the pilus genes (Fig. 3C).

Increases in transcript levels is a hallmark of exposure of *A. baumannii* to $CaCl_2$ and $MgCl_2$ alike (Fig. 3A). The transcriptional response is surprisingly specific and coordinated, inducing all of the type IV pilus genes operons (*pilMONPQ*, *pilA*, f*imU-pilVWXY1-pilE1-pilE2*, *pilGHIL-chpA*, *pilBCD*, *pilRS*) but also transformation-specific genes such as *yraN*, *dprA* or *comEA*. This suggests that sensing of $Ca^{2+}$ and $Mg^{2+}$ triggers the activation of a transcriptional regulator. A possible candidate would have been the chemosensory system Pil-chp, integrating sensing by solute binding protein, yet the known response regulators of the Pil-chp system (PilG, PilH) have no role in gene expression, and most importantly, deletion of *chpA, pilG,* or *pilH* did not influence *pilA* expression (28). Another candidate was the main two-component system of the Type IV pilus system, PilRS, possibly acting through the sensing of $Ca^{2+}$ by the type IV pilus adhesin PilY1. PilS directly senses the

decrease in periplasmic PilA levels and auto-phosphorylates itself. Transfers of the phosphate to PilR activates the transcription of *pilA* (28) and, according to work on *P. aeruginosa,* also of other Type IV pilus operons (42). $Ca^{2+}$ sensing by PilY1 could have stimulated PilA incorporation into the pilus, depleting periplasmic PilA, thus inducing a type IV pilus regulon. However, in a *pilY1* mutant that cannot sense $Ca^{2+}$ and also cannot assemble PilA into a functional pilus, $CaCl_2$ still induces *pilA* expression, indicating that it acts irrespective of the periplasmic levels of PilA sensed by PilRS (Fig 3E). The molecular sensor that can specifically interact with calcium and magnesium ions produce increased levels of mRNA of Type IV pilus genes remains unknown, and may be distinct from classical two-component system involving a transcriptional regulator. One possibility is that the transcriptional response reflects a post transcriptional mechanism. Indeed, the $CaCl_2$ effect on *pilA* transcription appears to be moderate (~3-fold increase in transcript) compared to the phenotypic effect (>1,000-fold increase in transformation rate). A similar situation exists in the growth phase variation of PilA protein expression which seems to exceed the 10-fold variation in its mRNA level (28). This may be the sign of a post-transcriptional regulation in which the increase in mRNA levels is not due to transcription but rather to the mRNAs being actively translated, thereby increase their stability and steady-state levels. This would be reminiscent of the control of expression of the minor pilins and the DNA uptake system proteins (ComEA, ComF, ComEC, ComM) in *L. pneumophila*, which are all under the post-transcriptional repression of a multi-target non-coding RNA *(23).* Supporting this hypothesis, a new sRNA regulator named ArpA (for *Acinetobacter* repressor of pilin) was found to target the Shine-Dalgarno sequence and the first 17 codon of *pilA* mRNA, blocking its translation and impacting natural transformation and twitching motility (43). Further work is needed to test whether ArpA, or additional sRNA, can simultaneously target in the other type IV pilus operons, *yraN*, *dprA* and *comEA*. If so, the expression of these sRNA might be modulated by calcium and magnesium ions and growth phase.

Interestingly, although $CaCl_2$ and $MgCl_2$ seemed identical in the activation of a transformation regulon, transformation efficiencies are about 10-times higher with $CaCl_2$ than with $MgCl_2$. This difference may rely on the calcium-binding loop of the non-pilin adhesin PilY1 (also named PilC1 in *Neisseria gonorrhoeae).* In *P. aeruginosa* the calcium binding site of PilY1 is required for twitching motility (34) and in *N. gonorrhoeae* the PilC1 calcium-binding mutant had a partial defect in natural transformation (38). Based on genetic and biochemical evidence, PilY1 is proposed to form a complex with minor pilins PilVWX which also interacts with PilE and FimU, acting as connector with the major pilin

PilA (37). A more recent model on the *M. xanthus* pilus, based on cryo-electron tomography and protein interaction networks also supports this macromolecular organization where PilY1 sits atop the minor pilin complex (44). AlphaFold predictions of PilY1 structure with PilX, PilV, PilW, FimU, FimT, PilE1, PilE2 and PilA of *A. baumannii* is consistent with the previously proposed models, with all minor pilins forming a helical bundle linking PilY1 to the major pilin PilA. Notably, PilVWX forms a complex beneath PilY1 while FimU-PilE1E2 contact the major pilin PilA. PilY1 contacts with the minor pilin complex by interacting directly with PilX through its the calcium-binding loop. Supporting this hypothesis, the PilY1-PilX interaction is more energetically stable with $Ca^{2+}$ coordinated in the loop. $Mg^{2+}$ may not be the primary ligand of the PilY1 calcium loop and weakly stabilize the interaction between PilX-and PilY1. In *P. aeruginosa*, it was proposed that the PilY1 C-terminal domain bound to $Ca^{2+}$ inhibits PilT-mediated pilus retraction, while the calcium-free PilY1 does not inhibit PilT and allows pilus retraction to proceed. The higher stability of the PilY1-PilX complex when bound to $Ca^{2+}$ support this hypothesis, with a more stable minor pilin complex preventing disassembly and pilus retraction. The interconversion between $Ca^{2+}$-bound and $Ca^{2+}$-free PilY1 was proposed to control the cycles of pilus extension and retraction necessary for twitching motility. This hypothesis could be extended here to natural transformation. If $Ca^{2+}$-binding causes less frequent disassembly, this will result in overall longer residence time of the DNA receptor FimT at the tip of the Type IV pilus (41), increasing the likelihood that DNA can be bound and imported in the periplasmic space.

The induction of natural transformation by $CaCl_2$ described here is fundamentally distinct from the use of $CaCl_2$ to artificially induce competence in *Escherichia coli*. In *E. coli*, high concentrations of $CaCl_2$ (millimolar range) are used to transiently alter membrane permeability, allowing passive entry of plasmid DNA that does not efficiently recombine with the chromosome. In contrast, the data support that $Ca^{2+}$ in *A. baumannii* functions as a biological signal, effective at submillimolar concentrations, specifically inducing the coordinated expression and activity of the natural transformation machinery to take up and recombine extracellular DNA. Calcium ions are abundant in host-associated environments (wounds, body fluids) and on medical surfaces frequently colonized by *A. baumannii*. Whether $Ca^{2+}$ contributes to regulation of natural transformation in these settings remains to be determined. However, $Ca^{2+}$ is also abundant and locally variable in soil and aquatic environments and may represent an ecologically meaningful cue. Yet, the environmental reservoirs of *A. baumannii* remain poorly defined and, as of today, the ecological relevance of calcium ions for this species seems difficult to assess. Future work investigating the

natural habitat of this species may benefit from including measurements of calcium ion
concentrations.

Importantly, independent of its ecological and clinical significance, the addition of $CaCl_2$ to
NaCl-free growth medium enables efficient natural transformation of *A. baumannii* in liquid
culture. $MgCl_2$, although less effective at stimulating transformation, improves bacterial
growth and thereby increases the number of transformants recovered. Based on these
observations, we established a robust protocol for natural transformation of *A. baumannii*,
which will greatly facilitate genome editing by homologous recombination.

## Material and Methods

**Bacterial strains, construction and growth conditions**

Bacterial strains and oligonucleotides used in this study are listed in the supplementary
tables S1 and S2, respectively. Genetic modifications were made in strain AB5075 (45) or
its ΔAbaR11 derivative, AB5075-T (46) and are listed in Table S2. Genetic modifications
were obtained using overlap extension PCR to synthesize a large chimeric DNA fragment
bordered by 2-kb fragments homologous to the insertion site and as previously described
(46). For strain Δ*pilY* and *pilY1*::D947A, we adapted the multiplex genome editing by
natural transformation (MuGENT) developed in *V. cholerae* (47). Briefly, this method relies
on the cotransformation of two or more PCR fragments by one transformable cell. One
PCR fragment carrying a selectable cassette (*aacC4* for apramycin or *tetC* for tetracycline)
that integrates into a neutral genomic locus (*att*Tn7), is co-transformed along with PCR
fragments carrying the desired mutation flanked by 2-Kb homology arms. The optimal yield
was obtained when co-transforming 800 ng of non-selectable fragment with 400 ng of
selectable fragment. The construction of the *pilA-sfGFP* fusion to follow *pilA* expression is
described in the supplementary information along with a new chromosomal expression
platform. Strains were grown either in lysogeny broth (LB) or tryptone medium (5g/L Bacto
Tryptone, Gibco). All experiments were performed at 37°C. Agarose soluble extract media
(ASE) was prepared by adding 0.2 g of agarose D3 (Euromedex) to 10 ml of tryptone
media. This suspension was vortexed for 5 min and subsequently centrifuged to sediment
the insoluble agarose particles. The supernatant was collected and filtered using a 0.22
µm filter.

**Construction of a chromosomal expression platform at the att*Tn7* locus**

We constructed the plasmid pAC-1 that carries the *sacB_aac4* selection/counter-selection cassette and a strong synthetic promoter (Pst) upstream of a reporter gene (*gfp* amplified from from pASG-1 (9)) using an overlap extension PCR into pJET1.2 following manufacturer instructions (CloneJET PCR cloning Kit, Thermofisher scientific). Strong terminators (5S ribosomal RNA and *filA* terminators) were added to obtain the pAC-2 plasmid carrying the $T_{5S}$-*sacB_aacC4_Pst-gfp*-$T_{filA}$ cassette. Chromosomal platforms were constructed in a two-step manner using transformation of overlap extension PCR fragments as described previously (46). First, the $T_{5S}$-*sacB_aacC4_Pst-gfp*-$T_{filA}$ cassette was inserted in the attTn7 locus by transformation of large chimeric DNA fragment carrying the $T_{5S}$-*sacB_aacC4_Pst-gfp*-$T_{filA}$ cassette flanked by 2 kb fragments that are homologous to the attTn7 insertion site. The resulting strain was then transformed with a chimeric PCR product carrying a fragment encompassing pilA promoter region and coding sequence and recombinants to replace the *sacB_aacC4_*Pst sequence. Transformants were counter-selected on minimal medium (M63) with 10% sucrose.

**Natural transformation conditions for strain constructions**

Strains were grown overnight in tryptone medium, followed by dilution in tryptone medium supplemented with $CaCl_2$ (0.68 mM) to obtain a final OD of 0.01. One hundred µl of culture were transferred to a 96-well plate along with 200 ng of transforming DNA and grown at 37 °C for a minimum of 7 hours, followed by plating on solid selective media (LB agar plates supplemented with apramycin, 50 µg/L or tetracycline 5 µg/L).

**Populational and single cell expression of *pilA*::*gfp***

Strains were grown for 16h in tryptone media and diluted to obtain a final optical density ($OD_{600}$; optical path 1 cm) of 0.01 in the different media tested. Cell density in cultures in 96-well plates ($Abs_{600}$) and fluorescence were monitored every 10 minutes over 8 hours of growth on a Tecan Infinite 200 Pro. The raw fluorescence data was smoothed using a 3-point moving average window and subsequently blank corrected using non-fluorescent wild type cells as control. $OD_{600}$ values were similarly corrected. Expression was calculated by taking the derivative of fluorescence relative to cell density at each time point ($dGFP/dT/Abs_{600}$).

Single cell expression was obtained by fluorescence microscopy of cultures at $OD_{600}$ of
0.01 (as for populational analysis above) grown for 3 hours in tryptone medium with or
without $CaCl_2$ (1 mM). Culture were placed on agar pad on glass slides and imaged with a
oil-immersion 60X objective with a EVOS M7000 microscope (Thermofisher Scientific)
under phase constrast and under LED illumination for detecting GFP (482/25 nm
Excitation; 524/24 nm Emission). Bacterial cells were segmented with Omnipose Cellpose
(v4.08) (48) and fluorescence signal was quantified with MicrobeJ (v5.10n) (49).

**Gene Expression Analysis and Expression Profiling by RNA-Seq**

Pellets from bacterial cultures in various media were stabilized and lysed as previously
described using the RNAsnap method (50), followed by RNA extraction as previously
described (23, 51). These samples were subsequently treated with DNase I and purified.
Ribosomal-depleted RNA were used to construct strand-specific (ASE condition) or
unstranded ($CaCl_2$, $MgCl_2$ conditions) libraries, which were sequenced on a Novaseq
instrument (Illumina) (GENEWIZ Germany GmbH). Raw read processing and differential
expression analyses were carried out using the Curare pipeline (52),  using fastp for read
trimming, Bowtie2 for alignment to the AB5075-UW reference genome
(GCF_031932345.1), and DESeq2 for differential gene expression analysis. Raw reads
are available at NCBI under Bioproject Accession number PRJNA1372483.

**Transformation assay using luminescence**

Transformation assay using luminescence were previously described (10). It is based on
the uptake and recombination of the pJET.Ab-pilMNOPQ::nLuc plasmid DNA. As this
plasmid is non-replicative in *A. baumannii*, DNA molecules which are internalized undergo
a double recombination event allowing the insertion of nanoluc gene in the pilMNOPQ
locus and subsequent expression of the NLuc luciferase enzyme. Bacteria from an
overnight culture (LB no salt, 10 g/L tryptone, 5 g/L yeast extract) were diluted at an $OD_{600}$
of 0.01 into tryptone medium supplemented with divalent cation when indicated. In a 96-
well plate, 100μl were mixed with 250ng of pJET.Ab-pilMNOPQ::nLuc plasmid DNA and
incubated overnight (unless indicated otherwise) at 37°C in a static incubator. The optical
density of the cell suspension at 600 nm ($Abs_{600}$) was measured on a Tecan microplate
reader. Subsequently, 80 μl of cells were mixed with 20 μl of the mix consisting of Nano-
Glo® Luciferase Assay Substrate diluted 1:50 in Luciferase Assay Buffer following the
manufacturer instructions (Nano-Glo® Luciferase Assay System, #N1120, Promega).

Following incubation at room temperature for 10 minutes, luminescence was measured in
a GloMax® Navigator Microplate luminometer (Promega). Relative luminescence units
(RLU) were calculated by dividing the luminescence values by the optical density values.

**Modeling and free energy analysis of AB5075 type IV pilus initiation complex**
The complete structure of the type IV pilus initiation complex comprising the major pilin
PilA, the minor pilins (PilX-PilV-PilW-FimU-FimT) and the adhesin PilY1 was modeled
using AlphaFold3 server (53). Five independent models were generated and evaluated
using the predicted TM-score (pTM), the interface predicted score (iPTM) and ranking
confidence metrics provided by Alphafold. The best model with the highest combined
confidence values was selected for analysis. Interaction free energy between PilY1-PilX
were estimated using the poisson-boltzmann model (PB8) gmx_MMPBSA v1.6 or the
generalized born model MMGBSA (GB8) with GROMACS 2023 and the CHARMM36 force
field energy with a ionic strength of 150mM as described in (54, 55). The binding free
energy $\Delta G\_bind$ (Kcal/mol) was extracted for each condition and average across replicate
minimized model.

# Acknowledgements

530 We warmly thank Adrien Ducret for assistance in microscopy image acquisition and
531 analysis.

# Funding

533 This work was supported by the French Agence Nationale de la Recherche (ANR), under
534 grant ANR-20-CE12-0004 (TransfoConflict). XC lab is funded by a grant Equipe FRM
535 (Fondation pour la Recherche Médicale) EQU202303016268.

## 536 **References**

1. Arnold BJ, Huang I-T, Hanage WP. 2022. Horizontal gene transfer and adaptive evolution in bacteria. Nat Rev Microbiol 20:206–218.

2. Johnston C, Martin B, Fichant G, Polard P, Claverys J-P. 2014. Bacterial transformation: distribution, shared mechanisms and divergent control. Nat Rev Microbiol 12:181–196.

3. WHO updates list of drug-resistant bacteria most threatening to human health. https://www.who.int/news/item/17-05-2024-who-updates-list-of-drug-resistant-bacteria-most-threatening-to-human-health. Retrieved 21 May 2024.

4. Perez F, Stiefel U. 2022. The Impact of Natural Transformation on the Acquisition of Antibiotic Resistance Determinants. mBio 13:e0033622.

5. Mazzamurro F, Touchon M, Charpentier X, Rocha EPC. 2025. Impact of Natural Transformation on the Acquisition of Novel Genes in Bacteria. Mol Biol Evol 42:msaf192.

6. Snitkin ES, Zelazny AM, Montero CI, Stock F, Mijares L, NISC Comparative Sequence Program, Murray PR, Segre JA. 2011. Genome-wide recombination drives diversification of epidemic strains of Acinetobacter baumannii. 33. Proceedings of the National Academy of Sciences 108:13758–13763.

7. Godeux A-S, Svedholm E, Barreto S, Potron A, Venner S, Charpentier X, Laaberki M-H. 2022. Interbacterial Transfer of Carbapenem Resistance and Large Antibiotic Resistance Islands by Natural Transformation in Pathogenic Acinetobacter. mBio 13:e02631-21.

8. Tuffet R, Carvalho G, Godeux A-S, Mazzamurro F, Rocha EPC, Laaberki M-H, Venner S, Charpentier X. 2024. Manipulation of natural transformation by AbaR-type islands promotes fixation of antibiotic resistance in Acinetobacter baumannii. Proc Natl Acad Sci U S A 121:e2409843121.

9. Godeux A-S, Lupo A, Haenni M, Guette-Marquet S, Wilharm G, Laaberki M-H, Charpentier X. 2018. Fluorescence-Based Detection of Natural Transformation in Drug-Resistant Acinetobacter baumannii. Journal of Bacteriology 200:e00181-18.

10. Mazzamurro F, Chirakadavil JB, Durieux I, Poiré L, Plantade J, Ginevra C, Jarraud S, Wilharm G, Charpentier X, P C Rocha E. 2024. Intragenomic conflicts with plasmids and chromosomal mobile genetic elements drive the evolution of natural transformation within species. PLoS Biol 22:e3002814.

11. Vesel N, Iseli C, Guex N, Lemopoulos A, Blokesch M. 2023. DNA modifications impact natural transformation of Acinetobacter baumannii. Nucleic Acids Res 51:5661–5677.

12. Hu Y, He L, Tao X, Meng F, Zhang J. 2019. High DNA Uptake Capacity of International Clone II Acinetobacter baumannii Detected by a Novel Planktonic Natural Transformation Assay. Front Microbiol 10:2165.

13. Dubnau D, Blokesch M. 2019. Mechanisms of DNA Uptake by Naturally Competent Bacteria. Annu Rev Genet 53:217–237.

14. Craig L, Forest KT, Maier B. 2019. Type IV pili: dynamics, biophysics and functional consequences. Nat Rev Microbiol 17:429–440.

15. Nguyen Y, Sugiman-Marangos S, Harvey H, Bell SD, Charlton CL, Junop MS, Burrows LL. 2015. Pseudomonas aeruginosa Minor Pilins Prime Type IVa Pilus Assembly and Promote Surface Display of the PilY1 Adhesin. J Biol Chem 290:601–611.

16. Guo S, Chang Y, Brun YV, Howell PL, Burrows LL, Liu J. 2024. PilY1 regulates the dynamic architecture of the type IV pilus machine in Pseudomonas aeruginosa. Nat Commun 15:9382.

17. Treuner-Lange A, Chang Y-W, Glatter T, Herfurth M, Lindow S, Chreifi G, Jensen GJ, Søgaard-Andersen L. 2020. PilY1 and minor pilins form a complex priming the type IVa pilus in Myxococcus xanthus. Nat Commun 11:5054.

18. Stedman MJM, Deselaers S, Braus SAG, Wang D, Balaguer MG, Gossert AD, Hospenthal MK. 2025. Molecular interplay between ComEC domains allows for selective degradation of the non-translocating strand during natural transformation. Nucleic Acids Res 53:gkaf932.

19. Mortier-Barrière I, Velten M, Dupaigne P, Mirouze N, Piétrement O, McGovern S, Fichant G, Martin B, Noirot P, Le Cam E, Polard P, Claverys J-P. 2007. A key presynaptic role in transformation for a widespread bacterial protein: DprA conveys incoming ssDNA to RecA. Cell 130:824–36.

20. Hardy L, Morales V, Rousseau CJ, Bergé M, Mazzamurro F, Plantade J, Dalia TN, Dalia AB, Rocha EPC, Charpentier X, Polard P. 2026. Two D-loop resolution systems enable natural genetic transformation in bacteria. bioRxiv https://doi.org/10.1101/2024.02.06.579203.

21. Nero TM, Dalia TN, Wang JC-Y, Kysela DT, Bochman ML, Dalia AB. 2018. ComM is a hexameric helicase that promotes branch migration during natural transformation in diverse Gram-negative species. Nucleic Acids Res 46:6099–6111.

22. Dalia AB, Dalia TN. 2019. Spatiotemporal Analysis of DNA Integration during Natural Transformation Reveals a Mode of Nongenetic Inheritance in Bacteria. Cell 179:1499-1511.e10.

23. Attaiech L, Boughammoura A, Brochier-Armanet C, Allatif O, Peillard-Fiorente F, Edwards RA, Omar AR, MacMillan AM, Glover M, Charpentier X. 2016. Silencing of natural transformation by an RNA chaperone and a multitarget small RNA. Proc Natl Acad Sci USA 113:8813–8818.

24. Meibom KL, Blokesch M, Dolganov NA, Wu C-Y, Schoolnik GK. 2005. Chitin induces natural competence in Vibrio cholerae. Science 310:1824–1827.

25. Taton A, Erikson C, Yang Y, Rubin BE, Rifkin SA, Golden JW, Golden SS. 2020. The circadian clock and darkness control natural competence in cyanobacteria. Nat Commun 11:1688.

26. Wilharm G, Piesker J, Laue M, Skiebe E. 2013. DNA uptake by the nosocomial pathogen Acinetobacter baumannii occurs during movement along wet surfaces. J Bacteriol 195:4146–4153.

27. Harding CM, Tracy EN, Carruthers MD, Rather PN, Actis LA, Munson RS. 2013. Acinetobacter baumannii Strain M2 Produces Type IV Pili Which Play a Role in Natural Transformation and Twitching Motility but Not Surface-Associated Motility. mBio 4.

28. Vesel N, Blokesch M. 2021. Pilus Production in Acinetobacter baumannii Is Growth Phase Dependent and Essential for Natural Transformation. Journal of Bacteriology 203.

29. Domingues S, Rosário N, Cândido Â, Neto D, Nielsen KM, Da Silva GJ. 2019. Competence for Natural Transformation Is Common among Clinical Strains of Resistant Acinetobacter spp. Microorganisms 7:30.

30. Yu Y, Mahdi R, Al-Hilfy Leon A, Vo N, Lofgren R, Mutabazi JL, Piepenbrink KH. 2025. Variation in type IV pilus stability modulates DNA uptake and biofilm formation. J Biol Chem 301:110787.

31. Traglia GM, Quinn B, Schramm STJ, Soler-Bistue A, Ramirez MS. 2016. Serum Albumin and Ca2+ Are Natural Competence Inducers in the Human Pathogen Acinetobacter baumannii. 8. Antimicrob Agents Chemother 60:4920–4929.

32.  Yarrington KD, Limoli DH. 2025. The type IV pilus steering committee: how Pil-Chp controls directional motility. Journal of Bacteriology 207:e00396-24.

33.  Kilmury SLN, Burrows LL. 2016. Type IV pilins regulate their own expression via direct intramembrane interactions with the sensor kinase PilS. Proceedings of the National Academy of Sciences 113:6017–6022.

34.  Orans J, Johnson MDL, Coggan KA, Sperlazza JR, Heiniger RW, Wolfgang MC, Redinbo MR. 2010. Crystal structure analysis reveals Pseudomonas PilY1 as an essential calcium-dependent regulator of bacterial surface motility. Proc Natl Acad Sci U S A 107:1065–1070.

35.  Herfurth M, Treuner-Lange A, Glatter T, Wittmaack N, Hoiczyk E, Pierik AJ, Søgaard-Andersen L. 2022. A noncanonical cytochrome c stimulates calcium binding by PilY1 for type IVa pili formation. Proceedings of the National Academy of Sciences 119:e2115061119.

36.  Ellison CK, Dalia TN, Klancher CA, Shaevitz JW, Gitai Z, Dalia AB. 2021. Acinetobacter baylyi regulates type IV pilus synthesis by employing two extension motors and a motor protein inhibitor. Nat Commun 12:3744.

37.  Nguyen Y, Sugiman-Marangos S, Harvey H, Bell SD, Charlton CL, Junop MS, Burrows LL. 2015. Pseudomonas aeruginosa Minor Pilins Prime Type IVa Pilus Assembly and Promote Surface Display of the PilY1 Adhesin. J Biol Chem 290:601–611.

38.  Cheng Y, Johnson MDL, Burillo-Kirch C, Mocny JC, Anderson JE, Garrett CK, Redinbo MR, Thomas CE. 2013. Mutation of the Conserved Calcium-Binding Motif in Neisseria gonorrhoeae PilC1 Impacts Adhesion but Not Piliation. Infect Immun 81:4280–4289.

39.  Ellison CK, Kan J, Dillard RS, Kysela DT, Ducret A, Berne C, Hampton CM, Ke Z, Wright ER, Biais N, Dalia AB, Brun YV. 2017. Obstruction of pilus retraction stimulates bacterial surface sensing. Science 358:535–538.

40.  Siryaporn A, Kuchma SL, O'Toole GA, Gitai Z. 2014. Surface attachment induces Pseudomonas aeruginosa virulence. Proceedings of the National Academy of Sciences 111:16860–16865.

41.  Ellison TJ, Ellison CK. 2025. Improved DNA binding to a type IV minor pilin increases natural transformation. Nucleic Acids Res 53:gkaf467.

42.  Kilmury SLN, Burrows LL. 2018. The Pseudomonas aeruginosa PilSR Two-Component System Regulates Both Twitching and Swimming Motilities. mBio 9:e01310-18.

43.  Hamrock FJ, Guest T, Daum MN, Connell O, Ershova AS, Hokamp K, Fleming AB, Gebhardt MJ, Westermann AJ, Kröger C. 2025. DNA uptake and twitching motility are controlled by the small RNA Arp through repression of pilin translation in Acinetobacter baumannii. bioRxiv 2025.07.19.665661.

44.  Treuner-Lange A, Chang Y-W, Glatter T, Herfurth M, Lindow S, Chreifi G, Jensen GJ, Søgaard-Andersen L. 2020. PilY1 and minor pilins form a complex priming the type IVa pilus in Myxococcus xanthus. Nat Commun 11:5054.

45.  Jacobs AC, Thompson MG, Black CC, Kessler JL, Clark LP, McQueary CN, Gancz HY, Corey BW, Moon JK, Si Y, Owen MT, Hallock JD, Kwak YI, Summers A, Li CZ, Rasko DA, Penwell WF, Honnold CL, Wise MC, Waterman PE, Lesho EP, Stewart RL, Actis LA, Palys TJ, Craft DW, Zurawski DV. 2014. AB5075, a Highly Virulent Isolate of Acinetobacter baumannii, as a Model Strain for the Evaluation of Pathogenesis and Antimicrobial Treatments. MBio 5:e01076-01014.

46. Godeux A-S, Svedholm E, Lupo A, Haenni M, Venner S, Laaberki M-H, Charpentier X. 2020. Scarless Removal of Large Resistance Island AbaR Results in Antibiotic Susceptibility and Increased Natural Transformability in Acinetobacter baumannii. Antimicrob Agents Chemother 64.

47. Dalia AB, McDonough E, Camilli A. 2014. Multiplex genome editing by natural transformation. Proceedings of the National Academy of Sciences 111:8937–8942.

48. Cutler KJ, Stringer C, Lo TW, Rappez L, Stroustrup N, Brook Peterson S, Wiggins PA, Mougous JD. 2022. Omnipose: a high-precision morphology-independent solution for bacterial cell segmentation. Nat Methods 19:1438–1448.

49. Ducret A, Quardokus EM, Brun YV. 2016. MicrobeJ, a tool for high throughput bacterial cell detection and quantitative analysis. Nat Microbiol 1:16077.

50. Stead MB, Agrawal A, Bowden KE, Nasir R, Mohanty BK, Meagher RB, Kushner SR. 2012. RNAsnap™: a rapid, quantitative and inexpensive, method for isolating total RNA from bacteria. Nucleic Acids Res 40:e156.

51. Durieux I, Ginevra C, Attaiech L, Picq K, Juan P-A, Jarraud S, Charpentier X. 2019. Diverse conjugative elements silence natural transformation in Legionella species. PNAS 116:18613–18618.

52. Blumenkamp P, Pfister M, Diedrich S, Brinkrolf K, Jaenicke S, Goesmann A. 2024. Curare and GenExVis: a versatile toolkit for analyzing and visualizing RNA-Seq data. BMC Bioinformatics 25:138.

53. Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard AJ, Bambrick J, Bodenstein SW, Evans DA, Hung C-C, O'Neill M, Reiman D, Tunyasuvunakool K, Wu Z, Žemgulytė A, Arvaniti E, Beattie C, Bertolli O, Bridgland A, Cherepanov A, Congreve M, Cowen-Rivers AI, Cowie A, Figurnov M, Fuchs FB, Gladman H, Jain R, Khan YA, Low CMR, Perlin K, Potapenko A, Savy P, Singh S, Stecula A, Thillaisundaram A, Tong C, Yakneen S, Zhong ED, Zielinski M, Žídek A, Bapst V, Kohli P, Jaderberg M, Hassabis D, Jumper JM. 2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3. Nature 630:493–500.

54. Hou T, Wang J, Li Y, Wang W. 2011. Assessing the performance of the MM/PBSA and MM/GBSA methods. 1. The accuracy of binding free energy calculations based on molecular dynamics simulations. J Chem Inf Model 51:69–82.

55. Wang S, Sun X, Cui W, Yuan S. 2022. MM/PB(GB)SA benchmarks on soluble proteins and membrane proteins. Front Pharmacol 13:1018351.