Epithelial-Mesenchymal Wnt Crosstalk Directs Planar Cell Polarity in the Developing Cochlea

Ippei Kishimoto[1], Abel P. David[2, 3], Kevin P. Rose[4], Balasubramanian Narasimhan[5], Bradley Efron[5], Sara E. Billings[1], Erin L. Su[1], Wuxing Dong[1], Taha A. Jan[6], Ronna Hertzano[4], Alan G. Cheng[1, *]

[1]Department of Otolaryngology-Head and Neck Surgery, Stanford University School of Medicine, Stanford, CA 94305, USA
[2]Department of Otolaryngology-Head & Neck Surgery, Northwestern Medicine, Chicago, IL 60611, USA
[3]Feinberg School of Medicine, Northwestern University, Chicago, IL 60611, USA
[4]Section on Omics and Translational Science of Hearing, Neurotology Branch, National Institute on Deafness and Other Communication Disorders, National Institutes of Health, Bethesda, MD 20892, USA
[5]Department of Statistics, Stanford University, Stanford, CA 94305, USA
[6]Department of Otolaryngology - Head and Neck Surgery, Epithelial Biology Center, Vanderbilt Center for Stem Cell Biology, Vanderbilt University Medical Center, Nashville, TN 37232, USA

Short title: Epithelial and mesenchymal Wnts regulate cochlear development

Abstract: 180 words
Text: 9,118 words
Figures: 6
Supplemental items: 11 figures, 1 table, 1 source data

*Corresponding author
Alan G. Cheng, M.D.
801 Welch Road
Department of Otolaryngology-HNS
Stanford, CA 94305
Phone: (650) 725-6500
Fax: (650) 721-2163
Email: aglcheng@stanford.edu

Abstract:

Cochlear hair cells are crucial for sound transduction and are precisely oriented by planar

cell polarity (PCP). Wnt proteins direct PCP in vertebrates, but their roles in regulating

cochlear PCP remain unclear. Here, we inhibited Wnt secretion by ablating Wntless in the

cochlear epithelium and found shortened cochlea, mild hair cell misorientation and

mislocalization of PCP proteins, Fzd6 and Dvl2, representing defects less severe than

classic PCP mutants. Computational cell-communication analysis predicted that

candidate Wnts from the cochlear epithelium (Wnt5a, Wnt7a, Wnt7b) and surrounding

periotic mesenchyme (Wnt5a) act on developing hair cells. Deletion of Wnt5a, Wnt7a,

and Wnt7b additively shortened the cochlea without causing hair cell misorientation or

mislocalization of PCP proteins. Moreover, deletion of Wnt5a alone in the periotic

mesenchyme failed to cause PCP defects. However, ablating both epithelial and

mesenchymal Wnts caused a severely shortened cochlea with apically malrotated hair

cells devoid of polarized core PCP proteins. Thus, Wnts serve as global instructive cues

directing cochlear outgrowth and hair cell polarization, with remarkable redundancy of

distinct Wnts across epithelial and mesenchymal compartments to ensure a fail-safe

developmental program.

Key words: cochlea, development, hair cells, Wnt, planar cell polarity

Introduction

The cochlea is a spiral-shaped sensory organ critical for sound transduction. As mechanotransducers of sound vibration, hair cells are precisely arranged both tonotopically along the length of the cochlea and radially to form three rows of outer hair cells (OHCs) and one row of inner hair cells (IHCs).

During embryonic development in mice, the cochlear duct initially forms as an outpouching from the ventral otocyst at embryonic day (E) 11 and subsequently elongates to become a two-and-a-half-turn organ[1, 2]. In the floor of the cochlear duct, terminal mitosis of prosensory cells begins in the apex and extends towards the base, with hair cell specification occurring in the opposite direction between E12.5 and E14.5[2, 3, 4]. Four rows of hair cells and interdigitating supporting cells are arranged in a checkerboard pattern and a precise radial orientation, ensuring that hair bundles, which house the mechanotransduction machinery, uniformly face laterally. This organized cochlear patterning is regulated by the planar cell polarity (PCP) pathway.

The PCP pathway is evolutionarily conserved from Drosophila to mammals, governing both convergent extension and hair cell polarity in the developing mouse cochlea[5, 6]. In humans, mutations underlying Bardet–Biedl syndrome[7] and Alstrom syndrome[8, 9] disrupt PCP signaling and cochlear hair cell development, resulting in hearing loss. In mammals, there are six groups of core PCP proteins-Frizzled (Fzd), Van Gogh (Vangl), Celsr1, Dishevelled (Dvl), Prickle (Pk), and Diego (Dgo)-all of which are asymmetrically expressed at opposite poles of cochlear hair cells and supporting cells[10, 11, 12, 13, 14, 15]. Deletion of Celsr1 or Vangl2 causes cochlear shortening and severe misorientation of hair cells[16, 17], whereas mutants of Vangl1 displayed relatively mild PCP

defects[16]. Concurrent deletion of Fzd3/Fzd6 or Dvl1/Dvl2 leads to striking PCP defects[12,]

[14], and deletion of other core PCP proteins in the presence of the Vangl2 looptail mutation

amplifies the severity[16], suggesting complementary and redundant roles among core PCP

proteins.

Wnt signaling plays multiple essential roles during development, including cell

differentiation, proliferation, cell polarity, and tissue patterning[18, 19]. The Wnt signaling

pathway can be broadly classified as canonical and non-canonical, where the former is

mediated by β-catenin and the latter by PCP signaling[20]. In the cochlea, Wnt/β-catenin

signaling is required for hair cell specification[21], radial patterning[22], and supporting cell

specification[23]. In vertebrates, although Wnts are known to regulate PCP signaling in

various organs such as the neural tube, muscles, and limbs[24, 25], whether this occurs in

the cochlea is unclear.

Among the 19 Wnt ligands in mammals, Wnt2b, Wnt4, Wnt7a, Wnt7b, Wnt9a, and

Wnt11 are expressed in the developing cochlear epithelium and Wnt5a is expressed in

both the cochlear epithelium and periotic mesenchyme (POM)[15, 26, 27]. Wnt7a mutant

cochleae show normal hair cell orientation[28]. While Wnt5a knockout cochleae display

defects in convergent extension, hair cell orientation, and localization of Vangl2, although

the defects were minor and penetrance was limited, with only one third of mutants

affected[29]. We and others have previously shown that concurrent ablation of Wnt

members in the cochlear epithelium (by deleting Wntless or Wnt5a and Wnt7b) caused a

significant shortening of the cochlea, mild misorientation of hair cells, and mislocalization

of a subset of core PCP proteins (Fzd6 and Dvl2) in hair cells[15, 30, 31]. Because the severity

of hair cell polarity defects observed in these studies is relatively mild in comparison to

other PCP mutants (e.g., Vangl2-looptail[16]), the requirement of Wnts for PCP signaling I

the cochlea has remained controversial[32, 33].

Here, we used computational analysis of single-cell transcriptomic data from the

developing cochlea and identified Wnt5a, Wnt7a, and Wnt7b in the cochlear epithelium

and Wnt5a in the surrounding POM as the top candidate Wnts that act on developing hair

cells. Deletion of multiple Wnts (Wnt5a, Wnt7a, and Wnt7b) from the cochlear epithelium

shortened the cochlear duct, without causing misorientation of hair cells or mislocalization

of core PCP proteins, suggesting a redundancy among Wnts. Inhibiting the secretion of

both epithelial and mesenchymal Wnts further shortened the cochlea and led to severely

malrotated hair cells and loss of polarization of all PCP proteins examined, with

phenotypes remarkably more severe than Wntless ablation in the cochlear epithelium

alone. Collectively, our study indicates that multiple secreted Wnts from both the epithelial

and mesenchymal compartments are required for cochlear outgrowth, hair cell orientation

and PCP signaling.

Results

Ablating Wntless from the cochlear epithelium disrupts cochlear outgrowth and

hair cell polarization.

Wntless (Wls) is a chaperone transmembrane protein required for secretion of Wnt

proteins, a process essential for all Wnts[34, 35]. We found that Wntless mRNA is expressed

throughout the developing cochlear epithelium (Fig. 1b, Supplementary Fig. 1c-c", e-e",

g-g"), corroborating previous results[15]. To ablate Wntless in the cochlear epithelium prior

to hair cell differentiation, we first established Emx2-Cre; Wls-flox mice. Emx2-Cre activity

starts in the otocyst as early as embryonic (E) 10.5 (Fig. 1a)[36]. In E16.5 *Emx2Cre/+;*

Wlsflox/flox (Emx2-Wls cKO) cochleae, there was almost no detectable Wls mRNA in the

cochlear epithelium, whereas that in the POM remained comparable to those in control

cochleae (Fig. 1c, Supplementary Fig. 1d-d", f-f", h- h"), indicating efficient and specific

deletion of Wls within the cochlear epithelium in Emx2-Wls cKO mice. At E18.5, Emx2-

Wls cKO cochleae are significantly shorter (~39%) than those from wildtype littermates

(2,856±381 and 4,653±220 μm, respectively, p < 0.0001) (Fig. 1d, e). Moreover, Emx2-

Wls cKO cochleae displayed significantly fewer (~29%) Myosin7A-positive hair cells

(mean 1,917±234 and 2,692±156 per cochlea, respectively, p < 0.01, Fig. 1f). These

findings suggest that inhibiting Wnt secretion in the cochlear epithelium leads to defects

in cochlear outgrowth and/or convergent extension. To assess hair cell orientation in

phalloidin-stained E18.5 cochleae (Fig. 1g, h, Supplementary Fig. 1i, j), we measured the

average and variability of orientation of each row of hair cells (circular mean and variance).

In control mice at E18.5, cochlear hair cell bundles face laterally with the most lateral row

of hair cells (the third row of OHCs) slightly rotated toward the apex (Fig. 1g-g',

Supplementary Fig. 1i-i'), consistent with previous reports[28, 37]. Compared to controls,

Emx2-Wls cKO cochlear hair cells exhibited significantly larger variance in the middle and

basal turns (circular variance ratio: 2.00, Supplementary Table 1), along with a mild, but

significantly different mean orientation of HCs (IHC, OHC1, and OHC3+, Fig. 1h-h', i,

Supplementary Fig. 1j-j', k, Supplementary Table 1). We also examined the molecular

basis of cochlear PCP by assessing the polarized expression of core PCP proteins in hair

cells. Hair cells display a complement of core PCP proteins[15] (Fig. 1j). In the control

cochlea, we found Fzd6, Vangl2, and Celsr1, to be localized to the medial junctions and

Dvl2 to the lateral junctions of OHCs (Fig. 1k-k''', m, o). By contrast, the Emx2-Wls cKO

cochlea lacked polarized expression of Fzd6 and Dvl2 while Celsr1 and Vangl2 remained

polarized (Fig. 1l-l''', n, p). This finding suggests that Wnt secretion from the cochlear

epithelium is required for establishing and/or maintaining the polarization expression of a

subset of, but not all, core PCP proteins. Taken together, our data suggest that disruption

of Wnt secretion from the cochlear epithelium prevents elongation of the organ and

causes a mild PCP defect (shortened cochlea, hair cell misorientation, and loss of

polarized expression of a subset of PCP proteins), corroborating previous reports[15, 31].

***In silico* analysis identified candidate Wnt ligands from the cochlear epithelium and**

periotic mesenchyme regulating planar cell polarization

To identify candidate Wnt ligands in the developing cochlea that could contribute

to hair cell polarity, we analyzed established single-cell transcriptomic data from the

embryonic cochleae. We integrated existing single-cell RNA sequencing data into two

categories, each comprised of both epithelial and mesenchymal cells: (1) early (E13.5

and E14.5)[38, 39] and late embryonic cochlea (E15.5 and E16.5)[39, 40]. From the analysis of

both the early and late embryonic cochlea datasets, we identified six cell clusters: hair

cells (HCs), cochlear roof, lesser epithelial ridge (LER), greater epithelial ridge (GER),

pro-sensory cells, and periotic mesenchyme cells (POM). These clusters were annotated

using previously described marker genes[39, 40].

We next computationally inferred cell-cell communication among different cell

clusters with a focus on Wnt ligand-receptor interactions in the developing cochlea using

the CellChat[41] algorithm. First, we combined the cochlear epithelial cells (cochlear roof,

LER, and GER cells), while excluding hair cells and prosensory cells. As a result, we

generated four new cell groups for the early and late embryonic cochlea datasets: 1) the

"Epithelial cells" group (comprising cochlear roof, LER, and GER cells); 2) the "POM"

group; 3) the "Prosensory cells" group; and 4) the "Hair cells" group, which includes both

IHCs and OHCs (Fig. 2a, b). We then used these newly defined cell groups to create

circle plots for each dataset, illustrating significant Wnt-receptor interactions among the

cell groups, with the width of the arrows reflecting the scale of these interactions based

on the calculated communication probability (Fig. 2c, d)[41]. In the early embryonic cochlea

dataset, significant bidirectional Wnt ligand-receptor interactions were detected among

all cell groups, along with autocrine signaling (Fig. 2c). In contrast, in the late embryonic

cochlea dataset, outgoing Wnts from the hair cell group interacting with any other group

were no longer significant (Fig. 2d). Furthermore, in both datasets, the strength of Wnt

ligand-receptor interactions involving the POM group was particularly notable (Fig. 2c, d).

To identify the Wnt subtype received by the hair cell and prosensory/supporting cell

groups, we analyzed the Wnt-ligand interactions in which these cells act as receptors and

created chord plots illustrating the significant Wnt ligand-receptor pairs for each dataset

(Fig. 2e, f). The values for the relative contribution, reflecting the calculated

communication probability of each Wnt ligand-receptor (Source data 1), and are

proportional to the width of the lines in the chord plots (Fig. 2e, f). We then categorized

and ranked the candidate Wnt ligands received by the hair cell and prosensory/supporting

cell groups separately based on whether they originate from the cochlear epithelium or

POM (Fig. 2g-j). In the early embryonic cochlea dataset, Wnt7b was identified as the

dominant Wnt from the cochlear epithelium, followed by Wnt7a and Wnt5a. In contrast,

for the late embryonic cochlea dataset, Wnt5a emerged as the dominant candidate Wnt

from the cochlear epithelium, followed by Wnt7b and Wnt7a (Fig. 2g, h). Among the Wnts

derived from the POM, Wnt5a was the most highly expressed across both datasets, with

Wnt5b following (Fig. 2i, j). In summary, the cell-cell communication analysis of both

datasets suggests Wnt5a, Wnt7a, and Wnt7b as candidate Wnts secreted from the

cochlear epithelium, with Wnt5a being the most robust among those derived from the

POM, that can contribute to Wnt signaling in hair cells during cochlear development.

Deletion of Wnt5a, Wnt7a, and Wnt7b from the cochlear epithelium shortens

cochlea and causes mild hair cell malrotation.

213   Because our computational analysis identified Wnt5a, Wnt7a, and Wnt7b as the

candidate Wnt ligands in the cochlear epithelium acting on hair cells and prosensory cells,

we hypothesized that concurrent deletion of these Wnt ligands from the cochlear

epithelium would phenocopy Emx2-Wls cKO cochlea. To test this hypothesis, we crossed

conditional KO animals of Wnt5a[42], Wnt7a[43], or Wnt7b[44] with Emx2-Cre mice, and

generated *Emx2[Cre/+]; Wnt5a[flox/flox]* (Emx2-Wnt5a cKO), *Emx2[Cre/+]; Wnt7a[flox/flox]* (Emx2-

Wnt7a cKO), and *Emx2[Cre/+]; Wnt7b[flox/flox]* (Emx2-Wnt7b cKO) mice. Previous studies

using *in situ* hybridization have demonstrated the expression of Wnt5a, Wnt7a, and

Wnt7b in the cochlear epithelium, particularly on the cochlear floor, at E14.5 and E16.5[15,

28, 29] (Fig. 3a). In E16.5 control cochlea, we confirmed expression of Wnt5a, Wnt7a, and

Wnt7b mRNA (Supplementary Fig. 2a, c, e, g, i, k), which were mostly deleted in the

cochlear epithelium of the Emx2-Wnt5a, Emx2-Wnt7a, and Emx2-Wnt7b cKO mice

(Supplementary Fig. 2b, d, f, h, j, l).

When Wnt5a, Wnt7a, or Wnt7b was individually ablated in the cochlear epithelium,

there was no significant difference in cochlear length compared to littermate controls (p =

0.5132, 0.9070, and 0.4882, respectively; Supplementary Fig. 3a-b, f-g, k-l). Additionally,

hair cell counts in these Emx2-Wnt5a and Emx2-Wnt7a cKO cochleae were unchanged

(p = 0.5722 and p = 0.5056, respectively), whereas Emx2-Wnt7b cKO cochleae had

mildly reduced hair cell counts (~9%; p < 0.05, Supplementary Fig. 3c, h, m). Furthermore,

the circular mean and variance of hair cell orientation in Emx2-Wnt5a and Emx2-Wnt7a

cKO cochleae were comparable to controls, while those in Emx2-Wnt7b cKO cochlea

were mildly abnormal (significant deviation of IHC, OHC1 and OHC2, Supplementary Fig.

4a-r, Supplementary Table 1). Lastly, Fzd6 and Dvl2 remained polarized in OHCs of

Emx2-Wnt5a, Emx2-Wnt7a, and Emx2-Wnt7b cKO cochleae (Supplementary Figure 3d-

e''', i-j''', n-o'''). These findings suggest that individual deletion of Wnt5a, Wnt7a, or Wnt7b

from the cochlear epithelium during development has a minimal effect on cochlear

extension and hair cell polarization, corroborating previous reports[15, 30].

Previously, Wnts were reported to have redundant functions during cochlear

development[30, 45]. To rule out this possibility, we generated double KO mice for Wnt

ligands: $Emx2^{Cre/+}$; $Wnt5a^{flox/flox}$; $Wnt7a^{flox/flox}$ (Emx2-Wnt5a; Wnt7a cKO), $Emx2^{Cre/+}$;

$Wnt5a^{flox/flox}$; $Wnt7b^{flox/flox}$ (Emx2-Wnt5a; Wnt7b cKO), and $Emx2^{Cre/+}$; $Wnt7a^{flox/flox}$;

$Wnt7b^{flox/flox}$ (Emx2-Wnt7a; Wnt7b cKO) mice. The cochlear length of Emx2-Wnt5a;

Wnt7b and Emx2-Wnt7a; Wnt7b cKO, but not Emx2-Wnt5a; Wnt7a cKO, mice were

significantly shorter than those of littermate controls (~20% and ~24%, p < 0.01 and p <

0.05, respectively) (Supplementary Fig. 5a-b, f-g, k-l). Hair cell counts in Emx2-Wnt5a;

Wnt7b and Emx2-Wnt7a; Wnt7b cKO cochleae were significantly lower than controls

(~20% and ~15%, p < 0.01 and p < 0.05, respectively), whereas hair cell counts in Emx2-

Wnt5a; Wnt7a cKO cochleae were comparable to littermate control cochleae (p = 0.6008,

Supplementary Fig. 5c, h, m). Emx2-Wnt5a; Wnt7a cKO cochleae showed normal hair

cell orientation (Supplementary Fig. 6a-f, 13), while Emx2-Wnt5a; Wnt7b cKO cochleae

exhibited more variable orientation of hair cells compared to controls (circular variance

ratio: 1.73, Supplementary Table 1). However, the average orientation of hair cells in

Emx2-Wnt5a; Wnt7b cKO cochleae were not significantly different from controls

(Supplementary Fig. 6g-r, 13, Supplementary Table 1). Hair cells in all three double cKO

cochleae still exhibited polarized expression of Fzd6 and Dvl2 (Supplementary Fig. 5d-

e''', i-j''', n-o'''), in contrast to the previously reported reduced expression in Emx2-Wnt5a;

Wnt7b cKO cochleae[30].

Finally, we generated and examined the *Emx2^Cre/+; Wnt5a^flox/flox; Wnt7a^flox/flox;*

Wnt7b^flox/flox (Emx2-Wnt5a; Wnt7a; Wnt7b cKO) mice. In these animals, cochlear length

was significantly lower than that of littermate controls (~30%, p < 0.001; Fig. 3b, c), and

that after concurrent deletion of Wnt5a and Wnt7a (p < 0.0001, Fig. 3e). Length of Emx2-

Wnt5a; Wnt7a; Wnt7b cKO cochlea is comparable to that of Emx2-Wnt5a; Wnt7b cKO,

Emx2-Wnt7a; Wnt7b cKO, and Emx2-Wls cKO cochleae (p > 0.05 for all, Fig. 3e). Hair

cell counts of Emx2-Wnt5a; Wnt7a; Wnt7b cKO cochleae were significantly reduced

compared to littermate controls (~16% and p < 0.01; Fig. 3d). Similarly, hair cell counts in

Emx2-Wnt5a; Wnt7a; Wnt7b cKO cochleae were significantly lower than those in any

single cKO (~17% and p < 0.01 for Wnt5a cKO, ~18% and p < 0.01 for Wnt7a cKO, and

~14% and p < 0.05 for Wnt7b cKO cochleae) or Emx2-Wnt5a; Wnt7a cKO cochleae

(~15% p < 0.05), yet comparable to those in Emx2-Wnt5a; Wnt7b cKO, Emx2-Wnt7a;

Wnt7b cKO, and Emx2-Wls cKO cochleae at E18.5 (p > 0.05 for all, Fig. 3f). These results

suggest that Wnt5a, Wnt7a, and Wnt7b in the cochlear epithelium are individually

dispensable yet contribute additively to cochlear outgrowth. By contrast, no misorientation

of the hair cells was detected (basal or middle turns) (Fig. 3g-i, Supplementary Fig. 7a-c,

Supplementary Table 1). Furthermore, Fzd6 and Dvl2 expression in hair cells remains

polarized (Fig. 3j-k'''). While the length of Emx2-Wnt5a; Wnt7a; Wnt7b cKO is similar to

that of Emx2-Wls cKO, hair cell orientation and core PCP protein expression were

remarkably normal.

In summary, these data indicate that the effects of Wnt5a, Wnt7a, and Wnt7b from

cochlear epithelium are additive for cochlear outgrowth, but not for hair cell orientation.

Furthermore, since core PCP protein expression remains polarized in all single, double,

and triple conditional KO cochleae, we postulate that additional Wnt ligands, either within

or outside the cochlear epithelium, may also contribute to planar polarization.

Deletion of Wnt5a from the POM does not cause cochlear PCP defects.

Our computational cell-cell communication analysis inferred that Wnt ligands from

the POM significantly interact with Wnt receptors on hair cells and prosensory/supporting

cells, particularly Wnt5a (Fig. 2e, f, i, j). Indeed, a previous study has reported Wnt5a

expression in the POM at E14.5[46]. To test the hypothesis that Wnt ligands from the POM

regulate cochlear outgrowth and hair cell polarity, we generated *Dermo1 (Twist2)-Cre;*

Wls^flox/flox mice. Dermo1-Cre mice exhibit Cre activity beginning around E9.5 in the

mesenchyme-derived tissues[47], including the POM but not the cochlear epithelium[48].

However, *Dermo1^Cre/+; Wls^flox/flox* (Dermo1-Wls cKO) were embryonically lethal (8 of 8

animals by E18.5) as previously reported,[49] precluding examination of their cochlea. We

next generated and examined *Dermo1[Cre/+]; Wnt5a[flox/flox]* (Dermo1-Wnt5a cKO) mice. In

control cochleae, Wnt5a mRNA was detected both in the cochlear epithelium and POM

(Fig. 4b-b', Supplementary Fig. 8a-a', c-c'), corroborating our previous study[15]. By

contrast, in Dermo1-Wnt5a cKO cochleae, *Wnt5a* mRNA signals in the POM were

undetectable while those in other regions (cochlear epithelium and spiral ganglion)

remained present (Fig. 4c-c', Supplementary Fig. 8b-b', d-d'), indicating efficient and

specific deletion of Wnt5a mRNA from the POM. For the cochlear PCP readouts, however,

Dermo1-Wnt5a cKO cochleae exhibited normal length, hair cell counts, hair cell

orientation (both circular variance and means), and polarized expression of the core PCP

proteins Fzd6 and Dvl2 (Fig. 4d-k''', Supplementary Fig. 8e-g, Supplementary Table 1).

These results suggest that deletion of Wnt5a alone from the POM does not disrupt

cochlear outgrowth, hair cell orientation and PCP signaling.

Inhibition of Wnt secretion in both the cochlear epithelium and periotic

mesenchyme causes severe planar polarization defects.

When Wnt secretion was prevented in the cochlear epithelium using Emx2-Wls

cKO mice, we have shown that hair cell orientation and polarization of two core PCP

proteins were disrupted, albeit to a markedly milder degree relative to other PCP

mutants[10, 12] (Fig. 1g-i, Supplementary Fig. 1i-k). Since Wnts from both the cochlear

epithelium and POM were predicted to significantly interact with Wnt receptors on hair

cells and prosensory/supporting cells (Fig. 2g-q), we generated *Sox9[CreERT2]; Wls[flox/flox]*

(Sox9-Wls cKO) mice to prevent Wnt secretion in both regions. Sox9 is expressed in both

the otic/cochlear epithelium and POM from E9.5 to E14.5, before becoming more

restricted later[50, 51]. Sox9-Wls cKO animals received tamoxifen (0.2 mg/g of body weight)

at E10.5-11.5 or E11.5-12.5 with survival rates of ~73% at E18.5, consistent with previous

reports[52, 53] (n=48, Fig. 5a, Supplementary Fig. 9a). In E18.5 Sox9-Wls cKO cochlea, we

found efficient and broad deletion of Wls in both the cochlear epithelium and POM

(Supplementary Fig. 10a- f').

After tamoxifen administration at E10.5-11.5, Sox9-Wls cKO cochleae were

dramatically shorter and contained significantly fewer hair cells than controls (~70% and

~60%, p < 0.0001 for both, Fig. 5b-d) and Emx2-Wls cKO cochleae (~38% and p < 0.0001,

and ~46% and p < 0.01, respectively). Sox9-Wls cKO cochlea displayed five to six rows

of hair cells throughout the cochlea (Fig. 5e-f, Supplementary Fig. 11a-b), consistent with

a defect in convergent extension seen in PCP mutants[54, 55]. Additionally, hair cells

appeared disorganized without a clear separation between IHCs and OHCs (Fig. 5f,

Supplementary Fig. 11b), possibly suggesting abnormal pillar cell development.

Furthermore, hair cells were severely malrotated with significantly higher variance and

more deviation than controls (circular variance ratio: 5.33, Fig. 5e-g, Supplementary Fig.

11a-c, Supplementary Table 1). Unlike Emx2-Wls cochleae, where only a subset of hair

cells was significantly malrotated and directed to the base, Sox9-Wls cochleae shows that

each row of hair cells is severely malrotated (Fig. 1g-i, 4e-g, Supplementary Fig. 1i-k,

11a-c, Supplementary Table 1). In contrast to PCP mutants where hair cells are randomly

malrotated[12, 16, 17], Sox9-Wls cochlear hair cells are rotated towards the apex, suggesting

a global instructive role for secreted Wnts.

In control cochleae, Celsr1, Vangl1, Vangl2, and Fzd6 are localized to the medial

junctions of hair cells, while Dvl2 is found at the lateral junctions (Fig. 5h, j, l, n, p). In

contrast, each of these core PCP proteins is mislocalized in hair cells in Sox9-Wls cKO

cochleae (Fig. 5i, k, m, o, p). Overall, compared to Emx2-Wls cKO cochleae, Sox9-Wls

cKO mice exhibited greater disruptions in cochlear duct elongation, hair cell orientation,

and polarization of core PCP proteins in hair cells.

To determine whether the reduced length and hair cell counts in Sox9-Wls cKO

cochleae were attributed to reduced proliferation, we performed Ki67 immunostaining and

EdU labeling. EdU (50 mg/kg, IP) was administered at E12.5, along with tamoxifen (0.2

mg/g) at E10.5-11.5. At E15.5, both control and Sox9-Wls cKO cochleae displayed Ki67-

positive and EdU-positive cells in both the cochlear epithelium and POM (Fig. 5r-s). No

significant difference was observed in the number of EdU- or Ki67-labeled Sox2-positive

cells in control and Sox9-Wls cKO cochleae ($p = 0.1015$ and $p = 0.4776$, respectively; Fig.

5t-u, Supplementary Fig. 11d-e). Additionally, no significant differences were detected in

the number of EdU- or Ki67-positive cells in the POM ($p = 0.7144$ and $p = 0.4975$,

respectively; Fig. 5r-s, Supplementary Fig. 11d-e). These findings suggest that the

cochlear shortening and reduced cochlear hair cell counts observed in Sox9-Wls cKO

mice are primarily due to disruptions in cochlear outgrowth and/or convergent extension.

To determine whether the regulation of cochlear outgrowth and PCP development

by Wnt secretion is time-dependent during cochlear development, we ablated Wntless

one day later (tamoxifen at E11.5-12.5, Supplementary Fig. 9a). Sox9-Wls cKO cochleae

were still significantly shorter (~26% and $p < 0.05$) and contained fewer hair cells (~14%

and $p < 0.05$) compared to controls at E18.5 (Supplementary Fig. 9b-d). However, when

compared an earlier injection of tamoxifen at E10.5-11.5, both the cochlear length (~60%

and p < 0.0001) and hair cell counts (~53% and p < 0.0001) were significantly higher

(Supplementary Fig. 9c-d). Additionally, hair cells remain significantly malrotated towards

the apex and showed a high variance in orientation (circular variance ratio: 3.28,

Supplementary Fig. 9e-j, Supplementary Table 1). However, core PCP proteins (Fzd6,

Celsr1, Vangl1, and Vangl2) were all polarized in hair cells (Supplementary Fig. 9k-s),

suggesting the overall effects were less severe than Wntless deletion earlier. These

results suggest that the effect of Wnts on the cochlear PCP development is highly time-

sensitive during the embryonic period.

Overall, these data suggest that secreted Wnt proteins, around E10.5-11.5, from

both the POM and cochlear epithelium are required for elongation of the cochlear duct,

orientation of hair cells, and polarization of core PCP proteins in hair cells during cochlear

development.

Discussion

During development, the mouse cochlea elongates to form a spiral organ that is

two and a half turns in length, in which sensory and non-sensory cells differentiate and

undergo precise radial patterning. These steps are temporally and spatially coordinated

and governed by the Notch, Wnt, and Shh signaling pathways[2]. PCP signaling critically

regulates cochlear elongation and sensory cell polarization. Previous studies have shown

that secreted Wnts in the cochlear epithelium are required for cochlear elongation,

sensory cell polarization, and polarization of PCP proteins[15, 31], although the severity of

the sensory cell polarization defects was remarkably mild, raising the possibility that other

factors co-regulate these steps. Here, based on computational analysis of single-cell

transcriptomes of the developing cochlear epithelium and surrounding POM, we predicted

that Wnt5a, Wnt7a and Wnt7b in the cochlear epithelium, and Wnt5a in the POM were

the most potent Wnts acting on the developing hair cells and prosensory/supporting cells.

Using combinations of transgenic mice, we selectively deleted Wnt members or Wnt

secretion in a spatiotemporally selective manner. We first demonstrated that these Wnt

members in both the cochlear epithelium and POM additively contribute to cochlear

outgrowth and convergent extension. By contrast, hair cell orientation and PCP signaling

based on the polarized expression of core PCP proteins are only mildly affected despite

deletion of all three Wnt members or Wntless in the cochlear epithelium. When Wnt

secretion was prevented in both the cochlear epithelium and POM, cochlear length, hair

cell orientation and PCP signaling were severely affected, suggesting that Wnt members

distributed across the epithelial and mesenchymal compartments are highly redundant in

regulating cochlear development.

<u>Wnts are required for cochlear hair cell planar cell polarity and cochlear outgrowth</u>

In our current study, inhibition of Wnt secretion in the cochlear epithelium during

development only caused mild hair cell misorientation and loss of polarization in a subset

of core PCP proteins, consistent with previous studies[15, 31]. This suggests that Wnt

proteins from the cochlear epithelium have a limited role in establishing hair cell polarity.

In contrast, Sox9-Wls cKO cochleae demonstrated severe hair cell misorientation and

loss of polarization of all core PCP proteins examined. Thus, Wnts from both the cochlear

epithelium and POM are critical in regulating hair cell polarity.

Wnts have previously been implicated to function as global cues for PCP[25]. In

Drosophila, Wingless and Wnt4 modulate intercellular Fzd–Vangl interactions, thereby

establishing directional cues for PCP axis orientation[56]. However, recent studies in

Drosophila have challenged the idea that secreted Wnts are required for core PCP protein

enrichment and planar polarity[57, 58]. In various vertebrate organs, the role of Wnt ligands

in regulating PCP is well established. For instance, Wnt5b and Wnt11 coordinate cardiac

remodeling through polarization of actomyosin activity in zebrafish. During chick muscle

development, Wnt11 directs elongation of myocytes along the antero-posterior axis of the

embryo[59]. In mice, Wnt5a establishes PCP in limb mesenchyme[60, 61] and Wnt5a and

Wnt5b direct node cell polarization in the developing heart[62], while Wnt5a induces

phosphorylation of Vangl2 in a dose-dependent manner[63]. However, the relationship

between Wnt signaling and PCP is not yet clear in the cochlea.

In the mouse cochlea, several studies have found that Wnt proteins in the cochlear

duct regulate the orientation and PCP signaling in hair cells[15, 30, 31]. However, in part

because deficiency of Wnt proteins results in hair cell defects that are relatively mild

compared to other PCP mutants, it has been suggested that Wnts do not play a central

role in determining cochlear PCP[32, 33]. Our results in this study clearly show that both

epithelial and mesenchymal Wnts contribute to cochlear outgrowth and hair cell planar

cell polarity, and not merely epithelial Wnts as previously suggested[15, 30, 31].

PCP mutants such as *Vangl2*[Lp/Lp] demonstrate convergent extension defects, with

shorter cochleae (~27% of wild-type littermates) without a significant reduction of hair

cells[10]. In contrast, ablation of Wntless in the cochlear epithelium (Emx2-Wls cKO) and in

both the cochlear epithelium and POM (Sox9-Wls cKO) lead to significantly shorter

cochleae (~38% and 70%, respectively) and a decrease in hair cells. Because the degree

of shortening is severe and more than accounted for by the convergent extension defect

alone, our study suggests that both epithelial and POM Wnt proteins are required for

cochlear outgrowth and convergent extension. When comparing the results from

transgenic mouse lines to delete one or multiple Wnts and epithelial or both epithelial and

mesenchymal Wnts, we conclude that multiple Wnt members additively contribute to

cochlear outgrowth (Fig. 6).

While cochlear PCP mutants are characterized by hair cell malrotation, different

degrees of defects have been reported, with Vangl2 and Celsr1 mutants being more

severe and the Vangl1 mutant milder[16, 17]. Moreover, core PCP proteins serve redundant

functions, with deletion of multiple members or the presence of the dominant negative

Vangl2-looptail mutation accentuating the polarity defects[12, 14, 16]. Several studies have

demonstrated that deletion of Wntless or multiple Wnts in the cochlear epithelium causes

mild hair cell polarity defects and loss of or decreased polarized expression of Fzd6 and

Dvl2, but not other core PCP proteins. When Wnt secretion is prevented in both cochlear

epithelium and periotic mesenchyme (using Sox9-Wls cKO at E10.5-11.5 and not E11.5-

12.5), hair cell polarity defects are severe and all core PCP proteins mislocalized. Thus,

our study demonstrates that secreted Wnt proteins are not only required for PCP

establishment in the developing cochlea (hair cell orientation and core PCP localization)

but also do so in an additive and time-dependent manner.

Dual regulation of hair cell orientation by PCP core proteins and Wnts

Cochlear hair cells are uniformly oriented with their bundles facing laterally. Prior

work on PCP mutants has characterized hair cell orientation to be highly variable with

random rotation in both clockwise and counterclockwise directions[10, 12, 17]. These results

suggest a local, instructive role for core PCP proteins in guiding the polarization of hair

cells and limiting variance in orientation. In support of this role, Vangl2 misexpression

increases the variance of hair cell orientation without affecting the average angle of their

rotation in the chicken basilar papilla[64].

The role of Wnt ligands in regulating hair cell orientation and PCP is less clear, as

Wntless deletion in the cochlear epithelium increases variance in hair cell orientation

without significantly affecting the average position, suggesting also a local instructive

role[15]. However, in the presence of the Vangl2-looptail mutation, Wntless deletion in the

cochlear epithelium leads to malrotation of hair cells towards the apex, suggesting that

Wnt ligands in the epithelium may serve as an instructive global cue[15]. In a similar vein,

inhibition of endogenous Wnts by inhibitors Sfrp1 or Wif1 in cultured mouse cochlea

directed hair cells to uniformly rotate unidirectionally[28]. In the current study, deletion of

Wls from both the cochlear epithelium and POM resulted in pronounced and uniform hair

cell rotation toward the apex, further supporting the notion that Wnt proteins provide an

instructive global cue for hair cell orientation. Together, we propose a model where hair

cell orientation and cochlear outgrowth are regulated by both core PCP proteins and Wnt

proteins from the epithelial and mesenchymal compartments in the developing cochlea.

In the model, core PCP proteins serve local instructive roles while Wnt proteins play a

more global role in guiding cell and tissue organization (Fig. 6).

In *Vangl2[Lp/Lp]* and other PCP mutants, hair cell malrotation is most severe in the

lateral-most most hair cells (i.e., abneural OHC3+), but the direction of malrotation is

random[10]. After broad Wntless deletion (Sox9-Wls cKO), hair cell malrotation is also most

severe in the lateral-most hair cells (Supplementary Table 1), but malrotation is almost

exclusively directed towards the apex. Hair cell malrotation towards the apex is also

observed in Emx2-Wls cKO in the presence of the *Vangl2[Lp/+]* allele[15], suggesting that

Wnts may act as a global instructive cue for hair cell orientation. It is possible that other

signaling pathways also contribute to cochlear patterning. For example, Wnt and FGF

signaling are both crucial in maintaining migratory polarity and gene expression domains

in the lateral line primordium in zebrafish[65] and regulating PCP during mouse limb

development[61]. Lastly, it is also possible that local shearing forces organize and pattern

hair cells and supporting cells differentially along the abneural-neural axis[32, 66].

<u>Wnt ligand-receptor interaction between epithelial and mesenchymal tissues</u>

Our study suggests that Wnt proteins from both the epithelium and POM regulate

the development of PCP in the cochlear epithelium. While Wnt ligands are classically

thought to act at short to medium ranges, they have been shown to travel remarkable

distances to mediate mesenchymal-epithelial crosstalk in other organ systems. For

example, Wnt9b in the Wolffian duct epithelium also regulates the development of

metanephric mesenchyme and mesenchymal to epithelial transitions[67]. Also,

mesenchymal cells secrete Wnt proteins to regulate stem cell proliferation and

differentiation in both the intestines and airway[68, 69, 70]. These studies illustrate the Wnt-

mediated mesenchymal-epithelial crosstalk during tissue development and homeostasis.

The distance and mechanisms by which Wnts are transported are likely context-

dependent[71, 72, 73], and warrants investigation given their significant roles in cochlear

development.

<u>Many Wnt members contribute to cochlear development</u>

Among the seven Wnt members expressed in the developing cochlea, our

computational cell-cell communication analysis predicts Wnt5a, Wnt7a, Wnt7b as the

most potent ligands to interact with corresponding receptors on hair cells. We and others

have found robust expression of Wnt5a mRNA in the cochlear epithelium flanking the

organ of Corti and the periotic mesenchyme from E12.5 to E16.5, higher in the base than

the apex[15, 30]. On the other hand, Wnt7a and Wnt7b mRNA are both broadly expressed

in the cochlear duct with no apparent apical-basal gradient. In agreement with prior

studies[15, 30], deletion of both Wnt5a and Wnt7b in the cochlear duct leads to significant

shortening of the cochlear duct and mild hair cell orientation defects. However, either

deletion alone has a minimal effect. Interestingly, deletion of both Wnt7a and Wnt7b in

the cochlear duct shortens the cochlea to the extent similar to that of deletion of

Wnt5a/Wnt7b, supporting the notion that Wnt proteins additively regulate cochlear

outgrowth. As Emx2-Wls cKO cochleae display more severe PCP defects than Emx2-

Wnt5a; Wnt7a; Wnt7b cochleae, we attribute this to either Wnt redundancy or incomplete

deletion in the latter model. Four other Wnt members (Wnt4, Wnt9a, Wnt11, and Wnt2b)

are also expressed at low levels in the developing cochlear duct[15] and may also contribute

to hair cell PCP signaling. According to our cell-cell communication computational

analysis, Wnt5a is deemed the most potent Wnt ligand; however, neither deletion of

Wnt5a in the POM (DermoCre-Wnt5a cKO) nor in the cochlear duct alone[15] causes

cochlear shortening or hair cell misorientation.

<u>Wnt signaling is not required for hair cell formation</u>

Our data demonstrate that even though we deleted Wnts or Wls either in the

cochlear epithelium or in both the cochlear epithelium and POM, hair cell formation still

occurs. This data suggests that secreted Wnts are not required for hair cell formation after

E10.5. Previous *in vivo* mouse studies deleting beta-catenin, the central mediator of

canonical Wnt signaling, show conflicting results. In mice where conditional deletion of

beta-catenin was performed using the CMV-CreER allele (tamoxifen at E11.5), Atoh1+

hair cells were absent at E14.5[21]. However, hair cells were observed at later time points

(E15.5-18.5) when other mouse lines were examined (Sox2-CreERT2 (tamoxifen at E12

or E13), Fgf20-Cre, Emx2-Cre)[21, 23, 74]. Together with the current study, these results

suggest that neither beta-catenin nor secreted Wnts are required for cochlear hair cell

formation.

<u>Summary</u>

In conclusion, secreted Wnts from the cochlear epithelium and POM are essential for

establishing planar polarity in cochlear hair cells and for cochlear outgrowth during

development. Additionally, multiple Wnt ligands, including Wnt5a, Wnt7a, and Wnt7b,

contribute to cochlear outgrowth, hair cell orientation and PCP protein localization, to

ensure precise cochlear length and hair cell patterning.

Materials and methods

Mice

The following mouse strains were used; *Wls-flox*[75](#012888, Jackson laboratory, Bar

Harbor, ME), *Wnt5a-flox*[42] (Strain #026626), *Wnt7b-flox*[44](#008467), Sox9-CreERT2[76]

(#018829) (All from the Jackson Laboratory), *Wnt7a-flox*[43], Emx2-Cre[36], and

Dermo1(Twist2)-Cre mice [77]. All the floxed mouse lines for the Wnts above mentioned

are maintained by homozygous sibling matings, while *Emx2$^{Cre/+}$* and *Dermo1$^{Cre/+}$* mouse

lines are maintained by breeding heterozygous mice. Tamoxifen was administered

intraperitoneally to dams carrying the Sox9-CreERT2 allele at a dose of 200 µg/g. For the

EdU assay, a single dose (50 mg/kg; Invitrogen) was given via intraperitoneal injection to

pregnant mice at E12.5, and then cochlear tissue was collected at E15.5. Embryos were

considered E0.5 at noon on the day when a plug was discovered. The protocol is

approved by the Animal Care and Use Committee of Stanford University School of

Medicine (#18606).

Genotyping

DNA templates were generated by incubating tissues (ear punch or tail clipping) in 180 µl

of 50 mM NaOH at 98°C for 1 h followed by the addition of 20 µl of 1 M TrisHCl (pH 8.0).

We used either GoTaq® Master Mixes (M712) or KAPA2G Fast HotStart®

ReadyMix™(KK5609) to amplify DNA fragments. The primers used were: Wls-flox (fwd,

5′-aggcttcgaacgtaactgacc-3′; rev, 5′-ctcagaactcccttcttgaagc-3′), Wnt5a-flox (fwd, 5′-

ggtgagggactggaagttgc-3′; rev, 5′-ggagcagatgtttattgccttc-3′), Wnt7a-flox (fwd, 5′-

cacagccacccctagagagctcaatt-3′; rev, 5′-atgctttgccaggga acaccc-3′), Wnt7b-flox (fwd1, 5′-

gccagaggccacttgtgtag-3′; fwd2, 5′-ggtagtccttcctgcccttt-3′; rev, 5′-gtgtgtcctggcctgatttt-3′),

Emx2-Cre and Sox9-CreERT2 (fwd, 5′-gagtaatagcgaccaatcatcaagcc-3′; rev1, 5′-

cgaacatcttcaggttctgcgg-3′; rev2, 5′-cttggaagcgatgacccagatatcgg-3′), and Dermo1-Cre (wt

fwd, 5′-cgtctcagctacgccttctc-3′; wt rev, 5′-tcacgaggagagatacactgg-3′; mt fwd 5′-

aacttcctctcccggagacc-3′; mt rev, 5′-ccggttattcaacttgcacc-3′).

Immunohistochemistry

The heads of E15.5 or E18.5 mouse embryos were halved and fixed with 4%

paraformaldehyde (PFA, Electron Microscopy Sciences) in phosphate-buffered saline

(PBS, pH 7.4) for 2 hours for the following markers: Myosin7a, Sox2, Celsr1, Phalloidin,

Ki67, and DAPI. Fixation with 10% Trichloroacetic acid (TCA, Sigma-Aldrich, 1 hour on

ice) in Milli-Q water was performed for these markers: Frizzled-6 (Fzd6), Dishevelled-2

(Dvl2), Vangl1, Vangl2, and ZO-1. After fixation, all tissues were stored in PBS at 4°C.

For whole-mount preparation of E15.5 and E18.5 mouse cochlea, the otic capsule

was dissected from the fixed tissues, followed by the removal of the lateral wall. For

cochlear sections, fixed otic capsule was treated with 10%, 20%, and 30% sucrose in

PBS at 4°C for 1 hour, 1 hour, and overnight, respectively. The samples were then

embedded in OCT for cryosectioning at a thickness of 10 µm. Both the whole-mount preps

and sections were incubated with 0.1% Triton X-100 in PBS for 1 hour at room

temperature. Following this, the samples were blocked with 5% donkey serum, 0.1%

Triton X-100, 1% bovine serum albumin (BSA, #BP1600–100, Thermo Fisher Scientific),

and 0.02% sodium azide (NaN3, #S2002–25G, Sigma-Aldrich,) in PBS (1 hour at RT).

Tissues were then incubated with primary antibodies overnight at 4°C. The primary

antibodies used were: Myosin7a (rabbit, 1:1000, 25-6790, Proteus Bioscience), Fzd6

(goat, 1:250, AF1526, R&D Systems), Dvl2 (rabbit, 1:250, 12037-1-AP, Proteintech,

Rosemont, IL), ZO-1 (mouse IgG1, 1:1000, 33-9100, Thermo Fisher Scientific), Ki67 (rat,

1:250, rat, 14-5698-82, Thermo Fisher Scientific), Vangl1 (rabbit, 1:1,000, HPA025235,

Sigma-Aldrich), Vangl2 (rabbit, 1:250, 21492-1-AP, Proteintech), and Celsr1 (rabbit,

1:3,000, gift from E. Fuchs, Rockefeller University). Following incubation with primary

antibodies, samples were washed with PBS three times (15 minutes each) and incubated

with the following secondary antibodies: Alexa Fluor 488, 546, and 647 (1:400 for all, A-

11055, A-10040, A-31571, Thermo Fisher Scientific) and DAPI (1:10,000, Invitrogen) at

room temperature for 2 hours. To label f-actin, phalloidin conjugated with Alexa546 (1:500,

A-22283, Thermo Fisher Scientific) was applied during secondary antibody incubation.

EdU detection was performed using the 647 Click-IT kit (C10340, Invitrogen). After

staining, cochlear epithelium was mounted in ProLong™ Gold Antifade Mountant

(P36930, Thermo Fisher Scientific) and coverslipped. Control and experimental samples

were always processed and imaged in parallel.

Fluorescent RNA *in situ* hybridization

The whole heads of E16.5 or E18.5 mouse embryos were halved and fixed with 4% PFA

in PBS for 16 to 24 hours at 4°C, treated with 10, 20, and 30% sucrose in PBS at 4°C (1

hour, 1 hour, and overnight, respectively). After that, tissues were embedded in OCT for

cryosections (10 µm). RNAScope and BaseScope assays were performed as follows:

tissue sections were hybridized with commercial probes from Advanced Cell Diagnostics

(ACDbio). These probes were specifically designed to detect deleted regions of the

following genes in that were deleted: Wls, Wnt5a, Wnt7a, and Wnt7b. The sections were

counterstained with anti-Myosin7a (rabbit, 1:1000, 25-6790, Proteus Bioscience) and

Sox2 (goat, 1:200, AF2018, R&D Systems) antibodies. The use of these antibodies was

done according to the manufacturer's instructions for fixed frozen sections. DAPI

(1:10,000, Invitrogen) was also used for nuclear staining. Briefly, sections were washed

in PBS for 5 mins once and then treated with H_2O_2 for 10 minutes. Next, the sections

were permeabilized using a target retrieval reagent (ACDbio) and proteinase prior to

hybridization. The following probes were used: DapB (Cat No. 310043), Polr2a (Cat No.

310451), Wls (Cat No. 405011), Wnt5a-2EJ (Cat No. 717251), Wnt7a-3zz-st-C1 (Cat No.

1277921-C1), and Wnt7b-2zz-st-C1 (Cat No. 1275011-C1) (all from Advanced Cell

Diagnostics). Control and cKO samples were processed and imaged in parallel.

Imaging and quantification

Immunofluorescence or *in situ* hybridization samples were imaged using a Zeiss LSM700

or LSM880 confocal microscope in conjunction with Zen software (Zeiss). Image analysis,

including measurements of cochlear length and hair cell orientation, was performed using

ImageJ software as described below (NIH).

Wholemount preparation and cochlear section were used to assess proliferation in

the cochlear epithelium and of periotic mesenchymal cells, respectively. For the former,

Sox2+ cells in the cochlear epithelium in the basal, middle, and apical regions were used

to calculate the percent Edu+ or Ki67+. For the POM, 20-50 DAPI+ cells from three areas

(lateral wall, perimodiolar, and basilar membrane) in the basal, middle, and apical regions

of the cochlea were used to calculate of the percent of Edu+ or Ki67+ cells. The basal,

middle, and apical regions were defined as 25, 50, and 70% length from the basal end of

the cochlear epithelium.

Hair cell orientation measurement

Whole mount preparation of the basal and middle turns of the right cochlea was used.

The orientation of individual hair cells was defined as previously described[15]. Specifically,

using ImageJ, a line was drawn parallel to the pillar cells (a reference line), and another

line was drawn from the base to the apex of bundles of individual hair cells (a hair bundle

line). Then, the angle between the hair bundle and reference lines was used to determine

the orientation of the hair cell (in degrees, counterclockwise direction is positive). Thirty

to 50 hair cells for each row from one cochlea from each mouse were measured (IHC,

OHC1, OHC2, OHC3, and additional rows more lateral to the OHC3 are denoted as

OHC3+). Three to six mice for each genotype were examined. Polar coordinate histogram

plots depicting hair cell orientation at each row and condition were generated and

statistics including the circular mean, mean resultant length, standard deviation, and

variance were calculated in using Python and the code is available on GitHub

(https://github.com/tahajanlab/published/tree/main/Kishimoto_et_al).

Inferred Cell-Cell Communication Analysis

To include both epithelial cells and mesenchymal cells in the cell-cell communication

analyses, we merged previously published E14.5/E16.5 epithelial-enriched data (GEO

Accession GSE137299[39]) with E13.5/E15.5 mesenchymal-enriched datasets (GEO

Accession GSE178931[38]; GEO Accession GSE217727[40]) in Seurat v5.3.0 merge function

resulting in combined raw count matrices[78]. The merged data was then log transformed,

normalized, and scaled for sequencing depth. Linear dimensionality reduction was then

performed via principal component analysis (PCA). For visualization, integration was

performed using Harmony[79] and non-linear dimensionality reduction was performed via

the uniform manifold approximation and projection (UMAP) algorithm. Cell types were

identified by canonical markers for mesenchymal cells (*Pou3f4*), prosensory cells (*Sox2*

and *Cdkn1b*), hair cells (*Pou4f3*), and the rest of the epithelial cells (*Epcam*). CellChat

was then used to infer cell-cell communication probability based on ligand and receptor

gene expression. Normalized count data was imported into CellChat v2.1.2 and the

CellChatDB mouse database was used[41]. Communication probability was calculated

based on the 10% truncated mean (average expression after discarding 10% from each

end of the data) for each ligand and receptor gene. The p-values were Benjamini-

Hochberg adjusted, and a cutoff of an adjusted p-value < 0.05 was considered statistically

significant. Chord plots were used to visualize directed ligand-receptor interactions for the

WNT and non-canonical WNT signaling pathways with a specific focus on hair cells and

prosensory cells as signal receivers. Bar graphs were also used to show relative

contributions of cell signaling from hair cells, prosensory cells, and epithelial cells.

Statistical analysis

For the comparison of cochlear length and hair cell counts, we performed either a two-

tailed unpaired Student's t-test, or a one-way ANOVA followed by Tukey's or Šídák's

multiple comparisons test. The Ki67- or EdU-positive ratios were evaluated using a two-

tailed unpaired Student's t-test. Student's t-test and ANOVA were conducted using

GraphPad Prism version 10 (GraphPad Software), with statistical significance set at P <

0.05. The N values indicate the number of hair cells in circular histograms and the number

of mice in other experiments.

Hair cell orientation analysis

The data comprised 11 individual experiment groups, each comparing the angle for hair

bundle orientation between control and cKO mice. Measurements were taken in a

patterned way for each mouse in each experiment group: from hair cells from four different

rows (IHC, OHC1, OHC2, and OHC3 rows) and at two cochlear turns (the basal and

middle turns), leading to 8 possible combinations.

A Bayesian mixed model analysis was run separately for each conditional KO

using the R package bpnreg v2.0.3[80] (Cremers, Jolien. n.d. Bpnreg: Bayesian Projected

Normal Regression Models for Circular Data. https://github.com/joliencremers/bpnreg)

with fixed effects treatment, i.e. control or conditional KO mice, and for cochlear turn

(basal and middle), and hair cells from four different rows (IHC, OHC1, OHC2, and OHC3

rows) along with random effects for the individual mice. Due to the limited number of mice

(n=6-8), the regression model was either a simple additive model or one containing the

interaction between cochlear turn and hair cell row. The bpnreg package respects the

circular nature of the angle measurements and yields samples from the posterior

distributions for the regression coefficients. Convergence of the underlying Markov Chain

was assessed via traceplots. The final model chosen for analysis (additive vs. interaction)

was based on Watanabe-Akaike criterion. Posterior samples from the selected model

were used to assess the difference in angles for each of the eight cochlear turn and hair

cell row combination between conditional KO and control groups. Each difference was

judged to be significant based on whether the posterior 95% credible intervals missed the

origin or not (Supplementary Table 1). The circular variance ratio was defined as the

circular variance of hair cell orientation in the KO group divided by the circular variance

of hair cell orientation in the control group.

Acknowledgments

We thank our lab members for insightful comments on the manuscript, W. Dong for

excellent technical support, E. Fuchs (Rockefeller University) for sharing the Celsr1

antibody. The *Wnt7a$^{flox/flox}$* mice were kindly shared by Jeremy Nathans (Johns Hopkins

University), *Emx2$^{Cre/+}$* mice by Angelika Doetzlhofer (Johns Hopkins University), and

Dermo1(Twist2)$^{Cre/+}$ mice by Sung-Ho Huh (University of Mississippi). This work was

supported by JSPS (Japan Society for the Promotion of Science) Overseas Research

Fellowships (IK), NIH UM1TR004921 (BN), K08DC019683, R21DC022058 (TAJ),

NIH/NIDCD Division of Intramural Research fund DC000094-01 (RH), R01DC021110,

R01DC016919 (AGC), the Stanford Initiative to Cure Hearing Loss.

Author contribution

I.K., E.L.S., W.D., and S.E.B. performed experiments and collected data; A.P.D., K.P.R.,

R.H., and T.A.J. performed computational analysis on single-cell data, B.N. and B.E., and

A.P.D. performed statistical analysis on hair cell orientation, I.K., S.E.B., and A.G.C.

designed experiments; I.K., A.P.D., K.P.R., B.N., and A.G.C. drafted the manuscript.

Code Availability

The following publicly available algorithms and software were used for data analysis

presented in this manuscript: Seurat (v5.3.0), CellChat (v2.1.2), and R (v4.5.2) package

bpnreg (v2.0.3).

Figure legend

Figure 1. Deletion of Wntless shortens the cochlea and causes mild hair cell

malrotation. a. Transgenic approach to delete Wntless in the cochlear epithelium. Cre

recombinase in the Emx2-Cre cochlear epithelium begins around embryonic day (E) 10.5.

b-c. Immunostaining and fluorescence *in situ* hybridization showing *Wls* mRNA in the

cochlear epithelium in controls (*Emx2$^{Cre/+}$; Wls$^{flox/+}$*), and no Wls mRNA signals in the

cochlear epithelium of *Emx2$^{Cre/+}$; Wls$^{flox/flox}$* (Emx2-Wls cKO) mice. **d.** Whole mount

preparation of E18.5 cochlea immunostained for Myo7A, showing shortened Emx2-Wls

cKO cochlea. **e.** Emx2-Wls cochleae are significantly shorter than controls at E18.5. **f.**

There are fewer Myo7A-positive hair cells per cochlea in Emx2-Wls cKO than controls.

g-h. Hair cells in Emx2-Wls cKO cochlea exhibit more variable orientations compared to

controls. **i.** Circular histograms depicting the distribution of hair bundle orientation in the

basal turn of control and Emx2-Wls cKO cochleae, with the latter showing significantly

larger circular variance and different average orientation in the IHC, OHC1, and OHC3+

rows. The OHC3+ row represents the third and more lateral rows of OHCs. **j.** Cartoon

illustrating the asymmetric localization of core PCP proteins in hair cells. Dvl2 is present

at the lateral pole of HCs, while Fzd6 is at the medial pole. Vangl1 and Vangl2 are present

at the lateral side of SCs, and Celsr1 is enriched at the junction between the medial side

of HCs and the lateral side of SCs. **k-l.** Immunostaining for Fzd6 and Dvl2, and ZO1 in

the basal turn of cochleae from control (k-k''') and Emx2-Wls cKO (l-l''') mice. Polarized

expression of Fzd6 and Dvl2 is observed at the medial and lateral poles of OHCs,

respectively, in the control (k''', white arrows) but not in the Emx2-Wls cKO cochlea (l''',

dashed white arrows). **m-p.** Immunostaining showing expression of Vangl2 and Celsr1 in

the medial pole of hair cells in control (m, o) and Emx2-Wls cKO cochleae (n, np). Scale

bar: 20 μm in (b, c), 200 μm in (d), 5 μm in (g, h), 10 μm in (m-p), 5 μm in (k-k'', l-l''), 2 μm

in (k''', l'''). Data are presented as mean ± S.D. and compared using an unpaired t-test. n,

number of hair cells measured. Circular mean and variance of hair cell orientation are

indicated with blue and red lines in circular histograms, respectively. A Bayesian mixed

model analysis used for comparison of circular mean of hair cell orientation between

control and cKO groups (see Materials and Methods section for more details).

*significantly different (for hair cell orientation analysis), **$p < 0.01$, ****$p < 0.0001$, ns =

not significant.

Figure 2. Cell-cell communication analysis of the embryonic cochlea. a-b. UMAP of

E13.5 periotic mesenchyme (POM) and E14.5 cochlear epithelium (early embryonic

cochlea dataset (a) and of E15.5 POM and E16.5 cochlear epithelium (late embryonic

cochlea dataset (b). **c-d.** Circle plot showing an inferred signaling network between Wnt

(canonical and non-canonical) ligands and receptors among cell groups from the early (c)

and late (d) developing cochlea datasets. The width of arrows is proportional to the

probability of communication between any two cell groups and autocrine signaling is

represented by a loop. In the early dataset, significant Wnt ligand-receptor interactions

are found among the four cell groups. Among the significant interactions, outgoing Wnt

ligand signals from hair cells are no longer significant in the late embryonic cochlea

dataset. Furthermore, the strength of interactions involving POM is robust in both datasets.

e-f. Chord plots for cell-cell communication from Wnt ligands and receptors in the early

(e) and late (f) developing cochlea datasets, with bands' width reflecting the scale of

contribution. **g-j.** Ranked Wnt ligands received by hair cells and prosensory cells in the

cochlear epithelium (g, h) or POM (i, j). In the early dataset, Wnt7b is the most potent Wnt

ligand emanating from the cochlear epithelium, followed by Wnt7a and Wnt5a (g). In the

late dataset, Wnt5a is the most potent Wnt, followed by Wnt7b and Wnt7a (h). Across

both datasets, Wnt5a is the most potent among Wnts derived from the POM followed by

Wnt5b (i, j). See Source Data file for details.

Figure 3. Deletion of multiple Wnts shortens the cochlea and causes mild PCP

defects. **a.** Schematic depicting the expression patterns of Wnt5a, Wnt7a, and Wnt7b in

the developing mouse cochlear epithelium. **B-C.** Whole mount preparation of E18.5

control and $Emx2^{Cre/+}; Wnt5a^{flox/flox}; Wnt7a^{flox/flox}; Wnt7b^{flox/flox}$ (Emx2-Wn5a; Wnt7a; Wnt7b

cKO) immunostained for Myo7A, with measurements showing that the latter are

significantly shorter. **d.** Emx2-Wnt5a; Wnt7a; Wnt7b cKO cochlea had significantly fewer

Myo7A-positive hair cells than littermate controls at E18.5. **e**. Measurements of the

lengths of cochleae in transgenic mice with deletion of one to multiple Wnts. The length

of Emx2-Wn5a; Wnt7a; Wnt7b cKO cochleae was significantly shorter than each of the

single or double cKO cochleae, and comparable to that of Emx2-Wls cKO cochleae. **f.**

Quantitative analysis of hair cells among control cochleae and cochleae with deletion of

one or multiple Wnts. Hair cell counts in Emx2-Wn5a; Wnt7a; Wnt7b cKO cochleae were

significantly lower than in each of single or Emx2-Wnt5a; Wnt7a cKO cochleae, and

comparable to those in Emx2-Wnt5a; Wnt7b, Emx2-Wnt7a; Wnt7b, or Emx2-Wls cKO

cochleae. Hair cell counts in Emx2-Wnt5a; Wnt7b cKO cochleae were significantly lower

than Emx2-Wnt5a cKO and Emx2-Wnt7b cKO cochleae ($p < 0.001$ and $p < 0.01$,

respectively. **g-h.** Whole mount preparation of E18.5 control (g, g') and Emx2-Wn5a;

Wnt7a; Wnt7b cochlea (h, h') immunostained for f-actin. **i.** Circular histograms illustrating

the distribution of hair bundle orientations in control and Emx2-Wn5a; Wnt7a; Wnt7b cKO

cochleae. No significant misorientation of the hair cells was detected (circular means and

variance). **j–k.** Polarized expression of Fzd6 and Dvl2 is observed at the medial and

lateral poles of OHCs, respectively, in the basal turn of both control (j''', white arrows) and

Emx2-Wn5a; Wnt7a; Wnt7b cKO cochleae (k''', white arrows). Scale bar: 200 μm in (b),

5 μm in (g-h, j-j'', k-k''), 2 μm in (j''', k'''). Data are presented as mean ± S.D. and analyzed

using an unpaired t-test (c, d) or one-way ANOVA followed by either Šídák's multiple

comparisons test (e-f). n, number of hair cells measured. Mean and S.D. of hair cell

orientation are indicated with blue and red lines in circular histograms, respectively. A

Bayesian mixed model analysis used for comparison of circular mean of hair cell

orientation between control and cKO groups. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p <$

0.0001, ns = not significant.

Figure 4. Wnt5a deletion from the periotic mesenchyme does not induce cochlear

PCP defects. a. Transgenic approach to delete Wnt5a in the POM. Cre recombinase in

the POM of Dermo1-Cre cochlea begins around embryonic day (E) 9.5. **d.** Whole mount

preparation of cochleae from E18.5 control and Dermo1-Wnt5a cKO embryos. **e-f.**

Cochlear lengths and hair cell counts are comparable between control and Dermo1-

Wnt5a cKO mice at E18.5. **g-h.** F-actin-stained cochleae from control (g) and Dermo1-

Wnt5a cKO (h) embryos at E18.5. **i** Circular histograms demonstrating comparable hair

bundle orientation in the basal turn of control and Dermo1-Wnt5a cKO cochleae. **j–j''', k–**

k'''. Polarized expression of Fzd6 and Dvl2 (white arrows) is observed at the medial and

lateral poles of OHCs, respectively, both in the control (j''') and the Dermo1-Wnt5a cKO

cochlea (k''').These results suggest that deletion of Wnt5a alone from the POM does not

disrupt cochlear PCP. CD cochlear duct; ST scala tympani; SV scala vestibule; SG spiral

ganglion. Scale bar: 20 µm in (b-c'), 5 µm in (g-h, j-k'''), 200 µm in (d). n, number of hair

cells measured. Mean and S.D. of hair cell orientation are indicated with blue and red

lines in circular histograms, respectively. A Bayesian mixed model analysis used for

comparison of circular mean of hair cell orientation between control and cKO groups. ns

= not significant.

Figure 5. Deletion of Wntless in the cochlear epithelium and periotic mesenchyme

causes severe cochlear shortening and PCP defects. a. Schematic of deleting

Wntless using the Sox9-CreERT2; Wntless-flox mice. Tamoxifen (0.2 mg/g) was

administered at E10.5 and E11.5 and cochleae were analyzed at E18.5. **b-c.**

Immunostaining and measurements of E18.5 control (*Sox9$^{CreERT2/+}$; Wls$^{flox/+}$*) and

Sox9$^{CreERT2/+}$; Wls$^{flox/flox}$ (Sox9-Wls cKO) cochleae, showing dramatic shortening of the

latter and significantly shorter than both control and Emx2-Wls cKO cochleae. **d.** Sox9-

Wls cKO cochleae had significantly fewer cells compared to control and Emx2-Wls cKO

cochleae. **e-f.** Whole mount preparation of basal turn of the E18.5 f-actin-stained cochlea

The Sox9-Wls cKO cochlea shows five to six rows of hair cells that are disorganized and

lack a separation between IHCs and OHCs. Notably, hair cells are mostly malrotated

toward the apex. **g** Circular histograms illustrating the distribution of hair bundle

orientation in control and Sox9-Wls cKO cochleae, which showed significantly larger

circular variance and different average orientation in each row relative to controls. **h-q.**

Immunostaining for core PCP proteins in the control and Sox9-Wls cKO cochleae for

Celsr1 (h, i), Vangl1 (j, k), Vangl2 (l, m), Fzd6 (n, o), and Dvl2 (p, q), showing a loss of

polarized expression of Celsr1, Vangl1, Vangl2, Fzd6, and Dvl2 in OHCs of Sox9-Wls

cKO cochleae. **r-s.** Sections of E15.5 control (r) and Sox9-Wls cKO (s) cochleae

illustrating Ki67- and EdU-positive cells in both the cochlear epithelium and periotic

mesenchyme. **t-u.** Quantitative assessment showing comparable proliferation in the

cochlear epithelium between the control and Sox9-Wls cKO cochleae. CD, cochlear duct;

LW, POM near the lateral wall; BM, POM near the basilar membrane; PM, perimodiolar

POM. Scale bar: 200 µm in (b), 5 µm in (e-f, h-q), 40 µm in (r-s). Data are presented as

mean ± S.D. and analyzed using one-way ANOVA followed by Tukey's multiple

comparisons test (c-d) and an unpaired t-test (t-u). n, number of hair cells measured.

Mean and S.D. of hair cell orientation are indicated with blue and red lines in circular

histograms, respectively. A Bayesian mixed model analysis used for comparison of

circular mean of hair cell orientation between control and cKO groups. *significantly

different (for hair cell orientation analysis), **$p < 0.01$, ****$p < 0.0001$, ns = not significant.

Figure 6. Working model of epithelial and mesenchymal Wnts coordinating to

regulate cochlear elongation, hair cell orientation, and PCP signaling. When only

mesenchymal Wnts are present, cochlear length is mildly shortened and hair cells mildly

misoriented and most core PCP proteins are present. In the absence of both epithelial

and mesenchymal Wnts, cochlear duct is severely shortened, and hair cells severely

malrotated toward the apex.

Supplementary Figure 1. In situ hybridization and hair bundle orientation in the

Emx2-Wls cKO cochlea. a-b. Negative (DapB) and positive (Polr2a) controls for *in situ*

hybridization in E16.5 cochleae. **c-h.** RNA scope *in situ* hybridization of Wls mRNA in the

basal, middle, and apical turn of the control (c-c'', e-e'', g-g'') and Emx2-Wls cKO (d-d'', f-

f'', h-h'') cochleae. Magnified views (white boxes) of the cochlear epithelium and POM are

shown in (c'-h') and (c''-h''), respectively. In Emx2-Wls cKO cochleae, Wls mRNA is

almost undetectable in the cochlear epithelium, while that in the POM remain comparable

to those in control cochleae. **i-j.** Whole mount preparation of the Emx2-Wls cKO and

control cochlea stained for f-actin. In the Emx2-Wls cKO cochlea, hair cells exhibit more

variable orientations compared to controls. **k.** Circular histograms depicting the

distribution of hair bundle orientation both control and Emx2-Wls cKO cochleae, showing

significantly large variance and the average rotation for the IHC, OHC1, and OHC3 rows

in the Emx2-Wls cKO cochleae. CD, cochlear duct; ST, scala tympani; SV, scala vestibule;

SG, spiral ganglion. Scale bar: 20 µm in (a-h''), 5 µm in (I, j). n, number of hair cells

measured. Circular mean and variance of hair cell orientation are indicated with blue and

red lines in circular histograms, respectively. A Bayesian mixed model analysis used for

comparison of circular mean of hair cell orientation between control and cKO groups.

*significantly different (for hair cell orientation analysis), ns = not significant.

Supplementary Figure 2. Expression of Wnt5a, Wnt7a, and Wnt7b mRNA in the

control and conditional knockout cochlea. a-d. *Wnt5a* mRNA is robustly expressed in

the epithelium and periotic mesenchyme of control cochlea (a, c) and is almost

undetectable in the epithelium of Emx2-Wnt5a cKO (b, d) cochleae. **e-h.** Wnt7a mRNA is

robustly expressed in the epithelium of the control cochlea (e, g) and is barely detectable

in the Emx2-Wnt7a cKO (f, h) cochleae. **i-l.** Wnt7b mRNA is robustly expressed in the

epithelium of the control cochlear (l, k) and is almost undetectable in the Emx2-Wnt7b

cKO (j, l) cochleae. Scale bar: 20 µm.

Supplementary Figure 3. Cochlear length and core PCP protein expression in

Wnt5a, Wnt7a, and Wnt7b cKO cochlea. a-b, f-g, k-m. Comparable lengths between

Myo7a-marked controls and cochleae from Emx2-Wnt5a (a), Emx2-Wnt7a (f), and Emx2-

Wnt7b cKO (k) mice at E18.5. **c, h, m** Hair cell counts show no significant differences

between Emx2-Wnt5a (b) and Emx2-Wnt7a cKO (g) mice and their littermate controls,

whereas those from Emx2-Wnt7b cKO (l) mice are slightly reduced compared to controls

at E18.5. **d-e''', i-j''', n-o'''** Polarized expression of Fzd6 and Dvl2 (white arrows) at the

medial and lateral poles of OHCs, respectively, in control cochleae and cKO cochleae for

Wnt5a (d''' and e''',) , Wnt7a (i''' and j'''), and Wnt7b (n''' and o'''). Data are presented as

mean ± S.D. and compared using an unpaired t-test. Scale bars: 200 µm in (a, f, k); 5 µm

in (d-d", e-e", i-i", j-j", n-n", o-o"); 2 μm in (d''', e''', i''', j''', n''', o'''). Data are presented as

mean ± S.D. and compared using an unpaired t-test. *p < 0.05, ns = not significant.

Supplementary Figure 4. Hair bundle orientation of the cochlea from Emx2-Wnt5a,

Emx2-Wnt7a, and Emx2-Wnt7b cKO mice. a-d, g-j, m-p. F-actin-stained cochleae from

Emx2-Wnt5a (a-d), Emx2-Wnt7a (g-j), and Emx2-Wnt7b cKO (m-p) mice at E18.5. **e, f,**

k, l, q, r. Circular histograms demonstrating no significant variance or average deviation

hair bundle orientation in Emx2-Wnt5a (e, f) and Emx2-Wnt7a (k, l) cKO cochleae. and

For Emx2-Wnt7b cKO (q, r) cochlea, there was no significant variance, but significantly

different circular means of hair cell orientation compared to controls was detected for the

IHC, OHC1, and OHC2 rows. Scale bar: 5 μm. n, number of hair cells measured. Circular

mean and variance of hair cell orientation are indicated with blue and red lines in circular

histograms, respectively. A Bayesian mixed model analysis used for comparison of

circular mean of hair cell orientation between control and cKO groups. *significantly

different (for hair cell orientation analysis), ns = not significant.

Supplementary Figure 5. Cochlear length and core PCP protein expression after

deletion of Wnts. a-b, f-g, k-l. Whole mount preparation of cochleae from Emx2-Wnt5a;

Wnt7a (a-b), Emx2-Wnt5a; Wnt7b (f-g), and Emx2-Wnt7a; Wnt7b cKO (k-l) mice and their

littermate controls at E18.5. The lengths of Emx2-Wnt5a; Wnt7b and Emx2-Wnt7a; Wnt7b

cKO, but not Emx2-Wnt5a; Wnt7a cKO, cochleae were significantly shorter than those of

controls. **c, h, m** Hair cell counts show no significant differences between Emx2-Wnt5a;

Wnt7a (c) and Emx2-Wnt7a; Wnt7b cKO (m) cochleae and their littermate controls,

whereas those from Emx2-Wnt5a; Wnt7b cKO (h) cochleae are slightly reduced

compared to controls at E18.5. **d-e''', i-j''', n-o'''** Polarized expression of Fzd6 and Dvl2

is observed at the medial and lateral poles of OHCs, respectively, in control cochlea and

cKO cochleae for Wnt5a and Wnt7a (d''' and e''', white arrows), Wnt5a and Wnt7b (i''' and

j''', white arrows), and Wnt7a and Wnt7b (n''' and o''', white arrows). Data are presented

as mean $\pm$ S.D. and compared using an unpaired t-test. Scale bars: 200 μm in (a, f, k);

5 μm in (d-d'', e-e'', i-i'', j-j'', n-n'', o-o''); 2 μm in (d''', e''', i''', j''', n''', o'''). **p < 0.01, ***p <

0.001, ****p < 0.0001, ns = not significant.

Supplementary Figure 6. Hair bundle orientation after deletion of Wnt members the

cochlear epithelium. a-d, g-j, m-p. F-actin-stained cochleae from Emx2-Wnt5a; Wnt7a

(a-d), Emx2-Wnt5a; Wnt7b (g-j), and Emx2-Wnt7a; Wnt7b cKO (m-p) mice at E18.5. **e, f,**

k, l, q, r. Circular histograms demonstrating the distribution of hair bundle orientation in

the cochlea from Emx2-Wnt5a; Wnt7a (e, f), Emx2-Wnt5a; Wnt7b (k, l), and Emx2-Wnt7a;

Wnt7b cKO (q, r) mice at E18.5. Emx2-Wnt5a; Wnt7a cKO cochleae show normal hair

cell orientation. Emx2-Wnt5a; Wnt7b cKO cochleae exhibit more variable orientations of

hair cells compared to controls (g-j), confirmed by significantly larger circular variance in

hair cell orientation, but the average orientation was not significantly different from

controls, with the exception of the IHC row of the basal turn. Similarly, hair cell variance

in Emx2-Wnt7a; Wnt7b cKO cochleae was not significantly different from controls, with

no significantly different circular mean except for the OHC3 row. Scale bar: 5 μm. n,

number of hair cells measured. Circular mean and variance of hair cell orientation are

indicated with blue and red lines in circular histograms, respectively. A Bayesian mixed

model analysis used for comparison of circular mean of hair cell orientation between

control and cKO groups. *significantly different, ns = not significant.

Supplementary Figure 7. Hair bundle orientation after deletion of three Wnt

members in the cochlear epithelium. Whole mount preparation of f-actin-stained

cochlea from control (a, a') and Emx2-Wn5a; Wnt7a; Wnt7b cKO (b, b') mice. **i** Circular

histograms illustrating the distribution of hair bundle orientations in the middle turn of both

control and Emx2-Wn5a; Wnt7a; Wnt7b cKO cochleae, showing no significant difference

in circular means and variance from controls. n, number of hair cells measured. Circular

mean and variance of hair cell orientation are indicated with blue and red lines in circular

histograms, respectively. A Bayesian mixed model analysis used for comparison of

circular mean of hair cell orientation between control and cKO groups. ns = not significant.

Supplementary Figure 8. In situ hybridization of Wnt5a mRNA in the Dermo1-Wnt5a

cKO cochlea. Wnt5a mRNA is robustly expressed in the cochlear duct and periotic

mesenchyme of the control cochleae (*Dermo1$^{Cre/+}$; Wnt5a$^{flox/+}$*) (a-a', c-c'), and notably

absent in the periotic mesenchyme of *Dermo1$^{Cre/+}$; Wnt5a$^{flox/flox}$* cochleae (Dermo1-

Wnt5a) (b-b', d-d') at E16.5. Wnt5a mRNA remain present in the cochlear epithelium and

spiral ganglion (b-b', d-d'), indicating effective deletion of Wnt5a mRNA specifically from

the POM. **e-g.** F-actin-stained middle turn of cochleae from control (e) and Dermo1-

Wnt5a cKO (f) embryos at E18.5. **i** Circular histograms demonstrating comparable hair

bundle orientation in the middle turn of control and Dermo1-Wnt5a cKO cochleae. CD

cochlear duct; ST scala tympani; SV scala vestibule; SG spiral ganglion. Scale bar: 20 µm

in (a-h"), 5 µm in (I, j). A Bayesian mixed model analysis used for comparison of circular

mean of hair cell orientation between control and cKO groups. ns = not significant.

Supplementary Figure 9. Deleting Wntless in the cochlear epithelium and periotic

mesenchyme at E11.5/12.5 causes mild PCP defects. a. Schematic of broadly deleting

Wntless using the Sox9-CreERT2; Wntless-flox mice. Tamoxifen (0.2 mg/g) was

administered at E11.5 and E12.5 and cochleae were analyzed at E18.5. **b.** Sox9-Wls cKO

cochlea is significantly shorter than controls. **c-d.** Length and hair cell counts of Sox9-Wls

cKO cochleae with tamoxifen administered at E11.5-12.5 were significantly lower than

controls, but higher than Sox9-Wls cKO cochleae with tamoxifen administered at E10.5-

11.5. **e-h.** Whole mount preparation of f-actin-stained cochlea showing hair cells

malrotated towards the apex in the Sox9-Wls cKO cochlea (arrows). **i-j.** Circular

histograms demonstrating the distribution of hair bundle orientation in control and Sox9-

Wls cKO cochleae. Significantly larger circular variance and different circular means of

hair cell orientation are detected in the basal and middle turns of Sox9-Wls cKO cochleae.

k-s. Control and Sox9-Wls cKO cochleae showed polarized expression of core PCP

proteins in (Fzd6 (k-l), Celsr1 (m-n), Vangl1 (p-q), and Vangl2 (r-s)). Scale bar: 200 µm

in (b), 5 µm in (e-h, k-s). Data are presented as mean ± S.D. and analyzed using one-way

ANOVA followed by Tukey's multiple comparisons test. n, number of hair cells measured.

Circular mean and variance of hair cell orientation are indicated with blue and red lines in

circular histograms, respectively. A Bayesian mixed model analysis used for comparison

of circular mean of hair cell orientation between control and cKO groups. *significantly

different (for hair cell orientation analysis), **p < 0.01, ***p < 0.001, ****p < 0.0001, ns =

not significant.

1004

1005   **Supplementary Figure 10. Loss of Wls mRNA expression in Sox9-Wls cKO**

1006   **cochleae. a-f'** In situ hybridization showing robust Wls mRNA expression in all turns of

1007   control (a-a', c-c', e-e') cochleae, and loss of expression in Sox9-Wls cKO (b-b', d-d', f-f')

1008   cochleae at E18.5. Scale bar: 50 μm. CD cochlear duct; ST scala tympani; SV scala

1009   vestibule; SG spiral ganglion.

1010

Supplementary Figure 11. Hair bundle orientation and proliferation in Sox9-Wls

cochlea after Wntless at E10.5 and E11.5. a-b'. F-actin-stained cochlea from Sox9-Wls

cKO (b, b') embryos showing five to six rows of hair cells, disorganized hair cells lacking

a clear separation between IHCs and OHCs, and malrotation of hair cells toward the apex.

c. Circular histograms illustrating the distribution of hair bundle orientation in IHCs and

OHCs in the middle turn of control and Sox9-Wls cKO cochleae. Sox9-Wls cKO cochleae

have significantly larger circular variance for combined IHC and OHCs and significantly

different circular means of hair cell orientation for each of IHC and OHC rows compared

to controls.   **d-e.** Whole-mount preparation of E15.5 cochlea labeled with EdU or

immunostained for Ki67, showing comparable degree of proliferation of Sox2-positive

cells in control and Sox9-Wls cKO organs. **f-g.** Proliferation of POM cells in the POM was

not significantly different between control and Sox9-Wls cKO cochleae at E15.5. Data are

presented as mean ± S.D. and analyzed using an unpaired t-test (d-e). n, number of hair

cells measured. Circular mean and variance of hair cell orientation are indicated with blue

and red lines in circular histograms, respectively. A Bayesian mixed model analysis used

for comparison of circular mean of hair cell orientation between control and cKO groups.

*significantly different (for hair cell orientation analysis), ns = not significant.

1028

References

1. Cantos R, Cole LK, Acampora D, Simeone A, Wu DK. Patterning of the mammalian
cochlea. *Proc Natl Acad Sci U S A* **97**, 11707-11713 (2000).

2. Driver EC, Kelley MW. Development of the cochlea. *Development* **147**, (2020).

3. Ruben RJ. Development of the inner ear of the mouse: a radioautographic study
of terminal mitoses. *Acta Otolaryngol*, Suppl 220:221-244 (1967).

4. Chen P, Segil N. p27(Kip1) links cell proliferation to morphogenesis in the
developing organ of Corti. *Development* **126**, 1581-1590 (1999).

5. Tarchini B, Lu X. New insights into regulation and function of planar polarity in the
inner ear. *Neurosci Lett* **709**, 134373 (2019).

6. Kelly M, Chen P. Shaping the mammalian auditory sensory organ by the planar
cell polarity pathway. *Int J Dev Biol* **51**, 535-547 (2007).

7. Ross AJ*, et al.* Disruption of Bardet-Biedl syndrome ciliary proteins perturbs planar
cell polarity in vertebrates. *Nat Genet* **37**, 1135-1140 (2005).

8. Jagger D*, et al.* Alstrom Syndrome protein ALMS1 localizes to basal bodies of
cochlear hair cells and regulates cilium-dependent planar cell polarity. *Hum Mol*
Genet **20**, 466-481 (2011).

9. Marshall JD, Maffei P, Collin GB, Naggert JK. Alstrom syndrome: genetics and
clinical overview. *Curr Genomics* **12**, 225-235 (2011).

10. Montcouquiol M, Rachel RA, Lanford PJ, Copeland NG, Jenkins NA, Kelley MW.
Identification of Vangl2 and Scrb1 as planar polarity genes in mammals. *Nature*
423, 173-177 (2003).

11. Montcouquiol M*, et al.* Asymmetric localization of Vangl2 and Fz3 indicate novel
mechanisms for planar cell polarity in mammals. *J Neurosci* **26**, 5265-5275 (2006).

12. Wang Y, Guo N, Nathans J. The role of Frizzled3 and Frizzled6 in neural tube
closure and in the planar polarity of inner-ear sensory hair cells. *J Neurosci* **26**,
2147-2156 (2006).

13. Etheridge SL*, et al.* Murine dishevelled 3 functions in redundant pathways with
dishevelled 1 and 2 in normal cardiac outflow tract, cochlea, and neural tube
development. *PLoS genetics* **4**, e1000259 (2008).

14. Wang J, *et al.* Regulation of polarized extension and planar cell polarity in the cochlea by the vertebrate PCP pathway. *Nat Genet* **37**, 980-985 (2005).

15. Najarro EH, Huang J, Jacobo A, Quiruz LA, Grillet N, Cheng AG. Dual regulation of planar polarization by secreted Wnts and Vangl2 in the developing mouse cochlea. *Development* **147**, (2020).

16. Torban E, *et al.* Genetic interaction between members of the Vangl family causes neural tube defects in mice. *Proc Natl Acad Sci U S A* **105**, 3449-3454 (2008).

17. Curtin JA, *et al.* Mutation of Celsr1 disrupts planar polarity of inner ear hair cells and causes severe neural tube defects in the mouse. *Curr Biol* **13**, 1129-1133 (2003).

18. Steinhart Z, Angers S. Wnt signaling in development and tissue homeostasis. *Development* **145**, (2018).

19. Xue C, Chu Q, Shi Q, Zeng Y, Lu J, Li L. Wnt signaling pathways in biology and disease: mechanisms and therapeutic advances. *Signal Transduct Target Ther* **10**, 106 (2025).

20. van Amerongen R. Alternative Wnt pathways and receptors. *Cold Spring Harb Perspect Biol* **4**, (2012).

21. Shi F, Hu L, Jacques BE, Mulvaney JF, Dabdoub A, Edge AS. beta-Catenin is required for hair-cell differentiation in the cochlea. *J Neurosci* **34**, 6470-6479 (2014).

22. Jansson L, *et al.* beta-Catenin is required for radial cell patterning and identity in the developing mouse cochlea. *Proc Natl Acad Sci U S A* **116**, 21054-21060 (2019).

23. Ebeid M, Kishimoto I, Roy P, Zaidi MAA, Cheng AG, Huh SH. beta-Catenin transcriptional activity is required for establishment of inner pillar cell identity during cochlear development. *PLoS Genet* **19**, e1010925 (2023).

24. Gao B. Wnt regulation of planar cell polarity (PCP). *Curr Top Dev Biol* **101**, 263-295 (2012).

25. Butler MT, Wallingford JB. Planar cell polarity in development and disease. *Nat Rev Mol Cell Biol* **18**, 375-388 (2017).

26. Munnamalai V, Fekete DM. Notch-Wnt-Bmp crosstalk regulates radial patterning in the mouse cochlea in a spatiotemporal manner. *Development* **143**, 4003-4015 (2016).

27. Noda T, Oki S, Kitajima K, Harada T, Komune S, Meno C. Restriction of Wnt signaling in the dorsal otocyst determines semicircular canal formation in the mouse embryo. *Dev Biol* **362**, 83-93 (2012).

28. Dabdoub A, *et al.* Wnt signaling mediates reorientation of outer hair cell stereociliary bundles in the mammalian cochlea. *Development* **130**, 2375-2384 (2003).

1097   29.   Qian D, *et al.* Wnt5a functions in planar cell polarity regulation in mice. *Dev Biol*
1098   **306**, 121-133 (2007).

1099   30.   Xie N, *et al.* Wnt7b acts in concert with Wnt5a to regulate tissue elongation and
1100   planar cell polarity via noncanonical Wnt signaling. *Proc Natl Acad Sci U S A* **121**,
1101   e2405217121 (2024).

1102   31.   Landin Malt A, Hogan AK, Smith CD, Madani MS, Lu X. Wnts regulate planar cell
1103   polarity via heterotrimeric G protein and PI3K signaling. *J Cell Biol* **219**, (2020).

1104   32.   Deans MR. Conserved and Divergent Principles of Planar Polarity Revealed by
1105   Hair Cell Development and Function. *Front Neurosci* **15**, 742391 (2021).

1106   33.   Kindt KS, Tarchini B. Not a hair out of place: Polarizing and orienting sensory hair
1107   cells. *Curr Top Dev Biol* **165**, 85-124 (2025).

1108   34.   Banziger C, Soldini D, Schutt C, Zipperlen P, Hausmann G, Basler K. Wntless, a
1109   conserved membrane protein dedicated to the secretion of Wnt proteins from
1110   signaling cells. *Cell* **125**, 509-522 (2006).

1111   35.   Wolf L, Boutros M. The role of Evi/Wntless in exporting Wnt proteins. *Development*
1112   **150**, (2023).

1113   36.   Ono K, *et al.* FGFR1-Frs2/3 signalling maintains sensory progenitors during inner
1114   ear hair cell formation. *PLoS Genet* **10**, e1004118 (2014).

1115   37.   Copley CO, Duncan JS, Liu C, Cheng H, Deans MR. Postnatal refinement of
1116   auditory hair cell planar polarity deficits occurs in the absence of Vangl2. *J*
1117   *Neurosci* **33**, 14001-14016 (2013).

1118   38.   Sun Y, *et al.* Single-cell transcriptomic landscapes of the otic neuronal lineage at
1119   multiple early embryonic ages. *Cell Rep* **38**, 110542 (2022).

1120   39.   Kolla L, *et al.* Characterization of the development of the mouse cochlear
1121   epithelium at the single cell level. *Nat Commun* **11**, 2389 (2020).

1122   40.   Rose KP, *et al.* Spatially distinct otic mesenchyme cells show molecular and
1123   functional heterogeneity patterns before hearing onset. *iScience* **26**, 107769
1124   (2023).

1125   41.   Jin S, *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat*
1126   *Commun* **12**, 1088 (2021).

1127   42.   Ryu YK, Collins SE, Ho HY, Zhao H, Kuruvilla R. An autocrine Wnt5a-Ror signaling
1128   loop mediates sympathetic target innervation. *Dev Biol* **377**, 79-89 (2013).

1129   43.   Dunlap KA, *et al.* Postnatal deletion of Wnt7a inhibits uterine gland morphogenesis
1130   and compromises adult fertility in mice. *Biol Reprod* **85**, 386-396 (2011).

1131   44.   Rajagopal J, *et al.* Wnt7b stimulates embryonic lung growth by coordinately
1132   increasing the replication of epithelium and mesenchyme. *Development* **135**,
1133   1625-1634 (2008).

1134

45.   Riccomagno MM, Takada S, Epstein DJ. Wnt-dependent regulation of inner ear morphogenesis is balanced by the opposing and supporting roles of Shh. *Genes Dev* **19**, 1612-1623 (2005).

46.   Liu W, Li L, Li G, Garritano F, Shanske A, Frenz DA. Coordinated molecular control of otic capsule differentiation: functional role of Wnt5a signaling and opposition by sfrp3 activity. *Growth Factors* **26**, 343-354 (2008).

47.   Sosic D, Richardson JA, Yu K, Ornitz DM, Olson EN. Twist regulates cytokine gene expression through a negative feedback loop that represses NF-kappaB activity. *Cell* **112**, 169-180 (2003).

48.   Huh SH, Warchol ME, Ornitz DM. Cochlear progenitor number is controlled through mesenchymal FGF receptor signaling. *Elife* **4**, (2015).

49.   Cornett B, Snowball J, Varisco BM, Lang R, Whitsett J, Sinner D. Wntless is required for peripheral lung differentiation and pulmonary vascular development. *Dev Biol* **379**, 38-52 (2013).

50.   Mak AC, Szeto IY, Fritzsch B, Cheah KS. Differential and overlapping expression pattern of SOX2 and SOX9 in inner ear development. *Gene Expr Patterns* **9**, 444-453 (2009).

51.   Veithen M*, et al.* Sox9 Inhibits Cochlear Hair Cell Fate by Upregulating Hey1 and HeyL Antagonists of Atoh1. *Cells* **12**, (2023).

52.   Sun MR, Steward AC, Sweet EA, Martin AA, Lipinski RJ. Developmental malformations resulting from high-dose maternal tamoxifen exposure in the mouse. *PLoS One* **16**, e0256299 (2021).

53.   Ved N, Curran A, Ashcroft FM, Sparrow DB. Tamoxifen administration in pregnant mice can be deleterious to both mother and embryo. *Lab Anim* **53**, 630-633 (2019).

54.   Montcouquiol M, Kelley MW. Development and Patterning of the Cochlea: From Convergent Extension to Planar Polarity. *Cold Spring Harb Perspect Med* **10**, (2020).

55.   Sutherland A, Keller R, Lesko A. Convergent extension in mammalian morphogenesis. *Semin Cell Dev Biol* **100**, 199-211 (2020).

56.   Wu J, Roman AC, Carvajal-Gonzalez JM, Mlodzik M. Wg and Wnt4 provide long-range directional input to planar cell polarity orientation in Drosophila. *Nat Cell Biol* **15**, 1045-1055 (2013).

57.   Ewen-Campen B, Comyn T, Vogt E, Perrimon N. No Evidence that Wnt Ligands Are Required for Planar Cell Polarity in Drosophila. *Cell Rep* **32**, 108121 (2020).

58.   Yu JJS*, et al.* Frizzled-Dependent Planar Cell Polarity without Secreted Wnt Ligands. *Dev Cell* **54**, 583-592 e585 (2020).

59. Gros J, Serralbo O, Marcelle C. WNT11 acts as a directional cue to organize the
elongation of early muscle fibres. *Nature* **457**, 589-593 (2009).

60. Gao B*, et al.* Wnt signaling gradients establish planar cell polarity by inducing
Vangl2 phosphorylation through Ror2. *Dev Cell* **20**, 163-176 (2011).

61. Gao B*, et al.* Coordinated directional outgrowth and pattern formation by
integration of Wnt5a and Fgf signaling in planar cell polarity. *Development* **145**,
(2018).

62. Minegishi K*, et al.* A Wnt5 Activity Asymmetry and Intercellular Signaling via PCP
Proteins Polarize Node Cells for Left-Right Symmetry Breaking. *Dev Cell* **40**, 439-
452 e434 (2017).

63. Yang W*, et al.* Wnt-induced Vangl2 phosphorylation is dose-dependently required
for planar cell polarity in mammalian development. *Cell Res* **27**, 1466-1484 (2017).

64. Sienknecht UJ, Anderson BK, Parodi RM, Fantetti KN, Fekete DM. Non-cell-
autonomous planar cell polarity propagation in the auditory sensory epithelium of
vertebrates. *Dev Biol* **352**, 27-39 (2011).

65. Aman A, Piotrowski T. Wnt/beta-catenin and Fgf signaling control collective cell
migration by restricting chemokine receptor expression. *Dev Cell* **15**, 749-761
(2008).

66. Cohen R*, et al.* Mechanical forces drive ordered patterning of hair cells in the
mammalian inner ear. *Nat Commun* **11**, 5137 (2020).

67. Carroll TJ, Park JS, Hayashi S, Majumdar A, McMahon AP. Wnt9b plays a central
role in the regulation of mesenchymal to epithelial transitions underlying
organogenesis of the mammalian urogenital system. *Dev Cell* **9**, 283-292 (2005).

68. Shoshkes-Carmel M*, et al.* Subepithelial telocytes are an important source of Wnts
that supports intestinal crypts. *Nature* **557**, 242-246 (2018).

69. Aros CJ*, et al.* Distinct Spatiotemporally Dynamic Wnt-Secreting Niches Regulate
Proximal Airway Regeneration and Aging. *Cell Stem Cell* **27**, 413-429 e414 (2020).

70. Maimets M*, et al.* Mesenchymal-epithelial crosstalk shapes intestinal
regionalisation via Wnt and Shh signalling. *Nat Commun* **13**, 715 (2022).

71. Stewart RA, Ramakrishnan AB, Cadigan KM. Diffusion and function of Wnt ligands.
PLoS Genet **15**, e1008154 (2019).

72. Mii Y*, et al.* Quantitative analyses reveal extracellular dynamics of Wnt ligands in
Xenopus embryos. *Elife* **10**, (2021).

73. Farin HF*, et al.* Visualization of a short-range Wnt gradient in the intestinal stem-
cell niche. *Nature* **530**, 340-343 (2016).

74. Jansson L, Kim GS, Cheng AG. Making sense of Wnt signaling-linking hair cell
regeneration to development. *Front Cell Neurosci* **9**, 66 (2015).

75.   Carpenter AC, Rao S, Wells JM, Campbell K, Lang RA. Generation of mice with a
1209   conditional null allele for Wntless. *Genesis* **48**, 554-558 (2010).

1210   76.   Kopp JL*, et al.* Sox9+ ductal cells are multipotent progenitors throughout
1211   development but do not produce new endocrine cells in the normal or injured adult
1212   pancreas. *Development* **138**, 653-665 (2011).

1213   77.   Yu K*, et al.* Conditional inactivation of FGF receptor 2 reveals an essential role for
1214   FGF signaling in the regulation of osteoblast function and bone growth.
1215   *Development* **130**, 3063-3074 (2003).

1216   78.   Hao Y*, et al.* Dictionary learning for integrative, multimodal and scalable single-cell
1217   analysis. *Nat Biotechnol* **42**, 293-304 (2024).

1218   79.   Korsunsky I*, et al.* Fast, sensitive and accurate integration of single-cell data with
1219   Harmony. *Nat Methods* **16**, 1289-1296 (2019).

1220   80.   Cremers J, Mulder KT, Klugkist I. Circular interpretation of regression coefficients.
1221   *Br J Math Stat Psychol* **71**, 75-95 (2018).

1222

1223

Figure 1

a Emx2$^{Cre/+}$; Wntless$^{fl/+}$
X
Wntless$^{fl/fl}$

Cre onset
E10.5

Harvest
E16.5 E18.5

Control
(Emx2$^{Cre/+}$; Wls$^{flox/+}$)

Emx2-Wls cKO
(Emx2$^{Cre/+}$; Wls$^{flox/flox}$)

Wntless mRNA Myosin7a Sox2 DAPI

b **c**

20 μm

E16.5 Base

d Control   Emx2-Wls cKO

E18.5   Myo7A

200 μm

e

Cochlear length (μm) ****

Control   Emx2-Wls cKO

f

Myo7a⁺ HC number per cochlea **

Control   Emx2-Wls cKO

g Control Apex ↔ Base

Base E18.5

h Emx2-Wls cKO F-actin

5 μm

g'

h' Apex ↔ Base

i IHC OHC1 OHC2 OHC3⁺

Control

n=110 n=111 n=110 n=109

Emx2-Wls cKO

n=148 n=130 n=128 n=111

Mean * * * ns Base

j

Lateral

Dvl2

Fzd6

Celsr1

Vangl2

Medial

Fzd6 Dvl2 ZO-1   Vangl2   Celsr1

k Fzd6 **k'** Dvl2 **k''** **k'''** **m** **o**

Control

E18.5

l **l'** **l''** **l'''** **n** **p**

Emx2-Wls cKO

Base 5 μm 2 μm Base 10 μm

Figure 2

a E13.5/14.5 Cochlea

POM
Cochlear roof
Hair Cells
LER   GER
Prosensory Cells
Lateral ↔ Medial

POM
Epithelial Cells
Prosensory Cells
Hair Cells

b E15.5/16.5 Cochlea

POM
Cochlear roof
Hair Cells
LER   GER
Prosensory Cells
Lateral ↔ Medial

POM
Epithelial Cells
Prosensory Cells
Hair Cells

c E13.5/14.5 Integrated Data
(Sun et al./Kolla et al.)

UMAP2
Epithelial Cells
Hair Cells
Prosensory Cells
POM
UMAP1

POM
Epithelial Cells
Prosensory Cells
Hair Cells

d E15.5/16.5 Integrated Data
(Rose et al./Kolla et al.)

UMAP2
POM
Epithelial Cells
Prosensory Cells
Hair Cells
UMAP1

POM
Epithelial Cells
Prosensory Cells
Hair Cells

e E13.5/14.5 WNT and ncWNT Signaling

Hair Cells
Prosensory Cells
POM
Epithelial Cells

f E15.5/16.5 WNT and ncWNT Signaling

Hair Cells
Prosensory Cells
POM
Epithelial Cells

g

h

POM
Epithelial Cells
Prosensory Cells
Hair Cells

i Epithelial Wnts
Wnt7b
Wnt7a
Wnt5a

j Epithelial Wnts
Wnt5a
Wnt7b
Wnt7a

k POM Wnts
Wnt5a
Wnt5b
Relative Contribution

l POM Wnts
Wnt5a
Wnt5b
Relative Contribution

Figure 3

Figure 4

Figure 5

Figure 6