Cognitive control networks in human and macaque

Valentina Mione[1, 2]

Freja Holm Kristensen[3]

Moataz Assem[2]

Urs Schüffelgen[4]

Søren Kyllingsbæk [3]

Mark Buckley[1]

Daniel J. Mitchell[2]

John Duncan[1, 2 +]

[1] Department of Experimental Psychology, University of Oxford, Anna Watts Building,

Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6GG, UK

[2] MRC Cognition and Brain Sciences Unit, University of Cambridge, 15 Chaucer Road,

Cambridge CB2 7EF, UK

[3] Department of Psychology, University of Copenhagen, Øster Farimagsgade 2A, 1353

Copenhagen K, Denmark

[4] Wellcome Centre for Integrative Neuroimaging (WIN), Department of Experimental

Psychology, University of Oxford, Oxford, UK

[+] Correspondence: john.duncan@mrc-cbu.cam.ac.uk

23   <u>Abstract</u>

A much-replicated finding in human brain imaging is a distributed "multiple-demand" or MD

system, increasing in activity for many kinds of cognitive demand, and centrally involved in

cognitive control. MD regions are proposed to encode a distributed mental model of critical

task events, bound together in the roles and relationships needed to direct action selection.

Though previous data hint at a corresponding network in the macaque, there has been no

direct comparison to human data. Here we used functional magnetic resonance imaging to

measure whole brain activation in a multi-step saccadic maze task, compared to a control

requiring similar moves but without goal-based decisions. Human data were a close match

to the canonical MD network, extended to include adjacent regions and in particular much

of the canonical dorsal attention network. Monkey data suggested correspondences in

dorsomedial frontal, lateral and medial parietal, insula/orbitofrontal and posterior temporal

cortex. In lateral frontal cortex there was just a single, largely dorsal activation patch, in

contrast to multiple distinct human patches. In macaque as in human, together with

previous data, our findings suggest an extended and strongly interconnected brain network

recruited by increased cognitive challenge.

<u>Introduction</u>

Among the most stable findings of human brain imaging is a "multiple-demand" or MD

system, defined by increased activation for a very wide range of cognitive demands [1] [2] [3].

With its broad recruitment, the MD system is thought to play a central role in attentional or

cognitive control [4] [3]. The high-resolution methods of the Human Connectome Project (HCP)

show nine distinct MD patches [5] [6], distributed across lateral and dorsomedial frontal cortex,

anterior insula, dorsal and medial parietal cortex, and posterior temporal cortex (Figure 1a).

Within these patches, based on the cortical parcellation of [7], Assem et al. [5] defined a set of

ten "core" MD parcels (Figure 1b), identified by the strongest converging response to

multiple cognitive demands, and strong interconnections indicated by resting state

correlations. Common activity for different demands, however, usually extends beyond

these core MD parcels, in particular to adjacent regions of dorsal attention (DA) and cingulo-

opercular (CO) networks [8] [6] (Figure 1b). For different demands, the precise activation

pattern shifts somewhat within the same nine MD patches, often with differential

recruitment of individual DA and CO parcels [6], but with strong overlap between demands

(Figures 1c & d).

Cognitive control must rest on an internal model of a task or situation, with parts combined

in the correct roles and relationships [9] [10]. It is the contents of this task model that indicate

correct behavior for a current goal. Studies using multivoxel pattern analysis show that,

across the human MD network, there is extensive coding of task-relevant information,

including stimuli, rules and responses [11]. Elsewhere, we have suggested that the widely

distributed parts of the MD network give access to many different kinds of information,

Figure 1. Human multiple-demand (MD) system. (a) 9-patch MD system based on mean activation across three dissimilar demands. Data from Human Connectome Project [5]. Top – inflated cortical surface; bottom – cortical flatmap. (b) Same data – average activation in individual cortical parcels from [7]. White outlines – core MD parcels [5]; green outlines – dorsal attention network [8]; blue outlines – cinguloopercular network [8]. (c) and (d) 9-patch MD system in independent data – (c) n-back working memory [6]; (d) mental rotation [12].

while strong interconnectivity allows different kinds of information to be integrated into a

task model [3].

A similar distributed network likely exists in the macaque. Electrophysiological studies,

certainly, show selective neural responses for many kinds of task event in lateral frontal,

dorsomedial frontal and parietal cortex [13] [14] [15] [16] [17] [18] [19] [20] [21]. Provisional support comes from

a handful of imaging studies. One comparison between pro- and antisaccades showed

increased activity in frontal eye field, anterior cingulate and intraparietal sulcus for the more

demanding antisaccade [22] [23]. A study of task switching suggested activation of lateral frontal

cortex, anterior cingulate, and a region of orbitofrontal cortex close to anterior insula [24].

Measures based on functional [25] and structural [26] connectivity have also been used to

estimate putative components of a macaque MD system, including regions of lateral frontal,

dorsomedial frontal, inferior parietal and medial parietal cortex. In the absence of a direct

comparison between species, however, it remains uncertain whether the macaque brain

contains a functional analogue to the distributed human MD system.

To address this uncertainty, we collected fMRI data from humans and macaques performing

a similar on-screen maze task. In the human, using HCP methods, we show the usual nine-

patch pattern of brain activity associated with increased cognitive demand. In the macaque,

we also observe widespread demand-related brain activity, and ask how well it matches the

human nine-patch pattern.

90   Results

Human

The human maze task is illustrated in Figure 2a. With a series of saccades, the task was to

navigate from a central start location (Figure 2a, white circle) to one of four possible goal

locations (Figure 2a, blue circles). Along the route, the current location was defined as the

current fixation point; permitted fixation locations were the intersections and corners of

maze paths, and each saccade moved just one step through the maze, i.e. from the current

location to one of the immediately adjacent locations. An initial display indicated the goal

for the current problem (Figure 2b, top), always one of the four shown in Figure 2a. At each

step towards this goal, a choice display indicated possible moves (Figure 2b). Of the

locations immediately adjacent to the current location, two were colored yellow, indicating

that they were open (available for selection), while the remainder were colored red,

indicating that they were blocked. A delay of 0.4 or 1.0 s preceded a go signal (Figure 2b,

blue dot in white circle), indicating that the next saccade could be made. The correct choice

Figure 2. (a) Human maze. White circle – initial fixation location; blue circles - possible goal
locations. (b) Human maze stimuli and timings. Upper: Example problem in the maze
condition (outer parts of maze not shown, though present on screen). On each step, yellow
dots marked open locations available for selection, with red locations blocked. On receipt of
the go signal (blue dot inside white circle), participants made a saccade to one of these
yellow locations. The task was to choose the shortest route to the goal. In this example, the
goal is reached directly in two steps. In other problems, longer routes were created by
enforcing some moves that did not move closer to the goal. Timings for each step were
identical. Lower: Control condition. (c) Monkey maze. All problems required just two steps
to the goal (brown arrows).   (d) Monkey maze stimuli and timings (upper – maze; lower -
control). Green locations – open; red locations – blocked.

was then one of the two open (yellow) locations, if possible moving closer to the goal. This

procedure was repeated for a series of steps, until the goal was finally reached and a reward

signal was received. Depending on the available choices at each step, the goal could be

reached directly, in just two steps, or less directly, in up to six steps including forced steps

away from the goal.

In a control task (Figure 2b, bottom), there was no goal and only a single location was open

(yellow) at each step. Again, the participant made a series of steps through the grid, but in

124   this case, simply following the route prescribed, until a reward signal was given when

125   reaching one of the (unmarked) goal locations after between 2 and 6 steps.

127   <u>Figure 3</u>. Behavioral data. (a) Human. Bars show 95% confidence intervals over
128   participants. (b) Monkeys A (top) and B (bottom). Bars show 95% confidence intervals over
129   sessions.
130

Behavioral data are shown in Figure 3a. Responses were recorded when the eye reached

one of the open or blocked locations, with saccades elsewhere ignored. For each step

through the maze, accordingly, there were five possible outcomes (correct, too fast, too

slow, move to blocked location, move to wrong open location). Data in Figure 3a are just for

steps in which the two open locations included one that was correct (towards the goal), one

that was not (away).   In both maze and control tasks, the majority of errors were either too

fast or slow, with few wrong choices among the indicated open and blocked locations.

Functional magnetic resonance imaging (fMRI) data were collected in blocks of problems of

a given type (maze or control). Each block lasted for 24 s, plus the time required to

complete the final problem. Every five consecutive blocks contained two blocks each of the

maze and control task, along with one additional block (not analysed) of a simple central

fixation task.

Whole brain data for the contrast of maze vs. control are shown in Figure 4. Figure 4a

shows whole-brain activation, on semi-inflated cortical surfaces for each hemisphere

(center), with cortical flatmaps to either side. Figure 4b shows average data for the two

hemispheres, related to the cortical parcellation of [7]. Colors for each parcel indicate mean

activation across all vertices within that parcel. Figure 4c shows mean activations for each of

12 cortical networks, defined by [8] using the [7] parcellation. At the left is mean activation for

the 10 core MD parcels defined by [5], all of which belong to the frontoparietal network (FPN)

from [8] (Figure 1c, yellow). Results for remaining FPN parcels are shown separately. In Figure

4b, colored outlines mark parcels of core MD (yellow), dorsal attention (DA, green) and

posterior multimodal networks (PM, lilac).

In broad outline, activations for this maze vs. control contrast strongly resembled the

canonical MD system (compare Figure 4a to Figure 1). The same six patches were observed

in lateral frontal cortex, anterior insula and dorsomedial frontal cortex. Maze activations

Figure 4. Contrast of maze minus control: Human data. (a) t values (thresholded at FDR<0.05) projected onto partially inflated cortical surface (center) and cortical flatmaps (sides). (b) Mean t value (for regions with FDR < 0.05) in individual cortical parcels from [7]. Yellow outlines – core MD parcels [5]; green outlines – dorsal attention network [8]; pink outlines – posterior multimodal network [8]. Parcel labels from [7]; some omitted to avoid clutter. (c) Mean values in the 12 cortical networks defined by [8]. Data are shown separately for core MD and other parcels from the frontoparietal network defined by [8]. Horizontal and vertical black lines show mean and 95% confidence intervals; violins show distribution across participants.

also encompassed the two parietal MD patches, though with substantial additional spread

on both lateral and medial surfaces. A posterior temporal activation was also a close match

to the posterior temporal patch from [5].

Parcel-wise results in Figure 4b confirm positive activations in all core MD parcels,

substantially extending into adjacent regions. Strong activations in particular, were seen in

many parcels of the DA network, with the next strongest average activity in PM (Figure 4c).

Additional activations extended across a large region of medial parietal cortex, a number of

dorsomedial frontal parcels adjacent to core area 8BM, and lateral frontal regions

immediately dorsal and postero-ventral to core MD regions. Strong activation was also seen

in the frontal eye field (Figure 4b, FEF).

For a closer match to the macaque task (see below), the analysis was repeated using data

just from two-step problems. The results (Supplementary Figure 1) were closely similar to

results from the main analysis, with the expected reduced power. For comparison, we also

examined data from a version of the task with manual rather than saccadic responses (see

Methods). Again, results were closely similar (Supplementary Figure 2).

<u>Macaque</u>

The monkey task was a simplified version of the human maze (Figure 2c). The goal was

always reached in two steps, the first moving to one of the four locations immediately

adjacent to the central start location, and the second reaching the goal. Again, animals

worked continuously through problems for the duration of each block, which lasted a

minimum of 24 s, plus the time required to complete the final problem.

Behavioral data are shown in Figure 3b. Both animals lower accuracy than humans, but an

otherwise similar distribution of errors. Again, in particular, there were few choices of

blocked locations, and many more correct than incorrect choices between the two open

locations, indicating strong control of behavior by the remembered goal.

For imaging analysis, because timing errors often indicated poor task engagement, we took

data just from problems that ended with saccade to an open or blocked location, made in

the correct time window. Imaging data for the contrast of maze vs. control are shown in

Figure 5. To increase power, data from the two hemispheres have been averaged, and are

<u>Figure 5</u>. Contrast of maze minus control: Monkey data. Left – cortical flatmap, with

projected t values (thresholded at FDR<0.05). Parcel labels according to [27]; some omitted to

avoid clutter. Right – partially inflated cortical surface.

shown on flatmap (Figure 5a) and semi-inflated surfaces (Figure 5b) of the left hemisphere.

Parcellation is according to the atlas of [27], projected onto the F99 surface [28].

In several respects, there was good correspondence between monkey and human data. On

the dorsomedial frontal surface, there was a large patch of activation centered on areas F6,

24c and 24c', corresponding to the monkey's preSMA and anterior cingulate. On the lateral

parietal surface, a patch of activation extended over parts of areas 7a (Opt/PG) and 7op,

extending into adjacent Tpt. In agreement with the human data, there was also a patch of

activation in medial parietal cortex, extending from 7m to adjacent parts of 23b/c and 31. In

the insula, there was activation in Iam, extending into adjacent regions of orbitofrontal

cortex. In occipitotemporal cortex there was a patch of activation in anterior V4v, tentatively

corresponding to the posterior temporal patch in human data.

Correspondence was less clear on the lateral surface of the frontal lobe. In the monkey,

contrasting with the multiple, clearly separable, activation patches in the human, there was

just one large posterior dorsal patch extending from F7 (dorsal premotor) through 8ad

(frontal eye field) to dorsolateral frontal cortex (9d, 46d) dorsal to the principal sulcus, and

within its dorsal bank. There was modest extension into the ventral bank of the sulcus (46v),

but not to the ventrolateral frontal surface. Differences between human and macaque were

also notable in dorsal parietal cortex, with strong activation of DA regions in the human, not

visible in the macaque.

Finally, there were further activation patches in the monkey not matched in human data.

Most conspicuously, in the monkey, there was a large activation patch in sensorimotor

cortex (3a/b, extending into ventral F1), and additional scattered activations close to the

temporal pole.

Finer breakdowns of the data are shown in Supplementary Figure 3.   Activations were

closely similar for left and right hemispheres (Supplementary Figure 3a). Except in lateral

frontal cortex, activations were stronger and more extensive for one of the two animals

(Supplementary Figure 3b); as expected, the combined analysis (Figure 5) revealed regions

common to the two animals. Of note, in one of the two animals, there was extensive dorsal

parietal activation, not visible in the combined data; while in the other, lateral frontal

activation spread ventrally, to include regions 45b and 44.

A variety of methods have been proposed for approximate warping of the macaque to the

human brain. Figure 6a shows macaque activations from Figure 5 warped by a method

based on resting state fMRI data [29], with human activations repeated for comparison (Figure

6b). With this approximate warping, lateral frontal activations show substantial overlap

Figure 6. (a) Monkey data: Contrast of maze minus control warped to human cortical surface. (b) Human data for comparison.

between species. As noted in the main analysis, dorsal parietal activation is absent in the

macaque data. In this warped version, insula/orbitofrontal activation in the macaque is

placed in human pOFC (top left of flatmap), perhaps reflecting limitations of the warping

method in this part of the brain.

Subcortical activation

In the human, cortical MD activity is generally accompanied by focal activations in caudate,

thalamus and cerebellum.[5] The same was true in the present human data (Figure 7a), and in

the monkey, regions of activation were also seen in cerebellum and caudate (Figure 7b).

Discussion

Though precise correspondences between brain networks in human and monkey are

uncertain, the whole-brain data of fMRI can provide a useful comparison. Here, we used

fMRI in human and monkey to compare brain activation during a multi-step spatial maze

task.

Figure 7. Contrast of maze minus control shown on horizontal slices (slice locations shown on right). (a) Human. (b) Monkey.

In humans, as expected, we found a highly specific, distributed pattern of brain activity, incorporating the canonical MD system, and extending to adjacent regions of DA and other networks. MD activations strongly resembled those broadly associated with increasing cognitive demand, while DA and visual activations were likely associated with the specific visuospatial content of the maze task.

In several respects, monkey data were a good match to human MD activations. Patches of activation were seen in dorsomedial frontal cortex, insula and adjacent orbitofrontal cortex, lateral and medial parietal cortex, and a small patch in occipitotemporal cortex. Conspicuous activation was also seen in lateral frontal cortex, though in contrast to several distinct patches in the human, monkey activations were seen only in a region extending from premotor cortex to the banks of the principal sulcus.

In human imaging, the MD system is defined by overlapping activation for varied cognitive demands. In the present work, in contrast, we used just a single task. There is some overlap between our putative monkey MD network and previous findings, including activations of intraparietal sulcus and anterior cingulate reported for antisaccades compared to

prosaccades [22][23], and of lateral frontal cortex, anterior cingulate and orbitofrontal cortex

close to insula for task switching compared to repeat [24]. Though a wider variety of tasks

would be desirable, these overlaps suggest that, in the macaque brain as in the human, an

extended MD network is recruited by diverse cognitive challenges.

Our results may be compared to previous estimates of a macaque MD system, based on

either functional or structural connectivity. In the human brain, MD regions show strong

functional connectivity, indicated by correlated activity time-courses in resting state data [5].

Previously, we searched for a similar network based on functional connectivity data from

anesthetised monkeys [25]. On the one hand, strong connectivity between regions of lateral

frontal, dorsomedial frontal and inferior parietal cortex was coarsely in line with current

findings. Often these strongly-connected regions were close to or overlapping with

activated regions in the maze task. At the same time, there were notable differences

between networks defined by these two methods. The functional connectivity network, for

example, included a larger region of premotor cortex, extending back into primary motor

cortex, and ventrally well beyond the principal sulcus. This network, furthermore, included

neither orbitofrontal/anterior insular nor medial parietal cortex. Plausibly, a network

underlying high-level cognitive control is only partially visible in time course correlations in

anesthetised animals. A second study identified macaque regions that most strongly

resemble human MD regions in terms of structural connectivity to major white matter tracts

[26]. Again there was coarse overlap with the current network, including regions in lateral

frontal, dorsomedial frontal, inferior parietal and medial parietal cortex, but again, this

coarse agreement was accompanied by notable differences. Once more, for example, the

structural connectivity network included a much larger region of lateral frontal cortex.

Plausibly, connectivity to major white matter tracts gives only a partial clue to functional

specialisation, especially for regions in close anatomical proximity.

In the present macaque data, data averaged across the two animals showed maze-related

lateral frontal activation that was largely restricted to dorsal regions, with minimal extension

onto the inferior convexity (ventrolateral frontal cortex, vlPFC). Others have suggested that

vlPFC plays a core role in cognitive and attentional control [30] [31] , and in a recent

electrophysiological study, we found that coding of both spatial and nonspatial task

information was stronger in ventral than dorsal frontal cortex [32]. In a spatial maze task too,

electrophysiological recordings show a strong role for vlPFC in rapid discrimination of

relevant sensory input, including initial goal and subsequent choice stimuli [33]. In

anesthetised macaques, the functional connectivity profile of vlPFC regions also shows

resemblance to the proposed MD network, including connectivity in dorsomedial frontal and

lateral parietal cortex [34] [25]. The absence of vlPFC activations in the current data may be a

false negative, linked to limited power of imaging data based just on a limited number of

sessions in only two animals. Separate examination of data for each animal (see

Supplementary Figure 3b) indeed suggests ventral extension of lateral frontal activity in one

of the two.

In the human, MD activation was accompanied by strong activation of the DA network,

including extensive regions of dorsal parietal cortex. In the monkey, the most consistent

parietal activation was ventral, though again, major dorsal extension was clearly visible in

one of the two animals (Supplementary Figure 3b). Again it is likely that, with more data,

macaque and human findings would show increased convergence.

The monkey data also showed regions not mirrored in the human. Most conspicuous was a

region of sensorimotor cortex. The imaging situation, of course, may often be different for

monkey and human. While humans lie still, monkeys do not, and possibly, the amount of

movement depends on task demands.

The existence of MD regions, recruited as a network to address diverse cognitive challenges,

does not imply homogeneity of function, either between regions or within a single region.

Between tasks, human data show partial functional specialisation of different MD regions [5].

Within a single patch, there are gradients of activation for any given task demand, with the

magnitude and direction of these gradients varying from one task to another [6]. In the

monkey, too, functional connectivity profiles for different regions of lateral frontal cortex

show strong family resemblances, but also quantitative variations [34]. Differing anatomical

connections likely give MD regions differential access to information of different kinds [30]; at

the same time, strong interconnectivity may bring substantial information sharing and co-

recruitment.

In putative MD regions defined here, electrophysiological studies show coding of many

different kinds of task-relevant information. This is amply documented in lateral frontal,

dorsomedial frontal, insula/orbitofrontal, premotor and lateral parietal cortex[13 14 15 16 17 18 19

20 35 36 21 37 38]. These results mirror human fMRI data, showing extensive MD encoding of a

task's stimuli, rules and responses [11]. In monkey as in human, we suggest that a distributed

MD network assembles information of many kinds into a distributed task model, needed to

represent task events and to choose appropriate behavior.

<u>Methods</u>

<u>Human: Participants</u>

39 participants took part, of whom two were not analysed owing to missing or low quality

structural scans, and three were excluded owing to poor performance (see below). The 34

participants (22 female) in the analysed sample had a mean age of 30.2 years, with normal

or corrected-to-normal vision, and with no history of neurological or psychiatric disorders.

Participants provided informed consent and were paid for their time. The experiment was

conducted in accordance with ethical approval from the Cambridge Psychology Research

Ethics Committee.

<u>Human: Maze task</u>

The experimental task was programmed in Matlab (The Mathworks Inc.) using Psychtoolbox

version 3.0.14 [39]. In two of four runs, the subject of the present report, participants

responded by making saccades to a sequence of target locations, while their gaze was

monitored using an eye tracker (EyeLink 1000; SR Research Ltd). In the other two runs (not

analysed here, except for comparison data in Supplementary Figure 2), tasks were identical

except that responses were made by button press. Saccadic runs were either the first two or

the last two, alternating across participants. In two participants, one of the saccadic runs

was missing due to technical issues. When responding using saccades, a location was

deemed fixated when gaze had remained within 1.6 degrees of visual angle (dva) of the

specified coordinate for 0.3 s.

The experiment used a blocked design, with each run containing 25 blocks, in five groups of

five. Each group contained two blocks of the maze task, two of control, and one (not

analysed) of fixation. The fixation block was always the third in the group, with one block

each of maze and control before the fixation block, one block each following it, and the

order of maze and control blocks counterbalanced across runs. Within each block, problems

were presented one after another for 24 s, after which the current problem was allowed to

finish before proceeding to the next block. Blocks comprised 1-6 problems (median 3),

depending on the number of required saccades and the performance of the participant.

Each run lasted approximately twelve minutes.

For every problem, participants were presented with a grey grid, whose width spanned

approximately 11.5 dva, on a white background (Figure 2a). At the start of each problem, a

circular yellow fixation dot (0.7 dva) was positioned at the center of the grid (Figure 2b). The

participant was required to fixate this dot for 0.3 s, whereupon it turned white, whilst

fixation was maintained for a further 1.2 s.

During this period, events diverged for maze and control tasks. In the maze task, for 0.6 s,

one of the four possible goal locations was highlighted by a blue circle (Figure 2b), then

disappeared for a further 0.6 s. The goal location was always one of the four shown in

Figure 2a. At the end of this 1.2 s period, two movement options appeared as yellow circles

at grid intersections immediately adjacent to the fixation location; the other adjacent

locations were marked by red circles, indicating them as blocked. The participant was to

maintain fixation for a further 0.4 or 1.0 s (equal probability), whereupon a small blue dot

(0.15 dva) appeared on the white fixated circle, signalling that a saccade should be made to

a yellow location within two seconds. The participant maintained fixation at the new

location for 0.6 s, before the choices appeared for the next step. Steps proceeded in this

manner until the goal was reached, after two, four, or six steps, whereupon the fixated circle

turned blue for 1 s. At each step, if possible, the participant was to choose a yellow option

that moved closer to the goal. In the simplest two-step routes, one of the two choice

options for the first step was a move one step closer to the goal, and options for the second

step included the goal itself (Figure 2a).   In more demanding routes, some steps had two

choice options that were equally attractive, either both towards or both away from the goal.

Sometimes these longer routes involved a move into the outer perimeter of the maze.

For control problems, no goal location was shown at problem onset (Figure 2b, bottom). For

1.2 s, the participant simply maintained fixation on the central white dot. At each

subsequent step, only one yellow circle was presented, cueing the move to be selected.

Accordingly, the participant was to follow the route prescribed by this sequence of cued

moves. Other details copied those for maze problems. The problem completed after 2, 4 or

6 moves, when the route had always led to one of the same four locations used as goals in

maze problems, and when this location was reached, the fixated circle turned blue.

In fixation blocks, neither the goal location nor choice locations appeared, requiring the

participant to maintain fixation on the central white circle for the entire problem, which was

matched in duration to the mean duration of either two-, four-, or six-step problems, until

the fixated circle turned blue.

In all block types, to maintain participant engagement, completing a problem was rewarded

by a pleasant sound, and animated stars that travelled from the fixated location to the

center of the grid. Every fourth problem had three stars, while other problems had one star,

although this was of no significance. The movement of the stars encouraged gaze to return

to the center, ready for the next problem, which started 0.5 s later. If any error was made

(failing to saccade to a yellow circle in time, making a saccade before the go cue appeared,

making a saccade to a red circle, or failing to choose the shortest available route to the goal),

the current problem aborted immediately, with a central red cross and a disappointed sound

(1 s), followed by the usual 0.5 s inter-problem interval. In the maze condition, three

participants consistently failed to select the shortest available route to the goal (on >45% of

problems, compared to <15% for other participants), so were excluded from all analyses.

Across problems of each condition, routes (defined by goal locations crossed with the

location of choice options) were presented in pseudorandom order. On average, combining

maze and control conditions, a run contained 14.1 two-step problems, 23.3 four-step

problems and 18.8 6-step problems (with additionally 14.5 fixation problems).

Participants practised a few problems of each type outside of the scanner, using manual

responses, until they and the experimenter were happy that they understood the task. In

the scanner, the eye tracker was calibrated, and the participant practised responding using

the button box and using saccades, before proceeding to the main experiment.

Human: MRI acquisition

Scanning was performed on a 3 Tesla Siemens Prisma MRI scanner, using a 32-channel head

coil. A T1-weighted structural volume was acquired using a 3D MPRAGE sequence (TR = 2.4

s, TE = 0.0022 s, flip angle = 8°, 0.8 mm isotropic voxels). A T2-weighted structural volume

was acquired using a 3D SPACE sequence (TR = 3.2 s, TE = 0.56 s, flip angle = 120°, 0.8 mm

isotropic voxels). During each of four task runs, T2*-weighted scans were acquired using a

gradient echo 2D EPI sequence (TR = 0.80 s, TE = 0.037 s, flip angle = 52°, 2 mm isotropic

voxels, 72 slices acquired with multiband factor 8). EPI runs were acquired with alternating

phase-encoding directions (AP/PA). Two spin echo volumes, with similar phase reversal,

were acquired between the second and third runs.

Human: MRI analysis

Data processing was based on the HCP minimal pre-processing pipelines ([40]; version 4.7.0),

called using automatic analysis software [41]. Per-subject processing steps are summarised

briefly here. Structural volumes were undistorted using the spin-echo volumes, and rigidly

aligned to MNI space. Subcortical structures and the cortical ribbon were segmented using

the structural volumes, and cortical surface meshes were extracted, using FreeSurfer ([42];

version 6.0.0). Functional volumes were aligned to the structurals, undistorted using the

spin-echo volumes, motion-corrected using FSL's MCFLIRT [43], and mapped to CIFTI

"grayordinate" space, where they were smoothed with a 2 mm full width half maximum

(FWHM) Gaussian kernel. Functional data were linearly detrended, and denoised using

spatial ICA+FIX [44] on concatenated runs, with a FIX-threshold of 20. Cortical surfaces were

registered to template space using multimodal surface matching ("MSMAll"; [45]).

To model task-based activations, for each run, a first-level general linear model (GLM) was

estimated using FSL's FEAT [46]. One regressor was defined for problems of each condition

(whether correctly completed or not), as boxcars from initial fixation to the onset of

feedback. These regressors were convolved with the canonical haemodynamic response

function (HRF) and its temporal derivative. A temporal high-pass filter with a cut-off period

of 200 s was applied to the data and the model. Activation due to goal-based decisions was

estimated from the parameter estimates for the maze condition's HRF-convolved regressor

contrasted against the control condition. Statistical significance was tested across

participants, using two-tailed, one-sample t-tests. This was initially performed at every

cortical vertex, then repeated for each cortical region in the HCP's multimodal parcellation [7],

after averaging across vertices and hemispheres for each region. In both analyses, multiple

comparisons were accounted for by controlling the false discovery rate (FDR = 0.05) [47].

Results were further summarised at a network level according to the networks defined by [8]

(excluding parcels whose network assignment differed across hemispheres).

<u>Monkey: Participants</u>

Subjects were two male rhesus monkeys (Macaca mulatta), each weighing 14 kg.

Experiments were performed in accordance with the Animals (Scientific Procedures) Act

1986 of the UK; all procedures were licensed by a Home Office Project License obtained

after review by Oxford University's Animal Care and Ethical Review committee, and were in

compliance with the guidelines of the European Community for the care and use of

laboratory animals (EUVD, European Union directive 86/609/EEC).

For each animal, an MRI-compatible head holder (Rogue Research) was sterotaxically

implanted on the skull. All surgical procedures were aseptic and carried out under general

anesthesia. At the end of the experiments, animals were deeply anesthetized with

barbiturate and then perfused through the heart with heparinized saline followed by 10%

formaldehyde in saline.

<u>Monkey: Maze task</u>

For monkeys, the task was a simplified version of the human maze. Problems were

presented on a grid of lines (Figure 2c), corresponding just to the inner 3 x 3 locations of the

human maze. In the maze condition, all routes were two-step, with one of the two choice

options at step 1 moving closer to the goal, and the goal then reached at step 2. In the

control condition, again, moves were forced by just a single available option at each step.

Timings for the maze condition are illustrated in Figure 2d. At the start of each problem,

subjects were given a green fixation point in the middle of the grid and had to fixate on this

point within 2 s. Once the fixation was initiated, a white diamond (cursor) appeared in the

location of the fixation. After 0.2 s a blue circle (final target) appeared at the final goal

location of the current problem and remained on the screen for 0.5 s while the monkeys

kept fixating in the center (except for a single session, when the target was shown for 0.4 s).

The final goal was only shown at this point and then disappeared, so the monkeys needed to

hold the location in mind throughout the remainder of the problem. After a 0.3 s delay, the

4 choice options of the first step were presented. 2 green dots and 2 red dots appeared

simultaneously at the 4 nodes next to the central one (where the monkeys were still

fixating). The red dots represented nodes that were blocked, and only one of the greens was

a correct choice, being closer to the final target location. The go signal to make a saccade

was the disappearance of the green fixation point, occurring after a delay randomly either

0.3 or 0.9 s. After the go signal, the monkey had 0.5 s to initiate a saccade to the correct

choice location and 0.05 s to complete the saccade. Once a saccade to the new location was

made, if the choice was correct, all the other old stimuli (initial fixation point, cursor, other

choices) disappeared while the monkeys had to hold fixation on the new location.

The second step of the problem had a similar structure: the cursor appeared when the

second fixation started, the second set of 3 choices (2 green 1 red) appeared after 0.3 s and

were presented for the same random delay as before, followed by the go signal. Using the

same timing as in step 1, the monkey needed to make a saccade towards the correct green

location. At this point, one of the choice options lay on the actual final goal location,

allowing the monkeys to complete the problem. If this happened, when the eye landed, the

cursor appeared at the fixated target location, joined after 0.05 s by brief (0.05 s)

reappearance of the blue target itself. A smoothie reward was then delivered (delivery

duration: ~0.05 s), the monkey could break fixation, and the problem ended with the

disappearance of all the stimuli (except the grid) from the screen and an inter-problem

interval of 1 s. Errors at any point during the problem (incorrect location selections, timing

errors, fixation breaks) led to an immediate error cue (large red circle) shown in the center

of the screen, and termination of the problem without reward.

Events for the control condition were similar, except that no target was shown at the start of

the problem. Again, each problem had two steps, each with only a single choice location,

and reward was delivered after the second correctly executed step. In an additional fixation

condition (not analysed here), each problem began with a green fixation point in the middle

of the grid. Fixation was required within 2 s, and once it was achieved, the fixation point

was surrounded by a white diamond. Maintained fixation was required for at least 2.05 s,

after which a blue circle appeared at fixation for 0.05 s, and reward was delivered.

As for the human, the experiment used a block design. For as long as the monkey kept

working, blocks of the maze condition alternated with blocks of a non-maze condition. Each

non-maze block was randomly selected to be either control or fixation, after which the next

block was again maze. The target length of each block was 30 s. When a problem was

completed, the block terminated if >24 s had already elapsed; otherwise a new problem

began and the block continued until this problem was finished.

The task was programmed in Psychopy Release 2021.2.3 (https://www.psychopy.org/).

Stimuli were presented on a 23 inch LCD screen (https://www.crsltd.com/tools-for-

functional-imaging/mr-safe-displays/boldscreen-23-lcd-for-fmri/), positioned 80 cm from the

monkey's eye. The grid measured 15 x 15 dva.   The fixation point and choice cues were

filled circles measuring 1.15 dva. The cursor, blue target, and error cue measured 2.3 dva.

Eye movement data were sampled at 1000 Hz using an MRI-compatible tracking system

(EyeLink 1000 Plus, SR Research; https://www.sr-research.com/). At each step, fixation was

required within a 7 x 7 dva window centered on the appropriate grid position. Reward (fruit

smoothie) was delivered through a Peristaltic Electric Operated Positive Displacement Pump

(Verderflex).

Monkey: MRI acquisition

Data were collected over 16 sessions in monkey A, and 18 in monkey B (5 sessions from

monkey B were excluded based on poor behavioral performance). Animals were head-fixed

in a sphinx position in an MRI-compatible chair. fMRI data were collected using a whole-

body 3T MRI scanner with a full-sized horizontal bore and a custom-made four-channel

phased-array local receive coil in conjunction with a radial transmit coil (H. Kolster;

Windmiller Kolster Scientific, Fresno, CA, USA). Whole-brain blood-oxygen-level-dependent

(BOLD) fMRI data were acquired using a gradient echo T2* echo planar imaging (EPI)

sequence with 1.5-mm3 isotropic voxel size resolution, 36 interleaved slices with no gap,

repetition time (TR) = 2.28 s, echo time (TE) = 30 ms, and flip angle = 90°. Proton density–

weighted images using a gradient-refocused echo (GRE) sequence (TR = 10 ms, TE = 2.52 ms,

and flip angle = 25°) were acquired as references for body motion artefact correction. T1-

weighted magnetization-prepared rapid gradient echo images (0.5-mm^3 isotropic voxel size

resolution, TR = 2.5 ms, TE = 4.01 ms, inversion pulse time = 1.1 s, and flip angle = 8°) were

acquired in separate anesthetized scanning sessions. In these sessions, monkeys were

scanned in sphinx position in an MRI-compatible stereotactic frame (Crist Instrument).

Anesthesia was induced via intramuscular injection of ketamine (10 mg/kg), xylazine (0.125

to 0.25 mg/kg), and midazolam (0.1 mg/kg).

Monkey: Preprocessing and reconstruction pipeline

Magnetic resonance images were preprocessed using tools from the FMRIB Software Library

(FSL) and the Magnetic Resonance Comparative Anatomy Toolbox (MrCat;

www.neuroecologylab.org). The GRE image was used to carry out the T2* EPI image

reconstruction by an offline-SENSE reconstruction method (Dr. Kolster, Windmiller Kolster

Scientific, Fresno, CA, USA).

<u>Monkey: MRI analysis</u>

Data were analysed in Matlab (The Mathworks Inc.), using automatic analysis software [48] to

call SPM functions (Statistical Parametric Mapping; www.fil.ion.ucl.ac.uk/spm). The T1-

weighted structural scan was spatially normalized to the 112RM-SL template [49] using

nonlinear warping, and segmented by tissue type (grey matter, white matter, CSF). The first

ten EPI volumes of each session were discarded, and then the remaining volumes were

rigidly realigned to each other, corrected for slice timing, and rigidly co-registered to the

structural volume. Although the head was fixed, limb movements caused changes in the

magnetic field, resulting in momentary spatial distortion and extreme intensity fluctuations.

Time points affected by such artefacts were defined as those with an apparent displacement

of the brain by > 1 mm, an apparent rotation of > 0.5 degrees around any axis, or an extreme

global intensity difference from the previous scan. (Global intensity differences were

calculated as the mean across voxels of the squared difference between one volume and the

next; this time-course was smoothed with Matlab's robust Lowess method, with a span of

10% of the data length; outliers were defined as exceeding four standard deviations of the

lower 90% of the distribution of residuals from the smoothed time-course.) EPI volumes

were finally warped to template space, using the parameters derived from the structural,

resliced to 1 mm isotropic resolution, and spatially smoothed using a 4 mm FWHM Gaussian

kernel.

Because animals often broke fixation when not working well, the analysis included just

problems that did not terminate with a timing error. For each session, the first-level GLM

included one regressor to model problems of the maze condition, another for the control

condition, and similar, but unanalysed, regressors to model excluded problems. These

regressors were constructed from boxcars spanning initial fixation to the onset of feedback

for each problem, convolved with SPM's canonical HRF. A separate regressor modelled

moments when any juice rewards were delivered, timed for 1 s from the recorded triggers,

and convolved with the HRF. To account for variance due to limb movements, the model

also included extensive unconvolved covariates of no interest. These included the six rigid-

body realignment parameters, their squares, and temporal derivatives of both, as well as the

mean signal from the CSF, and individual regressors to model any scan identified as a global

outlier (as defined above). A temporal high-pass filter with SPM's default cut-off period of

128 s was applied to the data and the model. Activation due to goal-based decisions was

estimated from the parameter estimates for the maze condition's HRF-convolved regressor

contrasted against the control condition. Statistical significance, per voxel, was tested across

sessions x hemispheres, using two-tailed t-tests, based on a repeated-measures ANOVA with

separate covariates to model each hemisphere of each animal, and a pooled error term that

assumed similar error covariance across voxels [50]. Multiple comparisons were accounted for

by controlling the false discovery rate (FDR = 0.05).

To warp macaque surface-based activations to the human cortical surface , we used the

transformation published by [29]. This required two steps: the activations were first warped

from the F99 surface to the Yerkes19 surface, using the transformation in RheMAP-Surf [51],

and then from the Yerkes19 surface to the human surface [29]. Both steps were performed

using Connectome Workbench version 1.5.0.

597
598

599   <u>Acknowledgments</u>

This work was supported by Wellcome Trust Grant 101092/Z/13/Z, Wellcome Trust Early

Career Award 101092/Z/13/Z, Medical Research Council UK Program MC_UU_00030/7, and

Carlsberg Foundation Grant CF22-1111. We thank Simon Baumann and Stuart Mason for

assistance with monkey training, and Makoto Kusunoki for surgery. For the purpose of open

access, the author has applied a Creative Commons Attribution (CC BY) licence to any Author

Accepted Manuscript version arising from this submission.

References

1   Duncan, J. & Owen, A. M. Common regions of the human frontal lobe recruited by

diverse cognitive demands. *Trends Neurosci.* **23**, 475-483 (2000).

2   Duncan, J. The multiple-demand (MD) system of the primate brain: Mental programs

for intelligent behaviour. *Trends in Cognitive Sciences* **14**, 172-179 (2010).

3   Duncan, J., Assem, M. & Shashidhara, S. Integrated intelligence from distributed brain

activity. . *Trends in Cognitive Sciences* **24**, 838-852 (2020).

4   Cole, M. W. & Schneider, W. The cognitive control network: Integrated cortical

regions with dissociable functions. *Neuroimage* **37**, 343-360, (2007).

5   Assem, M., Glasser, M. F., Van Essen, D. C. & Duncan, J. A domain-general cognitive

core defined in multimodally parcellated human cortex. *Cereb. Cortex* **30**, 4361-4380

(2020).

6   Assem, M., Shashidhara, S., Glasser, M. F. & Duncan, J. Basis of executive functions in

fine-grained architecture of cortical and subcortical human brain networks. *Cereb.*

Cortex **34**, bhad537 (2024).

7   Glasser, M. F. *et al.* A multi-modal parcellation of human cerebral cortex. *Nature* **536**,

171-178 (2016).

8   Ji, J. L. *et al.* Mapping the human brain's cortical-subcortical functional network

organization. *Neuroimage* **185**, 35-57 (2019).

9   Kimberg, D. Y. & Farah, M. J. A unified account of cognitive impairments following

frontal lobe damage: The role of working memory in complex, organized behavior. *J.*

Exp. Psychol. Gen. **122**, 411-428 (1993).

10   Duncan, J. Construction and use of mental models: Organizing principles for the

science of brain and mind. *Neuropsychologia* **207**, 109062 (2025).

11   Woolgar, A., Jackson, J. & Duncan, J. Coding of visual, auditory, rule, and response

632   information in the brain: 10 years of multivoxel pattern analysis. *J. Cogn. Neurosci.*

633   **28**, 1433-1454 (2016).

634   12   Faber D, Shields G, Duncan J, & Assem M. Precise topology of diverse cognitive

635   activations within the multiple demand core of the human brain. Program No.:

636   PSTR422.19/T1. 2024 Neuroscience Meeting Planner. Chicago, IL: Society for

637   Neuroscience (2024)

638   13   Amemori, K., Amemori, S. & Graybiel, A. M. Motivation and affective judgments

639   differentially recruit neurons in the primate dorsolateral prefrontal and anterior

640   cingulate cortex. *J. Neurosci.* **35**, 1939-1953 (2015).

641   14   Cavanagh, S. E., Towers, J. P., Wallis, J. D., Hunt, L. T. & Kennerley, S. W. Reconciling

642   persistent and dynamic hypotheses of working memory coding in prefrontal cortex.

643   *Nat Commun* **9**, 3498 (2018).

644   15   Chafee, M. W. & Goldman-Rakic, P. S. Matching patterns of activity in primate

645   prefrontal area 8a and parietal area 7ip neurons during a spatial working memory

646   task. *J. Neurophysiol.* **79**, 2919-2940 (1998).

647   16   Goodwin, S. J., Blackman, R. K., Sakellaridi, S. & Chafee, M. V. Executive control over

648   cognition: Stronger and earlier rule-based modulation of spatial category signals in

649   prefrontal cortex relative to parietal cortex. *J. Neurosci.* **32**, 3499-3515 (2012).

650   17   Johnston, K., Levin, H. M., Koval, M. J. & Everling, S. Top-down control-signal

651   dynamics in anterior cingulate and prefrontal cortex neurons following task

652   switching. *Neuron* **53**, 453-462 (2007).

18   Kadohisa, M., Watanabe, K., Kusunoki, M., Buckley, M. J. & Duncan, J. Focused

representation of successive task episodes in frontal and parietal cortex. *Cereb.*

Cortex **30**, 1779-1796 (2020).

19   Loonis, R. F., Brincat, S. L., Antzoulatos, E. G. & Miller, E. K. A Meta-Analysis Suggests

Different Neural Correlates for Implicit and Explicit Learning. *Neuron* **96**, 521-534

e527 (2017).

20   Miller, E. K. & Cohen, J. D. An integrative theory of prefrontal cortex function. *Annu.*

Rev. Neurosci. **24**, 167-202 (2001).

21   Salazar, R. F., Dotson, N. M., Bressler, S. L. & Gray, C. M. Content-specific fronto-

parietal synchronization during visual working memory. *Science* **338**, 1097-1100

(2012).

22   Ford, K. A., Goltz, H. C., Brown, M. R. & Everling, S. Neural processes associated with

antisaccade task performance investigated with event-related FMRI. *J. Neurophysiol.*

94, 429-440 (2005).

23   Ford, K. A., Gati, J. S., Menon, R. S. & Everling, S. BOLD fMRI activation for anti-

saccades in nonhuman primates. *Neuroimage* **45**, 470-476 (2009).

24   Premereur, E., Janssen, P. & Vanduffel, W. Functional MRI in Macaque Monkeys

during Task Switching. *J. Neurosci.* **38**, 10619-10630 (2018).

25   Mitchell, D. J. *et al.* A putative multiple-demand system in the macaque brain. *J.*

Neurosci. **36**, 8574-8585 (2016).

26   Karadachka, K. *et al.* Structural connectivity of the multiple demand network in

humans and comparison to the macaque brain. *Cereb. Cortex* **33**, 10959-10971

(2023).

27   Saleem, K. S. *et al.* High-resolution mapping and digital atlas of subcortical regions in

the macaque monkey based on matched MAP-MRI and histology. *Neuroimage* **245**

(2021).

28   Van Essen, D. C., Glasser, M. F., Dierker, D. L. & Harwell, J. Cortical parcellations of the

macaque monkey analyzed on surface-based atlases. *Cereb Cortex* **22**, 2227-2240

(2012).

29   Xu, T. *et al.* Cross-species functional alignment reveals evolutionary hierarchy within

the connectome. . *Neuroimage* **223**, 117346 (2020).

30   Passingham, R. E. & Wise, S. P. *The neurobiology of the prefrontal cortex: Anatomy,*

evolution, and the origin of insight. (Oxford University Press, 2012).

31   Trambaiolli, L. R. *et al.* Anatomical and functional connectivity support the existence

of a salience network node within the caudal ventrolateral prefrontal cortex. *Elife* **11**,

76334 (2022).

32   Kadohisa, M. *et al.* Frontal and temporal coding dynamics in successive steps of

complex behavior. *Neuron* **111**, 430-443 (2023).

33   Mione, V., Achterberg, J., Kusunoki, M., Buckley, M. J. & Duncan, J. Neural dynamics

of an extended frontal lobe network in goal-subgoal problem solving. *bioRxiv*, 2025-

2005 (2025).

34   Neubert, F. X., Mars, R. B., Thomas, A. G., Sallet, J. & Rushworth, M. F. Comparison of

human ventral frontal cortex areas for cognitive control and language with areas in

monkey frontal cortex. *Neuron* **81**, 700-713 (2014).

35   Mizuhiki, T., Richmond, B. J. & Shidara, M. Mode changes in activity of single neurons

in anterior insular cortex across trials during multi-trial reward schedules. *Neurosci.*

Res. **57**, 587-591 (2007).

36   Nakajima, T., Hosaka, R. & Mushiake, H. Complementary Roles of Primate Dorsal

Premotor and Pre-Supplementary Motor Areas to the Control of Motor Sequences. *J.*

Neurosci. **42**, 6946-6965 (2022).

37   Vallentin, D., Bongard, S. & Nieder, A. Numerical rule coding in the prefrontal,

premotor, and posterior parietal cortices of macaques. *J. Neurosci.* **32**, 6621-6630

(2012).

38   Yang, Y. P., Li, X. & Stuphorn, V. Primate anterior insular cortex represents economic

decision variables proposed by prospect theory. *Nat Commun* **13**, 717 (2022).

39   Kleiner, M., Brainard, D. & Pelli, D. What's new in Psychtoolbox-3? *Perception* **36**, 14-

14 (2007).

40   Glasser, M. F. *et al.* The minimal preprocessing pipelines for the Human Connectome

Project. *Neuroimage* **80**, 105-124 (2013).

41   Cusack, R. *et al.* Automatic analysis (aa): Efficient neuroimaging workflows and

parallel processing using Matlab and XML. *Frontiers in Neuroinformatics* **8** (2015).

42   Fischl, B. FreeSurfer. *Neuroimage* **62**, 774-781 (2012).

43   Jenkinson, M., Bannister, P., Brady, M. & Smith, S. Improved optimization for the

robust and accurate linear registration and motion correction of brain images.

Neuroimage **17**, 825-841 (2002).

44   Salimi-Khorshidi, G. *et al.* Automatic denoising of functional MRI data: combining

independent component analysis and hierarchical fusion of classifiers. *Neuroimage*

90, 449-468 (2014).

45   Robinson, E. C. *et al.* Multimodal surface matching with higher-order smoothness

constraints. *Neuroimage* **167**, 453-465 (2018).

46   Woolrich, M. W., Ripley, B. D., Brady, M. & Smith, S. M. Temporal autocorrelation in

univariate linear modeling of FMRI data. *Neuroimage* **14**, 1370-1386 (2001).

47   Genovese, C. R., Lazar, N. A. & Nichols, T. Thresholding of statistical maps in

functional neuroimaging using the false discovery rate. *Neuroimage* **15**, 870–878

(2002).

48   Cusack, R. *et al.* Automatic analysis (aa): efficient neuroimaging workflows and

parallel processing using Matlab and XML. *Front Neuroinform* **8**, 90 (2015).

49   McLaren, D. G. *et al.* A population-average MRI-based atlas collection of the rhesus

macaque. *Neuroimage* **45**, 52-59 (2009).

50   Henson, R. N. in *Brain mapping: An encyclopedic reference* Vol. 1 (ed A. W. Toga)

477-481 (Academic Press, 2015).

51   Zhou, Z.-X., & Xu, T. (2025). *RheMAP-Surf: Surface transformations among commonly*

used rhesus macaque brain templates (1.0). Retrieved

from: https://doi.org/10.5281/zenodo.15642802.

.

738