CRISPR-Cas9 and PiggyBac Mediated Genetic Modification of Sand Fly Vectors Targeting Olfactory and Non-Lethal Phenotypic Genes.

Authors

Rhodri T. M. Edwards[1], Luke Brandner Garrod[1], Tapan Bhattacharyya[1], Barbora Vomáčková Kykalová[2], Erich Loza Telleria[2], Matthew E. Rogers[1], Thomas Walker[3], Petr Volf[2] and Matthew Yeo[1†]

Affiliations

[1] Department of Infection Biology, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK
[2] Department of Parasitology, Faculty of Science, Charles University, Viničná 7, 128 44, Prague, Czech Republic.
[3] School of Life Sciences, University of Warwick, Gibbet Hill Campus, Coventry, CV4 7AL, UK

[†]To whom correspondence should be addressed:
Matthew Yeo, Ph.D.
Department of Infection Biology, London School of Hygiene and Tropical Medicine, Keppel Street, London WC1E 7HT, UK
Ph:
Email: matthew.yeo@lshtm.ac.uk

Keywords

Lutzomyia longipalpis, *Phlebotomus papatasi*, CRISPR, PiggyBac, Gene editing, Olfaction

Abstract

Leishmania spp. are protozoan parasites transmitted by female sand flies (Diptera) that cause a spectrum of devastating human pathologies affecting millions. Control of leishmaniases has proved immensely difficult: vector control strategies remain challenging and available chemotherapeutics imperfect. Gene editing of insect vectors offers prospects to interrupt disease transmission; for example by introducing antiparasite effectors or heritable modifications of genes implicated in fecundity. Here we present convergent evidence for successful gene editing using both CRISPR–Cas9 and PiggyBac approaches in two medically important sand fly species (*Lutzomyia longipalpis* and *Phlebotomus papatasi*). Targeted mutagenesis in G_0 and G_1 generations is supported by three independent lines of evidence, namely observable phenotypic changes, PCR-based detection assays, and confirmatory *in silico* algorithmic analysis of gene sequences which together provide a robust demonstration of targeted genome editing.

Introduction

CRISPR has transformed gene editing in insects of medical and veterinary importance due to a highly targeted and relatively simple application in comparison to previous methods (Dong et al. 2015). An important utility of CRISPR-Cas9 are the approaches to interrupt disease transmission. CRISPR-Cas9 based gene-drives that enable rapid spread of introduced heritable traits in fast-reproducing insect populations is a promising approach (Bier 2022). Two favoured applications of gene-drives are i) targeting genes that affect fecundity to suppress populations below the threshold required for parasite transmission (Hammond, Pollegioni, et al. 2021; Hammond et al. 2016), and ii) promoting population modification through the expression of introduced anti-parasite effectors that confer resistance to associated pathogens (Adolfi et al. 2020). Both approaches show great promise in laboratory and large-cage settings.

Despite rapid progress in gene editing methods in medically important mosquito vectors, including *Aedes* sp. (Anderson et al. 2023)*., Anopheles* sp. (D'Amato et al. 2024)*., and *Culex* sp. (Harvey-Samuel et al. 2023), advances in other medically important insect vectors remain limited, including sand flies that transmit the protozoan parasite *Leishmania*. Presently, control of sand flies, including *Lutzomyia longipalpis* and *Phlebotomus papatasi,* remains challenging (Kumari et al. 2025) due in part to their highly dispersed, low density terrestrial breeding sites (Balaska et al. 2021). Chemotherapeutics to treat leishmaniasis are imperfect and often not available to those in need (eBioMedicine 2023). Therefore, demonstration of gene editing of endogenous genes and the insertion of exogenous genetic cargo is an important step toward advancing sand fly biology and exploring new opportunities to control leishmaniasis, including gene drive development.

Currently there is a relative dearth of published research on the application of CRISPR-Cas9 in sand fly vectors. Optimisation of embryonic microinjection techniques for CRISPR-Cas9 editing has been described (Martin-Martin et al. 2018; Jeffries et al. 2018; Louradour et al. 2020). CRISPR-Cas9 mediated transgenesis has been demonstrated in a single landmark study targeting a *P. papatasi* transcription factor (NF-κB) and the Relish gene (*Rel*), a component of the immune deficiency (IMD) pathway (Louradour et al. 2019). In that report, recombinant Cas9 protein was directly injected together with sgRNAs derived by *in vitro* transcription to achieve targeted knockouts. The resulting *Rel* knockout colonies were difficult to maintain, and the number of injected embryo survivors was low, although efficiency of transgenesis was relatively high. To date, there are no published reports of successful genetic modification in *L. longipalpis*, the primary vector of visceral leishmaniasis in the Americas, using either CRISPR-mediated transgenesis or PiggyBac transformation.

Here we demonstrate the microinjection of CRISPR plasmid constructs and assess subsequent gene edits in two major sand fly vectors, providing the first demonstration of targeted CRISPR-induced mutagenesis in *L. longipalpis,* resulting in phenotypic mutation, as well as mutagenesis of olfactory genes involved in host seeking. Additionally, we report PiggyBac-mediated cargo insertion with subsequent inheritance in both *L. longipalpis* and *P. papatasi* following large-scale embryo microinjection. We provide convergent lines of evidence for successful mutagenesis in emergent transfected sand flies, namely targeted phenotypic changes, PCR-based detection assays, and confirmatory *in silico* algorithmic analyses of sequence data. Together, these findings provide robust support for the feasibility of gene editing in sand flies.

Methods

Identification of gene targets to affect

Candidate genes were chosen based on the likelihood of a phenotypic effect and non-lethality upon knockout. Genes with conserved function across multiple insect orders were selected using FlyBase (flybase.org) and VectorBase (vectorbase.org), and orthologues within *L. longipalpis* and *P. papatasi* identified. Orthologues were assessed via Clustal Omega (ver. 1.2. 3.) protein alignment against model organisms to infer gene ontology searching on AmiGO 2 (ver. 2. 5. 17) using template genomes lonJ1.6. and papI1.6. hosted on VectorBase. Briefly, candidate target genes likely to elicit a non-lethal phenotypic change were *ebony* (*e*), *vestigial* (*v*), *and rudimentary* (*r*). Separately candidate olfactory genes implicated in host seeking were chosen including *Orco*, *Gr2* and *Ir8a* (Supplementary Table S1).

Identification of gRNAS

ChopChop (Labun et al. 2021) was used to identify gRNA sequences using standard settings, which were selected based on efficiency scores (Doench et al. 2014). Functional gRNAs were assessed by an *in vitro* cleavage assay confirming activity. gRNAs were transcribed using the Engen Transcription kit (NEB, USA) following the manufacturer's protocols. Synthesised sgRNAs ability to cleave was assessed against amplified template DNA. Successful cleavage evidenced by digestion of template DNA was visualised by gel electrophoresis (Supplementary Figure S2).

CRISPR and piggyBac plasmids

For CRISPR induced knockouts the pDCC6 vector served as the backbone. The plasmid carries a gRNA and Cas9 binding scaffold expressed by a *Drosophila* U6 pol III promoter for RNA transcription and a human-codon-optimized Cas9 coding sequence (hCas9) promoted by an hsp70 pol II promoter (Gokcezade et al. 2014). pDCC6 was modified to produce a panel of plasmids targeting different genes (Supplementary Table S3). The dU6-2 (FBgn0266758) promoter was replaced with an endogenous U6 promoter (*L. longipalpis* and *P. papatasi*) via Gibson Assembly following manufacturer's instructions (Gibson Assembly® Cloning Kit, NEB, USA). The region 400bp upstream of the U6 start codon encompassed key promoter sequences including the pol III proximal sequence element A (PSEA) and TATA box within this region. Insertion of the U6 promoters was confirmed via PCR and Sanger sequencing. Adapted plasmids are here on referred to as Llon1-pDCC6 and Ppap1-pDCC6. gRNA's targeting genes of interest were inserted into the plasmid by Gibson Assembly to produce a panel of knockout plasmids (Supplementary Table S3). pDCC6 was a gift from Peter Duchek (Addgene plasmid # 59985 ; http://n2t.net/addgene:59985 ; RRID:Addgene_59985).

For piggyBac integration the Ubiq-Cas9.874W and pHome-T plasmids were selected. Ubiq-Cas9.874W contains a Ubi-63E promoter driving expression of hSpCas9 fused to GFP via a T2A self-cleaving peptide, and a fluorescent DsRed1 marker under control of the OpIE2 promoter. pHome-T expresses GFP under the synthetic promoter 3xP3 and RFP under control of the Ac5 promoter. Ubiq-Cas9.874W and pHome-T were gifts from Dr Roberto Galizi, (Keele University, UK) and Professor Tony Nolan (Liverpool School of Tropical Medicine, UK).

Two helper plasmids mhyPBase and ihyPBase, both hyperactive transposases, were delivered with piggybac plasmids containing a mammalian based transposase and Drosophila based hyperactive transposase respectively (Wright et al. 2013; Eckermann et al. 2018). Both hyperactive transposase plasmids were a gift from Dr Ernst Wimmer and Dr Hassan M.M. Ahmed (University of Gottingen, Germany).

Rearing of sand flies

Sand fly colony was maintained as described previously (Volf and Volfova 2011). Briefly, *L. longipalpis* and *P. papatasi* adults were housed in cages, with access to 30% sugar solution at 26 - 28°C, on a 14 hr:10 hr light-dark cycle. Blood-fed females were separated into small cages for 5-6 days to defecate, gravid females were aspirated into plastic pots to lay eggs 6-10 days post blood-feeding and removed once oviposition had occurred. Larvae hatched 6-10 days later, with the larval period typically lasting ~3 weeks followed by pupation for about 10 days Larval pots were placed within plastic boxes containing a base of moist sand (Lawyer et al. 2017).

Microinjections

A 10µl stock injection mixture was prepared prior to each injection session and incubated at 37°C for 10-20 minutes, before being placed on ice for 20 minutes prior to injections. 2µl was loaded into the needle (G-1 borosilicate glass capillaries, outer diameter 1mm; inner diameter 0.6mm, Narishige, Japan). For PiggyBac transformation lines, mixtures consisted of a PiggyBac construct and hyPBase transposase helper plasmid in a 1:2 ratio. Injection mixtures for UbiqCas9.874W and pHome-T plasmids were prepared in the same manner (Supplementary Table S4).

For CRISPR knockouts, plasmids (pDCC6 variants) targeting the same gene were inoculated simultaneously. Three *Rudimentary*-targeting, and three *vestigial*-targeting constructs were combined in an even ratio to derive a final concentration of 317.58ng/µL. Separately, six olfactory-targeting plasmids were combined into a single injection mixture containing an even ratio of each to a final concentration of 316.68ng/µL. Additionally four ebony pDCC6 constructs were combined for a concentration of 278.48ng/µL (Supplementary Table S4).

Injections were performed using a manual Narishige IM-9B microinjector and Narishige MMO-4 micromanipulator (Narishige, Japan) coupled to an inverted microscope (Leica DMi8, Germany). Each egg was injected between 1/4 and 1/3 of the length of the egg from the posterior pole (Jeffries et al. 2018) . Once injected, glass slides containing the eggs were transferred to a petri dish containing moist filter paper, and maintained at 25-28°C, 80% relative humidity. After ~1-3 days eggs were brushed from the glass slides into oviposition pots.

Identification of transgenic insects

A combination of methods were used to assess mutagenesis in transfected G0 survivors and subsequent G1 offspring. For emergent survivors transfected with plasmids inserting fluorescent cargo (PiggyBac plasmids) or targeting genes implicated in phenotypic change (CRISPR knockout plasmids) microscopy was performed. Secondly a PCR detection approach to identify mutagenesis was performed by T7

Endonuclease I Heteroduplex analysis. Thirdly a confirmatory *in silico* approach by Algorithmic deconvolution of Sanger sequence data was undertaken to predict indels.

Fluorescent microscopy

Emergent transgenic insects were imaged by fluorescence stereo microscope (Leica M205 FA, Germany). L1-L4 larvae were imaged to identify fluorescence for GFP (485nm) and EGFP (488nm). DsRed1 and RFP markers imaged at 558nm and 587nm respectively. Positive outcomes for injection survivors of *UbiqCas9.874W* PiggyBac transfections is fluorescent mosaicism indicative of plasmid integration and expression (Opie2 promoter of DsRed), and green fluorescent mosaicism indicating expression of Cas9 (Cas9 tagged with GFP, separated by T2A ribosome skipping element).

DNA extraction and PCR amplification.

50% of emergent larvae were harvested for DNA extraction (DNeasy Blood and Tissue Kit, Qiagen, Germany). Sand flies without phenotypic changes by microscopy were pooled in batches of 10. DNA was extracted from individual sand flies if there was visual evidence of phenotypic change such as changes to wing morphology (genes targeting wing development or pigmentation). The remaining 50% were left to pupate until adults emerged for subsequent backcrossing. PCR amplification of extracted genomic DNA was performed in 20µl Phusion HF Master mix according to the manufacturer's instructions (NEB, USA).

Outcrossing/backcrossing

G0 male and female *L. longipalpis* injected with wing-targeting CRISPR constructs were sibling crossed, as were *L. longipalpis* individuals transfected with CRISPR-based olfactory constructs (Supplementary Table S3).

Additionally, two rounds of PiggyBac UbiqCas9.874W microinjections were performed. For the first round (*L. longipalpis and P. papatasi*), 50% of larvae were allowed to pupate and eclose (the remaining 50% of larvae had DNA extracted). G0 adults were separated by sex and out crossed with wildtype adults. Subsequent G1 larvae were microscopically examined for fluorescent phenotypes and DNA extracted for sequencing if they did not survive to G1 adults. In a second round of UbiqCas9.874W microinjections, all *L. longipalpi*s larvae were allowed to pupate and eclose, with adult survivors outcrossed with wildtype.

Detecting mutagenesis by T7 Endonuclease I Heteroduplex assays

Mutagenesis was investigated using a T7 Endonuclease I Heteroduplex assay (T7EI, NEB, USA) according to the manufacturer's instructions. Initially PCR amplifies the target region through successive rounds of denaturation and annealing with heteroduplexes forming between amplicons containing a modified locus and those with an unmodified locus. The subsequent pool of amplicons encompasses homoduplexes of unmodified DNA and heteroduplexes of mutated DNA. T7 Endonuclease recognizes mismatches (heteroduplexes) inducing double-strand breaks (DSB) with fragments detected by Gel electrophoresis followed by subsequent Densitometric analysis of amplicons in Image Lab software (Bio-Rad, USA). Fragments generated by T7EI cleavage, were defined using the Lane Profile tool (background excluded) and the Analysis Table function to generate the Adjusted volume (background excluded volume) and the Band Percent (percent of the adjusted volume of the

selected band, relative to the other selected bands in the lane (% density of the sum of the Adjusted volume).

ICE analysis, algorithmic deconvolution analysis of sequence data to detect mutagenesis

Sanger sequencing chromatograms were analyzed using ICE (Inference of CRISPR Edits), an algorithmic trace-deconvolution method that estimates CRISPR editing outcomes from mixed amplicon populations by comparing edited samples to unedited control traces. ICE analysis was performed *in silico* using the Synthego Performance Analysis (ICE Analysis; v3.0, 2019). Briefly, Sanger files were uploaded together with the relevant guide RNA sequences. The software inferred editing efficiency and the spectrum of indels for knockout experiments. Workflow was used to assess intended sequence insertion/replacement events (up to ~200 bp).

Results

Microinjection Efficiency and Survival

A total of 7,623 sand fly embryos (*L. longipalpis* and *P. papatasi*) were injected with PiggyBac transposon constructs to integrate GPF or DsRed1 markers or CRISPR plasmids (Table 1) targeting wing-development, pigmentation, immunity, or olfactory genes (Table 2). Survival rates were comparable to those previously published, with adult survival as a percentage of eggs injected from 1.04 - 4.81%.

Evidence of PiggyBac Integration of Cas9 sequences

Microinjection with two different PiggyBac plasmids pHOME-T and UbiqCas9.874W into *L. longipalpis* and *P. papatasi* embryos resulted in successful genomic integration, detectable at the molecular level (via PCR detection and Sanger sequencing) in both the G_0 and G_1 generations, despite the absence of 3xP3-driven GFP or Actin5C driven RFP fluorescent marker expression in emergent individuals. In more detail, for pHOME-T, molecular screening identified genomic integration in G_0 pooled larval samples of both vector species. Germline transmission was also confirmed through analysis of G_1 progeny by Sanger sequencing. In *L. longipalpis*, 9 of 60 pooled G_1 larval samples were GFP-positive by PCR, with 7 of these 9 further validated by Sanger sequencing. In *P. papatasi*, 7 of 14 pooled G_1 larval samples were GFP-positive, with all seven confirmed by sequencing. Together, these results demonstrate both integration and inheritance of the pHOME-T cargo.

For UbiqCas9.874W, PCR and sequencing confirmed integration of Cas9, GFP, or DsRed1 sequences in G_0 larvae and adults of both sand fly species. Germline transmission was observed in *L. longipalpis*, where 33 of 52 G_1 pooled larval samples were PCR-positive for Cas9, and 15 of these were confirmed by sequencing (Supplementary Figure S5). For G0 *P. papatasi* injected with UbiqCas9.874W and IhyPBase (1,827 injected), two pooled samples of 10 larvae and two adult samples were PCR positive for Cas9, and two single adult samples were also positive for Cas9 via PCR and sequencing. No G_1 positives were detected for *P. papatasi*, in line with limited G_0 survival in this species. No DsRed1 or EGFP expression was detectable.

These data provide the first demonstration that PiggyBac cargo can integrate into sand fly genomes of both species tested and that exogenous cargo can be inherited through the germline. The current work validates these constructs for genetic manipulation in both *L. longipalpis* and *P. papatasi*.

Evidence for CRISPR-Cas9 Induced mutagenesis.

Phenotypic Mutagenesis

Evidence of mutagenesis affecting wing development (*L. longipalpis*) in three G1 individuals exhibited observable wing deformities, namely 1 with unilateral wing loss and the remaining two with pointed wings consistent with expected CRISPR disruptions (Figure 1). Importantly, phenotypes were not only observed visually but were also supported by DNA-level evidence of CRISPR mutagenesis, although the strength of molecular confirmation varied among individuals (Table 2). The individual with unilateral wing loss presented T7E1 heteroduplex cleavage (R2/R3 target) with subsequent support from ICE analysis (R3 target) comprising strong evidence of CRISPR targeted mutagenesis. Two additional flies (pointed wings) also demonstrated T7E1 mismatch evidence, but without supporting ICE analysis, due to sequence quality, and so a more moderate evidence level. Overall, phenotypic mutants had molecular support from at least one assay.

Molecular Evidence of CRISPR Mutagenesis

T7 Endonuclease I Heteroduplex Assays Reveal Evidence of CRISPR Editing

Sanger sequencing alone is not reliable to identify CRISPR-Cas9 modification where the dominant sequence can mask minority modified alleles. Therefore, to detect minority insertions and deletions generated by CRISPR editing, T7 endonuclease I (T7EI) assays were performed on PCR amplicons spanning targeted cut sites in *L. longipalpis* G_0 and G_1 individuals.

For wing-targeting constructs, amplicons were successfully obtained for all G_1 samples at the R3 (534 bp), V1 (577 bp), and V2 (567 bp) loci. Heteroduplex cleavage was observed in six R3 samples (R3C*, R3M*, R3N*, R3U*, R3V*, R3Z*) and in six V1 and nine V2 samples (Figure 2), each producing digestion fragments between 200–400 bp consistent with CRISPR-derived mismatches.

In olfactory-gene targeted flies, T7EI assays similarly provided evidence of mutagenesis. Among G_0 individuals, six Gr2 and four Ir8a amplicons showed digestion consistent with heteroduplex formation, while in G_1 samples, T7EI activity was detected across *Gr2, Ir8a,* and *Orco* loci. Notably, nine male and nine female G_1 samples exhibited heteroduplex digestion for *Orco*, with fragments between 200–600 bp.

Densitometric Quantification Indicates Low-to-Moderate Modification Levels

To estimate the proportion of modified alleles within each amplicon pool, densitometric analysis was applied to T7EI digests. For wing-gene targets, modification rates at R3 ranged from 0.36–6.13%, whereas V1 and V2 targets showed broader editing ranges, from 2.98–18.51% and 0.64–15.06%, respectively. Background digestion in controls remained consistently low (0.05–4.43%) and is highly suggestive that elevated cleavage intensities in experimental samples reflected genuine CRISPR editing rather than nonspecific nuclease activity.

Olfactory-gene knockouts showed similarly variable evidence of modification. In G_0 flies, densitometry estimated 0.68–7.52% modification at Gr2 and 0.48–3.91% at Ir8a, while G_1 samples demonstrated

wider ranges. Gr2 modification ranged from 0.19% to 57.64%, and Orco modification ranged from 0.44–9.95%, exceeding background digestion levels in named samples.

ICE Sequence Deconvolution Independently Supports CRISPR Editing

Inference of CRISPR Edits (ICE) analysis provided an orthogonal method to detect indels directly from Sanger sequencing traces.

For wing-targeting constructs, ICE predicted indels at the R2 cut site in 16 of 26 G_1 individuals, including those displaying wing deformities, and in two individuals at R3.

Olfactory gene analyses further support editing events. ICE detected indels in 8 of 11 G_0 individuals and in 17 of 34 G_1 individuals at Gr2, with 10 of these 17 showing matching T7EI cleavage, representing strong evidence of true gene editing. For Ir8a, 17 G_1 samples showed ICE predicted indels, and three were corroborated by T7EI. For Orco, 25 G_1 individuals carried ICE predicted indels, including one (G1-MA) with exceptionally strong support ($R^2 = 0.99$, KO-score = 1) and concordant T7EI digestion.

Together, the T7EI heteroduplex assays, densitometric quantification, and ICE-based sequence analysis provide strong molecular evidence of CRISPR-mediated mutagenesis in both G_0 and G_1 *L. longipalpis*. Although individual assays varied in sensitivity and specificity, the combined dataset reveals consistent patterns of indel formation across multiple target genes, with editing detected at wing-development loci (R3, V1, V2) and olfactory genes (Gr2, Ir8a, Orco). Crucially, numerous individuals exhibited concordance between two or more methods, including 18 instances where T7EI and ICE jointly identified indels, providing robust confirmation that CRISPR-Cas9 delivered via plasmid microinjection successfully induced heritable genome editing in sand flies.

Summary of Editing Outcomes

PiggyBac-Cas9 delivery using two different plasmid constructs (UbiqCas9.874W and pHOME-T) successfully produced germline-transmitting lines. The former verified Cas9 integration in G0 and G1 individuals, and the latter GFP expression in (G1), confirmed by Sanger sequencing. Wing-development genes showed both genotypic mutations and heritable phenotypic defects. For CRISPR - induced mutagenesis, olfactory genes (*Gr2, Ir8a, Orco*) showed clear evidence of CRISPR editing in G0 and G1 individuals (Table 2). Overall, these results demonstrate the first successful *in vivo* genetic modification of *L. longipalpis* and *P. papatasi* using PiggyBac-mediated integration of exogenous DNA, as well as the first reported CRISPR-Cas9 knockout mutagenesis in *L. longipalpis*.

Discussion

Functional genetic tools for phlebotomine sand flies significantly lag behind those available for other insects of medical importance, particularly mosquito vectors, which has hampered progress toward development of novel genetic vector control strategies for leishmaniasis. This is despite the continued epidemiological impact of the disease.

In recent years a plethora of publications have described CRISPR-Cas9 gene editing applied to insects of medical, veterinary and agricultural importance, stemming from initial demonstration in *Drosophila*

melanogaster (Gantz and Akbari 2018; Gratz et al. 2014; Gokcezade et al. 2014). Notable examples include Anopheline vectors of malaria, *Aedes* vectors of dengue and Zika viruses, and major crop pests *Plutella xylostella* and *Ceratitis capitata,* amongst many others (Gantz and Akbari 2018).

Early publications often described species-specific protocols exploring functional genomics in the context of gene knockouts (Kistler et al. 2015; Basu et al. 2015; Gantz et al. 2015; Hammond et al. 2016). More recently, refined homology-directed repair approaches have been developed in mosquito species to facilitate gene drives, enabling the spread of introduced heritable traits through populations at levels exceeding those achievable by Mendelian inheritance. Broadly, there are two main implementations of gene-drives applied to insects of medical importance, both interrupting disease transmission. The first, population suppression strategies, attempt to reduce the vector populations below that required to sustain pathogen transmission (Hammond et al. 2016; Simoni et al. 2020b)(Simoni et al. 2020a; Bernardini et al. 2018). Typically, the strategy introduces lethal or deleterious self-propagating traits affecting overall reproductive fitness leading to drastic reduction in local populations. An alternative approach, referred to as population replacement or modification, is transmission blocking, for example expression of antiparasite peptides that target vector genes implicated in establishing or transmitting infection to derive pathogen-resistant vectors (Gantz et al. 2015; Williams et al. 2020). Each approach has its benefits and drawbacks. Technical challenges remain (Hammond, Karlsson, et al. 2021; Garrood et al. 2021)(Beaghton et al. 2019); however, both approaches offer exciting potential to deliver tangible results and show great promise in laboratory settings as well as large-scale caged studies (Bier 2022; Hammond, Pollegioni, et al. 2021).

PiggyBac is the most widely used transposon system for genetic manipulations in insects, successfully applied to many different mosquito vectors to express fluorescent markers and anti-parasite peptides (Nolan et al. 2002; Grossman et al. 2000)(Ito et al. 2002; Nolan et al. 2002)(Perera et al. 2002)), in *Aedes aegypti* (Kokoza, Ahmed, Wimmer, & Raikhel, 2001), *Ae. albopictus* (Labbé, Nimmo, & Alphey, 2010), and *Ae. fluviatilis* (Rodrigues, Oliveira, Rocha, & Moreira, 2006). PiggyBac plasmids can carry CRISPR-Cas components as cargo for insertion into genomes (Li et al. 2017). Importantly, the technique has not been demonstrated in sand flies which is a significant omission for such a potentially useful tool.

In the current study, we establish a robust *in vivo* framework for genetic modification of *L. longipalpis* and *P. papatasi*. We provide the first evidence of PiggyBac-mediated transgenesis in *L. longipalpis* and *P. papatasi* and also CRISPR-Cas9-based targeted mutagenesis in *L. Longipalpis,* the major vector of visceral leishmaniasis in the Americas.

For PiggyBac constructs incorporating a ubiquitin-driven Cas9 cassette, Cas9 and GFP sequences were detected in both G0 and G1 individuals, together indicating successful germline transmission and establishing PiggyBac as a viable transformation system in sand flies. This finding addresses an important technical limitation highlighted by previous unsuccessful attempts described in the literature for *L. longipalpis* (Martin-Martin et al. 2018; Jeffries et al. 2018; Louradour et al. 2019). Building on this, we describe a functional approach to support future gene editing studies.

In parallel, the results provide multiple lines of evidence supporting CRISPR-Cas9-mediated mutagenesis that target both phenotypic marker genes and olfactory genes.

Molecular analyses by Sanger sequencing suggest that mosaicism is a dominant feature of CRISPR editing in sand flies, as expected given plasmid-based delivery during early embryogenesis. Overall,

standard Sanger sequencing alone proved insufficient to reliably detect low-frequency edits. However, the combined application of T7 endonuclease I assays, densitometric analysis, and in silico ICE deconvolution applied to Sanger sequences provided robust and reproducible evidence of targeted mutagenesis by application of CRISPR based approaches. Considering that rapid pre-cellular nuclear divisions occur shortly after oviposition in dipterans, reducing the 4-hour time window between collecting eggs and injecting them, if logistically possible, may help reducing mosaicism (Jeffries et al. 2018).

Targeting olfactory genes (*Gr2*, *Ir8a*, and *Orco*) implicated in chemosensory function, host-seeking behaviour, and the maintenance of vectorial capacity across Diptera, confirms that key behavioural pathways are, in principle, genetically accessible in sand flies. These findings provide a foundation for investigating the modulation of host seeking behaviour and exploring strategies to interrupt parasite transmission.

Low recovery of phenotypically visible mutants likely reflects a combination of mosaicism, developmental lethality or impaired adult fitness, together with the intrinsic challenges of sand fly embryo microinjection. Although phenotypic mutants were recovered at low frequency, observed wing malformations and pigmentation changes were consistent with predicted orthologous knockouts in mosquitoes. In this context, edits supported by molecular evidence were biologically consistent with disruption of vestigial (*vg*) and rudimentary (*r*).

In summary, this work establishes the first integrated approach for sand fly genetic modification, bridging bioinformatic target identification, the successful application of molecular tools incorporating piggyBac and CRISPR based approaches, and subsequent *in vivo* validation. The recent availability of improved assemblies for *P. papatasi* and *L. longipalpis* (Huang et al. 2024) will facilitate future targeting efficiency. Together, these results represent a critical step toward the application of advanced genetic control strategies for leishmaniasis vectors and will contribute to ongoing behavioural, ecological, and importantly, gene-drive development to facilitate vector control.

Acknowledgments.

We are grateful to Hassan M. M. Ahmed, Ernst Wimmer, Tony Nolan and Roberto Galizi for the donation of plasmids.

Author contributions

M.Y., conceptualised the study; R.T.M.E., L.B.G., B.V.K., and E.L.T. performed all experiments; M.Y., R.T.M.E., L.B.G., B.V.K., and E.L.T. performed data analysis; M.Y., and R.T.M.E., wrote the first draft of the manuscript; T. B. reviewed and edited an advanced draft of the manuscript; M.Y. T.W., M.E.R., and P.V provided resources and supervision; All authors contributed to analysing and compiling the data, writing and approving the final manuscript.

Funding

M.Y., R.T.M.E., and L.B.G. were supported by the Wellcome Trust, UK (208913/Z/17/Z); R.T.M.E., was additionally supported by and Medical Research Council LID PhD studentship (MR/N013638/1); P.V. and E.L.T. were supported by Czech Science Foundation (GACR, project No. 25-15318S).

References

Adolfi, Adriana, Valentino M. Gantz, Nijole Jasinskiene, et al. 2020. "Efficient Population Modification Gene-Drive Rescue System in the Malaria Mosquito Anopheles Stephensi." *Nature Communications* 11 (1): 5553.

Anderson, Michelle A. E., Estela Gonzalez, Joshua X. D. Ang, et al. 2023. "Closing the Gap to Effective Gene Drive in Aedes Aegypti by Exploiting Germline Regulatory Elements." *Nature Communications* 14 (1): 338.

Balaska, Sofia, Emmanouil Alexandros Fotakis, Alexandra Chaskopoulou, and John Vontas. 2021. "Chemical Control and Insecticide Resistance Status of Sand Fly Vectors Worldwide." *PLoS Neglected Tropical Diseases* 15 (8): e0009586.

Basu, Sanjay, Azadeh Aryan, Justin M. Overcash, et al. 2015. "Silencing of End-Joining Repair for Efficient Site-Specific Gene Insertion after TALEN/CRISPR Mutagenesis in Aedes Aegypti." *Proceedings of the National Academy of Sciences of the United States of America* 112 (13): 4038–4043.

Beaghton, Andrea K., Andrew Hammond, Tony Nolan, Andrea Crisanti, and Austin Burt. 2019. "Gene Drive for Population Genetic Control: Non-Functional Resistance and Parental Effects." *Proceedings. Biological Sciences / The Royal Society* 286 (1914): 20191586.

Bernardini, Federica, Roya Elaine Haghighat-Khah, Roberto Galizi, Andrew Marc Hammond, Tony Nolan, and Andrea Crisanti. 2018. "Molecular Tools and Genetic Markers for the Generation of Transgenic Sexing Strains in Anopheline Mosquitoes." *Parasites & Vectors* 11 (Suppl 2): 660.

Bier, Ethan. 2022. "Gene Drives Gaining Speed." *Nature Reviews. Genetics* 23 (1): 5–22.

D'Amato, Rocco, Chrysanthi Taxiarchi, Marco Galardini, et al. 2024. "Anti-CRISPR Anopheles Mosquitoes Inhibit Gene Drive Spread under Challenging Behavioural Conditions in Large Cages." *Nature Communications* 15 (1): 952.

Doench, John G., Ella Hartenian, Daniel B. Graham, et al. 2014. "Rational Design of Highly Active sgRNAs for CRISPR-Cas9-Mediated Gene Inactivation." *Nature Biotechnology* 32 (12): 1262–1267.

Dong, Shengzhang, Jingyi Lin, Nicole L. Held, Rollie J. Clem, A. Lorena Passarelli, and Alexander W. E. Franz. 2015. "Heritable CRISPR/Cas9-Mediated Genome Editing in the Yellow Fever Mosquito, Aedes Aegypti." *PloS One* 10 (3): e0122353.

eBioMedicine. 2023. "Leishmania: An Urgent Need for New Treatments." *EBioMedicine* 87

(January): 104440.

Eckermann, Kolja N., Hassan M. M. Ahmed, Mohammad KaramiNejadRanjbar, et al. 2018. "Hyperactive piggyBac Transposase Improves Transformation Efficiency in Diverse Insect Species." *Insect Biochemistry and Molecular Biology* 98 (July): 16–24.

Gantz, Valentino M., and Omar S. Akbari. 2018. "Gene Editing Technologies and Applications for Insects." In *Current Opinion in Insect Science*, vol. 28, 28. Preprint. https://doi.org/10.1016/j.cois.2018.05.006.

Gantz, Valentino M., Nijole Jasinskiene, Olga Tatarenkova, et al. 2015. "Highly Efficient Cas9-Mediated Gene Drive for Population Modification of the Malaria Vector Mosquito Anopheles Stephensi." *Proceedings of the National Academy of Sciences of the United States of America* 112 (49): E6736–43.

Garrood, William T., Nace Kranjc, Karl Petri, et al. 2021. "Analysis of off-Target Effects in CRISPR-Based Gene Drives in the Human Malaria Mosquito." *Proceedings of the National Academy of Sciences of the United States of America* 118 (22). https://doi.org/10.1073/pnas.2004838117.

Gokcezade, Joseph, Grzegorz Sienski, and Peter Duchek. 2014. "Efficient CRISPR/Cas9 Plasmids for Rapid and Versatile Genome Editing in Drosophila." *G3* 4 (11): 2279–2282.

Gratz, Scott J., Fiona P. Ukken, C. Dustin Rubinstein, et al. 2014. "Highly Specific and Efficient CRISPR/Cas9-Catalyzed Homology-Directed Repair in Drosophila." *Genetics* 196 (4): 961–971.

Grossman, G. L., C. S. Rafferty, M. J. Fraser, and M. Q. Benedict. 2000. "The piggyBac Element Is Capable of Precise Excision and Transposition in Cells and Embryos of the Mosquito, Anopheles Gambiae." *Insect Biochemistry and Molecular Biology* 30 (10): 909–914.

Hammond, Andrew, Roberto Galizi, Kyros Kyrou, et al. 2016. "A CRISPR-Cas9 Gene Drive System Targeting Female Reproduction in the Malaria Mosquito Vector Anopheles Gambiae." *Nature Biotechnology* 34 (1): 78–83.

Hammond, Andrew, Xenia Karlsson, Ioanna Morianou, et al. 2021. "Regulating the Expression of Gene Drives Is Key to Increasing Their Invasive Potential and the Mitigation of Resistance." *PLoS Genetics* 17 (1): e1009321.

Hammond, Andrew, Paola Pollegioni, Tania Persampieri, et al. 2021. "Gene-Drive Suppression of Mosquito Populations in Large Cages as a Bridge between Lab and Field." *Nature Communications* 12 (1): 4589.

Harvey-Samuel, Tim, Xuechun Feng, Emily M. Okamoto, et al. 2023. "CRISPR-Based Gene Drives Generate Super-Mendelian Inheritance in the Disease Vector Culex Quinquefasciatus." *Nature Communications* 14 (1): 7561.

Huang, Michelle, Sarah Kingan, Douglas Shoue, et al. 2024. "Improved High Quality Sand Fly Assemblies Enabled by Ultra Low Input Long Read Sequencing." *Scientific Data* 11 (1): 918.

Ito, Junitsu, Anil Ghosh, Luciano A. Moreira, Ernst A. Wimmer, and Marcelo Jacobs-Lorena. 2002. "Transgenic Anopheline Mosquitoes Impaired in Transmission of a Malaria Parasite." *Nature* 417 (6887): 452–455.

Jeffries, Claire L., Matthew E. Rogers, and Thomas Walker. 2018. "Establishment of a Method for Sand Fly Embryo Microinjection: The First Step towards Potential Novel Control Strategies for Leishmaniasis." *Wellcome Open Research* 3 (May): 55.

Kistler, Kathryn E., Leslie B. Vosshall, and Benjamin J. Matthews. 2015. "Genome Engineering with CRISPR-Cas9 in the Mosquito Aedes Aegypti." *Cell Reports* 11 (1): 51–60.

Kumari, Yasoda, Nayana Gunathilaka, and Deepika Amarasinghe. 2025. "A Comprehensive Review of Biological and Genetic Control Approaches for Leishmaniasis Vector Sand Flies; Emphasis towards Promoting Tools for Integrated Vector Management." *PLoS Neglected Tropical Diseases* 19 (1): e0012795.

Labun, Kornel, Maximilian Krause, Yamila Torres Cleuren, and Eivind Valen. 2021. "CRISPR Genome Editing Made Easy Through the CHOPCHOP Website." *Current Protocols* 1 (4): e46.

Lawyer, Phillip, Mireille Killick-Kendrick, Tobin Rowland, Edgar Rowton, and Petr Volf. 2017. "Laboratory Colonization and Mass Rearing of Phlebotomine Sand Flies (Diptera, Psychodidae)." *Parasite* 24 (November): 42. Originally published as *Établissement de Colonies de Laboratoire et élevage de Masse Des Phlébotomes (Diptera, Psychodidae).*

Li, Ming, Michelle Bui, Ting Yang, Christian S. Bowman, Bradley J. White, and Omar S. Akbari. 2017. "Germline Cas9 Expression Yields Highly Efficient Genome Engineering in a Major Worldwide Disease Vector,." *Proceedings of the National Academy of Sciences of the United States of America* 114 (49): E10540–E10549.

Louradour, Isabelle, Kashinath Ghosh, Ehud Inbar, and David L. Sacks. 2019. "CRISPR/Cas9 Mutagenesis in Phlebotomus Papatasi: The Immune Deficiency Pathway Impacts Vector Competence for Leishmania Major." *mBio* 10 (4). https://doi.org/10.1128/mBio.01941-19.

Louradour, Isabelle, Kashinath Ghosh, Ehud Inbar, David L. Sacks, Channa Aluvihare, and Robert A. Harrell 2nd. 2020. "Sand Fly (Phlebotomus Papatasi) Embryo Microinjection for CRISPR/Cas9 Mutagenesis." *Journal of Visualized Experiments : JoVE*, no. 165 (November). https://doi.org/10.3791/61924.

Martin-Martin, Ines, Azadeh Aryan, Claudio Meneses, Zach N. Adelman, and Eric Calvo. 2018. "Optimization of Sand Fly Embryo Microinjection for Gene Editing by CRISPR/Cas9." *PLoS Neglected Tropical Diseases* 12 (9): e0006769.

Nolan, Tony, Tom M. Bower, Anthony E. Brown, Andrea Crisanti, and Flaminia Catteruccia. 2002. "piggyBac-Mediated Germline Transformation of the Malaria Mosquito Anopheles Stephensi Using the Red Fluorescent Protein dsRED as a Selectable Marker." *The Journal of Biological Chemistry* 277 (11): 8759–8762.

Perera, O. P., R. A. Harrell II, and A. M. Handler. 2002. "Germ-Line Transformation of the

South American Malaria Vector, Anopheles Albimanus, with a piggyBac/EGFP Transposon Vector Is Routine and Highly Efficient." *Insect Molecular Biology* 11 (4): 291–297.

Simoni, Alekos, Andrew M. Hammond, Andrea K. Beaghton, et al. 2020a. "A Male-Biased Sex-Distorter Gene Drive for the Human Malaria Vector Anopheles Gambiae." *Nature Biotechnology* 38 (9): 1054–1060.

Simoni, Alekos, Andrew M. Hammond, Andrea K. Beaghton, et al. 2020b. "Author Correction: A Male-Biased Sex-Distorter Gene Drive for the Human Malaria Vector Anopheles Gambiae." *Nature Biotechnology* 38 (9): 1097.

Volf, P., and V. Volfova. 2011. "Establishment and Maintenance of Sand Fly Colonies." *Journal of Vector Ecology: Journal of the Society for Vector Ecology* 36 Suppl 1 (March): S1–9.

Williams, Adeline, Alexander Franz, William Reid, and Ken Olson. 2020. "Antiviral Effectors and Gene Drive Strategies for Mosquito Population Suppression or Replacement to Mitigate Arbovirus Transmission by Aedes Aegypti." In *Insects*, vol. 11, 11. no. 1. Preprint. https://doi.org/10.3390/insects11010052.

Wright, Jennifer A., Ryan C. Smith, Xianghong Li, Nancy L. Craig, and Peter W. Atkinson. 2013. "IPB7 Transposase Behavior in Drosophila Melanogaster and Aedes Aegypti." *Insect Biochemistry and Molecular Biology* 43 (10): 899–906.

Table 1. Survival for *L. longipalpis* and *P. papatasi* eggs injected with multiple plasmids. Counting larvae is difficult due to substrate and can result in undercounting in comparison to emergent sand flies, occasionally inferring adult survival as % of larvae being greater than 100% (indicated by *†*).

Species	Plasmid(s)	Injected eggs	Larval survival (%)	Adult survival Male	Adult survival Female	Adult survival as% of larvae	Adult survival as% of eggs injected
L. longipalpis	UbiqCas9.874W + IhyPBase	2058	75 **(3.64)**	27	15	56.0	2.04
P. papatasi	UbiqCas9.874W + IhyPBase	1827	44 **(2.41)**	29	13	95.45	2.30
L. longipalpis	pHome + IhyPBase	749	57 **(7.61)**	15	21	63.16	4.81
P. papatasi	pHome + IhyPBase	800	42 **(5.25)**	12	9	50.0	2.63
P. papatasi	pDCC6: Ebony1, Ebony2, Ebony3, Ebony4	1034	8 **(0.77)**	6	6	150†	1.16
L. longipalpis	pDCC6: R1, R2, R3, V1, V2, V3	579	6 **(1.04)**	3	3	100	1.04
L. longipalpis	pDCC6: Gr2-1, Gr2-2, Ir8a-1, Ir8a-2, Orco 12, Orco 34	576	10 **(1.74)**	9	2	110†	1.91

Table 2. Summary of evidence for CRISPR mutagenesis across gene targets (*Lutzomyia longipalpis* and *Phlebotomus papatasi*). Evidence types encompass phenotypic mutantagenesis, detection by T7 Endonuclease I heteroduplex cleavage assays, with subsequent densitometric quantification and *in silico* sequence analysis by ICE algorithmic indel prediction. Mutations were detected across both G0 and G1 generations depending on the target and method. Rudimentary, *Gr2*, and *Orco* demonstrate robust multi-method support indicative of mutagenesis.

Gene Target	Function / Class	Cut Sites / Constructs	Evidence of Mutagenesis	Generations	Comment
Rudimentary (r)	Wing development	R2, R3	ICE indels; T7E1; Wing phenotype mutants	G0, G1	Strong evidence across assays
Vestigial (vg)	Wing development	V1, V2, V3	T7E1 (V1,V2); limited ICE	G1	Sequencing limited V3 detection
Ebony (e)	Pigmentation	Ebony constructs	Darkened G0 pupae (phenotype)	G0	Phenotypic confirmation only
Gr2	Olfaction	Gr2-1, Gr2-2	ICE indels; T7E1	G0, G1	Strong olfactory evidence
Ir8a	Olfaction	Ir8a-1, Ir8a-2	ICE indels; T7E1	G0, G1	Moderate but consistent
Orco	Olfaction (co-receptor)	Orco-1, Orco-2	ICE indels; T7E1; High-confidence KO (G1-MA)	G0, G1	Most robust knockout evidence

Figure 1. G1 microinjection survivors with atypical wing phenotypes. Phenotypic characteristics of G1 male individual (sample G1-m6-Z) presenting a deformed wing phenotype, G1 female individual (sample G1-f3-C) presenting pointed wing tip phenotype, and G1 male (sample G1-m2-Z) undeveloped left wing. Scale bars 2.2mm.

Figure 2. *L. longipalpis* T7 Endonuclease I Heteroduplex assay and densitometric analysis (G1samples targeting V1 and V2 gRNAs regions). A) Samples V1X-V1Z followed by V2A-V2F, B) samples V2G-V2Q, and C) samples V2Q-V2Z. Asterisks (*) indicates PCR product and subsequent digestion with endonuclease. Lanes without asterisk are controls untreated with Endonuclease. For Densitometric analysis (an estimate of the proportion of modified alleles) blue boxes indicate boundaries of each lane, and pink lines indicate amplicons assessed within each lane.