Conformational ensembles of flexible multidomain proteins: How close are we to accurate and reliable predictions?

Santiago Rodriguez[1], Aurélie Fournet[1], Simon Bartels[2], Mátyás Pajkos[2], Ilinka Clerc[2], Laure Carrière[2], Aurélien Thureau[3], Cédric Montanier[4], Claire Dumon[4], Frédéric Allemand[1], Juan Cortés[2,*] and Pau Bernadó[1,*]

[1]*Centre de Biologie Structurale, Université de Montpellier, INSERM, CNRS, Montpellier, France.*

[2]*LAAS-CNRS, Université de Toulouse, CNRS, Toulouse, France.*

[3]*Synchrotron SOLEIL, L'Orme des Merisiers, Départementale 128, 91190 Saint-Aubin, France*

[4]*TBI, Université de Toulouse, CNRS, INRAE, INSA, Toulouse, France.*

[*]*Corresponding author: Juan Cortés (juan.cortes@laas.fr) and Pau Bernadó (pau.bernado@cbs.cnrs.fr)*

Abstract

Multidomain proteins connected by flexible linkers populate conformational ensembles that are challenging to characterize using conventional structural biology methods. In domain–linker–domain (DLD) proteins, linker-mediated inter-domain relative positions and orientations are functionally relevant, yet their dynamical behavior in solution normally remain poorly described. Small-angle X-ray scattering (SAXS) provides ensemble-averaged structural information for such systems; however, coupling with computational modeling is required to accurately describe the dynamic behavior of this family of proteins in solution. Here, we present a systematic evaluation of five ensemble-generation strategies applied to a set of eighteen proteins sharing the same two globular domains, connected by naturally occurring linkers of varying length and composition. Modeling methods based on different underlying principles are compared by assessing their agreement to experimental SAXS data, showing a large disparity and systematic structural biases among them. Furthermore, for each approach, we examine the effect of refinement against SAXS restraints and assess its capacity to describe the experimental data, as well as the induced biases in global dimensions and inter-domain distance distributions. This analysis underlines the importance of the initial conformational pool for deriving experimentally compatible ensembles. Overall, this work provides a high-quality benchmark for SAXS-driven ensemble modeling of flexible, multidomain proteins and establishes a framework for the critical interpretation of solution scattering data in systems with pronounced conformational heterogeneity.

Keywords — Multidomain proteins, Flexible linkers, Conformational ensembles, Small-angle X-ray scattering

1 Introduction

Many proteins exhibit multimodular architectures in which structured domains are connected by flexible or intrinsically disordered linkers [1, 2]. These domain–linker–domain (DLD) proteins are widespread across all kingdoms of life and play key roles in numerous biological processes [3–6]. The presence of flexible linkers enables functional adaptability by allowing domains to sample a wide range of relative orientations and distances, facilitating

molecular recognition, catalysis, and regulation [1, 6]. At the same time, this conformational heterogeneity poses significant challenges for structural characterization and quantitative modeling.

A prominent example of DLD architectures is found in enzymes involved in biomass degradation, where catalytic glycosyl hydrolase (GH) domains are often connected to carbohydrate-binding modules (CBMs) through flexible linkers (Figure 1). In these systems, the linker length and composition modulate the spatial relationship between catalytic and binding domains, directly influencing substrate targeting, enzymatic efficiency and overall function [7, 8]. As a result, GH–CBM chimeras have become important targets in enzyme engineering and biotechnology, motivating a deeper understanding of how linker properties shape their conformational ensembles in solution [7–9].

Traditional high-resolution structural biology techniques such as X-ray crystallography and cryo-electron microscopy have inherent limitations when applied to highly flexible or partially disordered proteins [10,11]. Although more adapted to the conformational plasticity of DLD [12–14], Nuclear Magnetic Resonance (NMR) often faces severe spectral overlap with large domains. As a consequence, flexible linkers often remain unresolved and the relative positions of the domains are unknown. In this context, small-angle X-ray scattering (SAXS) has emerged as a powerful technique that probes proteins in solution and reports on their ensemble-averaged structural properties [15, 16]. However, SAXS data are intrinsically low-resolution and do not provide unique structural solutions, making their interpretation strongly dependent on computational modeling [17–19]. Consequently, the generation of realistic conformational ensembles from sequence or structural information has become a central component in the analysis of SAXS data for flexible proteins [20, 21].

Molecular dynamics (MD) simulations using all-atom physics-based force field models have been widely used for the investigation of globular proteins. Although molecular force-fields and water models have dramatically improved in the recent years [22, 23], and new protocols have been developed to enhance the conformational sampling [24], MD simulations remain still limited to study large, highly flexible proteins [25]. Alternatively, a wide range of computational approaches has been developed for this purpose [26], including physics-based coarse-grained models [27, 28], structure-encoding libraries [29–31] and, more recently, methods derived from AlphaFold-based frameworks [32]. These approaches have demonstrated encouraging performance in various contexts, including community-wide benchmarks such as CASP [33]. Nevertheless, their general applicability and reliability for highly flexible, multidomain proteins remain uneven, and systematic evaluations focused on DLD architectures are still scarce.

Several key questions therefore remain open: How accurately do current ensemble generation methods reproduce experimental SAXS profiles for DLD proteins? To what extent and in which conditions can refinement against SAXS data improve the agreement between predicted and experimental observables? And do ensembles generated by different methods converge toward similar conformational descriptions after refinement, or do method-dependent differences persist? Addressing these questions is essential for establishing confidence in ensemble modeling strategies and for defining best practices in the interpretation of SAXS data.

In this work, we address these issues by systematically evaluating multiple computational approaches for generating conformational ensembles of GH11–CBM chimeras connected by linkers of diverse length and composition and for which we have recorded high-quality SAXS data. These profiles constitute an interesting benchmark set to assess the capacity of current ensemble generation methods to reproduce experimental SAXS profiles for DLD proteins. For each method, the quality of the resulting ensemble is evaluated by directly comparing with the experimental profile. Moreover, the structural impact of ensemble refinement against SAXS data is quantitatively assessed. By refining ensembles derived from distinct starting pools and comparing their resulting structural properties, we examine the consistency and robustness of the inferred conformational descriptions.

Beyond the specific protein family studied here, our results provide insights into the strengths and limitations of current ensemble modeling strategies for DLD proteins. Such understanding is critical not only for advancing fundamental studies of protein conformational behavior, but also for the rational design and optimization of multimodular enzymes with applications in biotechnology and industrial biocatalysis.

Figure 1: (A) Generic structure of the DLD proteins studied in this work. Cartoon representations correspond to the glycosyl hydrolase (GH; in green) and carbohydrate-binding module (CBM; in blue) domains, while the red dashed line represents the connecting linker, which varies in both length and composition across the different DLD systems. (B) Generic ensemble representation of a DLD protein. The relative pose (position and orientation) between the CBM and GH domains is defined by the linker.

2 Results

2.1 Protein system and linker diversity

For our benchmark set, we constructed a series of chimeric proteins consisting of the catalytic xylanase GH11 domain from *Neocallimastix patriciarum* (UniProt: P29127; PDB: 2C1F) connected to a CBM from *Cellulomonas fimi* (UniProt: P54865; PDB: 2XBD) through natural linker sequences extracted from the CAZy database [34]. From more than one thousand available protein fragments encompassing GH11–linker-CBM domains, we selected 18 linker sequences that were inserted between both globular domains for subsequent structural characterization. The selection was guided by sequence-based descriptors capturing key features of linker diversity, specially length and amino acid composition. This strategy ensured that sequences with diverse charge distribution, proline and glycine content or low complexity segments were represented in both short (10 residues) and long (>30 residues) linkers.

Table 1 presents the sequences and a set of descriptors of the selected linkers for the construction of the chimeras, which were named from DLD1 to DLD18. Physicochemical properties were computed using localCIDER [35]. The linkers sampled a broad range of sizes, from 10 to 88 residues. In terms of composition, while low complexity linkers were rich in glycines (DLD1, 4 and 8), prolines (DLD10 and 15), serines (DLD9 and 13) or asparagines (DLD16), others presented less compositional bias (DLD3, 7, 17 and 18). In terms of electrostatic charges, the vast majority of linkers were neutral, and only DLD3 (-2), DLD11 (+5) and DLD17 (-9) displayed a moderate global net charge. Thus, this panel of proteins captured the natural variability of GH11-associated linkers while providing a controlled system with identical folded domains.

2.2 Small-angle X-ray Scattering shows the flexibility of the DLD benchmark set

A plasmid was designed in order to facilitate the subsequent cloning of the 18 linkers between the GH and CBM domains (see Materials and Methods for details). The resulting chimeras were overexpressed

DLD	Linker sequence	Length	Hydrop.	Complex.	N.C.	κ	Rg	$D_{\max}$
#1	SGGGGGGGGC	10	4.35	0.21	0	-	26.65 ± 0.06	104
#2	SGNPTTPPPN	10	2.90	0.49	0	-	27.69 ± 0.05	118
#3	ICDCGHHKDD	10	3.33	0.57	-2	0.05	26.84 ± 0.06	103
#4	GTGSGGGGGGGGGGGC	16	4.24	0.11	0	-	27.26 ± 0.03	109
#5	SSSGGGGSSGGGSGSGS	17	3.91	0.22	0	-	27.58 ± 0.05	109
#6	GAAPNPTTPPPGGGGC	16	3.95	0.42	0	-	27.95 ± 0.06	114
#7	TGPVTSLDQLTSVRVQ	16	4.47	0.64	0	0.81	28.98 ± 0.06	116
#8	GSGGGGGGGGGGGGGGGG TGGC	22	4.20	0.06	0	-	28.91 ± 0.07	119
#9	EGGSSSGGGSSTSSSSGG GGTK	22	3.61	0.31	0	0.18	28.46 ± 0.07	117
#10	GGGTTPPPPPPPPPPPPG GGGC	22	3.55	0.17	0	-	30.06 ± 0.10	134
#11	TSAAPRTTTRTTTRTKSL PTNYNK	24	3.11	0.46	5	0.14	30.19 ± 0.06	128
#12	SSQQTSQQQTSQQQTSQQ QTSSQNNGGGNC	30	2.51	0.38	0	-	30.44 ± 0.07	126
#13	GSSGSTTSSSSSSSSSSS SSSSSSGSTGGQCGNC	34	3.81	0.20	0	-	29.39 ± 0.10	125
#14	GGGGSSSGGGSSSGGGSS SGGGSSSGGGGGTKSF	34	3.93	0.25	1	0.63	30.01 ± 0.07	131
#15	GSPPPPNPPPTNPPPTNP PPTNPPPSNPPPSGS	33	2.86	0.32	0	-	33.45 ± 0.11	136
#16	NATNNNNNNNNNNNNNNN NWNNNNNNNNNNNAGN	33	1.58	0.09	0	-	29.77 ± 0.07	125
#17	YTEANDITVTVDEDAIAQ LSEAEADAGTPEAVG	33	4.26	0.60	-9	0.15	32.19 ± 0.10	143
#18	GGGAASAPAQNNNQGNNF GNQQPQQQQNNNNQWNNW GQQQQQQNNWGQQQQQQN NWGQQQQQNNWGQQQNNW GQPQNAAAAGNGGGAN	88	2.38	0.41	-	0	33.06 ± 0.13	144

Table 1: Features of the selected linkers from the CAZy database. Hydropathy (Hydrop.) indicates the relative hydrophobicity of a sequence; higher values correspond to higher hydrophobicity. Complexity (Complex.) estimates the local compositional complexity of a sequence; lower values indicate simpler, low-complexity regions. N.C. indicates the total charge of the sequence. κ measures charge patterning in proteins; higher values indicate more segregated charges and a tendency toward compact conformations. Rg and $D_{\max}$ values, in Å, were derived from the analyses of the SAXS experiments.

in *Escherichia coli* and purified following previously developed procedures for this family of proteins (see methods section for details) [36, 37]. The purified proteins were submitted to size-exclusion chromatography coupled to small-angle X-ray scattering (SEC-SAXS) measurements. The concentrations used for the injections are shown in Figure S1. In all cases, the SEC profile showed a single peak that was treated using standard procedures to derive a final SAXS profile for subsequent analyses. All the curves displayed an excellent signal to noise and a complete absence of concentration-dependent phenomena, such as oligomerization and interparticle interactions (Fig. 1 and S1 in S.I.). These features underline the quality of this benchmark set to test present protein disorder modelling tools (see below).

The analyses of the resulting curves showed that these proteins presented a broad range of sizes, with Rg values spanning from 26.6 Å to 33.4 Å, with a slow but systematic increase with the length of the linker. Interestingly, DLD15, which has a proline-rich 33-residue long linker, exhibited a larger Rg value than DLD18 that has a much longer linker containing 88 residues. Moreover, another protein

with a proline-rich linker, DLD10, also presented a larger Rg than other proteins with the same length. Similarly to the Rg values, the maximum intramolecular distance, $D_{\max}$, derived from the pairwise distance distribution, $p(r)$, showed a gradual increase with the length of the linker, from 103 Å to 144 Å (see values in Table 1). Although is well known that the $D_{\max}$ values of flexible systems derived from SAXS experiments are not robust structural descriptors [38, 39], some dependence on the composition of the linker was observed. For instance, despite having the same number of residues, DLD10 presented a larger $D_{\max}$ (133 Å) than DLD8 and DLD9 (119 Å and 117 Å, respectively). This is in line with the larger Rg also observed for DLD10. This suggested that conformational features of the individual linkers affect the overall size of DLD proteins in solution.

The inspection of the Kratky plots (Fig. S2 in S.I.) indicated that our benchmark set of DLD proteins is highly flexible. A clear peak was identified for all proteins, but its maximum was displaced from the value expected for a rigid particle [40]. Moreover, the Kratky profile did not reach a value of 0 at large angles. These observations unambiguously indicate that the benchmark displays extensive flexibility. The $p(r)$ functions, with the absence of an isolated second peak corresponding to a dumbbell architecture, and smooth decays towards $D_{\max}$ values, substantiated the presence of high degree of disorder in these proteins (results not shown) [38].

In summary, the model-free analysis of the SAXS profiles demonstrated that SAXS curves measured for this family of proteins represented an excellent dataset to quantitatively validate computational methods adapted to the modelling of flexible multidomain proteins.

2.3 Variable performance of ensemble generation methods in reproducing SAXS data

We applied five computational methods for the generation of conformational ensembles of DLD proteins with the aim to evaluate their accuracy using the previously described SAXS dataset. Concretely, MoMA-FReSa [30, 31], CALVADOS3 [27], Mpipi-Recharged [28], bAIes [32] and BioEmu [41], which are all based on distinct principles, were used in this evaluation. MoMA-FReSa implements a stochastic sampling of disordered regions from local structural information extracted from a database of small protein fragments. CALVADOS3 and Mpipi-Recharged generate ensembles from MD simulations considering one-bead-per-residue coarse-grained (CG) potentials. bAIes also relies on MD simulations, but it uses an all-atom model based on a simplified amber99SB-ILDN force field [42] complemented with a bias potential derived from AlphaFold-predicted pairwise residue distance distributions [43]. Finally, BioEmu is a deep-learning-based approach, trained on a large dataset of MD simulations and experimental data, that generates conformational ensembles directly from protein sequences. For each of the methods, we generated conformational ensembles of similar size (around 10,000 conformations). It should be noted that computational requirements varied considerably depending on the method used. On the hardware employed in this study, MoMA-FReSa required only a few minutes on a multi-core CPU, whereas the coarse-grained MD approaches CALVADOS3 and Mpipi-Recharged typically required on the order of several hours. The all-atom bAIes simulations were substantially more demanding, with runtimes on the order of tens of hours on a single CPU core. BioEmu generated ensembles within a few hours on a modern GPU. Importantly, for MD-based methods, we ran several replicas to verify that the simulation reached convergence with respect to the SAXS data (see Methods section and Figures S3a-c). For the comparative analyses, we used the individual conformational ensemble that provided the best fit to SAXS profile.

Averaged simulated SAXS profiles were computed from each of the ensembles using Crysol [44] and directly compared with the experimental curves using the reduced χ^2-value to quantitatively evaluate the capacity of the different modelling approaches to capture the behaviour of DLD proteins in solution.

Figure 2: SAXS intensity profiles (black) for DLD3 (A), DLD10 (B) and DLD17 (C) compared with the averaged scattering profiles computed from the ensembles modelled with MoMA-FReSa (green), CALVADOS3 (blue), Mpipi-Recharged (cyan), bAIes (orange) and BioEmu (red). χ^2-values are shown in each panel. Point-by-point residuals are displayed below with the same colour code. Rg distributions computed for DLD3 (D), DLD10 (E) and DLD17 (F) using the ensembles computed using the different modelling strategies. Equivalent figures for the complete benchmark are provided in S.I. (see Figures S4, S5 and S6).

The performance of the distinct approaches exhibited an enormous disparity (Table 2). Despite its simplicity, MoMA-FReSa was the computational method that showed the best capacity to reproduce the experimental SAXS profiles. Indeed, MoMA-FReSa was the most accurate method in 14 out of the 18 DLD proteins of the benchmark, while CALVADOS3 was the best for the remaining four cases. For MoMA-FReSa, the χ^2-values ranged from 1.87 to 20.15, while this range varied from 3.59 to 42.56 for CALVADOS3. Conversely, the other three approaches, Mpipi, bAIes and BioEmu, exhibited a very poor capacity to describe the benchmark, with several χ^2-values above 100.

The excellent description of the structural behaviour of DLD proteins provided by the MoMA-FReSa ensembles indicated that the selected linkers present a high level of disorder. Note that this computational approach samples the conformational space in a sequence-dependent manner, but neither the electrostatic nor the hydrophobic long-range interactions are considered when building ensembles. Interestingly, DLD17, the protein for which MoMA-FReSa performed the worst, with $\chi^2 = 20.15$, compared with CALVADOS3, with $\chi^2 = 7.39$, encompassed the linker with the highest overall net charge, -9 (Table 1). We hypothesized that the linker experienced electrostatic interactions with the globular domains that were captured by CALVADOS3, but not by MoMA-FReSa. Similarly, CALVADOS3 performed notably

better than MoMA-FReSa for DLD18, which has the longest linker of the benchmark. Probably, the long linker of this DLD enabled more freedom to establish interactions between both globular domains than the other constructs of the benchmark, and these were better captured by a MD-based approach such as CALVADOS3.

The inspection of the comparison of the synthetic SAXS curves derived from the ensembles with the experimental data provided some clues for the disparity of results obtained for the five modelling approaches (Figure 2A-C, S5 and S6). While bAIes yielded profiles corresponding more extended particles than those present in solution, Mpipi and BioEmu yielded opposite results, generating more compact structures. Interestingly, these features occurred for all the cases of our benchmark, regardless of the composition and length of the linker. Less predictable results were observed for MoMA-FReSa and CALVADOS3, which adopted more or less compact conformations than the experimental one in a DLD-dependent manner.

To structurally evaluate the ensembles derived from the five methods, we computed their Rg distributions (Fig. 2D-F and S6). This analysis indicated strong conformational bias depending on the computational approach. On the one hand, for all DLD proteins, Mpipi and BioEmu exhibited a strong enrichment in highly compact conformations, with a highly populated peak corresponding to conformations with small Rg values. In both cases, however, extended conformations were also populated, although less frequently. On the other hand, bAIes built ensembles sampling a broad range of sizes, with an enrichment in extended conformations, while compact structures were scarcely sampled. MoMA-FReSa and CALVADOS3 built more balanced ensembles sampling a broad range of protein sizes. Interestingly, CALVADOS3 ensembles systematically showed an enrichment in more compact species than MoMA-FReSA, which did not show any special conformational bias, as expected from a random linker generation method. Similar features were observed when monitoring the inter-domain distances computed between the centers of mass (CoM) of the individual domains (Figure S7). Considering the distribution of MoMA-FReSa as the behaviour expected for a scenario without inter-domain interactions, Mpipi and BioEmu overpopulated conformations where the two domains were directly interacting. Interestingly, Mpipi displayed bimodal CoM distributions, suggesting an equilibrium between domain-interacting conformations with others in which the domains were free in solution. Conversely, conformations built with bAIes did not show contacts between domains.

Altogether, these analyses showed that modelling tools displaying a balanced population between compact and extended conformations provided the best accuracy to the DLD SAXS dataset. Conversely, these methodologies presenting systematic structural biases resulted in inaccurate ensembles that were unable to reproduce the behaviour in solution of our protein benchmark set.

2.4 Can SAXS-guided refinement rescue inaccurate (structurally biased) ensembles?

Conformational ensembles are often used as starting pools for data-driven refinement using distinct computational approaches [45]. Here, we have used the conformational ensembles generated by the five computational approaches as starting pools to fit the previously described experimental SAXS curves with the Ensemble Optimization Method (EOM) [46,47], the most popular approach to describe SAXS data of flexible systems. With this analysis, we aimed at exploring the capacity of experimental SAXS data to compensate for large structural disparities observed among the ensembles. The same EOM procedure, searching for the optimal 50-structure sub-ensemble, was applied in all cases (see Methods for details).

The results of this analysis, summarized in Table 2, revealed two distinct trends. High-quality ensembles, *i.e.* with low χ^2-values, were obtained from the pools generated by MoMA-FReSa, CALVADOS3

DLD	FReSa	FReSa +EOM	CALV3	CALV3 +EOM	Mpipi	Mpipi +EOM	bAIes	bAIes +EOM	BioEmu	BioEmu +EOM
#1	**6.01**	2.50	28.90	2.56	>100	54.22	76.07	29.27	76.03	2.90
#2	**8.76**	1.87	42.56	2.15	>100	55.41	85.48	35.63	>100	14.84
#3	**1.87**	1.18	3.59	1.10	27.17	12.95	16.12	7.57	9.74	1.13
#4	**4.61**	1.92	30.27	1.84	>100	53.56	80.19	29.80	96.76	1.65
#5	**8.33**	1.05	23.18	1.09	>100	17.84	>100	72.66	>100	8.24
#6	**2.95**	1.02	16.50	1.06	43.12	16.15	63.57	25.63	68.64	3.24
#7	**2.61**	1.25	16.56	1.40	92.45	31.01	67.44	24.35	75.55	2.38
#8	**4.15**	1.84	27.36	2.37	76.63	32.81	42.39	14.89	75.53	4.82
#9	**3.80**	1.20	15.73	1.24	80.45	36.10	71.56	19.62	>100	3.47
#10	**4.96**	1.08	27.42	1.20	62.69	25.60	42.23	13.55	60.31	1.15
#11	**4.68**	1.69	30.10	2.08	>100	45.72	72.14	21.16	>100	1.52
#12	**5.77**	1.82	8.51	1.40	>100	47.72	91.96	19.66	92.65	1.37
#13	5.59	0.97	**4.85**	0.97	>100	53.56	78.12	23.19	48.76	2.19
#14	**7.01**	2.24	32.02	3.62	>100	58.08	>100	26.60	>100	1.47
#15	**14.62**	1.50	39.38	1.63	63.05	22.70	70.09	13.17	>100	6.26
#16	3.97	1.56	**3.91**	1.09	>100	>100	>100	33.86	93.00	1.49
#17	20.15	1.95	**7.39**	1.93	>100	43.69	>100	18.83	>100	2.68
#18	10.96	1.02	**3.96**	0.91	67.96	33.37	50.66	10.64	5.29	0.96

Table 2: For each of the five ensemble generation methods tested, this table provides the χ^2-values from the comparison of simulated and experimental SAXS profiles. χ^2 values are provided for the ensembles before and after refinement using the EOM method. The best results for each protein before refinement are indicted in bold.

and, in some cases, by BioEmu. Concretely, from MoMA-FReSa and CALVADOS3 initial pools, sub-ensembles describing the SAXS curves with χ^2-values below 2.50 could be derived for all proteins of the benchmark set. Examples of these high quality EOM fits are displayed in Figures 3, S8 and S9. Importantly, when ensembles could be re-weighted to properly describe the data (χ^2-values < 4.0), the capacity of the three building approaches to describe the SAXS curves was similar, regardless to their initial distance to the behaviour of the protein in solution. In contrast, EOM could not find ensembles with the capacity to accurately describe the SAXS data with Mpipi-Recharged or bAIes, for which no χ^2-values below 7.6 could be derived with EOM with initial pools built with these two programs.

In the previous section, we have shown that Mpipi and bAIes built the most structurally biased ensembles of our benchmark, featuring highly compact and extended conformations, respectively. In contrast, MoMA-FReSa and CALVADOS3 provided the most balanced distribution of conformations in terms of Rg and distance between both domains. These observations unambiguously demonstrated that the properties of the original ensemble strongly affect the capacity of ensemble-refining programs to find an appropriate subgroup of conformations describing SAXS data. To have a more detailed picture of the effect of the structural biases of the initial pools to fit SAXS data, we analysed the Rg distribution of the EOM-fitted ensemble for the DLD proteins (Figure 3, SF). Note that only cases in which EOM was able to correctly fit the data ($\chi^2 < 4.0$) have been plotted [15, 20].

The analysis of the Rg distributions indicated that, even when the initial ensemble distributions did not fully agree with the SAXS measurements, a subset of conformations within the pool may still provide a good fit to the experimental profiles. This behavior was particularly evident for CALVADOS3 and BioEmu-generated ensembles across many of the DLD proteins in our benchmark set. For these two methods, the experimental data were able to correct the original structural bias. Conversely, for Mpipi-Recharged and bAIes, the generated ensembles did not adequately sample all the regions of the conformational space explored by the DLD proteins in solution (Figure S6). In both cases, the significantly reduced exploration of a region of the conformational space could not be compensated by the application

Figure 3: SAXS intensity profiles (black) for DLD3 (A), DLD10 (B) and DLD17 (C) compared with the EOM-fitted scattering profiles computed from the ensembles modelled with MoMA-FReSa (green) and BioEmu (red). χ^2-values are shown in each panel. Point-by-point residuals are displayed below with the same colour code. R_g distributions for DLD3 (D), DLD10 (E) and DLD17 (F) using the ensembles derived with EOM using the different modelling strategies. Equivalent figures for the complete benchmark are provided in the SI (see Figures S8-S10).

of SAXS data. These observations highlighted the need of an exhaustive exploration of the structural landscape in order to obtain a satisfactory description of experimental data.

2.5 Do different methods structurally converge after SAXS refinement?

Finally, we studied the similarity between SAXS-refined sub-ensembles derived from distinct original pools. This analysis provideed clues on the capacity of SAXS data in order to robustly define the conformational behaviour of proteins in solution. To gain a more intuitive understanding of the conformational differences between ensembles, we examined their corresponding R_g distributions. Figure 3D-F shows the results for three representative cases, DLD3, DLD10 and DLD17, while the distributions for the complete benchmark set are shown in Figure S10. Interestingly, the R_g distributions after the SAXS refinement were very similar, regardless of the conformational features of the original pool. This was particularly true for MoMA-FReSa and CALVADOS3, which presented astonishingly similar final R_g distributions with virtually equivalent fine structure. Although BioEmu also provided distributions globally similar to those of MoMA-FReSa and CALVADOS3 after EOM-based refinement, they tended to be more spiky and presented discontinuities, especially for very extended conformations. This can be observed for DLD17 (Figure 3F), where three R_g peaks were identified and the last one, at R_g around 47 Å, was detached of

the rest of the distribution. We attributed the differences in the R_g distribution derived from BioEmu ensembles to the larger structural distance between the original pool and that sampled by the protein in solution, as we have shown in the previous section (Table 2), and the lower density of conformations with large R_gs.

The similarity of the R_g distribution shapes and the position of the highly populated R_g values of the three refined ensembles suggested that the relative domain positions of the selected poses were consistent across the building strategies. In order to have a detailed perspective of this hypothesis, we analysed the CoM distribution for the SAXS-refined ensembles (Figure S11). Compared with the original distributions (Figure S7), the overall shapes of the CoM profiles derived from the three pools were very similar, suggesting that SAXS data were effective in selecting similar structures. When analysing these distributions in more detail, we observed that proteins with short linkers ($<$ 22 residues, from DLD1 to DLD7) exhibited larger disparities among the CoM distributions than those with large linkers. From DLD8, the three CoM distributions were very similar, with MoMA-FReSa and CALVADOS3 yielding virtually equivalent profiles for all cases. More disparities were observed for BioEmu CoM distributions. Again, we attributed the differences observed in BioEmu to the strong structural differences of its original ensemble with respect to MoMA-FReSa and CALVADOS3. All together, our analysis of the R_g and CoM distributions suggested that conformational biases of the original conformational pools slightly affect the structural features of the SAXS-refined ensemble.

3 Discussion

The present study underscores the substantial variability that still characterizes current approaches for modeling ensembles of disordered proteins [26]. Across the family of DLD proteins analyzed, performance, quantified through comparison with high-quality SAXS data, was strongly dependent on both the computational method employed and the specific linker sequence. The modeling tools tested, which represent distinct families of ensemble-generation approaches, systematically exhibited method-specific structural biases. In particular, BioEmu and Mpipi-Recharged produced highly compact ensembles, whereas bAIes consistently yielded overly extended conformations. In contrast, ensembles generated using MoMA-FReSA and CALVADOS3 were more balanced and less affected by pronounced structural biases. Among the approaches evaluated, MoMA-FReSA provided the most reliable overall performance, while also offering a very favorable balance between accuracy and computational efficiency. Importantly, the design of our protein dataset enabled us to relate method performance to specific structural features. Both linker composition and linker length emerged as critical determinants of accuracy in a clearly method-dependent manner. For example, CALVADOS3 systematically produced poor fits to SAXS data (large χ^2-values) for proteins encompassing proline-rich (DLD2, DLD10, and DLD15) or glycine-rich (DLD1, DLD8, and DLD14) linkers. This likely reflects limitations of the one-bead-per-residue representation in CALVADOS3, which appears insufficient to capture the distinctive conformational behaviour of these amino acids. In contrast, MoMA-FReSA, which is based on amino-acid-specific conformational sampling derived from high-resolution experimental structures [31], is less sensitive to linker sequence composition. Linker length also significantly influenced method performance. For the six longest linkers analyzed (33 residues or more), CALVADOS3 outperformed MoMA-FReSA in four cases (Table 2). We speculate that longer linkers may facilitate transient interdomain interactions, which are more readily captured by the CALVADOS3 force-field but not by MoMA-FReSA, resulting in relatively compact ensembles. When such intramolecular interactions play a significant role in shaping the conformational landscape, methods that explicitly include electrostatic and hydrophobic interactions are expected to yield more accurate ensemble descriptions. Overall, no single approach consistently reproduced SAXS data across the full set

of proteins examined, reinforcing the notion that ensemble prediction for flexible, multidomain systems remains a fundamentally challenging problem.

The use of SAXS-guided refinement further illustrates both the strengths and limitations of integrating experimental data with computational ensemble generation. In all cases, refinement resulted in a clear improvement in the agreement with experimental scattering profiles, demonstrating that SAXS data can effectively steer conformational ensembles toward more realistic distributions. However, the quality of the refined ensembles was strongly dependent on the generative method used to construct the initial conformational pool. Methods that produced physically plausible and structurally diverse starting ensembles, most notably MoMA-FReSA and CALVADOS3, benefited the most from refinement, yielding highly accurate ensembles with low χ^2-values. Interestingly, in some cases BioEmu ensembles, despite being initially far from the solution behavior of the proteins, were able to achieve good agreement with the SAXS data after refinement. This behavior was not observed for Mpipi-Recharged or bAIes. We attribute the relative success of BioEmu to its broad conformational sampling, which includes compact structures and, to a lesser extent, more expanded conformations (Figures 2D–F and S6). Such diversity appears to provide sufficient flexibility for refinement to effectively reweight the ensemble toward experimentally consistent states. Overall, our results indicate that a broad and representative sampling of the conformational landscape of disordered proteins is a prerequisite for effective SAXS refinement strategies. The conclusions presented here were obtained using EOM, a maximum-parsimony-based approach. Similar dependencies on the quality of the prior ensemble are expected for maximum-entropy-based refinement methods [48, 49], in which all conformations of the initial ensemble are reweighted to achieve agreement with experimental data.

The comparison of SAXS-refined ensembles with low χ^2-values derived from different initial conformational pools provided further insight into both the robustness and the limitations of current modeling and analysis strategies. Notably, ensembles generated using distinct modeling approaches converged upon SAXS refinement. Specifically, SAXS-refined ensembles originating from MoMA-FReSa and CALVADOS3 displayed remarkably similar R_g and CoM distributions, not only preserving the overall range of sampled radii of gyration but also reproducing comparable fine structural features (Figures 3D–F and S10). Strikingly, even BioEmu, whose original conformational pools differed substantially from those of MoMA-FReSa and CALVADOS3, yielded SAXS-refined ensembles with closely matching R_g distributions and similarly populated states. Together, these observations underscore the strong constraining power of SAXS data in defining robust distributions of global structural parameters for highly flexible proteins. However, the intrinsically low-resolution nature of SAXS data necessitates cautious interpretation of refined ensembles. While SAXS can reliably constrain overall molecular dimensions and interdomain CoM distances, it does not provide sufficient information to uniquely determine specific interdomain interactions or detailed local structural features [15, 20].

Beyond methodological considerations, the biological and biotechnological relevance of accurate ensemble modeling deserves particular emphasis. Flexible linkers play a central role in mediating the function of modular proteins by regulating domain spacing, orientation, and dynamics [1, 6]. Reliable ensemble predictions are therefore essential for understanding how these proteins operate in their native biological contexts. Furthermore, prediction accuracy also has direct implications for the engineering of multimodular enzymes for biotechnological applications, where linker design and domain organization can be exploited to optimize catalytic activity, specificity or stability. As demonstrated in this study, their functional and biotechnological importance contrasts sharply with the persistent difficulties in accurately predicting their structural and dynamical properties. In this regard, the integration of experimental data with physics-based modeling approaches emerges as a particularly promising strategy. Crucially, predicted ensembles should be systematically validated against experimental data that probe distinct and

complementary structural features. Such validation is necessary to ensure that the resulting ensembles faithfully capture both the underlying energetic landscape and the conformational preferences encoded in protein sequences.

4 Materials and Methods

Selection of linkers from natural diversity

Sequences of multimodular enzymes containing catalytic GH11 and CBM domains were retrieved from the CAZy database [34]. To ensure consistency, we selected only linkers flanked by a GH11 domain at the N-terminus and a CBM at the C-terminus. Sequences originating from eukaryotes and bacteria were retained, and an additional length constraint was imposed by selecting linkers between 10 and 100 amino acids. After redundancy filtering, this procedure yielded a total of 769 linker sequences.

For each linker, physicochemical properties were computed using localCIDER [35]. The resulting feature set was used to cluster the sequences via hierarchical agglomerative clustering (HAC) with Ward's linkage method [50]. Based on the resulting dendrogram, we selected 18 representative linker sequences that spanned the observed diversity in terms of length and physicochemical composition. This reduced yet diverse subset was chosen to capture the main structural and compositional trends present in natural GH11–CBM linkers, while remaining experimentally and computationally tractable for downstream analyses.

DNA Constructs

The synthetic genes for the glycosyl hydrolase (GH) catalytic module from *Neocallimastix patriciarum*, and for a fusion protein comprising the DLD16 linker and the carbohydrate-binding module (CBM) from *Cellulomonas fimi* were purchased at IDT® (Integrated DNA Technologies). Linker sequences corresponding to DLD1 to DLD17 (excluding DLD16) were ordered as Ultramer® Duplex DNA fragments at IDT®. Due to their length and complexity, the linker corresponding to DLD18 was purchased at GeneArt® (Life Technologies®).The GH gene was first cloned in-frame into the pDB-6His-ccdb plasmid using NcoI and NdeI restriction sites, generating the construct pDB-6His-GH-ccdb. The DLD16–CBM fusion was then inserted between the NdeI and BlpI sites to obtain the pDB-6His-GH-DLD16-CBM plasmid. The insertion and exchange of the different linker sequences (DLD1–18) were performed by an In-Fusion reaction between the NdeI and XhoI restriction sites located between the GH and CBM coding regions.

Protein production and purification

The pDB-6His-GH-DLDXX-CBM vectors were transformed in *E. coli* BL21(DE3) strain and grown in LB medium supplemented with 50 µg/mL of kanamycin until the OD reached 0.6. Protein expression was induced by the incorporation of IPTG (0.5 mM), and cells were grown for 3 h before harvesting them by centrifugation for 20 min at 6000g at 4°C. Pellets were resuspended in Tris-HCl 20 mM pH 7.5, NaCl 300 mM and 2 mM DTT (buffer A) and stored at -80°C. Cells were supplemented with a Complete® EDTA free tablet (Roche) and lysed by sonication. Insoluble proteins and cell debris were sedimented by centrifugation at 40000g at 4°C for 30 min. Supernatant was supplemented with imidazole to a 5 mM final concentration and loaded onto a 5 ml affinity column (HisTrap Excel, Cytiva) equilibrated with

buffer B (buffer A containing 5 mM imidazole). The column was washed with buffer B and proteins were eluted with a linear 0 to 100% gradient of buffer C (buffer A containing 0.5 M imidazole). The peak fractions were analyzed by SDS-PAGE. Fractions containing tagged GH-linker-CBM fusion protein were pooled and dialyzed overnight at 4 °C against buffer D (Na2HPO4/NaHPO4 50 mM pH 6, NaCl 50 mM, DTT 2 mM). The dialyzed protein was then loaded on a Superdex S75 16/60 (HiLoad 16/600 Superdex 75 pg, Cytiva) equilibrated with buffer E (Na2HPO4/NaHPO4 50 mM pH 6, TCEP 2 mM). Fractions containing the protein were analysed by SDS-PAGE and pooled. The purified proteins were concentrated to approximately 10 mg/mL with Vivaspin centrifuge concentrator (Sartorius stedim biotech) before structural analyses.

On fait le SAXS avec le même tampon que la SEC : Na2HPO4/NaHPO4 50 mM pH 6, TCEP 2 mM. On garde effectivement les prots sans sel.

Small-Angle X-ray Scattering measurements

Small-Angle X-ray Scattering (SAXS) experiments were performed on the purified DLD constructs at the SWING beamline of the SOLEIL Synchrotron (Saint-Aubin, France) [51], using an X-ray wavelength of 1.03 Å and a sample-to-detector distance of 2.00 m. Measurements were conducted at 20 °C using Size-Exclusion Chromatography coupled to SAXS (SEC-SAXS) in buffer E. A volume of 45 µL of sample was injected into a 3 mL Superdex 75 5/150 GL column (Cytiva, Uppsala, Sweden) pre-equilibrated with the same buffer and eluted at a flow rate of 0.2 mL/min. The acquired scattering data covered a momentum transfer range of 0.002 Å^{-1} < q < 0.5 Å^{-1}. Buffer scattering was recorded before the void volume of the column (1 mL). Data processing was carried out using CHROMIX [52], which automatically identified buffer and sample frames, subtracted the buffer signal, and averaged the scattering profiles. Final SAXS curves were analyzed using Primus from the ATSAS software suite [53].

Ensemble generation methods

Definition of rigid and flexible regions

Several of the ensemble-generation methods applied in this work require prior specification of rigid and flexible regions. Figure S12 illustrates the generic amino acid composition of the DLD constructs, in which the linker sequence varies across the 18 tested constructs. Based on this architecture, the segments MGSSHHHHHH, TTHM–linker–LESTGC, and SGGGTA were defined as flexible, while the remaining regions were treated as rigid. This partitioning of flexible and rigid residues was consistently applied in the MoMA-FReSA, CALVADOS3, and Mpipi-Recharged simulations.

MoMA-FReSa

MoMA-FReSA [30,31] requires as input an initial structure for the rigid domains together with a definition of which regions of the sequence are treated as rigid or flexible. Conformational ensembles are generated by sequentially sampling backbone dihedral angles within the flexible regions from distributions observed in naturally occurring coil conformations. We generated ensembles of 10,000 conformations for each of the 18 DLD constructs using the three-residue-based sampling strategy (TRS) strategy, where the identity and structure of the two neighboring residues are also taken into account.

CALVADOS3

We performed 1 μs coarse-grained molecular dynamics (CG-MD) simulations for each DLD construct using CALVADOS3 [27] with default parameters. The initial conformations were generated using MoMA-FreSa, with residues belonging to folded domains restrained and the remaining residues treated as flexible (Figure S12). All-atom structures were coarse-grained to a one-bead-per-residue representation using cg2all [54]. By default, CALVADOS3 employs Langevin dynamics, specifically the LangevinMiddleIntegrator implemented in OpenMM, with a time step of 10 fs. To avoid initial equilibration artifacts, the first 100 ns of each trajectory were discarded. Protein configurations were saved every 100 ps, and the resulting CG ensembles were back-mapped to all-atom representations using PULCHRA [55].

Mpipi-Recharged

We performed 1 μs CG-MD simulations using the Mpipi-Recharged force-field [28] for each DLD construction. Simulations were carried out using LAAMPS [56] (2 August 2023 version). Each starting conformation was originally generated by MoMA-FReSa, where the residues that take part of the folded domains were kept fixed and the remaining ones were set as flexible (see Figure S12). The all-atom starting structures were coarse-grained to a one-bead per residue representation using cg2all [54]. Then, the structure files were converted to the LAAMPS input format using in-house scripts. We performed local energy minimization of each starting structure using the steepest descendent algorithm, followed by 1 μs of NVT production using an implicit solvent model, already defined in the Mpipi-Recharged parameters (timestep: 20 fs, temperature: 293K, ionic strength: 150 mM). Snapshots were recorded every 100 ps. The generated ensembles were back-mapped to all-atom using PULCHRA [55].

bAIes

We obtained 100 ns of bAIes MD-simulations for each DLD construct [32]. In this case, the starting structures were obtained using AlphaFold 2 on ColabFold [57] together with the corresponding distogram for each protein, which is a mandatory input file for bAIes. Note that the AlphaFold 2 predicted structures of the GH11 and the CBM domains were extremely similar to the experimental structures ones, with RMSDs of 0.340 and 0.692 Å, respectively. In all cases, an energy minimization of each starting structure was carried out using the conjugate gradient algorithm followed by 100 ns of NVT production in vacuum (timestep of 1 fs and temperature of 293K). Snapshots were recorded each 10 ps. SAXS profiles were computed for the last 80 ns of the simulation.

BioEmu

BioEmu [41] is a parameter-free method. We ran BioEmu to generate ensembles of 10,000 conformations for each construct. Conformations exhibiting steric clashes were automatically filtered by the algorithm; consequently, the final ensemble size slightly varied across DLD systems.

Convergence assessment of MD-based methods

For CALVADOS3 simulations, we performed three independent replicas for each DLD, whereas five replicas were generated for Mpipi-Recharged. Convergence was assessed using the cumulative R_g as the primary criterion (see Figures S3a–c). The larger number of replicas for Mpipi-Recharged was motivated

by the larger variability observed across independent runs compared to CALVADOS3, and was introduced to ensure the reliability of the results. In all cases, simulations reached convergence for every DLD independently of the starting conformation.

For bAIes, we ran 5 independent replicas for the DLD16 and DLD18 constructions and we assessed the their convergence via the cumulative Rg (Figures S3a–c). In each case, one of the starting structures was obtained using AlphaFold 2 on ColabFold [57], while the rest of structures were generated using MoMA-FReSa. In both DLD and for all the starting structures, we used the same distogram retrieved from AlphaFold 2. Given the high degree of convergence of these two examples (Figure S3c, panels K and Q), and due to the higher computational cost compared with the other MD-based approaches, we ran a single trajectory for the rest of the DLD constructs using AlphaFold 2 predictions as starting structures.

Simulation of SAXS profiles

The SAXS profile for each conformation of all ensembles was calculated using Crysol [44], setting the number of spherical harmonics to 35 and using implicit hydrogens. Both before and after SAXS refinement, the χ^2 value between the theoretical and experimental scattering profiles was calculated as follows:

$$\chi^2 = \frac{1}{N-1} \sum_{i=1}^{N} \left(\frac{I_{\text{exp}}(q_i) - (a * I_{\text{calc}}(q_i) + b)}{\sigma(q_i)} \right)^2$$

where N is the number of experimental data points in the SAXS profile; $I_{\text{exp}}(q_i)$ and $I_{\text{calc}}(q_i)$ are the experimental and calculated SAXS intensities at momentum transfer q_i, respectively; and $\sigma(q_i)$ is the experimental error at q_i. Finally, a and b are the scaling factor and constant background offset applied to the theoretical SAXS curve to minimize χ^2.

Ensemble filtering/refinement (EOM)

The ensembles generated for each DLD were refined using the available SAXS data, using the program GAJOE from the EOM suite [46, 47]. In order to fit the calculated SAXS profiles to experimental data, we ran GAJOE using 200 cycles of 1,000 generations each, with a sub-ensemble size of 50 and disallowing repetitions.

CRediT authorship contribution statement

Santiago Rodriguez: Investigation, Validation, Visualization, Writing – original draft, Writing – review & editing; **Aurélie Fournet**: Investigation, Validation, Writing – review & editing; **Simon Bartels**: Investigation, Writing – review & editing; **Mátyás Pajkos**: Investigation, Software; **Ilinka Clerc**: Software; **Laure Carrière**: Investigation, Software; **Aurélien Thureau**: Investigation, Writing – review & editing; **Cédric Montanier**: Resources, Writing – review & editing, Funding acquisition; **Claire Dumond**: Resources, Writing – review & editing; **Fréderic Allemand**: Supervision, Writing – review & editing; **Juan Cortés**: Conceptualization, Supervision, Writing – original draft, Writing – review & editing, Funding acquisition; **Pau Bernadó**: Conceptualization, Supervision, Writing – original draft, Writing – review & editing, Funding acquisition.

Data availability

SAXS data will be deposited in the Small Angle Scattering Biological Data Bank (SASBDB; `https://www.sasbdb.org`) and the SAXS-refined ensembles will be deposited in the Protein Ensemble Database (PED; `https://proteinensemble.org`). Non-refined ensembles generated by the various methods will also be made publicly available upon publication of the final version of the manuscript.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank Nicolas Terrapon for his assistance with extracting data from the CAZy database. This work was supported by the French National Research Agency (ANR) under grant ANR-22-CE45-0003 (CORNFLEX project). It was also carried out in the framework of the COST Action ML4NGP [CA21160], supported by the COST (European Cooperation in Science and Technology) and the HORIZON-MSCA-SE project IDPfun2, funded by the European Union under grant agreement no. 101182949. The CBS is a member of France-BioImaging (FBI) and the French Infrastructure for Integrated Structural Biology (FRISBI), two national infrastructures supported by the French National Research Agency (ANR-10-INBS-04-01 and ANR-10-INBS-05, respectively).

References

[1] Apic, G., Huber, W., and Teichmann, S. A. Multi-domain protein families and domain pairs: comparison with known structures and a random model of domain recombination. *Journal of Structural and Functional Genomics* **4**, 2-3, 67–78 (2023). https://doi.org/10.1023/A:1026113408773.

[2] Levitt, M. (2009). Nature of the protein universe. *Proc. Natl. Acad. Sci. U S A* **106**, 27, 11079–11084.

[3] Zhang, M., Abrams, C., Wang, L., Gizzi, A., He, L., Lin, R., Chen, Y., Loll, P. J., Pascal, J. M., and Zhang, J. Structural basis for calmodulin as a dynamic calcium sensor. *Structure* **20**, 5, 911–923 (2012). https://doi.org/10.1016/j.str.2012.03.019.

[4] Luo, J., Zou, H., Guo, Y., and others. Src kinase-mediated signaling pathways and targeted therapies in breast cancer. *Breast Cancer Research 24*, 99 (2022). https://doi.org/10.1186/s13058-022-01596-y.

[5] Yoh, K., Ikeda, K., Horie, K., and Inoue, S. Roles of estrogen, estrogen receptors, and estrogen-related receptors in skeletal muscle: Regulation of mitochondrial function. *International Journal of Molecular Sciences* **24**, 3, 1853 (2023). https://doi.org/10.3390/ijms24031853.

[6] Lin, C.-Y. and Liu, J. C. Modular protein domains: An engineering approach toward functional biomaterials. *Current Opinion in Biotechnology 40*, 56–63 (2016). https://doi.org/10.1016/j.copbio.2016.02.011.

[7] Sammond, D. W., Payne, C. M., Brunecky, R., Himmel, M. E., Crowley, M. F., and others. Cellulase linkers are optimized based on domain type and function: Insights from sequence anal-

ysis, biophysical measurements, and molecular simulation. *PLOS ONE* **7**, 11, e48615 (2012). https://doi.org/10.1371/journal.pone.0048615.

[8] Badruna, L., Burlat, V., Roblin, P., Enjalbert, T., Lippens, G., Venditto, I., O'Donohue, M. J., and Montanier, C. Y. The jo-in protein welding system is a relevant tool to create cbm-containing plant cell wall degrading enzymes. *New Biotechnology 65*, 31–41 (2021). https://doi.org/10.1016/j.nbt.2021.07.004.

[9] Srivastava, A., Nagar, P., Rathore, S., and Adlakha, N. (2022). The linker region promotes activity and binding efficiency of modular lpmo towards polymeric substrate. *Microbiology Spectrum* **10**, 1, e0269721.

[10] Caffrey, M. A comprehensive review of the lipid cubic phase or in meso method for crystallizing membrane and soluble proteins and complexes. *Acta Crystallographica Section F: Structural Biology Communications* **71**, 1, 3–18 (2015). https://doi.org/10.1107/S2053230X14026843.

[11] Musselman, C. A. and Kutateladze, T. G. Characterization of functional disordered regions within chromatin-associated proteins. *iScience* **24**, 2, 102070 (2021). https://doi.org/10.1016/j.isci.2021.102070.

[12] Bernadó, P., Modig, K., Grela, P., Svergun, D. I., Tchorzewski, M., Pons, M., and Akke, M. Structure and dynamics of ribosomal protein l12: An ensemble model based on SAXS and NMR relaxation. *Biophysical Journal* **98**, 10, 2374–2382 (2010). https://doi.org/10.1016/j.bpj.2010.02.012.

[13] Varadan, R., Walker, O., Pickart, C., and Fushman, D. Structural properties of polyubiquitin chains in solution. *Journal of Molecular Biology* **324**, 4, 637–647 (2022). https://doi.org/10.1016/S0022-2836(02)01198-1.

[14] Bertini, I., Gupta, Y. K., Luchinat, C., Parigi, G., Peana, M., Sgheri, L., and Yuan, J. Paramagnetism-based nmr restraints provide maximum allowed probabilities for the different conformations of partially independent protein domains. *Journal of the American Chemical Society* **129**, 42, 12786–12794 (2007). https://doi.org/10.1021/ja0726613.

[15] Bernadó, P. and Svergun, D. I. Structural analysis of intrinsically disordered proteins by small-angle x-ray scattering. *Mol. BioSyst.* **8**, 151–167 (2012). https://doi.org/10.1039/C1MB05275F.

[16] Receveur-Brechot, V. and Durand, D. How random are intrinsically disordered proteins? a small angle scattering perspective. *Current Protein & Peptide Science* **13**, 1, 55–75 (2012). https://doi.org/10.2174/138920312799277901.

[17] Koch, M. H. J., Vachette, P., and Svergun, D. I. Small-angle scattering: a view on the properties, structures and structural changes of biological macromolecules in solution. *Quarterly Reviews of Biophysics* **36**, 2, 147–227 (2003). https://doi.org/10.1017/S0033583503003871.

[18] Putnam, C. D., Hammel, M., Hura, G. L., and Tainer, J. A. X-ray solution scattering (SAXS) combined with crystallography and computation: defining accurate macromolecular structures, conformations and assemblies in solution. *Quarterly Reviews of Biophysics* **40**, 3, 191–285 (2007). https://doi.org/10.1017/S0033583507004635.

[19] Bernadó, P., Shimizu, N., Zaccai, G., Kamikubo, H., and Sugiyama, M. Solution scattering approaches to dynamical ordering in biomolecular systems. *Biochimica et Biophysica Acta – General Subjects* **1862**, 2, 253–274 (2018). https://doi.org/10.1016/j.bbagen.2017.10.015.

[20] Cordeiro, T. N., Herranz-Trillo, F., Urbanek, A., Estaña, A., Cortés, J., Sibille, N., and Bernadó, P. Small-angle scattering studies of intrinsically disordered proteins and their complexes. *Current Opinion in Structural Biology 42*, 15–23 (2017). https://doi.org/10.1016/j.sbi.2016.10.011.

[21] Hub, J. S. Interpreting solution x-ray scattering data using molecular simulations. *Current Opinion in Structural Biology 49*, 18–26 (2018). https:/doi.org/10.1016/j.sbi.2017.11.002.

[22] Huang, J., Rauscher, S., Nawrocki, G., Ran, T., Feig, M., de Groot, B. L., Grubmüller, H., and MacKerell, A. D. J. Charmm36m: an improved force field for folded and intrinsically disordered proteins. *Nat. Methods* **14**, 1, 71–73 (2017). https://doi.org/10.1038/nmeth.4067.

[23] Robustelli, P., Piana, S., and Shaw, D. E. Developing a molecular dynamics force field for both folded and disordered protein states. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 21, E4758–E4766 (2018). https://doi.org/10.1073/pnas.1800690115.

[24] Laio, A. and Parrinello, M. Escaping free-energy minima. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 20, 12562–12566 (2002). https://doi.org/10.1073/pnas.202427399.

[25] Best, R. B. Computational and theoretical advances in studies of intrinsically disordered proteins. *Curr. Opin. Struct. Biol. 42*, 147–154 (2017). https:/doi.org/10.1016/j.sbi.2017.01.006.

[26] Clerc, I., Sagar, A., Barducci, A., Sibille, N., Bernadó, P., and Cortés, J. The diversity of molecular interactions involving intrinsically disordered proteins: A molecular modeling perspective. *Comput. Struct. Biotechnol. J. 19*, 3817–3828 (2021). https://doi.org/10.1016/j.csbj.2021.06.031.

[27] Cao, F., von Bülow, S., Tesei, G., and Lindorff-Larsen, K. A coarse-grained model for disordered and multi-domain proteins. *Protein Sci* **33**, 11, e5172 (2024). https://doi.org/10.1002/pro.5172.

[28] R. Tejedor, A., Aguirre Gonzalez, A., Maristany, M. J., Chew, P. Y., Russell, K., Ramirez, J., Espinosa, J. R., and Collepardo-Guevara, R. Chemically informed coarse-graining of electrostatic forces in charge-rich biomolecular condensates. *ACS Cent Sci* **11**, 2, 302–321 (2025). https://doi.org/10.1021/acscentsci.4c01617.

[29] Bernadó, P., Blanchard, L., Timmins, P., Marion, D., Ruigrok, R. W. H., and Blackledge, M. A structural model for unfolded proteins from residual dipolar couplings and small-angle x-ray scattering. *Proc. Natl. Acad. Sci. U.S.A.* **102**, 47, 17002–17007 (2005). https://doi.org/10.1073/pnas.0506202102.

[30] Pajkos, M., Clerc, I., Zanon, C., Bernadó, P., and Cortés, J. AFflecto: A web server to generate conformational ensembles of flexible proteins from alphafold models. *J Mol Biol* **437**, 15, 169003 (2025). https://doi.org/10.1016/j.jmb.2025.169003.

[31] Estaña, A., Sibille, N., Delaforge, E., Vaisset, M., Cortés, J., and Bernadó, P. Realistic ensemble models of intrinsically disordered proteins using a structure-encoding coil database. *Structure* **27**, 2, 381–391.e2 (2019). https://doi.org/10.1016/j.str.2018.10.016.

[32] Schnapka, V., Morozova, T. I., Sen, S., and Bonomi, M. Atomic resolution ensembles of intrinsically disordered proteins with AlphaFold. *Nature Communications*, (2026). https://doi.org/10.1038/s41467-026-69172-y.

[33] McBride, A. C., Yu, F., Cheng, E. H., Mpouli, A., Soe, A. C., Hammel, M., Montelione, G. T., Oas, T. G., Tsutakawa, S. E., and Donald, B. R. Predicting pose distribution of protein domains connected by flexible linkers is an unsolved problem. *Proteins: Structure, Function, and Bioinformatics* **94**, 1, 362–380 (2026). https://doi.org/10.1002/prot.70062.

[34] Drula, E., Garron, M.-L., Dogan, S., Lombard, V., Henrissat, B., and Terrapon, N. The carbohydrate-active enzyme database: functions and literature. *Nucleic Acids Res* **50**, D1, D571–D577 (2021). https://doi.org/10.1093/nar/gkab1045.

[35] Holehouse, A. S., Das, R. K., Ahad, J. N., Richardson, M. O. G., and Pappu, R. V. CIDER: Resources to analyze sequence-ensemble relationships of intrinsically disordered proteins. *Biophysical Journal* **112**, 1, 16–21 (2017). https://doi.org/10.1016/j.bpj.2016.11.3200.

[36] Montanier, C. Y., Fanuel, M., Rogniaux, H., Ropartz, D., Di Guilmi, A.-M., and Bouchoux, A. (2019). Changing surface grafting density has an effect on the activity of immobilized xylanase towards natural polysaccharides. *Scientific Reports* **9**, 1, 5763.

[37] Enjalbert, T., De La Mare, M., Roblin, P., Badruna, L., Vernet, T., Dumon, C., and Montanier, C. Y. (2020). Characterisation of the effect of the spatial organisation of hemicellulases on the hydrolysis of plant biomass polymer. *International Journal of Molecular Sciences* **21**, 12, 4360.

[38] Bernadó, P. Effect of interdomain dynamics on the structure determination of modular proteins by small-angle scattering. *European Biophysics Journal* **39**, 5, 769–780 (2010). https://doi.org/10.1007/s00249-009-0549-3.

[39] Heller, W. T. Influence of multiple well defined conformations on small-angle scattering of proteins in solution. *Acta Crystallographica Section D* **61**, 1 (Jan), 33–44 (2005). https://doi.org/10.1107/S0907444904025855.

[40] Durand, D., Vivès, C., Cannella, D., Pérez, J., Pebay-Peyroula, E., Vachette, P., and Fieschi, F. Nadph oxidase activator p67phox behaves in solution as a multidomain protein with semi-flexible linkers. *Journal of Structural Biology* **169**, 1, 45–53 (2010). https://doi.org/10.1016/j.jsb.2009.08.009.

[41] Lewis, S., Hempel, T., Jiménez-Luna, J., Gastegger, M., Xie, Y., Foong, A. Y., Satorras, V. G., Abdin, O., Veeling, B. S., Zaporozhets, I., Chen, Y., Yang, S., Foster, A. E., Schneuing, A., Nigam, J., Barbero, F., Vincent, S., Campbell, A., Yim, J., Lienen, M., Shi, Y., Zheng, S., Schulz, H., Munir, U., Sordillo, R., Tomioka, R., Clementi, C., and Noé, F. Scalable emulation of protein equilibrium ensembles with generative deep learning. *Science*, eadv9817 (2025). https://doi.org/10.1126/science.adv9817.

[42] Lindorff-Larsen, K., Piana, S., Palmo, K., Maragakis, P., Klepeis, J. L., Dror, R. O., and Shaw, D. E. Improved side-chain torsion potentials for the Amber ff99SB protein force field. *Proteins* **78**, 8, 1950–1958 (2010). https://doi.org/10.1002/prot.22711.

[43] Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., and others. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 7873, 583–589 (2021). https://doi.org/10.1038/s41586-021-03819-2.

[44] Franke, D., Petoukhov, M. V., Konarev, P. V., Panjkovich, A., Tuukkanen, A., Mertens, H. D. T., Kikhney, A. G., Hajizadeh, N. R., Franklin, J. M., Jeffries, C. M., and Svergun, D. I. *ATSAS 2.8*: a comprehensive data analysis suite for small-angle scattering from macromolecular solutions. *Journal of Applied Crystallography* **50**, 4, 1212–1225 (2017). https://doi.org/10.1107/S1600576717007786.

[45] Bonomi, M., Heller, G. T., Camilloni, C., and Vendruscolo, M. Principles of protein structural ensemble determination. *Current Opinion in Structural Biology 42*, 106–116 (2017). https://doi.org/10.1016/j.sbi.2016.12.004.

[46] Bernadó, P., Mylonas, E., Petoukhov, M. V., Blackledge, M., and Svergun, D. I. Structural characterization of flexible proteins using small-angle X-ray scattering. *J Am Chem Soc* **129**, 17, 5656–5664 (2007). https://doi.org/10.1021/ja069124n.

[47] Tria, G., Mertens, H. D. T., Kachala, M., and Svergun, D. I. Advanced ensemble modelling of flexible macromolecules using X-ray solution scattering. *IUCrJ* 2, 207–217 (2015). https://doi.org/10.1107/S205225251500202X.

[48] Bottaro, S., Bengtsen, T., and Lindorff-Larsen, K. Integrating molecular simulation and experimental data: A bayesian/maximum entropy reweighting approach. *Methods Mol. Biol. 2112*, 219–240 (2020). https://doi.org/10.1007/978-1-0716-0270-6₁5.

[49] Pesce, F. and Lindorff-Larsen, K. Refining conformational ensembles of flexible proteins against small-angle x-ray scattering data. *Biophys. J.* **120**, 22, 5124–5135 (2021). https://doi.org/10.1016/j.bpj.2021.10.003.

[50] Murtagh, F. and Legendre, P. Ward's hierarchical agglomerative clustering method: Which algorithms implement ward's criterion? *Journal of Classification* **31**, 3, 274–295 (2014). https://doi.org/10.1007/s00357-014-9161-z.

[51] Thureau, A., Roblin, P., and Perez, J. Biosaxs on the swing beamline at synchrotron soleil. *Journal of Applied Crystallography* **54**, 6, 1698–1710 (2021). https://doi.org/10.1107/S1600576721008736.

[52] Panjkovich, A. and Svergun, D. I. Chromixs: automatic and interactive analysis of chromatography-coupled small-angle x-ray scattering data. *Bioinformatics* **34**, 11, 1944–1946 (2018). https://doi.org/10.1093/bioinformatics/btx846.

[53] Manalastas-Cantos, K., Konarev, P. V., Hajizadeh, N. R., Kikhney, A. G., Petoukhov, M. V., Molodenskiy, D. S., Panjkovich, A., Mertens, H. D. T., Gruzinov, A., Borges, C., Jeffries, C. M., Svergun, D. I., and Franke, D. Atsas 3.0: expanded functionality and new tools for small-angle scattering data analysis. *Journal of Applied Crystallography* **54**, 1, 343–355 (2021). https://doi.org/10.1107/S1600576720013412.

[54] Heo, L. and Feig, M. One bead per residue can describe all-atom protein structures. *Structure* **32**, 1, 97–111.e6 (2024). https://doi.org/10.1016/j.str.2023.10.013.

[55] Rotkiewicz, P. and Skolnick, J. Fast procedure for reconstruction of full-atom protein models from reduced representations. *Journal of Computational Chemistry* **29**, 9, 1460–1465 (2008). https://doi.org/10.1002/jcc.20906.

[56] Thompson, A. P., Aktulga, H. M., Berger, R., Bolintineanu, D. S., Brown, W. M., Crozier, P. S., in't Veld, P. J., Kohlmeyer, A., Moore, S. G., Nguyen, T. D., Shan, R., Stevens, M. J., Tranchida, J., Trott, C., and Plimpton, S. J. LAMMPS - a flexible simulation tool for particle-based materials modeling at the atomic, meso, and continuum scales. *Computer Physics Communications 271*, 108171 (2022). https://doi.org/10.1016/j.cpc.2021.108171.

[57] Mirdita, M., Schuetze, K., Moriwaki, Y., Heo, L., Ovchinnikov, S., and Steinegger, M. Colabfold: making protein folding accessible to all. *Nature Methods* **19**, 6, 679–682 (2022). https://doi.org/10.1038/s41592-022-01488-1.

Supplementary Information

Conformational ensembles of flexible multidomain proteins: How close are we to accurate and reliable predictions?

Santiago Rodriguez[1], Aurélie Fournet[1], Simon Bartels[2], Mátyás Pajkos[2], Ilinka Clerc[2], Laure Carrière[2], Aurélien Thureau[3], Cédric Montanier[4], Claire Dumon[4], Frédéric Allemand[1], Juan Cortés[2,*] and Pau Bernadó[1,*]

[1] *Centre de Biologie Structurale, Université de Montpellier, INSERM, CNRS, Montpellier, France.*

[2] *LAAS-CNRS, Université de Toulouse, CNRS, Toulouse, France.*

[3] *Swing Beamline, Synchrotron SOLEIL, 91192 Gif sur Yvette, France.*

[4] *TBI, Université de Toulouse, CNRS, INRAE, INSA, Toulouse, France.*

[*] *Corresponding author: Juan Cortés (juan.cortes@laas.fr) and Pau Bernadó (pau.bernado@cbs.cnrs.fr)*

Figure S1. SAXS intensity profiles *versus* the momentum transfer measured for the 18 DLD proteins. The specific DLD number is shown at the bottom left of each panel.

Figure S2. Dimensionless Kratky representation for the 18 DLD proteins measured in this study. The vertical ($\sqrt{3}$) and horizontal (1.104) dashed lines indicate the maximum expected in the Kratky plot for a globular protein. The specific DLD number is shown in each panel.

Figure S3a. Cumulative Rg across simulations for the MD-based ensemble generation methods (displayed in the columns). Different colors in each panel represent a different starting conformation in the MD simulation. Note that the scales of the panels change according to the values obtained. The specific DLD number is shown in each row.

Figure S3b. Cumulative Rg across simulations for the MD-based ensemble generation methods (displayed in the columns). Different colors in each panel represent a different starting conformation in the MD simulation. Note that the scales of the panels change according to the values obtained. The specific DLD number is shown in each row.

Figure S3c. Cumulative Rg across simulations for the MD-based ensemble generation methods (displayed in the columns). Different colors in each panel represent a different starting conformation in the MD simulation. Note that the scales of the panels change according to the values obtained. The specific DLD number is shown in each row.

Figure S4. SAXS intensity profiles (black) for all DLDs constructs compared with the averaged scattering profiles computed from the ensembles modelled with MoMA-FReSa (green), CALVADOS3 (blue), Mpipi-Recharged (cyan), bAIes (orange) and BioEmu (red). χ^2-values are shown in each panel. The specific DLD number is shown in each panel.

Figure S5. Point-by-point residuals between the SAXS intensity profiles and the averaged theoretical scattering profiles from Figure S4. The specific DLD number is shown in each panel.

Figure S6. Rg distributions computed from the ensembles generated using the methods tested in this study for the 18 DLD proteins: MoMA-FReSa (green), CALVADOS3 (blue), Mpipi Recharged (cyan), bAIes (orange) and BioEmu (red). Note that the scales of the panels change according to the values obtained. The specific DLD number is shown in each panel.

Figure S7. CoM distance distributions computed from the ensembles generated using the methods tested in this study for the 18 DLD proteins: MoMA-FReSa (green), CALVADOS3 (blue), Mpipi Recharged (cyan), bAIes (orange) and BioEmu (red). Note that the scales of the panels change according to the values obtained. The specific DLD number is shown in each panel.

Figure S8. Theoretical scattering profiles fitted against the experimental SAXS curves (in black). MoMA-FReSa is displayed in green, CALVADOS3 in blue, and BioEmu in red. χ^2-values and the specific DLD number is shown in each panel.

Figure S9. Point-by-point residuals between the SAXS intensity profiles and the fitted theoretical scattering profiles from Figure S8. The specific DLD number is shown in each panel.

Figure S10. Rg distributions derived from the EOM analyses of the SAXS profile for each DLD protein using the ensembles computed using the different modelling strategies. Note that only distributions obtained with EOM fits with χ^2 values below 4 are shown. This corresponds to MoMA-FReSa (green) and CALVADOS3 (blue) in all cases, and BioEMu (red) for 14 cases. The specific DLD number is shown in each panel.

Figure S11. CoM distance distributions computed from the EOM-fitted ensembles for the 18 DLD constructs: MoMA-FReSa (green), CALVADOS3 (blue), and BioEmu (red). Note that the scales of the panels change according to the values obtained. The specific DLD number is shown in each panel.

MGSSHHHHHHKFTVGNGQNQHKGVNDGFSYEIWLDNTGGNGSMTLGSGATFKAEWNA
AVNRGNFLARRGLDFGSQKKATDYDYIGLDYAATYKQTASASGNSRLCVYGWFQNRGLN
GVPLVEYYIIEDWVDWVPDAQGKMVTIDGAQYKIFQMDHTGPTINGGSETFKQYFSVRQQ
KRTSGHITVSDHFKEWAKQGWGIGNLYEVALNAEGWQSSGVADVTLLDVY**TT**HM**-linker-
LE**STGC**SVTATRAEEWSDRFNVTYSVSGSSAWTVNLALNGSQTIQASWNANVTGSGSTRT
VTPNGSGNTFGVTVMKNGSSTTPAATCAG**SGGGTA**

His-Tag
GH (PDB: 2C1F)
Cloning site
CBM (PDB:2XBD)

Figure S12. Generic amino acid sequence for the 18 DLD constructs tested in this work. Residues in bold correspond to flexible amino acids, otherwise rigid.