Nanoliposomal Omega-3 Fatty Acids Promote Adult Hippocampal Neurogenesis through the

BDNF/TrkB Pathway in C57BL/6 Mice

Foltran RB*[1], Diaz G*[2,3,4], Stefani KM[5,6], Feliu MS[7], Impa Condori AR[7], Colapietro AA[5,6], Montagna

DR[8,9], Ambrosi V[2,3,4,10], Godoy MF[7], Guidi SM[2,3,10], Nanni M[11], Diaz SL[8,12]

*These authors contributed equally to this article

[1] Instituto de Fisiología, Biología Molecular y Neurociencias (IFIBYNE), UBA-CONICET, Buenos

Aires, Argentina.

[2] Instituto Tecnología de Alimentos, INTA, Buenos Aires, Argentina.

[3] Instituto de Ciencia y Tecnología de Sistemas Alimentarios Sustentables (ICyTeSAS) INTA –

CONICET, Buenos Aires, Argentina.

[4] Facultad de Farmacia y Bioquímica (FFyB), Cátedra de Bromatología, UBA. Buenos Aires,

Argentina.

[5] Instituto de Biología Celular y Neurociencias (IBCN), UBA-CONICET, Buenos Aires, Argentina.

[6] Universidad Argentina de la Empresa (UADE). Buenos Aires, Argentina.

[7] Facultad de Farmacia y Bioquímica (FFyB), Cátedra de Nutrición, UBA. Buenos Aires, Argentina.

[8] Instituto de Biociencias, Biotecnología y Biología traslacional (IB3), UBA-CONICET. Buenos

Aires, Argentina.

[9] Facultad de Ciencias Veterinarias (FVET), Cátedra de Química Biológica, UBA. Buenos Aires,

Argetina.

[10] Universidad de Moron, Escuela Superior de Ingeniería, Informática y Ciencias Agroalimentarias

(ESIICA, UM), Argentina.

[11] Universidad Nacional de Hurlingham (UNAHUR), Argentina.

[12] Facultad de Farmacia y Bioquímica (FFyB), Cátedra de Técnicas de Bioterio, UBA. Buenos Aires,

Argentina.

Abstract

Polyunsaturated fatty acids (PUFAs) are fundamental for different cellular and structural

processes, especially regarding the nervous system. However, its incorporation in food has many

bioaccessibility limitations, making it important to find new ways of its consumption. In this

work, nanoencapsulated PUFAs orally administrated to C57BL/6 elite male mice for 8 weeks

showed better bioavailability when compared to administration of free acids, also improving the

effects on dentate gyrus neuronal survival as well as on the proneurogenic elements of the Brain

derived neurotrophic factor biological pathway also in the hippocampus. In addition,

nanoencapsulated PUFAs increased expression of Fabp5, a relevant n-3 fatty acids transporter

in the brain. Altogether, our results would mean that the form of administration of the fatty

acids can alter not only how much and how preserved they reach the central nervous system,

but also have a differential impact in the diverse processes they contribute to.

Introduction

Polyunsaturated fatty acids (PUFAs) are indispensable structural and signaling components of

cellular architecture. Scientific interest in neurobiology focuses on omega-3 Polyunsaturated

fatty acids (n-3 PUFAs), specifically, eicosapentaenoic acid (20:5, n-3; EPA) and docosahexaenoic

acid (22:6, n-3; DHA). These molecules are far more than energy substrates: they are

preferentially esterified into neuronal membrane phospholipids. Within this lipid bilayer, their

high degree of unsaturation modulates critical physicochemical properties, i.e. membrane

fluidity, curvature, and the spatial organization of lipid rafts, essential for neurotransmission and

synaptic plasticity (1,2).

The physiological efficacy of n-3 PUFAs is dictated by their transition from the diet to the brain

parenchyma, a process restricted by the blood-brain barrier (BBB). Recent evidence identifies

the Major Facilitator Superfamily Domain-containing protein 2a (MFSD2A) as the primary

transporter for lysophosphatidylcholine-bound DHA, while the Fatty Acid-Binding Protein 5

(*Fabp5*) facilitates the uptake of non-esterified forms (3,4). Collectively, this coordinated

transport mechanism represents the physiological bottleneck that determines the

neuroprotective efficacy of lipids.

Furthermore, the modern "Western Diet" has shifted the systemic lipid profile toward a pro-

inflammatory state, characterized by an excessive n-6/n-3 ratio and high Saturated Fatty-Acids

(SFA) intake, which respectively competes with n-3 PUFAs for desaturation and elongation

pathways, and promote membrane lipid remodeling that limits n-3 PUFA incorporation (5).

Conventional n-3 incorporation in food, however, is hindered by the extreme susceptibility of

methylene-interrupted double bonds to oxidative stress. The resulting lipid peroxidation

products not only diminish biological functionality but can also induce cellular toxicity.

To overcome these bioaccessibility limitations, the use of fermented dairy matrices offers a

promising strategy as a carrier for omega-3 fatty acids, owing to their ability to optimize

bioactive stability and functionality. Furthermore, nanotechnology-specifically through

nanoemulsions and encapsulation-has been established as a superior strategy to enhance lipid

delivery (6). By protecting PUFAs from gastric degradation and optimizing intestinal

micellization, these systems significantly increase the concentration of active fatty acids in target

tissues (7).

A central objective of this research is to elucidate how this enhanced omega-3 fatty acids

delivery impacts adult hippocampal neurogenesis in a murine model. The dentate gyrus (DG)

maintains a neurogenic niche where multipotent stem cells expressing Nestin (Nes) transition

into mature, Calbindin-positive (Calb1) functional neurons (8). This continuum is orchestrated

by the Brain-Derived Neurotrophic Factor (BDNF) pathway, among other factors. The BDNF is

synthesized as a pro-protein, the 32 kDa pro-BDNF that may be cleaved to yield the 13.2-15.9

kDa mature BDNF (mBDNF). The pro-BDNF can release mBDNF after cleavage either intra- or

extra-cellularly, or be secreted as pro-BDNF without further modifications. By binding to TrkB

receptors, the mBDNF facilitates pro-neurogenic effects, whereas the pro-BDNF may bind

different receptors, being the most known the p75, through which pro-apoptotic actions are

promoted (see a rev. in (9)). While DHA is known to upregulate BDNF (10,11), the specific

molecular mechanisms by which nanoencapsulated n-3 PUFAs (DHA-EPA) modulate the

expression of markers of neuronal differentiation during this process remain under-investigated.

Therefore, the present study evaluates the effects of orally administered nanoencapsulated n-3

PUFAs (DHA-EPA), delivered via a fermented yogurt matrix, in male C57BL/6 mice over a chronic

period. The administration of the bioactive compounds was based on the Recommended Daily

Intake (RDI) of at least 250mg/day of EPA+DHA for healthy adults (excluding non-pregnant/non-

lactating adult females) established by international organizations (12), integrated into a

fermented dairy matrix to ensure its bioavailability and mimic an optimized human consumption

scenario. We hypothesize that this formulation increases systemic bioavailability, BBB transport

efficiency, and metabolic incorporation compared to free fatty acids, thereby enhancing

neuronal survival in the DG and robustly activating the BDNF signaling cascade.

Materials and Methods

Animals

Studies were performed on 51 male, C57BL/6 elite mice purchased at the Instituto de Medicina Experimental, Academia Nacional de Medicina, Buenos Aires, Argentina. Experiments on animals were conducted according to local regulations and were approved by the Institutional Ethical Committee (ResCD-2022-1246-E-UBA-DCT#FMED). Three-four-week-old mice, bred in barrier-conditions to maintain an SPF status, were transported in environmentally controlled conditions to our institute's animal facility. After arrival, mice were housed in 1284L Eurostandard Type II Long (365 mm × 207 mm × 140 mm) Tecniplast microisolator cages with filter tops (five to six animals per cage), with autoclavated aspen shavings as bedding and tissue paper as nesting material. Mice were maintained under controlled conditions, i.e., 22 ± 2°C room temperature, 40-70% relative humidity, 12–12 h light–dark cycle (lights on at 8 a.m.), pelleted food for rodents (Cooperación) and water ad libitum. Cages were changed twice a week. A period of acclimation of 2 weeks was left before the beginning of experiments, and therefore, mice were 5–6 week-old when treatments began.

Yogurt preparation and administration

Raw Materials and Starter Culture

Yogurt was manufactured using commercial pasteurized cow milk (2% w/v fat; *Mastellone Hnos. S.A., Buenos Aires, Argentina*). A lyophilized Direct Vat Set (DVS) thermophilic starter culture, YoFlex® YG-X16 (comprising *Streptococcus thermophilus* and *Lactobacillus delbrueckii* subsp. *bulgaricus*), was provided by Chr. Hansen (Hørsholm, Denmark). Prior to inoculation, the culture was activated by hydrating 3 g of the lyophilized powder in 1000 mL of milk at 45 °C according to the manufacturer's instructions.

Functional Ingredients and Yogurt Preparation

Omega-3 oil (MEG-3™ 2050EE) was sourced from DSM Nutritional Products. Three distinct

formulations were established based on the omega-3 delivery system:

• Control: Yogurt without oil supplementation.

• Free Acids: Yogurt supplemented with bulk omega-3 oil (250 mg / 200 mL milk).

• Nano Acids: Yogurt supplemented with a nanoencapsulated omega-3/soy lecithin

formulation. To achieve an equivalent dose of 250 mg of active omega-3 per 200 mL (provided

by Nanotica Agro SRL via an R&D agreement with INTA) was incorporated.

In both functional formulations, bioactive compounds were homogenized into the milk-sugar

base (75 g/L) prior to fermentation.

Yogurt Manufacturing

The mixtures were inoculated with the activated starter culture (2 mL/L) and aseptically

distributed into 200 g sterile polypropylene containers. Incubation was performed at 43 ± 1 °C

until a target pH of 4.5 was achieved. Samples were then cooled and stored at 4 °C for 28 days

for analysis.

Yogurt Administration

The different formulations of yogurt were administered to mice by mixing it with a palatable

cube. They were made with 10 g of powdered yeast, 66 g of powdered milk, 36 g of sugar, and

14 g of gelatin, for each ice tray (12 cubes) (13). These ingredients were mixed with 2 ml of each

of the three yogurts and with approximately 60 mL of tap water, forming a thick liquid

preparation, which was poured into an ice tray. The 3 cm × 3 cm × 4 cm cubes, weighing around

12 g, were kept at 4 °C and were prepared every 6 days. A yogurt containing cube was given per

cage, every day between 2 p.m. and 5 p.m. Mice voluntarily ate the cubes, immediately after

they were placed in the cages.

Experiments:

All mice were divided into 3 groups (control vs the two types of yogurt formulation) for 3 different experiments. For the first protocol, 5 mice per group were administered the yogurt cubes for 4 weeks. At the end of the treatment, all mice were deeply anesthetized (20 mg/kg xylazine and 200 mg/kg ketamine) and blood was obtained by intracardiac collection, from which serum was extracted to quantify fatty acids concentrations. After death confirmation, mice were decapitated, brains were removed, and both hippocampus (HC) were obtained, where the left hemi-HCs were prepared for protein extraction and Western blots and the right hemi-HCs were used for mRNA measurements by qRT-PCR.

For the second experiment, in order to study neuronal proliferation 18 mice (n = 6 per group) were administered yogurt cubes for 4 weeks. The thymidine analog 5-bromo-2'-deoxyuridine (BrdU; Sigma, B9285) was administered to mice at the end of the treatment, in order to label dividing cells. Animals received two ip injections of BrdU, 50 mg/kg each, at a 2 h interval. BrdU was dissolved in 0.9% NaCl at 50 °C. The next day, all mice were deeply anesthetized (20 mg/kg xylazine and 200 mg/kg ketamine) and transcardially perfused with 5 mL of 0.9% NaCl with heparin and 50 mL of 4% paraformaldehyde in 0.1 M phosphate-buffered saline (PBS, pH 7.4) for 15 min. Brains were recovered, post-fixed for 24 h at 4 °C, and cryoprotected with a 30% sucrose solution. Brains were then sliced into 35 μm thick coronal sections through the hippocampus (bregma −0.94 to −3.80) with a freezing microtome. Sections were stored at 4 °C in 0.1 M PBS with 0.1% sodium azide if not used immediately.

For the last trial, the yogurt cubes were administered for 8 weeks to 18 mice (6 per group) to study the survival of newborn neurons. Animals received two ip injections of BrdU, 50 mg/kg each, at a 2 h interval for two consecutive days at the end of the 4th week of treatment. Mice were then continued treatment for 4 more weeks, so that BrdU labeled 4 week-old neurons. At the end of the treatment, all mice were deeply anesthetized and transcardially perfused, as stated before.

Fatty acid profile determination

Serum fatty acid profile was determined by gas chromatography: Chromatograph: Perkin Elmer

Clarus 500; Column: Supelco SP 2560 100 m x 0.25 mm x 0.20 μm; FID detector 280 °C; Carrier

gas: helium. Previously, serum lipids were converted into methyl esters according to the Lepage

and Roy method as follows: 2 mL of methanol/toluene (4:1) were added to 200 μL of serum,

followed by 0.2 mL of acetyl chloride using a vortex mixer. The mixture was incubated at 100 °C

for one hour. Then, 5 mL of 6% potassium carbonate was added, and the samples were

centrifuged; 1 μL of the toluene layer was injected into the chromatograph. Fatty acid methyl

esters (FAMEs) were identified based on their retention times by comparison with the PUFA No.

2 (Animal Source) FAME analytical standard (SUPELCO 47015-U). The results (mean ±SD) were

expressed as the per cent (%) of total fatty acids content, with a quantification limit of 0.05%

(14).

Western Blotting

Tissue was homogenized with 250 μl of RIPA buffer (150 mM NaCl, 1% NP-40, 0.5% Sodium

deoxycholate, 0.1% SDS, 50 mM Tris) along with protease inhibitors (1:100) and centrifuged at

4°C for 30 min at 13000 (r/min). The supernatants were recovered and protein levels were

quantified by the Bradford protein assay. Samples (50 μgr in 5× loading buffer) were then loaded

into SDS-PAGE gels (12 or 15%) and transferred onto nitrocellulose membranes using the Mini-

PROTEAN® Tetra System (BIO-RAD) for 1 h.

Membranes were incubated for 1 h with blocking solution (5% milk in TBST) and then probed

overnight at 4°C with mouse anti-BDNF (1:2000; Icosagen; 327-100 clone 3C11), rabbit anti-p75

(1:700; Alomone Labs; ANT-007), rabbit anti-TrkB (1:700; Alomone Labs; ANT-019), and rabbit

anti-proBDNF (1:2500; Genocopoeia) in TBST. Actin was used as a loading control (1:2000; Santa

Cruz). Binding of primary antibodies was visualized with anti-mouse HRP-conjugated secondary

antibody (1:20000; Invitrogen) or anti-rabbit HRP-conjugated secondary antibody (1:20000;

Invitrogen). Membranes were developed using the ECL Plus Western blotting substrate (Thermo

Fisher Scientific) for chemiluminescence with the Gene Gnome® Imager. Densitometry was

carried out using ImageJ software (15). The signal of each protein is expressed after subtraction

of background signal and related to actin signal.

qRT-PCR

Hippocampal tissue was collected in 250 µL cold Trizol (Invitrogen, Paisley, UK) and

homogenized using plastic pestles for homogenizers. RNA was isolated according to the

manufacturer's protocol. After addition of chloroform, samples were centrifuged at 11,000 × g

for 15 min at 4 °C. The aqueous phase was transferred and RNA was precipitated with

isopropanol, washed with 75% ethanol and resuspended in 15 µL of RNase-free DEPC-treated

water. RNA concentration of the tissue was quantified with the Qubit 2.0 Fluorometer (RNA

Assay BR Kit, Invitrogen). To ensure genomic DNA removal, 1 µg of total RNA was treated with

1 U/µL of DNase I (Invitrogen) for 15 min. RNA purity and absence of residual DNA were verified

by qPCR using non-reverse-transcribed RNA as a negative control. Samples were stored at −80

°C until further use.

Reverse Transcription Quantitative Polymerase Chain Reaction (qRT-PCR) was carried out in a

StepOnePlus Real-Time PCR System (Applied Biosystems, CA, USA), and the data were analyzed

with StepOne software v2.2. The retrotranscription and amplification reactions were performed

in a total volume of 10 µl buffer solution containing 2 µl of template RNA, 5 µl of 2× iTaq universal

SYBR® Green reaction mix (2x), 0.125 µl iScript reverse transcriptase (Applied Biosystems), 2.08

µl RNase-free DEPC-treated water and 0.4 µl of each primer (400 µM). Primers for BDNF (Brain-

Derived Neurotrophic Factor), FABP5 (Fatty Acid Binding Protein 5), Calb1 (Calbindin) and Nes

(Nestin) were designed using Primer Express software (Applied Biosystems) and described in

table 1.

The thermal cycling profile consisted of:

-UNG pre-incubation: 50 °C for 2 min.

-Initial denaturation: 95 °C for 1 min.

-Amplification (40 cycles): 95 °C for 15 s and 60 °C for 1 min.

-Melt curve analysis: 95 °C (15 s), 60 °C (1 min), and a final ramp to 95 °C at 0.6 °C/s to confirm

amplicon specificity.

To ensure the robustness of the gene expression analysis, three independent biological

replicates were processed per experimental group, following the recommendations of the MIQE

guidelines (16). Each biological sample was analyzed in triplicate using qRT-PCR. This balanced

design allows for an accurate estimation of experimental error even with biological sample sizes

of n=3, ensuring that the observed differences possess true statistical and biological significance.

Target gene expression was normalized against β-actin (ACTB) as the internal reference gene.

The dynamic range was established using a six-point standard curve derived from three

independent biological replicates, each processed in technical triplicate (n=9 per dilution). R^2

values and amplification efficiencies were calculated based on mean Cq values, incorporating a

housekeeping gene (also analyzed in triplicate) for normalization. CT (threshold cycle) values

were calculated by subtracting the CT value of ACTB from the CT values. Relative expression was

calculated as E^−ΔCt, where E is the efficiency measured for each set of primers.

Gene Symbol	Accession No.	Primer Sequence (5' →3')	E (%)	R^2	Dynamic Range
Actb	NM_007393.5	**F:** ACCAGTTCGCCATGGATGAC **R:** TGCCGGAGCCGTTGTC	97.019	0.995	10^2 - 10^7
Bdnf	NM_001285418.1	**F:** CATAGACAAAAGGCACTGGAACTC **R:** TAAGGGCCCGAACATACGAT	92.000	0.980	10^2 - 10^7
Fabp5	BC002008.1	**F:** CGACAGCTGATGGCAGAAAA **R:** CAGGCCACCGTCTTGGAA	91.788	0.993	10^2 - 10^7

Calb1	NM_009788.4	**F:** GCTCCGCGCACTCTCAA	98.828	0.940	10^2 - 10^7
		R: TGACTGCAGGTGGGATTCTG			
Nes	NM_016701.3	**F:** GGTCACTGTCGCCGCTACTC	94.788	0.989	10^2 - 10^7
		R: AAGCGGACGTGGAGCACTA			

Table 1. Primer sequences and qPCR performance characteristics. Accession No. refers to the NCBI GenBank database. E **(%)** represents the amplification efficiency calculated from the slope of the standard curve ($E=[10^{(-1/slope)}-1]\times100$. R^2 denotes the coefficient of determination for the linear regression. **Dynamic Range** indicates the interval of template concentration where linearity was maintained. All assays were performed in technical triplicate using SYBR Green chemistry on a StepOnePlus™ Real-Time PCR System.

Immunofluorescence

Newborn cells in the dentate gyrus (DG) were revealed by immunostaining against BrdU as previously published (17). Free-floating sections were exposed to 2 N HCl for 1 h. Nonspecific binding was reduced by 1 h incubation in a saturation solution containing 2 g/L gelatin (Merck) and 0.25% triton in PBS. Sections were then incubated overnight at 4 °C with a mouse anti-BrdU antibody (1:1000; Developmental Studies Hybridoma Bank, concentrated). Next day, the tissue was incubated for 2 h with an anti-mouse-A488 (1: 500; Invitrogen). All primary and secondary antibodies were diluted in the saturation solution. Finally, sections were incubated with Hoechst (1: 10000, Invitrogen) to dye the nucleus of cells. The number of BrdU-labeled cells was quantified with a fluorescence microscope (Olympus IX81) with a 20× objective, on serial sections through the entire hippocampus. One out of six series was counted (180 µm intervals, 10 sections per series). BrdU-labeled cells were counted in the sub-granular zone of the granule-cell layer over the entire DG. Cells were considered to be BrdU⁺ when their nuclei were completely filled with fluorescence or showed clear patches of variable intensity (classic dilution pattern for survival assays).

Statistical Analysis

Group differences were analyzed using one-way analysis of variance (ANOVA). Assumptions of

normality and homogeneity of variances were assessed using the Shapiro–Wilk and Bartlett

tests, respectively. When variance heterogeneity was detected, statistical analyses were

performed using either Welch's one-way ANOVA or generalized least squares (GLS) models that

explicitly accounted for heteroscedasticity. For GLS analyses, models allowing different variance

structures were compared to homoscedastic models using likelihood ratio tests, and the final

model was selected by balancing goodness-of-fit and parsimony. Post hoc comparisons were

performed using Tukey's test following classical ANOVA and GLS models, whereas Games–

Howell tests were used following Welch's ANOVA. In fatty acid analyses, p-values were adjusted

for multiple testing using the Benjamini–Hochberg false discovery rate (FDR) procedure, applied

separately to individual fatty acids and fatty acid classes. Body weight was analyzed using a two-

way repeated-measures ANOVA with week as a within-subject factor and treatment as a

between-subject factor. Sphericity was assessed using Mauchly's test and Geenhouse-Geisser

corrections were applied when necessary. In all analyses, $p < 0.05$ was considered statistically

significant, whereas values between 0.05 and 0.1 were considered a trend.

Results

Body weight

Mice were weighed across every week of treatment to control for potential changes due to the

different treatments. Analyses showed an interaction between treatment and number of weeks

(p = 0.030), with no difference in treatment (p = 0.145) but significant changes across weeks (p

<.001). As expected, given their age, all mice gained some weight along the experimental

protocol but no differences were revealed between experimental groups.

Fatty acids profile

Individual serum levels of the SFA lauric (C12:0), myristic (C14:0) and palmitic (C16:0) were not

altered by the treatments (p = 0.2482, p = 0.5138, p = 0.5962), whereas the SFA stearic (C18:0)

and lignoceric (C24:0) showed a trend to higher values in the group that consumed the

nanoencapsulated version of omega-3 fatty acids in yogurt compared to the control group (p =

0.0697) (Table 2).

Both omega-3–fortified yogurt groups exhibited a trend to lower serum oleic acid (C18:1 n-9)

levels compared with the control group (p = 0.0697), whereas no effect was observed on

palmitoleic acid (C16:1 n-7) concentrations (p = 0.5962) (Table 2).

Regarding the essential PUFAs linoleic acid (C18:2 n-6) and α-linolenic acid (C18:3 n-3), no

significant differences were observed among groups (p = 0.7495, p = 0.5138). Likewise,

arachidonic acid (C20:4 n-6) levels were not significantly affected by the treatments (p = 0.5138)

(Table 2).

Eicosapentaenoic acid (EPA, C20:5 n-3) showed a trend toward higher serum concentrations in

the supplemented groups compared with the control (ANOVA p = 0.07; FDR-adjusted p = 0.1633)

(Figure 1.A; Table 2). The control group exhibited intra-group variability, whereas both omega-

3-enriched yogurt formulations resulted in more homogeneous and consistently elevated EPA

levels. Although these differences did not reach statistical significance after multiple-

comparison correction, the observed pattern suggests a potential treatment-related effect on

circulating EPA.

Docosahexaenoic acid DHA (C22:6 n-3) levels showed no significant differences in the free acids

group compared to the control group (p = 0.1973) (Figure 1.B). However, the nanoencapsulated

yogurt formulation produced a significant increase in serum DHA levels compared with both

control and free-fatty acids groups (p = 0,00007). As expected, the group consuming the

nanoencapsulated omega-3 yogurt exhibited a higher total n-3 fatty acid content (Figure 1.C)

and a lower n-6/n-3 ratio (Figure 1.D), relative to the other groups. This finding suggests that

nanoencapsulation may protect DHA from degradation during storage and digestion, thereby

enhancing its stability and facilitating intestinal absorption, as reflected by increased serum DHA

levels.

Figure 1. Omega-3 fatty acids profile and n-6/n-3 ratio in serum. Results are expressed as the

percentage (%) of total fatty acids content. (A-B) Individual % for DHA and EPA. (C) Total % of

fatty acids from the Omega-3 family. (D) Ratio of Omega-6/Omega-3 content in serum. Data are

expressed as mean ± SD, n =4-5/group. # 0.05 < p < 0.10; *p < 0.05; **p < 0.01; ***p < 0.001

Serum composition	Control	Free fatty acids	Nano fatty acids	Statistics	P value (FDR)
Lauric acid	0.07 ± 0.023	0.072 ± 0.026	0.052 ± 0.004	$F_2 = 2.538$	0.2482
Myristic acid	0.316 ± 0.056	0.355 ± 0.066	0.304 ± 0.036	$F_2 = 1.1$	0.5138
Palmitic acid	18.2 ± 0.765	19.2 ± 1.39	19 ± 1.72	$F_2 = 0.713$	0.5962
Stearic acid	8.48 ± 0.287	9.61 ± 0.972	9.46 ± 0.446[*]	$F_2 = 9.025$	0.0697
Lignoceric acid	0.166 ± 0.044	0.205 ± 0.031	0.284 ± 0.078[*]	$F_2 = 5.692$	0.0697
Oleic acid	17.3 ± 1.01	14.7 ± 1.94[#]	14.6 ± 1.39[*]	$F_2 = 5.262$	0.0697
Palmitoleic acid	3.71 ± 0.430	3.66 ± 0.512	3.37 ± 0.414	$F_2 = 0.786$	0.5962
Linoleic acid	20.9 ± 1.23	21.6 ± 1.97	21.5 ± 0.632	$F_2 = 0.374$	0.7495
Alfa-linolenic acid	0.622 ± 0.108	0.598 ± 0.045	0.684 ± 0.084	$F_2 = 1.24$	0.5138
Arachidonic acid	14 ± 1.72	12.3 ± 1	12.9 ± 2	$F_2 = 1.166$	0.5138
EPA	0.514 ± 0.195	0.738 ± 0.033	0.75 ± 0.052	$F_2 = 3.419$	0.1633
DHA	3.83 ± 0.243	4.2 ± 0.395	5.14 ± 0.262[***]	$F_2 = 25.39$	0.00007

Table 2: Fatty acid profile in serum of mice treated with different formulations of yogurt. Results

are expressed as the percentage (%) of total fatty acids content. Data are expressed as mean ±

SD, n =4-5/group. # 0.05 < p < 0.10; *p < 0.05; **p < 0.01; ***p < 0.001

Adult Hippocampal Neurogenesis

In order to analyze whether the fatty acid formulation differently altered the process of

neurogenesis, BrdU[+] cells were quantified by immunohistochemistry in the DG of mice, both at

24 h or 4 weeks after BrdU injection to respectively evaluate the proliferative and survival stages

of the neurogenesis process. No significant changes were found in the proliferation of new cells

(p = 0.67) but it should be noted that inspection of individual data points (Fig. 2A) revealed

considerable within-group variability in all the experimental groups. Regarding survival of the

newborn cells, we found a tendency to an increase in the mice treated with the

nanoencapsulated formulation compared to control (ANOVA p = 0.0921, Post-hoc p =

0.0953941) (Fig. 2B) (Table 3).

Figure 2: Adult neurogenesis process. (A) Proliferation of new cells in the dentate gyrus (DG) of

mice treated with fatty acids supplementation, 24 hs post injection of BrdU. (B) Survival of

newborn cells in the DG, 4 weeks post injection of BrdU. Data are expressed as mean ± SD, n

=5−6/group. # 0.05 < p < 0.10.

RNA analysis

To have an insight into mechanisms behind fatty acid fortification effects in adult neurogenesis,

the HC of mice were recovered after the 4 weeks of treatment and RNA expression of different

key molecules involved in fatty acid transport, regulation of neurogenesis, and markers of young

and mature neurons were analyzed.

As stated before, FABP5 plays a role in the uptake, transport, and metabolism of fatty acids, in

particular, of DHA. A tendency to a higher FABP5 expression in mice treated with the

nanoencapsulated yogurt formulation was detected (ANOVA p = 0.0573, Post hoc p = 0.0993)

(Fig. 3A) (Table 3).

On the other hand, BDNF has been linked to the regulation of adult hippocampal neurogenesis

which makes it a big candidate for being involved in the modulation of fatty acids effects in the

brain. Indeed, we found a significant increase in BDNF RNA after the treatment (p = 0.000766)

(Fig. 3B). Even though the neurotrophin expression levels were enhanced in both yogurt treated

groups compared to control (Free-acids p = 0.0006111; Nano-acids p = 0.0137006), expression

was higher after the free-fatty acids formulation (Free vs Nano p = 0.0287450) (Table 3).

Nestin is an intermediate filament protein that identifies neuronal progenitors and young

newborn neurons, while Calbindin is a calcium binding protein which serves as a marker of

mature, differentiated neurons. As a complementary approach to the immunohistochemistry

analysis, and to study the molecular profile of the tissue, we measured RNA expression of both

proteins. We found a significant increase in Nestin and Calbindin RNA levels after both fatty acid

supplementation treatments (ANOVA Nes p = 0.000271, Calb p = 0.000575). For Nestin, the free-

fatty acid formulation showed a higher expression (p = 0.0067782) (Fig. 3C) than the

nanoencapsulated acids formulation, whereas Calbindin levels were more enhanced in the

nanoencapsulated group (p = 0.0097060) (Fig. 3D) (Table 3). The increased expression of both

neuronal markers are in line with the proneurogenic effect of EPA-DHA chronic administration

revealed by immunohistochemistry.

Figure 3: RTqPCRs. RNA was extracted from the HC of mice from the different groups to analyze

the transcription levels of (A) FABP5, (B) BDNF, (C) Nestin and (D) Calbindin. Data are expressed

as mean ± SD, n =3/group. # 0.05 < p < 0.10; *p < 0.05; **p < 0.01; ***p < 0.001

 BDNF Pathway

 To deepen the understanding of the changes produced in the BDNF pathway, which has a very

 complex regulation, protein-expression of both BDNF isoforms and their receptors was analyzed

 in the HC of yogurt-treated mice. Regarding the survival pathway, a significant increase was

 observed in mBDNF levels (ANOVA p = 0.000191) only in the animals treated with the

 nanoencapsulated yogurt formulation compared to controls (p = 0.0006301) (Fig. 4A, F),

 whereas changes in the TrkB receptor between the treated groups and the control mice were

 not significant (p = 0.4678826) (Fig. 4B, E) (Table 3). On the other hand, for the apoptotic

 pathway, no significant changes were found for proBDNF (p = 0.2076) though inspection of

 individual data points revealed considerable within-group variability in the free fatty acids group

 (Fig. 4C, F). In addition, a tendency towards a decrease in the p75 receptor was observed in mice

 treated with the free- fatty acid yogurt (ANOVA p = 0.0568, Ctrl vs Free p = 0.0969230) (Fig. 4D,

 F) (Table 3), which would mean less activation of the cell death mechanisms. Results show that

 the nanoencapsulated formulation promotes the modulation of the neurotrophin towards the

 survival pathway, which is in accordance to the previous outcomes.

 Figure 4: Protein expression of molecules from the BDNF signaling pathway determined by

393   Western Blot and representative membranes. (A)mBDNF. (B)TrkB. (C)proBDNF. (D)p75 for

394   yogurt treated mice after 4 weeks. (E–F) Representative membranes showing signals for TrkB,

395   mBDNF, pro-BDNF, and p75 in Control and fatty acids-treated mice. Data are expressed as mean

396   ± SD, n =5–6/group. # $0.05 < p < 0.10$; *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

Paradigm of assay	Parameter measured	Statistical test	Comparison	Statistics	P value	Figure
Body weight	Body weight	Two-way repeated measures ANOVA	Treatment	$F_{2,48} = 2.01$	0.145	
			Week	$F_{3.12,149.77} = 620.92$	<.001	
			Treat:Week	$F_{6.24,149.77} = 2.38$	0.03	
Fatty acids profile	% of total Omega 3	One way ANOVA	Treatment	$F_2 = 21.43$	0.00016	**1A**
Fatty acids profile	% of DHA	One way ANOVA	Treatment	$F_2 = 25.39$	0.00007	**1B**
Fatty acids profile	% of DHA	Tukey post-test	Ctrl - Free		0.19728	
			Ctrl - Nano		0.00007	
			Free - Nano		0.00175	
Fatty acids profile	% of EPA	One way ANOVA	Treatment	$F_2 = 3.419$	0.1633	**1C**
Fatty acids profile	Omega6/Omega 3 ratio	One way ANOVA	Treatment	$F_2 = 11.564$	0.01103	**1D**
Fatty acids profile	Omega6/Omega 3 ratio	Tukey post-test	Ctrl - Free		0.214	
			Ctrl - Nano		0.019	

			Free - Nano		0.097	
Proliferation of new cells	N° of BrdU+ cells	One way ANOVA	Treatment	$F_2 = 0.414$	0.67	**2A**
Survival of newborn cells	N° of BrdU+ cells	One way ANOVA	Treatment	$F_2 = 2.807$	0.0921	**2B**
Survival of newborn cells	N° of BrdU+ cells	Tukey post-test	Ctrl - Free		0.88569	
			Ctrl - Nano		0.09539	
			Free - Nano		0.21169	
qRT-PCR	Fabp5 signal relative to actin	One way ANOVA	Treatment	$F_2 = 4.781$	0.0573	**3A**
qRT-PCR	Fabp5 signal relative to actin	Tukey post-test	Ctrl - Free		0.2399	
			Ctrl - Nano		0.0993	
			Free - Nano		0.2932	
qRT-PCR	BDNF signal relative to actin	One way ANOVA	Treatment	$F_2 = 29.79$	0.00077	**3B**
qRT-PCR	BDNF signal relative to actin	Tukey post-test	Ctrl - Free		0.00061	
			Ctrl - Nano		0.0137	
			Free - Nano		0.02874	
qRT-PCR	Nestin signal relative to actin	One way ANOVA	Treatment	$F_2 = 43.37$	0.00027	**3C**
qRT-PCR	Nestin signal relative to actin	Tukey post-test	Ctrl - Free		0.00021	
			Ctrl - Nano		0.01023	
			Free - Nano		0.00678	
qRT-PCR	Calbindin signal relative to actin	One way ANOVA	Treatment	$F_2 = 33.09$	0.00057	**3D**

qRT-PCR	Calbindin signal relative to actin	Tukey post-test	Ctrl - Free		0.02599	
			Ctrl - Nano		0.00046	
			Free - Nano		0.00971	
Western blot	mBDNF signal relative to actin	One way ANOVA	Treatment	$F_2 = 19$	0.00019	**4A**
Western blot	mBDNF signal relative to actin	Tukey post-test	Ctrl - Free		0.94531	
			Ctrl - Nano		0.00063	
			Free - Nano		0.00037	
Western blot	TrkB signal relative to actin	One way ANOVA	Treatment	$F_2 = 5.04$	0.0258	**4B**
Western blot	TrkB signal relative to actin	Tukey post-test	Ctrl - Free		0.17166	
			Ctrl - Nano		0.46788	
			Free - Nano		0.02127	
Western blot	proBDNF signal relative to actin	One way ANOVA	Treatment	$F_2 = 1.797$	0.2076	**4C**
Western blot	P75 signal relative to actin	One way ANOVA	Treatment	$F_2 = 3.678$	0.0568	**4D**
Western blot	P75 signal relative to actin	Tukey post-test	Ctrl - Free		0.09692	
			Ctrl - Nano		0.99148	
			Free - Nano		0.07893	

Table 3: P values and statistical information of all procedures performed.

Discussion

The results presented herein demonstrate that chronic administration of nanoencapsulated

PUFAs to adult mice increases the bioavailability of DHA that, in turns, may promote the survival

of the newborn neurons in the HC by enhancing the pro-neurogenic elements of the BDNF

pathway.

Although there are controversies regarding establishing a recommended dietary intake range

for the omega-6/omega-3 ratio (18–21), Zhang et al. (2024), in a UK Biobank large population

cohort study, revealed a strong association between the plasma omega-6/omega-3 PUFA ratio

and the risk of all-cause mortality, cancer, and cardiovascular disease (22). This supports dietary

interventions to increase omega-3 fatty acid levels and maintain a lower omega-6 to omega-3

fatty acid ratio as part of a healthy lifestyle.

In *Wistar* rats fed for 10 days a normocaloric but fat-unbalanced diet rich in saturated fats from

butter, supplementation with fish oil (EPA + DHA) or chia oil (ALA) increased circulating levels of

EPA and DHA compared to the non-supplemented group. These results indicate that the lipid

composition of food rapidly influences the serum fatty acid profile even in a diet rich in saturated

fats (14). Similarly, in the present rodent model, although PUFAs quantities ingested by different

animals may have varied depending on the amount of cube ingested, EPA showed a trend

toward higher serum concentrations in the supplemented groups, and the nanoencapsulated

yogurt formulation produced a significant increase in serum DHA levels. Consequently, the

group consuming the nanoencapsulated omega-3 yogurt exhibited a higher total n-3 fatty acid

content and a lower n-6/n-3 ratio, relative to the other groups (total n-3) and to the control

group (n-6/n-3 ratio). Indeed, in vitro studies have already demonstrated increased digestibility

and bioaccessibility of n-3 PUFAs when administered in nanoemulsions (7,23). Our finding

suggests that nanoencapsulation may protect DHA from degradation during storage and

digestion, thereby enhancing its stability and facilitating intestinal absorption, as reflected by

increased serum DHA levels. Nanoencapsulation systems might protect PUFAs from gastric

degradation and optimize intestinal micellization, significantly increasing their concentration (7).

The increased serum omega-3 levels in murine models stem from a tripartite synergy between

the PUFAs, the soy lecithin nanoliposomes, and the yogurt matrix. In the acidic environment

(pH≈4.5), nanoliposomes provide a robust physical barrier against oxidative and proton-induced

degradation (24). This is reinforced by a "protein-lipid corona", formed via hydrophobic and

electrostatic adsorption of caseins and whey proteins, which ensures steric stabilization and

prevents coalescence during storage and gastric transit (25). In the gastrointestinal tract, these

phosphatidylcholine-rich vesicles function as high-efficiency delivery vehicles. Unlike bulk oils,

they facilitate rapid micellization, effectively bridging the unstirred water layer to deliver

concentrated PUFAs to the enterocyte apical membrane (26,27). At the cellular level, soy

phospholipids upregulate FABP, optimizing intracellular trafficking (28). Furthermore, lecithin

provides essential precursors for chylomicron assembly, enhancing fatty acid re-esterification

and accelerating lymphatic exocytosis. Consequently, this integrated mechanism—from the

protective dairy-protein corona to facilitated transport—underpins the superior systemic

bioavailability observed (29).

Positive correlations between plasma and brain levels of DHA were already described (30,31).

Therefore, we could assume that the observed higher plasmatic levels of PUFAs may have a

correlate in the brain of treated mice. In the present study, the FABP-5, a key regulator of DHA

uptake in the brain, showed a trend to increase only in the group that received the

nanoencapsulated PUFAs. As this group showed the higher plasmatic levels of DHA, this result

could altogether show a better bioavailability and distribution of Omega-3 fatty acids under the

new formulation. In an elegant series of studies, Pan et al. demonstrated up-regulated

expression of brain microvascular FABP5 protein associated with increased brain DHA levels in

adult mice fed with a DHA-enriched diet for 21 days (32). Later they reported that mice knock-

out for the Fabp5 gene exhibited impaired performances in memory tasks, associated with a

reduction in endogenous brain DHA levels (33). These results, together with ours support the

hypothesis that increased plasmatic DHA promotes the expression of FABP-5 in the brain,

facilitating in turn the increase of DHA levels in the brain.

DHA has been inversely associated with cognitive decline in clinics (34). Indeed, populations with

a high consumption of fatty fish or fatty acid supplements display lower incidence of

neurodegenerative diseases compared to those consuming diets poor in these nutrients (35).

The process of neurogenesis offers a promising avenue in the search for therapies for aging and

neurodegenerative diseases, either by replacing lost neurons or by enhancing brain plasticity.

The generation of newborn neurons is well characterized in the adult hippocampus, particularly

in the DG, where several factors like neurotrophines, have been shown to facilitate the

proliferation and/or survival of newborn neurons (see rev in (36). We were not able to see

changes in the proliferation rate of DG neurons in adult mice receiving PUFAs, but mice treated

with the nanoencapsulated n-3 Omega fatty acids displayed a trend to an increase in the survival

of 4-week old neurons, compared with the control group. The enlarged neurogenesis was

accompanied by a significant increase of immature and mature markers of neuronal

differentiation like Nestin and Calbindin, respectively. Expression of both markers was also

enhanced in mice receiving the free acids, revealing that, globally, the n-3 omega fatty acids

appear to act at different stages of the neurogenic process, i.e, differentiation, and survival of

newborn neurons. Increased neuron survival has been already shown in a model of strong deficit

of PUFA where rats fed a fish-oil deficient diet for three generations were then, supplemented

with DHA for 7 weeks (37). Another report from rats fed a DHA and EPA-enriched diet during 4

weeks revealed an increase in hippocampal levels of the BDNF as measured by ELISA (10). Similar

results were observed for BDNF mRNA and protein levels in a transgenic mice model of

Alzheimer's disease after 2 months of a DHA-enriched diet (11) and in C57BL/6 mice chronically

supplemented with lysophosphatidylcholine-DHA or -EPA (38,39). Our results shed light on

these reports describing plastic effects of PUFAs in rodents, as we were able to go deep in the

analysis of the BDNF biological pathway. In this study, increased levels of total BDNF mRNA were

detected in both groups of mice that received the n-3 omega fatty acids compared to the control

groups. However, when the level of proteins were analyzed, mice receiving the free fatty acids

showed a sort of activation of the BDNF pro-apoptotic cascade, with a tendency towards higher

levels of the p-75 receptor and something similar for pro-BDNF compared to the other

experimental groups. In this line, a previous study in a mice model of Alzheimer disease found

that the administration of a DHA/EPA-enriched diet for 2 months decreased the expression of

pro-apoptosis factors like Bcl2, Caspase 3, and Caspase 9 (11). On the contrary, the

proneurogenic elements of this biological pathway, like mBDNF and its receptor TrkB, were

significantly increased after chronic administration of nanoencapsulated PUFAs, in synthon with

the increased survival of the newborn neurons detected in the DG.

Conclusions

In this work, nanoencapsulated PUFAs orally administrated to male mice for 4 or 8 weeks

showed better bioavailability when compared to administration of free acids, and this

potentially due to an increase of the specific transporter, the FABP-5 particularly in the HC.

Furthermore, the nanoformulation enhanced the survival of the DG newborn neurons, possibly

in response to an improved proneurogenic balance of BDNF actions. These positive effects can

be key to facilitate memory processes and also, to prevent cognitive decline that can occur at

advanced ages.

References

1.   McNamara RK, Carlson SE. Role of omega-3 fatty acids in brain development and
function: Potential implications for the pathogenesis and prevention of
psychopathology. Prostaglandins Leukot Essent Fatty Acids. 2006;75:329–349. doi:
10.1016/j.plefa.2006.07.010. Cited in PMID: 16949263.

2.   Rahim MA, Uzma M, Al-Asmari F, Koraqi H, Ramadan MF, Castro-Muñoz R, Zongo E. A
Comprehensive Review on Nutritional Traits, Extraction Methods, Oxidative Stability,
Encapsulation Technologies, Food Applications and Health Benefits of Omega Fatty
Acids. Food Sci Nutr. 2025;13. doi: 10.1002/fsn3.71008.

3. Mitchell RW, On NH, Del Bigio MR, Miller DW, Hatch GM. Fatty acid transport protein
expression in human brain and potential role in fatty acid transport across human brain
microvessel endothelial cells. J Neurochem. 2011;117:735–746. doi: 10.1111/j.1471-
4159.2011.07245.x. Cited in PMID: 21395585.

4. Iwao T, Takata F, Matsumoto J, Aridome H, Yasunaga M, Yokoya M, Kataoka Y, Dohgu S.
Aging decreases docosahexaenoic acid transport across the blood-brain barrier in
C57BL/6J mice. PLoS One. 2023;18. doi: 10.1371/journal.pone.0281946. Cited in PMID:
36795730.

5. Clemente-Suárez VJ, Beltrán-Velasco AI, Redondo-Flórez L, Martín-Rodríguez A,
Tornero-Aguilera JF. Global Impacts of Western Diet and Its Effects on Metabolism and
Health: A Narrative Review. Nutrients. 2023;15. doi: 10.3390/nu15122749. Cited in
PMID: 37375654.

6. Inapurapu SP, Ibrahim A, Kona SR, Pawar SC, Bodiga S, Bodiga VL. Development and
characterization of ω-3 fatty acid nanoemulsions with improved physicochemical
stability and bioaccessibility. Colloids Surf A Physicochem Eng Asp. 2020;606. doi:
10.1016/j.colsurfa.2020.125515.

7. Yang J, Ciftci ON. In vitro bioaccessibility of fish oil-loaded hollow solid lipid micro- A nd
nanoparticles. Food Funct. 2020;11:8637–8647. doi: 10.1039/d0fo01591a. Cited in
PMID: 32936172.

8. Kempermann G, Song H, Gage FH. Neurogenesis in the adult hippocampus. Cold Spring
Harb Perspect Biol. 2015;7:a018812. doi: 10.1101/cshperspect.a018812. Cited in PMID:
26330519.

9. Foltran RB, Diaz SL. BDNF isoforms: a round trip ticket between neurogenesis and
serotonin? J Neurochem. 2016;138:204–221. doi: 10.1111/jnc.13658.

10. Wu A, Ying Z, Gomez-Pinilla F. Dietary Omega-3 Fatty Acids Normalize BDNF Levels,
Reduce Oxidative Damage, and Counteract Learning Disability after Traumatic Brain
Injury in Rats. J Neurotrauma. 2004;21.

11. Che H, Zhou M, Zhang T, Zhang L, Ding L, Yanagita T, Xu J, Xue C, Wang Y. Comparative
study of the effects of phosphatidylcholine rich in DHA and EPA on Alzheimer's disease
and the possible mechanisms in CHO-APP/PS1 cells and SAMP8 mice. Food Funct.
2018;9:643–654. doi: 10.1039/c7fo01342f. Cited in PMID: 29292421.

12. FAO, WHO. Report of the Joint FAO/WHO Expert Consultation on the Risks and Benefits
of Fish Consumption. Rome; 2011.

13. Foltran RB, Stefani KM, Hocht C, Diaz SL. Neurochemical, behavioral, and neurogenic
validation of a hyposerotonergic animal model by voluntary oral consumption of PCPA.
ACS Chem Neurosci. 2020; doi: 10.1021/acschemneuro.9b00687.

14. Fernandez I, Giacomino MS, Condori AI, Godoy MF, Pellegrino N, Slobodianik N, Feliu
MS. Effect of n-3 fatty acids supplementation on serum lipid profile of rats. Revista
Chilena de Nutricion. 2021;48:170–178. doi: 10.4067/S0717-75182021000200170.

15. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
analysis. Nat Methods [Internet]. 2012;9:671–675. Cited in PMID: 22930834.

16. Bustin SA, Benes V, Garson JA, Hellemans J, Huggett J, Kubista M, Mueller R, Nolan T,
Pfaffl MW, Shipley GL, et al. The MIQE guidelines: Minimum information for publication
of quantitative real-time PCR experiments. Clin Chem. 2009;55:611–622. doi:
10.1373/clinchem.2008.112797. Cited in PMID: 19246619.

17. Diaz SL, Doly S, Narboux-Nême N, Fernández S, Mazot P, Banas SM, Boutourlinsky K,
Moutkine I, Belmer A, Roumier A, et al. 5-HT2B receptors are required for serotonin-
selective antidepressant actions. Mol Psychiatry. 2012;17:154. doi:
10.1038/mp.2011.159. Cited in PMID: 22158014.

18. Food and Agriculture Organization [FAO]. Fats and fatty acids in human nutrition:
report of an expert consultation. FAO Food Nutr Pap. 2010;91:1–166.

19. EFSA Panel on Dietetic Products N and A (NDA). Scientific Opinion on the Tolerable
Upper Intake Level of eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA) and
docosapentaenoic acid (DPA). EFSA Journal. 2012;10. doi: 10.2903/j.efsa.2012.2815.

20. Simopoulos AP. The importance of the omega-6/omega-3 fatty acid ratio in
cardiovascular disease and other chronic diseases. Exp Biol Med. 2008;233:674–688.
doi: 10.3181/0711-MR-311. Cited in PMID: 18408140.

21. Djuricic I, Calder PC. Beneficial outcomes of omega-6 and omega-3 polyunsaturated
fatty acids on human health: An update for 2021. Nutrients. 2021;13. doi:
10.3390/nu13072421. Cited in PMID: 34371930.

22. Zhang Y, Sun Y, Yu Q, Song S, Thomas Brenna J, Shen Y, Ye K. Higher ratio of plasma
omega-6/omega-3 fatty acids is associated with greater risk of all-cause, cancer, and
cardiovascular mortality: A population-based cohort study in UK Biobank. Elife.
2023;12:90132. doi: 10.7554/eLife.

23. Zhou Q, Lane KE, Li W. Evaluating the Stability and Digestibility of Long-Chain Omega-3
Algal Oil Nanoemulsions Prepared with Lecithin and Tween 40 Emulsifiers Using an In
Vitro Digestion Model. Foods. 2024;13. doi: 10.3390/foods13152407.

24. Ghorbanzade T, Jafari SM, Akhavan S, Hadavi R. Nano-encapsulation of fish oil in nano-
liposomes and its application in fortification of yogurt. Food Chem. 2017;216:146–152.
doi: 10.1016/j.foodchem.2016.08.022. Cited in PMID: 27596403.

25. Taneja A, Ye A, Singh H. Effects of the milk matrix on the bioavailability of vitamins and
minerals. In: McLean JK, Anderson AS, editors. Food-based approaches for a healthy
diet. Woodhead Publishing; 2010. p. 233–256.

26. Kim YM, Jang GH, Seok CH, Kim BH, Bae JW, Kim BH, Yoon MS. A self-emulsifying
omega-3 fatty acids delivery system for enhanced gastro-intestinal absorption in rats.
Food Science and Biotechnology 2022 31:13 [Internet]. 2022 [cited 2026 Mar
11];31:1631–1638. doi: 10.1007/s10068-022-01151-7.

27. Ahmmed MK, Ahmmed F, Tian H, Carne A, Bekhit AED. Marine omega-3 (n-3)
phospholipids: A comprehensive review of their properties, sources, bioavailability, and
relation to brain health. Compr Rev Food Sci Food Saf. 2020;19:64–123. doi:
10.1111/1541-4337.12510. Cited in PMID: 33319514.

28.   Lane KE, Li W, Smith C, Derbyshire E. The bioavailability of an omega-3-rich algal oil is
improved by nanoemulsion technology using yogurt as a food vehicle. Int J Food Sci
Technol. 2014;49:1264–1271. doi: 10.1111/ijfs.12455.

29.   Mozafari MR, Johnson C, Hatziantoniou S, Demetzos C. Nanoliposomes and their
applications in food nanotechnology. J Liposome Res. 2008;18:309–327. doi:
10.1080/08982100802465941. Cited in PMID: 18951288.

30.   Hashimoto M, Hossain S, Shimada T, Sugioka K, Yamasaki H, Fujii Y, Ishibashi Y, Oka JI,
Shido O. Docosahexaenoic acid provides protection from impairment of learning ability
in Alzheimer's disease model rats. J Neurochem. 2002;81:1084–1091. doi:
10.1046/j.1471-4159.2002.00905.x. Cited in PMID: 12065621.

31.   Orchard TS, Gaudier-Diaz MM, Phuwamongkolwiwat-Chu P, Andridge R, Lustberg MB,
Bomser J, Cole RM, Belury MA, Devries AC. Low sucrose, omega-3 enriched diet has
region-specific effects on neuroinflammation and synaptic function markers in a mouse
model of doxorubicin-based chemotherapy. Nutrients. 2018;10. doi:
10.3390/nu10122004. Cited in PMID: 30567351.

32.   Pan Y, Morris ER, Scanlon MJ, Marriott PJ, Porter CJH, Nicolazzo JA. Dietary
docosahexaenoic acid supplementation enhances expression of fatty acid-binding
protein 5 at the blood–brain barrier and brain docosahexaenoic acid levels. J
Neurochem. 2018;146:186–197. doi: 10.1111/jnc.14342. Cited in PMID: 29582413.

33.   Pan Y, Short JL, Choy KHC, Zeng AX, Marriott PJ, Owad Y, Scanlon MJ, Porte CJH,
Nicolazzo JA. Fatty acid-binding protein 5 at the blood–brain barrier regulates
endogenous brain docosahexaenoic acid levels and cognitive function. Journal of
Neuroscience. 2016;36:11755–11767. doi: 10.1523/JNEUROSCI.1583-16.2016. Cited in
PMID: 27852782.

34.   Mallick R, Basak S, Duttaroy AK. Docosahexaenoic acid,22:6n-3: Its roles in the structure
and function of the brain. International Journal of Developmental Neuroscience.
2019;79:21–31. doi: 10.1016/j.ijdevneu.2019.10.004. Cited in PMID: 31629800.

35.   Kalmijn S, Van Boxtel MPJ, Ocké M, Verschuren WMM, Kromhout D, Launer LJ. Dietary
intake of fatty acids and fish in relation to cognitive performance at middle age.
Neurology. 2004;62:275–280.

36.   Balu DT, Lucki I. Adult hippocampal neurogenesis: Regulation, functional implications,
and contribution to disease pathology. Neurosci Biobehav Rev. 2009;33:232–252. doi:
10.1016/j.neubiorev.2008.08.007. Cited in PMID: 18786562.

37.   Kawakita E, Hashimoto M, Shido O. Docosahexaenoic acid promotes neurogenesis in
vitro and in vivo. Neuroscience. 2006;139:991–997. doi:
10.1016/j.neuroscience.2006.01.021.

38.   Sugasini D, Thomas R, Yalagala PCR, Tai LM, Subbaiah P V. Dietary docosahexaenoic acid
(DHA) as lysophosphatidylcholine, but not as free acid, enriches brain DHA and
improves memory in adult mice. Sci Rep. 2017;7. doi: 10.1038/s41598-017-11766-0.
Cited in PMID: 28900242.

39.   Yalagala PCR, Sugasini D, Dasarathi S, Pahan K, Subbaiah P V. Dietary
lysophosphatidylcholine-EPA enriches both EPA and DHA in the brain: Potential

treatment for depression. J Lipid Res. 2019;60:566–578. doi: 10.1194/jlr.M090464.
Cited in PMID: 30530735.

631