1

4 Prioritization in working memory reduces interference via a beta-linked transformation of

5 the not-selected item

7 Jacqueline M. Fulvio[1*], Bradley R. Postle[1,2]

9 [1]Department of Psychology, University of Wisconsin—Madison, Madison, Wisconsin,
10 United States of America
11
12 [2]Department of Psychiatry, University of Wisconsin—Madison, Madison, Wisconsin,
13 United States of America
14

15 *Corresponding author

16 Email: jacqueline.fulvio@wisc.edu (JMF)

Abstract

We studied the effects of prioritization in a two-step retrocuing task in which participants hold two items in working memory, and the item not cued by the first cue cannot be dropped because it may be prioritized by the second cue. In Experiment 1, using a dense sampling procedure, we observed that recall performance oscillated at 15 Hz in the prioritization task, in comparison to 20 Hz in a matched task employing a neutral cue. Furthermore, the prioritized item was shielded from bias exerted by the uncued item, as well as from items from the previous trial. In Experiment 2, we recorded the EEG while participants performed variants of the two tasks. The prioritization cue uniquely triggered a phase reset at 15 Hz and an increase in oscillatory peaks at this frequency. Burst analysis ruled out bursting as a possible underlying factor. Time-resolved representational similarity analysis (RSA) revealed that the prioritization cue triggered representational transformations that were larger for the uncued item. The shielding effects of prioritization may arise from the transformation of the not-prioritized item into an "unprioritized" state that is implemented and maintained by a mechanism that cycles at 15 Hz.

Introduction

When multiple items are held in working memory, validly retrocuing the item that will subsequently be tested has consequences for behavior and for how this information is represented in the brain. Behaviorally, retrocuing results in higher accuracy and faster reaction times (as reviewed, e.g., by [1]), effects interpreted as evidence that the cued item has become "more accessible" [1]. Neurally, retrocuing an item improves its decodability (interpreted as evidence for a "strengthening" of its neural representation, e.g., [2, 3]), and transforms its representational geometry [4]. One question that motivates the current study relates to the potential "shielding" effects of prioritization. Intuitively, one might assume that in addition to becoming "stronger," the cued item might also become more resistant to the potentially interfering effects of other information being processed by the cognitive system, such as perceptual distractors. Empirically, however, evidence for such a "shielding" effect is mixed (as reviewed, e.g., by [1]).

Importantly for the present report, the majority of studies that have used a retrocuing procedure have done so with tasks that only require a single response, thereby rendering the uncued item an "irrelevant memory item" (IMI) that can be "dropped" from working memory as soon as the cue has been processed. For example, in a recurrent neural network (RNN) simulation of the retrocuing task from [4], after cuing, the IMI simply fades from working memory [5]. The focus in this report, in contrast, will be on a task in which the not-cued item cannot be dropped from working memory because it might still be needed later in the trial.

An example of a situation in which the not-prioritized item cannot be dropped

from working memory is the 2-back task, a continuous performance task in which each

newly presented item in a series must be judged as a match or nonmatch of the item

that had appeared two previously. While item n is on the screen, it serves as a

recognition probe for item $n - 2$; next it becomes an "unprioritized memory item" (UMI)

while $n + 1$ is being compared to $n - 1$; next it becomes the "prioritized memory item"

(PMI) against which $n + 2$ is compared, and finally it becomes an IMI. In

electroencephalography (EEG) data, the transition from probe to UMI is accompanied

by the reversal of an item's reconstruction by a multivariate inverted encoding model

(IEM), a phenomenon equivalent to significantly below-chance decoding [6]. In RNN

simulation, the UMI and PMI are represented in ring manifolds that are reversed relative

to one another, an item's geometry underdoing a scalar flip during the transition from

UMI to PMI [6], then, upon the transition to IMI status, collapsing to a scalar value

undifferentiable from 0 (i.e., "fading until no trace remains") [6]. These considerations

highlight two factors that provide important motivation for the present studies. First, they

highlight that the nature of possible shielding effects bestowed by prioritization may be

very different for tasks in which the UMI cannot be dropped from working memory (in

comparison to single-cue tasks like those summarized above). Second, in addition to

perceptual distractors, a potentially important source of interference in cannot-drop-the-

UMI tasks is, indeed, the UMI. To our knowledge, no previous studies of interference

between items concurrently held in working memory (what we will refer to as "within-

trial" interference) have examined the effects of prioritization.

The research presented here uses variants of a double serial retrocuing (DSR)

task, in which the presentation of two sample items is followed by an initial retrocue,

then an initial test of recall, then a second retrocue, then a second test of recall.

Although initial work with the DSR task suggested that an item might transition into an

"activity-silent" state upon being designated a UMI by the initial cue (e.g. [7, 8]), more

recent findings [3] indicate that the UMI can undergo a representational transformation

similar to what has been reported with the 2-back task [6]. Additionally, and of particular

relevance for this report, is the fact that previous work with the DSR task has also

implicated a possible role for oscillations in the beta band: beta-band dynamics relate to

the effects of transcranial magnetic stimulation (TMS) on behavior [9], and they make

selective contributions to decoding the identity of the UMI [8, 10]. Previous working

memory research has highlighted at least two distinct mechanisms operating in the beta

band. One is phase encoding, whereby each of two items in working memory is held at

a different phase angle in ongoing theta-band (~6 Hz) and beta-band (~25 Hz)

oscillations [11]. A second is that, in a manner analogous to the stopping of

skeletomotor actions [12, 13], bursts in the beta frequency range may provide a

mechanism for imposing top-down control on the processing of information in working

memory [14]. The research presented here was also designed to assess whether the

beta-band dynamics associated with prioritization in the DSR task [9, 10, 15] may

correspond to phase encoding [11], to bursty top-down control [14], or to a third

previously undescribed phenomenon.

Experiment 1

Participants (*n*=16) recalled the orientation of oriented gratings in a double serial

retrocuing (DSR) task in which two items were presented serially and the item to be

recalled was cued on two successive occasions with a digit referring to ordinal position

("1" or "2;" 100% informative cues, each with 50% probability for both retrocuing

occasions; **Fig 1A** see **Materials and Methods**). In a separate task, the same

participants performed a single recall on trials featuring an uninformative cue ("0")

accompanying the recall dial (**Fig 1B**). On each trial, the two memoranda could differ in

orientation by 0, 22.5, 45, or 67.5 degrees, and the post-cue to recall delay period was

densely-sampled in the range of 500 ms to 1200 ms.

Fig 1. Experiment 1 behavioral task schematics. A. Schematic of the double-serial retrocue (DSR)

task. Participants memorized the identity of two centrally presented oriented grating samples shown at

successive time points with an intervening delay ('*ISI*'). Partway through the delay (i.e., at the end of

Delay 1.1) an ordinal retrocue indicated which of the two items would be tested, after the ensuing *Delay*

1.2 of variable length, sampled uniformly from .5-1.2 s in 16.67 ms increments. At test (*Recall 1*),

participants rotated the recall dial with the mouse to the orientation of the tested sample item in memory.

After feedback, a second ordinal retrocue (*Cue 2*) indicating which of the two samples would be tested

during *Recall* 2 followed by a final delay. **B.** Schematic of the load-of-2 neutral cue (NEU) task, which

followed the same task structure as the DSR task through the first feedback interval, after which the trial

ended. Importantly, an uninformative '0' was presented during the cue interval, preventing participants

from anticipating which item would be tested at recall.

Recall accuracy

Mean absolute error (MAE) of recall did not differ between DSR *Recall 1* and NEU recall

(t(15) = 0.6046, p = .555), whereas the MAE of DSR *Recall 2* was significantly larger

than for DSR *Recall 1* and NEU recall (both *p*-values < .01; **Fig. 2A**). No difference was

observed between the MAE of DSR *Recall 2* for 'stay' versus 'switch' trials (t(15) =

0.0526, *p* = .959; **Figure A in S1 Fig**). Larger inter-item angular differences were

associated with larger errors, with a linear mixed effects model confirming a significant

main effect of current trial inter-item angular difference on performance (F(1,189) =

140.81, *p* < .001; **Fig. 2B**). When the model included both recall responses of the DSR

task, a significant main effect of task was also observed (F(1,189) = 9.9764, *p* = .002),

driven by the worse performance for *Recall 2* of the DSR task (see also **Figure B in S1**

Fig).

Fig 2. Experiment 1 recall accuracy. A. Mean absolute recall error for the two responses of the DSR

task and the single response of the NEU task. **B.** Same as **A.** separated by the current trial's inter-item

angular difference. **C.** Mean absolute error for DSR *Recall 1* and NEU recall as a function of which item

was tested (*Sample 1* vs. *Sample 2*). Error bars correspond to +/- 1 *SEM*. * symbols correspond to

significant differences between conditions at the Bonferroni-corrected alpha level = .0167 for 3 paired-

sample *t*-tests (**A.**) and the Bonferroni-corrected alpha level = .025 for two paired-sample *t*-tests and for

interaction *p* < .05 (**C.**).

Finally, comparison of MAE as a function of the presentation order of the

PMI/tested item revealed different patterns between NEU recall and DSR *Recall 1*: In

NEU, MAE was significantly smaller when *Sample 2* was tested (t(15) = 2.805, *p* =

.0133), whereas in DSR, MAE was significantly smaller when *Sample 1* was tested

(t(15) = -2.734, *p* = .0154); **Fig. 2C**). This pattern resulted in a significant performance

crossover between the two tasks (t(15) = -3.849, *p* = .0016).

Across- and within-trial bias

Linear mixed effects modeling revealed that response error was systematically

influenced by both the previous trial's tested item and the current trial's untested item,

with the latter effect dependent on the order of presentation (**S1 Table**). In both tasks,

the previous trial's tested item exerted an attractive bias on recall (NEU: β = 1.08°, p <

.001; DSR: β = 0.84°, p = 0.002), an effect that did not differ across task (β = 0.25°, p =

.517; **S2 Table**). The previous trial's untested item had no significant effect on recall.

Turning to within-trial effects, recall of *Sample 1* was attracted toward the value of

Sample 2 for NEU (β = 3.55°, p < .001) and for DSR (β = 1.66°) but recall of *Sample 2*

showed no significant bias in NEU (β = 0.08°, p = .818) or DSR (β = 0.50°, p = .177).

The three-way interaction between task, current-trial untested item, and order-of-

presentation was significant (β = –2.31°, p = .002; **S2 Table**), confirming that the effect

of prioritization was to reduce the influence of *Sample 2* on the recall of *Sample 1*.

Linear mixed effects modeling further revealed that inter-item angular difference

significantly modulated bias ($F(2,3977)$ = 7.16, p < .001; **S3 Table**) in the NEU task, and

this effect interacted with order of presentation ($F(2,3977)$ = 6.35, p = .002). Recall of

Sample 1 was significantly attracted toward *Sample 2* at small and medium angular

separations (22.5°: 3.68°, p < .001; 45°: 3.72°, p < .001), with attraction reduced to a

marginally significant level at the large angular separation (67.5°: 1.09°, p = .052; a

significant reduction from the bias at 22.5: β = -2.59°, p = .001). Recall of *Sample 2*, in

contrast, showed no significant bias relative to *Sample 1* at any angular separation

(22.5°: -0.42° p = .461; 45°: 0.01° p = .987; 67.5°: 0.63° p = .251). The main effect of

presentation order was significant ($F(1,3977)$ = 26.72, p < .001). In DSR, bias was small

and the pattern of bias did not vary significantly across inter-item angular differences

(F(2,3975) = 0.70, p = .499) nor was there an interaction of inter-item angular difference

interaction with order of presentation (F(2,3975) = 1.65, p = .192). Recall of *Sample 1*

was attracted to *Sample 2* at medium and large angular separations (45°: 1.76°, p =

.002; 67.5°: 1.42°, p = .013) but not at the small separation (22.5°: 0.81°, p = .155).

Recall of *Sample 2*, in contrast, showed no significant bias relative to *Sample 1* at small

and medium angular separations (22.5°: -0.71°, p = .208; 45°: 0.22°, p = .698), but

attraction at the large angular separation (67.5°: 1.67°, p = .003). Importantly, the

previous trial's target, which exerted a significant attractive bias on recall in the primary

analysis, did not significantly influence signed bias in either task. This indicates that the

previous trial's influence operated separately in absolute orientation space, pulling

responses toward the previous target's orientation, rather than by influencing

interactions between the current trial's items.

The task comparison model confirmed that NEU showed significantly greater

overall bias than DSR (F(1,7954) = 12.98, p < .001), and that the effect of angular

difference was significantly larger in NEU than DSR (interaction: F(2,7954) = 4.21, p =

.015; **S4 Table**).

Behavioral oscillations

Permutation-based FFT analysis revealed significant rhythmic fluctuation of mean

absolute error, at different frequencies, on both tasks (**Fig. 3A**). DSR showed significant

behavioral oscillatory peaks clustered around 15 Hz (14.06, 15.00, and 15.94 Hz), with

the dominant frequency at 15.00 Hz (amplitude = 0.34 degrees). The NEU task

exhibited significant behavioral oscillations with the dominant frequency at 20.62 Hz

(amplitude = 0.18 degrees; range = 19.69 and 20.62 Hz). Participant-level peak

frequency analysis confirmed that the peak oscillatory frequencies differed significantly

between conditions (*M* = 5.15 Hz; *95% CI*: [3.38, 6.93] Hz; *t*(15) = -6.18, *p* < 0.001).

From here forward, to relate them to the classical frequency band of EEG oscillation, we

will refer to these two frequencies as "low-beta" and "higher-beta," respectively.

Fig 3. Experiment 1 behavioral oscillations of mean absolute error. A. Demeaned mean absolute

recall error for DSR R1 (red) and the single response of the NEU task (blue) as a function of Delay 1.2

length. Error bands correspond to +/- 1 *SEM.* **B**: Fast Fourier Transform (FFT) amplitude spectra of mean

absolute error across delay periods for the DSR (red) and NEU (blue) tasks. Solid lines represent group-

averaged amplitudes (in degrees of error). Dashed lines show statistical significance thresholds (95th

percentile) derived from permutation testing. * symbols indicate peaks surviving Bonferroni-correction for

number of peaks exceeding statistical threshold. **C.** Polar plots depicting phase angles at the dominant

oscillatory frequency in the DSR task (15.00 Hz; left) and the NEU task (20.62 Hz; right). Individual

participants are shown as dots at unit radius. The black line represents the mean resultant vector, with its

length corresponding to the Phase Locking Value (PLV) and direction indicating the mean phase angle. *

denotes significant PLV.

Between-subject phase alignment was quantified for the dominant frequency of

each task. In the DSR task, participants exhibited phase locking of performance

fluctuations at 15 Hz (Phase Locking Value [PLV] = 0.628, permutation *p* = 0.0008). In

the NEU task, between-subject phase alignment was not significant at 20 Hz (PLV =

0.383, permutation p = 0.095). Rayleigh tests for non-uniformity confirmed these

findings (DSR: *z* = 6.31, *p* = 0.001; NEU: *z* = 2.34, *p* = 0.095; **Fig. 3B**). To test whether

between-subject phase alignment differed between the tasks, we conducted

permutation tests comparing PLVs. When comparing each task at its dominant

frequency (DSR at 15 Hz vs. NEU at 20.62 Hz), we found that the PLV difference was

not significant (ΔPLV = 0.245, p = 0.33). Thus, while DSR exhibited significant between-

subject phase alignment at 15 Hz and NEU did not at 20.62 Hz, the magnitude of phase

locking did not differ reliably between tasks.

When we applied the FFT analysis to inter-item bias, no significant oscillatory

peaks were observed in either task (all p > 0.05), indicating that, unlike MAE, inter-item

bias did not vary rhythmically (**S2 Fig**).

Interim discussion—Experiment 1

The results from Experiment 1 provided support for two effects of prioritization on

working memory. First, prioritization in the DSR task was associated with a reduction in

inter-item interference at recall: The biasing influence of *Sample 2* on the recall of

Sample 1 was markedly reduced. Furthermore, although small biases were present at

some angular separations, a critical distinction from NEU was that these biases were

uniform across separation magnitudes, whereas for NEU, attraction was strong when

items were similar and reduced when items were dissimilar. This suggests that

prioritization has a general effect of attenuating inter-item interference, reducing

sensitivity to the similarity between items. Second, prioritization in the DSR task was

associated with a lower frequency of behavioral oscillation of mean absolute error (15

Hz; "low-beta") relative to the NEU task (20 Hz; "higher-beta"). Furthermore, only in

DSR was the cue associated with phase-locking of the behavioral oscillation across

participants (although this effect did not differ significantly between the two tasks).

Coupled with the absence of evidence of rhythmic modulation of the strength of inter-

item bias, these results suggest that behavioral oscillations of MAE reflect fluctuations in

representational strength rather than periodic distortions in the remembered content.

This evidence for behavioral oscillations in the beta band is of particular interest in light

of previous work demonstrating an important role for beta-band dynamics in the

encoding of priority in working memory [8, 9, 10].

Rhythmicity is a prominent feature of neural systems supporting visual attention

[22, 23] and visual working memory [11, 24, 25]. One longstanding and influential

perspective holds that, when multiple items are held in working memory, the encoding of

individual items at different phases of a low-frequency oscillation provides a mechanism

for individuating items and minimizing interference between them (e.g., [11, 26]), and

may even be a basis for understanding capacity limitations [27, 28]. For human visual

working memory, one study recording the electroencephalogram (EEG) while subjects

held two items in mind has identified evidence for phase coding in the beta band (albeit

at a higher frequency than the low-beta and higher-beta that we observed in Experiment

1; [11]). Thus, one prediction that we planned to test in Experiment 2 was that phase

encoding of stimulus information in the low-beta band may be a mechanism underlying

the shielding effect of prioritization.

The idea of phase encoding in the beta band (c.f., [11]), together with Experiment

1's evidence for beta-clocked behavioral oscillations, raise an important question about

EEG dynamics observed during working memory: to what extent is this activity reflective

of one or more truly periodic processes versus activity that, when assessed at the level

of the single trial, is fundamentally bursty. In skeletomotor control, "beta bursts" have a

prominent role in stopping the execution of motor actions and, by analogy, they have

been proposed as a mechanism for imposing inhibitory control across many domains of

high-level cognition, including long-term memory, visual attention, visual working

memory, and language processing [14, 29]. Empirically, reanalysis of a dataset originally

interpreted as showing sustained, load-dependent activity in the alpha and beta bands

during the delay period of a working memory task has revealed that, at the single-trial

level, this activity was driven by oscillatory bursts [30]. Despite these considerations,

however, it is difficult for us to conceive of how a relatively sparse, stochastic, and

bursty phenomenon could underlie a behavioral oscillation derived from a time series

that was assembled via the after-the-fact sorting of trials, from shortest to longest.

Nonetheless, another important question for Experiment 2 was to assess whether, at

the single-trial level, activity in the low-beta and higher-beta frequency bands more

closely resembled persistent oscillating signals or stochastic bursting.

Experiment 2

294   A new group of participants (*n*=16) carried out the same two experimental tasks as

295   Experiment 1 with a fixed post-cue-to-recall delay period while the EEG was recorded.

Behavioral performance

Recall accuracy. Overall levels of mean absolute error (MAE) of recall were

comparable to what was observed in Experiment 1: MAE did not differ statistically

between DSR *Recall 1* and NEU recall ($t(15)$ = -2.0024, *p* = .064), nor for DSR *Recall 2*

versus NEU recall ($t(15)$ = 2.3483, *p* = .033 (Bonferroni-corrected α = .0167); **Fig. 4A**).

DSR *Recall 1* was superior to DSR *Recall 2* ($t(15) = 4.7537$, $p < .001$), and 'stay' and

'switch' trials from DSR *Recall 2* did not differ ($t(15) = -1.5845$, $p = .134$; **S3 Fig**). A

linear mixed effects (LME) model confirmed that larger inter-item angular differences

were associated with larger absolute errors (main effect: $F(1,189) = 57.837$, $p < .001$;

Fig. 4B).

Fig 4. Experiment 2 recall performance. A. Mean absolute recall error for the two responses of the

DSR task and the single response of the NEU task. **B.** Same as **A.** separated by the current trial's inter-

item angular difference. **C.** Mean absolute error for DSR *Recall 1* and NEU recall as a function of which

sample item was tested. Error bars correspond to +/- 1 *SEM.* * symbols correspond to significant

differences between conditions at the Bonferroni-corrected alpha level = .0167 for 3 paired-sample *t*-tests

(**A.**) and the Bonferroni-corrected alpha level = .025 for two paired-sample *t*-tests and for interaction *p* <

.05 (**C.**).

Assessment of MAE as a function of the presentation order indicated that, for

NEU, MAE was numerically larger when *Sample 1* was tested ($t(15) = 1.204$, $p = .247$),

for DSR, MAE was numerically smaller when was tested ($t(15) = -1.593$, $p = .132$; **Fig.**

4C), and the crossover interaction between the two was significant ($t(15) = -2.276$, $p =$

.038).

Across- and within-trial bias. Linear mixed effects modeling indicated that

the previous trial's tested item exerted a significant attractive bias on the current trial

(NEU: $\beta = 1.60°$, $p < .001$; DSR: $\beta = 0.63°$, $p = 0.04$; **S5 Table**), an effect that was larger

in the NEU task ($\beta = 0.97°$, $p = .032$; **S6 Table**). The previous trial's untested item had

no significant effect on recall in either task. Within-trial effects were dependent on the

order of presentation. The recall of *Sample 1* was attracted toward the value of *Sample*

2 for NEU (β = 4.10°, *p* < .001), but not for DSR (β = -0.39°, *p* = .364), whereas recall of

Sample 2 showed no bias from *Sample 1* in NEU (β = -0.72°, *p* = .131) but a significant

repulsive bias in DSR (β = -1.52°, *p* < .001; **S5 Table**). The three-way interaction

between task, current untested item, and order-of-presentation was significant (β = -

3.67°, *p* < .001; **S6 Table**).

Linear mixed effects modeling further revealed that in the NEU task, the effect of

inter-item angular difference on bias was marginal ($F(2,2828)$ = 2.57, *p* = .077; **S7**

Table), but inter-item angular difference significantly interacted with order of

presentation in recall bias ($F(2,2828)$ = 3.94, *p* = .019). Recall of *Sample 1* was

significantly attracted toward *Sample 2* at all angular separations (22.5°: 4.56°, *p* < .001;

45°: 3.21° *p* < .001; 67.5°: 2.30° *p* = .002) with a significant reduction at the largest

angular separation (β = -2.26°, *p* = .025). Recall of *Sample 2,* in contrast, exhibited

significant repulsion from *Sample 1* at the small angular separation (22.5°: -1.72° *p* =

.021) but no significant bias at medium and large angular separations (45°: 0.01° *p* =

.986; 67.5°: -0.20° *p* = .809; main effect of presentation order: $F(1,2828)$ = 37.99, *p* <

.001).

In the DSR task, there was a significant main effect of presentation order

($F(1,2827)$ = 6.74, *p* = .009) and a significant interaction between presentation order

and inter-item angular difference ($F(2,2827)$ = 4.67, *p* = .009). Recall of *Sample 1*

showed no significant bias relative to *Sample 2* at any angular separation (22.5°: -0.01°

p = .992; 45°: -0.40° *p* = .535; 67.5°: -0.57° *p* = .387). Recall of *Sample 2,* in contrast,

was significantly repulsed from *Sample 1* at small and medium angular separations

(22.5°: -2.33° p < .001; 45°: -2.03° p = .001) but no bias at large separations (67.5°:

0.75° p = .220; interaction: β = 3.64°, p = .004).

The previous trial's target did not significantly influence signed bias in either task

(DSR: p = .288; NEU: p = .157). The task comparison model confirmed that NEU

showed significantly greater overall bias than DSR ($F(1,5657)$ = 24.49, p < .001), and

that the effect of presentation order on bias differed significantly between the tasks

(interaction: $F(1,5657)$ = 8.22, p = .004; **S8 Table**): in NEU, presenting the untested item

first eliminated attraction, whereas in DSR, it induced repulsion. The effect of inter-item

angular difference did not differ significantly between tasks (interaction: $F(2,5657)$ =

0.69, p = .502).

In most regards, these results replicated the pattern from Experiment 1 that

prioritization shielded the cued item from the influence of other stimuli. In particular, the

attractive bias toward the previous trial and the attractive bias of *Sample 2* on recall of

Sample 1 were markedly smaller in DSR than in NEU. The angular difference analysis

further revealed that, as in Experiment 1, the previous trial's target did not influence

signed bias, confirming that between-trial and within-trial sources of recall bias operate

independently. However, several other findings diverged from Experiment 1. First, recall

of *Sample 2* showed greater repulsion from *Sample 1* in DSR vs. NEU. Second, for

DSR, recall of *Sample 2* showed repulsion from *Sample 1* for angular separations of

22.5° and 45°. Third, for NEU, recall of *Sample 1* showed a uniform pattern of attraction

across angular separations and a significant repulsion of *Sample 2* from *Sample 1* for

angular separations of 22.5°. We will discuss these effects in Interim Discussion—

Experiment 2.

EEG spectral analysis

Oscillatory components. For the DSR task, visual inspection of the full distribution of extracted peak frequencies pooled across all participants and channels revealed a post-cue increase in the number of peaks in the low-beta range (**Fig. 5A**), an effect confirmed by K-S test (D = .1367, p < .001). A follow-up Mann–Whitney U test found that the difference identified by the K-S test was not due to a shift in central tendency (z = 1.13, p = .260), indicating that the effect reflected differences in distributional shape. This was confirmed by a permutation test on Wasserstein distance between the two pooled distributions, which was significant (mean distance = 1.22 Hz, permutation p < .001). To assess whether these pooled differences were consistent across participants, we computed the Wasserstein distance between each participant's peak distributions in the two epochs. Participants consistently exhibited significant non-zero within-subject distributional differences (mean distance = 2.59 Hz; permutation p < .001).

Fig 5. EEG spectral properties. A. Distribution of all oscillatory peak frequencies identified across channels and participants from *Delay 1.1* (light blue) and *Delay 1.2* (orange) for the DSR task (left) and for the NEU task (right). **B.** Group-averaged mean number of low beta peaks (indigo circles) and higher-beta peaks (green squares) per channel from *Delay 1.1* and *Delay 1.2* for the DSR task (left) and the NEU task (right). **C.** Left: Mean aperiodic offset for *Delay 1.1* and *Delay 1.2* in the DSR task (red circles) and in the NEU task (blue squares). Right: Mean aperiodic exponent for *Delay 1.1* and *Delay 1.2* in the DSR task (red circles) and in the NEU task (blue squares). **D.** Group-averaged mean low-beta burst rate (indigo circles) and higher-beta burst rate (green squares) per channel from *Delay 1.1* data and *Delay 1.2* data for the DSR task (left) and the NEU task (right). Error bars correspond to +/- 1 *SEM.*

Results for the NEU task differed from DSR: the cue-locked change of the full

distributions of peak frequencies was a shift toward a greater number of peaks in the

alpha frequency range (**Fig. 5A**), an effect confirmed by K-S test (D = .0623, p = .024),

with no change in median peak frequency (z = 0.1675, p = .867). At the group level,

participants consistently exhibited non-zero Wasserstein distances between epochs

(mean distance = 2.31 Hz; permutation p < .001), indicating that the distributions of

peaks in the NEU task also changed in a consistent manner across participants, despite

no shift in central tendency.

We next focused our analysis on detection of oscillatory peaks in the two sub-

bands identified in the analysis of behavioral oscillations of Experiment 1: 14-16 Hz

('low-beta') and 19-21 Hz ('higher-beta'). Visual inspection of the results suggested an

interaction between task and epoch in the number of low-beta vs. higher-beta peaks

extracted (**Fig. 5B**). A three-way repeated measures ANOVA did not find a significant

three-way Beta band × Task × Epoch interaction (F(1,15) = 0.8873, p = .361). However,

a more focused analysis revealed task-specific patterns. In the DSR task, there was a

significant crossover pattern whereby the number of low-beta and higher-beta peaks

changed in opposite directions across the delay epochs (t(15) = 2.3586, p = 0.032), with

higher-beta decreasing significantly from *Delay 1.1* to *Delay 1.2* (mean change = -0.242

peaks/channel, t(15) = -2.41, p = 0.029); an effect most prominent at frontal and

occipital electrodes (**S4 Fig**) and low-beta showing a numeric increase (mean change =

0.178 peaks/channel, t(15) = 2.02, p = 0.061); a trend most prominent at occipital and

parietal electrodes (**S5 Fig**). The NEU condition, in contrast, showed stable patterns in

these two sub-bands across the two epochs (all p-values > 0.6).

Aperiodic spectral component. For the aperiodic offset, the ANOVA

revealed a significant main effect of Epoch, ($F(1,15) = 11.07$, $p = .005$), with offset

increasing from *Delay 1.1* ($M = 1.19$) to *Delay 1.2* ($M = 1.32$). This indicated elevated

broadband power during the post-cue portion of the delay period. There was no main

effect of Task ($F(1,15) = 0.23$, $p = .64$) and no Task × Epoch interaction ($F(1,15) = 3.56$,

$p = .08$; **Fig. 5C**). For the aperiodic exponent, spectral slopes were numerically steeper

during *Delay 1.2* versus *Delay 1.1*, ($F(1,15) = 3.63$, $p = .076$), with no effect of Task

($F(1,15) = 0.58$, $p = .46$), and no interaction ($F(1,15) = 2.39$, $p = .14$). Non-parametric

tests collapsing across tasks confirmed significant epoch effects for both offset

(Wilcoxon signed-rank, $z = -2.79$, $p = .005$, $r = .70$) and exponent ($z = -2.22$, $p = .026$, r

$= .56$).

Burst analysis. Many analyses of intra- and extracranial recordings have

shown that beta-band effects observed in trial-averaged electrophysiological data can

be shown to be due to bursting—rather than to a sustained oscillating signal—at the

single-trial level. Because the EEG effects relating to the behavioral oscillations from

Experiment 1 (**Fig. 5**) emerged from a spectral parameterization routine that is carried

out on trial-averaged power spectra, it was important that we assess whether these

effects might also be due to bursting in the low-beta and higher-beta frequency bands.

We carried out our burst analyses with routines that had previously been applied

to EEG data from a working memory task to demonstrate that averaged power

dynamics in the traditionally defined beta band (15-29 Hz) could be accounted for by

burst dynamics observed at the single-trial level [30]. Thus, to validate our approach, we

first applied a burst analysis to the 4 sec epochs of *Delay 1.2* from the NEU task, from

the P3 electrode, bandpass filtered to 15-29 Hz ("broadband beta"). This yielded burst

frequencies of NEU: of 0.91 bursts/sec (SD = .13); DSR: 0.89 bursts/sec (SD = .13).

These values are similar to the 1.21 bursts/sec (SD = .22) obtained by [30] (B.C.

Kavanaugh, personal communication). In the narrower "low-beta" (14-16 Hz) and

"higher-beta" (19-21 Hz) bands, however, bursts were sparse events that occurred

infrequently at any individual electrode (3-7% of trials per channel), although when

considering all channels simultaneously, in low-beta, at least one burst occurred on

49.5-65.3% of trials (depending on delay epoch and condition) and in higher-beta, on

66.9-76.3% of trials, with similar prevalences in the two tasks. Despite the rare

occurrence of bursts in these two frequency bands, the burst rate in both showed a

significant increase from *Delay 1.1* to *Delay 1.2*, albeit with no differences between the

DSR and NEU tasks, with values for *Delay 1.1* ranging from 0.086-0.13 bursts/sec and

values for Delay 1.2 ranging from 0.139-0.176 events/sec (**Fig. 5D**). Separate repeated-

measures ANOVAs (Task × Epoch) for each frequency band confirmed significant main

effects of Epoch (low-beta: $F(1,15) = 19.72$, $p < 0.001$; higher-beta: $F(1,15) = 16.66$, $p <$

0.001), with no main effects of Task (both p-values > 0.38) and no Task × Epoch

interactions (both p-values > 0.29). Within-task paired t-tests confirmed significant

increases from *Delay 1.1* to *Delay 1.2* for both bands in both tasks (all $p \leq 0.004$).

Comparing burst rates across frequency bands, higher-beta bursts were more frequent

than low-beta bursts during *Delay 1.1* (both tasks: $t(15) > 5.37$, $p < 0.001$), but this

difference diminished during *Delay 1.2* (DSR: $t(15) = 2.25$, $p = 0.04$; NEU: $t(15) = 3.45$,

$p = 0.004$), reflecting a proportionally larger increase in low-beta burst activity in the

early delay period following the cues.

Power. To characterize the phase-locking properties of beta-band activity, we

decomposed raw EEG power in the low-beta and higher-beta bands into induced (non-

phase-locked) and evoked (phase-locked) components. Across both tasks and both

frequency bands, power was almost exclusively induced, with evoked power

constituting only ~1-2% of total power (**S6 Fig**).

Induced power did not mirror the task-specific *Delay 1.1* to *Delay 1.2* crossover

pattern observed in the peak extraction analysis (**Fig A of S6 Fig**). In low-beta, there

was a main effect of Task ($F(1,15) = 5.79$, $p = .030$), with NEU showing greater induced

power than DSR across both epochs, but no Task × Epoch interaction ($F(1,15) = 1.72$, p

$= .210$). In higher-beta, there was a main effect of Epoch ($F(1,15) = 9.38$, $p = .008$), with

induced power increasing from *Delay 1.1* to *Delay 1.2* in both tasks with no interaction

($F(1,15) = 1.48$, $p = .242$). The absence of interactions indicates that raw power

measures—which conflate periodic oscillations with aperiodic activity—do not capture

the subtle, task-specific shifts in oscillatory peak distributions revealed by the spectral

parameterization. The epoch effect in higher-beta power may reflect the task-general

increases in burst rate and aperiodic offset reported above.

Evoked power, which reflects phase-locked oscillatory activity, revealed a

different pattern (**Fig B of S6 Fig**). Both frequency bands showed significant main

effects of Task (low-beta: $F(1,15) = 7.83$, $p = .014$; higher-beta: $F(1,15) = 20.11$, $p <$

$.001$) and Epoch (low-beta: $F(1,15) = 9.38$, $p = .008$; higher-beta: $F(1,15) = 13.20$, $p =$

$.003$), with no interactions (both p-values $> .37$). In both bands, NEU evoked power was

greater than DSR, particularly during the post-cue epoch. This is consistent with a more

stereotyped, phase-consistent neural response to the uninformative cue in the NEU

task, and more variable power timing across trials in response to the retrocue in the

DSR task.

Finally, to determine whether task differences reflected differential cue-related

modulation rather than baseline differences, we computed percent power change from

pre-cue to post-cue using *Delay 1.1* as baseline. Induced power showed no task

differences in percent change (low-beta: $t(15) = -1.25$, $p = .23$; higher-beta: $t(15) = -$

1.22, $p = .24$; **Fig C of S6 Fig**), indicating that both tasks exhibited similar cue-related

increases in non-phase-locked power. By contrast, evoked power revealed a significant

task difference in low-beta ($t(15) = -3.46$, $p = .004$): DSR exhibited a larger decrease in

phase-locked power (-48.1 ± 8.2%) compared to NEU (-24.9 ± 7.4%). Higher-beta

showed a similar numerical pattern but did not reach significance (DSR: -35.0 ± 9.6%;

NEU: -19.7 ± 7.5%; $p = .17$). This pattern is consistent with a more disruptive effect of

the informative retrocue (DSR) compared to the uninformative cue (NEU) on cross-trial

phase consistency.

Interim summary—EEG spectral analysis

Filtering the EEG data into two narrow bands identified in Experiment 1—"low beta" (14-

16 Hz) and "higher-beta" (19-21 Hz)—produced results that are broadly congruent with

the behavioral oscillations observed in Experiment 1. Specifically, in the DSR task, the

prioritization cue triggered a shift toward more prominent oscillatory activity in the low-

beta band (together with a shift away from higher-beta). In the NEU task, in contrast,

there was only evidence of a shift toward more prominent activity in the alpha band.

Analyses of the aperiodic component of the data revealed effects that were common to

both tasks. For broadband power (i.e., the offset parameter), there was an increase

from *Delay 1.1* to *Delay 1.2*, consistent with a ramping of neural activity leading up to

the response. For spectral slope (i.e., the exponent parameter) there was evidence,

albeit weaker, for a steeper slope during *Delay 1.2*, which would be consistent with a

shift in E:I balance toward greater relative inhibition. Beta-burst rates, although in the

expected range within the classically defined beta band, were too sparse in the two

narrow bands of theoretical interest to be candidate substrates of the behavioral

oscillations observed in Experiment 1. Nonetheless, consistent with the oscillatory

dynamics, burst rates increased from pre-cue to post-cue periods in both the low-beta

(14-16 Hz) and higher-beta (19-21 Hz) bands, albeit with no task differences. Finally,

raw power measures did not capture the task-specific oscillatory signatures identified

through spectral parameterization, although evoked power analyses revealed that the

informative retrocue in DSR was more disruptive to cross-trial phase consistency than

the uninformative cue in NEU.

Representational similarity

Time-course analyses. In the DSR task, onset of *Cue 1* triggered an abrupt

shift in the neural representation of both items (**Fig. 6A**). For the PMI, its representation

during both the *Cue 1* and *Delay 1.2* epochs was significantly different from (i.e., less

correlated with) the reference RDM (i.e., with its representation during *Delay 1.1;* both

p-values $\leqq$ .0027). After the initial transformation, the RSA time course was stable, in

that the PMI's similarity with respect to the *Delay 1.1* reference RDM did not differ

between the *Cue 1* and *Delay 1.2* epochs ($t(14) = 1.7488$, $p = .102$). For the UMI, its

representation during the *Cue 1* and *Delay 1.2* epochs was also significantly different

from the *Delay 1.1* reference RDM (both *p*-values ≦ .001). Unlike the PMI, however, the

representation of the UMI continued to change after cue offset, such that its correlation

with the reference RDM was higher during the *Cue 1* epoch than during the *Delay 1.2*

epoch (*t*(14) = 3.4866, *p* = .004). When comparing the two memory items directly, the

representational shift of the UMI were more pronounced than those of the PMI: during

the *Cue 1* epoch the UMI's representational dissimilarity relative to the *Delay 1.1*

reference RDM was greater than that for the PMI (*t*(14) = 2.1889, *p* = .046). This

difference persisted through *Delay 1.2* (*t*(14) = 2.3878, *p* = .032).

Fig 6. Time-resolved representational similarity. A. Time-resolved similarity traces for the correlation

of representation of the PMI (orange) and UMI (purple) at each time point spanning *Delay 1.1* through

*Delay 1.*2 with the representation of each item over the duration of *Delay 1.1.* **B.** Same as **A.** for the

tested (light orange) and untested (light purple) item representations of the NEU task. **C.** Topographic

importance maps depicting the relative contributions of each electrode's activity to the RSA traces in

panels **A.** and **B.** for the DSR task (right) and NEU task (left), during *Delay 1.1* (top row), cue epochs

(middle row), and *Delay 1.2* (bottom row). Warmer colors indicate stronger contribution to the overall

similarity estimates.

For the NEU task, variability of the RSA time course was visibly higher than for

the DSR task. Although both items exhibited changes in representational similarity

across the epochs (all *p*-values < .005), these time courses did not exhibit the abrupt

cue-locked transformations observed in the DSR task (**Fig. 6B**). Furthermore, the

degree of dissimilarity with respect to the reference RDM did not differ between the two

items in any of the epochs that were analyzed (all *p*-values > .596).

Inspection of the topographic importance maps for the RSA reveals similar

patterns during *Delay 1.1* for both memory items in both tasks, with prominent

contributions from a set of occipital-parietal electrodes and a separate set of frontal-

midline electrodes (**Fig. 6C**). These distributions changed only subtly during the cue

epoch, then more markedly during *Delay 1.2.* For the DSR task, for the PMI only, a

reversal of sign was observed at frontal midline electrodes for *Delay 1.2* versus earlier

epochs.

565   ***Spectral analyses—beta-band peak extraction at the task level.***

To assess whether the oscillatory dynamics of neural representations may also show

similarities with the behavioral oscillations revealed by Experiment 1, we next quantified

oscillatory patterns in the time-resolved RSA data. To do this, we first extracted beta-

frequency peaks from the PMI/tested and UMI/untested RSA time courses, separately in

the two narrow bands identified in behavioral oscillations (**Fig. 3**): 'low-beta' (14-16 Hz)

and 'higher-beta' (19-21 Hz). Next, to enable a direct comparison with the task-level

EEG spectral analysis (**Fig. 5**), we combined the results across the two items in each

task. Results for the DSR task revealed that the number of low-beta peaks and higher-

beta peaks increased from *Delay 1.1* to *Delay 1.2*, whereas for the NEU task the

number of low-beta peaks and higher-beta peaks decreased from *Delay 1.1* to *Delay*

1.2 (**Fig. 7A;** see **S1 File** for aperiodic component analyses). A three-way repeated

measures ANOVA with factors of Task (DSR vs. NEU), Frequency Band (low-beta vs.

higher-beta), and Epoch (*Delay 1.1* vs. *Delay 1.2*) revealed a significant Task x Epoch

interaction, ($F(1,14) = 17.441$, $p < .001$), indicating that this divergent pattern of

oscillatory peak counts between the tasks was significant. To clarify the bases of these

task-level differences, we conducted follow-up paired *t*-tests comparing the peak counts

for the DSR and NEU tasks within each epoch and band. These tests revealed

significantly more low-beta peaks in the time-resolved RSA of the DSR task than of the

NEU task during *Delay 1.2* (t(14) = 2.779, p = 0.015, d = 0.718), a pattern consistent

with the oscillatory patterns in the behavior. Moreover, the DSR and NEU tasks showed

divergent low-beta peak *shifts* from *Delay 1.1* to *Delay 1.2* (t(14) = 2.703, p = .0172, d =

.698), driven by the significant reduction in low-beta peaks across the epochs in the

NEU task (ΔM = -0.60, t(14) = -2.81, p = .014). Unexpectedly, these tests also revealed

significantly more higher-beta peaks in DSR representational similarity, relative to NEU,

during *Delay 1.2* (t(14) = 2.432, p = .029, $d.$ = .628). Although the higher-beta peak

shifts from *Delay 1.1* to *Delay 1.2* diverged across tasks at a level that approached

significance (t(14) = 2.073, p = .0572, d = .535), neither task exhibited a significant

change in the number of higher-beta peaks across epochs (both p-values $\geq$.096)

suggesting that the temporal evolution of task-related frequency modulation in

representational similarity was specific to the low-beta band.

Fig 7. RSA oscillatory characteristics. A. Group-averaged mean number of low-beta peaks (indigo

circles) and higher-beta peaks (green squares) per participant from *Delay 1.1* and *Delay 1.2* for the DSR

task (left) and the NEU task (right). **B.** Group-averaged mean number of low-beta peaks (upper row) and

higher-beta peaks (lower row) per participant from *Delay 1.1* data and *Delay 1.2* data for the DSR task

(left column) and the NEU task (right column), split by memory item with PMI/tested item results in purple

and UMI/untested item results in orange. **C.** Group-level phase consistency in *Delay 1.1* (pre-cue) and

Delay 1.2 (post-cue) periods for the DSR task (blue) and NEU task (red) in the low-beta band (14-16 Hz;

left) and the higher-beta band (19-21 Hz; right). * permutation p < .05; ** permutation p < .01. Error bars

correspond to +/- 1 *SEM* (**A.** & **B.**) and to bootstrapped 95% confidence intervals (**C.**).

Spectral analyses—beta-band peak extraction at the item level. To

determine whether the frequency-specific task effects illustrated in **Figure 7A** reflected

modulation of PMI representations, UMI representations, or both, we conducted

separate analyses for each item type. We extracted peak counts independently for PMI,

UMI, tested item, and untested item time courses and computed descriptive statistics

(**Fig. 7B;** see **S2 File** for item-specific aperiodic component analyses). In the DSR task,

PMI beta peak counts remained relatively stable across epochs for both low-beta (*Delay*

1.1: $M = .27$; *Delay 1.2: M = .20$) and higher-beta (*Delay 1.1*: $M = .2$; *Delay 1.2: M =*

.27). UMI beta peak counts, in contrast, increased from $M = 0.13$ to $M = 0.40$ peaks in

low-beta and from $M = .07$ to $M = .33$ peaks in higher-beta.

In the NEU task, beta peak counts declined for both the tested item and the

untested item: tested (low-beta from *Delay 1.1* ($M = .47$) to *Delay 1.2* ($M = .07$), and for

higher-beta from *Delay 1.1* ($M = .2$) to *Delay 1.2* ($M = 0$); untested (low-beta from *Delay*

1.1 ($M = .2$) to *Delay 1.2* ($M = 0.00$) and higher-beta *Delay 1.1* ($M = .2$) to *Delay 1.2* (M

$= .133$).

Spectral analyses—phase relations between items. To determine

whether the phase relationship between the representations of the two memory items

differed as a function of task, we quantified the phase consistency (r) in each task and

frequency band, which indicated the degree to which the cross-spectral phase angle

between the two RSA time courses was consistent across participants. In the low-beta

band (14–16 Hz), no significant Task × Epoch interaction was observed (permutation p

$= 0.58$; **Fig. 7C**), and no individual condition showed significant phase consistency (all

Rayleigh $p > 0.35$). In the higher-beta band (19–21 Hz), there was a significant Task ×

Epoch interaction (permutation p = 0.0079). Prior to the cue, there was significant phase

consistency between items in the DSR task (r = 0.481, Rayleigh z = 3.47, p = 0.028),

but not in the NEU task (r = 0.122, p = 0.81). Following the cue, this pattern reversed:

phase consistency in DSR decreased (r = 0.181, p = 0.62) and phase consistency in

NEU exhibited a marginal increase (r = 0.417, p = 0.072).

Interim discussion—Experiment 2

Behavioral results broadly replicated the evidence from Experiment 1 that

prioritization shields an item from the biasing influence of other information recently or

currently in working memory. One possible exception to this was the fact that, unique to

Experiment 2, recall of *Sample 2* showed greater repulsion from *Sample 1* in DSR than

in NEU. However, we speculate that this, together with other specific differences

between the two experiments, may reflect an interaction of prioritization with the longer

delay period of Experiment 2. For example, previous work has suggested that longer

delays [40], as well as "active, not passive, maintenance" [41] are associated with

greater separation between items.

Spectral parameterization of the EEG in the DSR task revealed a cue-locked

increase of oscillatory peaks in the low-beta band, together with a decrease in

oscillatory peaks in the higher-beta band, suggesting an oscillatory basis for the priority-

specific behavioral oscillations from Experiment 1. Burst analyses indicated that these

effects in these narrow frequency bands were unlikely to be due to changes of burst

dynamics at the single trial level.

Results from time-resolved RSA analyses provided two important insights. First,

in the DSR task, priority-related representational changes were more pronounced for

the uncued item (i.e., the UMI) than for the cued item (the PMI). This suggests that at

least some of the "shielding" effects of prioritization may result from representational

transformations of the item that is *not* needed for the impending action. Second,

although these analyses showed cue-related effects on phase dynamics, they failed to

provide evidence that a strengthening of phase encoding may be a basis for priority-

related shielding.

661 Discussion

The results from these two experiments provide evidence for a model in which the

prioritization of one among two items held in working memory prompts representational

transformations that are larger for the uncued item (i.e., the UMI) than for the cued item

(the PMI), and in which the encoding of UMI status is implemented, at least in part, by

dynamics in the beta band. Previous studies of the DSR task combining EEG and

transcranial magnetic stimulation (TMS) have shown that decoding of the UMI depends

on the beta band [8, 10], and this effect is associated with cue- and TMS-related

dynamics in the low beta band (i.e., 13-20 Hz; [10]). Furthermore, single pulses of TMS

(spTMS) delivered during *Delay 1.2* can produce behavioral evidence for the involuntary

retrieval of the UMI (indexed by an inflated false-alarm rate; [8, 42]), and the magnitude

of this behavioral effect is predicted by the magnitude of a spTMS-related decrease in

beta-band components derived from a decomposition of the EEG [9]. In the present

study, we have produced evidence that prioritization is associated with a behavioral

oscillation in the low-beta band and by an increase of oscillatory peaks in this same low-

beta band. Thus, our working model is that UMI status is encoded, and sustained, by a

process that cycles at a frequency in the low-beta band. Although this account lacks an

explicit mechanism, the RSA results from Experiment 2 do argue against the likelihood

that the mechanism might be the encoding of the two memory items at different phases

of a low-beta oscillation.

Although we do not yet understand the precise mechanism that implements the

state of deprioritization, we speculate that its dependence on oscillations at a low-beta

frequency links it to the natural frequency of thalamocortical circuits at posterior nodes

of the frontoparietal priority map. This follows from the results of several studies that

have recorded the EEG while spTMS is delivered to posterior parietal cortex (PPC):

spTMS of PPC produces a spectral perturbation that is strongest in β1 (13-20 Hz; in

comparison to spTMS of occipital cortex--alpha: 8-12 Hz--and spTMS of premotor

cortex--β2: 21-29 Hz; [43]); when measured as the global mean field amplitude (GMFA),

the amplitude of the response evoked by spTMS of PPC varies with pre-spTMS phase

in the beta band [44]; and spTMS-evoked phosphene perception depends on pre-

spTMS power in the beta band for spTMS of PPC (in comparison to its dependence on

pre-spTMS power in the alpha band for spTMS of occipital cortex; [45]). Assuming that

priority in working memory is encoded in the frontoparietal priority map, it may be that

the top-down signals that impose this priority status on stimulus representations in early

visual areas oscillate at the natural frequency of the PPC.

The behavioral and EEG data presented here point to an oscillatory mechanism

supporting prioritization: recall oscillates at 15 Hz, and the prioritization-specific increase

of oscillatory peaks in this narrow frequency band cannot be accounted for by the

sparse bursting rate that we observed. This suggests that prioritization is supported by a

mechanism that is fundamentally different from the beta bursts, also observed at

posterior electrodes, that implement inhibitory cognitive control (e.g., [14, 29]).

Importantly, however, our claim does not extend to the processes that underlie the more

general function of the maintenance of information in working memory. Indeed, when we

applied a burst analysis to the broader band of traditionally defined beta (15-29 Hz), we

observed a burst rate comparable to one that has been shown to underlie load-related

dynamics in the EEG [30].

Inspection of the time-resolved RSA suggests that both memory items undergo

their priority-based representational transformation almost instantaneously upon

presentation of the cue. This would be consistent with the possibility that the

transformations of the PMI and UMI are triggered by a common factor, as would occur

with the change of a control variable in a dynamical system. Also consistent with this

possibility is the fact that trial-by-trial variation in the efficacy of the geometric

transformation of the PMI and of the UMI are highly correlated, as measured with fMRI

and with EEG [46]. The time-resolved RSA indicates that the priority-related

transformation of the UMI is larger than that of the PMI, a finding consistent with

previous fMRI studies (e.g., [3, 47]). We speculate that this may reflect a rotation of the

UMI into an action-null subspace, whereas the PMI occupies an action-potent subspace

that is only minimally different from the representational state of both memory items

from portions of the trial when both are equally likely to be needed for the recall

response.

Materials & methods

Experiment 1

Participants

Eighteen neurologically healthy members of the University of Wisconsin–Madison

community participated in the behavioral experiment. Incomplete data sets from two

participants were excluded yielding a final sample of 16 data sets used in the analyses

(from 10 females, 2 unknown, 18-32 years, M = 22.8 years). The sample size was

chosen to match the sample size of [16] which reported the inter-item bias effects we

aimed to replicate and understand here. All participants had normal or corrected-to-

normal vision, and all reported having normal color vision. The research was approved

by the University of Wisconsin—Madison Health Sciences Institutional Review Board.

All participants gave written informed consent at the start of the session and received

monetary compensation in exchange for participation.

Experimental procedure

Double serial-retrocue task. The double serial-retrocue (DSR; **Fig. 1A**)

task began with the 200 ms presentation of an oriented grating at the center of the

display ("*Sample 1*"), followed by a 500 ms inter-stimulus fixation interval ("*ISI*"),

followed by the 200 ms presentation of a second oriented grating at the center of the

display ("*Sample 2*"), followed by 500 ms of fixation ("*Delay 1.1*"), followed by a 500 ms

presentation of *Cue 1* (the digit '1' or '2') whose value indicated with 100% validity which

of the two samples would be tested. This cue conferred on the cued item the status of

the prioritized memory item (PMI) and, by default, the uncued item took on the status of

unprioritized memory item (UMI). *Cue 1* was followed by a fixation period ("*Delay 1.2*")

of variable length, densely- and uniformly-sampled across a range from 500-1200 ms in

steps of 16.67 ms, implemented in the code as number of screen refreshes. *Delay 1.2*

was followed by *Recall 1* in which a circle with a line corresponding to its diameter (and

described to participants as a "dial") was presented. Using the mouse, participants

rotated the line to match the orientation of the cued item and clicked the mouse button

to lock in their response within a 3-sec window beginning with probe onset. Feedback

was displayed for the 1 second interval following the 3 second recall interval, the fixation

cross turning green for responses less than 20 deg from the sample orientation, yellow

for responses between 20 and 28 deg from the sample orientation, and red for

responses greater than 28 deg from the sample orientation. If no response was

recorded, the words "No response was recorded" appeared during the feedback

interval. The trial then continued with a 500 ms presentation of *Cue 2* (the digit '1' or '2')

whose value indicated with 100% validity which of the two samples would be tested

during *Recall 2*. The item designated by *Cue 2* took on the status PMI, and the uncued

item the status of irrelevant memory item (IMI; because it could no longer be cued on

that trial). *Recall 2* and feedback periods were identical to *Recall 1* and its feedback

period. After feedback to *Recall 2*, text appeared at the center of the display telling the

participant to press the spacebar key when ready to continue with the next trial. This

triggered the inter-trial interval, which varied between 2-3 s ("*ITI*"). Upon completion of

the block, text appeared on the display informing the participant of their average recall

error in degrees on that block. Over the course of each block, *Cue 2* indicated the same

item as had *Cue 1* on 50% of trials ("*stay*" trials; the remaining 50% were "*switch*" trials).

Neutral-cue task. The neutral-cue (NEU; **Fig. 1B**) task was identical to the

DSR task with two exceptions: (i) trials featured only one *Cue* and one *Recall* and

Feedback period; and (ii) the *Cue* was uninformative, appearing as '0' on each trial.

Throughout the text, we will refer to the items in the NEU task not as PMI and IMI, but

rather as "tested" and "untested."

Experimental stimuli and design

Participants were seated in a dimly lit room at a viewing distance of 50 cm from the

monitor. Sample stimuli were centrally presented black and white gratings appearing on

a mid-gray background with radii of 6°, spatial frequencies equal to 1 cycle/°, 50%

contrast, and random phase. The orientation of one of the sample items was selected

from a set of six base orientations (10, 40, 70, 100, 130, 160°), with a random jitter of

−3° to 3°. The orientation of the other sample item was selected to be offset from the

jittered base orientation from a set of inter-item angular differences (0, +/-22.5, +/-45, +/-

67.5°). The presentation order of the base orientation and offset orientation was

randomized and counter-balanced across trials. The centrally presented cues were

black digits, and the recall stimuli were black response dials (unfilled black circles with a

black line corresponding to the diameter of the circle) with the same radius and random

starting orientation. A fixation cross was presented during blank task intervals. The

cross was gray during the fixation and *ISI* intervals, white during the memory delay

periods, and green, yellow, or red during the feedback interval, depending on the

response error.

The experiment was completed in a single session, and all participants

completed four blocks of the DSR task followed by four blocks of the NEU task. (We

reasoned that participants were less likely to ignore DSR cues if they had not previously

performed a variant of the task with uninformative cues.) Prior to each set of four blocks,

participants completed 8-12 practice trials of the task in the presence of the

experimenter. During practice trials, the feedback display differed. After the participant

locked in their recall response, the recall stimulus remained on the display and changed

color according to the green/yellow/red feedback convention, and the original sample

grating reappeared under the recall stimulus to provide visual error feedback to the

participant during these trials.

Each block was comprised of 84 trials for a total of 336 trials per task, and

participants were encouraged to take breaks between blocks (and between trials if

necessary) before continuing. Any trial for which a response was not recorded within the

3-second window was repeated at the end of the block. Because we were primarily

interested in behavioral oscillations (and hence the manipulation of the duration of

Delay 1.2), we established the experimental design to ensure an equal number of trials

collected for each of 42 durations of *Delay 1.2*. This resulted in 8 trials collected at each

duration, of which there were 2 trials per inter-item angular difference (magnitude)

collected per task, per participant. Across those, the base orientations were randomly

sampled such that each base orientation was selected for an equal number of trials

across all possible base orientation values, but they were not equated across *Delay 1.2*

durations.

Behavioral analyses

Mean absolute error of recall. For both DSR and NEU tasks, we

quantified overall accuracy of recall by computing the average magnitude of the angular

difference between the participant's recall response value and the orientation of the

tested sample item, as a function of inter-item angular difference and, for the second

response of the DSR task, as a function of 'stay'/'switch' identity of *Cue 2*. We also

tested for differences in absolute recall error as a function of the target's presentation

order (i.e., as *Sample 1* or *Sample 2*). Paired-sample *t*-tests were used to test for

significant differences in mean absolute error of recall across task responses with

Bonferroni correction for the number of tests. A linear mixed effects model was used to

test for a significant effect of task (2 levels: DSR/NEU, coded as a categorical variable)

and inter-item angular difference magnitude (4 levels: 0, 22.5, 45, 67.5 deg). In the

model that is reported here, the task x inter-item angular difference interaction term was

omitted because this interaction was found to be non-significant and inclusion of the

term did not significantly improve the model fit. Subject was included as a random

effect, and a separate intercept was fitted for each.

Inter-item bias. Following [16], we quantified the bias exerted by the untested

item on the recall of the tested item as the mean signed recall error, with negative

signed errors indicating attraction and positive signed errors indicating repulsion. For

each trial, we computed the recall error, and if the response value was in the direction of

the untested item, the recall error was given a negative sign, and if the response value

was in the direction away from the untested item, the recall error was given a positive

sign. As with mean absolute error, we quantified the inter-item bias for both responses

during the DSR task and the single response during the NEU task, as well as separately

for the second response of the 'stay' and 'switch' trials of the DSR task, and as a

function of inter-sample item angular difference. Paired-sample t-tests were used to test

for significant differences in inter-item bias across task responses with Bonferroni

correction for the number of tests. A linear mixed effects model was used to test for a

significant effect of task (2 levels: DSR/NEU, coded as a categorical variable) and inter-

item angular difference magnitude (4 levels: 0, 22.5, 45, 67.5 deg) and their interaction

with subject included as a random effect and a separate intercept fitted to each.

Serial dependence. In addition to within-trial inter-item effects on recall, we

also tested for between-trial inter-item effects on recall. For each participant's data, the

response error on each trial was computed as the angular distance between the

presented and reported sample orientation. Following previous serial dependence work

[17-20], high-error trials (≥25 deg; M = 5.4%) were omitted from analysis. To remove

any global directional biases of individual participants, we subtracted their mean signed

error across trials from each trial. Serial dependence was quantified by sorting each

participant's demeaned errors according to the angular difference between the sample

direction on the current and previous trials. Here, we quantified serial bias exerted by

the previous trial's most recent PMI/tested item (i.e., with respect to the PMI designated

by *Cue 2* in DSR trials), as well as the bias exerted by the previous trial's IMI/untested

item. In these analyses, the first trial of each block was omitted. For instances in which

trials were removed due to large errors, the most immediately preceding non-large error

trial served as the previous trial. The sorted errors for each participant were then

smoothed by a 20°-wide median filter in 2° increments. (Similar results were obtained

with different-sized filters.)

We averaged the smoothed data within each condition and fit the averages of

each with a von Mises–based function of the following form:

$$y = a \, \sin(2x) \, e^{k\cos(2x)} \qquad\qquad (1)$$

where x is the relative orientation of the tested/untested item of the previous trial (in

radians), a is the amplitude of the curve, and k is the concentration parameter

determining the width of the tuning. The sine and cosine terms capture the circular

periodicity of orientation differences over 180°, allowing the function to model both

attractive and repulsive serial biases symmetrically around zero (analogous to a

derivative-of-von-Mises (dVM) formulation used in previous studies; e.g., [18]). Fitting

was achieved by minimizing the sum of squared errors using the *lsqcurvefit* MATLAB

routine, with amplitude a and concentration k parameters free to vary between –10° and

10° and between 0 and 10, respectively.

Statistical significance of the group-level dVM fits was assessed through a bias-

corrected and accelerated (BCa) bootstrapping procedure. On each of 100,000

iterations, we sampled participants' data with replacement and fit a dVM to the average

of the bootstrapped sampled data. The value of the amplitude parameter was retained

after each iteration, resulting in a distribution of the amplitude parameter of our sample.

We also computed the jackknife distribution of amplitude parameters by computing all of

the leave-one-sample-out dVM fits. The confidence interval and *p*-value of the

amplitudes of each group fit when compared to zero (i.e., no serial dependence) were

computed using the BCa_bootstrap MATLAB function [21], which utilized both the

bootstrapped and jackknife amplitude distributions.

Between-condition comparisons of dVM fit amplitudes were carried out using

permutation testing. On each of 100,000 iterations, we shuffled the group labels (e.g.,

'DSR' / 'NEU'; 'PMI' / 'IMI') and then assigned each dataset with one of the shuffled

labels. We then grouped the data by the shuffled labels and fit a dVM to the average of

each shuffled group. The value of the amplitude parameter was retained after each

iteration, resulting in a permutation distribution of the amplitude parameter. The *p*-value

of the original difference in amplitudes between the two groups was derived from the

permutation distribution as the proportion of the permutations greater than the original

difference.

Linear mixed effects modeling of inter-item effects. After the data

were collected, the separate analyses of inter-item influences on recall, as described in

the previous section, revealed interference from both current-trial and previous-trial

memory items (see **S3 file** for more detail). Because of this, we fit the data from each

task to a linear mixed effects model to simultaneously model the inter-item biases of

current and previous trial sample items on current trial recall. The dependent variable

was the signed recall error on the current trial, modeled as a function of (i.) the

orientations of the previous trial's tested item, (ii.) the previous trial's untested item, and

(iii.) the current trial's untested item. Specifically, the predictors were derived from the

difference between each item's orientation (in radians) relative to the current trial's

target orientation, computed using the *circ_dist.m* function to ensure proper angular

wrapping, and transformed using the sine of twice the angle, $\sin(2\theta)$, to capture

orientation-dependent modulations in a directionally signed space. An additional

categorical predictor ('NonTargetPresentation') coded whether the current trial's

untested item was presented before (1) or after (0) the target item, to assess whether

temporal order of presentation modulated inter-item bias. The model took the following

form:

Recall Error ~ $\sin(2^*\theta_{\text{prev\_tested}}) + \sin(2^*\theta_{\text{prev\_untested}})$

906              $+ \sin(2^*\theta_{\text{curr\_untested}}) * \text{NonTargetPresentation} + (1 \mid \text{Subject})$ (2)

Subject was included as a random effect, with random intercepts to account for inter-

individual variability. This model was selected based on nested model comparisons in

fitting the NEU data and then used to separately model both the NEU and DSR task

Recall 1 data.

A third model was fit to directly compare the within-trial biases of the two tasks

with the form:

Recall Error ~ Task* $(\sin(2^*\theta_{\text{prev\_tested}}) + \sin(2^*\theta_{\text{prev\_untested}})$

914              $+ \sin(2^*\theta_{\text{curr\_untested}}) * \text{NonTargetPresentation}) + (1 \mid \text{Subject})$ (3)

where Task was included as a categorical variable, with the DSR task dummy-coded to

be the reference task. This model included the three-way interaction between Task, the

current trial's untested item, and the presentation order of the current trial's untested

item to test whether the effect of prioritization on inter-item bias depended on temporal

order of presentation.

After model fitting, we extracted the fixed effects estimates and their

associated 95% confidence intervals. Standard errors were computed from the

confidence interval widths, and *t*-statistics were obtained as the ratio of each fixed-effect

estimate to its standard error. Corresponding two-tailed *p*-values were then derived from

the *t* distribution using the model's residual degrees of freedom. Importantly, positive

coefficients for the sine-transformed predictors would indicate attractive biases toward

them, whereas negative coefficients would indicate repulsive biases away from them.

To directly test whether the magnitude of inter-item angular difference modulated

bias, we conducted a supplementary analysis using a signed bias measure as the

dependent variable. This measure was computed by multiplying the signed recall error

by the sign of the angular difference between the current trial's untested and target

items, such that positive values indicate attraction toward the untested item and

negative values indicate repulsion away from it, maintaining consistency with the

primary LME analyses above. The 25% of trials with 0° angular difference were

excluded from this analysis because signed bias is undefined when the two items are

identical.

The signed bias was modeled as a function of (i.) the angular difference between

the current trial's items (22.5°, 45°, or 67.5°; treated as a categorical variable), (ii.) the

presentation order of the current trial's untested item, and (iii.) their interaction, while

controlling for previous trial influences using the sine-transformed orientation differences

of the previous trial's tested and untested items. The model took the following form:

Signed Bias ~ AngDiff * NonTargetPresentation + $\sin(2*\theta_{\text{prev\_tested}})$ +

942                        $\sin(2*\theta_{\text{prev\_untested}})$ + (1 | Subject)   (4)

943   Subject was included as a random effect with random intercepts to account for inter-

944   individual variability. This model was fit separately to the NEU and DSR task data.

945   A third model was fit to directly compare the effects of angular difference across tasks

946   with the form:

947   Signed Bias ~ Task * (AngDiff * NonTargetPresentation) +

948                    $\sin(2*\theta_{\text{prev\_tested}}) + \sin(2*\theta_{\text{prev\_untested}})$ + (1 | Subject)   (5)

where Task was included as a categorical variable, with the DSR task dummy-coded as

the reference level. This model tested whether the effect of angular difference on inter-

item bias, and its interaction with presentation order, differed between tasks.

Behavioral oscillations. Behavioral oscillations in recall were assessed by

analyzing mean absolute error (MAE) as a function of *Delay 1.2* duration (0.5–1.2 s, 42

discrete intervals). For each participant and duration, MAE values were averaged

across trials and smoothed with a two-point moving average. The resulting delay-

dependent MAE series was linearly detrended to remove slow drifts. To detect rhythmic

fluctuations in performance, a Fast Fourier Transform (FFT) was applied to each

participant's detrended MAE time series. Prior to transformation, data were multiplied by

a Hann window to reduce spectral leakage and zero-padding to the nearest power of

two was applied for computational efficiency and to interpolate the frequency spectrum.

The FFT yielded complex coefficients representing amplitude and phase values across

33 frequency bins spanning 0–30 Hz. Group-level spectra were obtained by averaging

participants' complex Fourier coefficients (real and imaginary components), preserving

the relative phase information across participants. This approach ensures that

consistent phase alignment across participants contributes to the group-level amplitude,

rather than being lost through power-only averaging. To assess statistical significance,

Delay 1.2 trial labels were randomly shuffled for each participant, after which delay-

dependent MAE was recomputed and the FFT was repeated. This procedure was

performed 10,000 times to generate participant-wise null spectra and corresponding

group-level amplitude distributions. Observed spectral peaks exceeding the 95th

percentile of the null distribution—corrected for multiple comparisons (Bonferroni

correction was applied across identified peaks)—were considered significant and would

indicate rhythmic modulation of behavioral performance at the frequencies of the

significant peaks.

For frequencies showing significant oscillatory peaks, between-participant phase

alignment was quantified using the Phase-Locking Value (PLV). To compute this, each

participant's phase angle at their peak frequency was extracted from the complex FFT

output, and PLV was computed as the absolute value of the mean complex phase

vector:

$$PLV = \frac{1}{N}\sum_{n=1}^{N} e^{i\theta_n} \qquad (6)$$

where N is the number of participants, θ_n is the phase angle for participant n, and

$i = \sqrt{-1}$. PLV values range from 0 (uniform phase distribution) to 1 (perfect phase

alignment). Statistical significance of phase alignment was assessed using both a

parametric Rayleigh test for non-uniformity and a non-parametric permutation test. For

the latter, PLV values were computed for each of the 10,000 permutation-based null

datasets, and the observed PLV was compared to the resulting null distribution

(α=0.05).

All analyses were conducted separately for the DSR and NEU tasks, allowing for

direct comparison of peak oscillation frequencies between tasks. Identical procedures

were repeated to assess oscillations in within-trial inter-item bias on recall.

Experiment 2

Participants

A separate sample of 16 neurologically healthy members of the University of Wisconsin–Madison community (10 females, 21-28 years, M = 23.7 years) participated in the EEG experiment, and all data were used in the final analyses. The sample size was chosen to match the sample size of Experiment 1. All participants had normal or corrected-to-normal vision, and all reported having normal color vision. The research was approved by the University of Wisconsin—Madison Health Sciences Institutional Review Board. All participants gave written informed consent at the start of the session and received monetary compensation in exchange for participation.

Experimental stimuli, procedure, and design

The experimental stimuli, procedure, and design were the same as those of Experiment 1, other than the following differences: (i.) *Delay 1.2* of both tasks had a fixed length of 4 s. (ii.) Participants completed four blocks of the DSR task followed by two blocks of the NEU task, with each block consisting of 60 trials. (iii.) A full-factorial design was ensured whereby each inter-item angular difference magnitude was selected for 30 trials, with each base orientation being selected for 5 of those 30 trials yielding 120 total trials of the NEU task. For the DSR task, we doubled the trials to accommodate the 'stay'/'switch' aspect of the task, yielding 240 total DSR task trials.

EEG recording and preprocessing

EEG was recorded from 63 electrodes with a BrainVision Recorder using active Ag/AgCl electrodes feeding into an ActiCHamp amplifier (BrainProdcuts GmbH). The electrodes were positioned according to the extended International 10–20 system.

Impedances were kept below 25 kΩ, and the data were sampled at 1000 Hz. An

electrode placed on the forehead served as ground, and all channels were referenced

online to channel Fpz and re-referenced offline to the average of all electrodes.

Data were processed offline using EEGLAB [31] with the FieldTrip toolbox [32],

and custom MATLAB scripts. The data were downsampled to 500 Hz and were

bandpass filtered between 1 and 100 Hz with a notch filter centered at 60 Hz.

Electrodes exhibiting excessive noise were removed and re-referenced to the average

of all remaining electrodes. The continuous data were then epoched to 500 ms prior to

Sample 1 through the feedback in the NEU task, and through the second feedback in

the DSR task. Automatic artifact epoch detection and rejection were then performed

using the EEGLAB *pop_autorej.m* procedure. The data then underwent dimensionality

reduction to 32 principal components, upon which an independent component analysis

was applied to identify and remove components containing eye movement artifacts and

muscle artifacts using ICLabel [33]. Finally, electrodes with excessive noise were

interpolated using the spherical spline method.

Behavioral analyses

We carried out analyses of the mean absolute error of recall and of inter-item bias

according to the procedures described in the corresponding sections of the Methods of

Experiment 1. Mirroring the reported results of Experiment 1, mean absolute error and

grand linear mixed effects modeling of inter-item biases are reported below; see **S3 File**

for separate analyses of inter-item bias due to current and previous trial memory items.

EEG spectral analysis

Broadband oscillatory peak extraction. For both tasks, we isolated

oscillatory peaks from *Delay 1.1* and *Delay 1.2* (i.e., the epoch preceding the cue and

following the cue), excluding the first 100 ms of both to minimize visual stimulus-related

effects. Because *Delay 1.1* was 500 ms in length, this yielded a window of 400 ms from

which to extract oscillatory peaks. To match this, analyses of *Delay 1.2* were restricted

to a 400 msec-long window that began 100 msec after the offset of the cue. We

computed the power spectral density for each epoch using multitaper Fast Fourier

Transform (mtmfft) with Hanning tapering across a frequency range of 3-40 Hz. Spectra

were calculated for individual trials and then averaged across trials for each participant.

To isolate oscillatory peaks from the background aperiodic (1/f) component, we

applied the Specparam algorithm (formerly FOOOF; [34]) with the following parameters:

3-40 Hz frequency range; 5-12 Hz peak width limits; 4 peaks per channel maximum;

0.15 minimum peak height; 2.0 peak threshold; fixed aperiodic mode. For each channel,

the algorithm fitted the aperiodic component and identified oscillatory peaks. For

visualization, we plotted histograms of all identified peaks across participants and

channels for each task and delay period. We used two-sample Kolmogorov-Smirnov (K-

S) Tests to assess whether the distributions of peak frequencies differed between

epochs. Because the K-S test is sensitive to any differences in distribution shape,

location, or spread, and detects the largest local discrepancy between distributions,

significant results would indicate that there were global distributional changes in

oscillatory peak frequencies between epochs. To test for differences in central tendency

(median) between the distribution of peaks from the two epochs (i.e., *Delay 1.1* vs.

Delay 1.2), we used Mann-Whitney U tests, for which significant results would indicate

that there were systematic shifts in the peak median frequency between epochs. Finally,

we calculated the Wasserstein distance (Earth Mover's Distance) between the

distributions of peaks from the two epochs. This measure quantifies the minimum cost

of transforming one distribution into the other, computed across the *entire* frequency

distribution, accounting for both the magnitude of differences and the distance over

which spectral energy must be displaced [35-36]. The Wasserstein distance was

calculated for the pooled data (i.e., all peaks extracted across participants) between all

peak frequencies from *Delay 1.1* and all peak frequencies from *Delay 1.2* using 200

quantile comparison points. We then shuffled the epoch labels and computed the

distance 5,000 times and the *p*-value was computed as the proportion of shuffled data

sets that showed a distance equal to or greater than the actual observed distance. We

also performed a participant-level analysis, computing the distance between epochs for

each participant, then shuffling the condition labels within that participant's data set

2,000 times to create a participant-specific null distribution. To assess whether within-

participant distributional shifts were consistent across participants, we randomly

sampled one value from each participant's null distribution and averaged across

participants (repeated 2,000 times), and compared the observed group mean distance

to this null distribution.

Beta band peak extraction. We next refined our peak extraction analysis

by focusing on the two ranges identified in the MAE behavioral oscillations in

Experiment 1: a 'low-beta' range of 14-16 Hz; and a 'higher-beta' range of 19-21 Hz.

The spectral analysis pipeline used for broadband peak extraction was applied to the

same *Delay 1.1* and *Delay 1.2* epoch data from the DSR and NEU tasks, extracting

peaks only in the specified low-beta and higher-beta ranges. For each channel and

participant, we extracted all peaks in each band and averaged the band counts

separately across channels. We then used a three-way repeated measures ANOVA with

factors: Task (DSR vs. NEU), Band (low-beta: 14-16 Hz vs. higher-beta: 19-21 Hz), and

Epoch (*Delay 1.1* vs. *Delay 1.2*) to quantify differences in beta peak density (in units of

peaks per channel). Follow-up analyses included task-specific paired *t*-tests comparing

peak densities across epochs within each band and a focused test of the Band × Epoch

crossover pattern identified in the DSR condition. For visualization, we plotted the

group-average beta peak density topographically across conditions. These topographic

maps are descriptive and were not subject to statistical testing due to sparse peak

distributions.

Aperiodic component analysis. In addition to analyzing oscillatory peaks,

we examined the aperiodic component of the power spectra, which reflects broadband

neural activity independent of narrowband oscillations. The Specparam algorithm

models the aperiodic component as a 1/f-like function characterized by two parameters:

the offset (reflecting broadband power) and the exponent (reflecting the slope of the

power spectral decay, which has been proposed as a proxy for the balance between

neural excitation and inhibition ("E:I balance"; [37-38]). For each participant, we

extracted the aperiodic exponent and offset for each channel, then averaged across

channels to obtain a single value per participant for each epoch and task. To test

whether aperiodic parameters changed across delay periods, and whether these

changes differed by task, we conducted 2 (Task: DSR, NEU) × 2 (Epoch: *Delay 1.1*,

Delay 1.2) repeated measures ANOVAs separately for offset and exponent. Prior to

analysis, we assessed normality of the difference scores using the Lilliefors and

Anderson-Darling tests. Because the exponent difference scores violated normality

assumptions in the DSR task (Lilliefors p = .005, kurtosis = 5.35), we supplemented the

parametric ANOVAs with non-parametric Wilcoxon signed-rank tests on the epoch

effects collapsed across tasks.

Sub-band burst rate analysis. To examine frequency-specific temporal

dynamics of beta bursts, we separately quantified burst rates in the low-beta (14-16 Hz)

and higher-beta (19-21 Hz) frequency bands. Bursts were detected using the Matlab

implementation of the SpectralEvents toolbox

(https://github.com/jonescompneurolab/SpectralEvents) with time-frequency analysis

conducted via Morlet wavelet decomposition (width = 5) across 13-30 Hz. Following

prior studies, findMethod = 1 was used as in [39], see also [30], which is agnostic to

event overlap, and an event threshold of 6× the median power across time and epochs

for each frequency bin of the time-frequency representation.

For each participant, bursts with a peak frequency falling within each sub-band of

interest were identified, and their rates (events per second) were calculated for the

same two temporal epochs used in the oscillatory peak analysis (see ***Broadband***

oscillatory peak extraction): the final 400 ms of *Delay 1.1* (pre-cue baseline period)

and a 400 ms window beginning 100 ms after cue onset during *Delay 1.2* (post-cue

period). Burst rates were averaged across all scalp electrodes to yield a single value per

participant, task, epoch, and frequency band.

We first computed two descriptive measures of trial prevalence to characterize

burst prevalence. First, we calculated the proportion of trials containing at least one

burst within each channel, then averaged these proportions across channels (per-

channel prevalence). Second, we calculated the proportion of trials containing at least

one burst anywhere across all channels (any-channel prevalence), which provides a

more direct estimate of how frequently bursts occurred during the epochs.

Statistical analyses utilized paired *t*-tests comparing low-beta and higher-beta

burst *rates* (events/sec) within each task and epoch, paired *t*-tests comparing *Delay 1.1*

and *Delay 1.*2 burst rates within each frequency band and task, and separate 2×2

repeated-measures ANOVAs (Task [DSR vs NEU] × Epoch [*Delay 1.1* vs *Delay 1.2*]) for

each frequency band to assess main effects and interactions.

Oscillatory power. To complement the spectral parameterization and burst

rate analyses, we examined oscillatory power in the low-beta (14-16 Hz) and higher-

beta (19-21 Hz) frequency bands. Time-frequency decomposition was performed using

Morlet wavelets (width = 5 cycles) as implemented in FieldTrip. Power was computed

over the timepoints spanning *Delay 1.1* through *Delay 1.2* (a total of 5000 ms sampled

at 250 Hz for 1250 timepoints) in 50 ms steps. We decomposed power into induced and

evoked components. Total power was computed by first applying wavelet decomposition

to each trial and then averaging across trials. Evoked power was computed by first

averaging the raw signal across trials to obtain the event-related potential and then

applying wavelet decomposition to the averaged signal. Induced power was calculated

as total power minus evoked power, isolating oscillatory activity that was not phase-

locked to stimulus onset.

Initially, the power values were not baseline-corrected because the period

preceding *Delay 1.1* contained stimulus presentation (*Sample 2*) and thus did not

provide an appropriate baseline. For statistical analysis, power was averaged across all

channels and across the two time windows used in the other spectral analyses

described above: the final 400 ms of *Delay 1.1* and the first 400 ms of *Delay 1.2*, offset

by 100 ms to avoid sample-evoked and cue-evoked transients, respectively. Induced

and evoked power were analyzed separately for each frequency band using 2 × 2

repeated measures ANOVAs with factors Task (DSR, NEU) and Epoch (pre-cue, post-

cue). Follow-up paired *t*-tests were carried out to examine simple effects the two factors

with a Bonferroni-corrected alpha-level = .025 (for two comparisons per effect/band).

Because strong task differences emerged in the non-baseline-corrected analysis,

we carried out a follow-up comparison of the power in the two tasks using *Delay 1.1* as

the baseline allowing us to address whether the power in the two tasks exhibited

differential cue-related changes in power, rather than just differing in absolute power

levels. For each participant, we computed the percent change in power from pre-cue to

post-cue as ((*post − pre*) / *pre*) × 100 separately for each frequency band and power

type (induced, evoked). Paired *t*-tests were carried out to test for differences in percent

power change between DSR and NEU at the Bonferroni-correct alpha-level = .025.

Time-resolved representational similarity analysis (RSA)

To examine how neural representations of prioritized and unprioritized memory items

(i.e., PMIs and UMIs) evolved over time, we computed a time-resolved representational

similarity analysis (RSA) separately for each item and participant. After the

preprocessed EEG data were downsampled to 250 Hz and baseline-corrected by

subtracting each channel's trial-averaged activity, for each participant, and timepoint, we

constructed representational dissimilarity matrices (RDMs) capturing the pairwise

dissimilarities (1 − Pearson's *r*) between condition-averaged ERP patterns across

electrodes. Conditions were defined by all 24 combinations of angular difference (0°, +/-

22.5°, +/-45°, +/-67.5°) and base orientation (10° – 160° in 30° steps), separately for

PMIs and UMIs in the DSR task. If there were no trials for a particular combination of

angular difference + base orientation in a priority condition (due to the removal of bad

trials during pre-processing), that row and column were omitted from the analysis. This

occurred for 1 row/column of 1 participant's analysis. Because the NEU task had half

the number of trials as the DSR task, row/column omissions were more frequent for the

tested and untested trial splits, and, as such, conditions were instead defined by the 6

base orientations, collapsing over the four angular difference conditions to ensure more

balance across participants. To track representational changes with priority, we first

generated two reference models for each task (one each corresponding to the

PMI/tested item and to the UMI/untested item) by computing an RDM from the 500

msec of Delay 1.1. Next, RDMs were computed at each timepoint spanning *Delay 1.1*

through *Delay 1.2* (in 7-sample (28 ms) windows) and correlated (Spearman's rho) with

the reference RDM. The resulting similarity values (i.e., the Spearman correlations)

were averaged across participants to produce group-averaged time-resolved

representational similarity time courses. Statistical comparisons of group-level

representational similarity across epochs (i.e., *Delay 1.1, Cue 1, Delay 1.*2) used paired-

sample *t*-tests on the within-participant averaged similarities within the epoch intervals

(i.e., similarities were averaged across the 28 ms windows falling within each epoch).

To reveal the importance (contribution) of each electrode's activity to the

representation at any point in the analysis interval, we repeated the procedure for the

original RSA analysis but for each electrode separately. In this iterative analysis, the

reference RDMs were computed as in the original analysis using data from all

electrodes during *Delay 1.1*. The timepoint-by-timepoint RDMs, however, only used data

from the single electrode being analyzed in the iteration. This analysis thus yielded a

similarity map across the scalp, such that electrodes with larger correlations were those

with greater importance/contribution to the representational similarity observed at each

timepoint in the original analysis. For visualization, the resulting importance maps were

averaged across participants to produce topographic maps of representational similarity

during the *Delay 1.1*, *Cue 1,* and *Delay 1.2* epochs.

Spectral analysis of RSA time courses

To characterize rhythmic fluctuations in representational similarity, we spectrally

decomposed the RSA time courses. First, each participant's PMI, UMI, tested, and

untested RSA time course was mean-centered. Subsequently, the spectral analysis of

the RSA was identical to the spectral analysis of the raw EEG (see section ***EEG***

spectral analysis), with the following exceptions: (1) Power spectral density was

computed using a standard FFT with an NFFT of 512 samples rather than multitaper

estimation with Hanning tapering. This is because the RSA time series represents a

single derived measure per participant rather than noisy multi-channel

electrophysiological recordings, which reduces the need for variance-bias tradeoffs

afforded by multitaper methods. (2) Because RSA yields a single time series per

participant (rather than multiple channels), peak counts reflect the number of peaks

detected in each participant's RSA trace, and aperiodic parameters were extracted

directly from each participant's spectrum without averaging across channels.

For the beta band peak extraction analysis, we computed peak counts separately

for the PMI and UMI in the RSA time courses in the DSR task, and for the tested and

untested items in the RSA time courses in the NEU task. To parallel the task-level EEG

analysis, we summed peak counts across PMI and UMI (for DSR) or across tested and

untested items (for NEU) before conducting the 3-way repeated measures ANOVA (Task

× Band × Epoch). We additionally conducted a 4-way repeated measures ANOVA

including Priority (PMI/UMI or tested/untested) as a factor to examine whether spectral

signatures differed by memory item status.

To directly test whether a task-specific beta band dissociation emerged following

the retrocue, we conducted a focused analysis of *Delay 1.2* using a 2 (Task: DSR, NEU)

× 2 (Band: low-beta, high-beta) repeated measures ANOVA. We followed up significant

interactions with simple effects tests comparing tasks within each band, and a crossover

test quantifying the difference of task effects across bands (i.e., $[DSR − NEU]_{low\text{-}beta}$ −

$[DSR − NEU]_{higher\text{-}beta}$). We also examined Task × Epoch interactions separately for each

band to determine whether cue-related changes in peak counts were task-specific.

For the aperiodic component analysis, we computed 2 (Task) × 2 (Epoch)

repeated measures ANOVAs for exponent and offset at the task level (averaging across

PMI/UMI or tested/untested), as well as 3-way ANOVAs (Task × Priority × Epoch) to

examine priority-specific effects.

To test for evidence of phase encoding of the representations of the two memory

items we computed cross-spectral phase consistency between the time-resolved RSA

time courses for each item pair (PMI/UMI in DSR; tested/untested in NEU). For each

participant, we computed cross-spectral phase using the fast Fourier transform (FFT)

with zero-padding to 512 points for the *Delay 1.1* and *Delay 1.2* epochs used in the beta peak extraction analysis and computed the phase difference between the two RSA traces from the angle of the cross-spectrum. Phase values were averaged across frequency bins within the two bands of interest: low-beta (14–16 Hz; 5 bins: 14.16, 14.65, 15.14, 15.62, 16.11 Hz–note that the addition of the final 16.11 Hz bin ensured the complete 2 Hz band would be captured) and higher-beta (19–21 Hz; 5 bins: 19.04, 19.53, 20.02, 20.51, 21.00 Hz), using circular averaging. To assess whether the phase relationships were consistent across participants, we computed the resultant vector length (r) and tested for non-uniformity using the Rayleigh test. We obtained bootstrapped standard errors for r by resampling participants with replacement (5,000 iterations). To test whether cue-related changes in phase consistency differed between tasks, we conducted a permutation test on the Task × Epoch interaction ($r_{DSRpost}$ − r_{DSRpre}) − ($r_{NEUpost}$ − r_{NEUpre}). The null distribution was generated by independently shuffling epoch labels (pre/post) within each task for each participant (10,000 permutations), and a two-tailed p-value was computed as the proportion of permuted interaction values with absolute magnitude greater than or equal to the observed value.

References

1. Hautekiet C, Niklaus M, Oberauer K. Susceptibility to visual interference in working memory: Different results depending on the prioritization mode? J Exp Psychol Hum Percept Perform. 2025;51(6):791–807. doi: 10.1037/xhp0001315

1264   2.   Sprague TC, Ester EF, Serences JT. Restoring latent visual working memory

1265   representations in human cortex. Neuron. 2016;91:694–707. doi:

1266   10.1016/j.neuron.2016.07.006

1267   3.   Yu Q, Teng C, Postle BR. Different states of priority recruit different neural

1268   representations in visual working memory. PLoS Biol. 2020;18:e3000769. doi:

1269   10.1371/journal.pbio.3000769

1270   4.   Panichello MF, Buschman TJ. Shared mechanisms underlie the control of

1271   working memory and attention. Nature. 2021;592:601–5. doi: 10.1038/s41586-

1272   021-03390-w

1273   5.   Piwek EP, Stokes MG, Summerfield C. A recurrent neural network model of

1274   prefrontal brain activity during a working memory task. PLoS Comput Biol.

1275   2023;19(10):e1011555. doi: 10.1371/journal.pcbi.1011555

1276   6.   Wan Q, Menendez JA, Postle BR. Priority-based transformations of stimulus

1277   representation in visual working memory. PLoS Comput Biol.

1278   2022;18(6):e1009062. doi: 10.1371/journal.pcbi.1009062

1279   7.   Lewis-Peacock JA, Drysdale AT, Oberauer K, Postle BR. Neural evidence for a

1280   distinction between short-term memory and the focus of attention. J Cogn

1281   Neurosci. 2012;24:61–79. doi: 10.1162/jocn_a_00140

1282   8.   Rose N, LaRocque JJ, Riggall AC, Gosseries O, Starrett MJ, Meyering E, et al.

1283   Reactivation of latent working memories with transcranial magnetic stimulation.

1284   Science. 2016;354:1136–9. doi: 10.1126/science.aah7011

9. Fulvio JM, Haegens S, Postle BR. Single-pulse TMS affects working-memory

performance via posterior beta-band oscillations. J Cogn Neurosci. 2024. doi:

10.1162/jocn_a_02194

10. Fulvio JM, Postle BR. Effects of TMS on the decoding and electrophysiology of

priority in working memory. bioRxiv. 2025. doi: 10.1101/2025.09.15.676374

11. Abdalaziz M, Redding ZV, Fiebelkorn IC. Rhythmic temporal coordination of

neural activity prevents representational conflict during working memory. Curr

Biol. 2023;33:1855–63. doi: 10.1016/j.cub.2023.03.088

12. Hervault M, Wessel JR. Common and unique neurophysiological processes that

support the stopping and revising of actions. J Neurosci. 2025;45(13). doi:

10.1523/JNEUROSCI.1537-24.2025

13. Diesburg DA, Greenlee JD, Wessel JR. Cortico-subcortical β burst dynamics

underlying movement cancellation in humans. eLife. 2021;10. doi:

10.7554/eLife.70270

14. Lundqvist M, Miller EK, Nordmark J, Liljefors J, Herman P. Beta: bursts of

cognition. Trends Cogn Sci. 2024. doi: 10.1016/j.tics.2024.03.010

15. Samaha J, Sprague TC, Postle BR. Decoding and reconstructing the focus of

spatial attention from the topography of alpha-band oscillations. J Cogn

Neurosci. 2016;28:1090–7. doi: 10.1162/jocn_a_00955

16. Bae GY, Luck SJ. Interactions between visual working memory representations.

Atten Percept Psychophys. 2017;79(8):2376–95. doi: 10.3758/s13414-017-1404-

8

17. Bliss DP, Sun JJ, D'Esposito M. Serial dependence is absent at the time of

perception but increases in visual working memory. Sci Rep. 2017;7(1):14739.

doi: 10.1038/s41598-017-15199-7

18. Fritsche M, Mostert P, de Lange FP. Opposite effects of recent history on

perception and decision. Curr Biol. 2017;27(4):590–5. doi:

10.1016/j.cub.2017.01.006

19. Samaha J, Switzky M, Postle BR. Confidence boosts serial dependence in

orientation estimation. J Vis. 2019;19(4):25. doi: 10.1167/19.4.25

20. Fulvio JM, Rokers B, Samaha J. Task feedback suggests a post-perceptual

component to serial dependence. J Vis. 2023;23(10):6. doi: 10.1167/jov.23.10.6

21. vanSnellenberg JX. BCa_bootstrap. MathWorks File Exchange; 2018

https://www.mathworks.com/matlabcentral/fileexchange/69119

22. Fiebelkorn IC, Kastner S. A rhythmic theory of attention. Trends Cogn Sci.

2019;23:87–101. doi: 10.1016/j.tics.2018.11.009

23. Buschman TJ, Miller EK. Serial, covert shifts of attention during visual search are

reflected by the frontal eye fields and correlated with population oscillations.

Neuron. 2009;63:386–96. doi: 10.1016/j.neuron.2009.06.020

24. Bastos AM, Loonis R, Kornblith S, Lundqvist M, Miller EK. Laminar recordings in

frontal cortex suggest distinct layers for maintenance and control of working

memory. Proc Natl Acad Sci USA. 2018;76:201710323. doi:

10.1073/pnas.1710323115

25. Lundqvist M, Rose J, Herman P, Brincat SL, Buschman TJ, Miller EK. Gamma

and beta bursts underlie working memory. Neuron. 2016;90:1–13. doi:

10.1016/j.neuron.2016.02.028

26. Siegel M, Warden MR, Miller EK. Phase-dependent neuronal coding of objects in

short-term memory. Proc Natl Acad Sci USA. 2009;106:21341–6. doi:

10.1073/pnas.0908193106

27. Lisman JE, Idart MA. Storage of 7 ± 2 short-term memories in oscillatory

subcycles. Science. 1995;313:1512–5. doi: 10.1126/science.7878473

28. Hakim N, Vogel EK. Phase-coding memories in mind. PLoS Biol.

2018;16:e3000012. doi: 10.1371/journal.pbio.3000012

29. Wessel JA, Anderson MA. Neural mechanisms of domain-general cognitive

control. Trends Cogn Sci. 2024;28(2):124–43. doi: 10.1016/j.tics.2023.09.008

30. Kavanaugh BC, Vigne MM, Thorpe R, Legere C, Acuff WL, Vaughan N, et al. The

association between oscillatory burst features and human working memory

accuracy. J Cogn Neurosci. 2026;38(2):281–98. doi: 10.1162/JOCN.a.87

31. Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-

trial EEG dynamics including independent component analysis. J Neurosci

Methods. 2004;134(1):9–21. doi: 10.1016/j.jneumeth.2003.10.009

32. Oostenveld R, Fries P, Maris E, Schoffelen JM. FieldTrip: Open source software

for advanced analysis of MEG, EEG, and invasive electrophysiological data.

Comput Intell Neurosci. 2011;2011:156869. doi: 10.1155/2011/156869

33. Pion-Tonachini L, Kreutz-Delgado K, Makeig S. ICLabel: An automated

electroencephalographic independent component classifier, dataset, and

website. NeuroImage. 2019;198:181–97. doi: 10.1016/j.neuroimage.2019.05.026

34. Donoghue T, Haller M, Peterson EJ, Varma P, Sebastian P, Gao R, et al.

Parameterizing neural power spectra into periodic and aperiodic components.

Nat Neurosci. 2020;23(12):1655–65. doi: 10.1038/s41593-020-00744-x

35. Rubner Y, Tomasi C, Guibas LJ. The earth mover's distance as a metric for

image retrieval. Int J Comput Vis. 2000;40(2):99–121. doi:

10.1023/A:1026543900054

36. Lipp A, Vermeesch P. The Wasserstein distance as a dissimilarity metric for

comparing detrital age spectra and other geological distributions. Geochronology.

2023;5(1):263–70. doi: 10.5194/gchron-5-263-2023

37. Gao R, Peterson EJ, Voytek B. Inferring synaptic excitation/inhibition balance

from field potentials. NeuroImage. 2017;158:70–8. doi:

10.1016/j.neuroimage.2017.06.078

38. Waschke L, Donoghue T, Fiedler L, Smith S, Garrett DD, Voytek B, et al.

Modality-specific tracking of attention and sensory statistics in the human

electrophysiological spectral exponent. eLife. 2021;10:e70068. doi:

10.7554/eLife.70068

39. Shin H, Law R, Tsutsui S, Moore CI, Jones SR. The rate of transient beta

frequency events predicts behavior across tasks and species. eLife.

2017;6:e29086. doi: 10.7554/eLife.29086

40. Chunharas C, Rademaker RL, Brady TF, Serences JT. An adaptive perspective

on visual working memory distortions. J Exp Psychol Gen. 2022;151(10):2300–

23. doi: 10.1037/xge0001191

41. Scotti PS, Hong Y, Leber AB, Golomb JD. Visual working memory items drift

apart due to active, not passive, maintenance. J Exp Psychol Gen.

2021;150(12):2506–24. doi: 10.1037/xge0000890

42. Fulvio JM, Postle BR. Cognitive control, not time, determines the status of items

in working memory. J Cogn. 2020;3(1):8. doi: 10.5334/joc.98

43. Rosanova M, Casali A, Bellina V, Resta F, Mariotti M, Massimini M. Natural

frequencies of human corticothalamic circuits. J Neurosci. 2009;29(24):7679–85.

doi: 10.1523/JNEUROSCI.0445-09.2009

44. Kundu B, Johnson JS, Postle BR. Prestimulation phase predicts the TMS-evoked

response. J Neurophysiol. 2014;112(8):1885–93. doi: 10.1152/jn.00390.2013

45. Samaha J, Gosseries O, Postle BR. Distinct oscillatory frequencies underlie

excitability of human occipital and parietal cortex. J Neurosci. 2017;37(11):2824–

33. doi: 10.1523/JNEUROSCI.3413-16.2017

46. Wan Q, Ardalan A, Fulvio JM, Postle BR. Representing context and priority in

working memory. J Cogn Neurosci. 2024;36(7):1374–94. doi:

10.1162/jocn_a_02166

47. Teng C, Postle BR. Investigating the roles of the visual and parietal cortex in

representing content versus context in visual working memory. eNeuro.

2024;11(2). doi: 10.1523/ENEURO.0270-20.2024

1393

Supporting information

S1 Fig. Experiment 1 recall accuracy. A. Mean absolute recall error for the two responses of the DSR task with the second recall accuracy reported separately for 'stay' and 'switch' trials and the single response of the NEU task. **B.** Same as **A.** separated by the current trial's inter-item angular difference. Error bars correspond to +/- 1 *SEM.* * symbols correspond to significant differences between conditions at the Bonferroni-corrected alpha level = .0083 for 6 paired-sample *t*-tests. See **Figure 2A & 2B** in the main text for the results collapsed over DSR *Recall 2* trial types. Splitting the DSR *Recall 2* responses according to whether the trial was a stay trial (same item cued for both recalls) or a switch trial (different items cued), revealed no difference in performance.

S1 Table. Linear mixed-effects model of inter-item biases in the DSR and NEU tasks (Experiment 1). Linear mixed-effects models were fit separately for the DSR and NEU tasks with identical fixed- and random-effects structures. Only *Recall 1* of the DSR task was used in the full LME analyses. Coefficients are expressed in radians. Coefficients involving sine-transformed predictors and their interactions are also expressed in degrees of response error per unit change in the sine-transformed orientation variable in parentheses. Sine-transformed predictors reflect directionally signed orientation-dependent biases with positive values indicating attractive biasing effects of the predictor and negative values indicating repulsive biasing effects of the predictor. For the current-trial non-target, simple effects are reported separately for trials in which the non-target was presented first versus second, derived from the interaction between presentation order and the non-target orientation term.

S2 Table. Joint DSR-NEU Linear mixed effects model fixed effects: task-dependent interactions only (Experiment 1). Linear mixed-effects model results comparing the NEU and DSR tasks. Only interactions involving task effects are shown, indicating how the inter-item biases differ between the two tasks. β estimates (in radians) are reported with their 95% confidence intervals (CI) and two-tailed p-values; β values converted to degrees are shown in parentheses for interpretability. Positive β values indicate that the bias associated with the predictor is more attractive in the NEU task relative to the

reference DSR task, whereas negative β values indicate that the bias is less attractive or more repulsive

in NEU relative to DSR. For interactions with the non-target order (categorical), a positive β indicates that

NEU trials with the target presented first are more attractive than the same category in DSR; a negative β

indicates that target first trials are less attractive or more repulsive. Main effects from this joint model are

omitted because they reflect the DSR reference condition and are fully reported in the task-specific

models above (**S1 Table**).

1427

S3 Table. Linear mixed effects model results for signed bias as a function of angular difference

and presentation order (Experiment 1). Linear mixed-effects models were fit separately for the DSR

and NEU tasks with identical fixed- and random-effects structures. The dependent variable was signed

bias, computed by multiplying the signed recall error by the sign of the angular difference between the

untested and target items, such that positive values indicate attraction toward the untested item and

negative values indicate repulsion. Trials with 0° angular difference were excluded. Coefficients are

expressed in radians with degree equivalents in parentheses. The reference level for Angular Difference

is 22.5°, and for Non-Target Order is non-target presented second. Coefficients for Angular Difference

reflect the change in bias relative to the 22.5° reference condition. Previous trial predictors are sine-

transformed to capture directionally signed orientation-dependent biases, with positive values indicating

attractive biasing effects and negative values indicating repulsive biasing effects.

1439

S4 Table. Joint DSR-NEU Linear mixed effects model fixed effects: task-dependent interactions

only (Experiment 1, Angular Difference Analysis). Linear mixed-effects model results comparing the

NEU and DSR tasks for the signed bias analysis. Only interactions involving task effects are shown,

indicating how inter-item biases differ between the two tasks. β estimates (in radians) are reported with

their 95% confidence intervals (CI) and two-tailed p-values; β values converted to degrees are shown in

parentheses for interpretability. Positive β values indicate that the signed bias (attraction toward the

untested item) is greater in the NEU task relative to the reference DSR task, whereas negative β values

indicate that the bias is smaller or more repulsive in NEU relative to DSR. The significant main effect of

Task indicates that overall attraction toward the untested item was greater in NEU than DSR. The

significant Non-Target Order × Task interaction indicates that the reduction in bias when the untested

item was presented first was greater in NEU than DSR. The significant Angular Difference: 67.5° × Task

interaction indicates that the reduction in bias at large angular separations was greater in NEU than DSR.

Main effects from this joint model are omitted because they reflect the DSR reference condition and are

fully reported in the task-specific models above (**S3 Table**).

1454

S2 Figure. Experiment 1 behavioral oscillations of mean inter-item bias. Left: Demeaned mean inter-

item bias for DSR *Recall 1* (red) and the single response of the NEU task (blue) as a function of Delay 1.2

length. Error bands correspond to +/- 1 *SEM.* Right: Fast Fourier Transform (FFT) amplitude spectra of

mean inter-item bias across delay periods for the DSR (red) and NEU (blue) tasks. Solid lines represent

group-averaged amplitudes (in degrees of error). Dashed lines show statistical significance thresholds

(95th percentile) derived from permutation testing.

1461

S3 Figure. Experiment 2 recall accuracy. A. Mean absolute recall error for the two responses of the

DSR task with the second recall accuracy reported separately for 'stay' and 'switch' trials and the single

response of the NEU task. **B.** Same as **A.** separated by the current trial's inter-item angular difference.

Error bars correspond to +/- 1 *SEM.* * symbols correspond to significant differences between conditions at

the Bonferroni-corrected alpha level = .0083 for six paired-sample *t*-tests (**A.**). As in Experiment 1,

splitting the DSR *Recall 2* responses according to whether the trial was a stay trial (same item cued for

both recalls) or a switch trial (different items cued), revealed no difference in performance. Compare with

Figure S1.

1470

S5 Table. Linear mixed-effects model of inter-item biases in the DSR and NEU tasks (Experiment

2). Linear mixed-effects models were fit separately for the DSR and NEU tasks with identical fixed- and

random-effects structures. Coefficients are expressed in radians. Coefficients involving sine-transformed

predictors and their interactions are also expressed in degrees of response error per unit change in the

sine-transformed orientation variable in parentheses. Sine-transformed predictors reflect directionally

signed orientation-dependent biases with positive values indicating attractive biasing effects of the

predictor and negative values indicating repulsive biasing effects of the predictor. For the current-trial

non-target, simple effects are reported separately for trials in which the non-target was presented first

versus second, derived from the interaction between presentation order and the non-target orientation

term. Compare to **S1 Table**.

1481

S6 Table. Joint DSR-NEU Linear mixed effects model fixed effects: task-dependent interactions

only (Experiment 2). Linear mixed-effects model results comparing the NEU and DSR tasks. Only

interactions involving task effects are shown, indicating how the inter-item biases differ between the two

tasks. β estimates (in radians) are reported with their 95% confidence intervals (CI) and two-tailed p-

values; β values converted to degrees are shown in parentheses for interpretability. Positive β values

indicate that the bias associated with the predictor is more attractive in the NEU task relative to the

reference DSR task, whereas negative β values indicate that the bias is less attractive or more repulsive

in NEU relative to DSR. For interactions with the non-target order (categorical), a positive β indicates that

NEU trials with the target presented first are more attractive than the same category in DSR; a negative β

indicates that target first trials are less attractive or more repulsive. Main effects from this joint model are

omitted because they reflect the DSR reference condition and are fully reported in the task-specific

models above (**S5 Table;** Compare to **S2 Table**).

1494

S7 Table. Linear mixed effects model results for signed bias as a function of angular difference

and presentation order (Experiment 2). Linear mixed-effects models were fit separately for the DSR

and NEU tasks with identical fixed- and random-effects structures. The dependent variable was signed

bias, computed by multiplying the signed recall error by the sign of the angular difference between the

untested and target items, such that positive values indicate attraction toward the untested item and

negative values indicate repulsion. Trials with 0° angular difference were excluded. Coefficients are

expressed in radians with degree equivalents in parentheses. The reference level for Angular Difference

is 22.5°, and for Non-Target Order is non-target presented second. Coefficients for Angular Difference

reflect the change in bias relative to the 22.5° reference condition. Previous trial predictors are sine-

transformed to capture directionally signed orientation-dependent biases, with positive values indicating

attractive biasing effects and negative values indicating repulsive biasing effects. Compare to **S3 Table.**

1506

1507 **S4 Fig. Topography of low beta (14-16 Hz) peak oscillations.** Left column: Mean number of extracted

1508 peaks in the low beta (14-16 Hz) band per channel for *Delay 1.1* in the DSR task (top row), the NEU task

1509 (middle row), and their difference (DSR-NEU, bottom row). Middle column: Same as the left-hand column,

1510 but for *Delay 1.2.* Right column: Within-task differences between the two delay periods (*Delay 1.1-Delay*

1511 *1.*2) for the DSR task (top row) and the NEU task (middle row).

1512

1513 **S5 Fig. Topography of high beta (19-21 Hz) peak oscillations.** Left column: Mean number of extracted

1514 peaks in the low beta (19-21 Hz) band per channel for *Delay 1.1* in the DSR task (top row), the NEU task

1515 (middle row), and their difference (DSR-NEU, bottom row). Middle column: Same as the left-hand column,

1516 but for *Delay 1.2.* Right column: Within-task differences between the two delay periods (*Delay 1.1-Delay*

1517 *1.*2) for the DSR task (top row) and the NEU task (middle row).

1518

S6 Fig. EEG power analyses. A. Left: Raw, uncorrected time-resolved induced (upper row) and evoked

(lower row) low-beta (14-16 Hz) power for the DSR task (blue) and NEU task (red) for the timepoints

spanning *Delay 1.1* ('pre-cue') through *Delay 1.2* ('post-cue'). Right: Mean power for the two tasks during

the final 400 ms of the *Delay 1.1* and the first 400 ms of *Delay 1.2* (beginning 100 ms after cue offset). **B.**

Same as **A.** for higher-beta (19-21 Hz) power. **C.** Percent change (post- - pre-cue)/pre-cue *100 in

induced power (left) and evoked power (right) for the two tasks and frequency bands. Error bars

correspond to +/- 1 *SEM.* * corresponds to significant differences at the Bonferroni-corrected alpha-level =

.025.

1527

1528 **S1 File. Aperiodic component analyses for beta-band peak extraction at the task level.**

1529

1530 **S2 File. Aperiodic component analyses for beta-band peak extraction at the item level.**

1531

1532   **S3 File. Individual analyses of within- and between-trial inter-item biases for Experiments 1 and 2.**

1533

1534   Figure 1.

1535

1536

1537

1538

1539   Figure 2

1540

1541

1542

1543   Figure 3.

1544

1545

1546

1547   Figure 4.

1548

1549

1550

1551

1552   Figure 5.

1553

1554

1555   Figure 6.

1556

1557

1558

1559   Figure 7.

1560

1561

1562