Hypanus brevis: a newly resurrected Eastern South Pacific stingray lineage revealed by integrative taxonomy

Alan Marín[1][*], Fabiola Zavalaga[2], Renato Gozzer-Wuest[3], Luis E. Santos-Rojas[1], Lorenzo E. Reyes-Flores[1], Ruben Alfaro[4], Philippe Béarez[5], Eliana Zelada-Mázmela[1]

[1] *Laboratorio de Genética, Fisiología y Reproducción, Facultad de Ciencias, Universidad Nacional del Santa, Chimbote, Perú*

[2] *Área Funcional de Investigaciones en Biodiversidad, Instituto del Mar del Perú - IMARPE, Callao, Perú*

[3] *Innovations for Ocean Action Foundation (I4OA), Lima, Peru*

[4] *Ecobiotech Lab S.A.C. Trujillo, Perú*

[5] *UMR 7209 BioArch, CNRS-MNHN, Muséum national d'Histoire naturelle, CP 56, 43 rue Buffon, 75005 Paris, France*

*Corresponding author
email amarin@uns.edu.pe

Abstract

Hypanus brevis (Garman, 1880) and *Hypanus dipterurus* (Jordan & Gilbert, 1880) are currently considered as a conspecific lineage of the "diamond stingray" from the Eastern Pacific. This taxonomic group has been the subject of nomenclatural disputes for about 145 years. To clarify the historical confusion surrounding this lineage, we employed an integrative taxonomic approach using specimens from the Eastern North and Eastern South Pacific (ENP and ESP). The genetic results, based on single and multilocus mitochondrial analyses, revealed a distinct evolutionary unit in the ESP. While morphological analyses detected subtle differences between ENP and ESP specimens, most characters exhibited significant overlap (e.g., disc shape, dentition patterns, body coloration), suggesting low evolutionary divergence. A calibrated molecular clock analysis estimated this divergence at approximately 3.09 Ma. In accordance with Garman's (1880) original description based on specimens from Paita (northern Peru), we formally resurrect *H. brevis* from synonymy with *H. dipterurus*. Our findings suggest an anti-tropical speciation pathway, with core populations of *H. brevis* and *H. dipterurus* restricted to the temperate waters of the ESP and ENP, respectively. Notably, a single, fixed COI haplotype was detected in all *H. brevis* specimens from the north-central Peruvian coast. This result may indicate a severe bottleneck event, raising concerns about the genetic health and long-term viability of this vulnerable species. Finally, we analyzed historical fishery data of *H. brevis* to infer its current population status, suggesting targeted conservation measures and precautionary management to prevent further loss of genetic diversity.

Keywords: Cryptic species, marine conservation, molecular phylogeny, Myliobatiformes, taxonomic resurrection

1. INTRODUCTION

Species of the genus *Hypanus* Rafinesque, 1818 (Myliobatiformes: Dasyatidae) are characterized by their wide pectoral fins that are fused to the head, a prominent row of thorns on nape and one or more thorns on shoulder in adults, a distinct but low ventral skin fold on tail (Last et al., 2016a). Most *Hypanus* species hold significant commercial value throughout their distribution range in the Atlantic and Pacific Oceans. They are either targeted by artisanal fisheries or frequently retained as valuable bycatch (Campos-León et al., 2025; Feitosa et al., 2021; Mejía-Falla et al., 2025). Due to their high market demand, unmanaged fishing practices, and their k-selected life history traits (Yamaguchi et al., 2021), half of the species in this genus are currently listed as threatened on The IUCN Red List (IUCN, 2025).

The genus *Hypanus* was considered a junior synonym of *Dasyatis*, until it was resurrected by Last et al. (2016b). It is now composed of eleven species (Fricke et al., 2025), including seven species from the Western Atlantic: *Hypanus americanus* (Hildebrand & Schroeder, 1928), *H. berthalutzae* Petean, Naylor & Lima, 2020, *H. geijskesi* (Boeseman, 1948), *H. guttatus* (Bloch & Schneider, 1801), *H. marianae* (Gomes, Rosa & Gadig, 2000), *H. sabinus* (Lesueur, 1824), and *H. say* (Lesueur, 1817); one species from the Eastern Atlantic: *Hypanus rudis* (Günther, 1870); and three species from the Eastern Pacific: *Hypanus dipterurus* (Jordan & Gilbert, 1880), *H. longus* (Garman, 1880), and *H. rubioi* Mejía-Falla, Navia, Cardeñosa & Tavera, 2025. However, a recent phylomitogenomic analysis suggested the existence of three additional evolutionary units within this genus (*H.* aff. *americanus*, *H.* aff. *guttatus*, and *H.* aff. *say*), warranting the need for a thorough taxonomic revision (Petean et al., 2024).

In the Eastern Pacific, the diamond stingray, *H. dipterurus*, is found from southern California (USA) to northern Chile (Almendras et al., 2025; Lamilla et al., 1995). This species is currently listed as Vulnerable on The IUCN Red List of Threatened Species, primarily due to

intense and largely unmanaged fisheries across most of its distribution range (Pollom et al.,

2020). The diamond stingray is one of the most captured species by Mexican artisanal

fisheries (Simental-Anguiano et al., 2022). In Colombia and Ecuador, its capture is limited to

bycatch (Calle-Morán & Béarez, 2020; Puentes et al., 2022), while in Peru, artisanal fisheries

target this species (Campos-León et al., 2025; González-Pestana et al., 2022). Despite its

intense exploitation, high demand, and signals of population declines (Pollom et al., 2020),

no formal management plans have been established in Peruvian stingray fisheries yet

(González-Pestana et al., 2022). Furthermore, a long-standing taxonomic confusion with a

junior synonym, *Dasyatis brevis* (Garman, 1880), may have skewed historical catch estimates.

The diamond stingray has faced a long period of confused nomenclature history, due to two

independent and nearly simultaneous descriptions published in 1880, that led to a confusing

synonymy (Miller et al., 2014). The first description was published in May 1880 by Jordan

and Gilbert, who named it *Dasybatis dipterurus* based on specimens collected from San

Diego Bay in California (Jordan & Gilbert, 1880). Then, in October of the same year,

Garman described *Trygon brevis*, based on two specimens from Paita, Peru (Garman, 1880).

Garman (1913: 396) synonymized the two species (*D. dipterurus* and *T. brevis*) under the

name *Dasybatus brevis*, considering wrongly that the date of publication of Jordan and

Gilbert's work was 1881, thus giving priority to his own description from 1880. This

inappropriate synonymization has led to long-standing ongoing confusion, dividing the

opinion of modern researchers (Miller et al., 2014; Nelson et al., 2004; Nishida & Nakaya,

1990) until Eschmeyer (1998) recognized the nomenclatural priority of *D. dipterurus* Jordan

& Gilbert, 1880 over *T. brevis* Garman, 1880. Ebert (2003), followed by Love et al. (2005)

and Mundy (2005) then reclassified the latter as a junior synonym of *Dasyatis dipterura*, and

finally Last et al. (2016b) reassigned the species to the genus *Hypanus* (currently *Hypanus*

dipterurus). Despite these taxonomic clarification efforts, the status of *H. dipterurus* and *D.*

brevis remains controversial, even 145 years after their initial descriptions.

Inaccurate taxonomic revisions and hidden diversity can lead to a cascade of errors resulting

in dire conservation consequences. This is particularly true for commercially exploited

batoids, due to their high vulnerability to overfishing (Hosegood et al., 2020; White & Last,

2012). Treating multiple distinct species as a single group, or inversely, a single species

managed as different units, can mask localized population declines and thus threaten the

sustainability of these batoids (Chatzispyrou & Koutsikopoulos, 2023; Siskey et al., 2019).

Furthermore, assessments of conservation status rely on the analysis of fisheries statistics

from landing reports and geographical distributions based on taxonomic revisions (IUCN,

2024). The use of flawed catch statistics and imprecise distribution ranges resulting from

taxonomic uncertainties can lead to inaccurate conservation status assessments (Marshall et

al., 2009).

DNA sequence analysis represents a powerful tool for achieving taxonomic clarity,

particularly in groups showing complex morphological identification, such as batoids, where

traditional methods may fall short (Cunha et al., 2026; Marín, 2025; Sales et al., 2019; White

et al., 2022). Genetic studies have played a crucial role in identifying hidden lineages within

morphologically identical ray species, leading to significant formal taxonomic revisions (Last

et al., 2016b; White et al., 2022). For example, a phylogenetic analysis of the southern

stingray, *Hypanus americanus*, revealed that it was paraphyletic (Petean et al., 2020). This

finding led to the discovery of a new species, the Lutz's stingray, *H. berthalutzae* by Petean et

al. (2020). Additionally, genetic studies have led to the reinstatement of names that were

previously treated as junior synonyms. These include the Indo-Pacific spotted eagle ray,

Aetobatus ocellatus (Kuhl, 1823), which was previously considered a junior synonym of *A.*

narinari (Euphrasen, 1790) (White et al., 2010); *Neotrygon trigonoides* (Castelnau, 1873),

previously synonymized with *N. kuhlii* (Müller & Henle 1841) (Borsa et al., 2013); and

Zearaja brevicaudata (Marini, 1933) (currently *Dipterurus brevicaudatus*), which was

confused with *Zearaja chilensis* (Guichenot, 1848) (currently *D. chilensis*) (Gabbanelli et al.,

2018), among others.

Most taxonomic and phylogenetic revisions including *H. dipterurus* samples have primarily

focused on populations from the Eastern North Pacific (hereafter referred to as ENP) (e.g.,

Last et al., 2016b; Naylor et al., 2012; Petean et al., 2020). On the other hand, the populations

of *H. dipterurus* from the Eastern South Pacific (hereafter referred to as ESP) have not yet

been assessed. Consequently, there is a significant knowledge gap regarding the connectivity

and gene flow among the populations from the northern and southern hemispheres.

Furthermore, a recent study suggested that *H. dipterurus* from the ESP may represent a

distinct evolutionary lineage (Petean et al., 2024). This study aims to evaluate the

evolutionary history and species boundaries of the Pacific diamond stingray, *H. dipterurus*.

First, we generated DNA sequences from *Hypanus* specimens collected in the ESP and

compared them with available conspecific sequences from the ENP populations. Then, we

provided a redescription of *H. dipterurus* from the ESP populations based on an integrative

taxonomic approach, which led to the formal resurrection of *H. brevis* (Garman, 1880).

Lastly, we evaluated historical and contemporary landing trends and discussed conservation

priorities and management strategies for sustainable *Hypanus* fisheries in the ESP.

2. MATERIALS AND METHODS

All stingray samples used in this study were obtained from Peruvian artisanal fisheries,

landings, wholesale markets, local retail markets, and from activities conducted aboard

research vessels. Consequently, no animals were sacrificed specifically for the purposes of

this study and no special permissions were required as the target species are not under

protection.

Specimens and tissue collection

The biological material used in this research was collected from January 2017 to January

2026. Four whole specimens of ESP *H. dipterurus* were collected and deposited in the

Scientific Marine Collection of the Peruvian Marine Research Institute (IMARPE) under the

catalog numbers IMARPE-020476 to IMARPE-020479. These include a juvenile caught as

bycatch off Redondo Beach in Miraflores district, collected at the landing site "*Muelle de*

Pescadores" in Chorrillos district (Peru). Three additional commercial specimens, caught at

San José Beach in Lambayeque (Peru), were bought in a wholesale fish market located in

Chiclayo city (Lambayeque, Peru) (see Table 1 and Figure 1). A small piece of tissue was

excised from the oral epithelium of the buccal cavity of each sampled specimen using a

sterile scalpel and preserved in ethanol 96 % for further DNA analysis. We also collected

tissue samples from whole and incomplete specimens of ESP *H. dipterurus* and *H. longus*

sourced from commercial fisheries, landings, aboard research vessels of IMARPE, and a

wholesale fish market in the Lambayeque, Ica, and Tumbes regions (Peru). Additionally,

tissue samples from *H. longus* were obtained from a local market in Huaquillas city

(Ecuador). Small pieces of muscle from fillets, fins, and torsos were collected and preserved

in commercial ethanol 96 % in 1.5 mL Eppendorf microtubes and stored at -15 °C until DNA

extraction. Tissue samples obtained during IMARPE research activities were labeled with the

prefix SV (without voucher specimens).

186 **TABLE 1** List of samples analyzed in this study. GenBank and BOLD accessions are written between
187 parentheses and brackets, respectively. ESP: Eastern South Pacific; ENP: Eastern North Pacific; Eastern
188 Tropical Pacific: ETP.

Species name (Population)	Specimen Code	(GenBank)/[BOLD]		Collection date (MM/DD/YY)	Collection site	Geographic coordinates	Depth (m)	Country	Reference
		COI	16S rRNA						
Hypanus dipterurus (ESP)	ITPPL-073	[ITPPL073-14]	–	2014	Chorrillos (Lima)	–	–	PERU	Unpublished
	SF33	(MH194454)	(PZ027606)	01/24/2017	Tumbes (Tumbes)	–	–	PERU	Marín et al. (2018)
	SV-0063	(PZ027578)	(PZ027607)	06/08/2024	San José (Lambayeque)	6.9 S, 80.016667 W	–	PERU	This study
	SV-0146	(PZ027579)	(PZ027608)	06/03/2025	San José (Lambayeque)	6.767000 S, 80.547833 W	51	PERU	This study
	SV-0151	(PZ027580)	(PZ027609)	06/03/2025	San José (Lambayeque)	6.926667 S, 80.241333 W	53	PERU	This study
	SV-0153	(PZ027581)	(PZ027610)	06/03/2025	San José (Lambayeque)	6.871667 S, 80.393833 W	52	PERU	This study
	IMARPE-020476	(PZ027582)	(PZ027611)	10/25/2025	Miraflores (Lima)	–	10	PERU	This study
	IMARPE-0204777	(PZ027583)	(PZ027612)	11/11/2025	San José (Lambayeque)	–	–	PERU	This study
	IMARPE-0204778	(PZ027584)	(PZ027613)	11/11/2025	San José (Lambayeque)	–	–	PERU	This study
	IMARPE-0204779	(PZ027585)	(PZ027614)	11/11/2025	San José (Lambayeque)	–	–	PERU	This study
	HbLy-02	(PZ027586)	(PZ027615)	11/11/2025	San José (Lambayeque)	–	–	PERU	This study
	HbLy-05	(PZ027587)	(PZ027616)	11/11/2025	San José (Lambayeque)	–	–	PERU	This study
	HbLy-06	(PZ027588)	(PZ027617)	11/11/2025	San José (Lambayeque)	–	–	PERU	This study
	HbLy-07	(PZ027589)	(PZ027618)	11/11/2025	San José (Lambayeque)	–	–	PERU	This study
	HbLy-08	(PZ027590)	(PZ027619)	11/11/2025	San José (Lambayeque)	–	–	PERU	This study
	HbLy-09	(PZ027591)	(PZ027620)	11/28/2025	San José (Lambayeque)	–	–	PERU	This study
	HbLy-10	(PZ027592)	(PZ027621)	11/28/2025	San José (Lambayeque)	–	–	PERU	This study
	HbLa-01	(PZ027593)	(PZ027622)	11/28/2025	Puerto Chicama (La Libertad)	–	–	PERU	This study
	HbLa-02	(PZ027594)	(PZ027623)	11/28/2025	Puerto Chicama (La Libertad)	–	–	PERU	This study
	HbPi-01	(PZ027595)	(PZ027624)	01/−−/2026	Pisco (Ica)	–	–	PERU	This study
	HbPi-02	(PZ027596)	(PZ027625)	01/−−/2026	Pisco (Ica)	–	–	PERU	This study
	HbPi-03	(PZ027597)	(PZ027626)	01/−−/2026	Pisco (Ica)	–	–	PERU	This study
	HbPi-04	(PZ027598)	(PZ027627)	01/−−/2026	Pisco (Ica)	–	–	PERU	This study
	HbPi-05	(PZ027599)	(PZ027628)	01/−−/2026	Pisco (Ica)	–	–	PERU	This study
	HbPi-06	(PZ027600)	(PZ027629)	01/−−/2026	Pisco (Ica)	–	–	PERU	This study
	HbPi-07	(PZ027601)	(PZ027630)	01/−−/2026	Pisco (Ica)	–	–	PERU	This study
	HbPi-08	(PZ027602)	(PZ027631)	01/−−/2026	Pisco (Ica)	–	–	PERU	This study
Hypanus dipterurus (ENP)	MFC048	(GU440302)	–	09/27/2001	California	–	–	USA	Unpublished
	GN1598	Mitogenome (MZ381286)		06/16/2006	Gulf of California	25.916667 N, 110.416667 W	–	MEXICO	Petean et al. (2024)
	GN5379	Mitogenome (MZ381283)		10/24/2008	Baja California Sur	27.330833 N, 112.258333 W	–	MEXICO	Petean et al. (2024)
	GN5266	Mitogenome (MZ381284)		–	Baja California Sur	25.831111 N, 111.327222 W	–	MEXICO	Petean et al. (2024)
	GN5196	Mitogenome (MZ381285)		–	Baja California	28.985833 N, 113.548056 W	–	MEXICO	Petean et al. (2024)
	GN1598	[KERRI346-08]	–	2009	Gulf of California	23.8744 N, 101.554 W	–	MEXICO	Unpublished
	SEMAR-139	[SEMAR139-10]	–	05/19/2010	Manzanillo Bay	19.064 N, 104.305 W	–	MEXICO	Unpublished
	LPZSF31	(MT311551)	–	2018	–	–	–	MEXICO	Unpublished
Hypanus longus (ETP)	GN1599	Mitogenome (MT326628)		06/16/2006	Gulf of California	25.831111 N, 111.327222 W	–	MEXICO	Petean et al. (2020)
	GN5281	Mitogenome (MT326629)		10/24/2008	Gulf of California	23.045833 N, 109.6925 W	–	MEXICO	Petean et al. (2020)
	GN5282	Mitogenome (MT326630)		10/24/2008	Gulf of California	23.045833 N, 109.6925 W	–	MEXICO	Petean et al. (2020)
	GN5283	Mitogenome (MT326631)		10/24/2008	Gulf of California	23.045833 N, 109.6925 W	–	MEXICO	Petean et al. (2020)
	HlHu-01	(PZ027603)	(PZ027632)	06/02/2017	Huaquillas (El Oro)	–	–	ECUADOR	This study
	SV-0092	(PZ027604)	(PZ027633)	04/25/2024	Nuevo Paraíso (Tumbes)	3.762233 S, 80.821883 W	–	PERU	This study
	SV-0166	(PZ027605)	(PZ027634)	10/24/2024	Punta Pico (Tumbes)	3.727633 S, 80.805683 W	12.8	PERU	This study

FIGURE 1 Sampling sites of *Hypanus* stingrays from the Eastern Pacific coast. Panel (a): Inverted purple triangles pinpoint sampling locations of *H. dipterurus* from the Eastern North Pacific (ENP) population with DNA sequences available in GenBank. Red triangles pinpoint sampling locations of *H. dipterurus* from the Eastern South Pacific (ESP) population collected in this study. Inverted pink triangles denote sampling sites of *H. dipterurus* samples whose DNA sequences were retrieved from the GenBank and BOLD databases. Panel (b): Inverted blue triangles indicate sampling locations of *H. longus* from the ENP population whose DNA sequences were retrieved from GenBank. Upright blue triangles indicate the sampling locations of *H. longus* from the ESP population collected in this study. Yellow circles represent global occurrences for *H. dipterurus* (a) and *H. longus* (b) as reported in the Global Biodiversity Information Facility (GBIF) database.

Molecular identification and data analyses

DNA isolation

Total genomic DNA was isolated using the cetyltrimethylammonium bromide (CTAB) protocol described by Zuccarello and Lokhorst (2005) with two modifications. In the organic extraction phase, the original chloroform:isoamyl alcohol (24:1) mixture was replaced with a phenol:chloroform:isoamyl alcohol in a final ratio of 25:24:1. During the subsequent DNA precipitation step, an additional 1/10 volume of sodium acetate (3 M, pH 5.2) was added to enhance precipitation efficiency. DNA quantification was estimated in an Epoch spectrophotometer (BioTek Instruments, Winooski, VT, USA). The genomic DNA was diluted to a final concentration of 50 ng/µL and used in PCR amplifications.

PCR amplification

A partial fragment (707 bp) of the mitochondrial COI gene was obtained using the universal fish primer set FishF1 (5'-TCAACCAACCACAAAGACATTGGCAC-3') and FishR1 (5'-TAGACTTCTGGGTGGCCAAAGAATCA-3') (Ward et al., 2005). For the amplification of the mitochondrial 16S rRNA gene fragment (632 bp), we used the universal primers 16Sar-L (5'-CGCCTGTTTATCAAAAACAT-3') and 16Sbr-H (5'-CCGGTCTGAACTCAGATCACG T-3') (Palumbi et al., 1991). All PCR products were amplified in a total reaction volume of 10 µL with the following master mix composition: 0.1 µL of *Taq* polymerase 5 U/µL (HiMedia, MBT060, Maharashtra, India), 0.9 µL of 10x PCR Buffer, 0.76 µL of $MgCl_2$ (25 mM), 0.5 µL of dNTPs (2.5 mM), 0.1 µL each primer (50 µM), 1 µL of genomic DNA (50 ng/µL), and 6.54 µL of ultrapure H_2O. Two different PCR amplification conditions were used, depending on the DNA marker. For the COI marker, it consisted of an initial denaturation for 5 min at 95 °C, followed for 35 cycles of denaturation step 30 s at 94 °C, annealing for 40 s optimized from a range of 54-60 °C, and elongation for 60 s at 72 °C. A final extension step was performed for 10 min at 72 °C. For the 16S rRNA, the initial denaturation was 5 min at 95 °C, followed for 35 cycles of denaturation for 45 s at 94 °C, annealing for 45 s at 58 °C, and elongation for 90 s at 72 °C. The final extension lasted 5 min at 72 °C. Successful PCR amplifications were verified in a 1.5 % agarose gel electrophoresis.

DNA sequencing

Prior to DNA sequencing, PCR products were purified using FastAP Thermosensitive Alkaline Phosphatase and Exonuclease I (Thermo Fisher Scientific, USA). Purified amplicons were sequenced bi-directionally using the same primer sets and the BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems, Waltham, Massachusetts, USA) on a ABI PRISM 3500 Genetic Analyzer (Applied Biosystems, Hitachi, Foster City, CA,

USA). All sequencing reactions were performed at the Laboratory of Genetics, Physiology,

and Reproduction of the National University of Santa. Electropherograms were visually

revised using MEGA 7 (Kumar et al., 2016), consensus sequences were obtained by aligning

forward and reverse complemented sequences, and primer positions were removed.

Estimation of genetic diversity and population differentiation

We used Arlequin v3.5 (Excoffier & Lischer, 2010) to quantify genetic variation within each

analyzed *H. dipterurus* population (ENP and ESP) and to estimate diversity metrics. These

include haplotype number, haplotype diversity "*Hd*", nucleotide diversity "π". The

relationships among haplotypes resulting from the COI and COI + 16S rRNA datasets were

assessed using Median-joining networks in HaplowebMaker tool (Spöri & Flot, 2020). To

assess deviations from selective neutrality and infer historical demographic events, we

calculated Fu's F_S and Tajima's D in both datasets (COI and COI + 16S rRNA) using

Arlequin. Statistical significance was evaluated with 10,000 coalescent simulations, with

threshold significance levels set at $P < 0.02$ (F_S) and $P < 0.05$ (D).

To quantify the degree of genetic divergence between the *H. dipterurus* samples collected

from ENP and ESP populations, we used the Analysis of Molecular Variance (AMOVA)

implemented in Arlequin. Pairwise fixation indices (Φ_{CT}, Φ_{SC}, and Φ_{ST}) were estimated using

the K2P distance model (Kimura, 1980) for the COI and concatenated COI + 16S rRNA

datasets. Accordingly, total variance was grouped into the following components: Φ_{CT}:

variation among groups, Φ_{SC}: variation among populations within groups, and Φ_{ST}: variation

among all populations. Two populations were defined: 1) the ENP population, comprising 8

DNA sequence samples sourced from GenBank, whose geographic origin indicated

California in the USA, and Gulf of California and Manzanillo Bay in Mexico; 2) the ESP

population, represented by 27 DNA sequences of which 25 sequences were determined in

this study (Ica region n = 8, Lambayeque region n = 14, La Libertad region n = 2, and Lima

region n = 1). Two additional Peruvian sequences were sourced from GenBank (Tumbes

region n = 1) and BOLD (Lima region n = 1) databases.

Bayesian phylogenetic analysis

The phylogenetic relationships among the DNA sequences of *Hypanus* species were inferred

using two independent Bayesian analyses based on single COI locus dataset and a

concatenated COI + 16S rRNA multilocus dataset. The partitioned Bayesian approach on the

multilocus dataset was conducted to enhance the phylogenetic signal of the 16S rRNA gene

and improve the reliability of species delimitation (Nylander et al., 2004). Although a

preliminary single-locus analysis of the 16S rRNA gene recovered ESP *H. dipterurus* and

ENP *H. dipterurus* as separate clades, the relatively low genetic distance (K2P: 1.7 %)

indicates a recent divergence between these two clades. Such shallow genetic divergences are

often associated with limited phylogenetic signal (Philippe et al., 2011). This may result in

weakly supported topology or overly confident statistical nodal support (Chen et al., 2017).

The single (COI) and multilocus (COI + 16S rRNA) datasets consisted of 55 and 50

sequences, respectively. These included 25 sequences for ESP *H. dipterurus* (Peru) and three

sequences for *H. longus* (Ecuador and Peru) determined in this work. They were analyzed

along with all available conspecific sequences from Mexico and the USA that were retrieved

from BOLD and GenBank databases. Additionally, four sequences of *H. say* (sister taxon of

H. dipterurus) and *H. americanus* (sister taxon of *H. longus*), and two sequences of *H.*

sabinus (sister taxon to all *Hypanus* species) (Mejía-Falla et al., 2025; Petean et al., 2024;

Rosenberger, 2001), were retrieved from the databases and included in the analyses. Two

bluespotted stingrays (*Neotrygon indica* Pavan-Kumar, Kumar, Pitale, Shen & Borsa, 2018

and *N. kuhlii*, subfamily Neotrygoninae) and the Daisy stingray (*Fontitrygon margarita*

(Günther, 1870), subfamily Urogymninae) were used as outgroups. DNA sequences from the

COI and 16S rRNA genes were aligned independently and trimmed using MEGA 7. Pairwise

genetic distances were calculated using the Kimura 2-parameter model (K2P, Kimura, 1980),

as implemented in MEGA 7. We assessed signals of nucleotide saturation in the protein-

coding COI dataset using DAMBE 7 (Xia, 2018). Next, the best-fit model of evolution for

each aligned matrix were assessed with jModelTest2 (Darriba et al., 2012) under the

corrected Bayesian Information Criterion (BICc). The best evolutionary model for the COI

and 16S rRNA gene datasets were HKY + G and HKY + I, respectively.

The Bayesian phylogenetic analyses were conducted in MrBayes 3.2.7 (Ronquist et al., 2012),

as implemented on the CIPRES Science Gateway 3.3 server (Miller et al., 2010). The

analyses were conducted with the data partitioned by gene and codon position. In both

datasets, the COI and the concatenated COI + 16S rRNA, the setup parameters consisted of

two independent runs of four Markov chains each, conducted for 10,000,000 generations.

Trees were sampled every 1,000 generations and the first 25 % of the sampled trees were

discarded as burn-in. Quality check of output files consisted on an Average Standard

Deviation of Split Frequencies (ASDSF) below 0.01 and Effective Sample Sizes (ESS) for all

parameters exceeding 200. The ESS were verified in Tracer v1.7.2 (Rambaut et al., 2018).

The consensus trees were visualized with FigTree version 1.4.4. (http://tree.bio.ed.ac.uk/soft

ware/figtree/).

304 Calibrated molecular dating analysis

To estimate the divergence times among target *Hypanus* species, we conducted a relaxed

molecular dating analysis based on the multilocus (COI + 16S rRNA) dataset. Since the two

previous phylogenetic analyses (single and multilocus) consistently recovered ESP *H.*

dipterurus and ENP *H. dipterurus* as distinct monophyletic clades, for the final molecular

dating we applied a more general model across the partitions: the GTR + I + G. To ensure

taxonomic consistency with previous related studies, the following constraints were enforced

a priori: root, crown *Hypanus*, ENP *H. dipterurus* monophyly, ESP *H. dipterurus* monophyly,

ENP *H. dipterurus*/ESP *H. dipterurus* sister taxa, Panama Isthmus node 1 (*H. say*/*Hypanus*

clade ancestor), Panama Isthmus node 2 (*H. longus/H. americanus*), and Dasyatinae node

(Aschliman et al., 2012).

Node ages were calibrated using and "offsetlognormal" or "uniform" distribution based on

established fossil or geological events. These include a) Root Dasyatidae: calibrated with and

offsetlognormal(94.0, 100.0, 0.5) million years ago (Ma), to establish the minimum age of the

Dasyatidae family based on *Dasyatis* fossil teeth record from the Cenomanian stage (Late

Cretaceous)   (Landemaine,   1991).   b)   The   Dasyatinae   node: calibrated   via

offsetlognormal(57.4, 64.9, 0.5) Ma, based on the *Dasyatis*/*Urogymnus* split age estimated by

Aschliman et al. (2012) and applied following the updated classification of Last et al. (2016b).

c) Crown *Hypanus*: calibrated with offsetlognormal(33.0, 40.0, 1.0) Ma, which corresponds

to the divergence age of the *D. brevis* (currently *H. dipterurus*) during the Early Oligocene

estimated by Aschliman et al. (2012), aligned with the taxonomic revision by Last et al.

(2016b). d) Two transisthmian splits between *H. americanus*/*H. longus* and *H. say*/ENP *H.*

dipterurus: each calibrated with offsetlognormal(3.1, 3.5, 0.5) Ma. We applied a lognormal(-

5.2, 1.0) prior to the mean clock rate to model the typical slow mitochondrial mutation rate of

0.2-0.3 % per million years of chondrichthyan taxa (Martin et al., 1992).

The molecular clock analysis was conducted using the same Bayesian settings as in the

phylogenetic inference described in the previous subsection, with the exception that the

MCMC run was increased to 50,000,000 generations to reach stationary.

Morphometric and meristic analysis

Morphometric and meristic analyses were conducted on four complete, fresh specimens of *Hypanus* (Table 1), following primarily Petean et al. (2020), with additional measurements taken from Jordan and Gilbert (1880). All body measurements were taken in a straight line using a caliper with a precision of 0.1 mm; measurements exceeding 170 mm were taken using dividers and read with a steel ruler with a precision of 1 mm. After measurements, specimens were fixed in 10 % formalin, rinsed, preserved in 70 % ethanol, and deposited in the IMARPE Scientific Collection under the catalog numbers IMARPE-020476, IMARPE-020477, IMARPE-020478, and IMARPE-020479. Comparative morphometric data were obtained from the original descriptions of *Trygon brevis* by Garman (1880), considering the two syntypes under the collection number MCZ Ichthyology S-422, one female and one male from Paita, Peru; and from Lamilla et al. (1995). We compared with morphometric data of two syntypes of *Dasybatis dipterurus* from Jordan and Gilbert (1880) description; and with those given by Beebe and Tee-Van (1941) and Nishida and Nakaya (1990). Measurements are expressed in millimeters (mm) and in percentages (%) of the disc width (% DW). In some cases, measurements originally expressed in inches were converted to millimeters. When necessary, measurements were also standardized from proportions of total length (TL) or disc length (DL) to ensure comparability. Taxonomic validity and nomenclature follow Fricke et al. (2025), while collection codes are based on Sabaj (2020). All specimens from the IMARPE fish collection were also used for species identification through genetic analyses.

Fishery data analysis

A database of chondrichthyan landings from 1996 to 2024, geolocated by fishing zones, was provided by the IMARPE, in response to a request for access to public information under Peruvian transparency provisions. In total, the database comprised 182,282 landing records,

classified by geolocation (latitude and longitude), fishing gear, common and scientific names,

year, and landed volume. Records of batoid species, particularly those classified as *Dasyatis*

brevis and the generic scientific designation *Hypanus* sp., were filtered to describe small-

scale fishing activity targeting species under analysis. Consequently, both labels were

extracted for descriptive purposes. Additionally, fishing areas of the small-scale fisheries

were classified into four zones by latitude: Tropical North, North, Center, and South, based

on IMARPE's classification available in the Peruvian Atlas of Small-Scale Fisheries (Marín-

Soto et al., 2017). Additionally, the distance to the coast of the reported catches was

estimated using the Fenix R Studio package (Marín-Abanto, 2018). The data were used to

estimate trends in landings and to show the marine areas within the Peruvian Sea with the

highest occurrence of this species' fishing activity. Furthermore, to discover its economic

value, annual average off-vessel prices and total monetary landing values were estimated

using open-access data collected and stored by IMARPE (2025). Nearly 2,108 records of

nominal off-vessel prices for the research target species from 2009 to 2024 were obtained.

The nominal off-vessel prices were converted from Peruvian soles (PEN) to US dollars

(USD) using monthly PEN/USD exchange rates published by the Peruvian Central Bank

(BCRP, 2025a). Additionally, nominal prices were converted to real prices by deflating them

using the consumer price index (CPI), published monthly by the BCRP from January 1991

onwards, with December 2021 as the baseline (i.e., IPC = 100) (BCRP, 2025b).

3. RESULTS

DNA sequencing

We successfully obtained genomic DNA, PCR products, and DNA sequences from all

specimens collected from the ESP populations. These include 25 sequences for *H. dipterurus*

and three sequences for *H. longus*. After the removal of the primer sequences, the DNA

sequences for the COI and 16S rRNA markers were 655 bp and 590 bp in length, respectively.

All DNA sequences generated in this research were assigned the following GenBank

accessions: from PZ027578 to PZ027605 for the COI gene and from PZ027606 to PZ027634

for the 16S rRNA gene.

Genetic diversity indexes and neutrality tests

Hd for both the single locus and multilocus datasets, was high (0.8571 and 0.8333,

respectively) in *H. dipterurus* samples from the ENP population. Conversely, the *Hd* in the

ESP population was zero for the COI dataset due to the fixation of a single haplotype, while

the combined COI + 16S rRNA dataset yielded an *Hd* of 0.1477. π followed a similar pattern,

being higher in the ENP (COI: 0.003461, COI + 16S rRNA: 0.000906), and notably lower in

the ESP population (COI: 0.0, COI + 16S rRNA: 0.000134) (see Table 2). Neutrality test for

the ENP yielded negative values for both the Fu's F_S and Tajima's *D*. Although these values

were not significant in all cases, their negative trend deviates from expectations of a neutral

population model and may reflect a recent population expansion. Neutrality test from the ESP

could not be calculated for the COI gene due to the lack of polymorphism. However, the

combined COI + 16S rRNA dataset, which contained two haplotypes, resulted in a non-

significant Fu's F_S (-0.3167, P = 0.1648) and a Tajima's *D* of 0.000 (P = 1.000). While these

values suggest no significant departures from mutation-drift equilibrium, this is likely an

artifact of the extremely low genetic diversity (e.g., one COI haplotype for 27 specimens),

which limits the statistical power to detect demographic shifts. This lack of variation is

consisted with a founder effect associated with a recent speciation event or a severe historical

bottleneck.

TABLE 2 Summary results of genetic diversity indices and Fu's *F s*and Tajima's *D* neutrality tests for single locus dataset (COI) and multilocus dataset (COI + 16S rRNA) of *H. dipterurus*. ENP: Eastern North Pacific population, ESP: Eastern South Pacific population. Genetic diversity includes: number of haplotypes (h), haplotype diversity (*Hd*), and nucleotide diversity (π). Sample size: n. N.A.: Not Applicable.

Population	Marker	n	h	$Hd \pm$ S.E.	π	Fu's F_S*	Tajima's D**
ENP (Mexico- USA)	COI	8	5	0.8571 ± 0.1083	0.003461± 0.002537	-1.40322 (P = 0.08330)	-1.06428 (P = 0.16190)
	COI + 16S rRNA	4	3	0.8333 ± 0.2224	0.000906 ± 0.000897	-0.88730 (P = 0.08860)	-0.70990 (P = 0.28510)
ESP (Peru)	COI	27	1	0.0000 ± 0.0000	0.0000 ± 0.0000	0.0000 (P = N.A.)	0.0000 (P = 1.0000)
	COI + 16S rRNA	26	2	0.1477 ± 0.0888	0.000134 ± 0.000219	-0.31678 (P = 0.16480)	0.0000 (P = 1.0000)

414 * P < 0.02 significant.
415 ** P < 0.05 significant
416

Haplotype network analysis

The median-joining haplotype network analyses for *H. dipterurus* samples from the ENP and

ESP populations are presented in Figure 2. Our haplotype network findings from the single

locus and concatenated multilocus datasets revealed a clear genetic divergence between both

the ENP and ESP populations. In the COI haplotype network (top panel in Figure 2a), five

haplotypes were observed in samples from the ENP population (Mexico n = 7; USA n = 1),

while a single private haplotype represented the ESP population (Peru, n = 27). The branch

connecting the unique haplotype of the ESP haplogroup (H1b) to the closest haplotype from

the ENP (H1d, Gulf of California) displayed a genetic distance of 18 stepwise mutations. The

largest genetic distance (23 nucleotides) between both haplogroups was detected between

haplotypes H1b (Peru) and H5d (Gulf of California). The haplotype network obtained using

the concatenated COI + 16S rRNA dataset (bottom panel in Figure 2a) resulted in a dominant

haplotype H1b for the ESP population (Peru, n = 24), while a single satellite haplotype H2b

was found in only two individuals. These two ESP haplotypes were separated by a single

mutational step, represented by a deletion that was observed in the 16S rRNA gene from two

specimens collected in Pisco (Ica region, GenBank accessions PZ027624 and PZ027626).

The ESP haplogroup was distanced by 30 mutational steps to closest haplotype H1d in the

ENP haplogroup, which comprised a total of three haplotypes from four specimens collected

in Mexico. This substantial genetic distance and lack of shared haplotypes between the two

haplogroups (ESP and ENP) strongly suggest that they represent isolated evolutionary units.

On the other hand, the haplotype networks of both datasets (COI and COI + 16S rRNA) from

H. longus suggest the existence of a single lineage across the Eastern Pacific (top and bottom

panels in Figure 2b, respectively). A unique dominant haplotype was detected in each

population (ENP: Mexico, ESP: Ecuador and Peru), with three and five stepwise mutations

between each haplotype in the single locus and multilocus datasets, respectively.

443
FIGURE 2 Median joining network analysis illustrating the genetic relationships and geographical distribution
of two *Hypanus* lineages using DNA sequences from samples collected in the Eastern North Pacific (ENP) and
Eastern South Pacific (ESP) populations. Panel (a): Haplotype network of *H. dipterurus* based on COI
sequences (above) and concatenated sequences from COI + 16S rRNA (below). Panel (b): Haplotype network
of *H. longus* based on COI sequences (above) and concatenated sequences from COI + 16S rRNA (below).
Circles (nodes) represent unique haplotypes, where its size is proportional to the frequency of that haplotype in
the sampled population. Perpendicular bars across the lines connecting haplotypes represent single mutational
steps. Colors denote the geographic origin of the samples. In the ENP population: yellow color for the Gulf of
California (GC) and green for Manzanillo Bay (MB), both in Mexico. Electric lime color for California (CA) in
the USA. In the ESP population: raspberry pink for Huaquillas (HU) in Ecuador. Purple color for the Ica region
(IC), electric blue color for the Lambayeque region (LY), neon blue for the La Libertad region (LL), pale blue
for the Lima region (LI), lavender blue for the Tumbes region (TU), all located in Peru.

456
457

Analysis of Molecular Variance (AMOVA)

The AMOVA results (Table 3) revealed significant differences in the molecular data for both datasets. For the single locus (COI) dataset, a vast majority of the variation (97.54 %) occurred between both *H. dipterurus* groups (ENP vs ESP), while only a minor variation was observed within populations (2.49 %). Similarly, in the combined multilocus (COI + 16S rRNA) dataset, a higher level of variation (99.25 %) was detected between groups. These results strongly suggest a genetic isolation based on geographic location of the ENP and ESP groups. Fixation indices further supported the genetic divergence between ENP and ESP populations. The Φ_{CT} values, which measure differentiation among groups, were notably high at 0.97 (P = 0.02) for COI and 0.99 (P = 0.16) for the combined data. Additionally, consistently high Φ_{ST} values (0.98 and 0.99, P = 0.00) confirm complete genetic isolation and lack of gene flow between the analyzed groups.

TABLE 3 Results of the Analysis of Molecular Variance (AMOVA) and fixation indices (Φ_{CT}, Φ_{SC}, and Φ_{ST}) based on single locus (COI) and multilocus (COI + 16S rRNA) datasets from DNA sequences of *Hypanus dipterurus*. Two groups were compared: ENP (Eastern North Pacific: Mexico and USA) population group and ESP (Eastern South Pacific: Peru) population group.

Dataset	Source of Variation	d.f.	Variance Components	% of Variation	Fixation Indices	P-value
COI (Single)	Among groups	1	9.46466 Va	97.54	$\Phi_{CT} = 0.97536$	0.02053
	Among populations within groups	6	-0.00257 Vb	-0.03	$\Phi_{SC} = -0.01077$	0.65396
	Within populations	20	0.24167 Vc	2.49	$\Phi_{ST} = 0.97510$	0.00000
COI + 16S rRNA (Combined)	Among groups	1	15.22730 Va	99.25	$\Phi_{CT} = 0.99249$	0.16618
	Among populations within groups	4	-0.00976 Vb	-0.06	$\Phi_{SC} = -0.08466$	0.36364
	Within populations	24	0.12500 Vc	0.81	$\Phi_{ST} = 0.99185$	0.00000

Fixation indices significant at P < 0.05

Bayesian phylogenetic analysis

The Bayesian phylogenetic analyses of both, the single (Figure 3a) and multilocus (Figure 3b) datasets, further supports the conclusions drawn from the haplotype network and AMOVA findings. In both phylogenetic trees, DNA sequences from ENP and ESP *H. dipterurus* populations were recovered in two discrete monophyletic clades at the top of the trees with strong nodal support. The sister relationship between both *H. dipterurus* populations exhibited a COI genetic distance ranging from 3.6 % (± 0.008) to 4.6 % (± 0.010), with an average value of 4 % (± 0.009) (Table S1). This genetic distance range is substantially higher than the interspecific genetic threshold reported in congeneric species (Petean et al., 2024). Our phylogenetic findings, combined with the haplotype network and AMOVA results, provide robust evidence that *H. dipterurus* populations from ENP and ESP constitute distinct evolutionary lineages.

Conversely, the results of the Bayesian phylogenetic analyses support the existence of a single Pacific lineage for *H. longus*. In both Bayesian phylogenetic trees, corresponding to the COI and concatenated COI + 16S rRNA dataset, all *H. longus* samples were recovered within a single monophyletic group with high statistical support (Figure 3). They displayed low COI intraspecific genetic distance, ranging from 0 to 0.6 % (± 0.003), which is lower than the standard interspecific DNA barcoding gap reported in species from this genus (Petean et al., 2024).

FIGURE 3 Bayesian phylogenetic trees showing the evolutionary relationships among stingrays in the genus *Hypanus*. Panel (a) shows the phylogenetic tree constructed using a single locus COI marker (substitution model: HKY + G) and the tree from Panel (b) utilized the concatenated information from the COI + 16S rRNA genes (substitution model: HKY + I). Each label shows the species scientific name, voucher name, geographical origin (when available), and GenBank (between parentheses) or BOLD (between brackets) accessions numbers. The numbers next to each node indicate the Bayesian Posterior Probability. To distinguish *H. dipterurus* from the Eastern North Pacific (ENP) and Eastern South Pacific (ESP) populations, the clade grouping all samples from the ESP population is highlighted in blue color and shaded in sky blue color. The *H. longus* clade is highlighted in red color. *Hypanus sabinus*, the sister taxa to all *Hypanus* group, was used as an internal outgroup. Two bluespotted stingrays (*Neotrygon kuhlii* and *N. indica*) and the Daisy stingray (*Fontitrygon margarita*) were used as external outgroups.

Calibrated molecular clock analysis

The calibrated Bayesian molecular clock analysis dated the most recent common ancestor (MRCA) of the Dasyatidae lineage (subfamilies Dasyatinae, Neotrygoninae, and Urogymninae) at approximately 120.89 Ma, during the Early Cretaceous (Figure 4). The major Dasyatinae clade divergence occurred around the Cretaceous-Paleogene (K-Pg) boundary, at approximately 64.88 Ma. The crown *Hypanus* radiation started about 39.98 Ma, during the Late Eocene. The divergence of the common ancestor of the Atlantic *H. americanus* and *H. say* occurred approximately 24.62 Ma during the Early Miocene. The most recent *Hypanus* divergences occurred during the Late Pliocene. These include two transisthmian divergence events that resulted in the formation of the geminate species *H. say* (Atlantic) and *H. dipterurus* (Pacific) (3.42 Ma), and *H. americanus* (Atlantic) and *H. longus* (Pacific) (3.43 Ma). Finally, the most recent speciation event occurred within the diamond stingray *H. dipterurus*, when the ESP *H. dipterurus* lineage branched off from its geminate from the ENP, at approximately 3.09 Ma (Figure 4).

Based on the findings of this research, which strongly support that *H. dipterurus* populations

from the ENP and ESP represent isolated evolutionary units, we propose that the ESP lineage

be formally recognized as *H. brevis*. This taxonomic assignment follows Garman's (1880)

original description based on specimens from Paita, northern Peru. The revalidation of *H.*

brevis increases the number of recognized species to 12 and brings the total number of

distinct *Hypanus* evolutionary units to 15 (Table 4), including the three molecular

evolutionary units (*H.* aff. *americanus, H.* aff. *guttatus, and H.* aff. *say*) identified by Petean

et al. (2024).

TABLE 4 Current accepted species and latest taxonomic changes in members of the genus *Hypanus*.
Distribution: EP (Eastern Pacific), ESP (Eastern South Pacific), ENP (Eastern North Pacific), ETP (Eastern
Tropical Pacific), WA (Western Atlantic), EA (Eastern Atlantic), ECA (Eastern Central Atlantic), WCA
(Western Central Atlantic), WNA (Western North Atlantic), WSA (Western South Atlantic), Gulf of Mexico
(GM), NB (Northern Brazil), SB (Southern Brazil), and CS (Caribbean Sea).

Taxonomy	Common name	Description Year	Distribution	Original name	Latest taxonomic change	Reference	IUCN status
Genus							
Hypanus Rafinesque, 1818	Stingrays	1818	EP, WA, EA	*Hypanus* Rafinesque, 1818	Resurrected from synonymy with *Dasyatis*	Last et al. (2016b)	2 as CR (18.1 %) 1 as EN (9.1 %) 3 as VU (27.3 %) 3 as NT (27.3 %) 1 as LC (9.1 %) 1 as NE (9.1 %)
Validated species by the present work							
Hypanus brevis (Garman, 1880)	Peruvian diamond stingray	1880	ESP	*Trygon brevis* Garman, 1880	Resurrected from synonymy with *H. dipterurus*	This study	VU (needs reassessment)

Validated species in the literature							
Hypanus guttatus (Bloch & Schneider, 1801)	Longnose stingray	1801	WCA to SB	*Raja guttata* Bloch & Schneider, 1801	Moved from the genus *Dasyatis* (*D. guttata*)	Last et al. (2016b)	NT
Hypanus say (Lesueur, 1817)	Bluntnose stingray	1817	WA	*Raja say* Lesueur, 1817	Moved from the genus *Dasyatis* (*D. say*)	Last et al. (2016b)	NT
Hypanus rudis (Günther, 1870)	Smalltooth stingray	1870	ECA	*Trygon rudis* Günther, 1870	Moved from the genus *Dasyatis* (*D. rudis*)	Last et al. (2016b)	CR
Hypanus dipterurus (Jordan & Gilbert, 1880)	Diamond stingray	1880	ENP	*Dasybatis dipterura* Jordan & Gilbert, 1880	Moved from the genus *Dasyatis* (*D. dipterura*)	Last et al. (2016b)	VU (needs reassessment)
Hypanus longus (Garman, 1880)	Longtail stingray	1880	ETP	*Trygon longa* Garman, 1880	Moved from the genus *Dasyatis* (*D. longa*)	Last et al. (2016b)	VU
Hypanus sabinus (Lesueur, 1824)	Atlantic stingray	1824	WNA	*Trygon sabina* Lesueur, 1824	Moved from the genus *Dasyatis* (*D. sabina*)	Last et al. (2016b)	LC
Hypanus americanus (Hildebrand & Schroeder, 1928)	Southern stingray	1928	WNA, WCA and CS	*Dasyatis americana* Hildebrand & Schroeder, 1928	Moved from the genus *Dasyatis* (*D. americana*)	Last et al. (2016b)	NT
Hypanus geijskesi (Boeseman, 1948)	Wingfin stingray	1948	WCA, NB	*Dasyatis geijskesi* Boeseman, 1948	Moved from the genus *Fontitrygon*	Petean et al. (2024)	CR
Hypanus marianae (Gomes, Rosa & Gadig, 2000)	Brazilian large-eyed stingray	2000	WSA (Brazil)	*Dasyatis marianae* Gomes, Rosa & Gadig, 2000	Moved from the genus *Dasyatis* (*D. marianae*)	Last et al. (2016b)	EN
Hypanus berthalutzae Petean, Naylor & Lima, 2020	Lutz's stingray	2020	WSA (Brazil)	*Hypanus berthalutzae* Petean, Naylor & Lima, 2020	New discovery. It was split from *H. americanus*	Petean et al. (2020)	VU
Hypanus rubioi Mejía-Falla, Navia, Cardeñosa & Tavera, 2025	Pointed-nose Pacific stingray	2025	ETP	*Hypanus rubioi* Mejía-Falla, Navia, Cardeñosa & Tavera, 2025	New discovery	Mejía-Falla et al. (2025)	NE
Molecular evolutionary units (aff.)							
Hypanus aff. *americanus*	–	–	WNA, WCA, CS	–	–	Petean et al. (2024)	–
Hypanus aff. *guttatus*	–	–	WCA	–	–	Petean et al. (2024)	–
Hypanus aff. *say*	–	–	GM	–	–	Petean et al. (2024)	–

Resurrection of *Hypanus brevis* (Garman, 1880)

Trygon brevis Garman, 1880: 171. Description.

Dasibatis brevis Garman (1883: 70).

Dasybatus brevis. In part Garman (1913: 396). Specimens of Paita, Peru, pl. 32, figs. 5-6.

Dasyatis brevis Chirichigno (1974: 75, figs. 67-68); Lamilla et al. (1995: 23-27); Chirichigno

& Vélez (1998: 65, figs. 11-12); Chirichigno & Cornejo (2001: 67).

Hypanus dipterurus. Almendras et al. (2025), specimens of Peru and Chile.

Syntypes collected in Paita, Peru (MCZ Ichthyology S-422; Georeferenced as 5° 9' S, 81° 8'

W or -5.1583333°, -81.141666°; collection date May 5, 1872; collector Hassler expedition).

Held at the Museum of Comparative Zoology, Harvard University.

https://mczbase.mcz.harvard.edu/guid/MCZ:Ich:S-422

Although Garman (1880) did not mention collection numbers, it is assumed that there are at

least two syntypes. He indicated Paita (Piura, Peru) as the locality of origin for at least one of

the specimens (female) mentioned in his description and confirmed the provenance of the two

measured specimens (male and female) as Peru (Garman, 1883). He made no reference to

type material from California.

Diagnosis

A large-sized diamond-shaped species of *Hypanus* (attaining at least 900 mm DW) with the

following combination of characters: body slightly wider than long; snout pointed but not

elongated, rostral length 18.5–20.7 % DW; anterolateral margins of disc nearly straight,

forming an angle of 124–130°; disc length 82.5–87.0 % DW; internostril width 8.5–9.7 %

DW; oral papillae five, arranged in a transverse row on the floor of the mouth, with three

well-developed medial papillae and one minute lateral papilla on each side; tail whiplike, less

than 1.25 times disc width (1.4–1.5 times disc length), bearing a long, slender caudal spine

and a short ventral caudal fin fold (25.7–34.6 % DW); dorsal disc generally smooth in

juveniles, adults with scattered scapular denticles and enlarged denticles anterior to the

caudal spine; coloration brown to olive-brown dorsally, lacking spots.

Description

Disc wider than long, disc length equivalent to 82.5–87 % of disc width (DW). Anterolateral

margins of disc nearly straight, posterior margins convex. Snout pointed but not protruding,

forming an angle of 124–130°. Upper and lower jaws with small, blunt teeth. Oral papillae

five, arranged in a single transverse row on the floor of the mouth; three medial papillae large

and well developed, and one-minute lateral papilla on each side, widely separated from the

medial group. Pelvic fins broad, with truncate margins, slightly rounded apices, and rounded

angles.

Tail moderately long, with a length equivalent to 119–123 % of disc width; broad and

slightly depressed at the base, progressively tapering toward the caudal spine. Dorsal and

ventral folds present on tail; dorsal fold low and short, ventral fold higher and longer than

dorsal, representing 25.73–34.63 % of disc width. Ventral fold originates at level of caudal

spine basis, dorsal fold at level of caudal spine tip. Both folds terminate at approximately the

same level along the tail, from where the tail begins to compress, subsequently narrowing

into a filament.

Dorsal surface of disc smooth. The largest specimen (IMARPE-20479), a female of 540 mm

DW, exhibits an irregular row of scattered denticles extending from the nuchal region to the

origin of the caudal spine, as well as a short row on each shoulder, with up to five denticles

per side. Juvenile specimens lack denticles. Additionally, a large incomplete specimen

collected at ~6° S (SV-0063) showed more than 14 prominent, broad-based denticles on the

tail prior to the caudal spine (see Figure 5).

Comparison

Hypanus brevis is distinguished from its Eastern Pacific congeners *H. longus* and *H. rubioi* by having a very short tail, measuring 1.4–1.5 times disc length vs 2.4 times in *H. longus* and 2.4–3.1 times in *H. rubioi*. Tail length in *H. brevis* is similar to that of *H. dipterurus*, which ranges from 1.1 to 1.4 times disc length (Jordan & Gilbert 1880; Beebe & Tee-Van 1941; Mejía-Falla et al., 2025). *Hypanus brevis* shares with *H. longus* a disc that is wider than long, with disc length values ranging from 82.5 to 87.0 % DW in *H. brevis* and 81.6 to 86.8 % DW in *H. longus*. In contrast, *H. brevis* differs from *H. dipterurus*, in which disc length represents a higher proportion of disc width, ranging from 88.9 to 94.9 % DW. In *H. rubioi*, disc length is slightly greater, with values ranging from 89.6 to 98.2 % DW, resulting in a disc that is nearly as long as wide. *Hypanus brevis* does not have a pointed nor elongated rostrum, with rostral length ranging from 18.5 to 20.7 % of disc width (DW), compared to 29.5-32.1 % DW in *H. rubioi*. In this respect, *H. brevis* shares a similar rostral length with *H. longus* (16.4-19.5 % DW) and *H. dipterurus* (15.9–21.8 % DW). However, the rostral angle is markedly wide in *H. brevis* (124–130°) and *H. dipterurus*, slightly less wide in *H. longus*, whereas it is acute in *H. rubioi*. *H. brevis* shares with *H. dipterurus* a small ventral caudal fin fold, 25.7-34.6 % DW and 28.4-33.3 % DW, respectively. In contrast, *H. longus* and *H. rubioi* have a longer ventral caudal fin fold, measuring 45–52.4 % DW and 38–60.8 % DW, respectively. The internostril width is smaller in *H. brevis* (8.5–9.7 % DW) compared to *H. dipterurus* (13.6-18.6 % DW). In contrast, values for *H. longus* (9.4–10.4 % DW) and *H. rubioi* (9.1–10.7 % DW) are similar to, though slightly higher than, those observed in *H. brevis*.

Color in life

Dorsal surface uniformly brown, dark brown, or olive-brown, becoming slightly reddish near

the margins (see Figure 6); ventral surface white, with the posterior margins of the pectoral

and pelvic fins brown. Juveniles exhibit a slightly more violaceous coloration on the dorsal

surface.

FIGURE 6 *Hypanus brevis* (Garman, 1880) Pictures illustrating the coloration observed in recently caught/live
specimens. Panels (a) and (b): female specimens (SV-0146 and SV-0151, respectively) caught in San José
(Lambayeque region, Peru); (c): specimen collected in Pisco (Ica region, Peru). Photographic credits: (a) and (b)
F. Zavalaga; (c) J. Quiñones (used with written permission).

TABLE 5 Morphometric and meristic characteristics of *Hypanus brevis* (Garman, 1880) based on specimens collected in the southern Pacific Ocean off Peru, expressed in millimeters and as percentages of disc width (%DW). Data for the syntypes are taken and compared from Garman (1880), specimens MCZ S-422 from Paita, Peru. Additional comparisons are made with data from Garman (1913); Lamilla et al. (1995); and from the syntypes of *H. dipterurus* Jordan and Gilbert (1880), from Beebe and Tee-Van (1941), and Nishida and Nakaya (1990). *Data corrected from the original publication. (–) (?) Data not reported.

Species	This study, *Hypanus brevis* n=4								*H. brevis* (Garman, 1880) (Syntypes)				*H. brevis* (Lamilla et al., 1995)				*H. dipterurus* (Jordan & Gilbert, 1880)				*H. dipterurus* (Beebe & Tee-Van, 1941)			*H. dipterurus* (Nishida & Nakaya, 1990)	
Museum catalogue N°	IMARPE-20479		IMARPE-20478		IMARPE-20477		IMARPE-20476		MCZ S-422		MCZ S-422		IZUA-PM 1433		w/n		No.1		No.2		24,943		24,995	LACM 31759-16	
Measurements	mm	%DW	mm	%DW	mm	%DW	mm	%DW	mm	%DW	mm	%DW	mm	%DW	mm	%DW	mm	%DW	mm	%DW	mm	%DW	mm	mm	%DW
Total length TL	1140	211.11	614	206.04	658	204.98	483	200.41	-	-	-	-	510	205.60	-	-	-	-	-	-	1257	175.44	702	466.9	207.4
Angle of the rostrum (°) AR	130	-	128	-	129	-	124	-	-	-	-	-	125	-	-	-	-	-	-	-	-	-	-	-	-
Disc width DW	540	100.00	298	100.00	321	100.00	241	100.00	457.2	100.00	233.7	100.00	248	100.00	600	100.00	231.1	100.00	237.5	100	716.49	100.00	-	225.1	100.0
Disc length DL	470	87.04	253	84.90	265	82.55	206	85.48	431.8	94.44	205.8	88.06	218	87.90	505	84.20	222.2	94.87	215.9	90.91	672.50	93.86	331.34	200.1	88.9
Mouth to tail MtT	367	67.96	255	85.57	222	69.16	167	69.29	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pre-caudal length PCL	462	85.55	253	84.90	267	83.18	205	85.06	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Snout to cloaca length StC	420	77.77	220	73.83	240	74.77	187	77.59	-	-	-	-	194	78.20	435	72.50	155.57	66.40	151.1	63.63	-	-	-	176.5	78.4
Cloaca to caudal tip length CtCT	732	135.55	387	129.87	430	133.96	288	119.50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pre-spine length PSL	605	112.04	320	107.38	347	108.10	265	109.96	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	253.7	112.7
Caudal length CL	663	122.77	360	120.81	398	123.99	289	119.92	584.2	127.77	304.8	130.42	315	127.00	-	-	311.15	132.82	315	132.6	734.09	102.46	370.66	-	-
Caudal width at spine CSW	16	2.96	8	2.68	7	2.18	6	2.49	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tail origin to spine origin length TSL	160	29.63	85	28.52	93	28.97	71	29.46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Caudal width at P2 end CW	39	7.22	18.5	6.21	17	5.30	14	5.81	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Caudal height at pelvic fin end CH	24	4.44	12	4.03	12	3.74	9.5	3.94	-	-	-	-	5	2.00	13	2.20	-	-	-	-	-	-	-	-	-
Caudal length postventral fold CLPF	347	64.26	202	67.79	224	69.78	149	61.83	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Dorsal caudal fold height DFH	10	1.85	3.5	1.17	5	1.56	3	1.24	-	-	-	-	1.1	1.90	8.8	1.50	3.9	1.69	4.1	1.73	-	-	-	-	-
Dorsal caudal fold length DFL	78	14.44	36.5	12.25	34	10.59	38	15.77	-	-	-	-	31.4	12.70	76.5	12.80	30	12.81	33.5	14.09	-	-	-	-	-
Ventral caudal fold height VFH	16	2.96	4	1.34	8	2.49	3.5	1.45	-	-	-	-	4.6	1.90	10.6	1.80	5	2.11	4.9	2.06	-	-	-	-	-
Ventral caudal fold length VFL	187	34.63	77	25.84	88	27.41	62	25.73	-	-	-	-	66.7	26.90	160	26.70	66.7	28.46	78.8	33.18	-	-	-	74.9	33.3
Pelvic fin posterior margin P2PM	107	19.81	40	13.42	63	19.63	32.5	13.49	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelvic fin length P2L	104	19.26	52	17.45	57	17.76	42	17.43	-	-	-	-	40	16.10	103	17.20	-	-	-	-	-	-	-	-	-
Pelvic fin anterior margin P2AM	93	17.22	48	16.11	54	16.82	42	17.43	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Pelvic fin width P2W	112	20.74	38	12.75	55	17.13	33	13.69	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Clasper internal length CIL	-	-	8	2.68	-	-	9	3.73	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Clasper external length CEL	-	-	9	3.02	-	-	9	3.73	-	-	-	-	16.7	6.70	-	-	-	-	-	-	-	-	-	-	-
Head length HDL	190	35.19	122	40.94	130	40.50	103	42.74	-	-	-	-	105*	42.30*	-	-	71.12	30.36	72.32	30.45	-	-	-	94.9	42.2
Preorbital length POBL	100	18.52	57	19.13	60	18.69	50	20.75	-	-	-	-	45	18.10	124	20.70	46.7	19.92	38.9	16.36	155.87	21.75	77.22	35.8	15.9
Pre-oral length POL	87	16.11	47	15.77	51	15.89	41	17.01	-	-	-	-	40	16.10	85.8	14.30	-	-	-	-	123.19	17.19	54.76	-	-
Pre-nasal length PNL	68	12.59	37	12.42	40	12.46	33	13.69	-	-	-	-	31.7	12.80	69	11.50	-	-	-	-	91.76	12.81	44.93	-	-
Internostril width INW	46	8.52	27	9.06	29	9.03	23.5	9.75	-	-	-	-	22	8.90	50.6	8.40	34.4	14.7	32.4	13.63	133.24	18.60	-	-	-
Nasal curtain length NCL	28	5.19	14	4.70	14	4.36	13	5.39	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nasal curtain width NCW	56	10.37	28	9.40	33	10.28	25	10.37	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Mouth width MW	46	8.52	27	9.06	27	8.41	21	8.71	-	-	-	-	22.9	9.20	52.8	8.80	24.4	10.44	23.7	10	54.05	7.54	35.1	-	-
Mouth length to widest part of disc MWD	300	55.56	160	53.69	174	54.21	130	53.94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Distance from mouth to base of tail DMtT	400	74.07	205	68.79	225	70.09	165	68.46	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Eye length EYL	23	4.26	12	4.03	13	4.05	10	4.15	-	-	-	-	10.9	4.40	17.5	2.92	-	-	-	-	-	-	-	15.1	6.7
Eye width EW	10	1.85	4	1.34	5.5	1.71	4	1.66	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Interorbital length IOL	80	14.81	43	14.43	42	13.08	36	14.94	-	-	-	-	34	13.70	122	20.30	28.9	12.33	24.8	10.45	138.27	19.30	71.60	22.1	9.8
Interspiracle length ISL	91	16.85	49	16.44	57	17.76	43	17.84	-	-	-	-	48	19.40	110	18.30	-	-	-	-	-	-	-	-	-
Spiracle length SPL	41	7.59	24	8.05	24	7.48	20	8.30	-	-	-	-	18	7.30	22.5	3.80	-	-	-	-	-	-	-	-	-
Spiracle width SW	22	4.07	9	3.02	11	3.43	10	4.15	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Width of branchial area (in front)	155	28.70	82	27.52	92	28.66	68	28.22	-	-	-	-	-	-	-	-	71.1	30.36	66.93	28.18	-	-	-	-	-
First gill slit width 1GW	20	3.70	9	3.02	11	3.43	8	3.32	-	-	-	-	7.5	3.00	21.6	3.60	-	-	-	-	-	-	-	-	-
Third gill slit width 3GW	22	4.07	10	3.36	13	4.05	9	3.73	-	-	-	-	9	3.60	25	4.20	-	-	-	-	-	-	-	-	-
Fifth gill slit width 5GW	16	2.96	9	3.02	9	2.80	7	2.90	-	-	-	-	6.5	2.60	19.9	3.30	-	-	-	-	-	-	-	-	-
Distance between first gill slits DB1GS	121	22.41	68	22.82	71	22.12	56	23.24	-	-	-	-	52.1	21.00	130	21.70	-	-	-	-	167.18	23.33	77.22	-	-
Distance between third gill slits DB3GS	108	20.00	60	20.13	62	19.31	47	19.50	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Distance between fifth gill slits DB5GS	83	15.37	45	15.10	49	15.26	39	16.18	-	-	-	-	35	14.10	88.7	14.80	-	-	-	-	100.56	14.03	49.14	-	-
Distance between the first and fifth gill slits	87	16.11	46	15.44	49	15.26	37	15.35	-	-	-	-	-	-	-	-	43.34	18.5	38.86	16.36	-	-	-	-	-
Spine length SL	69	12.78	43	14.43	37	11.53	34	14.11	-	-	-	-	31.5	12.70	92.6	15.40	-	-	72.3	30.46	-	-	-	-	-
Distance from disk to dorsal fold	245	45.37	112	37.58	105	32.71	83	34.44	-	-	-	-	-	-	-	-	100	42.69	114.4	48.18	-	-	-	-	-
Distance from root of tail to ventral fold	138	25.56	70	23.49	75	23.36	62	25.73	-	-	-	-	-	-	-	-	84.4	36.05	81	34.09	-	-	-	-	-
Distance of spine from disk	135	25.00	67	22.48	82	25.55	59	24.48	-	-	-	-	-	-	-	-	77.8	33.21	70.17	29.55	-	-	-	-	-
Counts																									
Cross-series of teeth in the upper jaw	>10	-	>11	-	>12	-	>10	-	-	-	-	-	-	-	-	-	8	-	-	-	20	-	-	-	-
Cross-series of teeth in the lower jaw	20	-	16	-	17	-	>9	-	-	-	-	-	-	-	-	-	10	-	-	-	26	-	-	-	-
Papillae on floor of mouth	5	-	5	-	5	-	5	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	5	-
Sex	female		male		female		male		female		male		male		female		female		female		male		female	male	
Maturity	adult		juvenile		juvenile		juvenile		?		young		?		?		?		?		?		?	?	

Size

The largest known specimen of *H. brevis* is an incomplete mature individual of approximately 900 mm DW (SV-0063, present study), collected off Peru at ~6° S (Figure 5h). A male specimen 690 mm DW collected in Chorrillos, Lima, Peru, on 25 March 1999 (MNHN-ICOS-02627; Figure 5g) weighed 13800 g. In the present study, the specimens examined measured between 241 and 540 mm DW.

Distribution

Hypanus brevis and *H. dipterurus* have long been considered synonyms and widely distributed along the coastline of the ETP. However, recent evidence suggests that *H. brevis* is restricted to the southeastern Pacific (ESP). In South America, *H. dipterurus* has been reported in commercial fisheries off Ecuador (Calle-Morán & Béarez, 2020; Martínez-Ortiz & García-Domínguez, 2013) and Colombia (Mejía-Falla & Navia, 2019). These records may correspond to the northernmost occurrence of *H. brevis*; however, further genetic analyses are required to determine whether the diamond stingrays captured in these countries belong to *H. brevis* or *H. dipterurus*. To date, *H. brevis* has been genetically confirmed only from specimens collected in Peru. Southward, several studies have reported the presence of *H. brevis* south to Antofagasta, northern Chile (Almendras et al., 2025; Lamilla et al., 1995), which likely represents the southernmost limit of its distribution.

Common name

To differentiate *H. brevis* from its sister species, *H. dipterurus* (commonly known as the Diamond stingray), we propose the common name "Peruvian diamond stingray". This name is fitting because the original description is based on specimens collected from northern Peru.

As for the Spanish common name, the vernacular term "*batea*" is widely used in Peru as the

commercial name for this particular species, and we suggest keeping it unchanged.

Fishery

Over 29 years, a total of 135 thousand tons of chondrichthyans were recorded in the

IMARPE landing database. These figures account for 19 % of the batoid group, with less

than 1 % potentially attributable to *H. brevis* (top panel of Figure 7). Notably, 98 % of these

volumes were originally labeled by IMARPE as *Dasyatis brevis*, while the remaining were

identified as *Hypanus* sp. Between 2015 and 2024, IMARPE recorded an annual catch of

approximately 46.2 tons of *H. brevis* (*sd* of 12.3 tons), corresponding to an annual economic

value of approximately 62,000 USD (*sd* of ± 16,000; see Figure S1). The majority of the

landings were reported from northern regions of the country (44 %; averaging 20.4 ± 6.6 tons

per year) and from the southern regions (37 %; averaging 16.8 ± 9.8 tons per year; bottom

panel of Figure 7). During this period, 78 % of landings originated from gillnets, with the

remaining percentage from unspecified fishing gear types other than gillnets (e.g., beach

seines and purse seines as reported by González-Pestana et al., 2022).

The fishery for *H. brevis* is primarily coastal, with 32 % of reported catches occurring during

2015-2024 within the first nautical mile (nm) off the Peruvian coastline and an additional

44 % between 1-5 nm. The remaining catches, reported far from the 5 nm, are likely still

within the continental shelf, as most of these catches occur around the *Lobos de Afuera*

Islands (Figure 8).

FIGURE 7 Peruvian Chondrichthyan species documented in the IMARPE landing data from 1996 to 2024 (top panel) and landings potentially attributable to *Hypanus brevis* within the same period (bottom panel).

FIGURE 8 The occurrence of landings of *Hypanus brevis* in the Peruvian Sea from 2015 to 2024, classified within a 0.1° latitude by 0.1° longitude grid.

4. DISCUSSION

Allopatric speciation is a major evolutionary model that has shaped the population history of most aquatic species, including elasmobranchs (Ball et al., 2016; Gayford & Jambura, 2025). This evolutionary mechanism has been a primary driver of species richness in batoids (Staggi et al., 2025), which comprises more than a half of all elasmobranch diversity (Ebert et al., 2017). The speciose nature of this elasmobranch group, coupled with its highly conserved morphology (dorsoventrally flattened bodies and countershaded coloration), have led to a long history of inaccurate taxonomic descriptions (Chatzispyrou & Koutsikopoulos, 2025; Last et al., 2016b).

This issue is particularly evident in the genus *Hypanus*, whose taxonomic composition has been subjected to major controversy, with significant revisions performed since the last decade (Last et al., 2016b). Most of these taxonomic revisions (see Table 4) were accomplished by means of integrative taxonomic approaches, including DNA information from complete mitogenomes or partial gene sequences (Last et al., 2016b; Mejía-Falla et al., 2025; Petean et al., 2020, 2024). Nevertheless, the species diversity of *Hypanus* is far from taxonomic clarity, primarily due to unresolved species complexes and hidden lineages, warranting additional revisions (Petean et al., 2024).

The findings of the present research resolve a taxonomic debate spanning nearly one and a half century, originating with contemporaneous descriptions published by Jordan and Gilbert (1880) and Garman (1880). By providing robust molecular and morphological evidence, this study resurrects the Peruvian diamond stingray, *H. brevis* (ESP), from synonymy with *H. dipterurus* (ENP). The resurrection of *H. brevis* as a distinct evolutionary unit from the ESP was supported by the results of different molecular analyses conducted herein. The Bayesian phylogenetic analysis recovered all specimens of *H. brevis* in a distinct monophyletic clade. A mean distance of 4 % between the COI gene from *H. dipterurus* and *H. brevis* reinforce its

validation as a different species. The species status of both Pacific diamond stingrays was

further supported by AMOVA results. The high and significant Φ_{ST} values confirm that the

vast majority of genetic variation is partitioned between groups (ENP vs ESP), indicating

complete genetic isolation and lack of gene flow between them. Moreover, results of the

haplotype network analyses revealed a complete lineage sorting, with no haplotypes shared

between these two sister species.

Morphologically, *H. brevis* and *H. dipterurus* appear difficult to distinguish. Both species

share a generally similar disc shape, proportional measurements, dentition patterns, and

external coloration, resulting in considerable overlap in the characters traditionally used for

identification. Subtle differences have been suggested in previous studies; however, these

traits often show intraspecific variation and may be influenced by ontogeny or preservation

condition, reducing their diagnostic reliability. Consequently, the identification of these

species based solely on external morphology remains challenging and may lead to

misidentifications in the absence of comprehensive comparative material or additional lines

of evidence.

Garman (1880, 1883) and Lamilla et al. (1995) reported the presence of five oral papillae in

H. brevis. The present study likewise confirms five papillae on the floor of the mouth,

arranged in a transverse row, consisting of three well-developed medial papillae and one-

minute lateral papilla on each side, widely separated from the medial group. The small size

and inconspicuous nature of the lateral papillae may hinder their proper recognition.

Therefore, although the number of papillae appears consistent, careful examination is

required when assessing this character.

Dermal ornamentation in *H. brevis* has been recognized since the earliest descriptions as a

highly variable character, potentially associated with body size and age. Garman (1880,

1883), based on a large female, described the presence of three large, erect, broad-based

denticles located anterior to the caudal spine, as well as a rough tail bearing smaller tubercles.

A similar pattern was observed in the present study on a large incomplete specimen of *H.*

brevis (SV-0063), approximately 900 mm DW, whose tail exhibited more than 14 well-

developed denticles (Figure 5h). Garman (1880, 1883) also noted short rows of one to four

tubercles on the shoulder region, suggesting that their number likely increases with age,

following a pattern comparable to that observed in closely related Atlantic species.

With respect to coloration, early descriptions already noted differences among *Hypanus*

specimens along a latitudinal gradient. Jordan and Gilbert (1880) described specimens as

light brown, slightly marbled with darker pigmentation, lacking distinct spots, and with a

blackish tail. Subsequently, Beebe and Tee-Van (1941) observed that specimens from ENP

exhibit a darker overall coloration, ranging from dark brown to nearly black. In contrast,

specimens from the ESP display brown to olive, or slightly reddish tones. Nevertheless, these

differences in pigmentation should be interpreted with caution, given the uncertainty

surrounding the identity of the northern specimens. Broader comparative studies will be

necessary to assess the taxonomic consistency of this trait and to rule out the influence of

environmental or ontogenetic factors.

Regarding body size, specimen SV-0063 represents the largest documented record of *H.*

brevis directly observed by the authors, reaching approximately 900 mm disc width (DW)

(Figure 5h). Chirichigno and Cornejo (2001) reported maximum total lengths of up to 1.8 m

TL. Although these larger sizes were not verified through examined material, they suggest

that *H. brevis* may attain greater dimensions than currently represented in scientific

collections.

The scarcity of large specimens in reference collections likely reflects logistical difficulties in

capturing, handling, and preserving large rays rather than true rarity in nature. As a result, *H.*

brevis remains poorly represented in museum collections and historically understudied.

Similar limitations have been reported for other Eastern Pacific chondrichthyans, where

insufficient material constrains robust taxonomic and biological assessments (Becerril-García

et al., 2022).

It has been proposed that the population history of the Pacific *Hypanus* lineages started

during the Pliocene, with the formation of the Isthmus of Panama (Rosenberger, 2001). This

major geological event occurred about three to four Ma (Thacker, 2017) and led to the

allopatric speciation of three Pacific *Hypanus* lineages: *H. dipterurus*, *H. longus*, and *H.*

rubioi, which branched off from their Atlantic transisthmian geminates: *H. say*, *H.*

americanus, and *H. guttatus*, respectively (Mejía-Falla et al., 2025; Petean et al., 2024;

Rosenberger, 2001). To get further insights into their evolutionary history, we employed the

combined information from the COI and 16S rRNA genes to estimate their divergence times.

Regrettably, the lack of 16S rRNA sequences in public databases for the recently discovered

H. rubioi prevented us from dating its transisthmian divergence.

Our calibrated phylogenetic results dated the divergence time between *H. say* (Western

Atlantic) and *H. dipterurus* (ENP) at approximately 3.42 Ma, while the subsequent split

between *H. dipterurus* and *H. brevis* (ESP) was estimated at 3.09 Ma. These findings suggest

that the monophyletic amphi-American lineage formed by these three *Hypanus* species

emerged by an initial transisthmian vicariance (Jordan, 1908) followed by a secondary

antitropical intra-Pacific radiation (Thacker, 2017). This two-step speciation model can be

attributed to geological and oceanographic shifts that occurred during the gradual shoaling of

the Central American Seaway (CAS), which lasted for approximately 12 million years

(Lessios, 2008). These environmental drivers include intensified Pacific upwellings, cooler

nutrient-rich deep waters, and salinity divergence (Carrillo-Briceño et al., 2018; O'Dea et al.,

2016). During the Pliocene and Pleistocene, the Eastern Tropical Pacific was marked by

dramatic changes in marine systems resulting in cold water upwelling zones (Ludt, 2021).

These lowered temperature zones during glacial periods potentially served as corridors

enabling marine taxa to cross the Equator (Burridge, 2002; Ludt, 2021). After the glacial

dispersal, warm tropical interglacial waters acted as a thermal barrier, pushing the ancestral

Pacific populations toward the poles and leading to allopatric speciation (Lindberg, 1991).

The proposed "tropical gap" between *H. brevis* and *H. dipterurus* is currently based on a lack

of confirmed occurrences in Central America. Our study analyzed genetically verified

samples from the core populations of both species: *H. brevis* from the ESP (Peru) and *H.*

dipterurus from the ENP (Mexico and southern USA). These locations represent typical

temperate to subtropical waters, as defined by established marine ecoregions (Spalding et al.,

2007). Although previous studies have reported low catch frequencies of *H. dipterurus* in

Central America (e.g., Panama, Morales-Saldaña et al., 2025), these records may represent

misidentifications. It is also likely that infrequent sightings of *H. dipterurus* in central

American waters are merely transient range shifts triggered by climatic phenomena (Cerutti-

Pereyra et al., 2024). In this light, further research is needed to confirm the actual geographic

distribution boundaries of these stingrays.

The mean genetic distance of 4 % (see Table S1) observed between the COI gene of *H.*

dipterurus and *H. brevis* represents a substitution rate of 1.22 % per million years. This rate is

relatively high for elasmobranchs, which are known for their low mitochondrial mutation

rates, estimated at a range of 0.2 to 0.3 % per million years (Martin et al., 1992). This can be

a result of the rapid divergence between *H. dipterurus* and *H. brevis* following the closure of

the CAS. These geminate species were forced to adapt to major shifting environments,

triggering fast microevolutionary changes to maintain fitness in their new habitats (Knowlton

& Weigt, 1998). Another explanation could be associated with the small effective population

size that occurs when species experience a founder effect. Genetic drift can be more powerful

in small populations, leading to the fast fixation of new mutations (Slatkin & Excoffier,

2012).

A founder event could be also a possible speciation pathway in *H. brevis*. This event would

also explain the low intraspecific diversity in *H. brevis*, where a single, fixed haplotype was

detected in the COI gene. However, the divergence span of 3.09 Ma, suggested by our

molecular dating analysis, is more than enough for new mutations to accumulate and

establish different haplotypes (Dos Reis et al., 2015; Martin et al., 1992). The fixation of a

single COI haplotype across a wide latitudinal gradient (~3°S to 14°S), from the tropical

waters of Tumbes to the temperate-cold waters of Pisco, suggests high connectivity among

the Peruvian populations of *H. brevis*. High levels of gene flow among these populations are

likely facilitated by the northward movement of the Humboldt Current System (HCS) (Thiel

et al., 2007).

The low genetic variation observed among the Peruvian samples of *H. brevis* may be also the

result of genetic erosion driven by repetitive, high-intensity El Niño-Southern Oscillation

(ENSO) events that have occurred since the Late Pliocene (Oldeman et al., 2021). This

timeline coincides with the divergence age of *H. brevis* of approximately 3.09 Ma (Figure 4).

During extreme ENSO events, collapse of primary productivity in the HCS leads to mass

mortalities, which may have caused severe population bottleneck events in the Peruvian

diamond stingray, purging ancestral genetic variation (Arntz et al., 1991; Parouffe et al.,

2025). Alongside ENSO events, another possible driver of genetic erosion in Peruvian

populations of *H. brevis* is the highly selective fishing pressure that has targeted this

particular species for several decades. Myliobatiform rays, including *H. dipterurus*, are

known for their marked sexual size dimorphism, with females reaching larger sizes than

males (Smith, 2005). Peruvian artisanal fisheries frequently use gillnets selectively targeting

large individuals, which can lead to a loss of maternally-transmitted mitochondrial diversity

and a reduced capacity to recover from depletion (Sadler et al., 2023).

The reduced genetic diversity of *H. brevis* is of particular conservation concern given that

low genetic variation is often associated with a reduced adaptive potential to climatic changes

(Reed & Frankman, 2003). While our genetic results indicate that *H. dipterurus* from the

ENP showed classic signs of population expansion, *H. brevis* from the ESP, contrastingly,

may have experienced a recent population bottleneck. This warrants immediate research on

the ESP populations of the recently resurrected Peruvian diamond stingray lineage. These

studies should conduct population genetic structure analyses and the estimation of effective

population size using additional nuclear markers in more intensive geographic sampling. The

results from these studies will determine whether the observed low mitochondrial variation in

H. brevis reflects a critical population bottleneck or widespread connectivity within the

Peruvian populations. Ultimately, these data will clarify the population's adaptive potential

and enable the identification of management units needed for robust conservation strategies.

Conservation and management considerations

The taxonomic separation of *H. brevis* from *H. dipterurus* indicates that both species possess

a more restricted distribution than previously understood when they were considered as a

single species. Based on the available information, Peru serves as a significant hub for the

fishing of *H. brevis* by small-scale operators, with certain reports indicating the presence of

this species in northern Chile (Almendras et al., 2025; Bustamante et al., 2023) and Ecuador,

including the Galápagos Islands (Alcivar-Holguin & Pilligua-Holguin, 2025; Bermello-Vera,

2024; Grove et al., 2022; Victor et al., 2022). Moreover, further research is required to

validate the distribution range and population structure, and to ascertain the exploitation

status of the stock(s). *H. dipterurus* is a species that exhibits localized depletion due to its

intrinsic life-history traits and excessive fishing pressure (Smith et al., 2008). Given their

close relatedness, the latter underscores the need to protect the two main fishing spots in Peru

(see Figure 8), in a way to prevent future collapses of *H. brevis*, which exhibits remarkable

mitochondrial homogeneity (see a discussion of area-based management mechanisms in

Campos-León et al., 2025).

Considering the relative importance of the Peruvian fishery as the sole focus of regular

fishing efforts throughout the years, it is a positive signal the fact that Peruvian landings have

not experienced a sharp decline over the past three decades (Figure 7). Nonetheless, this

should not be interpreted in isolation as an indicator of sustainable fishing practices. While

catches may have remained relatively stable and provided a relatively consistent supply to

markets over time, the unit of fishing effort may have increased, and/or the average size of

caught individuals may have decreased. IMARPE maintains a comprehensive monitoring

program for artisanal fisheries that likely contains relevant data that could thereby facilitate

answering some of the relevant research questions to inform specific management provisions.

Therefore, it is advisable to disclose existing biological and fisheries data and to develop

plans for the collection of supplementary data (Campos-León et al., 2025; González-Pestana

et al., 2022). Revision of the available datasets may lead to an update of the life-history traits,

as until now most have been estimated solely through sampling *H. dipterurus* individuals

within the ENP (Mariano-Meléndez, 1997; Restrepo-Gómez et al., 2020; Smith, 2005; Smith

et al., 2008), while fewer studies have been conducted on the ESP *H. brevis*. With some

exceptions, the life-history parameters of *H. brevis* remain largely undocumented (Garcia-

Yarihuamán & Mantarí-Gavilano, 2021; González-Pestana et al., 2022; Sánchez-Rea, 2020;

Silva-Garay et al., 2018).

Finally, *H. dipterurus* is classified as a vulnerable species in its most recent assessment by the

IUCN Red List of Threatened Species, considering that "[i]n Peru, directed gillnet fisheries

landings for this species have seen a decline of 50 % in annual landed biomass between 1997

and 2015 [...]" (Pollom et al., 2020). Our results, covering a longer period than the IUCN

evaluation, do not show a declining landing trend. However, this should not be construed as

an indicator of sustainable management, given that the fishery is undermanaged. Therefore,

the absence of a clear definition of the exploitation status should not justify failure to

implement management measures; instead, a more precautionary approach must be adopted,

acknowledging the significant uncertainty and recognizing that the genetic diversity of the

reclassified *H. brevis* is impoverished, serving as a potential proxy for population health that

indicates conservation concerns. Consequently, it calls for further analyses to identify

management strategies that prevent additional losses of genetic variation. Future IUCN

evaluations should consider *H. brevis* as an independent species.

1038

1039 AUTHOR CONTRIBUTIONS

Alan Marín: conceived and designed the study, supervised the project, designed the

molecular experiments, analyzed the data, conducted the population

genetics/phylogenetic/evolutionary analyses, developed figures, wrote the first draft of the

manuscript.

Fabiola Zavalaga: conceived and designed the study, conducted field work, conducted and

led the morphometric analysis, analyzed the data, developed figures, wrote the original draft

(morphometric sections).

Renato Gozzer-Wuest: conceived and designed the study, conducted the fisheries analysis,

analyzed the data, developed figures, wrote the original draft (fisheries and conservation

sections).

Luis E. Santos-Rojas: conducted field work, performed the molecular experiments, analyzed

the data.

Lorenzo E. Reyes-Flores: conducted field work, performed the molecular experiments,

analyzed the data.

Ruben Alfaro: conducted field work, performed the molecular experiments, analyzed the

data.

Philippe Béarez: conceived and designed the study, conducted and led the morphometric

analysis, analyzed the data, developed figures, reviewed and edited the manuscript, wrote the

original draft (morphometric sections).

Eliana Zelada-Mázmela: conceived and designed the study, reviewed and edited the

manuscript, provided laboratory facilities for molecular analysis, administered the project,

responsible for funding acquisition.

All authors read, revised, and approved the final version of the manuscript.

1063

ACKNOWLEDGEMENTS

The authors thank Luis Bancayán for providing photographic material and tissue samples of

H. longus from Tumbes, Peru. We are grateful to R. Robertson and J. Quiñones for providing

photographs of *H. americanus*, *H. brevis*, *H. sabinus*, and *H. say*. Special thanks are also

extended to fisheries biologist Andrey Moreno for his assistance with the photographic

documentation of *H. brevis* specimens deposited at IMARPE, as well as for his support in

specimen handling.

1071

1072 **CONFLICT OF INTEREST STATEMENT**

1073 The authors declare no conflict of interest.

1074

1075

1076

DATA AVAILABILITY STATEMENT

All DNA sequences generated in the present study are available in the GenBank database under accession numbers from PZ027578 to PZ027634.

ORCID

Alan Marín https://orcid.org/0000-0001-7842-133X

Fabiola Zavalaga https://orcid.org/0000-0001-8680-4795

Renato Gozzer-Wuest https://orcid.org/0000-0002-1156-4552

Luis E. Santos-Rojas https://orcid.org/0000-0002-3756-6759

Lorenzo E. Reyes-Flores https://orcid.org/0000-0002-1993-8010

Ruben Alfaro https://orcid.org/0000-0001-8749-1647

Philippe Béarez https://orcid.org/0000-0003-0397-2393

Eliana Zelada-Mázmela https://orcid.org/0000-0002-5813-0954

REFERENCES

Alcivar-Holguin, N. A., & Pilligua-Holguin, I. A. (2025). Distribución temporal de batoideos capturados en Jaramijó-Manabí durante el 2024-2025. Universidad Laica Eloy Alfaro de Manabí, Manta Ecuador.

Almendras, D., Villafaña, J. A., Bustamante, C., Contreras, I., Campoy, A. N., Dufflocq, P., & Rivadeneira, M. M. (2025). New evidence confirms the presence of the diamond stingray *Hypanus dipterurus* (Jordan & Gilbert 1880) in Chile and extends its southern range. *Journal of Fish Biology*, *106*(4), 1245-1250.

Arntz, W. E., Tarazona, J., Gallardo, V. A., Flores, L. A., & Salzwedel, H. (1991). Benthos communities in oxygen deficient shelf and upper slope areas of the Peruvian and Chilean Pacific coast, and changes caused by El Niño. *Geological Society, London, Special Publications*, *58*(1), 131-154.

Aschliman, N. C., Nishida, M., Miya, M., Inoue, J. G., Rosana, K. M., & Naylor, G. J. (2012). Body plan convergence in the evolution of skates and rays (Chondrichthyes: Batoidea). *Molecular Phylogenetics and Evolution*, *63*(1), 28-42.

Ball, R. E., Serra-Pereira, B., Ellis, J., Genner, M. J., Iglésias, S., Johnson, A. F., ... & Griffiths, A. M. (2016). Resolving taxonomic uncertainty in vulnerable elasmobranchs: are the Madeira skate (*Raja maderensis*) and the thornback ray (*Raja clavata*) distinct species? *Conservation Genetics*, *17*(3), 565-576.

BCRP. (2025a). Data mensual, Serie de tiempo del tipo de cambio nominal. Retrieved from https://estadisticas.bcrp.gob.pe/estadisticas/series/mensuales/tipo-de-cambio-nominal

BCRP. (2025b). Data mensual, Serie de tiempo de inflación. Retrieved from https://estadisticas.bcrp.gob.pe/estadisticas/series/mensuales/inflacion.

Becerril-García, E. E., Arauz, R., Arellano-Martínez, M., Bonfil, R., Ayala-Bocos, A., Castillo-Géniz, J. L., Carrera-Fernández, M., Charvet, P., Chiaramonte, G., Cisneros-Montemayor, A. M., Concha, F., Espinoza, M., Ehemann, N. R., Estupiñán-Montaño, C., Fuentes, K., Galván-Magaña, F., Graham, R., Hacohen-Domené, A., Hazin, F., ...Godard-Codding, C. A. G. (2022). Research priorities for the conservation of chondrichthyans in Latin America. *Biological Conservation*, *269*, 109535. https://doi.org/10.1016/j.biocon.2022.109535

Beebe, W. & Tee-Van, J. (1941). Eastern Pacific expeditions of the New York Zoological Society. XXVIII. Fishes from the tropical eastern Pacific. Part 3. Rays, Mantas and Chimaeras. *Zoologica, Scientific Contributions of the New York Zoological Society*, *26* (3), 245–280.

Bermello-Vera, K. S. (2024). Distribución y caracterización de la fauna acompañante en la pesca industrial de *Merluccius gay*i Merluza durante el año 2017–2022. La Libertad. UPSE, Matriz. Facultad de Ciencias del Mar. 98p.

Borsa, P., Arlyza, I. S., Chen, W. J., Durand, J. D., Meekan, M. G., & Shen, K. N. (2013). Resurrection of New Caledonian maskray *Neotrygon trigonoides* (Myliobatoidei: Dasyatidae) from synonymy with *N. kuhlii*, based on cytochrome-oxidase I gene sequences and spotting patterns. *Comptes Rendus. Biologies*, *336*(4), 221-232.

Burridge, C. P. (2002). Antitropicality of Pacific fishes: molecular insights. *Environmental Biology of Fishes*, *65*(2), 151-164.

Bustamante, C., Acuña, E., Tapia-Jopia, C., & Vargas-Caro, C. (2023). Actualización del Plan de Acción Nacional para la conservación y manejo de tiburones de Chile. Informe Final Proyecto del Fondo de Investigación Pesquera y de Acuicultura (FIPA N°2021-24). Universidad de Antofagasta.

Calle-Morán, M. D., & Béarez, P. (2020). Updated checklist of marine cartilaginous fishes from continental and insular Ecuador (Tropical Eastern Pacific Ocean). *Cybium*, *44*(3), 239-250.

Campos-León, S., Hernández-Muñoz, S., Huerta-Beltrán, B. L., Zavalaga, F., Cevallos-Garcias, A., & Ehemann, N. (2025). Peruvian Chondrichthyes biodiversity, marketing and conservation: a synthesis of current knowledge. *Journal of Fish Biology*. https://doi.org/10.1111/jfb.70227

1160 Carrillo-Briceño, J. D., Carrillo, J. D., Aguilera, O. A., & Sanchez-Villagra, M. R. (2018).
1161   Shark and ray diversity in the Tropical America (Neotropics)—an examination of
1162   environmental and historical factors affecting diversity. *PeerJ*, *6*, e5313.
1163
1164 Cerutti-Pereyra, F., Drunkard, E., Espinoza, M., Finucci, B., Galván-Magaña, F., Hacohen-
1165   Domené, A., … & Chin, A. (2024). Vulnerability of Eastern Tropical Pacific
1166   chondrichthyan fish to climate change. *Global change and biology, 30*(7), e17373.
1167
1168 Chatzispyrou, A., & Koutsikopoulos, C. (2023). Tracing patterns and biodiversity aspects of
1169   the overlooked skates and rays (Subclass Elasmobranchii, Superorder Batoidea) in
1170   Greece. *Diversity*, *15*(1), 55.
1171
1172 Chen, M. Y., Liang, D., & Zhang, P. (2017). Phylogenomic resolution of the phylogeny of
1173   laurasiatherian mammals: exploring phylogenetic signals within coding and noncoding
1174   sequences. *Genome Biology and Evolution*, *9*(8), 1998-2012.
1175
1176 Chirichigno, N. (1974). *Clave para identificar los peces marinos del Perú*. Informe Instituto
1177   del Mar del Perú (IMARPE), Callao, Perú. 387 pp.
1178
1179 Chirichigno, N., & Vélez, J. (1998). *Claves para identificar los peces marinos del Perú*. 2ª
1180   edición. Instituto del Mar del Perú (IMARPE), Callao, Perú. 496 pp.
1181
1182 Chirichigno, N., & Cornejo, R.M. (2001). *Catálogo comentado de los peces marinos del
1183   Perú*. Instituto del Mar del Perú (IMARPE), Callao, Perú. 314 pp.
1184
1185 Cunha, Y. T. C., Leite, R. D., Pereira da Silva, L. G., Cruz, V. P., Nunes, J. L. S., Rotundo,
1186   M. M., ... & Sales, J. B. D. L. (2026). A comprehensive overview of evolutionary
1187   relationships and geographic distributions of cownose rays: more valid taxa needing
1188   protective measures. *Reviews in Fish Biology and Fisheries*, *36*(1), 37.
1189
1190 Dos Reis, M., Donoghue, P. C., & Yang, Z. (2016). Bayesian molecular clock dating of
1191   species divergences in the genomics era. *Nature Reviews Genetics*, *17*(2), 71-80.
1192
1193 Ebert, D.A. (2003). Sharks, Rays, and Chimaeras of California. University of California Press.
1194
1195 Ebert, D. A., Bigman, J. S., & Lawson, J. M. (2017). Chapter Two - Biodiversity, life history,
1196   and conservation of Northeastern Pacific Chondrichthyans. In S. E. Larson, & D. Lowry
1197   (eds), *Advances in marine biology* (Vol. 77, pp. 9-78). Academic Press.
1198
1199 Eschmeyer, W.N. (editor). 1998. Catalog of fishes. Spec. Publ. No. 1, Center for Biodiversity
1200   Research and Information. California Academy of Sciences, San Francisco. 483 pp.
1201
1202 Feitosa, L. M., Queiroz, A. P., Labonne, M., Dressler, V. L., & Lessa, R. P. (2021). Habitat
1203   use and nursery evaluation for the longnose stingray *Hypanus guttatus* (Bloch &
1204   Schneider, 1801) using vertebral microchemistry. *Journal of Fish Biology*, *99*(5), 1538-
1205   1549.
1206
1207 Fricke, R., Eschmeyer, W. N., & Fong, J. D. (2025). Eschmeyer's Catalog of Fishes:
1208   Genera/Species/References. (http://researcharchive.calacademy.org/research/ichthyology/c
1209   atalog/fishcatmain.asp). Electronic version accessed 09 dec 2025.

1210
1211 Gabbanelli, V., de Astarloa, J. M. D., Gonzalez-Castro, M., Vazquez, D. M., & Mabragaña, E.
1212   (2018). Almost a century of oblivion: Integrative taxonomy allows the resurrection of the
1213   longnose skate *Zearaja brevicaudata* (Marini, 1933) (Rajiformes; Rajidae). *Comptes*
1214   *Rendus Biologies*, *341*(9-10), 454-470.
1215
1216 García-Yarihuamán, L. R., & Mantarí-Gavilano, P. J. (2021). Hábitos alimentarios de los
1217   batoideos de importancia comercial de Pisco-Ica, Mayo–Septiembre 2019. Undergraduate
1218   thesis. Universidad Nacional San Luis Gonzaga. Ica, Perú Available at:
1219   https://repositorio.unica.edu.pe/server/api/core/bitstreams/ee5ad01d-a16c-4f9f-9add-9be0
1220   280382e1/content.
1221
1222 Garman, S. (1880). New species of selachians in the museum collection. *Bulletin of the*
1223   *Museum of Comparative Zoology*, *6*(11), 167-172.
1224
1225 Garman, S. (1883). The American species of the genus *Dasibatis*. *In* D. S. Jordan & C. H.
1226   Gilbert (Eds.), Synopsis of the fishes of North America. *Bulletin of the United States*
1227   *National Museum*, *16*, 65-72.
1228
1229 Garman, S. (1913). The Plagiostomia. (Sharks, Skates, and Rays). *Memoirs of the Museum of*
1230   *Comparative Zoology*, *36*, 1-515.
1231
1232 Gayford, J. H., & Jambura, P. L. (2025). Drivers of diversification in sharks and rays
1233   (Chondrichthyes: Elasmobranchii). *Frontiers in ecology and evolution*, *12*, 1530326.
1234
1235 González-Pestana, A., Velez-Zuazo, X., Alfaro-Shigueto, J., & Mangel, J. C. (2022). Batoid
1236   fishery in Peru (1950-2015): Magnitude, management and data needs. *Revista de Biología*
1237   *Marina y Oceanografía*, *57* (Especial), 217-233.
1238
1239 Grove, J. S., Long, D. J., Robertson, D. R., & Victor, B. C. (2022). List of fishes of the
1240   Galapagos Archipelago, Ecuador. *Journal of the Ocean Science Foundation, 39*, 14-22.
1241
1242 Hosegood, J., Humble, E., Ogden, R., De Bruyn, M., Creer, S., Stevens, G. M., ... &
1243   Carvalho, G. (2020). Phylogenomics and species delimitation for effective conservation of
1244   manta and devil rays. *Molecular ecology*, *29*(24), 4783-4796.
1245
1246 Jordan, D. S. (1908). The law of geminate species. *The American naturalist, 42*(494), 73-80.
1247
1248 Knowlton, N., & Weigt, L. A. (1998). New dates and new rates for divergence across the
1249   Isthmus of Panama. *Proceedings of the Royal Society of London. Series B: Biological*
1250   *Sciences*, *265*(1412), 2257-2263.
1251
1252 IMARPE. (2025). INFOMAR: Proyecto Pesca Artesanal - Sistema de información de precios
1253   playa, mareas y oleajes. Retrieved from http://infomar.imarpe.gob.pe:8080/Acceso/Login?
1254   ReturnUrl=%2F
1255
1256 iNaturalist. (2026). iNaturalist Research-grade Observations. iNaturalist.org. Occurrence
1257   dataset https://doi.org/10.15468/ab3s5x accessed via GBIF.org on 2026-01-10.
1258   https://www.gbif.org/occurrence/1945421174
1259

IUCN Standards and Petitions Committee. (2024). Guidelines for Using the IUCN Red List Categories and Criteria. Version 16. Prepared by the Standards and Petitions Committee. Downloadable from https://www.iucnredlist.org/documents/RedListGuidelines.pdf.

IUCN (2025). The IUCN Red List of Threatened Species. Version 2025-2. https://www.iucnredlist.org

Jordan, D. S., & Gilbert, C. H. (1880). Notes on a collection of fishes from San Diego, California. *Proceedings of the United States National Museum, 3*, 23-24.

Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *Journal of molecular evolution, 16*(2), 111-120.

Lamilla, J., Pequeño, G., & Kong, I. U. (1995). *Dasyatis brevis* (Garman, 1880) segunda especie de Dasyatidae registrada para Chile (Chondrichthyes, Myliobatiformes). *Estud. Oceanol, 14*, 23-27.

Landemaine, O. (1991). Sélaciens nouveaux du Crétacé supérieur du Sud-Ouest de la France. Quelques apports à la systématique des élasmobranches. *Saga, 1*, 1-45.

Last, P. R., Manjaji-Matsumoto, B. M., Naylor, G. J. P., & White, W. T. (2016a). Chapter 25 Stingrays: family Dasyatidae. In P. R. Last, W. T. White, M. R. de Carvalho, B. Séret, M. F. W. Stehmann, & G. J. P. Naylor (Eds.), *Rays of the World* (pp. 522-618). CSIRO Publishing, Clayton South, Australia.

Last, P. R., Naylor, G. J., & Manjaji-Matsumoto, B. M. (2016b). A revised classification of the family Dasyatidae (Chondrichthyes: Myliobatiformes) based on new morphological and molecular insights. *Zootaxa, 4139*(3), 345-368.

Lessios, H. A. (2008). The great American schism: divergence of marine organisms after the rise of the Central American Isthmus. *Annual Review of Ecology, Evolution, and Systematics, 39*(1), 63-91.

Lindberg, D. R. (1991). Marine biotic interchange between the northern and southern hemispheres. *Paleobiology, 17*(3), 308-324.

Love, M. S., Mecklenburg C. W., Mecklenburg T. A., & Thorsteinson L. K. (2005). Resource inventory of marine and estuarine fishes of the west coast and Alaska: A checklist of North Pacific and Arctic Ocean species from Baja California to the Alaska-Yukon border. U. S. Department of the Interior, U. S. Geological Survey, Biological Resources Division, Seattle.

Ludt, W. B. (2021). Missing in the middle: a review of equatorially disjunct marine taxa. *Frontiers in Marine Science, 8*, 660984.

Marín, A. (2025). The first complete mitogenome of the endangered spotted eagle ray, *Aetobatus ocellatus*, confirms species and family taxonomic status. *Genetica, 153*(1), 34.

Marín, A., Serna, J., Robles, C., Ramírez, B., Reyes-Flores, L. E., Zelada-Mázmela, E., ... & Alfaro, R. (2018). A glimpse into the genetic diversity of the Peruvian seafood sector: Unveiling species substitution, mislabeling and trade of threatened species. *PLoS One*, *13*(11), e0206596.

Marín-Abanto, P. M. (2018). Fenix Package. Retrieved from https://github.com/PabloMBooster/fenix/blob/master/DESCRIPTION

Marín Soto, W., Medina Cruz, A., Castillo Mendoza, G., Estrella Arellano, C., Guardia Otárola, A., Guevara Carrasco, R., ... & Bertrand, A. (2017). Atlas de la pesca artesanal del mar del Perú. IMARPE-IRD, Lima, Perú, 183 pp.

Marshall, A. D., Compagno, L. J., & Bennett, M. B. (2009). Redescription of the genus *Manta* with resurrection of *Manta alfredi* (Krefft, 1868) (Chondrichthyes; Myliobatoidei; Mobulidae). *Zootaxa*, *2301*(1), 1-28.

Martin, A. P., Naylor, G. J., & Palumbi, S. R. (1992). Rates of mitochondrial DNA evolution in sharks are slow compared with mammals. *Nature*, *357*(6374), 153-155.

Martínez-Ortiz J. & García-Domínguez M. (2013). Guía de campo de condrictios del Ecuador: quimeras, tiburones y rayas / Chondrichthyes of Ecuador field guide. Chimaeras, sharks and rays. 246 p. Manta: Ministerio de Agricultura, Ganadería, Acuicultura y Pesca.

Mejía-Falla, P. A., & Navia, A. F., (2019). Checklist of marine elasmobranchs of Colombia. *Universitas Scientiarum*, *24*(1), 241-276.

Mejía-Falla, P. A., Navia, A. F., Cardeñosa, D., & Tavera, J. (2025). New Species of the Genus *Hypanus* (Dasyatidae) from the Eastern Tropical Pacific Ocean. *Ichthyology & Herpetology*, *113*(1), 44-60.

Miller, M. A., Pfeiffer, W., & Schwartz, T. (2010). Creating the CIPRES Science Gateway for inference of large phylogenetic trees. *2010 Gateway Computing Environments Workshop (GCE), New Orleans, USA* (pp. 1-8).

Miller, E. F., Dossett, W. H., & Rankin, J. L. (2014). Clarifying the northern extent of the diamond stingray (*Dasyatis dipterura*) in Southern California. Bulletin, Southern California Academy of Sciences, *113*(3), 232-235.

Morales-Saldaña, J. M., Guzmán, H. M., Vega, A. J., Robles, Y. A., Montes, L. A., & Kyne, P. M. (2025). A review of the status of sharks, rays and chimaeras of Panama to guide research and conservation. *Aquatic Conservation: Marine and Freshwater Ecosystems*, *35*(4), e70122.

Mundy, B. C. (2005). Checklist of the fishes of the Hawaiian Archipelago. *Bishop Museum Bulletins in Zoology*, *6*, 1-703.

Naylor, G. J. P., Caira, J. N., Jensen, K., Rosana, K. A. M., White, W. T., & Last, P. R. (2012). A DNA sequence–based approach to the identification of shark and ray species and its implications for global elasmobranch diversity and parasitology. *Bulletin of the American Museum of Natural History*, *367*, 1-262.

Nelson, J. S., Crossman, E. J., Espinosa-Pérez, H., Findley, L. T., Gilbert, C. R., Lea, R. N., & Williams, J. D. (2004). *Common and scientific names of fishes from the United States, Canada and Mexico.* American Fisheries Society. Special Publication 29, Bethesda, Maryland. 192 pp.

Nishida, K., & Nakaya, K. (1990). Taxonomy of the genus *Dasyatis* (Elasmobranchia, Dasyatidae) from the North Pacific. In: Pratt Jr., H. L., S. H. Gruber and T. Taniuchi (eds.): *Elasmobranchs as living resources: advances in the biology, ecology, systematics, and the status of the fisheries.* NOAA (National Oceanic and Atmospheric Administration) Tech. Rept. NMFS 90:327-346.

Nylander, J. A., Ronquist, F., Huelsenbeck, J. P., & Nieves-Aldrey, J. (2004). Bayesian phylogenetic analysis of combined data. *Systematic biology*, *53*(1), 47-67.

O'Dea, A., Lessios, H. A., Coates, A. G., Eytan, R. I., Restrepo-Moreno, S. A., Cione, A. L., ... & Jackson, J. B. (2016). Formation of the Isthmus of Panama. *Science advances*, *2*(8), e1600883.

Oldeman, A., Baatsen, M., von Der Heydt, A., Dijkstra, H., Tindall, J., Abe-Ouchi, A., …& Williams, C. (2021). Reduced El Niño variability in the mid-Pliocene according to the PlioMIP2 ensemble. *Climate of the Past, 17*(6), 2427-2450.

Palumbi, S. R., Martin, A. P., Romano, S., McMillan, W. O., Stice, L., Grabowski, G. (1991). The simple fool's guide to PCR. Department of Zoology-University of Hawaii, Special Publication, Honolulu.

Parouffe, A., Dewitte, B., Paulmier, A., & Garçon, V. (2025). ENSO modulates aerobic habitat across varying hypoxia tolerance levels in the Southeast Pacific throughout the twenty-first century. *Scientific Reports*, *15*(1), 20670.

Petean, F. F., Naylor, G. J., & Lima, S. M. (2020). Integrative taxonomy identifies a new stingray species of the genus *Hypanus* Rafinesque, 1818 (Dasyatidae, Myliobatiformes), from the Tropical Southwestern Atlantic. *Journal of Fish Biology*, *97*(4), 1120-1142.

Petean, F. F., Yang, L., Corrigan, S., Lima, S. M., & Naylor, G. J. (2024). How many lineages are there of the stingrays genus *Hypanus* (Myliobatiformes: Dasyatidae) and why does it matter? *Neotropical Ichthyology*, *22*, e230046.

Philippe, H., Brinkmann, H., Lavrov, D. V., Littlewood, D. T. J., Manuel, M., Wörheide, G., & Baurain, D. (2011). Resolving difficult phylogenetic questions: why more sequences are not enough. *PLoS biology*, *9*(3), e1000602.

Pollom, R., Bizzarro, J., Burgos-Vázquez, M.I., Cevallos, A., Velez-Zuazo, X., Avalos, C., Espinoza, M., González, A., Herman, K., Mejía-Falla, P.A., Navia, A.F., Pérez Jiménez, J.C., & Sosa-Nishizaki, O. (2020). *Hypanus dipterurus*. The IUCN Red List of Threatened Species 2020: e.T60152A80677563. https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T 60152A80677563.en. Accessed on 22 November 2025.

Puentes, V., Mejía-Falla, P. A., Ramirez, J. G., Manjarrés-Martínez, L. M., Rguez-Baron, J. M., Zapata, L. A., ... & Navia, A. F. (2022). Sharks and marine batoids management in Colombia: Policy instruments, management duty and implications for their populations and stakeholders. *Marine Policy*, *145*, 105264.

Rambaut, A., Drummond, A. J., Xie, D., Baele, G., & Suchard, M. A. (2018). Posterior summarization in Bayesian phylogenetics using Tracer 1.7. *Systematic biology*, *67*(5), 901-904.

Reed, D. H., & Frankham, R. (2003). Correlation between fitness and genetic diversity. *Conservation biology*, *17*(1), 230-237.

Restrepo-Gómez, D. C., Cruz-Escalona, V. H., Peterson, M. S., Mejía-Falla, P. A., & Navia, A. F. (2020). Effects of age, maturity stage, sex and seasonality on the feeding strategies of the diamond stingray (*Hypanus dipterurus*) in the southern Gulf of California. *Marine and Freshwater Research, 72*(4), 469-480.

Robertson, D. R., & Van Tassell, J. (2023). Shorefishes of the Greater Caribbean: online information system. Version 3.0 Smithsonian Tropical Research Institute, Balboa, Panamá.

Rosenberger, L. J. (2001). Phylogenetic relationships within the stingray genus *Dasyatis* (Chondrichthyes: Dasyatidae). *Ichthyology & Herpetology*, *2001*(3), 615-627.

Sabaj, M. H. (2020). Codes for natural history collections in ichthyology and herpetology. *Ichthyology & Herpetology*, *108*(3), 593-669.

Sadler, D. E., Watts, P. C., & Uusi-Heikkilä, S. (2023). The riddle of how fisheries influence genetic diversity. *Fishes*, *8*(10), 510.

Sales, J. B. L., de Oliveira, C. N., dos Santos, W. C. R., Rotundo, M. M., Ferreira, Y., Ready, J., ... & da Silva Rodrigues-Filho, L. F. (2019). Phylogeography of eagle rays of the genus *Aetobatus*: *Aetobatus narinari* is restricted to the continental western Atlantic Ocean. *Hydrobiologia*, *836*(1), 169-183.

Sánchez-Rea, A. B. (2020). Caracterización morfológica de las estructuras dérmicas de rayas (Chondrichthyes: batoidea) desembarcadas en el litoral peruano. Undergraduated thesis. Universidad Científica del Sur. Lima, Peru Available at: https://repositorio.cientifica.edu.pe/handle/20.500.12805/1333.

Silva-Garay, L., Pacheco, A. S., & Vélez-Zuazo, X. (2018). First assessment of the diet composition and trophic level of an assemblage of poorly known chondrichthyans off the central coast of Peru. *Environmental Biology of Fishes*, *101*(10), 1525-1536.

Simental-Anguiano, M. D., Torres-Rojas, Y. E., Galván-Magaña, F., & Tripp-Quezada, A. (2022). Importance of Shellfish in the Diet of Two Ray Species (*Rhinoptera steindachneri* and *Hypanus dipterurus*) in the Upper Gulf of California. *Journal of Shellfish Research*, *41*(2), 291-299.

Siskey, M. R., Shipley, O. N., & Frisk, M. G. (2019). Skating on thin ice: Identifying the need for species-specific data and defined migration ecology of Rajidae spp. *Fish and Fisheries*, *20*(2), 286-302.

Slatkin, M., & Excoffier, L. (2012). Serial founder effects during range expansion: a spatial analog of genetic drift. *Genetics*, *191*(1), 171-181.

Smith, W. D. (2005). Life history aspects and population dynamics of a commercially exploited stingray*, Dasyatis dipterura* (Doctoral dissertation, San Francisco State University).

Smith, W. D., Cailliet, G. M., & Cortés, E. (2008). Demography and elasticity of the diamond stingray, *Dasyatis dipterura*: parameter uncertainty and resilience to fishing pressure. *Marine and Freshwater Research, 59*(7), 575-586.

Spalding, M. D., Fox, H. E., Allen, G. R., Davidson, N., Ferdaña, Z. A., Finlayson, M. A. X., ... & Robertson, J. (2007). Marine ecoregions of the world: a bioregionalization of coastal and shelf areas. *BioScience*, *57*(7), 573-583.

Spöri, Y., & Flot, J. F. (2020). HaplowebMaker and CoMa: two web tools to delimit species using haplowebs and conspecificity matrices. *Methods in Ecology and Evolution*, *11*(11), 1434-1438.

Thacker, C. E. (2017). Patterns of divergence in fish species separated by the Isthmus of Panama. *BMC Evolutionary Biology*, *17*(1), 111.

Thiel, M., Macaya, E., Acuña, E., Wolf, E. A., Bastias, H., Brokordt, K., …& Vega, A. (2007). The Humboldt Current System of northern and central Chile. *Oceanography and Marine Biology: An Annual Review*, *45*, 195-344.

Victor, B. C., Grove, J. S., Long, D. J., Robertson, D. R., Keith, I., Bensted-Smith, W., & Salinas-de-León, P. (2024). List of fishes of the Galapagos archipelago, Ecuador (version 2.0). *Journal of the Ocean Science Foundation*, *41*, 54-111.

Ward, R. D., Zemlak, T. S., Innes, B. H., Last, P. R., & Hebert, P. D. (2005). DNA barcoding Australia's fish species. *Philosophical transactions of the royal society B: biological sciences*, *360*(1462), 1847-1857.

White, W. T. Last, P. R., Naylor, G.J.P., Jensen, K., & Caira, J. (2010). Clarification of *Aetobatus ocellatus* (Kuhl, 1823) as a valid species, and a comparison with *Aetobatus narinari* (Euphrasen, 1790) (Rajiformes: Myliobatidae). *CSIRO Marine and Atmospheric Research Paper*, *32*, 141.

White, W. T., & Last, P. R. (2012). A review of the taxonomy of chondrichthyan fishes: a modern perspective. *Journal of Fish Biology, 80*(5), 901-917.

White, W. T., O'Neill, H. L., & Naylor, G. J. (2022). Taxonomy and diversity of extant elasmobranchs. In W. T. White, H. L. O'Neill, & G. J. Naylor (eds), *Biology of sharks and their relatives* (pp. 31-57). CRC Press.

1506 Xia, X. (2018). DAMBE7: New and improved tools for data analysis in molecular biology
1507   and evolution. *Molecular biology and evolution*, *35*(6), 1550-1552.
1508
1509 Yamaguchi, A., Furumitsu, K., & Wyffels, J. (2021). Reproductive biology and embryonic
1510   diapause as a survival strategy for the East Asian endemic eagle ray *Aetobatus narutobiei*.
1511   *Frontiers in Marine Sciences 8*:768701
1512
1513 Zuccarello, G. C., & Lokhorst, G. M. (2005). Molecular phylogeny of the genus *Tribonema*
1514   (Xanthophyceae) using rbc L gene sequence data: monophyly of morphologically simple
1515   algal species. *Phycologia*, *44*(4), 384-392.

1516
1517
1518

Supplementary material

1519

1548 **FIGURE S1** Box-plot with the off-vessel 2009-2024 prices of *Hypanus brevis* (top-panel) and the landing
1549 income attributable to this species during the same period (bottom-panel). The horizontal dashed line represents
1550 the average 2015-2024 annual income.

TABLE S1 Group interspecific mean distances of a partial COI fragment of five *Hypanus* species from the
Pacific (*H. brevis*, *H. dipterurus*, and *H. longus*) and Atlantic Oceans (*H. americanus* and *H. say*). Analyses
were conducted using the Kimura 2-parameter model. Interspecific mean distances are shown below the
diagonal, while standard error estimates are shown above the diagonal. The last column shows the mean
intraspecific distances and their standard error estimates between parenthesis. ESP: Eastern South Pacific; ENP:
Eastern North Pacific; WA: Western Atlantic; WNA: Western North Atlantic.

1563

Species (n: sample size)	*H. brevis* (ESP)	*H. dipterurus* (ENP)	*H. say* (WA)	*H. longus* (ESP)	*H. americanus* (WNA)	Mean intraspecific distance (SD)
H. brevis (n = 27)	-	±0.0090	±0.0118	±0.0178	±0.0153	0.0000
H. dipterurus (n = 8)	0.0397	-	±0.0110	±0.0145	±0.0151	0.0035 (±0.0015)
H. say (n = 4)	0.0650	0.0578	-	±0.0145	±0.0149	0.0013 (±0.0012)
H. longus (n = 7)	0.0984	0.1063	0.1063	-	±0.0066	0.0033 (±0.0019)
H. americanus (n = 4)	0.1014	0.1046	0.1073	0.0249	-	0.0026 (±0.0018)

1564