GFI1 as a novel regulator of γδ T cell development and the IL-17/IFNγ lineage commitment

Jennifer Fraszczak[1,*], David Obwegs[2,3], Kaifee Arman[1], Tanja Muralt[1,#], Bavanitha Thurairajah[4,5], Ève Mallet Gauthier[5,6,7], Irah L. King[4,5], Heather J. Melichar[5,6], Sagar[2] and Tarik Möröy[1,7,8,*]

[1]Institut de recherches cliniques de Montréal (IRCM), Montréal, Canada.

[2]Department of Medicine II (Gastroenterology, Hepatology, Endocrinology, and Infectious Diseases), Freiburg University Medical Center, Faculty of Medicine, University of Freiburg, Freiburg, Germany

[3]Faculty of Biology, University of Freiburg, Freiburg, Germany

[4]Meakins-Christie Laboratories, Research Institute of the McGill University Health Centre, Montreal, Canada

[5]Department of Microbiology and Immunology, McGill University, Montreal, Canada

[6]Rosalind and Morris Goodman Cancer Institute of the McGill University, Montreal, Canada

[7]Département de microbiologie, infectiologie et immunologie, Faculty of Medicine, Université de Montréal, Montréal, Canada

[8]Department of Medicine, Division of Clinical and Translational Research, McGill University, Montreal, Canada

[#]present address: Department for BioMedical Research (DBMR), University of Bern, Bern, Switzerland

[*]**Corresponding authors:**

Tarik Möröy and Jennifer Fraszczak

Institut de recherches cliniques de Montréal (IRCM)

110 des Pins West Avenue

Montréal, Québec, H2W 1R7, Canada.

Tel.: 1 (514) 987-5776

Tarik.Moroy@ircm.qc.ca, Jennifer.Fraszczak@ircm.qc.ca

Abstract

GFI1 is a DNA-binding zinc finger transcription factor regulating the commitment of

hematopoietic precursors to myeloid and lymphoid lineages. Here we report that GFI1 is

expressed in γδ T cells and restricts the cellularity of RORγt$^+$Vγ6$^+$ γδT cells that produce high

levels of IL-17A, while promoting the expansion of Vγ1$^+$ and Vγ4$^+$ γδT cells. Absence of GFI1

results in a pronounced bias toward the production of γδT17 cells, commencing post-birth.

Additionally, we observe an expansion of RORγt$^+$/MAF$^+$ cells within the thymic DN1e

population of GFI1-deficient mice. The DN1e population, along with other DN subsets in GFI1

knock-out (KO) mice, exhibits a distinctive γδT17 cell-specific transcriptomic profile.

Specifically, DN1, DN3, and γδ T cells lacking GFI1 show upregulation of the B-ZIP

transcription factor MAF, which regulates genes critical for γδT cells, such as *Il17a, Il22*, and *Blk*

that are all induced in *Gfi1* deficient cells. The *Maf* gene is occupied by GFI1 in DN pre-T cells

at cognate binding sites in its promoter region, suggesting that GFI1 acts as a direct repressor of

Maf. We conclude that GFI1 functions as a novel regulator of Vγ6$^+$ γδT17 precursor cells

restricting their peripheral expansion by acting upstream of a MAF dependent regulatory

network.

Highlights

- GFI1 controls the expansion of γδ T cells in peripheral lymphoid organs and in barrier

tissues.

- GFI1 specifically restricts expansion of γδ T cells secreting IL-17 by acting upstream of a

MAF dependent transcriptional regulatory network.

- GFI1 plays a cell intrinsic role in controlling the generation of γδT cells through the

repression of *Maf*.

- GFI1 controls a MAF$^+$/RORγt$^+$ γδT cell precursor cell population within the DN1e cell

subset.

Introduction

GFI1 is a 50kD transcription factor that contains an N-terminal SNAG domain and six C-terminal C_2H_2-type zinc fingers, which facilitate DNA binding to target genes[1-4]. The SNAG domain of GFI1 interacts with the histone demethylase KDM1A (LSD1) and recruits it with CoRest to target genes[5-8]. GFI1 and LSD1 are also part of the Nucleosome remodeling and deacetylase complex (NuRD)[9,10], which enables transcriptional repression[5]. *Gfi1* was first identified in retrovirally induced murine T cell lymphoma[11,12], but later, through the analysis of *Gfi1* deficient mice[13], it was recognized that GFI1 controls the proliferation of hematopoietic stem cells[14] as well as lymphoid/myeloid[15] and monocyte/granulocyte lineage decisions in progenitor cells[2,16]. Moreover, GFI1 controls pre-T cell numbers in the thymus and the T cell differentiation process which begins at the early thymic progenitor (ETPs) stage. GFI1 is required for the survival and proliferation of the very early DN1 and DN2 pre-T cells and absence of GFI1 leads to the accumulation of an abnormal population between the DN1-DN2 stages, described as a block of differentiation[17]. In addition, GFI1 also plays a role in pre-TCR selection, where it regulates the proliferation and differentiation of DN3 cells into DN4 cells and is involved in major histocompatibility complex (MHC)-restricted positive and negative selection processes, highlighting its influence on the development of CD4 and CD8 single-positive T cells[17]. GFI1 has been shown to regulate pre-T cell differentiation through NOTCH1 signaling[18], which is essential for the maintenance of ETPs and T cell lineage commitment and gene expression of lung Krüppel-like factor (KLF2) or inhibitor of DNA binding 1 and 2 (Id1 and Id2)[17]. GFI1 also regulates Foxo1 expression in thymic DP cells to maintain a specific transcriptional program in DP cells to assure the formation of mature T cells[19]. GFI1 also restrains the generation of regulatory T cells and plays a role in their functions[20-22]. In mature CD4[+] T cells, GFI1 is induced to maintain an active Th2 program[22,23] by either preventing the degradation of GATA3 or by blocking the production of IL-17 regulated by RORγt[24-28]. GFI1 also inhibits the Th1 program in activated CD4[+] T cells[29] and reins the Th9 polarisation[30]. In mature CD8[+] T cells, GFI1 has recently been shown to control the formation of effector CD8[+] T cells[31] and the persistence of memory CD8[+] T cells[32] during infection.

To date, all studies on the function of GFI1 in T cell differentiation or peripheral T cell activity focused on the αβ T cell receptor-expressing populations, while a potential role for GFI1 in γδ T cells has not been investigated. Murine γδ T cells develop in waves starting in the fetal

thymus at E14 with Vγ5[+]Vδ1[+] dendritic epidermal T cells (DETC) followed by Vγ6[+] and Vγ4[+] γδ

T cell subsets around fetal stage E16 homing to specific tissues like the lungs or the tongue[33].

Around birth, while DETC and Vγ6[+] γδ T cell production declines drastically, new subsets of γδ

T cells emerge that seed the intestine, liver, and peripheral lymphoid organs, respectively[34].

Different γδ T cell subsets have been identified[34-36] but the two main subpopulations are γδT1

cells[37], and γδT17 cells[38]. It is known that most γδT17 cells carry either the Vγ4 or the Vγ6

chain[54], with Vγ6[+] γδ T cells being produced almost exclusively at the fetal stage and the Vγ4[+] γδ

T cells at the fetal and perinatal stages[56-58]. Recent reports suggest that γδT17 and γδT1 cells can

develop in the thymus *via* several pathways. One of these pathways is independent of the TCR

and is initiated in two subsets of the DN1 thymic precursors called DN1d and DN1e[39,40,41]. From

these subsets both IFNγ[+] and IL17[+] γδ T cells can emerge[41]. The second pathway to generate

γδT17 and γδT1 cells in the thymus involves DN3 cells, where a strong TCR signal favors the

IFNγ-producing lineage whereas a weak signal is associated with the development of γδT17

lineage[33,42-45]. Each γδ T cell subset has specific functions that are not only critical for the

immune innate response against many pathogens[35,46], but also play a role in autoimmune

disorders or inflammation[47-51]. Moreover, clinical trials are also currently under way to test

whether γδ T cells are effective in cancer immunotherapy[52] creating a need to better understand

their development and function. The role that transcription factors plays in establishing γδ T cells

fate has been challenging and current knowledge remains limited.

Here we demonstrate that GFI1 is expressed in γδ T cells and that genetic ablation of

GFI1 leads to a significant expansion of RORγt[+]Vγ6[+]γδT cells producing IL17 with a

concomitant decrease in Vγ1[+] and Vγ4[+] γδT cells in adult mice. This bias toward the γδT17

subtype starts after birth in GFI1 knock-out (KO) mice and remains in place in the adult thymus,

where a Vγ6[+] subset secreting IL-17 becomes the predominant population among γδ T cells. In

addition, GFI1 KO mice have an increased number of c-MAF[+]/RORγt[+] cells in the DN1e subset,

which has been described to contain precursors for γδT17 cells[40,41]. *Gfi1*-deficient DN

thymocytes and γδ T cells show an upregulation of the B-Zip transcription factor c-MAF, which

is known to regulate the production of γδT17 cells suggesting that GFI1 controls the cellularity of

γδ T cells and their progenitors though the regulation of c-MAF.

Results

***Gfi1* deficient mice show an expansion of γδ T cells in barrier tissues and in peripheral**

lymphoid organs.

To gain more insight into a potential role of GFI1 in the immune compartment of barrier

tissues, we performed single cell RNA-sequencing (scRNA-seq) of lung cells from GFI1 WT and

KO mice. We noticed that the UMAP representation showed two clusters (clusters 5 and 8) of

Cd4[-], *Cd8a*[-], *Cd8b*[-], *Trdc*[+] cells within the *Cd3e*[+] lymphocyte compartment that were

overrepresented in GFI1 KO mice (**Fig. 1A-C & Suppl. Fig.1A & 1B**). These two clusters were

annotated as γδ T cells while all other clusters were conventional CD4[+] or CD8[+] T cells (**Fig. 1A-**

C & Suppl. Fig.1A & 1B). Flow cytometry confirmed the presence of an increased proportion of

γδ T cells in the lungs, the spleen, lymph nodes (LN) as well as in the intestine lamina propria of

GFI1 KO mice (**Fig. 1D-F** & **Suppl. Fig. 1C**). This observation stood in stark contrast to the

numbers of CD4[+] T cells, which are known to be reduced in *Gfi1* deficient mice[17,18,53].

Accordingly, the proportion of cells from GFI1 KO mice belonging to the CD4[+] T cell clusters

(clusters 7, 10 and 0) were reduced compared to WT mice (**Fig. 1A-C & Suppl. Fig.1A & 1B**),

which was also seen by flow cytometry (**Suppl. Fig. 1D**). The same observations were made with

vav-cre GFI1[flox/flox] mice where GFI1 is deleted solely in hematopoietic cells. As in the germline

GFI1 KO mice, γδ T cells from *vav-cre GFI1*[flox/flox] mice were expanded in number and frequency

compared to controls (**Fig. 1 G-I**), while CD4[+] T cells were reduced (**Suppl. Fig. 1E**). This

excluded the non-hematopoietic environment as a potential cause for γδ T cell expansion in GFI1

KO mice.

Specific expansion of γδT17 cells in GFI KO mice

To determine if the expansion of γδ T cells in GFI1 KO mice was specific to a γδ T cells

subtype, we further examined the key features of expanded γδ T cells in GFI1 KO lungs.

Expanded lung γδ T cells showed transcriptomic features of gdT17 cells indicated by the

expression of *Blk*, *Maf*, and *Rorc* (**Fig. 2A**). We analysed these cells further and found that a large

proportion of the expanded γδ T compartment in GFI1 KO lacked CD27 (**Fig. 2B, Suppl. Fig.**

2A), a marker associated with the γδT1 subtype[54]. Consistent with this observation, we also

found that up to 50% of *Gfi1* deficient γδ T cells are RORγt[+] in the lung, spleen, and LN (**Fig.**

2C-E). Similar data were obtained with *vav-cre GFI1flox/flox* mice where increased frequencies and

numbers of RORγt$^+$ γδ T cells were also evident compared to controls (**Suppl. Fig. 2B-D**). The

upregulation of RORγt was specifically seen in γδ T cells and was not found in *Gfi1*-deficient

CD4$^+$ T cells (**Fig. 2C, Suppl. Fig. 2B**).

To test the functionalities of GFI1 KO γδ T cells, we cultured cells from the spleen, LN,

and lung of WT and GFI1 KO mice and treated them with PMA and ionomycin in the presence

of Golgi plug. We found that large proportions of *Gfi1* deficient γδ T cells from spleen, LN and

lungs produced IL17A (**Fig. 2F & 2G**), which led to significantly increased numbers of IL17A$^+$

γδ T cells in the KO culture when compared to WT controls (**Fig. 2H**). Moreover, GFI1 KO γδ T

cells were also able to produce IL22 (**Suppl. Fig. 2E-G**), a cytokine often produced together with

IL17A by γδT17 cells[55]. These findings suggest that the absence of GFI1 leads specifically to the

expansion of γδT17 cells, which produce IL17 after stimulation.

Vγ6-expressing γδT17 cells expand in the absence of GFI1.

To characterize further the subtype of γδT17 cells, we performed a scRNA-seq analysis

on sorted γδ T cells from spleens of WT and GFI1 KO mice. We annotated the clusters based on

our previously published study[59], and 14 clusters were identified in splenic γδ T cells (**Fig. 3A &**

Suppl. Fig. 3A), with 6 clusters (0, 1, 3, 4, 5 and 12) expressing genes associated with the Vγ6$^+$

γδ T cell subtype such as *Trgv6*, *Cxcr6, Scart1*, and *Rorc*[60] (**Fig. 3B & 3C**). These 6 clusters

represented almost the totality of the splenic γδ T cells sorted from GFI1 KO mice (**Fig. 3C-E**).

Consistent with the previous observations, we found that the most upregulated genes in GFI1 KO

splenic γδ T cells compared to WT controls were associated with the mature Vγ6$^+$ γδT17 cells

like *Scart1*, *Maf*, *Blk* and *Ccr2*[60], while genes associated with exchanging and adaptive-like γδ T

cells such as *Tcf7* and *Lef1* were downregulated (**Fig. 3F**), which was in agreement with the

expansion of Vγ6$^+$ γδ T cells in the spleen and in the lung of GFI1 KO mice compared to controls

determined by flow cytometry (**Fig. 3G & 3H**).

Stimulation of sorted splenic Vγ1$^-$Vγ4$^-$Vγ5$^-$ γδ T cells (meaning most likely Vγ6) with

either IL1β and IL23 or IL7 also showed that cells from the GFI1 KO mice were more responsive

than WT cells as shown by the higher production of IL17A (**Sup. Fig. 3B**) or the increased

phosphorylation of STAT3 (**Sup. Fig. 3C**). This indicated that most of the cells from GFI1 KO

mice were mostly constituted of functional Vγ6$^+$ γδ T cells, capable of responding to cytokine

stimulation. Unexpectedly, the expansion of Vγ6$^+$ γδ T cells in the spleen started only after birth

(**Fig. 3I**), at a point in development when the generation of this subtype is not supposed to be

maintained, suggesting that GFI1 controls and possibly restricts, the expansion of functional

Vγ6$^+$ γδT17 cells after birth in the periphery.

Defects in γδ T cell production are detected in the thymus of adult mice.

To determine whether the γδ T cell phenotype observed in GFI1 KO mice was due to a

developmental defect, we analyzed γδ T cells from the thymus, where γδ T cells originate and

diversify[61]. Even though, absolute numbers of γδ T cells were similar in GFI1 KO and WT adult

thymus, both numbers and frequencies of IL17A$^+$/RORγt$^+$ cells were significantly increased in

Gfi1 deficient mice compared to WT controls (**Fig. 4A-F**). Again, similar data were obtained

with *vav-cre Gfi1$^{fl/fl}$* mice (**Suppl. Fig. 4A-C**). scRNA-seq analysis of sorted thymic γδ T cells

from WT and GFI1 KO mice revealed 14 clusters with 5 clusters (clusters 6, 9, 10, 11 and 13)

expressing genes specific to the Vγ6$^+$ γδT17 subtype such as *Trgv6*, *Scart1* and *Maf* and one

cluster with genes specific to Vγ4$^+$ γδT17 cell subtype (cluster 14) (**Fig. 4G-4I and Suppl. Fig.**

4D)[60,62]. Clusters 6, 9, 10, 11 and 13 form a larger cluster containing only Vγ6$^+$ γδT17 cells

identified as Trgv6$^+$ and cluster 14 contains only Vγ4$^+$ γδT17 cells identified as Trgv4$^+$ (**Fig. 4I**).

Interestingly, the cells found in the Trgv6$^+$ cluster were exclusively from GFI1 KO mice

and represented the majority of its thymic γδ T population while the cluster Trgv4$^+$ was only

found in WT thymus (**Fig. 4I-K**). Consistent with this data, the most upregulated genes in GFI1

KO thymic γδ T cells were again genes associated to the Vγ6 subset such as *Scart1* and *Cxcr6*

(**Suppl. Fig. 4E**), which is consistent with the over representation of the of Vγ6 γδT17 subtype

already seen in the thymus of GFI1 KO mice. This bias towards the Vγ6 γδ T subset in adult

GFI1 KO thymus was also confirmed by flow cytometry by using the specific antibody against

the Vγ6 chain with a tendency but no significantly higher number of Vγ6$^+$ γδ T cells found in

adult thymocytes lacking GFI1 compared to WT thymus (**Fig. 4L**). We also noticed an increased

number of CD24$^-$ γδ T cells in adult thymocytes lacking GFI1 compared to control mice (**Fig.**

4M), making the gating on Vγ chain populations easier in Gfi1 deficient γδ T cells (**Fig. 4N**).

Again, we could confirm a higher proportion and cell count of Vγ1$^-$Vγ4$^-$Vγ5$^-$ (most likely

meaning Vγ6$^+$) cells in CD24$^-$γδ T cells lacking GFI1 in adult mice (**Fig. 4N-P**). The bias was

also confirmed by staining for CD44 and CD45RB[33,43], known to distinguish the different subsets

of γδ T cells, with significantly increased numbers of CD44$^+$CD45RB$^-$ cells (known as γδT17)

within the CD24$^-$ γδ T cell subset in GFI1 KO thymus (**Suppl. Fig. 4F, G**). As observed in the

spleen and lungs, the over representation of Vγ6$^+$ γδ T cells in the GFI1 KO thymus also started

after birth (**Suppl. Fig. 4H-K**) with higher frequencies of Vγ1$^-$Vγ4$^-$Vγ5$^-$ (most likely meaning

Vγ6$^+$) cells in CD24$^-$γδ T cells of the GFI1 KO young mice and a slight but significant increased

number of this population at day 8-10, also seen at D18-20 after birth. This data suggested that

the absence of GFI1 leads to a γδ T cell developmental defect with biased production of Vγ6$^+$ γδ

T cells over the other γδ T cell subtypes.

GFI1 plays a cell intrinsic role in γδT cells.

To determine whether the loss of GFI1 can directly impact γδ T cell development, we

evaluated GFI1 expression levels in these cells. First, flow cytometric analyses of cells from a

GFI1 reporter mouse which carries an EGFP cDNA "knock-in" in the *Gfi1* locus placing it under

the control of the *Gfi1* promoter[63] indicated that γδ T cells from spleen, LN, thymus, and lung

express higher levels of *Gfi1* than conventional CD4$^+$ T cells (**Fig. 5A**). Additionally, a higher

expression of *Gfi1* mRNA in sorted splenic and lung γδ T cells compared to naïve CD4$^+$ T cells,

where GFI1 is poorly expressed[53,63] was also confirmed by q-RT-PCR (**Fig. 5B**). *Gfi1* expression

in thymic γδ T cells was even higher than in DN1 but lower than in DN3 pre-T cells (**Fig. 5C**),

while *Gfi1b* mRNA, which was used as a negative control, was not detected in γδ T cells (**Suppl.**

Fig. 5A).

Next, we wished to determine whether the bias towards the γδT17 subtype in GFI1 KO

mice was due to the intrinsic lack of GFI1 in these cells or a consequence of the specific

environment created by the absence of GFI1 in the hematopoietic compartment. To clarify this,

we transplanted CD45.1/2$^+$ WT bone marrow (BM) cells[64,65] into non-irradiated CD45.2$^+$ GFI1

KO mice and analysed the animals after 4-5 weeks. This enabled us to study the impact of the

GFI1 KO environment on WT γδ T cells generated during adulthood (**Fig. 5D**). We were able to

identify a small pool (between 15-20%) of WT hematopoietic cells identified as CD45.1/2$^+$ cells

amongst CD45.2$^+$ GFI1 KO deficient cells representing 80-85% of the hematopoietic cells in the

spleen and lung of the recipients (**Fig. 5E & 5F**). As a control, we also transplanted CD45.1/2$^+$

WT BM cells into non-irradiated CD45.2$^+$ WT recipient mice. However, as expected, the

frequencies of CD45.1/2$^+$ WT donor cells in non-irradiated CD45.2$^+$ WT recipients were very

low in the organs analysed (**Fig. 5E & 5F**). The frequencies of WT donor splenic γδ T cells in

GFI1 KO and WT recipients were similar, while the frequencies of the endogenous GFI1 KO

host γδ T cells were still elevated compared to WT donor or recipient γδ T cells (**Fig. 5G and**

Suppl. Fig. 5B). We obtained the same results in lungs and thymus of the GFI1 KO recipient

mice when they were transplanted with WT BM cells (**Fig. 5G and Suppl. Fig. 5B**). We found

that the splenic, thymic, and lung WT γδ T cells in the KO recipient mice did not show an

overexpression of RORγt or overproduction of IL17A when compared to the endogenous GFI1

KO host γδ T cells (**Fig. 5H, I and Suppl. Fig. 5C, D**), even though a slight increase of IL17A

and RORγt could be seen compared to recipient WT γδ T cells in the spleen (**Fig. 5H, I and**

Suppl. Fig. 5C, D). Moreover, the distribution of the different Vγ chain subtypes was similar

between the WT γδ T donors in the GFI1 KO recipient and the WT γδ T cells in all analysed

organs (**Fig. 5J and Suppl. Fig. 5E**). This suggested that the GFI1 KO environment has no or

very little effect on γδ T cells and cannot explain the preferential bias toward the γδT producing

IL17 in *Gfi1* deficient mice.

Identification of a potentially precursor population in adult GFI1 KO mice

γδ T cells are thought to differentiate from progenitors located in the thymic DN

compartment, which includes DN1, DN2, DN3 and DN4 subsets[33]. We set out to determine

whether DN cells from GFI1 KO mice may have gained the ability to preferentially produce Vγ6$^+$

γδ T cells. To test this, we cultured total DN cells from adult WT and GFI1 KO thymi on OP9-

DL1 cells (**Suppl. Fig. 6A**), by excluding γδ T cells, NK cells, myeloid cells, B cells, and

conventional T cells (**Suppl. Fig. 6B, C**). Under these conditions, we observed that GFI1 KO DN

cells can give rise to higher proportions of the CD24$^-$CD44$^+$CD45RB$^-$ γδ T cells (known as

γδT17) in the CD45$^+$ cells compared to WT DN cells (**Fig. 6A, B**). These data suggested that the

bias towards the γδT17 subset in the absence of GFI1 has its origin in early precursors of γδ T

cell differentiation, which are not yet fully committed to the γδ T cell lineage.

It is already known that most γδ T cells differentiate from thymic DN2 or DN3 cells[40,56],

but recently DN1d and DN1e subsets have also been shown to generate γδ T cells[40,41]. The DN2

and DN3 subsets are strongly decreased in GFI1 KO mice compared to WT animals and the

DN1a/b subpopulations are almost absent (**Fig. 6C**). Even though DN1d and DN1e subsets were

significantly decreased in numbers in GFI1 KO mice, their frequencies were increased in *Gfi1*

deficient mice compared to control animals (**Suppl. Fig 6D**).

To gain more insight into the origin of the expanded γδT17 cell population in GFI1 KO

mice, we performed a scRNA-seq analysis of pools of DN1, DN2, and DN3 cells from adult

GFI1 WT and KO thymus. Our analysis showed that these cells fall into 9 distinct clusters (**Fig.**

6D & Suppl. Fig. 6E & 6F), which could be annotated as DN1, DN2 and DN3 cells and their

subtypes[41,62,66] (**Fig. 6E & Suppl. Fig. 6E**). Interestingly, clusters 6 and 8 contained populations

of the DN1d and DN1e subsets, that were identified as *Rorc*[+]DN1d and *Rorc*[+]DN1e, express

genes associated to the γδT17 subtype such as *Maf*, *Rorc*, *Blk,* and *Cxcr6* and were largely found

in adult GFI1 KO thymus (**Fig. 6D-H**). Another cluster (cluster 5) identified as *Il2rb*[+]DN1e was

also overrepresented in GFI1 KO thymus compared to WT thymus (**Fig. 6D-G**). These data show

that DN1d and DN1e cells were still present in adult GFI1 KO mice, with *Rorc*[+]DN1e cells being

the most expanded population in GFI1 KO DN population (**Fig. 6G**). This correlated with the

high *Gfi1* expression levels found in DN1e cells compared to other DN1 cells (**Suppl. Fig. 6G**).

The most up regulated genes in the pool of DN1-DN3 cells from adult GFI1 KO thymus

compared to control DN1-DN3 cells contained genes related to Vγ6[+] γδT17 cells, such as *Maf*

and *Scart1* (**Fig. 6I**), confirming that a γδT17 program was already active in DN1 cells from

GFI1 KO mice. We confirmed by flow cytometry, the higher proportions of c-MAF[+] cells in

GFI1 KO and in *vav-cre Gfi1*[fl/fl] DN1e, DN2 and DN3 cells (**Suppl. 6H**) and identified an

expanded c-MAF[+]RORγt[+] DN1e population in GFI1 KO and in *vav-cre Gfi1*[fl/fl] thymus as

compared to controls (**Fig. 6J & K**). This c-MAF[+]RORγt[+] DN1e population overlapped with the

abnormal DN1 population that we had described previously[17] (**Fig. 6K**), and which expressed

high level of CD44 and low level of CD25. This suggested that GFI1 restricts or controls a DN1e

population bearing the potential to give rise to Vγ6[+] γδT17 cells.

Characterisation of the DN1 MAF[+]/RORγt[+] and DN3 compartments in GFI1 KO mice

To determine if this DN1e subpopulation that was expanded in GFI1 KO mice can

preferentially differentiate into Vγ6[+] γδT17 cells, we performed a trajectory inference analysis

using Partition-based Graph Abstraction (PAGA) by combining the scRNA-seq data obtained

with DN1-3 cells and the Trgv6[+] cluster from WT and GFI1 KO thymic γδ T cells (**Fig. 7A-C**).

PAGA can infer developmental trajectories from scRNA-seq data by calculating the probability

of transition between cells[67]. This provides indications about relationships and connections of

cells and their precursors. Using this method, a potential trajectory was obtained from cluster 7

containing *Rorc*⁺DN1e, almost exclusively found in GFI1 KO thymus, to the three clusters

containing the Vγ6⁺γδT17 cells identified as Trgv6⁺ (clusters 11, 12 and 13) that represented

almost the totality of γδ T cells in GFI1 KO thymus. This indicated that the *Rorc*⁺DN1e subset

enriched in GFI1 KO mice is developmentally linked to the γδT17 cells expressing the Vγ6

chain.

The cluster containing DN1-DN2 cells (cluster 10) was connected to the different DN3a

and/or DN3b clusters (clusters 2 and 5) but was not linked to the Trgv6⁺ clusters (clusters 11, 12

and 13) containing Vγ6⁺γδT17 cells (**Fig. 7A-C**), reinforcing the idea that the expanded Vγ6⁺

γδT17 population in GFI1 KO thymus does not originate from the DN2 or DN3 compartment but

most likely from the DN1e subset. The analysis of *Cd2-cre*; Gfi1$^{flox/flox}$ mice, where a deletion of

Gfi1 occurs after the DN1 stage in the DN2-3 precursors during T cell development didn't show

changes in numbers for γδ T cells negative for Vγ1 and Vγ4 chain expression, expressing RORγt⁺

or producing IL17A compared to control mice (**Suppl. Fig. 7A-H**). One of the differences was

decreased numbers of Vγ1⁺ γδ T cells in the thymus and in the spleen of the *Cd2-cre* Gfi1$^{flox/flox}$

mice compared to control mice, meaning that GFI1 could have an impact on the production of

this particular γδT cell subset when deleted at the DN2/3 stage (**Suppl. Fig. 7B-D**). We also

observed a slightly increased proportion of RORγt⁺ cells in the spleen and a slightly increased

frequency of IL17A⁺ cells in the thymus, but nothing comparable to what was found in the total

GFI1 KO mice or in the *vav-cre Gfi1$^{fl/fl}$* mice. This strongly suggests that the DN2/3 T cell

precursors are not the source of the accumulated Vγ6⁺ γδT cell population seen in mice lacking

GFI1. This was also confirmed when we cultured DN2/3 precursors on OP9-DL4 cells in the

presence of IL7 and FLT3 (**Suppl. Fig. 7I-J**). After several days of coculture, WT DN2/DN3

cells differentiated as expected mostly into DP cells, while GFI1 KO DN2/DN3 cells were still

mostly found in the DN compartment and did not differentiate into DP cells (**Fig. 7D-F**). Both

WT and GFI1 KO DN2/DN3 cells were able to produce γδ T cells, but DN2/DN3 from GFI1 KO

thymus produced a smaller amount of γδ T cells compared to WT DN2-3 cells (Fig. **7D-F**)

reducing the numbers of the CD24⁻γδT cell subtypes (**Fig 7G**). The γδ T cells differentiated from

DN2/3 lacking GFI1 did not show a preference to generate higher proportions of CD24⁻Vγ1⁻Vγ4⁻

γδ T cells over the other subtypes but produced all γδ T cell populations in similar proportions to

WT DN2-3 cells (**Fig. 7D, 7E & 7G**). We further tested whether a developmental link exists
between GFI1 KO DN1 cells and the Vγ6$^+$ γδ T cells by also culturing sorted DN1 cells from WT
or GFI1 KO mice on OP9-DL4 cells in the presence of IL7 and FLT3 (**Suppl. Fig. 7I-J**). As
expected, WT DN1 cells were able to progress though the different stages of T cell
differentiation, while GFI1 KO DN1 remained mostly in the DN1 stage after 8 or 18 days of
culture (**Fig. 7 H-K**). Only very few cells developed into γδ T cells from WT DN1 and GFI1 KO
DN1 cells, with a peak of γδT cell production after 12 days for WT precursors, while GFI1 KO
DN1 cells produce most of the γδ T cells earlier in the coculture before being lost (**Fig. 7L**). We
gated the cells surviving in the coculture on CD24$^-$γδT cells and found that the GFI1 KO did
indeed produce higher proportions of Vγ1$^-$Vγ4$^-$ γδ T cells (most likely meaning Vγ6$^+$) compared
to WT DN1 cells (**Fig. 7M**). However, the number of cells produced was very low (**Fig. 7M**)
suggesting that GFI1 KO DN1 cells most likely require other signals to either generate the Vγ6$^+$
γδ T cells or to fully support this biased γδ T cell population leading to their accumulation in
adult Gfi1 deficient mice. Our data indicate that the expanded γδT17 cell population in GFI1 KO
mice are unlikely to originate from the DN2-DN3 cells. Instead DN1 subsets, particularly DN1e
may contribute to this bias. However, this interpretation remains tentative, as additional
experiments are required to confirm the developmental origin of these cells.

GFI1 can directly regulate *Maf*.
The scRNA-seq analysis indicted that the expression of *Maf* was increased in clusters from GFI1
KO cells and flow cytometry data confirmed this for the c-MAF protein, known to be a regulator
of γδT17 cell differentiation and maintenance[68]. We also confirmed overexpression of MAF in
GFI1 KO Vγ1$^-$Vγ4$^-$ thymic γδ T cells and also in Vγ1$^+$ thymic γδ T cells compared to WT control
cells (**Fig. 8A & 8B**). However, this upregulation of MAF was only associated with a higher
expression or RORγt in the Vγ1$^-$Vγ4$^-$ thymic γδ T cells (**Fig. 8A & 8B**). We observed similar
data in splenic γδ T cells, with an upregulation of c-MAF in GFI1 KO Vγ1$^-$Vγ4$^-$ and Vγ1$^+$ splenic
γδ T cells, associated with a strong RORγt upregulation only in Vγ1$^-$Vγ4$^-$ splenic γδ T cells (**Fig.
8C & 8D**). We also found that total DN3 cells from GFI1 KO mice that have a very different
transcriptome profile compared to WT DN3 cells showed up regulation of *Maf* (**Suppl. Fig. 8A**),
which was also confirmed by flow cytometry for DN2 and DN3 cells (**Fig. 8E**).

In addition, we also found higher expression of c-MAF specifically in the GFI1 KO DN1e

cells compared to WT cells (**Fig. 8E**), but not in DN1d or DN1a/b cells. This correlated with the

high expression of GFI1 shown in DN1e cells (**Suppl. Fig. 6G**). This data suggested that *Maf* is a

potential direct GFI1 target. To test this hypothesis, we interrogated available ChIP-Seq data

from MLL-ENL lymphoblastic cells and observed several peaks with a GFI1 binding motif

binding in the *Maf* 5' region and peak for GFI1 (**Fig. 8F**). Binding of GFI1 on *Maf* promoter was

also found in other available ChIP-Seq data with different cell types (**Suppl. Fig. 8B**). A query of

the JASPAR database confirmed the location of several GFI1 binding motifs of the *Maf* gene

from the transcription start site up to several kb upstream (**Fig. 8F**). ChIP-qPCR with purified

WT and GFI1 KO DN cells and primers covering these putative binding sites indicated binding

of GFI1 located in regions where the primer pairs were designed (**Fig. 8G**). As controls, we used

GFI1 KO DN cells, where no enrichment was found in this region and the   promoter, where

GFI1 does not bind, confirming occupation of the Maf promoter by GFI1 in WT DN pre-T cells

(**Fig. 8G**). This suggested that GFI1 directly represses the expression of *Maf* and with this the

expansion of Vγ6$^+$ γδ T cells.

Discussion

In this study, we describe expression of GFI1 in γδ T cells and their precursors and

present evidence for a novel regulatory role of this transcription factor in restricting γδ T cell fate

and cellularity across various tissues, including thymus, spleen, lung, and lymph nodes. Our

findings demonstrate that genetic ablation of GFI1 results in a substantial expansion of

RORγt$^+$Vγ6$^+$γδT cells producing IL17 and a concurrent decrease in Vγ1$^+$ and Vγ4$^+$ γδT cells.

This bias toward the γδT17 subtype emerges postnatally in GFI1 KO mice and persists in the

adult thymus, where the Vγ6$^+$ subset secreting IL17 becomes the predominant population among

γδ T cells. Furthermore, GFI1 KO mice exhibit an increased number of c-MAF$^+$/RORγt$^+$ cells in

the DN1e subset, which has been described as containing precursors for γδT17 cells[40,41,62]. *Gfi1*

deficient DN thymocytes and γδ T cells exhibit an upregulation of the bZip transcription factor c-

MAF, which is known to regulate γδT17 fate[68]. We further demonstrate that GFI1 occupies the

promoter region of the *Maf* gene at a cognate GFI1 binding site, suggesting that GFI1 controls

the cellularity of γδ T cells and their progenitors through the potential regulation of a c-MAF-

dependent transcriptional network.

The specific expansion of only Vγ6$^+$ γδT cells without affecting the fate of other γδT

subtypes (Vγ4$^+$), which could also adopt a γδT17 profile[33,57,69], supports the idea of an intrinsic

effect of GFI1 in γδ T cell development. In addition, the experiments with *vav-cre Gfi1$^{fl/fl}$* mice,

which delete *Gfi1* only in hematopoietic cells, also ruled out that the non-hematopoietic

environment such as epithelial cells in GFI1 KO mice are causing the specific expansion of Vγ6$^+$

γδT17 cells seen in these animals. Transplantation experiments of WT cells into GFI1 KO

animals did not reproduce the strong bias towards the γδT17 subtype observed in GFI1 KO mice,

even though we could observe a slight increase of both IL17A and RORγt frequencies in donor

WT γδ T cells in the spleen of the GFI1 KO recipients. This effect could be due to the

transplantation itself or may suggest that the environment of the GFI1 KO spleen could have a

slight impact on the fate of Vγ4$^+$ γδ T, which is the other γδ T subset capable of becoming γδT17

cells.

However, it remained unclear from which cell population the expansion of Vγ6$^+$ γδT17 in

GFI1 KO mice originates. We could show that Vγ6$^+$ γδT17 cells start to increase in numbers in

GFI1 KO mice only after birth. It is thus possible that *Gfi1*-deficient precursors from the fetal

thymus for Vγ6$^+$ γδT17 cells continue to produce these cells when they reach adult tissues,

whereas their WT counterparts would cease their production. Transplantation of adult BM cells

from GFI1 KO mice did not recapitulate the expansion of Vγ6$^+$ γδT17 cells in recipient animals

(data not shown), which excluded the presence of a γδ T precursor within the adult thymic DN

pre-T cell population. It is thus more likely that precursor cells from fetal thymus give rise to a c-

MAF$^+$/RORγt$^+$ cells in the DN1e subset in the adult thymus where they are under the restriction

of GFI1 in WT mice. These cells were seen in flow cytometry analyses in WT mice, albeit at

very low numbers. It is conceivable that these cells expand into Vγ6$^+$ γδT17 cells in the thymus

when GFI1 is absent. Both scRNA-seq and flow cytometry show this expanded population of c-

MAF$^+$/RORγt$^+$ cells in the DN1e pre-T cells from GFI1 KO mice very clearly. The transcription

pattern of this DN1e subpopulation in GFI1 KO mice supports this notion and suggests that a

DN1e subset exists with a specific transcriptional signature which is maintained after birth in the

adult thymus that is associated with Vγ6$^+$ γδ T cells producing IL-17. Although these data point

to a fetal origin of the expanded Vγ6$^+$ γδT17 cell population in GFI1 KO mice, further

experimentation with fetal and adult pre-T cells are necessary to firmly support such a

conclusion.

The decrease of Vγ1$^+$ and Vγ4$^+$ γδ T cells that we observed in GFI1 KO mice alongside

the expansion of Vγ6$^+$ γδT17 could be a consequence of the decreased cell numbers in the DN3

subset. Lower numbers of Vγ1$^+$ and Vγ4$^+$ γδ T cells were observed in fetal and adult thymus of

GFI1 KO mice as compared to WT controls, in contrast to Vγ6$^+$ γδ T cells, which expanded only

after birth. It is thus likely that separate precursors exist for the Vγ1$^+$ and Vγ4$^+$ γδ T cells and the

Vγ6$^+$ subset. In addition, DN3 cells lacking GFI1 have a distinct transcriptomic profile from their

WT counterparts, which could also affect the production of Vγ1$^+$ and Vγ4$^+$ γδ T cells, since they

develop from this thymic DN population. Surprinsingly, an upregulation of MAF was also

observed in the thymic and splenic Vγ1$^+$ γδ T cells but not in the Vγ4$^+$ γδ T cells, the other

population where MAF can be expressed. We didn't observe a differential expression of GFI1

(data not shown) between the different γδ T cell populations which could have explained this

difference in MAF regulation. Most likely, additional regulatory or compensatory pathways exist

that regulate MAF expression in different γδ T cell subtypes in addition to GFI1.

A DN1-like (CD117$^-$CD24$^-$CD25$^-$CD44$^+$) precursor subpopulation (DN1e) was more

recently identified to be programmed or pre-committed to become Tγδ17 cells independently of

TCR signaling[40,41]. These cells express *Rorc, Sox4, Tcf7, Tcf12, Maf, Il7r, Scart2*, and *Blk* and

are predisposed to become CCR6[+] IL17A-producing cells[41]. c-MAF promotes the expression of

genes essential for Tγδ17 cell identity, such as *Rorγt* and *Blk*, which are key players in the Tγδ17

program, and which are also found to be increased in expression in GFI1 KO γδT cells[68,70]. Our

data suggest that lineage committed precursors and γδT17 cells are not only controlled by TCF1

and HEB/SOX13[71,72], which regulate c-MAF, but also by GFI1, which acts upstream of c-MAF.

This would be coherent with the model that lineage fate can be prewired cell-intrinsically and is

not necessarily specified by antigen receptor signals[41]. This also lends further support to the view

that the MAF[+]/ RORγt[+] cell population that we found expanded in thymic DN1e cells from GFI1

KO mice represents the precursor for the expanded IL17 producing Vγ6[+] γδ T cells. However,

further experimentations would be needed to support this hypothesis, because the coculture of

DN1 cells from GFI1 KO mice on OP9-DL4 layer or in FTOC (data not shown) was not robust

enough to support the expansion of Vγ6[+] γδ T, even though a bias was observed towards this

subtype. It may suggest that other signals in the periphery such as the cytokine environment

specifically found in GFI1 KO mice may be required to support the intrinsically biased γδ T

populations to expand in *Gfi1* deficient mice.

It has previously been shown that c-MAF is highly expressed in RORγt[+] γδ T cells, which

are γδT17 cells, but not in other γδ T cell subsets like γδT1 cells[68]. c-MAF functions by

promoting the expression of key genes in the type 17 program (e.g., *Rorc, Il17a, Blk*) and

possibly also *Il23R* and antagonizes negative regulators like TCF1, which are required to

upregulate *Ifng*. c-MAF helps γδ T cells adapt to specific tissue microenvironments by regulating

metabolic pathways and integrates signals from IL1β, IL23 and interactions with epithelial cells

to fine-tune γδ T cell responses[73]. These reports identified c-MAF as a universal and essential

regulator of γδT17 cell differentiation and function but still lacked insights into the molecular

mechanisms that control c-MAF expression.

Our results with ChIP-qPCR and ChIP-seq data from the public domain showed that GFI1

occupies the *Maf* promoter region. This observation together with our findings that c-MAF

expression is upregulated in all tested *Gfi1* deficient γδ T cells indicates that GFI1 acts as an

upstream negative regulator of c-MAF. It is therefore conceivable that the direct transcriptional

repression of the *Maf* gene by GFI1 is necessary to maintain and restrict the development of Vγ6

γδT17 cells and that absence GFI1 leads to the developmental defect in these γδ T cells seen in

GFI1 KO animals. The upregulation of c-MAF and c-MAF target genes such as *Il17, Blk* support

such a role of GFI1, but further studies are needed to establish a direct mechanistic link.

It is also known that GFI1 antagonizes RORγt expression through GATA3[3,23,70] and that -

c-MAF induces expression of RORγt and synergizes with RORγt to regulate IL17-producing

cells. It is thus conceivable that a GFI1/c-MAF/RORγt regulatory circuit exists that controls the

expression of IL17A and ensures the cellularity and functionality of γδT17 cells. GFI1 has been

shown to regulate Notch/IL7 signaling[18], both important for γδ T cells, suggesting that it could be

involved at different levels, possibly also in γδ T cell development. Our data do not exclude

contribution from these pathways and further experiments would be required to decide whether

the link between Notch/IL7 signaling and GFI1 bears any role in γδ T cells.

The ability of c-MAF to fine-tune IL17 production has been highlighted as a potential

therapeutic target for controlling γδ T cell-mediated inflammation without compromising their

protective roles[74]. Strategies to modulate c-MAF activity or its interaction with RORγt are being

explored as therapeutic options to balance inflammation and immunity. Our findings which place

GFI1 upstream of c-MAF may point to new possibilities to interfere with this regulatory circuit

since GFI1 is in a complex with LSD1 for which small molecule inhibitors exist[75].

Materials and Methods

Mice

The generation of GFI1 KO, GFI1-EGFP, and GFI1-fl/fl mice was previously
described[13,28,63]. C57BL6/J CD45.2 congenic mice (B6.SJL-*Ptprc^a Pepc^b*/BoyJ) and *Vav-cre*
mice (B6.Cg-Tg(VAV1-cre)1Graf/MdfJ) were obtained from Jackson labs (Bar Harbour, MN)**.**
C57BL6/J CD45.1/2 mice were a gift from Dr. Javier Di Noia, IRCM, Montreal and hCD2-iCre
mice (B6.Cg-Tg(CD2-icre)4Kio/J) were a gift from Dr. André Veillette, IRCM, Montreal. All
mice were maintained on the C57BL6/J background. Mice were housed and bred in the IRCM
specific pathogens-free facility in a 12 h dark / light cycle with food and drink at libitum. All
mouse work was reviewed and approved by the animal protection committee at the IRCM
(protocols #2020-09), according to the guidelines of the Canadian Council of Animal Care.

Tissue dissociation

Lungs from adult mice (at least 5 weeks old) were perfused with 10 ml of PBS *via* the
right ventricle and then were cut into small pieces with scissors and a scalpel. Lung pieces were
dissociated and digested in a buffer with collagenase IV (1 mg/ml) (Gibco) and DNase I (0.2
mg/ml) (Sigma) in RPMI medium for 30 min at 37°C. Cells were aspirated several times with a
syringe and needle (18 gauge) every 10 min to further the dissociation and digestion process.
Lung cell suspensions were filtered and washed in PBS. Thymi, spleen, and lymph nodes from
adult mice (at least 5 weeks old) were mechanically dissociated to obtain a cell suspension which
were filtered. Then, red blood cell lysis buffer (Sigma) was added to the lung, thymic, splenic or
LN cell pellet for 4 min and washed before staining or sorting.

Bone marrow chimera

Bone marrow (BM) cells from CD45.1/2 mice were obtained by flushing femurs and
tibiae with cold PBS using a syringe with 23G needle. The red blood cells were subsequently
lysed, and the cells were washed in PBS before being injected by the tail vein in 100 µl of PBS
containing 2x10^6 BM cells into either CD45.2 WT or GFI1 KO recipient mice. 4 or 5 weeks after
the transplantation, the recipient mice were sacrificed, and splenic, thymic, and lung cells were
analyzed by flow cytometry.

OP9-DL1 and OP9-DL4 culture

534
OP9-DL1 and OP9-DL4 cells were a gift from the lab of Dr. Zúñiga-Pflücker[76]. OP9-DL1
cells were plated in 96 flat-bottom well plate at a density of 3000 cells in 100 uL of OP9 medium
(AMEM, 20% FBS, 1% Pen/strep, 2.2g/L sodium bicarbonate, 50 μM β-mercaptoethanol). The
day after, the same numbers of WT and GFI1 KO sorted DN1, DN2-DN3 or total DN cells were
co-cultured on OP9-DL1 or OP9-DL4 monolayers in DMEM-glutamax medium containing 10%
of FBS, 1% Pen/strep, 50 μM β-mercaptoethanol, 10 mM Hepes with Flt3 (5 ng/mL, Peprotech)
and IL7 (1 ng/mL, Peprotech). Thymocytes were transferred to a new OP9-DL1 or OP9-DL4
monolayer with fresh cytokines every 3 days.

Flow cytometry and cell sorting

For the analysis of cells from spleen, lung, lymph nodes and thymus from mice by flow
cytometry, a Fortessa (BD Biosciences) flow cytometer and FlowJo software were used. Cell
sorting was performed on a FACSAria III Sorter (BD Biosciences). For IL17A and IFNγ
intracellular staining, cells were cultured for at least 3h with GolgiPlug (BD Biosciences), PMA
(50μM, Sigma) and ionomycin (1mM, Sigma) and stained using the eBioscience cytokine
intracellular staining kit (Invitrogen). For IL22 staining[77], cells were first cultured with IL23 (20
ng/mL, R&D Systems) overnight before stimulation with PMA (50μM, Sigma) and ionomycin
(1mM, Sigma) in the presence of GolgiPlug (BD Biosciences) for the last 4 hours. Intracellular
staining for the transcription factors RORγt and c-MAF was also done using eBioscience FoxP3
intracellular kit (Invitrogen). For phosphoSTAT3 intracellular staining, after the surface staining,
cells were stimulated for 15 minutes at 37°C with either IL23 (20ng/mL, R&D Systems) or IL7
(10 ng/mL, Peprotech) and washed with cold PBS. Cells were subsequently fixed for 15 minutes
at 37°C with Biolegend fixation buffer[78] before being permeabilized with the Biolegend True-
phos Perm buffer (425401) for 30 minutes at 4°C. After washing, the cells were incubated with
Phospho-STAT3 antibody for 30 min at 4°C. For the splenic γδ T cell culture, sorted splenic γδ T
cells were cultured *in vitro* in DMEM medium containing 10% FBS, 1% Pen/Strep, 10 mM
Hepes and 50μM β-mercaptoethanol in the presence or not of IL23 (20 ng/mL, R&D Systems)
and IL1β (10 ng/mL, Biolegend) for 16 hours and GolgiPlug (BD Biosciences) was added for the
last 4 hours, before cytokine staining. All antibodies directly conjugated to a fluorochrome or to
biotin were purchased from BioLegend, eBiosciences, or BD Biosciences (Table 1).

PCR

RNA extraction was performed using Tri-Reagent (Life Technologies) for cell numbers between 1×10^6 and 3×10^6 or using the RNeasy Micro kit from QIAGEN for low numbers of cells following the company's instructions (RNeasy Micro kit; QIAGEN). RT-PCR was performed using Superscript II (Invitrogen). RNA was incubated with oligodeoxy thymidine (Invitrogen), EDTA (Invitrogen), and dNTP (Invitrogen) for 10 min at 65°C and then DTT (Invitrogen), Superscript buffer 5× (Invitrogen), RNaseOUT buffer (Invitrogen), and the enzyme Superscript II (Invitrogen) were added and the reaction was subjected to PCR cycles (step 1, 25°C for 10 min; step 2, 42°C for 60 min; step 3, 70°C for 15 min) to generate cDNA for real-time quantitative PCR. Real-time quantitative PCR was performed in triplicates on the ViiA7 real-time PCR machine (Applied Biosystems) in SYBR Green Master Mix (Applied Biosystems) containing specific primers for mouse. The primers used are *Gfi1 Forward*: GTGCCATCAGAGGAGGTGAA, *Gfi1 Reverse*: TCTGATAACCTGCGGCCAAT; *Gfi1b Forward*: CAGTCCTACGCCCACCTTGG, *Gfi1b Reverse*: AGTGTCCGAGTGGATGAGCA, and *Gapdh Forward*: ACTGAGCAAGAGAGGCC, *Gapdh Reverse*: TATGGGGGTCTGGGATGGAA.

Single cell RNA-seq sequencing

The single-cell RNA sequencing on lungs was performed using 10x Genomics platform with the Chromium Next GEM Single Cell 3 Kit v3.1. Total lung cells from adult mice (between 5 and 8 weeks old) were sorted on live cells using Hoecsht to stain the dead cells. Then, cells were loaded on a Chromium controller on a chip G to target a recovery of 10,000 cells per sample and library were constructed following the manufacturer recommendations. Library size distribution was assessed on a 2100 bioanalyzer (Agilent Technologies) using a High Sensitivity DNA Kit and libraries were quantified by qPCR. Equimolar libraries were sequenced in paired-end reads (PE 28-91), on a Novaseq 6000 system (Illumina), with an average depth of 25,000 reads per cell. The single-cell RNA sequencing on sorted γδ T cells (from the spleen or the thymus DN1-DN2 of adult mice) or on sorted DN1-DN3 cells was performed using 10x Genomics platform with the Chromium Next GEM Single Cell 3 v3.1 Kit and the 3' Feature barcode kit for cell surface protein. To accelerate the sorting step, splenic cells or thymic cells were preselected to remove B cells (CD19), myeloid cells (Gr1, CD11b and CD11c) and NK

cells (NK1.1 and CD49b) using biotinylated antibodies and Mojo magnet selection (Biolegend),

before FACS staining. CD8$^+$ cells were also partially removed from the thymus using Mojo

Magnet selection (Biolegend). Then, WT or GFI1 KO cells were stained with TotalSeq-B oligo-

conjugated antibodies (Totalseq-B0302 & Totalseq-B0303, Biolegend) to hashtag the samples in

order to reduce costs and decrease technical variability during the incubation with the

fluorochrome coupled antibodies for the FACS sorting. Sorted cells were loaded on a Chromium

controller on a chip G to target a recovery of 20,000 cells per sample (10,000 cells per condition

WT or KO) and library were constructed following the manufacturer recommendations. Library

size distribution was assessed on a 2100 bioanalyzer (Agilent Technologies) using a High

Sensitivity DNA Kit and libraries were quantified by qPCR. Equimolar libraries were sequenced

in paired-end reads (PE 100), on a Novaseq X system (Illumina). An average depth of 25,000 or

5,000 reads per cell was targeted for gene expression or hashtag libraries, respectively. Duplicatat

were used for all the single cell RNA-sequencing experiments, meaning 2 WT and 2 GFI1 KO

samples for each cell types.

Single cell RNA-seq analyses

scRNAseq data was analyzed using python (v3.12.4) and the package scanpy (v1.10.4).

Cells with <500 reads, <200 genes or >20% mitochondrial genes were removed as outliers.

Importantly, ribosomal genes, predicted genes with *Gm*-identifier and mitochondrial genes were

excluded for subsequent analysis. The samples were demultiplexed using hashsolo[79]. The

datasets were normalized using the normalize_total function with a target sum of 10000, followed

by log1p transformation and scaling with a maximum value of 10. The top 2000 highly variable

features were calculated using the flavor "seurat_v3" and the raw counts. Dimensionality

reduction was performed with default settings, while the resolution for clustering using the

Leiden algorithm was set to 0.5 for the DN dataset, 0.8 for the thymic γδ T cells and lung datasets

and 1 for the splenic γδ T cells dataset. To characterize the clusters, differential gene expression

analysis was done using the rank_genes_groups function. Graph abstraction was performed using

the function paga in scanpy with default parameters using clusters as partitions[80].

Chromatin immunoprecipitation (Ch-IP)

Ch-IP for GFI1 was performed on 10×10^6 fresh cells. Briefly, cells were cross-linked

with 1% formaldehyde for 10 min before quenching with 125mM glycine for 5 min. Cells were

initially lysed in cell lysis buffer (5mM PIPES, 85mM KCl, 0.5% Nonidet P-40, freshly added

protease inhibitors mixture, and 0.8mM PMSF) for 15 min on ice with occasional vortexing.

After centrifugation at 4°C at 4000 rpm for 5 min, the pellet was resuspended in nuclei lysis

buffer (50mM Tris [pH 8], 10mM EDTA, 0.1% SDS, freshly added protease inhibitors mixture,

and 0.8mM PMSF) and incubated on ice for 20 min with occasional vortexing. The lysed cells

were sonicated using a Covaris E220 to generate 200–600-bp fragments. One volume of

immunoprecipitation dilution buffer (300mM NaCl and 2% Triton X-100) was added to

sonicated nuclei lysates and was immune-precipitated with 5μg of anti-Gfi1 (AF3540; R&D

Systems). Beads were added to harvest protein☐DNA complexes. After de-crosslinking, the

eluate was digested with RNase A and proteinase K. The chromatin☐associated DNA was

extracted using QIAquick PCR Purification Kit (Qiagen). The primers for the ChIP-PCR are the

following: *Gfi1 Promoter Forward*: AATTCGGGAGCTGAAGGCAA, *Gfi1 Promoter Reverse*:

AGAGGGGCACACAGTTTAGC; *Maf-1 Forward*: CTCTCGAACTCCGAGGAAAGC, *Maf-1*

Reverse: GAGCGGCAGCGATTAGCAT; *Maf-2 Forward*: TGGTCAATAATGACAAGCGCA,

Maf-2   Reverse: ATACAAATCGGAACCCCGGC;   *Maf-3   Forward*:

TCAATCAGTGTAGAAAACTGAACC, *Maf-3 Reverse*: ACAAGTCCGGAAACCCAGAT; ;

Il6 Promoter   Forward: GGCGGAGCTATTGAGACTGT, *Il6 Promoter Reverse:*

AAACCGGCAAGTGAGCAGAT and *Chr2 Forward*: TGGGCATATCCCTGGAGCTT, *Chr2*

Reverse: GGCCATCCCACAGTCACAAC.

Statistical analysis

Multiple paired *t*-test or paired *t*-test was used to calculate p values where indicated. A p

value ≤0.05 was considered as statistically significant. Survival curves were analyzed by log-rank

Mantel–Cox test using GraphPad Prism (GraphPad Software, La Jolla, CA). The p values are as

follows: *p < 0.05, **p < 0.01, ***p < 0.005, and ****p < 0.001.

Data availability and resources

The primary read files and the raw counts for all single-cell sequencing datasets reported

in this paper are available to download from the Gene Expression Omnibus under accession

number GSE294462. The analysis of sc-RNA-seq data and codes for each Figure containing sc-
RNA-seq data are also deposited on GEO. Codes are available upon request. All data supporting
the findings of this study are available within the paper and its Supplementary Information.
Antibodies are listed on a separate table under supplementary information.

Authors' Disclosures

The authors have nothing to disclose.

Authors' Contributions

J. Fraszczak: Conceptualization, data curation, formal analysis, validation, investigation,
visualization, methodology, writing original draft, revising, and editing. D. Obwegs: Single cell
RNA-seq analysis and interpretation. K. Arman: ChIP-PCR experiments and analysis. T. Muralt:
data curation, formal analysis. Sagar: Single cell RNA-seq analysis and interpretation, writing,
and editing. I King: support with materials, resources, and reading manuscript. B. Thurairajah:
support with materials, resources. H. Melichar: support with reading manuscript. E. Mallet
Gauthier: support with experiments. T. Moroy: Conceptualization, resources, data curation,
supervision, funding acquisition, investigation, visualization, writing original draft, project
administration, and editing.

Acknowledgments

The authors thank the IRCM animal technicians, the IRCM flow cytometry facility staff
for help with cell sorting, the IRCM molecular genomics facility staff for performing single cell
RNA-seq experiment and advice, the IRCM bioinformatic facility for processing single cell
RNA-seq data. The authors thank Mathieu Lapointe for the mouse genotyping and all the Dr
Moroy's lab members for advice. This work was supported by the Canadian Institutes for Health
Research (CIHR) through a Foundation grant (FDN - 148372) and by the IRCM Foundation.

Figure Legends

Fig. 1: Phenotype of γδ T cells in adult GFI1 KO mice

A & **B**: UMAP representation based on gene expression profiling by sc-RNA-seq depicting 11

clusters in the CD3$^+$ lymphocyte compartment in the lung tissue of WT and GFI1 KO mice

(**A**) and the defined subsets labeled in different colors (**B**). **C**: Bar plots showing the distribution

of the defined populations in the indicated genotypes. **D**: Representative plots of lung, splenic

and LN cells from WT and GFI1 KO mice analyzed by flow cytometry and stained to identify γδ

T cells (CD45$^+$CD4$^-$γδT$^+$) after gating on live cells (according to the FSC/SSC) and singlets. **E** &

F: Cell numbers (**E**) and frequencies in CD45$^+$ cells (**F**) of γδ T cells (CD4$^-$γδT$^+$) in the lung,

spleen and LN of WT and GFI1 KO mice. **G**: Representative plots of lung, splenic and LN cells

from control and *vav-cre GFI1$^{flox/flox}$* mice analyzed by flow cytometry and stained to identify γδ

T cells (CD45$^+$CD4$^-$γδT$^+$) after gating on live cells (according to the FSC/SSC) and singlets. **H** &

I: Cell numbers (**H**) and frequencies in CD45$^+$ cells (**I**) of γδ T cells (CD4$^-$γδT$^+$) in the lung,

spleen and LN of control and *vav-cre GFI1$^{flox/flox}$* mice. Adult mice meaning mice older than 5

weeks. Statistical testing was performed with paired *t*-test using Prism v9 (GraphPad). *p*-Values

are shown as * < 0.05, ** < 0.01, *** < 0.005, and **** < 0.001 where each statistical

significance was found.

Fig. 2: Expansion of the RORγt$^+$Il-17a$^+$ γδT subset in adult GFI1 KO mice.

A: UMAP representation showing expression of *Blk, Maf and Rorc* genes in the CD3$^+$

lymphocyte compartment in the lung tissue of WT and GFI1 KO mice. Color represents the

normalized expression of the gene. **B:** Frequencies of CD27$^+$ and CD27$^-$ cells in splenic, LN and

lung CD45$^+$CD4$^-$γδT$^+$ from WT and GFI1 KO mice. **C:** Representative normalized histogram of

RORγt$^+$ cells in either splenic, LN or lung CD45$^+$CD4$^-$γδT$^+$ and CD45$^+$CD4$^+$γδT$^-$ from WT and

GFI1 KO mice. **D:** Frequencies of RORγt$^+$ cells in splenic, LN and lung CD45$^+$CD4$^-$γδT$^+$

comparing WT and GFI1KO mice. **E:** Splenic, LN and lung CD45$^+$CD4$^-$γδT$^+$RORγt$^+$ cell counts

in WT and GFI1 KO mice. **F:** Representative plots of lung, splenic and LN cells from WT and

KO mice treated with PMA, ionomycin and Golgi Plug for 3.5 hours and analysed by flow

cytometry. The cells were gated on singlets CD45$^+$CD4$^-$γδT$^+$ before gating for IFNγ and IL17A.

G: Frequencies of IL17A$^+$ in splenic, LN and lung CD45$^+$CD4$^-$ γδT$^+$ comparing WT and GFI1

KO mice. **H**: Cell numbers of splenic, LN and lung CD45$^+$CD4$^-$γδT$^+$IL17A$^+$ in WT and GFI1 KO

mice. Adult mice meaning mice older than 5 weeks. Statistical testing was performed with

multiple paired *t*- test for **B** and paired *t*-test for **D-E** and **G-H** using Prism v9 (GraphPad). *p*-

Values are shown as * < 0.05, ** < 0.01, *** < 0.005 and **** < 0.001 where each statistical

significance was found.

Fig. 3: Expansion of Vγ6$^+$ γδT17 cells in the periphery of adult GFI1 KO mice.

A: UMAP representation based on gene expression profiling by sc-RNA-seq depicting 14

clusters in sorted splenic γδ T cells of WT and GFI1 KO mice. **B**: Dot plot showing the key

differentially expressed genes in each cluster shown in Fig. 3A. Color represents the scaled mean

expression of the gene in the respective cluster, and dot size represents the fraction of cells in the

cluster expressing the gene. **C** and **D**: UMAP representation of sorted splenic γδ T cells of WT

and GFI1 KO mice colored by cell type (**C**) or genotype (**D**). **E**: Bar plots showing the

distribution of cell types found in splenic γδ T cells in the indicated genotypes. **F**: Differential

gene expression showing key up- or down- regulated genes in GFI1 KO splenic γδ T cells versus

WT splenic γδ T cells. **G**: Representative plots of Vγ5$^+$ and Vγ6$^+$ in γδT$^+$ cells from WT and

GFI1 KO spleen and lung after gating on live cells (according to FSC/SSC) and singlets. **H**:

Vγ1$^+$, Vγ4$^+$, Vγ5$^+$ or Vγ6$^+$ γδT$^+$ cell counts in the spleen and lung of adult WT and GFI1 KO

mice. **I**: Cell numbers of Vγ1$^-$Vγ4$^-$Vγ5$^-$ γδT$^+$ cells at different time point in WT and GFI1 KO

spleen, with adult meaning mice older than 5 weeks. Adult mice meaning mice older than 5

weeks. Statistical testing was performed with multiple Mann-Whitney tests using Prism v9

(GraphPad). *p*-Values are shown as * < 0.05, ** < 0.01, *** < 0.005, and **** < 0.001 where

each statistical significance was found.

Fig. 4: Expansion of Vγ6$^+$ γδT17 cells in the thymus of adult GFI1 KO mice.

A: Representative plots of thymic CD45$^+$CD4$^-$γδT$^+$ cells in WT and GFI1 KO mice after gating

on live cells (according to FSC/SSC) and singlets. **B**: γδ T cell counts (CD45$^+$CD4$^-$γδT$^+$, left) and

frequencies in CD45$^+$ cells (CD4$^-$γδT$^+$, right) in WT and GFI1 KO thymus. **C**: Representative

plots of thymic γδ T cells (CD45$^+$CD4$^-$γδT$^+$) treated with PMA, ionomycin and Golgi Plug for

3.5 hours and stained for IL17A and IFNγ. The cells were gated on singlets CD45$^+$CD4$^-$γδT$^+$

before gating for IFNγ and IL17A. **D**: CD45$^+$CD4$^-$γδT$^+$IL17$^+$ cell counts (left) and IL17A$^+$ cell

frequencies in CD45$^+$CD4$^-$γδT$^+$ (right) in WT and GFI1 KO thymus. **E**: Representative

normalised histogram for RORγt and control isotype (Ctrl) in CD45$^+$CD4$^-$γδT$^+$ from WT and

GFI1 KO thymic cells. **F**: CD45$^+$CD4$^-$γδT$^+$RORγt$^+$ cell counts (left) and RORγt$^+$ cell frequencies

in CD45$^+$CD4$^-$γδT$^+$ (right) in WT and GFI1 KO thymus. **G**: UMAP representation based on gene

expression profiling depicting 14 clusters in sorted thymic γδ T cells of WT and GFI1 KO mice.

H: Dot plot showing the key differentially expressed genes in each cluster shown in Fig. 4G.

Color represents the scaled mean expression of the gene in the respective cluster, and dot size

represents the fraction of cells in the cluster expressing the gene. **I** and **J**: UMAP representation

of sorted thymic γδ T cells of WT and GFI1 KO mice colored by cell type (**I**) or genotype (**J**). **K**:

Bar plots showing the distribution of each cell type found in thymic γδ T cells in the indicated

genotypes. **L**: Thymic Vγ6$^+$ γδT$^+$ cell counts in adult WT and GFI1 KO mice based on cytometry

staining. **M**: Cell numbers of CD24$^-$ γδT$^+$ in adult thymus of WT and GFI1 KO mice or control

(*vav-cre GF1$^{wt/wt}$*) and *vav-cre GFI1$^{flox/flox}$* mice. **N**: Representative plots of CD24 expression in

thymic γδ T cells and Vγ chains expression in thymic CD24$^-$ γδ T cells from adult WT and GFI1

KO mice. **O & P**: Vγ chain frequencies (**O**) and Vγ chain cell counts (**P**) in thymic CD24$^-$ γδ T

cells from adult WT and GFI1 KO mice or control (*vav-cre GF1$^{wt/wt}$*) and *vav-cre GFI1$^{flox/flox}$*

mice. Adult mice meaning mice older than 5 weeks. Statistical testing was performed with paired

t-test or multiple paired *t*-test using Prism v9 (GraphPad). *p*-Values are shown as * < 0.05, ** <

0.01, *** < 0.005, and **** < 0.001 where each statistical significance was found.

Fig. 5: GFI1 intrinsically controls the generation of γδ T cells.

A: GFP$^+$ cells indicating *Gfi1* expression in either γδ T cells (CD45$^+$CD4$^-$γδT$^+$) or CD4$^+$ T cells

(CD45$^+$CD4$^+$γδT$^-$) from spleen, LN, lung, and thymus from GFI1:GFP reporter mice[63] (het) and

controls (wt). **B & C**: Relative *Gfi1* expression in γδ T and CD4$^+$ T cells from Spleen, LN and

Lung (**B**) and in thymic γδ T, DN1, DN2 and DN3 cells (**C**) measured by RT-PCR. *Gapdh* was

used as a housekeeping gene. **D**: Scheme representing the transplantation strategy of CD45.1/2$^+$

WT BM cells into either CD45.2$^+$ WT or GFI1 KO recipient mice. **E**: Representative plots

showing the CD45.1/2$^+$ donor cells in the spleen, thymus and lung of non-irradiated WT and KO

recipients. **F**: Frequencies measured by flow cytometry of either WT CD45.1/2$^+$ donor cells or

CD45.2$^+$ recipient cells in the spleen, lung and thymus of transplanted WT and GFI1 KO

recipient mice. **G**: Frequencies measured by flow cytometry of either WT donor γδ T cells (in

CD45.1/2$^+$ cells) or endogenous recipient γδ T cells (in CD45.2$^+$ cells) in the spleen, lung and

thymus of transplanted WT and GFI1 KO recipient mice. **H**: IL17A frequencies in either donor

WT γδ T cells (CD45.1/2$^+$ cells) or endogenous recipient γδ T cells (in CD45.2$^+$ WT or in

CD45.2$^+$ GFI1 KO cells) in the spleen, lung and thymus of transplanted WT and GFI1 KO

recipient mice. **I**: RORγt frequencies in either WT donor γδ T cells (CD45.1/2$^+$ cells) or

endogenous recipient γδ T cells (in CD45.2$^+$ WT or in CD45.2$^+$ GFI1 KO cells) in the spleen,

lung and thymus of transplanted WT and GFI1 KO recipient mice. **J**: Vγ1$^-$Vγ4$^-$ frequencies in

either WT donor γδ T cells (CD45.1/2$^+$ cells) or endogenous recipient γδ T cells (in CD45.2$^+$ WT

or in CD45.2$^+$ GFI1 KO cells) in the spleen, thymus and lung of transplanted WT and GFI1 KO

recipient mice. Statistical testing was performed with multiple paired *t*- test using Prism v9

(GraphPad). *p*-Values are shown as * < 0.05, ** < 0.01, *** < 0.005, and **** < 0.001 where

each statistical significance was found.

Fig. 6: Precursors in thymus of adult mice

A: Representative plots of WT or GFI1 KO thymic DN progenitors cultured on OP9-DL1 in the

presence of FLT3 and IL7 for 12-14 days. After gating on singlets CD45$^+$γδT$^+$CD24$^-$ cells,

mature CD24$^-$ γδT subsets were distinguished according to the expression of CD44 and CD45RB.

B: Frequencies of each indicated subset from Fig. 6A. **C**: DN cell counts in the thymus of WT

and GFI1 KO mice or control (*vav-cre GF1$^{wt/wt}$*) and *vav-cre GFI1$^{flox/flox}$* mice. DN1a/b cells were

characterized as CD4$^-$CD8$^-$γδT$^-$CD44$^+$CD25$^-$CD24$^+$Ckit$^+$, DN1d cells were characterized as CD4$^-$

CD8$^-$γδT$^-$CD44$^+$CD25$^-$CD24$^+$Ckit$^-$, DN1e cells were characterized as CD4$^-$CD8$^-$γδT$^-$

CD44$^+$CD25$^-$CD24$^{-/low}$Ckit$^-$, DN2 cells were characterized as CD4$^-$CD8$^-$γδT$^-$CD44$^+$CD25$^+$ and

DN3 cells were characterized as CD4$^-$CD8$^-$γδT$^-$CD44$^-$CD25$^+$. **D, E and F**: UMAP representation

based on gene expression profiling of pooled DN1, DN2 and DN3 from WT and GFI1 KO mice

colored by cluster (**D**), cell type (**E**) and genotype (**F**). **G**: Bar plots showing the distribution of

each cell type found in the pool of DN1, DN2 and DN3 cells in the indicated genotypes. **H**:

UMAP representation showing expression of *Maf*, *Rorc*, *Blk and Cxcr6* genes. Color represents

the normalized expression of the gene. **I**: Differential gene expression showing key up- or down-

regulated genes in GFI1 KO DN1-3 cells versus WT DN1-3 cells. **J**: Frequencies of c-

MAF$^+$RORγt$^+$ cells in DN1, DN2 and DN3 cells from WT and GFI1 KO mice or control and

vav-cre GFI1$^{flox/flox}$ mice. **K**: Representative plots showing in pink c-MAF$^+$RORγt$^+$DN1 cells in

the DN1 subsets characterized by the expression of c-Kit and CD24 after gating on lin$^-$CD4$^-$CD8$^-$

γδT$^-$CD44$^+$CD25$^{-/low}$ from WT and GFI1 KO thymus. (Lin$^-$ = CD19, CD11b, CD11c, NK1.1).

Adult mice meaning mice older than 5 weeks. Statistical testing was performed with multiple

paired *t*- test using Prism v9 (GraphPad). *p*-Values are shown as $* < 0.05$, $** < 0.01$, $*** < 0.005$,

and $**** < 0.001$ where each statistical significance was found.

Fig. 7: Potential of differentiation of precursors in thymus of adult mice

A & B: UMAP representation based on gene expression profiling by sc-RNA-seq of thymic

DN1, DN2, DN3 and γδ T cells from WT and GFI1 KO mice colored by cluster (**A**) and cell type

(**B**). **C:** Partition based Graph Abstraction (PAGA) graph after running PAGA on thymic DN1-3

pre-T cells and Trgv6$^+$ thymic γδ T cells from WT and KO at cluster resolution. **D &E**:

Representative plots showing the DP (CD45$^+$CD4$^+$CD8$^+$) population, the γδ T cells (CD45$^+$CD4$^-$

CD8$^-$γδT$^+$) and the CD24$^-$γδ T subsets according to the expression of Vγ chains after gating on

live singlets CD45$^+$CD4$^-$CD8$^-$γδT$^+$CD24$^-$ cells from sorted WT (**D**) and GFI1 KO (**E**) DN2/3

cells cultured on OP9-DL4 as described in Supp. Fig. 7I. **F**: Cell counts of generated DP cells

(CD45$^+$CD4$^+$CD8$^+$) and γδ T cells (CD45$^+$CD4$^-$CD8$^-$γδT$^+$) from WT and GFI1 KO DN2-3

cultured on OP9-DL4 as described in Supp. Fig. 7I at different time points of the coculture. **G**:

Cell counts and frequencies of the indicated subsets based on the expression of Vγ chains of

mature CD24$^-$ γδ T cells after coculture of sorted DN2-3 cells from WT or GFI1 KO mice on

OP9-DL4 cells as described in Supp. Fig. 7I. **H-K**: Representative plots showing the DN subsets

based on CD44 and CD25 expression after gating on CD45$^+$CD4$^-$CD8$^-$γδT$^-$ cells, DP

(CD45$^+$CD4$^+$CD8$^+$) population, the γδ T cells (CD45$^+$CD4$^-$CD8$^-$γδT$^+$) and the CD24$^-$γδ T subsets

according to the expression of Vγ chains after gating on live singlets CD45$^+$CD4$^-$CD8$^-$γδT$^+$CD24$^-$

cells from sorted WT (**H-J**) and GFI1 KO (**I-K**) DN1 cells cultured on OP9-DL4 as described in

Supp. Fig. 7 at different time points D8 (**H-I**) and D18 (**J-K**). **L**: γδ T cell counts generated from

WT or GFI1 KO DN1 cultured on OP9-DL4 as described in Supp. Fig. 7 at different time points.

M: Cell counts and frequencies of the indicated subsets of mature CD24$^-$ γδ T cells after

coculture of sorted DN1 cells from WT or GFI1 KO mice on OP9-DL4 cells as described in

Supp. Fig. 7I. Adult mice meaning mice older than 5 weeks. Statistical testing was performed

with paired *t*-test and multiple paired *t*- test using Prism v9 (GraphPad). *p*-Values are shown as *

< 0.05, ** < 0.01, *** < 0.005, and **** < 0.001 where each statistical significance was found.

Fig. 8: GFI1 regulation in adult mice

A: RORγ and c-MAF MFI in the indicated subsets of thymic γδ T cells based on Vγ1 and Vγ4

expression from WT and GFI1 KO mice. **B**: c-MAF and RORγt MFI quantification for the

indicated thymic γδ T cell subsets comparing WT and GFI1 KO mice. **C:** RORγ and c-MAF MFI

in the indicated subsets of splenic γδ T cells based on Vγ1 and Vγ4 expression from WT and

GFI1 KO mice. **D**: c-MAF and RORγt MFI quantification for the indicated splenic γδ T cell

subsets comparing WT and GFI1 KO mice. **E**: c-MAF MFI quantification comparing WT and

GFI1 KO mice for the indicated thymic DN progenitors. **F**: Peaks for GFI1 on the *Maf* promoter

region from the MLL-ENL ChIP-Seq (GSE31657). **G**: ChIP-PCR on different potential binding

sites (arrowheads) on *Maf* promoter for GFI1 identified in Fig. 8F with WT and GFI1 KO DN

cells. *Gfi1* promoter (*Gfi1-P*) was used as positive control and *Il6 promoter* region was used as

negative control. Statistical testing was performed with multiple paired *t*-test using Prism v9

(GraphPad). *p*-Values are shown as * < 0.05, ** < 0.01, *** < 0.005, and **** < 0.001 where

each statistical significance was found.

Supplementary data
856
Sup. Fig. 1: Phenotype of γδ T cells in adult GFI1 KO mice

A: Violin plots showing the number of reads and genes as well as the percentage of
mitochondrial genes quantified per cell in the lung tissue in the scRNA-seq data. **B:** Dot plot
showing the key differentially expressed genes in each cluster shown in Fig. 1A. Color represents
the scaled mean expression of the gene in the respective cluster, and dot size represents the
fraction of cells in the cluster expressing the gene. **C**: Representative plots on small intestine (SI)
and colon lamina propria cells showing γδ T cells and CD4 T cells after gating on live (according
to FSC/SSC) singlets CD45$^+$CD3$^+$ cells. SI and colon lamina propria cells were isolated
following published protocols[81]. **D & E**: CD4$^+$ T cell counts in the spleen, LN, and lung of WT
and GFI1 KO mice (**D**) or control and *vav-cre GFI1$^{flox/flox}$* mice (**E**). Adult mice meaning mice
older than 5 weeks. Statistical testing was performed with paired *t*-test using Prism v9
(GraphPad). *p*-Values are shown as * < 0.05, ** < 0.01, *** < 0.005, and **** < 0.001 where
each statistical significance was found.

Sup. Fig. 2: Expansion of the RORγt$^+$IL17a$^+$ γδT subset in adult GFI1 KO mice.

A: Representative plots showing the expression of CD27 on gated live singlets CD45$^+$γδT$^+$CD4$^-$
cells from WT and GFI1 KO spleen, LN and lung. **B**: Representative normalized histograms of
RORγt$^+$ cells in either singlets CD45$^+$CD4$^-$γδT$^+$ or CD45$^+$CD4$^+$γδT$^-$ splenic, LN or lung cells
from control and *Vav-cre Gfi1$^{flox/flox}$* mice. **C**: Frequencies of RORγt$^+$ cells within singlets
CD45$^+$CD4$^-$ γδT$^+$ cells from splenic, LN and lung compared to control and *vav-cre GFI1$^{flox/flox}$*
mice. **D**: splenic, LN and lung CD45$^+$CD4$^-$γδT$^+$RORγt$^+$ cell counts in control and *vav-cre*
GFI1$^{flox/flox}$ mice. **E**: Representative plots showing the expression of IL22 and IL17A in gated
singlets CD45$^+$CD4$^-$γδT$^+$ cells in the spleen, LN and lung from WT and GFI1 KO mice. Cells
were stimulated with IL23 (20ng/mL) overnight and treated with PMA, ionomycin and Golgi
Plug for the last 4 hours before staining. **F**: Percentages of IL17A$^+$IL22$^+$ cells in γδ T cells from
the spleen, LN and lung of WT and GFI1 KO mice. **G**: IL17A$^+$IL22$^+$γδT$^+$ cell counts in the
spleen, LN and lung of WT and GFI1 KO mice. Adult mice meaning mice older than 5 weeks.
Statistical testing was performed with paired *t*-test using Prism v9 (GraphPad). *p*-Values are
shown as * < 0.05, ** < 0.01, *** < 0.005 and **** < 0.001 where each statistical significance
was found.

Sup. Fig. 3: Expansion of Vγ6$^+$ γδT17 cells in the periphery of adult GFI1 KO mice.

A: Violin plots showing the number of reads and genes as well as the percentage of mitochondrial genes quantified per cell in the splenic γδ T cells in the scRNA-seq data. **B**: Scheme showing *in vitro* culture of sorted splenic Vγ1$^-$Vγ4$^-$Vγ5$^-$ γδ T cells from WT and GFI1 KO mice in the presence of IL1β (10 ng/mL) and IL23 (20ng/mL) for 16 hours and the frequencies of IL17A$^+$ cells in the cultured cells after adding Golgi Plug for the last 4 hours. **C**: Representative normalized histogram of PhosphoSTAT3 in gated WT and GFI1 KO Vγ1$^-$Vγ4$^-$ Vγ5$^-$ γδ T stimulated or not with IL7 (10 ng/mL) or IL23 (20 ng/mL) for 15 min and the measured mean fluorescence (MFI) for each condition. Adult mice meaning mice older than 5 weeks. Statistical testing was performed with multiple paired *t*- test **G** using Prism v9 (GraphPad). *p*-Values are shown as * < 0.05, ** < 0.01, *** < 0.005, and **** < 0.001 where each statistical significance was found.

Sup. Fig. 4: Expansion of Vγ6$^+$ γδT17 cells in the thymus of adult GFI1 KO mice.

A: Cell numbers (left) and frequencies in CD45$^+$ cells (right) of γδ T cells (CD4$^-$γδT$^+$) in the thymus of control and *vav-cre GFI1$^{flox/flox}$* mice. **B**: Representative normalized histogram of RORγt$^+$ cells in either CD45$^+$CD4$^-$γδT$^+$ or CD45$^+$CD4$^+$γδT$^-$ total thymocytes from control and *vav-cre, Gfi1*$^{flox/flox}$ mice. **C**: Frequencies of RORγt$^+$ cells in thymic CD45$^+$CD4$^-$γδT$^+$ comparing control and *vav-cre, Gfi1*$^{flox/flox}$ mice (left) and thymic CD45$^+$CD4$^-$γδT$^+$RORγt$^+$ cell counts in control and *vav-cre, Gfi1*$^{flox/flox}$ mice (right). **D:** Violin plots showing the number of reads and genes as well as the percentage of mitochondrial genes quantified per cell in the thymic γδ T cells in the scRNA-seq data. **E**: Differential gene expression showing key up- or down- regulated genes in GFI1 KO thymic γδ T cells versus WT splenic γδ T cells. **F**: Representative plots showing CD44 and CD45RB expression in gated thymic CD45$^+$CD4$^-$γδT$^+$CD24$^-$ cells from WT and GFI1KO mice. **G**: Cell counts of thymic CD45$^+$CD4$^-$γδT$^-$CD24$^-$CD44$^+$CD45RB$^-$ (γδT17), CD45$^+$CD4$^-$γδT$^-$CD24$^-$CD44$^+$CD45RB$^+$ and CD45$^+$CD4$^-$γδT$^-$CD24$^-$CD44$^-$CD45RB$^+$ cells in WT and GFI1 KO mice. **H**: Representative plots of CD24 expression in γδ T cells and Vγ chains expression in gated CD24$^-$ γδ T cells from D8-10 and D18-D20 old WT and GFI1 KO mice. **I**: Cell numbers of thymic CD24$^-$ γδT$^+$ cells in D8-10 and D18-D20 old WT and GFI1 KO mice. **J & K**: Vγ chain frequencies (**J**) and Vγ chain cell counts (**K**) in thymic CD24$^-$ γδ T cells from

D8-10 and D18-D20 old WT and GFI1 KO mice. $V\gamma^- = V\gamma1^-V\gamma4^-V\gamma5^-$. Statistical testing was

performed with multiple paired *t*-test for **G**, multiple unpaired *t*-test for **I**, **J** and **K** or paired *t*-test

for **A** and **C**, using Prism v9 (GraphPad). *p*-Values are shown as * < 0.05, ** < 0.01, *** <

0.005, and **** < 0.001 where each statistical significance was found.

Sup. Fig. 5: GFI1 intrinsically controls the generation of γδ T cells.

A: *Gfi1b* mRNA expression in γδ T cells from thymus, spleen, LN, and lung. *Gapdh* was used as

the housekeeping gene. **B**: Representative plots showing CD45.1/2$^+$ donor γδ T cells and

CD45.2$^+$ recipient γδ T cells in the spleen, thymus and lung of non-irradiated WT and KO

recipients after gating on live, singlets cells. **C and D**: Representative plots showing IL17A$^+$ (C)

and RORγt$^+$ (D) cells previously gated on CD45.1/2$^+$ donor γδ T cells or CD45.2$^+$ recipient (WT

and KO) γδ T cells from the spleen, thymus and lung of non-irradiated WT and KO recipients

after gating on live, singlets cells. **E**: Representative plots showing Vγ1 and Vγ4 expression in

CD45.1/2$^+$ donor γδ T cells or in CD45.2$^+$ recipient (WT and KO) γδ T cells from the spleen,

thymus and lung of non-irradiated WT and KO recipients after gating on live, singlets cells.

Sup. Fig. 6: Precursors in thymus of adult mice

A: Scheme showing the culture of sorted thymic DN cells from WT and KO mice on an OP9-

DL1 layer in the presence of Flt3 (5 ng/mL) and IL7 (1 ng/mL) for 12-14 days. **B**: Representative

plots showing the purity of the thymic DN population before selection (total thymocytes) and

after selection. **C**: Representative plots showing the purity of the thymic DN population after

selection from WT and GFI1 KO thymi. **D**: DN cell frequencies in the thymus of WT and GFI1

KO mice or control and *vav-cre GFI1*$^{flox/flox}$ mice. DN1a/b cells were characterized as CD4$^-$CD8$^-$

γδT$^-$CD44$^+$CD25$^-$CD24$^+$Ckit$^+$, DN1d cells were characterized as CD4$^-$CD8$^-$γδT$^-$CD44$^+$CD25$^-$

CD24$^+$Ckit$^-$, DN1e cells were characterized as CD4$^-$CD8$^-$γδT$^-$CD44$^+$CD25$^-$CD24$^{-/low}$Ckit$^-$, DN2

cells were characterized as CD4$^-$CD8$^-$γδT$^-$CD44$^+$CD25$^+$ and DN3 cells were characterized as

CD4$^-$CD8$^-$γδT$^-$CD44$^-$CD25$^+$. **E**: Dot plot showing the key differentially expressed genes in each

cluster shown in Fig. 6C. Color represents the mean expression of the gene in the respective

cluster, and dot size represents the fraction of cells in the cluster expressing the gene. **F**: Violin

plots showing the number of reads and genes as well as the percentage of mitochondrial genes

quantified per cell in the thymic DN cells in the scRNA-seq data. **G**: GFP$^+$ cells indicating Gfi1

expression in the indicated thymic DN populations from GFI1:GFP reporter mice (het) and

controls (wt). **H**: c-MAF % in DN1a/b, DN1d, DN1e, DN2 and DN3 cells for WT and GFI1 KO

mice or control and *vav-cre GFI1^{flox/flox}* mice analyzed by flow cytometry. Statistical testing was

performed with multiple paired *t*- test **G** using Prism v9 (GraphPad). *p*-Values are shown as * <

0.05, ** < 0.01, *** < 0.005 and **** < 0.001 where each statistical significance was found.

Sup. Fig. 7: Potential of differentiation of precursors in thymus in adult mice

A: Representative plots of splenocytes and thymocytes from control and *Cd2-cre* Gfi1^{flox/flox} mice

analyzed by flow cytometry and stained to identify γδ T cells (CD45^+CD4^-γδT^+) after gating on

live cells (according to the FSC/SSC) and singlets. **B**: Cell numbers and frequencies in CD45^+

cells of γδ T cells (CD4^-γδT^+) in the spleen and in the thymus of control and *Cd2-cre* Gfi1^{flox/flox}

mice. **C & D**: Cell counts of Vγ1^+ γδ T cells, Vγ4^+ γδ T cells and Vγ1^-Vγ4^- γδ T cells in the

spleen (**C**) and in the thymus (**D**) of control and *Cd2-cre* Gfi1^{flox/flox} mice. **E:** Representative

plots of splenic and thymic cells from control and *Cd2-cre* Gfi1^{flox/flox} mice treated with PMA,

ionomycin and Golgi Plug for 3.5 hours and analysed by flow cytometry. The cells were gated on

singlets CD45^+CD4^-γδT^+ before gating for IFNγ and IL17A. **F:** Representative normalized

histogram of RORγt^+ cells in either splenic or thymic CD45^+CD4^-γδT^+ and CD45^+CD4^+γδT^- from

control and *Cd2-cre* Gfi1^{flox/flox} mice. **G**: Frequencies of IL17A^+ cells in γδT^+ cells and

CD45^+CD4^-γδT^+IL17A^+ cell counts in the spleen and thymus of control and *Cd2-cre* Gfi1^{flox/flox}

mice. **H**: Frequencies of RORγt^+ cells in γδT^+ cells and CD45^+CD4^-γδT^+RORγt^+ cell counts in

the spleen and thymus of control and *Cd2-cre* Gfi1^{flox/flox} mice. **I**: Scheme showing the culture of

sorted thymic DN1 or DN2/3 cells from WT and KO mice on an OP9-DL4 layer in the presence

of Flt3 (5 ng/mL) and IL7 (1 ng/mL) for maximum 20 days. **J**: Representative plots showing the

gating strategy to sort DN1 and DN2/DN3 for the coculture with the OP9-DL4 cells described in

Suppl. Fig. 7I. Statistical testing was performed with paired *t*- test **G** using Prism v9 (GraphPad).

p-Values are shown as * < 0.05, ** < 0.01, *** < 0.005 and **** < 0.001 where each statistical

significance was found

Sup. Fig. 8: GFI1 regulation in adult mice

A: Differential gene expression showing key up- or down- regulated genes in GFI1 KO DN3a,

DN3a→b and DN3b cells versus WT cells. **B**: Peak for GFI1 on *Maf* promoter from ChIP-seq

analysis from Mast cells (GSE42518), ILC2 cells (GSE50806) and Th1 cells (GSE70893).

REFERENCES

1. Moroy, T. & Khandanpour, C. Growth factor independence 1 (Gfi1) as a regulator of lymphocyte development and activation. *Semin Immunol* **23**, 368-378 (2011).
2. Fraszczak, J. & Moroy, T. The transcription factors GFI1 and GFI1B as modulators of the innate and acquired immune response. *Adv Immunol* **149**, 35-94 (2021).
3. Phelan, J.D., Shroyer, N.F., Cook, T., Gebelein, B. & Grimes, H.L. Gfi1-cells and circuits: unraveling transcriptional networks of development and disease. *Curr Opin Hematol* **17**, 300-307 (2010).
4. van der Meer, L.T., Jansen, J.H. & van der Reijden, B.A. Gfi1 and Gfi1b: key regulators of hematopoiesis. *Leukemia* **24**, 1834-1843 (2010).
5. Saleque, S., Kim, J., Rooke, H.M. & Orkin, S.H. Epigenetic regulation of hematopoietic differentiation by Gfi-1 and Gfi-1b is mediated by the cofactors CoREST and LSD1. *Mol Cell* **27**, 562-572 (2007).
6. Thambyrajah, R.*, et al.* GFI1 proteins orchestrate the emergence of haematopoietic stem cells through recruitment of LSD1. *Nat Cell Biol* **18**, 21-32 (2016).
7. Zweidler-Mckay, P.A., Grimes, H.L., Flubacher, M.M. & Tsichlis, P.N. Gfi-1 encodes a nuclear zinc finger protein that binds DNA and functions as a transcriptional repressor. *Mol Cell Biol* **16**, 4024-4034 (1996).
8. Grimes, H.L., Chan, T.O., Zweidler-McKay, P.A., Tong, B. & Tsichlis, P.N. The Gfi-1 proto-oncoprotein contains a novel transcriptional repressor domain, SNAG, and inhibits G1 arrest induced by interleukin-2 withdrawal. *Mol Cell Biol* **16**, 6263-6272 (1996).
9. Helness, A.*, et al.* GFI1 tethers the NuRD complex to open and transcriptionally active chromatin in myeloid progenitors. *Commun Biol* **4**, 1356 (2021).
10. Wang, Y.*, et al.* LSD1 is a subunit of the NuRD complex and targets the metastasis programs in breast cancer. *Cell* **138**, 660-672 (2009).
11. Schmidt, T., Zornig, M., Beneke, R. & Moroy, T. MoMuLV proviral integrations identified by Sup-F selection in tumors from infected myc/pim bitransgenic mice correlate with activation of the gfi-1 gene. *Nucleic Acids Res* **24**, 2528-2534 (1996).
12. Zornig, M., Schmidt, T., Karsunky, H., Grzeschiczek, A. & Moroy, T. Zinc finger protein GFI-1 cooperates with myc and pim-1 in T-cell lymphomagenesis by reducing the requirements for IL-2. *Oncogene* **12**, 1789-1801 (1996).
13. Karsunky, H.*, et al.* Inflammatory reactions and severe neutropenia in mice lacking the transcriptional repressor Gfi1. *Nat Genet* **30**, 295-300 (2002).
14. Zeng, H., Yucel, R., Kosan, C., Klein-Hitpass, L. & Moroy, T. Transcription factor Gfi1 regulates self-renewal and engraftment of hematopoietic stem cells. *EMBO J* **23**, 4116-4125 (2004).
15. Fraszczak, J.*, et al.* Threshold Levels of Gfi1 Maintain E2A Activity for B Cell Commitment via Repression of Id1. *PLoS One* **11**, e0160344 (2016).
16. Fraszczak, J. & Moroy, T. The role of the transcriptional repressor growth factor independent 1 in the formation of myeloid cells. *Curr Opin Hematol* **24**, 32-37 (2017).
17. Yucel, R., Karsunky, H., Klein-Hitpass, L. & Moroy, T. The transcriptional repressor Gfi1 affects development of early, uncommitted c-Kit+ T cell progenitors and CD4/CD8 lineage decision in the thymus. *J Exp Med* **197**, 831-844 (2003).
18. Phelan, J.D.*, et al.* Growth factor independent-1 maintains Notch1-dependent transcriptional programming of lymphoid precursors. *PLoS Genet* **9**, e1003713 (2013).

19. Shi, L.Z.*, et al.* Gfi1-Foxo1 axis controls the fidelity of effector gene expression and
developmental maturation of thymocytes. *Proc Natl Acad Sci U S A* **114**, E67-E74 (2017).
20. Shi, L.Z.*, et al.* Inhibitory role of the transcription repressor Gfi1 in the generation of
thymus-derived regulatory T cells. *Proc Natl Acad Sci U S A* **110**, E3198-3205 (2013).
21. Zhu, J.*, et al.* Down-regulation of Gfi-1 expression by TGF-beta is important for
differentiation of Th17 and CD103+ inducible regulatory T cells. *J Exp Med* **206**, 329-
341 (2009).
22. Kurebayashi, Y.*, et al.* PI3K-Akt-mTORC1-S6K1/2 axis controls Th17 differentiation by
regulating Gfi1 expression and nuclear translocation of RORgamma. *Cell Rep* **1**, 360-373
(2012).
23. Ichiyama, K.*, et al.* Gfi1 negatively regulates T(h)17 differentiation by inhibiting
RORgammat activity. *Int Immunol* **21**, 881-889 (2009).
24. Spooner, C.J., Cheng, J.X., Pujadas, E., Laslo, P. & Singh, H. A recurrent network
involving the transcription factors PU.1 and Gfi1 orchestrates innate and adaptive
immune cell fates. *Immunity* **31**, 576-586 (2009).
25. Spooner, C.J.*, et al.* Specification of type 2 innate lymphocytes by the transcriptional
determinant Gfi1. *Nat Immunol* **14**, 1229-1236 (2013).
26. Zhu, J.*, et al.* Growth factor independent-1 induced by IL-4 regulates Th2 cell
proliferation. *Immunity* **16**, 733-744 (2002).
27. Zhu, J., Yamane, H., Cote-Sierra, J., Guo, L. & Paul, W.E. GATA-3 promotes Th2
responses through three different mechanisms: induction of Th2 cytokine production,
selective growth of Th2 cells and inhibition of Th1 cell-specific factors. *Cell Res* **16**, 3-10
(2006).
28. Zhu, J., Jankovic, D., Grinberg, A., Guo, L. & Paul, W.E. Gfi-1 plays an important role in
IL-2-mediated Th2 cell expansion. *Proc Natl Acad Sci U S A* **103**, 18214-18219 (2006).
29. Suzuki, J.*, et al.* Gfi1, a transcriptional repressor, inhibits the induction of the T helper
type 1 programme in activated CD4 T cells. *Immunology* **147**, 476-487 (2016).
30. Friesen, L.*, et al.* Cutting Edge: The Expression of Transcription Inhibitor GFI1 Is
Induced by Retinoic Acid to Rein in Th9 Polarization. *J Immunol* **209**, 1237-1242 (2022).
31. Ojo, O.A.*, et al.* Gfi1 controls the formation of effector-like CD8(+) T cells during
chronic infection and cancer. *Nat Commun* **16**, 4542 (2025).
32. Chaudhry, M.Z.*, et al.* GFI1-driven transcriptional and epigenetic programs maintain
CD8(+) T cell stemness and persistence. *Nat Immunol* **26**, 894-907 (2025).
33. Sumaria, N., Martin, S. & Pennington, D.J. Developmental origins of murine gammadelta
T-cell subsets. *Immunology* **156**, 299-304 (2019).
34. Ribot, J.C., Lopes, N. & Silva-Santos, B. gammadelta T cells in tissue physiology and
surveillance. *Nat Rev Immunol* **21**, 221-232 (2021).
35. Hu, Y.*, et al.* gammadelta T cells: origin and fate, subsets, diseases and immunotherapy.
Signal Transduct Target Ther **8**, 434 (2023).
36. Sumaria, N.*, et al.* Perinatal thymic-derived CD8alphabeta-expressing gammadelta T cells
are innate IFN-gamma producers that expand in IL-7R-STAT5B-driven neoplasms. *Nat
Immunol* **25**, 1207-1217 (2024).
37. Buus, T.B., Odum, N., Geisler, C. & Lauritsen, J.P.H. Three distinct developmental
pathways for adaptive and two IFN-gamma-producing gammadelta T subsets in adult
thymus. *Nat Commun* **8**, 1911 (2017).
38. Serre, K. & Silva-Santos, B. Molecular Mechanisms of Differentiation of Murine Pro-
Inflammatory gammadelta T Cell Subsets. *Front Immunol* **4**, 431 (2013).

39. Bhandoola, A., von Boehmer, H., Petrie, H.T. & Zuniga-Pflucker, J.C. Commitment and developmental potential of extrathymic and intrathymic T cell precursors: plenty to choose from. *Immunity* **26**, 678-689 (2007).

40. Oh, S*., et al.* Distinct subpopulations of DN1 thymocytes exhibit preferential gammadelta T lineage potential. *Front Immunol* **14**, 1106652 (2023).

41. Spidale, N.A*., et al.* Interleukin-17-Producing gammadelta T Cells Originate from SOX13(+) Progenitors that Are Independent of gammadeltaTCR Signaling. *Immunity* **49**, 857-872 e855 (2018).

42. Munoz-Ruiz, M*., et al.* TCR signal strength controls thymic differentiation of discrete proinflammatory gammadelta T cell subsets. *Nat Immunol* **17**, 721-727 (2016).

43. Sumaria, N., Grandjean, C.L., Silva-Santos, B. & Pennington, D.J. Strong TCRgammadelta Signaling Prohibits Thymic Development of IL-17A-Secreting gammadelta T Cells. *Cell Rep* **19**, 2469-2476 (2017).

44. Zarin, P., Chen, E.L., In, T.S., Anderson, M.K. & Zuniga-Pflucker, J.C. Gamma delta T-cell differentiation and effector function programming, TCR signal strength, when and how much? *Cell Immunol* **296**, 70-75 (2015).

45. Anderson, M.K. & Selvaratnam, J.S. Interaction between gammadeltaTCR signaling and the E protein-Id axis in gammadelta T cell development. *Immunol Rev* **298**, 181-197 (2020).

46. Lopez, R.D. Human gammadelta-T cells in adoptive immunotherapy of malignant and infectious diseases. *Immunol Res* **26**, 207-221 (2002).

47. Kang, I., Kim, Y. & Lee, H.K. Double-edged sword: gammadelta T cells in mucosal homeostasis and disease. *Exp Mol Med* **55**, 1895-1904 (2023).

48. Gray, E.E., Suzuki, K. & Cyster, J.G. Cutting edge: Identification of a motile IL-17-producing gammadelta T cell population in the dermis. *J Immunol* **186**, 6091-6095 (2011).

49. Ramirez-Valle, F., Gray, E.E. & Cyster, J.G. Inflammation induces dermal Vgamma4+ gammadeltaT17 memory-like cells that travel to distant skin and accelerate secondary IL-17-driven responses. *Proc Natl Acad Sci U S A* **112**, 8046-8051 (2015).

50. Spidale, N.A*., et al.* Neonatal-derived IL-17 producing dermal gammadelta T cells are required to prevent spontaneous atopic dermatitis. *Elife* **9**(2020).

51. Frascoli, M*., et al.* Skin gammadelta T cell inflammatory responses are hardwired in the thymus by oxysterol sensing via GPR183 and calibrated by dietary cholesterol. *Immunity* **56**, 562-575 e566 (2023).

52. Revesz, I.A., Joyce, P., Ebert, L.M. & Prestidge, C.A. Effective gammadelta T-cell clinical therapies: current limitations and future perspectives for cancer immunotherapy. *Clin Transl Immunology* **13**, e1492 (2024).

53. Pargmann, D*., et al.* Differential impact of the transcriptional repressor Gfi1 on mature CD4+ and CD8+ T lymphocyte function. *Eur J Immunol* **37**, 3551-3563 (2007).

54. Ribot, J.C*., et al.* CD27 is a thymic determinant of the balance between interferon-gamma- and interleukin 17-producing gammadelta T cell subsets. *Nat Immunol* **10**, 427-436 (2009).

55. Liang, S.C*., et al.* Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J Exp Med* **203**, 2271-2279 (2006).

56. Fiala, G.J., Gomes, A.Q. & Silva-Santos, B. From thymus to periphery: Molecular basis of effector gammadelta-T cell differentiation. *Immunol Rev* **298**, 47-60 (2020).

57. Haas, J.D*., et al.* Development of interleukin-17-producing gammadelta T cells is restricted to a functional embryonic wave. *Immunity* **37**, 48-59 (2012).

58. Shibata, K*., et al.* Identification of CD25+ gamma delta T cells as fetal thymus-derived naturally occurring IL-17 producers. *J Immunol* **181**, 5940-5947 (2008).

59. du Halgouet, A*., et al.* Multimodal profiling reveals site-specific adaptation and tissue residency hallmarks of gammadelta T cells across organs in mice. *Nat Immunol* **25**, 343-356 (2024).

60. Tan, L*., et al.* Single-Cell Transcriptomics Identifies the Adaptation of Scart1(+) Vgamma6(+) T Cells to Skin Residency as Activated Effector Cells. *Cell Rep* **27**, 3657-3671 e3654 (2019).

61. Parker, M.E. & Ciofani, M. Regulation of gammadelta T Cell Effector Diversification in the Thymus. *Front Immunol* **11**, 42 (2020).

62. Sagar*, et al.* Deciphering the regulatory landscape of fetal and adult gammadelta T-cell development at single-cell resolution. *EMBO J* **39**, e104159 (2020).

63. Yucel, R., Kosan, C., Heyd, F. & Moroy, T. Gfi1:green fluorescent protein knock-in mutant reveals differential expression and autoregulation of the growth factor independence 1 (Gfi1) gene during lymphocyte development. *J Biol Chem* **279**, 40906-40917 (2004).

64. Mercier, F.E., Sykes, D.B. & Scadden, D.T. Single Targeted Exon Mutation Creates a True Congenic Mouse for Competitive Hematopoietic Stem Cell Transplantation: The C57BL/6-CD45.1(STEM) Mouse. *Stem Cell Reports* **6**, 985-992 (2016).

65. Gray, E.E*., et al.* Deficiency in IL-17-committed Vgamma4(+) gammadelta T cells in a spontaneous Sox13-mutant CD45.1(+) congenic mouse substrain provides protection from dermatitis. *Nat Immunol* **14**, 584-592 (2013).

66. Porritt, H.E*., et al.* Heterogeneity among DN1 prothymocytes reveals multiple progenitors with different capacities to generate T cell and non-T cell lineages. *Immunity* **20**, 735-745 (2004).

67. Wolf, F.A*., et al.* PAGA: graph abstraction reconciles clustering with trajectory inference through a topology preserving map of single cells. *Genome Biol* **20**, 59 (2019).

68. Zuberbuehler, M.K*., et al.* The transcription factor c-Maf is essential for the commitment of IL-17-producing gammadelta T cells. *Nat Immunol* **20**, 73-85 (2019).

69. Yang, T*., et al.* RORgammat(+) c-Maf(+) Vgamma4(+) gammadelta T cells are generated in the adult thymus but do not reach the periphery. *Cell Rep* **42**, 113230 (2023).

70. Tanaka, S*., et al.* Sox5 and c-Maf cooperatively induce Th17 cell differentiation via RORgammat induction as downstream targets of Stat3. *J Exp Med* **211**, 1857-1874 (2014).

71. In, T.S.H*., et al.* HEB is required for the specification of fetal IL-17-producing gammadelta T cells. *Nat Commun* **8**, 2004 (2017).

72. Fahl, S.P*., et al.* The E protein-TCF1 axis controls gammadelta T cell development and effector fate. *Cell Rep* **34**, 108716 (2021).

73. Chen, X*., et al.* Differential metabolic requirement governed by transcription factor c-Maf dictates innate gammadeltaT17 effector functionality in mice and humans. *Sci Adv* **8**, eabm9120 (2022).

74. Bialy, S. & Bogunia-Kubik, K. Uncovering the mysteries of human gamma delta T cells: from origins to novel therapeutics. *Front Immunol* **16**, 1543454 (2025).

75. Maiques-Diaz, A. & Somervaille, T.C. LSD1: biologic roles and therapeutic targeting. *Epigenomics* **8**, 1103-1116 (2016).

1171   76.   Schmitt, T.M. & Zuniga-Pflucker, J.C. Induction of T cell development from
1172   hematopoietic progenitor cells by delta-like-1 in vitro. *Immunity* **17**, 749-756 (2002).
1173   77.   Rizk, J.*, et al.* The cIAP ubiquitin ligases sustain type 3 gammadelta T cells and ILC
1174   during aging to promote barrier immunity. *J Exp Med* **220**(2023).
1175   78.   de Araujo, P.S., Rosseneu, M.Y., Kremer, J.M., van Zoelen, E.J. & de Haas, G.H.
1176   Structure and thermodynamic properties of the complexes between phospholipase A2 and
1177   lipid micelles. *Biochemistry* **18**, 580-586 (1979).
1178   79.   Bernstein, N.J.*, et al.* Solo: Doublet Identification in Single-Cell RNA-Seq via Semi-
1179   Supervised Deep Learning. *Cell Syst* **11**, 95-101 e105 (2020).
1180   80.   Wolf, F.A., Angerer, P. & Theis, F.J. SCANPY: large-scale single-cell gene expression
1181   data analysis. *Genome Biol* **19**, 15 (2018).
1182   81.   Meli, A.P.*, et al.* Bcl-6 expression by CD4(+) T cells determines concomitant immunity
1183   and host resistance across distinct parasitic infections. *Mucosal Immunol* **16**, 801-816
1184   (2023).
1185

Figure 1: Mice lacking GFI1 show an expansion of γδ T cells

Figure 2: subtype of γδT cells

Figure 4: Thymus Vγ6

Figure 5: GFI1 intrinsically controls the generation of γδT cells

Figure 6: DN

Figure 7: Precursors

A Clusters

B Cell types

- DN1b-c/DN2
- DN3a
- DN3a → DN3b
- DN3b
- Il2rb⁺DN1e
- Rorc⁺DN1d
- Rorc⁺DN1e
- Trgv6⁺

C

D DN2/3 WT D8

E DN2/3 KO D8

F ●WT ●KO

DP cells

γδT cells

G ●WT ●KO

CD24⁻ γδT cells

CD24⁻ γδT cells

H DN1 WT D8

I DN1 KO D8

J DN1 WT D18

K DN1 KO D18

L ●WT ●KO   γδT cells

M CD24⁻ γδT cells   ●WT ●KO

CD24⁻ γδT cells

Figure 8: GFI1 regulation