Title

Identification of molecular determinants governing KIF4 cargo recognition of NTCP and their role in HBV/HDV infection

Authors

Sameh A. Gad[1,2*], Doaa A. Abo Elwafa[1,3,4], Masaaki Toyama[1], Mitsunori Ikeguchi[5], Ikuo Shoji[6], Kazuaki Chayama[7,8], Masanori Isogawa[1], Takaji Wakita[1], Hussein H. Aly[1#]

Affiliation

[1]Department of Virology II, National Institute of Infectious Diseases; Japan Institute for Health Security; 1-23-1 Toyama; Shinjuku-ku: Tokyo 162-8640; Japan

[2]Department of Microbiology and Immunology; Faculty of Pharmacy; Minia University; Minia 61519; Egypt

[3]Graduate School of Medical and Pharmaceutical Sciences, Chiba University, Japan

[4]Department of Medical Microbiology and Immunology, Faculty of Medicine, South Valley University, Egypt

[5]Computational Life Science Laboratory, Graduate School of Medical Life Science, Yokohama City University, Yokohama, Japan

[6]Division of Infectious Disease Control; Center for Infectious Diseases; Kobe University Graduate School of Medicine; 7-5-1 Kusunoki-cho; Chuo-ku Kobe 650-0017; Japan

[7]Hiroshima Institute of Life Sciences; Hiroshima; Japan

[8]RIKEN Center for Integrative Medical Sciences; Yokohama; Japan

Running title

KIF4 Recognition of NTCP regulating HBV/HDV entry.

[#]**Address correspondence to:** Hussein H. Aly, hussein.a@jihs.go.jp

[*] **Present address:** Department of Virology, Graduate School of Medical and Dental Sciences, Niigata University, Niigata, Japan. (Recent affiliation)

Abstract word count: 232, Text word count: 4514

Abstract

The surface availability of the sodium taurocholate co-transporting polypeptide (NTCP), which serves as entry receptor for Hepatitis B virus (HBV) and Hepatitis D virus (HDV), is a key determinant of HBV/HDV infectivity. We previously identified the Kinesin motor protein KIF4 as a key regulator of NTCP trafficking to the plasma membrane. However, the molecular mechanism by which NTCP is recognized as a KIF4 transport cargo has remained undefined. Here, we identified a leucine-rich motif, LLLI (aa 298–301), within the C-terminal transmembrane domain of NTCP as an essential determinant for KIF4 transport. Co-immunoprecipitation analyses revealed its importance for NTCP interaction with KIF4. This motif is conserved in another established KIF4 cargo, integrin β1 (ITGB1), and mutation of the corresponding residues similarly disrupted ITGB1–KIF4 interaction, indicating a shared cargo recognition mechanism employed by KIF4. Functionally, disruption of the NTCP leucine-rich motif markedly impaired NTCP surface localization without affecting total protein expression, resulting in severe attenuation of HBV and HDV infection. Structural modeling and computational protein–protein interaction analysis indicated that the mutation does not alter NTCP structure or NTCP/preS1 binding affinity, supporting a trafficking-specific defect. Substitution mutations identified L298 and L299 as the minimal residues required for interaction with KIF4. Our findings define a recognition motif that regulates KIF4-dependent cargo identification and surface trafficking. It also underscores the potential of targeting KIF4-mediated NTCP recognition and transport to inhibit HBV and HDV entry.

Importance

Hepatitis B and D viruses require the hepatocyte surface receptor NTCP for entry. We identified a conserved leucine-rich motif (LLLI, aa 298–301) in NTCP that is necessary for its recognition by the kinesin motor KIF4; disrupting this motif selectively impairs NTCP surface trafficking and markedly attenuates HBV/HDV infection without reducing total NTCP or intrinsic receptor function. Importantly, we identified a similar motif in integrin β1 (ITGB1; aa 747–750) that is required for interaction with KIF4, revealing a similar cargo selection mechanism for different proteins. Scientifically, these findings define a motif-derived cargo-selection determinant implied by KIF4, advancing a general principle for kinesin-dependent cargo recognition. More broadly, this work highlights a possible host-directed antiviral approach that targets cargo-selection interfaces and provides a conceptual framework for manipulating kinesin-mediated trafficking of clinically relevant membrane proteins beyond NTCP, with implications for receptor regulation and membrane protein therapeutics.

Introduction

Chronic hepatitis B virus (HBV) infection remains a major global health burden, affecting nearly 1.5 million new infections/year with 254 million individuals suffering from chronic HBV worldwide. Chronic HBV patients may progress to complications from the progressive liver disease such as cirrhosis, and hepatocellular carcinoma (1). Clinical outcomes of HBV patients worsen substantially in the presence of hepatitis D virus (HDV), a defective satellite virus that requires HBV envelope proteins for propagation. HDV coinfection accelerates liver inflammation, increases the risk of fulminant hepatitis, and markedly elevates the likelihood of end-stage liver disease (2). A pivotal step in the life cycle of both HBV and HDV is their shared dependence on the sodium taurocholate co-transporting polypeptide (NTCP), a bile acid transporter that functions as the essential entry receptor on hepatocytes (3). Understanding how NTCP is trafficked and presented on the cell surface is therefore central to elucidating the mechanisms governing HBV/HDV infection (4).

Kinesin family member 4 (KIF4) is a microtubule-based motor protein involved in intracellular transport, chromosome segregation, and cytoskeletal organization (5). Beyond its canonical roles in mitosis and nuclear dynamics, emerging evidence indicates that KIF4 also functions in cytoplasmic vesicular trafficking and regulates the transport and surface localization of specific membrane proteins (6-8). Our previous findings revealed that KIF4 interacts with NTCP and facilitates its anterograde transport to the plasma membrane, thereby enabling the receptor's availability for HBV/HDV attachment and viral entry (4). These observations positioned KIF4 as a previously unrecognized host factor that

modulates viral susceptibility by controlling NTCP surface expression (4, 9).

Although kinesins share highly conserved motor domains, cargo specificity is primarily

determined by divergent C□terminal tail domain that engage adaptor proteins, recognize

discrete cargo□binding motifs, and integrate regulatory inputs such as post□translational

modifications and scaffolding complex assembly (10-13). For many kinesins, such as

KIF1A, KIF5, and KIF3 family members, structural studies have identified short linear

motifs or coiled-coil regions that mediate selective cargo recognition (11, 12, 14, 15).

However, the molecular basis by which KIF4 identifies and transports its physiological

cargoes remains poorly defined. Although, we clearly demonstrated the interaction between

KIF4 and NTCP by pull-down assays and intracellular localization studies, recombinant

KIF4 and NTCP failed to exhibit a direct physical association in vitro. This observation

suggests that the KIF4–NTCP interaction is likely facilitated by specific adaptor proteins

that mediate or regulate this association within the cellular environment (4). Only limited

information exists regarding KIF4's tail-domain interactome, and no detailed determinants

have been identified for its cargo selection of membrane proteins such as NTCP. This gap

in knowledge reduces our ability to understand how KIF4-dependent cargo trafficking is

regulated and how it may be selectively modulated.

Defining the molecular determinants that govern KIF4–cargo recognition, particularly in

the context of NTCP, holds substantial therapeutic relevance. Because NTCP surface

availability is a rate-limiting step for HBV/HDV entry (3, 4), elucidating how KIF4

recognizes and transports NTCP could reveal new intervention points for antiviral drug

development (4, 9). Targeting the specific interaction interface—rather than globally

inhibiting kinesin activity—offers the possibility of selectively disrupting viral entry while

preserving essential cellular transport processes. A mechanistic understanding of this cargo

recognition pathway may therefore enable the rational design of small molecules or

peptides that block NTCP trafficking, providing a novel host-directed strategy to prevent

HBV/HDV infection.

In this study, we identify the molecular basis by which the kinesin motor protein KIF4

recognizes (NTCP and ITGB1) as a transport cargo and regulates its delivery to the

hepatocyte plasma membrane. We show that KIF4-mediated transport requires the

leucine-rich motif (LLLI) present at the C-terminal transmembrane domain of both

membrane proteins. Disruption of this motif abolishes the interaction between KIF4 and its

cargo (NTCP or ITGB1), markedly reduces the surface expression of both proteins without

affecting total protein levels. As a result, suppression of NTCP surface expression severely

impaired HBV and HDV entry. Together, these findings define a previously unrecognized

kinesin cargo-recognition motif that controls NTCP surface localization and represents a

critical regulatory node in HBV and HDV infection.

Materials and Methods

Cell lines and peptides

HepG2 were cultured with DMEM/F-12 + GlutaMax medium (Gibco) supplemented

with 10% fetal bovine serum (FBS), 10 mM HEPES, 100 unit/ml penicillin, 100 μg/ml

streptomycin, and 5 μg/ml insulin (4). HEK293FT cells were cultured in DMEM, high

glucose supplemented with 10% FBS, sodium pyruvate (1 mM), MEM non-essential amino

acids (1X), penicillin 100 unit/ml penicillin, 100 μg/ml streptomycin. All cell lines used in

this work were incubated at 37 °C in the presence of 5% CO_2 (9). Fluorescent HBV preS1

probe, a peptide spanning the 2-48 amino acids of the L-HBsAg preS1 region with

myristoylation and 6-carboxytetramethylrhodamine (TAMRA) labeling at N- and C-termini,

respectively, was synthesized by Scrum, Inc.

Plasmid Construction

The N-terminal 6×Myc-tagged KIF4 expression construct in the pIRESpuro3 vector was

kindly provided by Dr. Toru Hirota (Cancer Institute of the Japanese Foundation for Cancer

Research, JFCR) (16). C-terminally HA-tagged full-length NTCP (GenBank Accession No.

NM_003049.4), cloned into the CSII-EF-MCS vector at the *XhoI* (5′) and *XbaI* (3′) sites

and provided by Dr. Hiroyuki Miyoshi (RIKEN BioResource Research Center, Japan), was

subcloned into the pcDNA3.1 expression vector. The NTCP-HA open reading frame (ORF)

was excised by *XhoI* and *XbaI* and ligated into *XhoI/XbaI*-digested pcDNA3.1 to generate

full-length NTCP-HA in pcDNA3.1.

Generation of full-length NTCP-HA alanine-substitution mutants

157 The full-length NTCP-HA construct carrying four alanine substitutions (L298A, L299A,

158 L300A, and I301A) was generated by introducing these mutations into wild-type (WT)

159 NTCP-HA using a two-step PCR approach.

160 Two overlapping PCR fragments were first amplified from WT NTCP-HA using the

161 following primer pairs:

- **Fragment 1:**

Forward 5′-TAACGCTCGAGATGGAGGCCCACAAC-3′

Reverse   5′-

CTCATAGCACCAAAATATGGCGGCGGCGGCGGCCCCTTCTCCAAGCTGGAAA

ATC-3′

• **Fragment 2:**

Forward   5′-

GATTTTCCAGCTTGGAGAAGGGGCCGCCGCCGCCGCCATATTTTGGTGCTATG

AG-3′

Reverse 5′-GGCCCTCTAGACTAAGCGTAATCTGGAAC-3′

A third PCR reaction was then performed using the two overlapping amplicons as templates,

with the forward primer of Fragment 1 and the reverse primer of Fragment 2. The resulting

full-length mutated NTCP-HA fragment was digested with *XhoI* and *XbaI* and ligated into

XhoI/XbaI-digested pcDNA3.1.

Generation of NTCP truncated constructs

Two N-terminal NTCP truncations, NTCP-HA (aa 1–233) and NTCP-HA (aa 1–117),

were generated by PCR amplification of WT NTCP-HA using:

• **Forward primer** (both constructs):

5′-TGCCAGGTACCATGGAGGCCCACAAC-3′

• **Reverse primers:**

NTCP-HA (aa 1–233): 5′-

TCGGCTCTAGACTAAGCGTAATCTGGAACATCGTATGGGTAGCCAATAAAAGG

CATCAGGG-3′

NTCP-HA (aa 1–117): 5′-

TCGGCTCTAGACTAAGCGTAATCTGGAACATCGTATGGGTAGTTCATGTCCCC

CTTCATGG-3′

PCR products were digested with *KpnI* and *XbaI* and subcloned into *KpnI/XbaI*-digested

pcDNA3.1.

A C-terminal region construct NTCP-HA (aa 234–349) was similarly generated using:

• Forward 5′-TGCCAGGTACCATGTTTCTGCTGGGTTATG-3′

• Reverse 5′-

TCGGCTCTAGACTAAGCGTAATCTGGAACATCGTATGGGTAGGCTGTGCAAGG

GGAGCAGT-3′

and inserted into *KpnI/XbaI*-digested pcDNA3.1.

Generation of NTCP-HA (aa 234–349) constructs carrying alanine-substitution

mutations

The truncated NTCP-HA (aa 234–349) construct carrying the quadruple mutation

(L298A/L299A/L300A/I301A) was generated using the same overlapping PCR strategy

employed for full-length NTCP. Two PCR fragments were amplified from WT NTCP-HA

(aa 234–349) using the primer pairs:

• **Fragment 1:**

Forward 5′-TGCCAGGTACCATGTTTCTGCTGGGTTATG-3′

Reverse 5′-

CTCATAGCACCAAAATATGGCGGCGGCGGCGGCCCCTTCTCCAAGCTGGAAA

ATC-3′

• **Fragment 2:**

Forward 5′-

GATTTTCCAGCTTGGAGAAGGGGCCGCCGCCGCCGCCATATTTTGGTGCTATG

AG-3′

Reverse 5′-GGCCCTCTAGACTAAGCGTAATCTGGAAC-3′

The overlap-extension PCR product was digested with *KpnI* and *XbaI* and subcloned into

KpnI/XbaI-digested pcDNA3.1.

Generation of NTCP-HA (aa 234–349) single alanine-substitution mutants

Single alanine-substitution mutants of NTCP-HA (aa 234–349) (L298A, L299A, L300A,

or I301A) were generated using the same overlap-extension PCR strategy, with WT NTCP-

HA (aa 234–349) as the template.

For each mutant, two PCR reactions were performed as follows:

• **L298A mutant:**

PCR reaction 1:

Forward: 5′-TGCCAGGTACCATGTTTCTGCTGGGTTATG-3′

Reverse: 5′-

CTCATAGCACCAAAATATGGCAATGAGGAGGGCCCCTTCTCCAAGCTGGAAA

ATC-3′

PCR reaction 2:

Forward: 5′-

GATTTTCCAGCTTGGAGAAGGGGCCCTCCTCATTGCCATATTTTGGTGCTATGA

G-3′

Reverse: 5′-GGCCCTCTAGACTAAGCGTAATCTGGAAC-3′

• **L299A mutant:**

PCR reaction 1:

Forward: 5′-TGCCAGGTACCATGTTTCTGCTGGGTTATG-3′

Reverse: 5′-

CTCATAGCACCAAAATATGGCAATGAGGGCAAGCCCTTCTCCAAGCTGGAAA

ATC-3′

PCR reaction 2:

Forward: 5′-

GATTTTCCAGCTTGGAGAAGGGCTTGCCCTCATTGCCATATTTTGGTGCTATGA

G-3′

Reverse: 5′-GGCCCTCTAGACTAAGCGTAATCTGGAAC-3′

• **L300A mutant:**

PCR reaction 1:

Forward: 5′-TGCCAGGTACCATGTTTCTGCTGGGTTATG-3′

Reverse: 5′-

CTCATAGCACCAAAATATGGCAATGGCGAGAAGCCCTTCTCCAAGCTGGAAA

ATC-3′

PCR reaction 2:

Forward: 5′-

GATTTTCCAGCTTGGAGAAGGGCTTCTCGCCATTGCCATATTTTGGTGCTATGA

G-3′

Reverse: 5′-GGCCCTCTAGACTAAGCGTAATCTGGAAC-3′

• **I301A mutant:**

PCR reaction 1:

Forward: 5′-TGCCAGGTACCATGTTTCTGCTGGGTTATG-3′

Reverse: 5′-

CTCATAGCACCAAAATATGGCGGCGAGGAGAAGCCCTTCTCCAAGCTGGAAA

ATC-3′

PCR reaction 2:

Forward: 5′-

GATTTTCCAGCTTGGAGAAGGGCTTCTCCTCGCCGCCATATTTTGGTGCTATGA

G-3′

Reverse: 5′-GGCCCTCTAGACTAAGCGTAATCTGGAAC-3′

For each mutant, a third PCR reaction was performed using the products of the two

reactions as templates, with the forward primer from PCR reaction 1 and the reverse primer

from PCR reaction 2. The final PCR products were digested with *KpnI* and *XbaI* and

ligated into *KpnI*/*XbaI*-digested pcDNA3.1.

Plasmid transfection

Plasmid transfections in HEK293FT cells were performed using Lipofectamine 2000,

whereas transfections in HepG2 cells were carried out using Lipofectamine 3000, in

accordance with the manufacturer's instructions.

Immunoblot analysis

Protein expressions were analyzed by immunoblotting as previously described (17). The

primary antibodies used were mouse monoclonal anti-Myc (Santa Cruz Biotechnology, sc-

40), anti–E-cadherin (Santa Cruz Biotechnology, sc-8426), and anti–β-actin (Sigma-Aldrich,

A5441), as well as rabbit polyclonal anti-HA (Sigma-Aldrich, H6908). To assess NTCP

expression, protein samples were treated with Peptide-N-Glycosidase F (PNGase F; 250 U)

prior to SDS–PAGE to remove N-linked glycans from glycosylated NTCP, as previously

reported (9).

Isolation of the cell surface proteins

280 Cell surface proteins were isolated by surface biotinylation followed by streptavidin–

281 agarose affinity purification, as previously described (4). The recovered cell surface

282 fractions were subsequently analyzed by immunoblotting as described above. E-cadherin

283 (CDH1) was used as an internal loading control to verify the enrichment and integrity of

284 the surface protein fraction (18).

Co-immunoprecipitation (Co-IP) assay

HEK293FT cells were co-transfected with the indicated HA-tagged NTCP or HA-tagged

ITGB1 expression plasmids together with a Myc-tagged KIF4 expression plasmid at a 1:1

DNA ratio. At 72 h post-transfection, cells were harvested and lysed, and protein complexes

were subjected to co-immunoprecipitation (co-IP) using a mouse monoclonal anti-Myc

antibody (Santa Cruz Biotechnology, sc-40). Mouse normal IgG was used in parallel as a

negative control to assess assay specificity. Following immunoprecipitation, the recovered

complexes were analyzed by immunoblotting to detect co-immunoprecipitated HA-tagged

NTCP or HA-tagged ITGB1 in the pull-down fraction. Cell lysis and co-IP procedures were

performed using the Pierce Co-Immunoprecipitation Kit (Thermo Fisher Scientific, 26149)

in accordance with the manufacturer's instructions.

HBV preS1 peptide binding assay

Cell surface attachment of the N-terminally myristoylated and C-terminally TAMRA-labeled HBV preS1 peptide was assessed in HepG2 cells transiently expressing hNTCP as previously described (9).

HBV infection assay

HBV viral inoculum was prepared from the culture supernatant of HepAD38.7-Tet cells as previously described (19). HepG2 cells transiently expressing hNTCP were inoculated with HBV at a multiplicity corresponding to 10,000 genome equivalents (GEq) per cell in the presence of 4% PEG8000. Following a 16-h incubation, unbound viral particles were removed by extensive washing, and cells were maintained in culture for an additional 7 days. HBV infection was assessed by quantifying secreted HBeAg using ELISA and intracellular HBV pgRNA levels by real-time RT-PCR.

HDV infection assay

HDV used in infection experiments was generated from the culture supernatant of Huh7 cells co-transfected with pSVLD3 and pT7HB2.7 plasmids, as previously described (20, 21). HepG2 cells transiently expressing hNTCP were inoculated with HDV at 100 GEq per cell in the presence of 5% PEG8000. After a 16-h incubation, unbound virus was removed by extensive washing, and the cells were maintained in culture for an additional 6 days. HDV infection was evaluated by detection of HDAg using immunofluorescence analysis.

Immunofluorescence (IF) assay

Immunofluorescence staining was performed as previously described (4). Briefly, cells were fixed with 4% paraformaldehyde and permeabilized with 0.3% Triton X-100. The

samples were then incubated with a rabbit anti-HDAg primary antibody (developed by Scrum), followed by incubation with an Alexa Fluor 555–conjugated secondary antibody. Nuclei were counterstained with DAPI. Fluorescence signals were examined and imaged using a fluorescence microscope.

Structural prediction and comparative analysis of NTCP

The three-dimensional structures of wild-type NTCP and the NTCP mutants were predicted using AlphaFold 3.0.0 installed on a local server. The default databases were used, including the Google DeepMind–curated versions of BFD, MGnify (v2022_05), UniProt (2021_04), UniRef90 (2022_05), PDB, and PDB_seqres (2022_09_08). The wild-type NTCP sequence was obtained from UniProt (Q14973). For the mutants L298A, L299A, L300A, I301A, and the quadruple mutant L298A/L299A/L300A/I301A, the corresponding residues in the sequence were modified accordingly. For each prediction, a single random seed was used, and among the five generated output models, the structure with the highest confidence score was selected. RMSD calculations were performed using the Cα atoms of residues 22–309.

Amino acid sequence alignment

Multiple sequence alignments were performed for the C-terminal 116 amino acids of NTCP (aa 234–349) in comparison with human ITGB1, as well as for the NTCP C-terminal transmembrane sequence (aa 294–305) across HBV-susceptible species. Alignments were generated using the Clustal Omega algorithm (EMBL-EBI web server) with default parameters. Graphical representations of sequence conservation and alignment annotations were prepared using ESPript 3.2 (22-24).

340 **Statistical analysis**

341 All experiments were performed in triplicate, and data represent the mean ± standard

342 deviation (SD) from three independent experiments. Statistical significance was assessed

343 using an unpaired two-tailed Student's *t* test (**, P < 0.01; ***, P < 0.001; NS, not

344 significant).

345

346 **Results:**

347 **The C-terminal Domain of NTCP is crucial for interaction with KIF4**

348 We previously demonstrated that KIF4 mediates NTCP trafficking to the hepatocyte cell

349 surface(4). To further clarify the mechanism by which KIF4 recognizes and interacts with

350 its cargo, we constructed expression plasmids encoding C-terminal HA-tagged full-length

351 NTCP, as well as NTCP truncation mutants containing either the first 117 or 233 amino

352 acids (Fig. 1A). Each NTCP construct was separately co-transfected with a Myc-tagged

353 KIF4 expression plasmid into HEK293FT cells and incubated for 72 hours, after which co-

354 immunoprecipitation (co-IP) assays were performed. Immunoprecipitation with the anti-

355 Myc antibody resulted in efficient recovery of KIF4, while no KIF4 was detected in

356 samples processed with the isotype control antibody, thereby validating the specificity of

357 the assay (Fig. 1B). Notably, co-immunoprecipitation analysis demonstrated that KIF4

358 selectively associates with the full-length NTCP, whereas truncated NTCP variants failed to

359 associate with KIF4 (Fig. 1B), indicating that the interaction between NTCP and KIF4

360 requires the C-terminal region of NTCP.

361 **A leucine-rich motif (aa 298–301) in the C-terminal region of NTCP is essential for its**

362 **interaction with KIF4.**

363   In addition to NTCP, KIF4 has been reported to mediate the intracellular transport of

364 other host proteins, including human integrin β1 (ITGB1) (25). This raises the possibility

365 that shared functional determinants within these cargo proteins mediate their recognition by

366 KIF4. To identify candidate motifs within the NTCP C-terminal region that may be

367 responsible for KIF4 interaction, we aligned the amino acid sequence of the NTCP C-

368 terminal 116 residues (aa 234–349) with human ITGB1 sequence (Fig. 2A) (Fig. S1). The

369 alignment revealed that the NTCP C-terminal transmembrane domain (C-TMD; aa 281–

370 311) contains a leucine-rich motif (LLLI; aa 298–301), which is also present within the C-

371 TMD of ITGB1 (aa 729–751) at positions 747–750. This conservation suggested that the

372 LLLI motif might function as a determinant for KIF4 recognition. To experimentally

373 evaluate this hypothesis, we generated expression constructs encoding C-terminal HA-

374 tagged NTCP fragments comprising aa 234–349, either wild-type or carrying an alanine-

375 substitution mutation (AAAA) in place of the LLLI motif (Fig. 2B). Each construct was

376 individually co-transfected with a Myc-tagged KIF4 expression plasmid into HEK293FT

377 cells for 72 h, followed by co-IP analysis. Immunoprecipitation with anti-Myc antibody

378 efficiently recovered KIF4, whereas the isotype control yielded no detectable signal,

379 validating the specificity of the assay (Fig. 2C). Both full-length NTCP and the wild-type

380 NTCP (aa 234-349) fragment co-immunoprecipitated with KIF4. In contrast, the AAAA

381 mutant failed to associate with KIF4 (Fig. 2C). These results suggest that the leucine-rich

382 LLLI motif (aa 298–301) within the NTCP C-terminal region is essential for its interaction

383 with KIF4 and likely constitutes a key determinant for KIF4-dependent cargo recognition.

A conserved leucine-rich motif within the C-terminal region of ITGB1 (aa 747–750) is critical for KIF4-mediated recognition

To further assess the functional importance of the leucine-rich motif (LLLI; aa 747–750) located within the C-TMD of ITGB1 in mediating ITGB1–KIF4 interaction, we constructed expression plasmids encoding C-terminal HA-tagged full-length ITGB1 in either WT form or containing alanine mutations (AAAA) substituting the LLLI motif (Fig. 3A). Each construct was individually co-transfected with a Myc-tagged KIF4 expression vector into HEK293FT cells for 72 h, followed by co-IP analysis. Immunoprecipitation with an anti-Myc antibody efficiently recovered KIF4, whereas no signal was obtained with the isotype control antibody, confirming assay validity (Fig. 3B). Remarkably, WT full-length ITGB1 co-immunoprecipitated with KIF4, whereas the AAAA mutant did not (Fig. 3B). These results indicate that the leucine-rich LLLI motif (aa 747–750) within the ITGB1 C-TMD is crucial for KIF4-dependent recognition and interaction with ITGB1.

The C-terminal leucine-rich motif (aa 298–301) of NTCP is crucial for KIF4-mediated NTCP surface transport and efficient NTCP-dependent HBV/HDV infection

Given that NTCP is transported to the hepatocyte cell surface along microtubules by KIF4 to facilitate HBV and HDV entry (4), we investigated the functional importance of the C-terminal leucine-rich motif of NTCP (aa 298–301) in KIF4-mediated surface NTCP trafficking and HBV/HDV infection. To this end, we created expression plasmids encoding C-terminally HA-tagged full-length NTCP, either as the wild-type (NTCP^{WT}) or with alanine substitutions in the LLLI motif (NTCP^{AAAA}). HepG2 cells transiently expressing NTCP^{WT} or NTCP^{AAAA} for 72 h were exposed to biotinylation-based surface labeling assay

to selectively tag and purify cell surface protein fractions. Although total NTCP expression

levels were comparable between the two constructs (Fig. 4A, input), NTCP^{AAAA} exhibited

markedly reduced surface localization relative to NTCP^{WT}, which predominantly localized

to the cell membrane (Fig. 4A, surface). Consistent with impaired surface expression,

NTCP^{AAAA} expressing cells displayed significantly reduced binding of the HBV preS1

peptide ($P < 0.001$) (Fig. 4B and C). Furthermore, HBV infection was strongly attenuated

in NTCP^{AAAA} expressing cells, as evidenced by a significant reduction in intracellular HBV

pgRNA levels ($P < 0.001$) (Fig. 4D and E) and HBeAg secreted in the cell culture

supernatant ($P < 0.001$) (Fig. 4F) compared with NTCP^{WT} expressing cells. Similarly, HDV

infection was substantially decreased in NTCP^{AAAA} expressing cells, as demonstrated by a

marked reduction in HDAg positive cells ($P < 0.001$) (Fig. 4G and H). Importantly, the

NTCP^{AAAA} substitution did not adversely affect cell viability relative to NTCP^{WT} (Fig. 4I),

excluding cytotoxic effects. Collectively, these results demonstrate that the NTCP C-

terminal leucine-rich motif (aa 298–301) is essential for KIF4-mediated NTCP surface

transport and is consequently critical for efficient HBV and HDV entry.

Structural modeling demonstrates that the NTCP^{AAAA} substitution mutation does not

directly alter NTCP's binding affinity for the preS1 domain

Additionally, we conducted an *in silico* structural analysis using AlphaFold3 (26) to

predict the three-dimensional structures of NTCP^{WT}, the single alanine-substitution mutants

NTCP^{L298A}, NTCP^{L299A}, NTCP^{L300A}, and NTCP^{I301A}, and the quadruple mutant NTCP^{AAAA}.

The overall predicted folds were nearly identical across all variants, with root mean squared

deviations (RMSDs) of ≤ 0.3 Å relative to NTCP^{WT} (Fig. 5A, a-f). A close-up view of the

mutated C-terminal region likewise revealed no discernible local structural perturbations

(Fig. 5A, g). Because cell surface expressed NTCP mediates HBV entry through binding to

the preS1 domain of the large hepatitis B surface protein (LHBs) on hepatocytes (3), we

next assessed whether the AAAA substitution directly affects NTCP–preS1 interaction. To

this end, we employed DDMut-PPI (27), a deep learning–based model that predicts changes

in protein–protein interaction binding free energy upon multiple amino acid substitutions

(28). The predicted affinity change ($\Delta\Delta G^{\text{Affinity WT}\rightarrow\text{L298A, L299A, L300A, I301A}}$) was -1.11

kcal/mol (root mean squared error: 1.33 kcal/mol), predicting no substantial reduction in

the intrinsic binding affinity between NTCP and the HBV preS1 domain (Fig. 5B and C).

Collectively, these *in silico* analyses support the conclusion that the impaired ability of the

NTCP$^{\text{AAAA}}$ mutant to support HBV and HDV infection results from defective surface

trafficking rather than from a direct disruption of NTCP–preS1 binding.

440   L298 and L299 of NTCP constitute the minimal critical residues required for

441   mediating the interaction between NTCP and KIF4

As our results indicated that the NTCP leucine-rich motif (aa 298–301) is required for its

interaction with KIF4 and subsequent surface transport, we next sought to define the

minimal residues within this motif that mediate the interaction. To this end, we generated

expression plasmids encoding C-terminal HA-tagged NTCP fragments spanning aa 234–

349, either wild-type or carrying a single alanine substitution at L298, L299, L300, or I301

(Fig. 6A). Each construct was co-transfected with a Myc-tagged KIF4 expression plasmid

into HEK293FT cells for 72 h, followed by co-IP analysis. Immunoprecipitation with an

anti-Myc antibody efficiently recovered KIF4, whereas the isotype control yielded no

detectable signal, validating the assay specificity (Fig. 6B). The wild-type NTCP fragment

(aa 234–349), as well as the L300A and I301A variants, robustly co-immunoprecipitated

with KIF4. In contrast, fragments carrying L298A or L299A exhibited absent or markedly

reduced co-IP signals. As expected, the AAAA mutant fragment used as a negative control

failed to associate with KIF4 (Fig. 6B), further validating the co-IP specificity. Taken

together, these findings indicate that NTCP residues L298 and L299 represent the minimal

and essential determinants required for efficient NTCP–KIF4 interaction.

Discussion

ATP-dependent intracellular transport of cargo proteins along microtubule networks by

motor proteins is a fundamental process required for cellular homeostasis. Microtubule-

based motors, primarily kinesins and dyneins, are ATP-driven transporters that mediate the

directional trafficking of diverse intracellular cargos. Kinesins generally drive anterograde

transport toward the cell periphery, whereas dyneins and certain C-terminal kinesins

facilitate retrograde movement toward the perinuclear region (10). A growing body of

evidence demonstrates that these motor proteins play critical roles in viral infection by

directly interacting with viral components, thereby promoting viral replication and

propagation (29-32). In addition, we and others have shown that viruses can indirectly

exploit the kinesin transport machinery by engaging host proteins that are transported by

motor proteins and subsequently co-opted to support viral entry or replication (4, 25, 33).

Notably, the motor protein KIF4 interacts with and anterogradely transports several host

proteins, including human sodium taurocholate co-transporting polypeptide (NTCP), to the

hepatocyte cell surface, where NTCP serves as an entry receptor for HBV and HDV. KIF4 has also been reported to transport integrin β1 (ITGB1) and L1 cell adhesion molecule (L1CAM)–containing vesicles in developing cortical neurons (4, 7, 25). These observations raise the question of how KIF4 selectively recognizes and transports specific host cargos. Structurally, KIF4 contains an N-terminal ATPase motor domain responsible for microtubule engagement and a C-terminal cargo-binding tail that mediates cargo recognition (10). In the present study, we used NTCP and ITGB1 as model cargos to interrogate the molecular basis of KIF4-mediated transport.

Through a combination of amino acid sequence alignment and biochemical analyses, we identified a conserved C-terminal leucine-rich motif in both NTCP (aa 298–301) and ITGB1 (aa 747–750) that is essential for cargo recognition and transport mediated by KIF4. NTCP is a liver-specific, basolaterally expressed transporter that mediates the hepatic uptake of bile acids and functions as the bona fide entry receptor for HBV and HDV viruses (3, 34). In this study, alanine substitution of the NTCP leucine-rich motif (aa 298–301) disrupted normal NTCP surface localization, markedly reduced binding of a fluorescent HBV preS1 probe, and significantly attenuated HBV and HDV infection (Fig. 4). These findings underscore the functional importance of this motif in KIF4-mediated NTCP trafficking to the cell surface.

Although several NTCP orthologs from different species support HBV binding and infection (35, 36), sequence alignment of the C-terminal transmembrane sequence (aa 294–305) revealed that the putative leucine-rich motif (aa 298–301) is not strictly conserved among HBV-susceptible species (Fig. S3). The fact that this motif is not fully conserved

among different species supporting HBV infection suggests it does not play a direct role in mediating NTCP binding to the HBV preS1 domain. Instead, it may possess a function unique to the host cell, most likely in regulating the intracellular trafficking of NTCP, rather than directly facilitating viral binding to NTCP. Consistent with this interpretation, *in silico* structural modeling using AlphaFold supported that alanine substitution of this motif does not induce detectable global or local conformational changes in NTCP (Fig. 5). Comparison with the experimentally determined human NTCP cryo-EM structure (PDB: 7PQG) showed that all AlphaFold3 models most closely resembled the inward-facing conformation, with an RMSD of approximately 1.5 Å (34), and no mutation-associated structural perturbations were predicted. This apparent structural insensitivity may reflect inherent limitations of machine-learning–based prediction, which tends to favor conformations represented in the training data, or alternatively suggests that NTCP may adopt distinct, context-dependent conformational states influenced by membrane environment or interacting partners that are not captured by isolated *in silico* modeling. In parallel, DDMut-PPI analysis predicted only a minimal change in binding free energy between NTCP and the HBV preS1 domain upon introduction of the AAAA mutation, further supporting the conclusion that this motif does not directly contribute to intrinsic NTCP–preS1 binding affinity. While these computational predictions cannot substitute for direct biochemical binding measurements, they are consistent with prior studies identifying alternative NTCP residues as the primary determinants of HBV preS1 engagement and viral internalization (37-41).

Taken together with the comparable steady-state expression levels of NTCP WT and the AAAA mutant (Fig. S2) and the markedly reduced surface localization of the AAAA

variant, indicative of impaired motor-dependent transport, these data support a trafficking defect—rather than an intrinsic loss of receptor function—as the primary basis for the reduced NTCP-mediated HBV and HDV infectivity observed for the AAAA mutant. Collectively, our combined biochemical, structural, and computational analyses support a model in which the NTCP leucine-rich motif is dispensable for direct viral binding but is essential for KIF4-dependent NTCP recognition and surface transport, thereby indirectly regulating HBV and HDV entry.

Integrin β1 (ITGB1/CD29) is a transmembrane protein that forms functional heterodimers with multiple α-integrin subunits, including ITGA2 and ITGA3, thereby generating receptors that are essential for cellular homeostasis. Through binding to extracellular matrix (ECM) components such as laminin, collagen, and fibronectin, ITGB1-containing integrins physically link the ECM to the intracellular cytoskeleton and regulate key cellular processes, including adhesion, proliferation, migration, and gene expression (42, 43). Beyond these physiological functions, ITGB1 heterodimers also serve as entry factors for a variety of viruses (43-47) and contribute to tumor invasion and metastasis by enabling cancer cells to adhere to the basement membrane of target tissues (48). Collectively, these properties position ITGB1 as an attractive antiviral and anticancer target. In this context, elucidating the molecular mechanisms determining KIF4-dependent ITGB1 transport may provide a strategy to selectively disrupt ITGB1 surface delivery without broadly impairing other KIF4-dependent cellular functions, thereby offering a potential avenue for the development of targeted antiviral and anticancer therapies.

Here, we provide the first mechanistic insight into how the motor protein KIF4

selectively recognizes specific host cargo proteins for directed trafficking to the cell surface.

Our findings reveal that surface protein delivery is a highly organized and tightly regulated

process, in which defined sequence determinants enable selective engagement with distinct

motor proteins to ensure precise spatial localization. This work advances our understanding

of motor-driven secretory pathways and highlights cargo-specific recognition as a key

principle governing intracellular transport to the plasma membrane.

Acknowledgement

We gratefully acknowledge Dr. John Taylor at the Fox Chase Cancer Center, the USA for

providing pSVLD3 plasmid, Toru Hirota at JFCR, Japan for providing Myc-tagged KIF4;

Dr. Hiroyuki Miyoshi at RIKEN, Japan, for providing HA-tagged NTCP. This study was

supported by Grants in aid for Scientific Research C, 23K06575 and the Research Program

on Hepatitis B from AMED (grants number: 26fk0310532s0102; and 26fk0310528s0102)

to Hussein H. Aly, and the Research Program on Hepatitis B from AMED (grant number:

25fk0310532h0001) to Kazuaki Chayama.

<u>Figure legends</u>

FIG 1

KIF4 interacts with the C-terminal region of NTCP. (A) Schematic representation of C-terminally HA-tagged full-length NTCP (aa 1–349) and C-terminal truncation mutants comprising the N-terminal 117 (aa 1–117) or 233 (aa 1–233) amino acids. **(B)** HEK293FT cells were co-transfected with expression plasmids encoding Myc-tagged KIF4 and C-terminally HA-tagged full-length NTCP or the indicated NTCP truncation mutants at a 1:1 ratio. At 72 h post-transfection, cells were lysed, and 10% of each lysate was analyzed by immunoblotting to detect Myc-KIF4 (*Input; uppermost panel*), HA-NTCP (*Input; middle panel*), and β-actin (loading control) (*Input; lowermost panel*). The remaining lysates (90%) were subjected to co-immunoprecipitation using either an isotype control antibody or anti-Myc antibody to pull down Myc-KIF4. Immunoprecipitates were analyzed by immunoblotting for Myc-KIF4 (*IP; upper panel*) and co-immunoprecipitated HA-NTCP (*IP; lower panel*). Data are representative of three independent experiments.

FIG 2

KIF4 recognizes NTCP via a leucine-rich motif (aa 298–301) at the C-terminal end of NTCP. (A) Schematic representation of the NTCP truncation mutant comprising the C-

terminal 116 residues (aa 234–349) and full-length human ITGB1 (aa 1–798). Black boxes

denote the C-terminal transmembrane domains (CT-TMDs) of NTCP (aa 281–311; upper)

and ITGB1 (aa 729–751; lower). The shared leucine-rich motif within the CT-TMDs is

highlighted in yellow. **(B)** Schematic representation of C-terminally HA-tagged NTCP

truncation mutant (aa 234–349) either WT or carrying alanine substitutions at residues

L298, L299, L300, and I301. **(C)** HEK293FT cells were co-transfected with expression

plasmids encoding Myc-tagged KIF4 and C-terminally HA-tagged full-length NTCP or the

NTCP (aa 234–349) truncation mutant, either WT or carrying the indicated alanine

substitutions, at a 1:1 ratio. At 72 h post-transfection, cells were lysed, and 10% of each

lysate was analyzed by immunoblotting to detect Myc-KIF4 (*Input; uppermost panel*), HA-

NTCP (*Input; middle panel*), and β-actin (loading control) (*Input; lowermost panel*). The

remaining lysates (90%) were subjected to co-immunoprecipitation using either an isotype

control antibody or anti-Myc antibody to pull down Myc-KIF4. Immunoprecipitates were

analyzed by immunoblotting for Myc-KIF4 (*IP; upper panel*) and co-immunoprecipitated

HA-NTCP (*IP; lower panel*). Data are representative of three independent experiments.

FIG 3

KIF4 selectively recognizes ITGB1 via a leucine-rich motif (amino acids 747–750) at

the C-terminal end of ITGB1. **(A)** Schematic representation of C-terminally HA-tagged

full-length human ITGB1 (aa 1–798) either WT or harboring alanine substitutions at

residues L747, L748, L749, and I750. **(B)** HEK293FT cells were co-transfected with

expression plasmids encoding Myc-tagged KIF4 and C-terminally HA-tagged full-length

ITGB1 either WT or carrying the indicated alanine substitutions, at a 1:1 ratio. At 72 h

post-transfection, cells were lysed, and 10% of each lysate was analyzed by

immunoblotting to detect Myc-KIF4 (*Input; uppermost panel*), HA-ITGB1 (*Input; middle*

panel), and β-actin (loading control) (*Input; lowermost panel*). The remaining lysates

(90%) were subjected to co-immunoprecipitation using either an isotype control antibody

or anti-Myc antibody to pull down Myc-KIF4. Immunoprecipitates were analyzed by

immunoblotting for Myc-KIF4 (*IP; upper panel*) and co-immunoprecipitated HA-ITGB1

(*IP; lower panel*). Data are representative of three independent experiments.

FIG 4

The NTCP C-terminal leucine-rich motif (aa 298–301) is required for KIF4-dependent

surface trafficking and HBV/HDV entry. (A) HepG2 cells were transfected with

plasmids expressing C-terminally HA-tagged full-length NTCP WT (NTCP^{WT}) or carrying

the L298A, L299A, L300A, I301A substitutions (NTCP^{AAAA}). At 72 h post-transfection, the

cells were surface-biotinylated or PBS-treated at 4°C for 30 min. prior to cell lysis. Ten

percent of each lysate was analyzed by immunoblotting to detect total NTCP (*Input; upper*

panel), and β-actin (loading control) (*Input; lower panel*). The remaining lysates were

incubated with streptavidin (SA) beads to isolate biotinylated surface proteins. Eluted

fractions were analyzed by immunoblotting to detect surface NTCP (*Surface; upper panel*)

and CDH-1 (loading control for surface fraction) (*Surface; lower panel*). **(B)** HepG2 cells

ectopically expressing NTCP^{WT} or NTCP^{AAAA} were incubated with 40 nM C-terminal

TAMRA-labeled and N-terminally myristoylated preS1 peptide (preS1 probe) for 30 min at

37°C; Red and blue signals indicate the preS1 probe binding and nuclei, respectively. **(C)**

Quantification of relative fluorescence intensity from panel B. **(D)** Schematic diagram

depicting the scheme for plasmid transfection and subsequent HBV infection in HepG2;

HepG2 cells were transfected with Mock, NTCP^{WT}, or NTCP^{AAAA} plasmids for 72 h,

followed by HBV inoculation (10,000 GEq/cell) in the presence of 4% PEG8000 for 16 h.

After washing out free viral particles, the cells were cultured for an additional 7 days,

followed by detection of different HBV markers. **(E)** Intracellular HBV pgRNA levels were

quantified by RT-qPCR and presented as fold changes, relative to NTCP^{WT} transfected cells.

(F) secreted HBeAg levels in the culture supernatant was measured by ELISA and

presented as (A/CO) values. **(G)** HepG2 expressing NTCP^{WT} or NTCP^{AAAA} were

inoculated with HDV (100 GEq/cell) in the presence of 5% PEG8000 for 16 h. After

washing and removal of the free viral particles, the cells were cultured for 6 days followed

by detection of intracellular HDAg by IF; Red and blue signals indicate HDAg and nuclei,

respectively. **(H)** Quantification of fluorescence intensity from panel G. **(I)** Cell viability

was also assessed by MTT assay. All assays were performed in triplicate, and data from

three independent experiments were included. Data are presented as mean ± SD. ***, $P <$

0.001; NS, not significant.

FIG 5

***In silico* structural modeling suggests the NTCP^{AAAA} substitution mutation does not**

alter NTCP/preS1 binding. (A) Panels A (a–f) show the predicted human NTCP structures

of the wild type and the single alanine mutants; L298A, L299A, L300A, and I301A, and

the quadruple mutant L298A/L299A/L300A/I301A, respectively. Panel A (g) overlays the structures at the mutation sites. The colors for L298A, L299A, L300A, I301A, and L298A/L299A/L300A/I301A are magenta, yellow, tint, cyan, and green, respectively. **(B)** Panel B (a, b) shows the predicted human NTCP and PreS1 structures of the wild type (a) and the quadruple mutant L298A/L299A/L300A/I301A (b), respectively. **(C)** Predicted effects of single and quadruple alanine substitutions within the NTCP leucine-rich motif on binding free energy between NTCP and the HBV preS1 domain. Individual $\Delta\Delta G^{Binding}$ values (kcal/mol) are shown for the single mutants (L298A, L299A, L300A, I301A), along with the predicted $\Delta\Delta G^{Binding}$ (kcal/mol) for the quadruple mutant (L298A/L299A/L300A/I301A).

FIG 6

NTCP residues L298 and L299 are the minimal determinants essential for interaction with KIF4. (A) Schematic representation of C-terminally HA-tagged NTCP truncation mutant (aa 234–349) either WT, carrying single alanine substitutions at residues L298, L299, L300, or I301, or quadruple alanine substitutions (L298A/L299A/L300A/I301A). **(B)** HEK293FT cells were co-transfected with expression plasmids encoding Myc-tagged KIF4 and C-terminally HA-tagged NTCP (aa 234–349) truncation mutant, either WT or carrying the indicated alanine substitutions, at a 1:1 ratio. At 72 h post-transfection, cells were lysed, and 10% of each lysate was analyzed by immunoblotting to detect HA-NTCP variants (*Input; uppermost panel*), Myc-KIF4 (*Input; middle panel*), and β-actin (loading control) (*Input; lowermost panel*). The remaining lysates (90%) were subjected to co-

670 immunoprecipitation using either an isotype control antibody or anti-Myc antibody to pull

671 down Myc-KIF4. Immunoprecipitates were analyzed by immunoblotting for Myc-KIF4

672 (*IP; upper panel*) and co-immunoprecipitated HA-NTCP (*IP; lower panel*). Data are

673 representative of three independent experiments.

References

1. Burki T. 2024. WHO's 2024 global hepatitis report. Lancet Infect Dis 24:e362-e363.
2. Yang S, Ye Q, Yang Y, Ma Z. 2025. Clinical impact of hepatitis delta virus coinfection on liver fibrosis
in hepatitis B patients: a population-based study. Virol J 22:273.
3. Yan H, Zhong G, Xu G, He W, Jing Z, Gao Z, Huang Y, Qi Y, Peng B, Wang H, Fu L, Song M, Chen P, Gao
W, Ren B, Sun Y, Cai T, Feng X, Sui J, Li W. 2012. Sodium taurocholate cotransporting polypeptide is a
functional receptor for human hepatitis B and D virus. Elife 1:e00049.
4. Gad SA, Sugiyama M, Tsuge M, Wakae K, Fukano K, Oshima M, Sureau C, Watanabe N, Kato T,
Murayama A, Li Y, Shoji I, Shimotohno K, Chayama K, Muramatsu M, Wakita T, Nozaki T, Aly HH.
2022. The kinesin KIF4 mediates HBV/HDV entry through the regulation of surface NTCP localization
and can be targeted by RXR agonists in vitro. PLoS Pathog 18:e1009983.
5. Sheng L, Hao SL, Yang WX, Sun Y. 2018. The multiple functions of kinesin-4 family motor protein
KIF4 and its clinical potential. Gene 678:90-99.
6. Martinez NW, Xue X, Berro RG, Kreitzer G, Resh MD. 2008. Kinesin KIF4 regulates intracellular
trafficking and stability of the human immunodeficiency virus type 1 Gag polyprotein. J Virol
82:9937-50.
7. Peretti D, Peris L, Rosso S, Quiroga S, Caceres A. 2000. Evidence for the involvement of KIF4 in the
anterograde transport of L1-containing vesicles. J Cell Biol 149:141-52.
8. Zhou J, Scherer J, Yi J, Vallee RB. 2018. Role of kinesins in directed adenovirus transport and
cytoplasmic exploration. PLoS Pathog 14:e1007055.
9. Gad SA, Abo Elwafa DA, Hussein SA, Fukano K, Kato T, Shoji I, Chayama K, Muramatsu M, Isogawa M,
Wakita T, Merk D, Aly HH. 2025. RXR agonist S169 inhibits HBV/HDV entry in vitro by disrupting
KIF4-dependent NTCP trafficking. Antiviral Res 240:106214.
10. Hirokawa N, Noda Y, Tanaka Y, Niwa S. 2009. Kinesin superfamily motor proteins and intracellular
transport. Nat Rev Mol Cell Biol 10:682-96.
11. Verhey KJ, Hammond JW. 2009. Traffic control: regulation of kinesin motors. Nat Rev Mol Cell Biol
10:765-77.
12. Jiang X, Ogawa T, Yonezawa K, Shimizu N, Ichinose S, Uchihashi T, Nagaike W, Moriya T, Adachi N,
Kawasaki M, Dohmae N, Senda T, Hirokawa N. 2023. The two-step cargo recognition mechanism of
heterotrimeric kinesin. EMBO Rep 24:e56864.
13. Sharma P, Cline HT. 2010. Visual activity regulates neural progenitor cells in developing xenopus CNS
through musashi1. Neuron 68:442-55.
14. Chui SS, Lo SM, Charmant JP, Orpen AG, Williams ID. 1999. A chemically functionalizable
nanoporous material. Science 283:1148-50.

15.   Gelfond D, Borowitz D. 2013. Gastrointestinal complications of cystic fibrosis. Clin Gastroenterol
711   Hepatol 11:333-42; quiz e30-1.

712   16.   Takahashi M, Wakai T, Hirota T. 2016. Condensin I-mediated mitotic chromosome assembly requires
713   association with chromokinesin KIF4A. Genes Dev 30:1931-6.

714   17.   Aly HH, Suzuki J, Watashi K, Chayama K, Hoshino S, Hijikata M, Kato T, Wakita T. 2016. RNA Exosome
715   Complex Regulates Stability of the Hepatitis B Virus X-mRNA Transcript in a Non-stop-mediated
716   (NSD) RNA Quality Control Mechanism. J Biol Chem 291:15958-74.

717   18.   Londino JD, Gulick DL, Lear TB, Suber TL, Weathington NM, Masa LS, Chen BB, Mallampalli RK. 2017.
718   Post-translational modification of the interferon-gamma receptor alters its stability and signaling.
719   Biochem J 474:3543-3557.

720   19.   Ibrahim MK, Abdelhafez TH, Takeuchi JS, Wakae K, Sugiyama M, Tsuge M, Ito M, Watashi K, El Kassas
721   M, Kato T, Murayama A, Suzuki T, Chayama K, Shimotohno K, Muramatsu M, Aly HH, Wakita T. 2021.
722   MafF Is an Antiviral Host Factor That Suppresses Transcription from Hepatitis B Virus Core Promoter.
723   J Virol 95:e0076721.

724   20.   Kuo MY, Chao M, Taylor J. 1989. Initiation of replication of the human hepatitis delta virus genome
725   from cloned DNA: role of delta antigen. J Virol 63:1945-50.

726   21.   Sureau C, Guerra B, Lee H. 1994. The middle hepatitis B virus envelope protein is not necessary for
727   infectivity of hepatitis delta virus. J Virol 68:4063-6.

728   22.   Ma H, Kim AS, Kafai NM, Earnest JT, Shah AP, Case JB, Basore K, Gilliland TC, Sun C, Nelson CA,
729   Thackray LB, Klimstra WB, Fremont DH, Diamond MS. 2020. LDLRAD3 is a receptor for Venezuelan
730   equine encephalitis virus. Nature 588:308-314.

731   23.   Sievers F, Higgins DG. 2018. Clustal Omega for making accurate alignments of many protein
732   sequences. Protein Sci 27:135-145.

733   24.   Robert X, Guillon C, Gouet P. 2025. FoldScript: a web server for the efficient analysis of AI-generated
734   3D protein models. Nucleic Acids Res 53:W277-W282.

735   25.   Heintz TG, Heller JP, Zhao R, Caceres A, Eva R, Fawcett JW. 2014. Kinesin KIF4A transports integrin
736   beta1 in developing axons of cortical neurons. Mol Cell Neurosci 63:60-71.

737   26.   Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore L, Ballard AJ,
738   Bambrick J, Bodenstein SW, Evans DA, Hung CC, O'Neill M, Reiman D, Tunyasuvunakool K, Wu Z,
739   Zemgulyte A, Arvaniti E, Beattie C, Bertolli O, Bridgland A, Cherepanov A, Congreve M, Cowen-Rivers
740   AI, Cowie A, Figurnov M, Fuchs FB, Gladman H, Jain R, Khan YA, Low CMR, Perlin K, Potapenko A,
741   Savy P, Singh S, Stecula A, Thillaisundaram A, Tong C, Yakneen S, Zhong ED, Zielinski M, Zidek A,
742   Bapst V, Kohli P, Jaderberg M, Hassabis D, Jumper JM. 2024. Accurate structure prediction of
743   biomolecular interactions with AlphaFold 3. Nature 630:493-500.

744   27.   Zhou Y, Myung Y, Rodrigues CHM, Ascher DB. 2024. DDMut-PPI: predicting effects of mutations on
745   protein-protein interactions using graph-based deep learning. Nucleic Acids Res 52:W207-W214.

746   28.   Zhang L, Struss A, Qin M, Dombrowski A, Li T, Viebahn S, Mehlhorn D, Grefen C. 2025. A
747   phosphoinositide-mediated switch of GET pathway receptor dimerization in Arabidopsis. Proc Natl
748   Acad Sci U S A 122:e2514354122.

749   29.   Kelkar SA, Pfister KK, Crystal RG, Leopold PL. 2004. Cytoplasmic dynein mediates adenovirus binding
750   to microtubules. J Virol 78:10122-32.

751   30.   Kelkar S, De BP, Gao G, Wilson JM, Crystal RG, Leopold PL. 2006. A common mechanism for
752   cytoplasmic dynein-dependent microtubule binding shared among adeno-associated virus and
753   adenovirus serotypes. J Virol 80:7781-5.

754   31.   Strunze S, Engelke MF, Wang IH, Puntener D, Boucke K, Schleich S, Way M, Schoenenberger P,
755   Burckhardt CJ, Greber UF. 2011. Kinesin-1-mediated capsid disassembly and disruption of the
756   nuclear pore complex promote virus infection. Cell Host Microbe 10:210-23.

757   32.   Malikov V, da Silva ES, Jovasevic V, Bennett G, de Souza Aranha Vieira DA, Schulte B, Diaz-Griffero F,
758   Walsh D, Naghavi MH. 2015. HIV-1 capsids bind and exploit the kinesin-1 adaptor FEZ1 for inward

movement to the nucleus. Nat Commun 6:6660.

33. Ravindran MS, Engelke MF, Verhey KJ, Tsai B. 2017. Exploiting the kinesin-1 molecular motor to
generate a virus membrane penetration site. Nat Commun 8:15496.

34. Goutam K, Ielasi FS, Pardon E, Steyaert J, Reyes N. 2022. Structural basis of sodium-dependent bile
salt uptake into the liver. Nature 606:1015-1020.

35. Xu Z, Zhao K, Wang J, Zhang L, Yin J, Chen N, Chen S, Zhao G, Wang M, Xin T, Zhu C, Cheng X, Xia Y.
2025. Screening of different species reveals cat hepatocytes support HBV infection. PLoS Pathog
21:e1013390.

36. Chen F, Wettengel JM, Gegenfurtner F, Moosmuller J, Bunse T, Jeske SD, Hagen P, Ni Y, Urban S,
Protzer U. 2025. Identification of NTCP animal orthologs supporting hepatitis B virus binding and
infection. J Virol 99:e0183324.

37. Yan H, Peng B, He W, Zhong G, Qi Y, Ren B, Gao Z, Jing Z, Song M, Xu G, Sui J, Li W. 2013. Molecular
determinants of hepatitis B and D virus entry restriction in mouse sodium taurocholate
cotransporting polypeptide. J Virol 87:7977-91.

38. Matsuzawa K, Ekimoto T, Kobayashi C, Shionoya K, Mifune J, Morita T, Takeuchi JS, Park SY, Ikeguchi
M, Sureau C, Kawaguchi A, Watashi K. 2026. Structural mapping of NTCP distinguishes its dual
functionality as a hepatitis B virus receptor and bile acid transporter. PLoS Pathog 22:e1013824.

39. Takeuchi JS, Fukano K, Iwamoto M, Tsukuda S, Suzuki R, Aizaki H, Muramatsu M, Wakita T, Sureau C,
Watashi K. 2019. A Single Adaptive Mutation in Sodium Taurocholate Cotransporting Polypeptide
Induced by Hepadnaviruses Determines Virus Species Specificity. J Virol 93.

40. Liu H, Zakrzewicz D, Nosol K, Irobalieva RN, Mukherjee S, Bang-Sorensen R, Goldmann N, Kunz S,
Rossi L, Kossiakoff AA, Urban S, Glebe D, Geyer J, Locher KP. 2024. Structure of antiviral drug
bulevirtide bound to hepatitis B and D virus receptor protein NTCP. Nat Commun 15:2476.

41. Zakrzewicz D, Leidolf R, Kunz S, Muller SF, Neubauer A, Leiting S, Goldmann N, Lehmann F, Glebe D,
Geyer J. 2022. Tyrosine 146 of the Human Na(+)/Taurocholate Cotransporting Polypeptide (NTCP) Is
Essential for Its Hepatitis B Virus (HBV) Receptor Function and HBV Entry into Hepatocytes. Viruses
14.

42. Liu S, Calderwood DA, Ginsberg MH. 2000. Integrin cytoplasmic domain-binding proteins. J Cell Sci
113 (Pt 20):3563-71.

43. Fu R, Jordan P, Engels Z, Weihs JA, Murle J, Chi H, Burbano de Lara S, Helm B, Klohn M, Hu J,
Freistaedter A, Boettler T, Binder M, Klingmuller U, Steinmann E, Lozach PY, Tubiana T, Lemon SM,
Dao Thi VL. 2025. Integrin beta 1 facilitates non-enveloped hepatitis E virus cell entry through the
recycling endosome. Nat Commun 16:5403.

44. Shuai L, Wang J, Zhao D, Wen Z, Ge J, He X, Wang X, Bu Z. 2020. Integrin beta1 Promotes Peripheral
Entry by Rabies Virus. J Virol 94.

45. Izmailyan R, Hsao JC, Chung CS, Chen CH, Hsu PW, Liao CL, Chang W. 2012. Integrin beta1 mediates
vaccinia virus entry through activation of PI3K/Akt signaling. J Virol 86:6677-87.

46. Kaminsky PM, Keiser NW, Yan Z, Lei-Butters DC, Engelhardt JF. 2012. Directing integrin-linked
endocytosis of recombinant AAV enhances productive FAK-dependent transduction. Mol Ther
20:972-83.

47. Rivera-Serrano EE, Gonzalez-Lopez O, Das A, Lemon SM. 2019. Cellular entry and uncoating of naked
and quasi-enveloped human hepatoviruses. Elife 8.

48. Hood JD, Cheresh DA. 2002. Role of integrins in cell invasion and migration. Nat Rev Cancer 2:91-
100.

Supplementary figure legends

FIG S1

Sequence alignment of the NTCP C-terminal region with human ITGB1. Alignment of

the C-terminal 116 amino acids of NTCP (aa 234–349) with human ITGB1. Conserved

residues and regions of sequence similarity are marked or highlighted.

FIG S2

Protein expression of NTCP and ITGB1 (WT *vs* quadruple alanine mutant) in

HEK293 FT cells. HEK293FT cells were transfected with mock plasmid or plasmids

expressing C-terminally HA-tagged full-length NTCP WT, NTCP carrying

L298A/L299A/L300A/I301A substitutions, C-terminally HA-tagged full-length ITGB1 WT,

or ITGB1 harboring L747A/L748A/L749A/I750A substitutions. At 72 h post-transfection,

cells were lysed and analyzed by immunoblotting to detect NTCP or ITGB1 (*upper panel*)

and β-actin (*lower panel*; loading control). Data are representative of three independent

experiments. Data are representative of three independent experiments.

FIG S3

Sequence alignment of NTCP residues 294–305 within the C-terminal transmembrane

region across HBV-susceptible species. The C-terminal transmembrane segment (aa 294–

305) of NTCP orthologs from human (*Homo sapiens*), cat (*Felis catus*), rabbit (*Oryctolagus*

cuniculus), ferret (*Mustela putorius furo*), woodchuck (*Marmota marmota*), big brown bat

(*Eptesicus fuscus*), aardvark (*Orycteropus afer afer*), horse (*Equus caballus*), chinese tree

shrew (*Tupaia chinensis*), whale (*Balaenoptera acutorostrata scammony*) was shown.

Residues corresponding to the putative leucine-rich motif identified in human NTCP are

highlighted in purple.

Fig. 1

Fig. 2

A

Truncated NTCP
(aa 234-349)

281 PLFFFPLLYMIFQLGEG**LLL**IAIFWCYEKFFK 311 aa

C-terminal TMD

Full length
Human ITGB1
(aa 1-798)

729 IIPIVAGVVAGIVLIGLA**LLL**IW 751 aa

C-terminal TMD

B

Truncated NTCP
(aa 234-349)

HA Tag

WT 281 PLFFFPLLYMIFQLGEG**LLL**IAIFWCYEKFFK 311 aa
298   301

L298A, L299A,
L300A, I301A   PLFFFPLLYMIFQLGEG**AAAA**IFWCYEKFFK

C

Myc-KIF4A

NTCP
(aa 1-349)

NTCP
(aa 234-349)

β-actin

Input

Myc-KIF4A

NTCP
(aa 1-349)

NTCP
(aa 234-349)

IP

	WT	L298A, L299A, L300A, I301A

| Control-IgG | + | − | − | − |
| Myc-IgG | − | + | + | + |

Fig. 3

Fig. 4

A

Surface Fractionation

Surface
HA-NTCP — KDa
— 35
CDH-1 — 100

Input
HA-NTCP — 35
β-actin — 55

Biotinylation: − + +
NTCP^WT: − + −
NTCP^AAAA: − − +

B

DAPI | preS1 Probe | Merge
NTCP^WT
NTCP^AAAA

C

Fluorescent area (fold)
***
NTCP^WT NTCP^AAAA

D

HBV infection timeline
HBV
HA-NTCP plasmid
72 h | 16 h | 7 d
Plasmid Transfection | HBV | Wash | Detection

E

HBV pgRNA (fold)
*** ***
Mock NTCP^WT NTCP^AAAA

F

HBeAg (A/Co)
**** ****
Mock NTCP^WT NTCP^AAAA

G

DAPI | HDAg | Merge
NTCP^WT
NTCP^AAAA

H

Fluorescent area (fold)
***
NTCP^WT NTCP^AAAA

I

Cell viability (fold)
NS
NTCP^WT NTCP^AAAA

Fig. 5

NTCP (PDB ID:8RQF) mutation	Average distance (Å)	Individual ΔΔG^{Binding} kcal/mol	Predicted ΔΔG^{Binding} kcal/mol
A L298A; A L299A; A L300A; A I301A	4.56	{'A L298A': -0.24567555, 'A L299A': -0.29362273, 'A L300A': -0.26692277, 'A I301A': -0.30670142}	-1.11

Fig. 6