**Colitis-induced visceral pain recruits central neurotensin neurons that modulate colonic**

**sensitivity**

**Authors**

Yu-Ting Cheng[1,2], Natalie MacKinnon-Booth[1], Yingfu Jiao[1], Jordan R. Robbins[1], Murillo Duarte-

Silva[3], Perry E. Mitchell[3], Yuchu Liu[1], Omer Barkai[1,2], Keunjung Heo[1,2], Biyao Zhang[1], Bruna Lenfers

Turnes[1,2], Meenakshi Rao[3] and Clifford J. Woolf[1,2]*

**Affiliations**

[1]F.M. Kirby Neurobiology Center, Boston Children's Hospital, Boston, MA, USA

[2]Department of Neurobiology, Harvard Medical School, Boston, MA, USA

[3]Division of Gastroenterology, Department of Pediatrics, Boston Children's Hospital and Harvard

Medical School, Boston, USA

*Correspondence: clifford.woolf@childrens.harvard.edu

**Abstract**

Inflammatory bowel disease produces debilitating visceral pain that remains a major clinical challenge. Notably, many patients experience persistent pain even after the inflammation resolves, indicating a sustained sensitization of central neural circuits that drives enduring pain. The brainstem parabrachial nucleus integrates interoceptive signals from the gastrointestinal tract to elicit both pain perception and affective responses. Using activity-dependent mapping and an RNAscope assay, we identified a neurotensin (NT)-expressing neuronal population in the lateral PBN ($PBN_L$) that is selectively activated during dextran sulfate sodium-induced colitis. *In vivo* neural activity recordings demonstrate that $PBN_L$ NT neurons encode colon-derived nociceptive signals in an intensity-dependent manner. Silencing these neurons attenuates colonic reflexes evoked by luminal distension and normalizes aberrant gastrointestinal transit and nociceptive licking behavior in colitic mice. Pharmacological blockade of NT signaling alleviates colitis-associated hypersensitivity. These findings identify a central neural population that encodes visceral inflammation and regulates peripheral organ function, and pinpoints neurotensin as a promising therapeutic target to treat colitis-induced visceral pain.

## Introduction

Pain associated with inflammatory bowel disease (IBD) affects up to 10 million people worldwide and arises from the activation of nociceptor afferents by inflammatory responses mediated by a compromised epithelial lining of the gut barrier[1-3]. Clinical hallmarks of colitis include spontaneous abdominal pain, visceral hypersensitivity, and altered bowel function, such as diarrhea and dysmotility[4,5]. Current pharmacological treatments primarily target mucosal inflammation; however, tissue healing verified by endoscopy or histopathology, does not reliably produce symptomatic relief, with 30–50% of patients continuing to report persistent pain even after resolution of their inflammatory flare[6,7]. This suggests that a central neural sensitization may contribute to the maintenance of visceral pain independently of ongoing peripheral pathology[8-10].

The brainstem parabrachial complex serves as a key integrative hub that relays interoceptive and exteroceptive signals from the periphery to forebrain regions to elicit pain and affective perception[11-15]. Within this complex, the lateral parabrachial nucleus ($PBN_L$) primarily receives nociceptive input from the spinal dorsal horn via the anterolateral pathway and is engaged in acute and chronic pain states[16-18]. Genetic and circuit-level dissection of $PBN_L$ neuronal populations, together with an analysis of their neuropeptide signaling has begun to reveal a powerful leverage mechanism for modulating pain and its associated emotional disturbances[19-22]. However, despite accumulating evidence implicating the role of $PBN_L$ in somatosensory pain[23-26], the identity of those neuronal ensembles that encode visceral nociceptive signals, and if and how these influence organ function and visceral pain-related behavior, are unclear.

We identified a population of neurotensin (NT)-expressing neurons within the $PBN_L$ that is selectively activated during dextran sulfate sodium (DSS)-induced colitis and encodes colon-derived nociceptive information in an intensity-dependent manner. Silencing the $PBN_L$ NT neurons abrogated the exaggerated colonic reflexes evoked by pressure distension, while blockade of neurotensin signaling normalized the diarrhea and pain-associated behaviors induced by the colitis. These findings provide evidence for a central neural modulation of peripheral visceral function and pinpoint $PBN_L$ NT neurons as a mechanistically defined node linking visceral inflammatory pain and affective states, with promising implications for treating colitis-induced visceral pain.

## Results

### Activation of nociceptors by colitis drives spontaneous nocifensive behaviors

We modeled experimental colitis by administering 3% DSS into the drinking water of mice[27,28] for 4 days and measured spontaneous pain-related behaviors together with abdominal allodynia for 2 weeks to evaluate the temporal profile of altered nociceptive behavioral responses (**Fig. 1a**). Colon inflammation was confirmed by an infiltration of CD45-positive immune cells into distal colon tissues 4 days after DSS exposure (D4) and 3 days after the DSS withdrawal (D7), which progressively resolved over 2 weeks (**Fig. 1b,c**). Additionally, we monitored a reduction of mouse body weight and water intake in DSS-treated WT mice compared to regular $H_2O$ drinking water controls, which gradually recovered on D14 (**Extended Data Fig. 1a,b**).

We implemented a supervised machine learning algorithm to automatically screen for pain-related behavior signatures relevant to the DSS-induced colitis[29,30]. This analysis revealed a significant increase in "lower body licking" encompassing abdominal and peri-anal licking during the active inflammation, when compared with mice receiving regular $H_2O$ drinking water (**Fig. 1d**). A licking bout was defined as contact between the snout and the abdominal or peri-anal region sustained for at least two consecutive video frames. To interpret model features underlying licking classification, we analyzed the 10 most informative features using SHAP (Shapley additive explanations). Spatial features contributed most strongly to the model performance, with the distance between the snout/neck and the tail base ranked as the top feature, reflecting close proximity of the head to the lower body during licking behavior (**Extended Data Fig. 2a**). Model performance improved with increasing training data, as shown by the F1-score learning curve, and reached a plateau near 0.8 after approximately 6,000 labeled frames (**Extended Data Fig. 2b**). To further optimize classification performance, we evaluated precision, recall, and F1-score across a range of discrimination thresholds on the validation dataset. This analysis identified an optimal operating threshold centered around ~0.6, which was used for subsequent behavioral quantification (**Extended Data Fig. 2c**).

A correlation between time spent licking and the stool-based disease activity index (DAI), a score calculated as the sum of two components: stool consistency and fecal blood[28], demonstrated a positive linear relationship in the DSS mice (**Fig. 1e**), suggesting that lower body licking reflects the severity of colitis in mice. To measure colitis-induced mechanical allodynia, which is analogous to the feature of abdominal tenderness on clinical examination, we performed an up-down von Frey assay on the skin of the lower abdomen to calculate the

50% withdrawal threshold. We observed a profound decrease in the abdominal response
threshold during the DSS water treatment, which gradually recovered after DSS removal to
thresholds found in regular $H_2O$ regimen control mice (**Fig. 1f**). To examine the functional
relevance of the licking behavior during the 4-day DSS water regimen to pain, we used Nav1.8-
Cre mice that express either an inhibitory hM4Di receptor for chemogenetic inhibition, or
tdTomato as a control, in dorsal root ganglion nociceptor neurons that trigger nocifensive
reflexes in response to noxious stimuli (**Extended Data Fig. 3**). Mice where Nav1.8-lineage
nociceptors were temporarily silenced by activating hM4Di on receiving 0.1 mg/kg of
deschloroclozapine (DCZ)[31] once per day over the 4 days of DSS exposure, displayed a marked
reduction of nociceptive licking behavior and an attenuation of body weight loss (**Fig. 1g and**
**Extended Data Fig. 1a**). This analgesic-like effect persisted on D5 after the cessation of
Nav1.8 nociceptor silencing, but subsided by D7, when the behavior was indiscernible between
Nav1.8-tdTom and WT controls with DSS treatment. This reveals that the behavioral licking
signature tracks colitis severity and is triggered by Nav1.8-lineage nociceptors.
150
In parallel, we used an acute, mechanically induced visceral pain model by introducing graded
mechanical distension in the mouse distal colon (**Extended Data Fig. 4a,b**). To produce an
acute noxious colonic stimulus, a balloon was inserted and connected via a bifurcated tube to
an air-pressurized syringe and a manometer to enable controlled distension and pressure
monitoring. Acute colon distension at a pressure of 40 and 80 mmHg elicited pressure
intensity-dependent spontaneous behaviors, including abdominal licking, abdominal squashing
and hunching (**Extended Data Fig. 4c**), which were effectively diminished following Nav1.8-
nociceptor chemogenetic silencing by exposure to DCZ (**Extended Data Fig. 4d-g**).
159
Additionally, we performed a cFOS analysis in the $PBN_L$ of mice that received a colon balloon
placement either as a sham, or with a pressure distension of 80 mmHg to identify central
visceral pain-activated neurons. We found a substantial increase of cFOS-positive neurons in
$PBN_L$ following 30-min of stimuli and that the signal was significantly reduced by
chemogenetic silencing of the Nav1.8-lineage neurons (**Extended Data Fig. 4h,i**), suggesting
that peripheral Nav1.8$^+$ nociceptors are the active driver of visceral pain hypersensitivity and
trigger transmission of visceral pain signals into the central nervous system (CNS) to generate
adaptive behavioral responses.
168
169 **<u>Colitis drives a $PBN_L$ neural activation mediated by peripheral nociceptors and</u>**

 **inflammation**

Neurons in the mouse parabrachial nucleus are molecularly diverse with multiple subsets of

cell types that reside within distinct spatial niches[20-22]. We performed activity-dependent

mapping in PBN neurons using cFOS expression to determine the neural activation pattern

along the $PBN_L$, a critical hub that receives multiple afferent inputs from the periphery and

projects to the brain to generate sensory and affective responses. Following the 4-day DSS

colitis induction paradigm we observed a significant increase in cFOS-positive signals in the

$PBN_L$ region compared to regular $H_2O$ controls (**Fig. 2a-c**). To verify whether this activation

in the $PBN_L$ is driven by primary sensory neurons, we utilized the Nav1.8-hM4Di mice for a

temporary chemogenetic silencing of nociceptors, and TRPV1-DTA mice for a selective

diphtheria toxin-based permanent ablation (DTA) of peripheral nociceptors. We observed a

considerable reduction of the colitis-induced cFOS activation following either temporary or

permanent silencing of Nav1.8 and TRPV1-lineage neurons, respectively (**Fig. 2a-f**).

To selectively capture neurons activated during DSS-induced colitis, we employed an activity-

dependent genetic labeling strategy by crossing TRAP2 mice with Rosa-lxl-tdTomato reporter

mice. TRAP2-tdTom mice were subjected to 4-day DSS treatment followed by an

administration of 4-hydroxytamoxifen (4-OHT). This approach enabled permanent labeling of

FOS-expressing neurons that were active during the defined colitis window, allowing

identification and analysis of DSS colitis-activated neuronal populations (**Fig. 2g**). We

observed a comparable increase in the tdTomato-positive signal relative to baseline as the

cFOS staining, and this increased FOS-driven activation was noticeably diminished in mice

that received ketorolac injections, a nonsteroidal anti-inflammatory drug, to dampen the DSS-

induced local inflammation in the colon (**Fig. 2h-i**). These results indicate that the central

activation of $PBN_L$ neurons is driven by peripheral inflammation and nociceptor inputs.

**PBN$^{colitis}$ neurons tune abdominal sensitivity**

Next, we sought to probe what function the activated $PBN_L$ ensemble may have in mediating

colitis pain and its associated affective phenotypes. To gain functional access to the cFOS-

driven population, we utilized TRAP2 mice to functionally label the colitis-activated neural

population. The TRAP2 mice first received a stereotaxic injection of AAVs encoded with Cre-

dependent DREADDS bilaterally into $PBN_L$. Colitis-activated $PBN_L$ (PBN$^{colitis}$) neurons were

tagged by treating the mice with DSS water followed by an injection of 4-OHT to induce iCre

recombination specifically in the FOS-expressing population (**Fig. 2j**). Quantification of the

signal overlap between TRAPed and FOS-positive cells induced by DSS colitis revealed a

~60% TRAPing efficiency for the TRAPed and FOS-positive population (**Fig. 2k-m**). We then

asked whether PBN$^{colitis}$ neuronal activity is sufficient to modulate basal mechanical thresholds

in the abdominal region, a measure of the referred cutaneous hypersensitivity associated with

visceral pain (**Extended Data Fig. 5a**). We found that chemogenetic silencing (hM4Di) or

activation (hM3Dq) of the PBN$^{colitis}$ neurons bidirectionally tuned abdominal mechanical

sensitivity across a range of von Frey forces following DCZ administration when compared

with vehicle controls (**Extended Data Fig. 5b,d**).

When the same cohort of mice was assessed for basal hindpaw mechanical sensitivity, neither

silencing nor activation of PBN$^{colitis}$ neurons produced any observable change in the paw

withdrawal responses (**Extended Data Fig. 5c,e**). Hindpaw zymosan-activated PBN$_L$

(PBN$^{Zymosan}$) neurons modified the paw withdrawal response to von Frey stimuli (**Extended**

**Data Fig. 7a,b**) and thermal sensitivity and preference across a temperature gradient

(**Extended Data Fig. 7d-f**), but displayed a negligible modulatory effect on abdominal

sensitivity (**Extended Data Fig. 7c**). These results support a modality-specific role of the PBN$_L$

in visceral versus somatic nociceptive processing.

**PBN$^{colitis}$ neurons gate behavioral responses towards visceral nociception**

To evaluate the impact of PBN$^{colitis}$ neural activity in modulating colon pain-associated

behaviors, TRAP2 mice with Cre-dependent inhibitory hM4Di AAVs received a colonic

infusion of 0.05% capsaicin to activate TRPV1-lineage nociceptors in the distal colon to initiate

pain (**Fig. 2o**). We identified several distinct spontaneous behaviors, including lower body

licking, jumping and adnominal squashing immediately following the low-dose colonic

capsaicin exposure compared to vehicle treated controls (**Fig. 2n**). These spontaneous behavior

bouts were effectively reduced by silencing of PBN$^{colitis}$ neurons with DCZ, as opposed to the

increase in behavior bouts in vehicle treated mice (**Fig. 2p**). In parallel, we identified the

development of tactile allodynia over the abdominal region in the colonic capsaicin-treated

mice and that mechanical threshold returned close to baseline in response to PBN$^{colitis}$ neural

silencing (**Fig. 2q**).

**PBN$^{colitis}$ neurons promote aversive states**

To assess whether PBN$^{colitis}$ neurons influence affective and motivational valences we

implemented a three-chamber task where mice were allowed to freely explore a completely

shielded chamber ("hidden zone") or an open chamber ("open zone"). We then performed a

conditioning paradigm by pairing the injection of DCZ in mice in the hidden zone, and vehicle

in the open zone, in a set of mice that had previously received a stereotaxic delivery of Cre-

dependent hM3Dq AAV for chemogenetic activation, or mCherry AAV as controls (**Extended**

**Data Fig. 6a**). Real-time tracking of the mouse trajectory demonstrated that PBN[colitis]

activation drove an aversive response, as revealed by a robust reduction of the time spent in

the hidden zone when that chamber was paired with DCZ treatment during the conditioning

sessions (**Extended Data Fig. 6b,c**). In parallel, we also conducted a two-bottle choice test to

monitor motivational changes towards an appetitive incentive (**Extended Data Fig. 6d**) and

found that a strong taste aversion developed in mice after PBN[colitis] activation conditioning, as

indicated by significantly reduced consumption of sucrose when paired with DCZ treatment,

compared with the control group (**Extended Data Fig. 6e,f**).

**PBN$_L$ neurotensin neurons encode visceral pain induced by colonic insults**

Cell type-specific populations reside in different subregions of the PBN$_L$ and form distinct

projection patterns and circuit functions[12,21,32]. We divided colonic stimulation induced cFOS

expression in the PBN$_L$ region into two distinct subregions, the external lateral PBN (elPBN)

and the dorsal lateral PBN (dlPBN), based on established anatomical boundary standards[12,20,21].

We identified enriched cFOS-positive signals within the elPBN region following DSS-induced

colitis, which comprised ~67.5% of cFOS-positive signals within the elPBN compared to

~32.5% within the dlPBN (**Extended Data Fig. 8**), indicating preferential activation of neural

ensembles within elPBN under the condition of colitis. We then focused on the elPBN

subregion to identify the neuronal cell types recruited during colitis using an established spatial

transcriptomic dataset[20,21]. The elPBN region primarily encompasses a *Calca*-expressing

population along with other gene markers, including *Tac1*, *Penk*, *Crh and Nts*. To focus on

those elPBN neurons that send projections to sensory and pain-regulating brain regions, we

used *Crh* and *Nts* markers to evaluate signal overlap in our cFOS dataset. We collected PBN$_L$

from mice that had either received 3% DSS drinking water for 4 days to induce colitis or 20

μL of 5 mg/mL zymosan into the left hindpaw for 6 hours to induce plantar skin inflammation[33]

(**Fig. 3a**). We determined the signal overlap between the two marker genes and *Fos* expression

by performing RNAscope on serial sections of PBN$_L$ (**Fig. 3b**). *Nts$^+$* neurons displayed a ~65%

overlap with *Fos$^+$* signals in neurons during DSS colitis, with only a ~30% signal overlap with

*Crh$^+$* neurons, and a ~25% overlap between *Nts$^+$* and *Fos$^+$* signals in neurons following

hindpaw zymosan (**Fig. 3c**), suggesting a preferential role of neurotensin-expressing PBN$_L$

neurons in encoding visceral over somatic pain.

Using a neurotensin genetic driver of Cre mice (NT-Cre), we next sought to delineate the *in*
*vivo* calcium dynamics of the $PBN_L$ NT neurons using fiber photometry in response to the
visceral pain elicited by noxious distension with a pressurized colon balloon (**Fig. 3d-f**). We
inserted a balloon with a bifurcated ending to an air-pressured syringe and a manometer to
monitor balloon distension pressure within the colon simultaneously (**Fig. 3d**) and introduced
pressures of 20, 40 and 80 mmHg into the balloon to mechanically distend the colonic lumen
with a 5-sec inflation to the designated pressure and a 5-sec deflation back to the baseline
pressure. We observed an encoding of the $PBN_L$ NT neural population to colonic distension in
a pressure intensity-dependent manner (**Fig. 3g-i**), providing direct evidence for their role in
the central processing of visceral afferent insults.

**$PBN_L$ neurotensin neurons control visceral reflex and behavioral responses to colonic**
**insults**
We next sought to determine whether central NT-positive neurons play a functional role in
colitis-induced symptoms. We utilized two different AAV-based silencing strategies to explore
the impact of reduced $PBN_L$ NT activity on the aberrant behavior and altered GI function
triggered by DSS-colitis. We injected an AAV vector that encoded (1) Cre-dependent hM4Di
(2) Cre-dependent TeLC (tetanus toxin light chain) or (3) Cre-dependent GFP into the bilateral
$PBN_L$ of NT-Cre mice and evaluated behavioral outcomes over a 4-day DSS treatment regimen
(**Fig. 4a-c**). Temporary silencing elicited by M4 muscarinic receptor binding to the DREADD
agonist versus long-term silencing mediated by a tetanus toxin-based synaptic blockade were
implemented to compare their analgesic profiles on colitis-induced licking behavior (**Fig. 4d**).
We found a robust attenuation of nociceptive licking behavior in mice that received 0.1 mg/kg
of DCZ (once per day for 4 days) to silence $PBN_L$ NT activity during DSS treatment, and that
this analgesic effect persisted beyond the chemogenetic silencing period when compared to the
GFP control group (**Fig. 4e**). Long-term silencing of NT release in the $PBN_L$ by tetanus toxin
substantially reduced the colitis-induced licking behavior with only a marginal increase in
licking bouts at D7 (**Fig. 4e**).

Licking bouts in mice are modulated either by a local inflammation-induced sensitization
driven by Nav1.8-lineage nociceptors or through a body-axis projection that controls
gastrointestinal (GI) physiology[34,35]. To test whether central NT neural activity modulates GI

function, we delivered DCZ to NT-Cre mice that had received a Cre-dependent hM4Di or GFP

AAV in the $PBN_L$ during the DSS treatment, followed by an oral gavage of carmine dye at D7

to measure gastrointestinal transit time (**Fig. 4f**). In the NT-silenced group we found a

normalization of the colitis-accelerated gut motility, as revealed by cumulative stool number

over time and the latency to the first red fecal pellet expelled by the mice (**Fig. 4g,h**). We

measured local colon sensitivity in response to a series of controlled distensions using a close-

loop balloon barostat device in both the NT-silenced and control groups (**Fig. 4i**). Distension

intensity-dependent changes of the colon reflex were present in control mice, with a large

increase of the response when the distension pressure surpassed 60 mmHg. Importantly, these

evoked responses were largely dampened in NT-silenced mice (**Fig. 4j,k**). We conclude that

$PBN_L$ NT neurons serve as a central hub to both sense visceral hypersensitivity and impact

associated colonic reflex and behavioral outputs.

**$PBN_L$ neurotensin neurons protect against colitis-induced pathology**

To determine whether silencing $PBN_L$ NT neurons attenuate the histopathological outcomes of

DSS-induced inflammation, we administered a calibrated dose of DCZ into the mouse drinking

water to achieve a sustained silencing of $PBN_L$ NT neurons that matches the multi-day

progression of the pathology (**Extended Data Fig. 10a**). Longitudinal assessment of the stool-

based disease activity index (DAI) revealed that chronic NT-silencing significantly improved

stool consistency and fecal bleeding, and reduced DAI compared to a non-silenced control

group, as the disease progressed toward D7 (**Extended Data Fig. 10b**). Analysis of the colon

revealed that silencing $PBN_L$ NT neurons reduced neutrophil infiltration and preserved crypt

integrity and overall colon length (**Extended Data Fig. 10c-e**). Fecal lipocalin-2, an

inflammatory marker of mucosal immune activation, was significantly elevated at D7 under

DSS colitis but was markedly reduced in the NT-silencing group (**Extended Data Fig. 10h**).

These protective effects appeared to occur independently of overt systemic immune alterations,

as evidenced by a lack of weight change in spleen and mesenteric lymph nodes (**Extended

Data Fig. 10f,g**). These data indicate that $PBN_L$ NT neurons can exert a protective influence

against colitis-mediated pathology.

**$PBN_L$ neurotensin neurons exert a restricted modulation of somatic inflammatory pain**

We next investigated whether $PBN_L$ NT neurons differentially modulate distinct peripheral

sensory modalities. An intraplantar injection of zymosan was administered to induce

inflammatory pain in the hindpaw. Somatic hypersensitivity was analyzed by measuring

contact luminance of the hindpaws via a transillumination recording platform[30]. A reduction in
contact luminance is a proxy for tactile allodynia as zymosan-induced inflammation reduces
plantar surface contact (**Extended Data Fig. 9a,b**). Contrary to the effects observed in the
colitis model, silencing of $PBN_L$ NT neurons did not significantly attenuate the zymosan-
induced hypersensitivity throughout the three-day post-injection period (**Extended Data Fig.**
**9c**).

**<u>Blocking neurotensin signaling attenuates colitis-induced pain-related behavior</u>**
We next aimed to investigate the downstream brain targets that receive neural projections from
the $PBN_L$ NT neurons by conducting an AAV-based projection mapping, and identified several
regions known to regulate pain, including the bed nucleus of the stria terminalis (BNST),
central amygdala (CeA), ventral posteromedial (VM) thalamus and dorsolateral periaqueductal
gray (dlPAG) (**Extended Data Fig. 11a**). We then evaluated whether blockade of downstream
terminal NT release could provide any therapeutic relief of colitis-associated pain-related
behaviors. To confirm a functional connectivity between colitis-activated $PBN_L$ neurons and
the terminal release of NT in the CeA, we delivered a Cre-dependent hM3Dq into the $PBN_L$
region of TRAP2-Cre mice, followed by a 4-OHT injection during the 4-day DSS treatment to
induce the labeling of $PB^{colitis}$ by a FOS driver (**Extended Data Fig. 11b**). In the CeA, we
delivered a G protein–coupled receptor activation–based neurotensin sensor (GRAB-NT1.0)[36]
followed by an optic fiber implant to monitor NT release dynamics *in vivo* (**Extended Data**
**Fig. 11c**). We detected a robust surge in NT release in the CeA region after activation of $PB^{colitis}$
by DCZ treatment, as compared to the baseline signal in the vehicle treatment group (**Extended**
**Data Fig. 11d,e**), confirming NT signaling in the PBN-CeA pathway. To test whether blocking
neurotensin receptor type 1 (NTSR1) signaling in the CeA region affects DSS-colitis
behaviors, we implanted cannulas bilaterally in the CeA for selective NTSR1 antagonist,
SR48692, infusions (0.3 μL/day at 20 nM) over 4-day DSS treatment (**Extended Data Fig.**
**11f**). This produced an effective attenuation of colitis-induced licking behavior bouts as well
as a reduction in diarrhea, as evidenced by the normalization of the stool water content
(**Extended Data Fig. 11g-i**). These results highlight CNS neurotensin signaling as a key
modulator that calibrates colitis-induced visceral pain and associated gastrointestinal
pathological changes.

**Discussion**

Understanding the central neural modules that encode and regulate visceral sensation has been
hindered by the diffuse and heterogeneous nature of the peripheral innervation of viscera and
the absence of robust pain-related behavioral reflexes that allow direct functional
assessment[4,6,37]. We implemented a supervised machine-learning framework that integrates
pose estimation with body-part contact detection to extract subtle and disease-relevant
behaviors. We identified a distinct lower body licking behavior that emerged selectively in
mice with colitis. Notably, this behavioral signature closely tracked the temporal dynamics of
the colonic inflammation and was functionally dependent on Nav1.8-lineage nociceptors.

By combining FOS-based activity mapping with TRAP to identify activation-defined neuronal
populations, we found that DSS-induced colitis preferentially recruits neurons within the
external lateral PBN, a region implicated in encoding aversive salience and the affective
valuation of bodily threat[13,24,38]. Silencing this colitis-activated parabrachial ensemble
(PBN$^{colitis}$) markedly attenuated the behavioral responses to an acute intracolonic capsaicin
challenge. In contrast to somatosensory pain, visceral pain is commonly characterized by
emotional distress and is therefore thought to substantially engage limbic and affective
circuits[39]. Consistent with this, we found that activation of PBN$^{colitis}$ neurons robustly drives
aversive behavior as assessed by both conditioned place and conditioned taste paradigms,
indicating that this neural ensemble engaged during visceral inflammation promotes a negative
valence.

The enrichment of FOS-positive neurons within the elPBN in response to colitis prompted us
to define the molecular identity of the colitis-responsive cell types in this region. We
discovered that neurotensin-expressing neurons exhibited a selective overlap with colitis-
induced FOS activation, with minimal overlap with the FOS signals elicited by a zymosan-
induced somatic inflammation in the hindpaw, supporting preferential engagement of the PBN$_L$
NT population by visceral rather than somatic afferent inputs. Using *in vivo* fiber photometric
recordings, we further demonstrated that PBN$_L$ NT ensemble activity increased robustly in
response to graded colorectal distension when intraluminal distension exceeded 40 mmHg.
This threshold-dependent profile parallels the force-response characteristics observed in most
colon-innervating dorsal root ganglion neuron subtypes[40,41], supporting the notion that PBN$_L$
NT neurons encode high-threshold visceral nociceptive signals.

PBN$_L$ NT neurons constitute a distinct subset of the excitatory population within the elPBN region and partially overlap with *Calca*, *Oprm1* and *Tac1* markers, which have been implicated in pain modulation. We found that silencing the PBN$_L$ NT population, using either chemogenetics, or toxin-directed impairment of synaptic transmission, generated a profound analgesic effect, as evidenced by a marked reduction in colitis-associated licking behavior. Notably, this analgesic effect was more sustained than that achieved by targeting Nav1.8-lineage nociceptors, indicating that PBN$_L$ NT neurons operate at a higher-order integrative node in the visceral pain pathway. We reasoned that the improvement of the nocifensive behavior is contributed by a normalization of local hypersensitivity in the sensitized colon. Importantly, we identified a significant reduction in the colonic hyperreflex triggered by barostat distension with PBN$_L$ NT silencing, particularly at pressures exceeding the 40 mmHg threshold associated with visceral pain. In addition, PBN$_L$ NT silencing also restored gastrointestinal transit, emphasizing that this neuronal population integrates visceral sensory inputs to coordinate both pain-related behaviors and peripheral organ-level physiological responses.

Neurotensin has been implicated in the regulation of fundamental homeostatic processes, including feeding, social valence and pain[42-44], and has also been characterized as a peripheral proinflammatory mediator, as evidenced by an elevated expression of its receptor, NTSR1 in the colon in experimental colitis models[45-47]. We hypothesized that central neurotensin signaling contributes to the visceral nociceptive and autonomic sequelae of colitis. Using AAV-based circuit tracing, we revealed that PBN$_L$ NT neurons project to affective brain regions, with a particularly dense innervation of the central amygdala (CeA). Activation of this parabrachial source evoked a robust surge in NT release in the CeA, as detected by a neurotensin-based GRAB sensor. Importantly, pharmacological blockade of NTSR1 signaling within the CeA effectively ameliorated colitis-associated licking behavior and diarrhea, indicating that central NT signaling acts as a supraspinal modulator of disease severity, potentially through the amplification of those aversive and autonomic pathways that feed back onto peripheral visceral inflammatory and nociceptive processes[34,35,48]. By integrating machine learning-assisted behavioral decoding, molecular profiling, *in vivo* neural activity recording, and circuit-specific manipulations, our study provides evidence for a central circuit mediated modulation of peripheral visceral function. These findings offer a mechanistic framework for understanding those visceral disorders arising from dysregulated brain-body interactions, and

pinpoint neurotensin in the CNS as a potential therapeutic target that can be leveraged to treat

colitis-induced visceral pain.

**Materials and Methods**

<u>**Animals**</u>

All animal husbandry and procedures involving mice were conducted in strict accordance with

guidelines set forth by the Boston Children's Hospital Institutional Animal Use and Care

Committee. Mice at 6-8 weeks of age with the strains of C57BL/6J (B6, JAX #000664), $Fos^{2A\text{-}iCreER}$

(TRAP2, JAX #030323), Nav1.8-Cre (JAX #036564), NT-Cre (JAX #017525), R26-

LSL-hM4Di/mCitrine (JAX #026219), Ai14(RCL-tdTomato)-D (JAX #007914), TRPV1-Cre

(JAX #017769) and ROSA-DTA (JAX #009669) were obtained from the Jackson Laboratory.

Adult mice of 8-12 weeks of age for both male and female mice of heterozygous alleles were

used for this study. Induction of iCre-mediated recombination in the TRAP2 mice was

performed by an i.p. injection of 0.1 mL of 10 mg/mL of 4-OHT dissolved in corn oil, with

experiments beginning ~2 weeks after the induction.

<u>**Pharmacological reagents**</u>

Dextran sulfate sodium salt (MP Biomedicals #0216011080) was dissolved in regular drinking

water at a concentration of 3% to induce colitis, 4-hydroxytamoxifen (Sigma-Aldrich #H6278)

was prepared in corn oil and ethanol for the induction of $Cre^{ER}$ translocation,

deschloroclozapine (Hello Bio #HB9126) was prepared in saline solution as an agonist for

DREADDs experiments, ketorolac (TargetMol #T1212) was prepared in saline, capsaicin

(Tocris #0462) was dissolved in ethanol, tween80 and saline to induce acute nociceptive

responses, zymosan A from saccharomyces cerevisiae (Sigma-Aldrich #Z4250) was prepared

in saline solution for intraplantar injection to induce hindpaw inflammation and pain. SR48692

(Sigma-Aldrich #SML0278), a selective NTSR1 antagonist, was prepared at 20 nM in a

solution of saline with 2% dimethylsulfoxide (DMSO). Carmine powder (Sigma-Aldrich

#C1022) was prepared at 6% w/v with methylcellulose to measure gut transit time.

<u>**Adeno-associated virus (AAV) reagents**</u>

For chemogenetic manipulation of $PBN_L$ neurons the following AAV vectors were used:

AAV2/1-hSyn-DIO-hM4Di-mCherry (BCH Viral Core; $3.51×10^{13}$ gc/mL) for silencing,

AAV2/1-hSyn-DIO-hM3Dq-mCherry (BCH Viral Core; $2×10^{13}$ gc/mL) for activation and

AAV2/1-hSyn-DIO-mCherry (BCH Viral Core; $3.48×10^{13}$ gc/mL) as fluorescent controls. For

permanent silencing of $PBN_L$ neurons AAV2/8-hSyn-FLEX-TeLC-GFP (BCH Viral Core;

$1.56 \times 10^{13}$ gc/mL) was used to block synaptic transmission. For fiber photometric recording

AAV2/8-hSyn-FLEX-jGCaMP8s-WPRE (BCH Viral Core; $4.16 \times 10^{13}$ gc/mL) was used to

monitor neural calcium activity and AAV2/1-hSyn-FLEX-H2b-GFP (BCH Viral Core; $3 \times 10^{13}$

gc/mL) for fluorescent controls. To detect neurotensin release AAV2/1-hSyn-GRAB-NT1.0

(Addgene plasmid #180006 from Y. Li, viral vector packaged by BCH Viral Core; $2.78 \times 10^{13}$

gc/mL) was used.

**Stereotaxic injections and surgical procedures**

*Intracranial AAV vector injections*

Intracranial injections were conducted with aseptic procedures on a digital stereotaxic platform

for small animals. Mice were anesthetized with 4% isoflurane and maintained with ~1.5%

throughout the entire surgery on a feedback-controlled heating pad at a body temperature of

~37°C. We injected AAVs using a 33 gauge, 5 μL microsyringe (Hamilton #65460-03) at a

rate of 50 nL/min for 250 nL volume in one injection site. $PBN_L$ was targeted at coordinates

relative to bregma:  AP= –5.20 mm, ML= ±1.43 mm, DV= –3.55 mm. CeA was targeted at

coordinates: AP= 1.15 mm, ML= ±2.83 mm, DV= −4.17 mm. Mice were allowed to recover

and waited for at least 3 weeks for the virus to express before experiments were conducted.

*Optic fiber and cannula implantation*

The optic fiber was placed ~0.3 mm above the AAV injection site to monitor bulk calcium

activity using fiber photometric recording. For pharmacological experiments, cannulae (22GA,

PlasticsOne) were implanted bilaterally above the targeted brain region for repeated infusions

of drug into the mouse brain. All implants were tightly anchored onto the skull and secured

with Metabond (Parkell) and dental cement. Implanted mice were allowed to recover at least 2

weeks before experiments.

**Behavior assays**

*Spontaneous behavior recording*

Mice were individually placed in a $18 \times 18 \times 15$ cm (length x width x height) black acrylic

chamber closed on all sides (BlackBox Bio) except for an NIR camera (Basler acA2000-

50gmNIR GigE, Basler AG, Ahrensburg, Germany) was positioned 30 cm beneath the glass

surface to track whole body movement from a bottom-up perspective[30]. Image brightness

features were captured using LED light reflected changes in pixel intensity acquired at a frame

rate of 25 Hz with $1000 \times 1000$ pixels dimensions. Mice were habituated in the recording

chamber 1 hour per day for three days before the start of the experiment. Recordings were
performed in both experiment and control groups on the same day for 30 min in each session
during the animal's light cycle.

<u>Automated behavior screening</u>

Lower body licking behavior was quantified using BAREfoot, a supervised machine learning
framework that integrates pose estimation with light-based measurements of body-part
pressure and spatial distance[29]. Training datasets for the lower body licking classifier were
generated through manual annotation of video recordings by experienced scorers, after which
new classifiers were trained on these labeled data. After testing and validating classifier
performance, automated detection and quantification of lower body licking behaviors bouts
was performed. Manual annotation of spontaneous behavior signatures, including peri-anal
licking (repetitive licking around the anal port), abdomen licking (repetitive licking on a fixed
region of the lower abdomen), abdomen squashing (contact with pressing of the lower abdomen
against the floor), hunching (shortening the distance between fore and hindlimbs) and jumping
(leaping off the floor with two hindlimbs), were labeled by three experienced independent
investigators on a frame-by-frame level using a customized MATLAB scoring platform.

*Paw luminance measurement*

Hind paw coordinates were tracked using DeepLabCut[49], and 23 × 23 squared pixels centered
on these locations were extracted from the FTIR frames. Tactile allodynia was assessed by
measuring paw luminance, defined as the average pixel brightness within each extracted ROI.

*Abdominal and plantar von Frey assay*

Mice were habituated in von Frey chambers on a meshed floor for three days (1 hour per day),
including the day of the test, before the mechanical sensitivity was evaluated. To evaluate the
reflexive sensitivity in response to mechanical stimuli, a logarithmically increasing set of 8 von
Frey filaments (Stoelting Co, #58011) ranging from 0.02 to 4.0 grams were applied
perpendicularly to the plantar hindpaw or lower abdomen with a sufficient force to gently bend
the monofilament. We employed the up-down method to measure the 50% withdrawal
threshold on the mice lower abdomen or plantar hindpaw[50]. The response frequency was
measured as the number of positive rapid withdrawal of the lower abdominal region or the
hindpaw in response to 10 von Frey stimulations at an interval of at least 30 sec between each
stimulus. All the behavior tests were conducted in a blinded fashion to either the experimental
vs control group, or the injection of treatment vs vehicle group.

*Colorectal distension*

Mice were briefly anesthetized with ~1.5% of isoflurane to receive a lubricated polyethylene
balloon insertion through the rectal port as previously described[51]. A colonic balloon with a
diameter of ~2 mm was gently placed at the distal colon segment ~20 mm from the mouse anal
verge and connected to a pressure-equalization tubing with a manometer (Cole Parmer)
positioned at the bifurcation to simultaneously monitor distension pressure. A pressured tape
was attached to the base of the tail to secure the placement of the colonic balloon for freely
behavior recording. For each distension trial, the balloon was inflated to the designated
pressure of 40 mmHg or 80 mm Hg within 5 seconds and a 5-sec deflation back to the baseline
pressure for a total of 30-sec distension. Each behavioral recording consisted of alternation of
30-second distension and 30-second rest, continuously over a 30-minute recording session.

*Conditioned motivation tests*

To evaluate the impact of colitis-activated $PBN_L$ neurons on motivational behaviors in mice,
we conducted conditioned place motivation and conditioned taste motivation tasks. For
conditioned place motivation, mice were habituated for 1 hour in a three-chamber apparatus
separated with a central corridor that allowed free access to left or right compartment, including
one shielded compartment with a meshed floor (hidden zone) and the other open compartment
with a smooth acrylic floor (open zone). On day 1, mice were placed in the central corridor to
freely explore both compartments for 30 min to determine the place preference by tracking the
time spent across the entire apparatus using Ethovision XT (Noldus). On conditioning day 2
and 3, mice were confined to each compartment and paired with an i.p. injection of 0.1 mg/kg
DCZ in the hidden zone, or vehicle in the open zone for 1 hour. The conditioning occurred
twice a day (9 AM and 5 PM) with alternating compartments for a period of 2 days. On day 4,
mice were re-introduced back into the central corridor to freely explore for 30 min to record
the time spent in each compartment. For conditioned taste motivation, mice were single-housed
and habituated to the home cages and water bottles for three days before the experiment. Prior
to each day of conditioning, mice were restricted to water access overnight and the
consumption amount was measured during the light cycle (9 AM to 5 PM). On day 1 and day
3, mice were allowed to have free access to regular drinking water in the home cage for 30
min, followed by an i.p. injection of vehicle, then returned to the home cage until 8 hours were
timed to measure the consumption amount of solution. On day 2 and day 4, mice were given
free access to a 10% sucrose solution in the water bottle for 30 min, followed by an i.p. injection
of 0.1 mg/kg DCZ, then returned to the home cage until 8 hours were timed. On day 5, mice
were given two bottles that contained regular water and 10% sucrose in each, for 8 hours to

measure the taste preference. The preference index of sucrose over water during the two-bottle

test on day 5: (amount of sucrose consumption) / (amount of water + sucrose consumption).

*Thermal gradient ring (TGR)*

Mice were habituated for 30 min in the thermal gradient ring (Ugo Basile #35550) with the

platform temperature pre-set to 10°C on one side and 55°C on the opposite side at an increment

of 5°C along the circular ring. During the test, each mouse was allowed to freely explore along

the TGR setup for 1 hour and the occupancy duration as well as distance traveled in each

temperature zone were recorded using the ANY MAZE software.

*Hotplate assay*

Thermal nociceptive thresholds were assessed using a hotplate assay. Mice were first

acclimated to the device for 1 hour before the experiment. The hotplate (BIOSEB) was preset

and maintained at a constant temperature of 52°C degree. Each mouse was placed within a

transparent plexiglas cylinder on the heated surface, and the latency to exhibit the first

nocifensive flinching behavior was recorded. A maximum cut-off time of 30 seconds was

implemented to prevent plantar damage. Each mouse was tested in three trials with at least 10-

minute between trials and the average latency is calculated.

*Gastrointestinal (GI) transit assay*

To assess total gastrointestinal transit time, mice were orally gavaged 300 μL of 6% carmine

red powder dissolved in 0.5% methylcellulose as previously described method[52]. Mice were

then placed individually in clean home cages to measure the number of expelled stools. The

GI transit time was defined as the interval between oral gavage and the first observation of a

carmine-colored fecal pellet with the observation conducted every 10 minutes to ensure

temporal accuracy.

**Physiological recordings**

*In vivo fiber photometry recording in awake behaving mice*

Optical fibers (400 μm core diameter, RWD) were implanted two weeks following Cre-

dependent AAV delivery of GCaMP8s or GRAB-NT1.0. During implantation, real-time

photometric signals were monitored to optimize fiber placement depth to maximize

fluorescence signal intensity. Photometric recordings were acquired using a fiber photometry

system (Doric Lenses) controlled by Doric Neuroscience Studio software. Excitation light at

405 nm (isosbestic, activity-independent control) and 465 nm (activity-dependent signal) was

delivered, and emitted fluorescence signals were recorded at a sampling rate of 120 Hz. To

enable temporal alignment with behavioral recordings, the fiber photometry system generated

a synchronization light pulse every 90 s, which was captured concurrently by a behavior video

camera operating at 30 Hz. Raw photometry signals were processed by averaging every 12

consecutive samples, yielding an effective sampling rate of 10 Hz. To directly compare neural

activity and behavior response to stimulus onset, fiber photometry and video-derived

behavioral signals were resampled onto a shared time base. Photometry data were exported to

MATLAB (MathWorks) for further processing and analysis. Fluorescence signals were

expressed as $\Delta F/F$, calculated as $\Delta F/F = (F - F_0)/F_0$, where F represents the 465 nm activity-

dependent signal and $F_0$ denotes the fitted isosbestic control signal derived from the 405 nm

channel. Z-scored $\Delta F/F$ values were used for synchronization and subsequent analyses.

*Distension barostat recording*

For assessment of colonic sensitivity, colorectal distension was performed using a close-loop,

programmed balloon distension barostat (G&J Electronics Inc). Mice were briefly anesthetized

with 3% isoflurane followed by an i.p. injection of either vehicle or DCZ. Mice were then

loosely positioned in a cylindrical restrainer, and a lubricated flexible balloon catheter was

gently guided into the anus and advanced into the distal colon. The catheter was secured by

taping it to the tail and the restrainer was tightened to minimize movement. Mice were allowed

to recover fully from anesthesia and habituate in the restrainer for 30 min prior to recording

experiment. The barostat operated through a closed-loop feedback system and utilized an

internal pressure transducer to monitor intracolonic pressure change in real-time. The

distension protocol is consisted of balloon inflation to graded pressures of 20, 40, 60, and 80

mmHg for 10 sec per distension pressure, with a 3-min inter-stimulus interval between

distensions. Each pressure series was repeated three times. The barostat control signal

indicating balloon distension was synchronized with the intracolonic pressure recordings

obtained from the probe sensor to define distension timing and pressure levels. All raw data

were acquired and processed using LabChart software (ADInstruments). To isolate dynamic

pressure fluctuations, the intracolonic pressure signal was processed offline using a high-pass

Butterworth filter (cutoff frequency, 0.1 Hz) to remove low-frequency background drift. The

root mean square (RMS) amplitude of the filtered signal was calculated within predefined

analysis windows and stored as a derived signal. For each distension epoch, the area under

curve (AUC) of the RMS signal was calculated relative to the pre-distension baseline and used

as an integrated measure of intracolonic pressure activity. Identical filtering parameters and

analysis windows were applied across all experimental sessions.

**Colon histopathology and stool composition analyses**

*Colitis severity evaluation*

Colon tissues were prepared as swiss-rolls, fixed, and sectioned for Hematoxylin & Eosin (H&E) staining to assess the severity of colonic inflammation. Colitis severity was assessed using a semi-quantitative scoring system based on three parameters: crypt dropout, submucosal expansion, and neutrophil infiltration. Each parameter was scored on a scale from 0 to 3, where 0 represented absence of pathology and 3 indicated severe involvement. The scores from all three parameters were summed to yield a total histology score ranging from 0 to 9, with higher scores reflecting greater tissue damage and inflammatory severity. Disease activity was further assessed using a stool-based disease activity index (DAI), which was calculated as the sum of two components: stool consistency and fecal blood. Stool consistency was scored as follows: formed and soft=1, loose or semi-formed=2, diarrhea=3. Fecal blood was evaluated using a Hemoccult card (Beckman Coulter) to detect occult blood and scored as follows: occult blood=1, visible blood=2, rectal bleeding=3. For stool composition measurements, mice were transferred to clean cages and allowed free access to food and water. Stool output was continuously collected for 1 h. The total mass of fecal material collected during this period was recorded as the wet weight per mouse. Collected fecal pellets were then air-dried for 14 days until a stable mass was achieved, at which point the dry weight was recorded. Percentage of fecal water content = [(wet−dry)/wet]x100.

*Fecal lipocalin-2 (LCN2/NGAL) quantification by ELISA*

Fecal lipocalin-2 levels were quantified using a sandwich ELISA (DuoSet™ Mouse Lipocalin-2/NGAL, R&D Systems/Bio-Techne; DY1857) according to the manufacturer's instructions. Frozen fecal pellets were weighed and resuspended at 100 mg/mL in PBS containing 0.1% Tween-20, vigorously homogenized by vortexing, and clarified by centrifugation at maximum speed for 1 min. The resulting supernatant was collected, stored at −80 °C, and subsequently used for ELISA analysis.

## **Immunohistochemistry**

Mice were anesthetized with an i.p. injection of 200 mg/kg pentobarbital and perfused transcardially with an ice-cold 1xPBS, followed by 4% paraformaldehyde (PFA) in 1xPBS. The brain, DRG and colon tissues were post-fixed for 4-6 hrs at 4°C and transferred to 30% sucrose in 1xPBS for 2 days. Tissues were frozen in O.C.T. (Tissue-Tek) and sectioned with a cryostat at 40-μm thickness. Sliced tissues were mounted on histology slides and immersed in blocking solution containing 5% normal donkey or goat serum and 0.3% Triton X-100 in 1xPBS for 1 hr at 4°C. The tissue slides were then incubated with primary antibodies at 4°C

overnight. On the next day, tissues underwent extensive wash with clean 1xPBS followed by
incubation with appropriate secondary antibody conjugated with AlexaFluor for 2 hrs at room
temperature. After washing with 1xPBS for 5 min three times, stained slides were coverslipped
with ProLong Diamond Antifade Mountant (Thermo Fisher, P36970). Histological images
were acquired using either a Leica SP8 or Zeiss LSM 980 confocal microscope at a resolution
of $1024 \times 1024$ pixels, using 10× or 20× air objectives with z-stack step sizes of 2–5 μm.
Immunofluorescence signals were quantified using ImageJ (NIH, v1.53f). For each histological
section, 8-10 consecutive z-slices were acquired and combined using an average-intensity z-
projection. Projected images were background-subtracted and thresholded to generate binary
masks of fluorescent signals. Automated particle/cell counting was performed to quantify the
number of detected fluorescent cells. The following primary antibodies were used: anti-cFOS
(Cell Signaling, #2250), anti-RFP (Rockland #48710), anti-Nav1.8 (Alamone Labs #ASC028),
anti-CD45 (eBioscience #14-0451-82), anti-GFP (Thermo Fisher #A10262); secondary
antibodies were used: Alexa Fluor$^{TM}$ 555-conjugated donkey anti-rabbit (Invitrogen #A31572),
Alexa Fluor$^{TM}$ 594-conjugated goat anti-rabbit (Invitrogen #A11012), Alexa Fluor$^{TM}$ 488-
conjugated goat anti-mouse (Invitrogen #A11001), Alexa Fluor$^{TM}$ 488-conjugated goat anti-
chicken (Invitrogen #A11039).
692

**<u>Fluorescence in situ hybridization</u>**
Mice underwent transcardial perfusion with 4% PFA and brains were post-fixed overnight and
30% sucrose for 48 hrs. Tissues were fresh-frozen in O.C.T. compound (Thermo Fisher
Scientific) and stored at −80 °C until cryoslicing. Coronal sections of 20 μm thickness covering
the entire PBN region were serially collected and mounted onto Superfrost Plus slides (Thermo
Fisher Scientific) and stored at −80 °C for processing. RNAscope in situ hybridization was
performed using the RNAscope Fluorescent Multiplex V2 Kit (Advanced Cell Diagnostics)
following the manufacturer's instructions. Slides were dehydrated through graded ethanol
(50%, 70%, and 100%) and treated with hydrogen peroxide for 10 min, washed, and incubated
with Protease IV for 30 min at room temperature. The following probes from Advanced Cell
Diagnostics were used: Mm-Fos (316921), Mm-Nts (420441) and Mm-Crh (316091). Sections
were incubated for probe hybridization and signal amplification steps in a HybEZ oven (ACD
Bio) at 40 °C. Fluorescent signal detection was achieved using TSA Plus HRP reagents and
Opal 520, 570, and 620 dyes (Akoya Biosciences). HRP reagents were applied for 15 min and
Opal dyes for 30 min at 40 °C with 15min HRP blocking steps between each channel and wash
buffer rinses between all incubations. FISH images covering serial sections of the PBN$_L$

between bregma −5.0 mm and −5.4 mm were analyzed using ImageJ. Image brightness and contrast were adjusted using consistent thresholding parameters to optimize visualization of each fluorophore. Cellular boundaries were defined based on DAPI labeling, and individual mRNA puncta (5–10 pixels in diameter) were quantified within each cell. Cells containing >5 puncta of Fos, Nts, or Crh mRNA were classified as marker-positive. This threshold was applied consistently across all analyzed sections to determine the overlapping of gene-expressing populations.

### Statistical analysis

All statistical analyses were performed using GraphPad Prism. The specific statistical tests applied, exact p values, and sample sizes (n) are reported in the corresponding figures and figure legends. Data distribution was assessed where appropriate, and statistical tests were selected accordingly. Unless otherwise stated, all statistical tests were two-tailed, and statistical significance was defined as $p < 0.05$.

### Acknowledgments

We thank the members of the Woolf laboratory for inputs and discussions. We thank the IDDRC Animal Behavior and Physiology Core, funded by NIH/NICHD P50HD105351 and Cellular Imaging Core, funded by NIH P50 HD105351 and Viral Core, funded by NIH 5P30EY012196 at Boston Children's Hospital; Dr. Mark L. Andermann for helpful comments and Dr. Fan Wang for general support. This study is supported by the National Institutes of Health grants R01AT011447 and R35NS105076 (C.J.W.), R01DK135707 and R01DK130836 (M.R), and the William Randolph Hearst Fund (Y.T.C). Schematics and illustrations were adapted from BioRender.com.

### Author Contributions

Y.T.C and C.J.W. conceived and designed the study. Y.T.C performed all the experiments and analyzed the data, with assistance from N.M.B. (AAV injections and histology), Y.J. (behavior scoring and RNAscope), J.R.R. (behavior scoring and histology), M.D.S. (GI transit assay and colon histopathology), P.E.M. (barostat recording), Y.L. (fiber photometry), O.B. (automatic behavior screening), K.H. (histology), B.Z. (behavior scoring) and B.L.T. (behavior assays). M.R. provided resources and oversight of the colon histopathology, GI transit and barostat recording experiments. Y.T.C and C.J.W. prepared the figures and wrote the manuscript with inputs from all the authors.

**Disclosures**

C.J.W. is a founder of Nocion Therapeutics, Quralis, and BlackBox Bio, and serves on the scientific advisory boards of Lundbeck Pharma, Axonis, Niroda, Mimetic Medicine, Tenvie Therapeutics, and Tafalgie Therapeutics. M.R. collaborates with Fzata and her spouse is a Novo Nordisk employee.

**Fig 1. Nocifensive behaviors elicited in response to DSS-induced colitis.** (a) Schematic of

experimental timeline to evaluate spontaneous behavior and abdominal sensitivity in mice

treated with 4-day of 3% dextran sodium sulfate (DSS) in their drinking water. (b) Histological

verification of immune cell cluster ($CD45^+$, green) invasion into the lamina propria of the

mouse colon 3 days post 4-day DSS treatment (right) compared with baseline (left). Scale bar,

20 mm. (c) Quantification of $CD45^+$ clusters at D0 (n=22 from 4 mice), D4 (n=14 from 3 mice),

D7 (n=22 from 4 mice) and D14 (n=15 from 3 mice). One-way ANOVA with Dunnett's

multiple comparisons. (d) Example of a mouse with DSS-colitis engaged lower body licking

behavior. (e) Linear regression analysis between the stool-based disease activity index and time

spent licking over a 30-min recording session in mice exposed to DSS (n=25 mice). Light

shaded areas represent 95% confidence band. (f) 50% cutaneous mechanical threshold of the

mouse lower abdomen using an up-down von Frey test ($H_2O$ n=9 mice, DSS n=9 mice). Light

blue shade: 3% DSS drinking water from D1-D4. (g) Automatic licking behavior quantified

using a supervised machine learning classifier in WT mice with $H_2O$ (n=3), WT mice with 4-

day DSS (n=6), Nav1.8-tdTom mice with 4-day DSS (n=3-6) and Nav1.8-hM4Di mice with 4-

day DSS (n=6). In two Nav1.8 groups, the chemogenetic agonist deschloroclozapine (DCZ,

0.1 mg/kg) was injected i.p. at D1, 2, 3 and D4, once per day (two-way ANOVA with Tukey's

multiple comparisons).

**Fig 2. PBN<sup>colitis</sup> neurons modulate behavioral responses to nociception in the colon.** (a)
Experimental scheme to measure FOS activation in the lateral parabrachial nucleus (PBN$_L$) in
WT and Nav1.8-hM4Di mice with 3% DSS drinking water. (b) Histological images of FOS
antibody staining in PBN$_L$ region at baseline, with exposure to 3% DSS and after Nav1.8
chemogenetic silencing. scp, superior cerebellar peduncle. Scale bar, 200 μm. (c)
Quantification of FOS-positive signals (baseline n=3 mice, DSS n=6 mice, Nav1.8-hM4Di n=4
mice, one-way ANOVA with Tukey's multiple comparisons). (d) Experimental scheme to
measure PBN$_L$ FOS activation in TRPV1$^{Cre}$(-)-DTA and TRPV1$^{Cre}$(+)-DTA mice with DSS.
(e) Histological images of FOS staining in PBN$_L$ region of TRPV1$^{Cre}$(-)-DTA and
TRPV1$^{Cre}$(+)-DTA mice with DSS. Scale bar, 200 μm. (f) Quantification of FOS-positive
neurons (TRPV1$^{Cre}$(-) n=2 mice, TRPV1$^{Cre}$(+) n=3 mice, two-tailed unpaired t-test). (g)

Experimental scheme to quantify $PBN_L$ FOS activation using TRAP2-tdTom mice under DSS treatment with i.p. injections of vehicle or 5 mg/kg ketorolac from D1-D4. (h) Histological images of tdTom$^+$ cells in $PBN_L$ region at baseline, 3% DSS with vehicle injections and 3% DSS with ketorolac injections. Scale bar, 200 μm. (i) Quantification of tdTom$^+$ cells (baseline n=5 mice, DSS+vehicle n=7 mice, DSS+ketorolac n=7 mice, one-way ANOVA with Tukey's multiple comparisons). (j) Experimental strategy to functionally label DSS colitis-activated parabrachial neurons (PBN$^{colitis}$) using TRAP2-Cre$^{ER}$ mice injected with Cre-dependent AAV vectors encoding chemogenetic actuators (hM4Di or hM3Dq) bilaterally into $PBN_L$ followed by an i.p. injection of 4-hydroxytamoxifen (4-OHT, 10 mg/kg) to drive expression of FOS-positive cells after 4-day of 3% DSS treatment. (k) Histological verification of signal overlap between TRAPed PBN$^{colitis}$ cells (magenta) and FOS RNAscope (green) under DSS colitis. Scale bar, 200 μm. White arrowheads indicate overlapping signals. (l) Pie chart depicting overlap between TRAP$^+$ and FOS$^+$ neurons. (m) Quantification of signal overlaps in TRAP$^+$ versus FOS$^+$ population (n=5 mice). (n) Quantification of spontaneous pain-related behaviors in response to an intracolonic 0.05% capsaicin infusion over a 60-min recording session (vehicle n=4 mice, capsaicin n=5 mice, multiple unpaired t-tests using two-stage step-up method (Benjamini, Krieger and Yekutieli)). (o) Experimental timeline to evaluate the effect of silencing PBN$^{colitis}$ neurons on capsaicin-induced spontaneous and evoked nocifensive responses. (p) Percentage time difference of capsaicin-induced abdominal licking, peri-anal licking and jumping after vehicle or 0.1 mg/kg DCZ to silence PBN$^{colitis}$ activity in a 30-min recording session (vehicle n=4 mice, DCZ n=5-7 mice). (q) Cutaneous threshold of mouse lower abdomen using up-down von Frey test (vehicle group n=4 mice, DCZ group n=4 mice, one-way ANOVA with Tukey's multiple comparisons).

**Fig 3. PBN_L neurotensin neurons encode colon-derived nociceptive signals.** (a) Schematic
of colitis-induced visceral pain in mice exposed to 4-day of 3% DSS drinking water (top) and
somatosensory inflammatory pain induced by an injection of 5 mg/mL zymosan into plantar
surface of the left hindpaw (bottom). (b) PBN_L sections with RNAscope detecting DSS colitis-
activated FOS and neurotensin (NT) (left), DSS FOS and corticotropin-releasing hormone
(CRH) (middle), and contralateral PBN_L sections with left hindpaw zymosan-activated FOS
and NT (right). Scale bar, 200 μm. (c) Quantification of signal overlap between FOS DSS and
NT (red bar), FOS DSS and CRH (orange bar), FOS Zymosan and NT marker (green bar). N
= 17 sections from 4 mice, 13 sections from 4 mice and 24 sections from 5 mice. (d) Schematic
of fiber photometric recordings of neural activity in PBN_L NT neurons evoked by intracolonic
balloon distension. (e) Histological verification of AAV-transfected jGCaMP8s cells and fiber
implant track in PBN of an NT-Cre mouse. Scale bar, 300 μm. (f) Example z-scored ΔF/F of
GCaMP8s signal in NT neurons in response to colorectal distension (CRD) of 80 mmHg. (g)
Heatmap responses of PBN_L NT neural activity to 10-sec CRD at 20, 40, 80 mmHg and GFP
controls in response to CRD at 80 mmHg (GFP n=30 trials from 3 mice, CRD 20 n=22, CRD
40 n=24, CRD 80 n=40 trials from 5 mice). (h) CRD-triggered average z-scored ΔF/F of NT
neurons. Light shaded areas represent s.e.m. (i) Area under curve (AUC) measured using

867 average signal during the 5 sec prior to CRD onset as baseline, from (h). One-way ANOVA

868 with Tukey's multiple comparisons.

**Fig 4. PBN_L neurotensin neurons control behavior and visceral sensitivity induced by colonic insults.** (a) Schematic of two different silencing (hM4Di: modified human M4 muscarinic receptor for chemogenetic silencing, TeLC: tetanus toxin light chain-mediated synaptic impairment) and a control AAV GFP injection to measure causal function of PBN_L NT neurons. (b) Histological verification of Cre-dependent AAV injections into bilateral PBN_L of an NT-Cre mouse, scp, superior cerebellar peduncle. Scale bar, 200 μm. (c) Quantification of mCherry⁺ cells in the PBN_L at A-P axis 5.0, 5.2 and 5.4 mm relative to bregma (n=3 mice). (d) Diagram illustrating two different silencing strategies (1) virally mediated Gi-coupled muscarinic receptors induce an efflux of potassium ions for temporary neural hyperpolarization (2) virally mediated tetanus toxin light chain blocks a presynaptic fusion protein for long-term

abrogation of neurotransmitter release. (e) Licking behavior quantified using a supervised machine-learning classifier in NT-Cre mice with bilateral injections of a Cre-dependent AAV that encodes GFP (n=7 mice), or hM4Di (n=7 mice) and TeLC (n=3 mice) in mice subjected to 4-day 3% DSS treatment. In both the GFP and hM4Di groups, the chemogenetic agonist (DCZ; 0.1 mg kg$^{-1}$, i.p.) was injected once daily on D1-D4. Two-way ANOVA with Dunnett's multiple comparisons. (f) Experimental scheme to evaluate gastrointestinal transit using a carmine-red dye assay to measure latency of the first appearance of dye-colored stool following oral administration on D7. GFP and hM4Di mice underwent 4-day DSS treatment and received DCZ injection once per day from D1-D4. (g) Quantification of total stool number 30 min after an oral gavage of carmine dye followed by stool count every 10-min until 400 min (GFP n=5 mice, hM4Di n=7 mice). Light shaded areas represent s.e.m. (h) Quantification of latency to first carmine-dyed stool (GFP n=5 mice, hM4Di n=7 mice. Two-tailed unpaired t-test). (i) Schematic of colonic reflex response to graded distensions by an inserted balloon connected to a close-loop, calibrated barostat system. (j) CRD-triggered average intracolonic pressure responses to distension pressures of 20, 40, 60 and 80 mmHg in GFP (red, n=8 trials from 4 mice) or hM4Di (green, n=14 trials from 7 mice) mice. Light shaded areas represent s.e.m. (k) Quantification of normalized AUC (average of 5-sec AUC following CRD onset / average of 5-sec AUC prior to CRD onset) in response to CRD pressure at 20, 40, 60 and 80 mmHg in GFP (red, n=4 mice) and hM4Di (green, n=7 mice) group. Two-way ANOVA with Šídák's multiple comparisons.

**References**

Bielefeldt, K., Davis, B. & Binion, D. G. Pain and inflammatory bowel disease. *Inflamm Bowel Dis* **15**, 778–788 (2009). https://doi.org/10.1002/ibd.20848

Coates, M. D. *et al.* Abdominal Pain in Inflammatory Bowel Disease: An Evidence-Based, Multidisciplinary Review. *Crohns Colitis 360* **5**, otad055 (2023). https://doi.org/10.1093/crocol/otad055

Yang, D. *et al.* Nociceptor neurons direct goblet cells via a CGRP-RAMP1 axis to drive mucus production and gut barrier protection. *Cell* **185**, 4190–4205 e4125 (2022). https://doi.org/10.1016/j.cell.2022.09.024

Drewes, A. M. *et al.* Gastrointestinal pain. *Nat Rev Dis Primers* **6**, 1 (2020). https://doi.org/10.1038/s41572-019-0135-7

Rao, M. An increasingly complex view of intestinal motility. *Nat Rev Gastroenterol Hepatol* **17**, 72–73 (2020). https://doi.org/10.1038/s41575-019-0249-0

Defaye, M., Nasser, Y. & Altier, C. Mechanisms of chronic abdominal pain in inflammatory bowel disease and implications for treatment. *Nat Rev Gastroenterol Hepatol* (2025). https://doi.org/10.1038/s41575-025-01139-8

Takahashi, K. *et al.* Post-inflammatory Abdominal Pain in Patients with Inflammatory Bowel Disease During Remission: A Comprehensive Review. *Crohns Colitis 360* **3**, otab073 (2021). https://doi.org/10.1093/crocol/otab073

Latremoliere, A. & Woolf, C. J. Central sensitization: a generator of pain hypersensitivity by central neural plasticity. *J Pain* **10**, 895–926 (2009). https://doi.org/10.1016/j.jpain.2009.06.012

Woolf, C. J. Central sensitization: implications for the diagnosis and treatment of pain. *Pain* **152**, S2–S15 (2011). https://doi.org/10.1016/j.pain.2010.09.030

Zeng, X., Powell, R. & Woolf, C. J. Mechanism-based nonopioid analgesic targets. *J Clin Invest* **135** (2025). https://doi.org/10.1172/JCI191346

Alexandre, C. *et al.* Nociceptor spontaneous activity is responsible for fragmenting non-rapid eye movement sleep in mouse models of neuropathic pain. *Sci Transl Med* **16**, eadg3036 (2024). https://doi.org/10.1126/scitranslmed.adg3036

Chiang, M. C. *et al.* Divergent Neural Pathways Emanating from the Lateral Parabrachial Nucleus Mediate Distinct Components of the Pain Response. *Neuron* **106**, 927–939 e925 (2020). https://doi.org/10.1016/j.neuron.2020.03.014

Han, S., Soleiman, M. T., Soden, M. E., Zweifel, L. S. & Palmiter, R. D. Elucidating an Affective Pain Circuit that Creates a Threat Memory. *Cell* **162**, 363–374 (2015). https://doi.org/10.1016/j.cell.2015.05.057

Palmiter, R. D. The Parabrachial Nucleus: CGRP Neurons Function as a General Alarm. *Trends Neurosci* **41**, 280–293 (2018). https://doi.org/10.1016/j.tins.2018.03.007

Rodriguez, E. *et al.* A craniofacial-specific monosynaptic circuit enables heightened affective pain. *Nat Neurosci* **20**, 1734–1743 (2017). https://doi.org/10.1038/s41593-017-0012-1

Choi, S. *et al.* Parallel ascending spinal pathways for affective touch and pain. *Nature* **587**, 258–263 (2020). https://doi.org/10.1038/s41586-020-2860-1

Yarmolinsky, D. A. *et al.* Differential modification of ascending spinal outputs in acute and chronic pain states. *Neuron* **113**, 1223–1239 e1225 (2025). https://doi.org/10.1016/j.neuron.2025.01.031

Huang, T. *et al.* Identifying the pathways required for coping behaviours associated with sustained pain. *Nature* **565**, 86–90 (2019). https://doi.org/10.1038/s41586-018-0793-8

Goldstein, N. *et al.* A parabrachial hub for need-state control of enduring pain. *Nature* **647**, 689–697 (2025). https://doi.org/10.1038/s41586-025-09602-x

Nardone, S. *et al.* A spatially-resolved transcriptional atlas of the murine dorsal pons at single-cell resolution. *Nat Commun* **15**, 1966 (2024). https://doi.org/10.1038/s41467-024-45907-7

Pauli, J. L. *et al.* Molecular and anatomical characterization of parabrachial neurons and their axonal projections. *Elife* **11** (2022). https://doi.org/10.7554/eLife.81868

Yao, Z. *et al.* A high-resolution transcriptomic and spatial atlas of cell types in the whole mouse brain. *Nature* **624**, 317–332 (2023). https://doi.org/10.1038/s41586-023-06812-z

Ma, Q. A functional subdivision within the somatosensory system and its implications for pain research. *Neuron* **110**, 749–769 (2022). https://doi.org/10.1016/j.neuron.2021.12.015

Chiang, M. C. *et al.* Parabrachial Complex: A Hub for Pain and Aversion. *J Neurosci* **39**, 8225–8230 (2019). https://doi.org/10.1523/JNEUROSCI.1162-19.2019

Li, Y. *et al.* Tachykinin signaling in the right parabrachial nucleus mediates early-phase neuropathic pain development. *Neuron* **113**, 605–619 e606 (2025). https://doi.org/10.1016/j.neuron.2024.11.021

Condon, L. F. *et al.* Parabrachial Calca neurons drive nociplasticity. *Cell Rep* **43**, 114057 (2024). https://doi.org/10.1016/j.celrep.2024.114057

Chassaing, B., Aitken, J. D., Malleshappa, M. & Vijay-Kumar, M. Dextran sulfate sodium (DSS)-induced colitis in mice. *Curr Protoc Immunol* **104**, 15 25 11–15 25 14 (2014). https://doi.org/10.1002/0471142735.im1525s104

Wirtz, S. *et al.* Chemically induced mouse models of acute and chronic intestinal inflammation. *Nat Protoc* **12**, 1295–1309 (2017). https://doi.org/10.1038/nprot.2017.044

Barkai, O. *et al.* A machine learning tool with light-based image analysis for automatic classification of 3D pain behaviors. *Cell Rep Methods* **5**, 101145 (2025). https://doi.org/10.1016/j.crmeth.2025.101145

Zhang, Z. *et al.* Automated preclinical detection of mechanical pain hypersensitivity and analgesia. *Pain* **163**, 2326–2336 (2022). https://doi.org/10.1097/j.pain.0000000000002680

Nagai, Y. *et al.* Deschloroclozapine, a potent and selective chemogenetic actuator enables rapid neuronal and behavioral modulations in mice and monkeys. *Nat Neurosci* **23**, 1157–1167 (2020). https://doi.org/10.1038/s41593-020-0661-3

Liu, S. *et al.* Divergent brainstem opioidergic pathways that coordinate breathing with pain and emotions. *Neuron* **110**, 857–873 e859 (2022). https://doi.org/10.1016/j.neuron.2021.11.029

Jain, A. *et al.* Nociceptor-immune interactomes reveal insult-specific immune signatures of pain. *Nat Immunol* **25**, 1296–1305 (2024). https://doi.org/10.1038/s41590-024-01857-2

Sammons, M. *et al.* Brain-body physiology: Local, reflex, and central communication. *Cell* **187**, 5877–5890 (2024). https://doi.org/10.1016/j.cell.2024.08.050

Meerschaert, K. A. & Chiu, I. M. The gut-brain axis and pain signalling mechanisms in the gastrointestinal tract. *Nat Rev Gastroenterol Hepatol* **22**, 206–221 (2025). https://doi.org/10.1038/s41575-024-01017-9

36 Wang, H. *et al.* A tool kit of highly selective and sensitive genetically encoded
neuropeptide sensors. *Science* **382**, eabq8173 (2023).
https://doi.org/10.1126/science.abq8173

37 Gebhart, G. F. & Bielefeldt, K. Physiology of Visceral Pain. *Compr Physiol* **6**, 1609–
1633 (2016). https://doi.org/10.1002/cphy.c150049

38 Campos, C. A., Bowen, A. J., Roman, C. W. & Palmiter, R. D. Encoding of danger by
parabrachial CGRP neurons. *Nature* **555**, 617–622 (2018).
https://doi.org/10.1038/nature25511

39 Van Oudenhove, L. *et al.* Common and distinct neural representations of aversive
somatic and visceral stimulation in healthy individuals. *Nat Commun* **11**, 5939
(2020). https://doi.org/10.1038/s41467-020-19688-8

40 Wolfson, R. L. *et al.* DRG afferents that mediate physiologic and pathologic
mechanosensation from the distal colon. *Cell* **186**, 3368–3385 e3318 (2023).
https://doi.org/10.1016/j.cell.2023.07.007

41 Gao, X. *et al.* Calcium imaging in population of dorsal root ganglion neurons
unravels novel mechanisms of visceral pain sensitization and referred somatic
hypersensitivity. *Pain* **162**, 1068–1081 (2021).
https://doi.org/10.1097/j.pain.0000000000002096

42 Gazit Shimoni, N. *et al.* Changes in neurotensin signalling drive hedonic
devaluation in obesity. *Nature* **641**, 1238–1247 (2025).
https://doi.org/10.1038/s41586-025-08748-y

43 Li, H. *et al.* Neurotensin orchestrates valence assignment in the amygdala. *Nature*
**608**, 586–592 (2022). https://doi.org/10.1038/s41586-022-04964-y

44 Guo, R. *et al.* Arrestin-biased allosteric modulator of neurotensin receptor 1
alleviates acute and chronic pain. *Cell* **188**, 4332–4349 e4321 (2025).
https://doi.org/10.1016/j.cell.2025.04.038

45 Castagliuolo, I. *et al.* Neurotensin is a proinflammatory neuropeptide in colonic
inflammation. *J Clin Invest* **103**, 843–849 (1999).
https://doi.org/10.1172/JCI4217

46 Gui, X., Liu, S., Yan, Y. & Gao, Z. Neurotensin receptor 1 overexpression in
inflammatory bowel diseases and colitis-associated neoplasia. *World J
Gastroenterol* **19**, 4504–4510 (2013). https://doi.org/10.3748/wjg.v19.i28.4504

47 Law, I. K. *et al.* Neurotensin--regulated miR-133alpha is involved in
proinflammatory signalling in human colonic epithelial cells and in experimental
colitis. *Gut* **64**, 1095–1104 (2015). https://doi.org/10.1136/gutjnl-2014-307329

48 Udit, S., Blake, K. & Chiu, I. M. Somatosensory and autonomic neuronal regulation
of the immune response. *Nat Rev Neurosci* **23**, 157–171 (2022).
https://doi.org/10.1038/s41583-021-00555-4

49 Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body
parts with deep learning. *Nat Neurosci* **21**, 1281–1289 (2018).
https://doi.org/10.1038/s41593-018-0209-y

50 Gonzalez-Cano, R. *et al.* Up-Down Reader: An Open Source Program for Efficiently
Processing 50% von Frey Thresholds. *Front Pharmacol* **9**, 433 (2018).
https://doi.org/10.3389/fphar.2018.00433

51 Christianson, J. A. & Gebhart, G. F. Assessment of colon sensitivity by luminal
distension in mice. *Nat Protoc* **2**, 2624–2631 (2007).
https://doi.org/10.1038/nprot.2007.392

52    Muppirala, A. N. *et al.* Tachykinin signaling defines distinct populations of glia in
the     enteric     nervous     system.     *Neuron*     (2026).
https://doi.org/10.1016/j.neuron.2025.11.030
1093