Rapid modular evolution of antiviral repertoires in P2 phages and their P4 satellites: shuttling, swapping and mixing

Jorge Moura de Sousa[1*], Florence Depardieu[2], Marie Touchon[1], Raphaël Laurenceau[1,2], Kristen Curry[3], Alice Maestri[1], Florian Tesson[4], Arthur Loubat[4], Rayan Chikhi[3], Jean Cury[4], Aude Bernheim[4], David Bikard[2] and Eduardo P.C. Rocha[1*]

[1] Institut Pasteur, Université de Paris Cité, CNRS UMR3525, Microbial Evolutionary Genomics, Paris, France
[2] Institut Pasteur, Université de Paris Cité, CNRS UMR3525, Synthetic Biology, Paris, France
[3] Institut Pasteur, Université de Paris Cité, CNRS UMR3525, Sequence Bioinformatics, Paris, France
[4] Institut Pasteur, Université de Paris Cité, CNRS UMR3525, Molecular Diversity of Microbes, Paris, France

*Corresponding authors: jorge-andre.sousa@pasteur.fr; eduardo.rocha@pasteur.fr

ORCID IDs:
Jorge Moura de Sousa: 0000-0003-3530-8550
Florence Depardieu: 0000-0002-2680-0589
Marie Touchon: 0000-0001-7389-447X
Raphael Laurenceau: 0000-0002-1326-9928
Kristen Curry: 0000-0001-8375-397X
Alice Maestri: 0000-0002-2176-8831
Florian Tesson: 0000-0003-4038-1154
Arthur Loubat: 0009-0006-4647-4561
Rayan Chikhi: 0000-0003-1099-8735
Jean Cury: 0000-0002-6462-8783
Aude Bernheim: 0000-0003-0212-777X
David Bikard: 0000-0002-5729-1211
Eduardo P. C. Rocha: 0000-0001-7704-822X

Abstract

Interactions between bacteria, bacteriophages, and their satellites are shaped by a myriad of defence and counter-defence mechanisms. Here, we identified and characterized the defence hotspots of thousands of P2-like phages and P4-like satellites to elucidate the origins and evolutionary dynamics of defence systems. Both P4 and P2 encode a broad diversity of recognizable defence systems. Defences are a substantial, yet likely underestimated, share of the elements' pangenomes, as shown by novel antiviral functions discovered in P4 loci lacking known defence genes. Defence loci are very rapidly swapped, without pseudogenization, suggesting defences are replaced before becoming non-adaptive. This intense local recombination melds components of distinct systems into novel functional chimeras. Systems swap so rapidly that many elements with identical core genes have completely different defences. Surprisingly, despite P4 and P2's concomitant replication and packaging, they almost never exchange

45   defence genes. In contrast, near identical defence systems can be found in distinct types

46   of MGEs and in cryptic chromosomal locations. Our findings highlight P4 and P2 as

47   mobile platforms driving the modular diversification of bacterial antiviral repertoires.

48   Hence, bacterial defences change quickly by phage and satellite turnover, and by the

49   quick swap of defences within these elements.

Introduction

Microbial ecosystems are shaped by complex interactions between bacteria and bacteriophages (phages). These interactions range from the direct antagonism between virulent phages and bacteria[1] to the more nuanced continuum between antagonism and commensalism of temperate phages and their hosts, where the interests of both bacteria and phage can be aligned[2,3]. One key example of the latter is the presence of antiviral defences in prophages, which protect their host bacteria from predation by other phages, thus ensuring the survival of both[4,5]. Prophages are not the only mobile genetic elements (MGEs) that provide antiviral functions to their hosts. Anti-MGE defence systems are frequently encoded by MGEs present in bacterial genomes[6], possibly because of competition, and/or cooperation, between MGEs[7]. This is exemplified by phage satellites, non-autonomous mobile elements (hitchers) that hijack (helper) phages to mobilize between bacteria[8,9]. Some P4-like satellites encode antiviral defences against competitors of P2-like helper phages, protecting both helper and bacterial host[10]. Hence, the associations between satellites, phages and bacteria evolve in a mutualism-parasitism continuum, where partners incur in costs and benefits associated with the presence of the other elements[8].

The diversity of known defence systems is rapidly increasing. Defence genes have one of the highest rates of gain/loss across bacterial genomes[11], and in bacteria, their acquisition is typically associated with the mobility of the MGEs carrying them[12]. But little is known about the evolutionary dynamics of defence genes in MGEs themselves. Where do these genes come from, and how stable are they? What is the fate of these genes, and that of the MGEs carrying them, once defences are no longer useful? These questions are intricately linked not only with the ecology of MGEs (who is defending whom, when and against what), but also with their genomic organization, which for some elements like phages and satellites, can be tightly conserved[9,13,14]. The accumulation of accessory genes in phages is constrained by the size of their genomes, which must fit within their capsids[15], and the very modular genomic organization of both satellites and phages[9,14]. Although prophages often have accessory genes useful to their bacterial hosts, their high gene density suggests they rarely have pseudogenes and is thought to explain the low

frequency of transposable elements in their genomes, in contrast with plasmids[16].

Hence, the systematic detection of defence genes in prophages and satellites reflects

the selective importance of these functions for their ecology and evolution. At a broader

level, understanding the prevalence, localization and evolutionary dynamics of antiviral

genes can highlight the rules that govern the genomic plasticity and genomic organization

of MGEs, as well as their ecological associations and interactions with bacterial hosts

and with other MGEs.

Several defence systems were previously detected in P2-like phages and P4-like

satellites, in a single hotspot near the *cos* site of each element[10]. Here, we delimit

satellite and phage genomes to identify additional hotspots and to account for MGE

similarity when studying the evolutionary dynamics of antiviral genes within and across

P4 satellites and P2 helpers. To extend the diversity of our dataset, we coupled the

detection and precise delimitation of thousands of P4-like satellites[17] and P2-like

prophages in complete bacterial genomes with thousands of additional P4-like satellites

detected in metagenomes to generate the most abundant and diverse genomic

collection of satellites and helper phages to date. The extensive dataset of P4 and P2

genomes allowed to uncover the quick turnover of defence genes encoded by P4 and P2,

the genetic exchanges that underly this turnover, and how they shape defence

repertoires.

RESULTS

Defence systems' loci are pervasive in a few hotspots of P2 and P4

We identified the P2-like prophages (P2 for short) and P4-like satellites (P4 for short) in the complete genomes of *Escherichia coli, Klebsiella pneumoniae, Salmonella enterica*. We also used the P4-like contigs detected in Logan, a repository of assembled sequence data from tens of millions of public sequencing experiments[18], which greatly expanded the number and genomic diversity of known P4 satellites (Fig S1, see Methods). The 3085 P4 (from complete bacterial genomes and metagenomes) and 1511 P2 (from complete bacterial genomes) elements in our dataset were dereplicated and genetically delimited at the nucleotide level by identification of their *att* sites (see Methods). We then identified the defence systems in P4 and P2 using DefenseFinder[19] and PADLOC[20]. We also searched for anti-defence systems, for which we only find two types (Anti_RM and Anti_CRISPR) distributed in 28 P2. Most of these P2 have a single anti-antiviral system, but one has five distinct systems. We detected no anti-defence systems in P4. Given the low numbers of these anti-defence systems, we decided to exclude them from the main analysis.

We found a very broad diversity of known types of defence systems, which occurred in 57% of P4 and in 49% of P2 (Fig 1A and B, Fig S2A for PADLOC classification). Some types of systems were detected in both satellites and phages (e.g., AbiU or PD-T7-2), whereas others were found in only one of them (e.g., AbiP2 in P2 and Hachiman in P4). We also identified individual genes that are usually associated with defence systems, but that do not form a complete known system (see the two rightmost columns in Fig 1A for systems that were partially detected by DefenseFinder). We will refer to these as putative defence systems and will explore them in more detail in a subsequent section. If we include both complete and putative systems, the proportion of elements that encode defence-associated genes increases to 73% (P4) and 81% (P2) (Fig 1B). Hence, a large majority of these two types of MGEs encode genes associated with antiviral activity.

Even if this is by far the largest known dataset of P4 and P2, the pangenome saturation curves show that we are far from having uncovered the full genetic diversity of these elements (Fig 1C). Known defence genes correspond to a sizeable fraction of these

pangenomes (orange lines): 34% for P4 and 17% for P2. P4 and P2 account for 11-16% of

the genetic diversity of the defence systems in their hosts' species (Fig S3). Moreover, up

to 50% of the total defence systems of some genomes are encoded in their P4 or P2

elements (Fig 1D, Fig S4). This suggests that P4 and P2 are important contributors to the

defence repertoire of bacteria.

Most defence systems in P4 are in a hotspot between the integrase and Psu-Delta-Sid,

whilst many of the defence systems in P2 are in a hotspot between the primase (GpA) and

the portal (GpQ) (Fig 1E). These are the two previously described hotspots[10]. However,

other less populated defence hotspots exist between the primase and one of the *att* sites

of P4 (7% of the systems in P4), between the integrase and the *att* site of P2 (3% of the

systems in P2), between the integrase and the primase in P2 (8% of the systems in P2)

and within the late genes of P2. Although it is atypical for integrated elements to mobilize

genes between *att* sites and the integrase, this was previously observed for defence

systems of PICI[21]. An additional P2 defence hotspot was found downstream of GpL (16%

of the systems in P2), consistent with previous descriptions of a hotspot of accessory

genes within the tail genes in the classical P2 phage[22]. Moreover, some of these P2

defence hotspots correspond to known integration sites of horizontally transferred genes

in P2[23], and to the hotspots of accessory genes described recently in more than 1000 P2-

like phages[24]. In either P4 or P2, some defence systems were consistently found in a

single hotspot, whilst others were found across genomic locations of the elements (Fig

S5A). Of the P2 with complete defence systems, 10% have multiple defence systems

across different hotspots (Fig S2C, see also genomes in Fig 1E for examples). This fraction

is much lower for P4 (1%), suggesting that the accumulation of multiple defence systems

occurs more often in the larger genomes of P2. These results confirm that whilst defence

systems are very abundant and diverse in P4 and P2, representing a large fraction of their

accessory genes, they are encoded in a small number of genomic locations.

163
164
165 **Figure 1. Abundance, distribution and localization of defence systems in P4-like satellites and in P2-**

166 **like prophages. A.** Number of the different types of complete (two leftmost columns) and putative (two

167 rightmost columns) defence systems detected by DefenseFinder in P4 (1st and 3rd column) and P2 (2nd and

168 4th column). Darker colours in the heatmap correspond to higher numbers of systems. **B.** The proportion of

169 P4 and P2 elements with defence systems deemed complete (orange, as detected by DefenseFinder or

PADLOC) or putative (light orange, detected by DefenseFinder). **C.** The pangenome (black) of P4-like satellites (top) and P2-like phages (bottom) with the relative contributions of the core (blue for P4, red for P2) and the defence-associated genes (orange). **D.** Fraction of systems (detected by DefenseFinder) of the host encoded in its P4 and P2 elements. The proportions of complete defence systems encoded by P4 in hosts that have at least one P4 (leftmost panel), encoded by P2 in hosts that have at least one P2 (middle panel), or encoded either by P4 or P2 in hosts that have at least one of either element (rightmost panel) are shown in the x-axis. The size of the circles corresponds to the frequency of each (x,y) pair. **E.** Distribution of defence systems in hotspots of P4-like satellites (top) and P2-like prophages (bottom). Hotspots are defined as locations among successive core gene families and are indicated by the edges in the graphs (% indicates the proportion of defence systems at these locations). When the core genes are not successive, it reflects cases where either one of the core genes is missing, or the cores genes are arranged differently from the prototypical organization of their respective element. Colour codes in genetic loci and graphs coincide; orange indicating complete and light orange indicating putative defence systems. Grey indicates genes that are neither core nor defence-associated.

Rapid turnover of defence systems

We then enquired on the turnover of the defence systems, i.e. how frequently closely related elements differ in the systems they encode. The phylogenies of P4 and P2 elements, based on their core genes (see Methods), show highly similar elements often carrying unrelated defences (Fig 2A and D). This can best be appreciated by focusing on the parts of the trees with many very similar elements and inspecting the diversity of systems they contain. To quantify this effect, we compared the gene repertoire relatedness (with wGRR, see Methods) of core genes and of defence genes between pairs of elements with defence systems. Consistent with the phylogenies, we observed an overwhelmingly large proportion of pairs of the same type of elements with non-homologous defence systems (99% in P4 and 99% in P2 pairs have wGRR defence<0.05). Since this includes pairs of very similar elements (i.e., with highly similar core genes) (Fig 2B and E, right panels), it shows that defence genes change very rapidly in both P4 and P2. We observed only a few pairs of dissimilar elements with similar defence systems suggesting that convergent acquisition of defences is rare (Fig 2B and E, left panels). These patterns remain qualitatively unchanged when comparing full genomes instead of core genes and when including the putative defence systems (Fig S6). If defence systems evolved in linkage with the core genomes of P4 and P2, we would expect a linear relationship between core and defence similarity, with deviations corresponding to

potential events of horizontal transfer of antiviral genes. Instead, the association between the evolution of defence systems and the core genes is almost null.

We then quantified the genetic linkage between defence genes and each individual core gene (Fig 1E). The Spearman correlations between the wGRR of defence genes and the sequence similarity of each core gene in pairs of P4 (Fig 2C and Fig S7A) and P2 (Fig 2F and Fig S7B) were all very low (≤0.2), in contrast to the correlations between core genes themselves. In summary, similar elements rarely encode similar defence genes and the latter show little genetic linkage with core genes. Both indicate very rapid turnover of defence systems in the genomes of P4 and P2.

Figure 2. Relationship between antiviral repertoires and the core genomes of P4 and P2. A. Phylogeny of core genes of P4. Bootstraps of at least 90 are indicated with a circle (the larger the circle, the higher the bootstrap value). Colours in the innermost circle indicate the bacterial host of each P4 (white spaces

indicate P4 from metagenomes, without an assigned host). Note that since these are dereplicated P4

genomes, some P4 might be representative of a group of elements with multiple bacterial hosts (we

consider only the host of the representative genome). The ten outward circles represent the presence

(orange) or absence (white) of the ten most frequent defence systems in P4 (complete defence systems,

from DefenseFinder). The outermost circle indicates whether each P4 element has any other complete

defence system, apart from the ten most frequent in the family of elements (black shows cases where

another defence system is present). **B.** Left, number of pairs of P4 genomes with similar defence genes

(wGRR≥0.95) in function of the similarity between core genes (x-axis). Right, number of pairs of P4 genomes

with distinct defence genes (wGRR≤0.05) in function of the similarity between core genes (x-axis). **C.**

Spearman correlation values between the wGRR of defence genes and the bitscore values from BlastP of

individual core genes of P4-like satellites, for the set of all pairs of P4s. Colours of a stronger shade indicate

higher Spearman correlation values. **E.** Same as in A, but for P2 genomes and their defence systems. **F.**

Same as in **B**, but for P2 genomes. **G.** Same as in **C**, but for the defence and core genes of P2-like phages.

Pseudogenes of defence systems are rare in P4 and P2

Turnover of defence systems could occur in two ways. They could be first inactivated by

mutations or indels and subsequently replaced (Deletion-reacquisition model).

Alternatively, functional systems could be swapped by others before becoming defective

(Swap model). The first scenario suggests that systems are lost because they are not

selected for anymore, i.e., they are nearly neutral or even deleterious (costs outweigh

gains). The second scenario only requires that the new defence system is at least as

adaptive as the old one in the current population context. These models have different

testable predictions. The deletion-reacquisition model predicts the frequent presence of

pseudogenes of defence systems whereas the swap model predicts the opposite. We

assessed the fraction, type, and past function of pseudogenes in P4 and P2 using

PseudoFinder[25] (see Methods). The vast majority of P4 and P2 have at least one

pseudogene (94% and 98%, respectively, Fig 3), showing that defective genes remain long

enough to be identifiable in large numbers in these genomes. We note that, in some

cases, several identified pseudogenes can originate from a single gene. Although

pseudogenes are frequently found in defence-associated loci (Fig S8), only a very small

percentage of them had sequence similarity with defence-associated genes (less than

1% in P4 and c.a. 2% in P2). Many more had sequence similarity to core genes (24% in P4,

typically Ash pseudogenes) and especially to unknown function genes (Fig 3 insets, see

File S1 for PFAM annotations). These results suggest that defence systems are frequently

swapped before being inactivated. Moreover, most pseudogenes in P4 and P2 are

detected as very small intergenic sequences (c.a. 90% in either P4 or P2, Fig S9), which

confirms that the putative defence systems detected in these elements are intact, likely

functional genes.

260
261
262

263 **Figure 3. Quantification and characterization of pseudogenes in P4-like satellites and P2-like**

264 **prophages.** Histograms show the distribution of pseudogenes detected by PseudoFinder per P4-like

265 satellite (left) and P2-like prophage (right). The charts as insets show the proportion of the pseudogenes

266 annotated as defence (orange), core of each element (blue and red for P4 and P2, respectively), or with

267 other (or no) annotations (grey).

268
269

P4 hotspots contain numerous novel or chimeric defence systems

We found many genes of defence systems that lacked the typical cognate components

in the defence hotspots (Fig S5B). These could be inactivated systems, but our

observation that pseudogenes of defence systems are rare make this hypothesis unlikely.

Instead, they might be functional novel systems resulting from re-assembly of others[26,27].

To test this hypothesis, we selected 11 candidate systems based on their different

characteristics (Fig 4, see File S2 for a description of the putative systems): some are

potential variants of known systems (e.g., a Lamassu system missing a known effector);

others are mixtures of genes from different systems (e.g., two genes from the PD Lambda
2 system, and another from the Rst_gop_cll system); and others are potential
minimizations of known systems (e.g., a Thoeris system with a single ThsA gene). We
synthesized and cloned each putative system in a low copy plasmid under the control of
a pTet promoter, which were introduced into *E. coli* MG1655. An empty plasmid was
transformed to the same background as a control. We then challenged these strains with
a collection of 28 virulent phages to test the antiviral protection of each putative defence
system (see Methods, Fig S10, File S2). 8 out of 11 provided protection against several
virulent phages (Fig 4, top 11 rows). Failure to observe protection can arise from multiple
causes. The most likely explanations are: (i) the defence system is functional but does
not target any of the 28 phages included in the present panel; or (ii) the system is
functional in its native host but is not properly expressed, assembled, or active in the
heterologous *E. coli* host used here. Interestingly, a system similar to the Lamassu
variant, which did not provide protection against the phages tested here, was recently
shown to be provide an anti-phage function[28]. The high success rate we obtained in
detecting resistance against this limited phage panel strongly suggests that most of the
putative systems not recognized as complete by either DefenseFinder or PADLOC are
functional.
297
Some P4 (27%) and P2 (19%) lack identifiable defence genes (Fig 1B) but have genes of
unknown function at their main defence loci. We hypothesised that some of these could
be novel antiviral systems. We selected at random 8 P4 satellites with genes of unknown
function to test their anti-phage phenotypes, as above (Fig 4, bottom 8 rows, Fig S10). 6
out the 8 conferred protection against multiple virulent phages. phages. As previously,
this high success rate indicates that most of these unannotated loci are likely *bona fide*,
functional defence systems. We named the novel functional systems according to deities
from Lusitanian mythology, while waiting for a uniform standardised system of
nomenclature, and they are described in detail in File S3.
307
308 When possible, we characterized the novel systems using available data, sensitive
309 sequence similarity detection programs (wjth HHPred), and structural similarity (with
310 ESMFold and FoldSeek, File S3). The Crouga system has some evidence of being an RM-

like system, with one of its proteins containing PFAM domains (ResIII, DEAD and Helicase C) associated with other RM systems[29]. The Bandua system has two proteins with domains typically associated with immune functions (a nuclease and a kinase, respectively). Neither protein is recognized by current antiviral annotation pipelines. The Aernus system has one protein annotated with kinase and nuclease domains, and a second one annotated as a mannitol repressor. It is unknown at this point if the latter has any impact in antiviral capabilities. The single protein in the Toga system has several hits with protein domains associated (through FoldSeek) with the Gao_Ppl[30] and the SMC defence systems[31], albeit not being recognized by any of the current models for either system. The Reve system is another single gene system, with ATPase and KAP family P-loop domains[32]. Finally, the last system we tested has two proteins, one with pentapeptide repeats and an IRK potassium channel domain, which could be involved in membrane depolarization, while the other protein contains a Nucleotidyltransferase like domain and a HEPN-like domain, suggesting it acts on host or phage RNA. A system with similar components was described by others during the course of this study as Hailong[33], which is currently not detected by DefenseFinder or PADLOC. Hence, and despite being part of the non-defence annotated genes in P4, we refer to it as Hailong to avoid introducing redundant nomenclature.

The novel systems described above, together with the unannotated Hailong system, are clear evidence that at least some genes of unknown function in P4 have antiviral phenotypes, with most (e.g., Bandua, Aernus, Toga and Reve) providing strong protection against Drexlerviridae and/or Siphoviridae phages). We thus wondered whether other genes in the defence-less elements could have similar properties. For this, we collected all the genes of unknown function from P2 and P4 detected only in complete genomes, to use the information on their larger chromosomal context, and analysed them using a recent machine learning approach to detect antiviral domains (see Methods). We observed a significant signal for the association of antiviral defences in 30% of the proteins of unknown function in P4 (62%, with the method that considers the genomic context), and in 15% of the unknown proteins in P2 (32%, with the method that considers the genomic context) (Fig S11). We note that all the confirmed novel systems are part of those predicted to have an antiviral function, due to their genetic context. The

computance predictions, together with experimental validations, demonstrate that
P4-like satellites constitute major loci for both established and yet-to-be-discovered
functional defence systems. Moreover, many of the proteins of unknown function that
cluster in P4- and P2-associated defence hotspots are predicted to encode antiviral
activities.

Figure 4. Putative and unknown defence systems in P4 provide protection against phage infection. On the left are the genomes of P4 with putative defence systems (light orange genes, top 11 rows) and the genomes of P4 with genes of unknown function in their defence loci (grey genes, bottom 8 rows). For the former, the names of the genes, as well as the complete systems with which they are associated, are shown in front of each genome. The regions synthetized in plasmids are boxed (exact sequences in Datasets E1 and E2). Heatmap of the phage resistance phenotype shows the mean fold resistance of two independent replicates of each putative and unknown system against a panel of 28 *E. coli* phages. The degree of protection is indicated by the colour (log) scale, with darker red colours indicating higher levels of protection, relative to the same bacterial background with an empty plasmid. Only the infections where both replicates show increased protection, and the mean of the two replicates is ≥ 10-fold difference, are shown as coloured squares. Asterisks indicate smaller plaque morphologies, relative to the infection of the strain with the empty plasmid. Plaque-forming units were measured for each phage on cells harbouring either a control plasmid or a defence system (Fig S10). Fold-resistance was calculated as the ratio between these two values. All systems were expressed from pTet promoter in the presence of anhydrotetracycline (aTc, 0.5 µg/ml) and measured at 37°C, except for T7-like phages, which were measured at room temperature. The raw data for the ratios system/control are shown in File S4.

P4 and P2 exchange defences with other MGEs but not with each other

If defence systems are constantly swapped in both P4 and P2, where do these genes

come from? To address this question, we searched for closely related homologs (>85%

identity, see Methods) of defence genes from either P4 or P2 in 32798 bacterial and in

27022 phage genomes. For 6% of defence genes in P4, and for 2% of the defence genes

in P2, we found at most one homolog, indicating that some of these defence genes are

likely variants specific to the P4 and P2 in our dataset (Fig S12). Many of the remaining P4

and P2 defence genes have homologs in multiple genomic contexts (Fig 5A). Similar

results were found when including the putative novel defence systems from the first

section (File S5, Fig S13). To avoid comparing elements that might have similar systems

because of recent common ancestry, we excluded the pairs of very similar (wGRR>=0.95)

P4 or P2 elements. Nevertheless, the analysis reveals that most homologs of defence

genes of P4 (76%) and P2 (47%) are in bacterial regions predicted to be respectively P4-

like satellites and P2 phages. These include P4 and P2 in bacterial hosts (e.g.,

Citrobacter, *Enterobacter* or *Pectobacterium*, Fig S14) where these elements'

delimitation in bacterial chromosomes was not possible for lack of information on their

att sites (called P2* and P4*, see Methods). These elements had hence not been included

in our focal dataset of phages and satellites. P2 and P4 often co-occur in genomes and

are induced and replicated at the same time. One could thus expect that they would

exchange defence systems. Surprisingly, we found only one case of closely related

defence genes between P4 and P2, even when including all putative defence systems (Fig

S15). Hence, hitchers and helpers have not recently exchanged defence systems.

The highly similar homologs of P4 and P2 defence genes found in bacterial regions that

are not identified as P4 or P2-like could be the result of recent genetic exchanges with

other MGEs. We found no homologs in the other families of satellites known in

Enterobacteria (PICI and cfPICI[9,17],File S5). Both P4 and P2 defence homologs were found

in (non-P2 like) prophages, albeit this was much more frequent for P2, which also has

defence homologs in 31 temperate and 11 virulent phage genomes (all P2-like).

Chromosomal regions with integrative conjugative elements[34], which are known to

encode antiviral systems[35], account for very few of the homologs of P4 or P2 defence

genes. P4 and especially P2 defence systems have relatively more homologs in plasmids.
Some of the latter could correspond to phage-plasmids (P-Ps), since they are plasmid
replicons identified as phages by geNomad[36], although they are not part of the current
dataset of known P-Ps[37]. The MGEs and bacterial regions with homologs of defence genes
of P4 and P2 are sometimes within bacteria that are phylogenetically distant from the
hosts of P4 and P2, such as *Pseudomonas* or *Vibrio* (Fig S14). These results suggest that
exchange of defence systems tends to occur between similar types of elements, but can
also involve distinct MGEs.

Genetic exchanges with other MGEs may allow to introduce novel types of defences in P2
or P4, which can then be exchanged within each group of elements. For instance, the
phylogeny of a variant of the AVS_V system, detected in P2 from *K. pneumoniae*, includes
homologs in non-P2 prophages and uncharacteristic genomic regions from different
bacterial hosts, that form a separate clade (Fig 5B). Intermingled with the two main
clades are two terminal branches, corresponding to an uncharacteristic genomic region
in *Proteus* and a single plasmid in *Duffyella*. The phylogeny suggests the acquisition of
the system by P2 was recent. It was then followed by the dissemination of this variant of
AVS_V within *K. pneumoniae* in P2 phages. Another defence system detected in P2, Lit,
is mostly found in plasmids and uncharacteristic genomic regions. While being rare in
prophages, a homolog is found in a single P2-like prophage (Fig S16A). Given its
singularity, and the scarcity of this type of systems in P2 (Fig 1A), this likely illustrates a
recent acquisition of the Lit system by this P2.

Exchange of defence systems between different types of MGEs also broadens the
mechanisms of dissemination of the former within bacterial species. For instance, the
phylogeny of a variant of the PD-T7-2 A is consistent with a unique, recent exchange of
this system between P4 and plasmids (whose direction cannot be ascertained from this
data alone). The system in the plasmids has diversified in two separate genetic variants
that disseminated in *K. pneumoniae* (Fig 5C). The phylogeny of the other gene of the
system (PD-T7-2 B) shows yet another variant in P4 that is disseminating the defence
system in *Escherichia* and *Klebsiella* hosts, with a transfer to plasmids in the latter (Fig
S16B). Hence, events of exchange between MGEs show the establishment and further

diversification of defence homologs in different MGEs, allowing the spread of defence

systems within and across species through different mechanisms of horizontal gene

transfer.

Figure 5. Highly similar homologs of defence genes of P4 and P2 are found in completely different

genomic contexts. A. Graph based representation of the number of instances where homologs of defence

genes (from complete defence systems) in P4 and P2 are found in different genomic contexts. These

include other MGEs (plasmids, phage-plasmids, conjugative integrated elements, prophages and virulent

phages) or regions of the bacterial genomes that are uncharacteristic. Thickness of the arrows represents

the proportion of homologous matches detected between defence genes from P4 or P2 and the other

different genomic contexts (thicker arrows correspond to a higher proportion of matches). The exact values

for each combination are shown in File S5. **B.** Phylogeny of the AVS_V gene, comprised of the gene found

as a defence system in a focal P2 in *Klebsiella*, and its homologs detected in either other P2s (red circles at

the tree leaves), plasmids (green circles), prophages (violet circles) or uncharacteristic genomic regions

(grey circles). The genes corresponding to each of the elements (or to the segments that include the

uncharacteristic genomic regions as grey circles) are shown in front of the tree, with larger plasmids

truncated for visibility. **C.** Phylogeny of the PD-T7-2-A gene, comprised of the gene found in a PD-T7-2

defence system in a focal P4 in *Klebsiella*, and its homologs detected in either other P4s (blue circles at the

tree leaves), plasmids or uncharacteristic genomic regions. The genes corresponding to each of the

elements (or to the segments that include the uncharacteristic genomic regions) are shown in front of the

tree, with larger plasmids truncated for visibility.

Cryptic genomic islands have homologs of defence genes from P4 and P2

A large fraction of homologs of P4s or P2s identifiable defence systems are found in

bacterial underline{uncharacteristic genomic regions} (UGRs) that do not correspond to known

MGEs. Their genomic context (20 genes each side) often contained tRNAs genes, which

are well-known points of integration of MGEs[38] (Fig S17A). Functional annotation with

EggNOG Mapper and PFAM revealed that some of these genes are typical from P4 or

prophages (Fig S17C and D, see blue and red dots). These regions also contained many

pseudogenes of proteins typical of MGEs (Fig S17A and B). Since these results suggested

that some UGRs could be defective satellites or phages, we systematically annotated

them using the core markers of P4 and P2 (see Methods) and PseudoFinder. In total, 745

URGs encode highly similar homologs of bacterial antiviral genes detected in P4s (Fig 6A,

File S5) and many of them (27%) have at least one P4-specific marker(s). All 745 UGRs

have at least one pseudogene, and about half have pseudogenes (typically intergenic)

homologous to a P4 core gene. Conversely, in the 775 UGRs that encode highly similar

homologs of bacterial defence genes detected in P2s, P2-like markers are found in ca.

20% of them. Although almost all (>98%) of these UGRs have at least one pseudogene,

only 5% of them correspond to a P2-like core gene. The above observations remain

qualitatively similar when including the putative defence systems (Fig S18). UGRs with P4

or P2 core markers are thus very likely defective versions of these elements. Within them,

we found numerous pseudogenes of mobility functions, but the neighbouring defence

genes remained almost identical to those of intact elements, and are thus likely

functional (Fig 6B-C). These results mean that defective P4 and P2 either engaged in

genetic exchanges of defence systems with their functional counterparts, or are non-

mobile satellites and prophages with stabilized (and conserved) defence genes.

Moreover, and consistent with the absence of genetic exchanges between hitcher and

helper (Fig 5), homologs of defence genes in P4 and P2 are also not found in defective

elements of the opposite type.

Almost 50% of UGRs with homologs to P4 defences and 80% of the UGRs with homologs

to P2 defences lack any P4- and P2-specific markers, considering both the genes and

pseudogenes within these regions. They are thus cryptic genomic islands, completely

distinct from P4/P2, and of unknown mobility. The detection of defence homologs in such

dissimilar islands is consistent with the observed putative exchanges of P4 and P2

defences with MGEs of different types, where we do not expect to find large regions of

homology (Fig 5).

493
494
Figure 6. Quantification of pseudogenes and P4-like/P2-like markers in uncharacteristic genomic

regions with homologs to defence genes encoded by P4 and P2. A. Characterization of uncharacteristic

genomic regions (URGs) that encode homologs to defence genes found in P4 (left bars, blue) and URGs

that encode homologs to defence genes found in P2 (right bars, red). We quantified the proportion of these

URGs that encode P4/P2 markers predicted as pseudogenes by PseudoFinder (last two bars). **B.** Examples

of P4-like satellites, and the homology of their defence systems with URGs that contain incomplete (likely

defective) P4-like satellites. In each genomic comparison, the coloured arrows indicate P4-like core genes

that were detected as intergenic pseudogenes in the URGs. **C.** Examples of P2-like prophages, and the

homology of their defence systems with URGs that contain incomplete (likely defective) P2-like prophages.

In each genomic comparison, the coloured arrows indicate P2-like core genes that were detected as

intergenic pseudogenes in the genomic regions.

DISCUSSION

The systematic detection of defence genes in prophages and satellites is a strong statement on the selective importance of antiviral functions for the ecology and evolution of these MGEs. Given the prevalence of P4 and P2 in Enterobacteria[17], and the large proportion of their accessory genes that we show here to be dedicated to defence, these MGEs constitute a significant fraction the bacterial repertoire of defences. Yet, the elements' pangenome dynamics show that the sampling of their natural diversity is not yet exhausted, even when considering, for the first time, the vast diversity of P4 elements in metagenomes using Logan[18]. Our experimental validation of anti-phage activity for 13 novel systems encoded by P4 confirms that the contribution of defence functions to the pangenomes of P4 and P2, as well as to the antiviral repertoire of their bacterial hosts, will keep surprising us.

The diversity of antiviral functions in P4 and P2 is driven by events of genetic swaps at defence-specific hotspots, enlightening the underlying processes of natural selection in MGE-defence associations. A novel system will fix in a population of MGEs if it has adaptive value. Targeted phages may, over time, adapt to resist this novel system, reducing the selective advantage of retaining it in the MGE genome. It has remained unclear if this process typically runs until the defence systems become neutral (or even deleterious, e.g., due to autoimmunity[39]) or if defences are replaced beforehand. Our observation of rapid replacement of the systems and the lack of defence-associated pseudogenes at these loci suggest that recombination quickly swaps functional systems without the latter passing by a period of inactivation. Although many small intergenic regions predicted as pseudogenes are in defence loci, they are likely remnants of an intense process of recombination. What little may remain of antiviral genes in these regions lost any recognisable homology with known defences, even if they could potentially give rise to novel defence genes[40]. These findings have two implications. First, defence systems may be lost when they are still adaptive: it suffices that the novel system is more adaptive for the swap to be fixed in the population by natural selection. Second, it suggests that systems spend little time, if any, as neutral components of the MGE genome.

P4 and P2 exchange their defence systems with elements of the same type and, less

frequently, with other MGEs. Yet, they do not exchange with each other. This is puzzling

because P4 and P2 frequently co-occur within bacterial hosts, and are co-induced,

replicate and package their DNA simultaneously[41]. This should provide plenty of

opportunities for genetic exchanges[9,42], particularly since phage induction, replication

and recombination are mechanistically concomitant[43,44]. Three factors may explain this

result. First, exchanges by homologous recombination require high sequence identity

that is lacking between P2 and P4[45]. Yet, we do observe exchanges between these

elements and other unrelated MGEs, like plasmids. Second, because P4 and P2 often co-

occur in genomes, there might be selection against the presence of redundant defences

in the two elements. Different types of defences in P2 and P4 would maximize the range

of anti-phage defence of the consortium. Identical defences would be redundant, and

one set could be easily lost by genetic drift. Third, if P2 have defence systems to prevent

P4 infection, or to avoid the hijacking of P2 particles by P4, these systems might be

counter-selected in P4. Conversely, P4 might carry defence systems that increase the

fitness of P4 by decreasing the ability of P2 to use its own capsids. If they exist, such

genes are not expected to be acquired by P2 phages.

Despite the lack of exchanges between P2 and P4, we found very similar defence systems

in distinct MGEs or genomic islands, suggesting that genetic exchanges between very

different elements is possible. It is difficult to estimate the direction of these exchanges,

although in some cases the defence was very likely acquired by P4 or P2 from other

MGEs. When a phage or a satellite acquires a defence gene from a plasmid, the system

can now be exchanged more easily with other satellites or phages[46]. These exchanges

can also contribute to the flow of antiviral genes between phylogenetically distant

bacteria, if the MGEs involved in these exchanges have different host ranges. We also

found defence homologs in defective MGEs. It will be interesting to investigate if this

corresponds to their domestication, resulting in the stabilization of the defence genes in

bacterial genomes, or if defective elements still actively swap their defences with other

MGEs.

Defence genes in P4 and P2 have evolutionary histories that are strikingly different from the rest of the elements' genomes. Antiviral defences, in accordance with the idea that they are "guns for hire", change genomic contexts of MGEs as quasi-independent entities[47]. Yet, they do so in ways that resemble the change of bacterial chromosome contexts by MGEs: frequent recombination breaks genetic linkage between defence genes and the core genes of the elements, like MGEs are unlinked from the bacterial chromosome lineage[48]; defence genes integrate predictable (hotspot) locations, like MGEs in bacterial chromosomes[49] (through a presumably different mechanism); the modularity of defence systems can promote the emergence of hybrids through recombination, resembling the modularity of the evolution of prophages[50,51]; defence genes are a large fraction of the satellite and the phage's pangenomes, similar to the contribution of MGEs to bacterial pangenomes[52]; and defence genes provide accessory functions to both phages and satellites, which in turn do the same to their bacterial hosts[2]. The key difference is that defence systems are not known to encode mobilization functions. Instead, our results suggest that P2 and P4 hotspots function as factories of novel systems by recombination processes that still need to be elucidated. It also remains to clarify how mutations and natural selection act on these chimeras to eventually lead to their fixation as functional hybrid systems. In summary, our work shows that P2 and P4 can be seen as genetic platforms that accumulate, diversify, and then disseminate bacterial defence systems across microbial populations.

MATERIALS AND METHODS

Genomic and metagenomic datasets

We retrieved all the complete genomes of the NCBI non-redundant RefSeq database (ftp://ftp.ncbi.nlm.nih.gov/genomes/refseq/, last accessed in May 2023), which consists of 32799 complete bacterial genomes. The metagenomes used for the detection of P4-elements were retrieved by aligning a query set of 538 clustered P4 protein sequences to Logan v1.1 contigs using DIAMOND2[53], as described in the Logan manuscript[18]. We also retrieved from GenBank a dataset with 27022 complete bacteriophage genomes (retrieved March 2024). The lifestyle of each of these bacteriophages was inferred using Bacphlip[54].

Identification and delimitation of genomic regions of P4-like satellites or P2-like prophages in complete bacterial genomes

P4-like satellites were initially detected from the RefSeq bacterial genomes as described previously[17]. This resulted in 3437 P4-like genomes of Type A or B. All prophages encoded by bacterial genomes were detected using geNomad (version 1.5.2, default parameters)[36]. In total, we detected 88288 prophages in the RefSeq bacterial genomes. We then designed a MacSyFinder[55] model corresponding to a prototypical P2-like prophage. The components of the model include the GpQ-L operon, which is conserved in P2-like prophages[56], as well as an integrase and a primase (GpA). This model was directly applied to the prophages detected by geNomad, resulting in 8165 P2-like prophages of Type A, B or C. The latter two types of P2-like prophages are often missing either the integrase, or the primase. We speculate that these two components are either less conserved than the profiles used for the GpQ-L operon, or excluded by geNomad during the predicted prophage region. The proteomes of these elements were retrieved by extracting all the genes within the farthest core components. This general set of non-att delimited 3437 P4-like satellites and 8165 P2-like prophages constitutes what we name, respectively, the broad P4* and P2* categories.

A stricter definition of the genomic regions corresponding to P4 and P2 was achieved by

identifying *att* sites (attL and attR) at the borders of these elements in three bacterial

species, *Escherichia coli* (Ec), *Klebsiella pneumoniae* (Kp), and *Salmonella enterica* (Se),

which constitute the most abundant hosts of P4 and P2 elements[17]. To do so, we started

by identifying genomic spots containing P2 or P4 core genes across bacterial

chromosomes. The genomic positions of P4 and P2 elements described above were

mapped relative to persistent genes in the three host species (Ec: 2527 genomes, Kp:

1510 genomes, and Se:1256 genomes) to facilitate MGE delimitation and comparison

across genomes. The pangenome of each species (i.e., the full bacterial species gene

repertoire, not to be confused with the P2/P4 pangenome) was computed using the

pangenome module of PanACoTA v1.4.1-dev[57], applying a minimum sequence identity

and coverage threshold of 80%. Persistent gene families—those present in a single copy

in at least 90% of genomes—were extracted using the corepers module of PanACoTA.

Intervals were defined as the regions between two consecutive persistent genes within

genomes. P2 and P4 core genes are not persistent across species and are confined to

these intervals, providing an upper bound for their genomic positions. In total, 6,008

intervals containing P2 or P4 core genes (excluding integrases) were identified across the

three species (2258 in *Ec*, 2,66 in *Kp*, and 1384 in *Se*). Intervals from genomes of the same

species were grouped into genomic spots if flanked by the same pair of persistent gene

families, allowing comparison of equivalent regions across genomes. The number of

genomic spots occupied by P2 and P4 was generally low in each species (30 in *Ec*, 16 in

Kp, and 21 in *Se*).

P2 and P4 integrate the bacterial chromosome using a Tyrosine recombinase (integrase),

which mediates site-specific recombination between the host *attB* site and the MGE *attP*

site. Integration duplicates part of *attB* at the element edges, generating *attL* and *attR*[58],

which can be identified by sequence similarity within genomic spots. For each spot

described above, the shortest interval observed in the species was extracted as the

ancestral MGE-free state containing *attB*; when this assumption does not hold, *att* sites

cannot be reliably identified. Each such interval was then compared using sequence

alignments with longer intervals containing P2 or P4 elements, expected to harbour short

repeats of *attB* at their edges. To ensure at least two significant sequence alignments per

comparison and to account for possible integrations within persistent genes, intervals
were extracted including the flanking persistent genes and oriented so that these genes
point in the same direction, restricting the search to the positive strand for computational
efficiency. Sequence similarity between the shortest interval (query, *attB*) and each
longer interval (subject, candidate *attL/attR*) within the same spot was screened using
BLASTn of BLAST 2.9.0+[59] (-evalue 0.001 -strand plus). Alignments with ≥80% identity
were retained, and redundant hits fully contained within longer alignments were
discarded. Pairs of hits on the subject aligning to overlapping regions of the query were
interpreted as duplications associated with integration. The overlapping region on the
query defines the putative *attB* site, and the corresponding regions on the subject define
attL and *attR*. Pairs of *att* sites flanking all P4/P2 core genes were then selected to
precisely predict element boundaries, resulting in the delimitation of 2175 elements in
Ec, 2230 in *Kp*, and 1357 in *Se,* corresponding to 96 % of intervals containing P2/P4 core
genes. This approach does not rely on genomic context, such as the presence or position
of tyrosine recombinase genes (unlike methods such as TIGER[60]). Although integrases
were not considered while detecting P4 and P2 spots, they are present in nearly all
delimited elements – with only 76 exceptions - and in most cases (5638), the integrase is
located within 2kb of an element boundary, demonstrating the robustness of our method.
Att site pairs (sequences), element size, and integrase presence are provided in File S6.
674
From the P4/P2 intervals with an *att* identification, we further filtered the elements
according to their size (<20kb for P4 and <50kb for P2) and discarded P4 that were lacking
more than two core genes (i.e, only considered P4 of Type A or B, according to
SatelliteFinder). Delimiting and filtering the elements in the 3 main bacterial species,
resulted in a total of 2168 P4-like satellites and 3574 P2-like prophages originating from
complete bacterial genomes.
681
682 Identification and delimitation of genomic regions of metagenomes that correspond to
683 P4-like satellites
684
685 To detect P4-like satellites in the metagenomic contigs, the latter were initially processed
686 as described previously in[18]. The 7 P4 core genes were detected in a set of 210895 "P4-

687 like contigs". An initial step of dereplication of data was performed in these full contigs,

688 using MMSeqs2[61] with an identity cutoff of 0.999 and a (default) minimum of 80%

689 coverage. This resulted in 16205 unique "P4-like contigs". The collection of *att* sites

690 detected for the P4-like satellites in the 3 main bacterial species (described above) were

691 then used to subsequently delimit the regions in each contig that correspond to a P4-like

692 satellite. Briefly, we used blastn with the *blastn-short* option to detect the *att* sites in each

693 contig, retaining matches with ≥ 70% identity. To consider possible variation in *att* sites,

694 we did not enforce the match with both a specific *attL* and *attR* sequence, but instead

695 searched for either one of these sequences (i.e., two *attL* or two *attR* sequences would

696 suffice). We then extracted the shortest regions that correspond to nucleic sequences

697 that are surrounded by two of these *att* sequences, requesting a minimum of 5kb and a

698 maximum of 20kb in length for the whole *att*-defined region, according to the expected

699 distribution of sizes of P4-like elements. This resulted in 6039 unique P4-like replicons.

700 To confirm that these regions correspond to the expected P4-like satellites with enough

701 components, we first used Prodigal to extract the proteins, and then used SatelliteFinder

702 to characterize the putative P4s. This resulted in a total of 6026 P4-like satellites, where

703 most sequences (6001) correspond to P4-like satellites of Type A or B (Fig S1). This

704 suggests that our delimitation of P4-like regions using the known *att*-sites, whilst likely

705 conservative and resulting in the discarding of some valid elements, is appropriate to

706 properly define the regions of P4-like satellites.

708 Dereplication of P4 and P2 genomes

710 To dereplicate the final dataset used for the analyses, we merged the nucleotide

711 sequences from the *att*-defined P4-like satellites *in E. coli*, *K. pneumoniae* and *S. enterica*

712 with all the *att*-defined P4-like contigs from the metagenomes. We then clustered these

713 replicons using MMSeqs2 with the same identity cutoff as before (0.999) but with a

714 stricter 99% coverage threshold, to avoid the merging of elements that have only one

715 gene that is distinct. A similar approach was done for the *att*-defined P2-like prophages

716 in the 3 main bacterial species. This resulted in a final dataset of 3085 dereplicated P4-

717 like satellites and 1511 dereplicated P2-like prophages, which we use for all the analysis

718 described in the manuscript.

Identification of defence genes and systems

The detection of defence genes and systems were done with DefenseFinder version 2.0.0 (defense-finder-models database v2.02)[19], using the default options. We included the flag -a to detect the anti-defence systems as well, albeit none were detected in P4-like satellites, and very few in P2-like prophage (see main text for details). We used PADLOC version 2.0.0[20] to complement the detection of complete defence systems by DefenseFinder. To retrieve genes associated with putative defence systems, we extracted the best hit for each gene from the *.hmmer.tsv output file of DefenseFinder.

To identify the gene repertoire of defence systems in the bacterial hosts of P4 (restricted to P4 detected in complete bacterial genomes) and P2 elements, we identified the defence systems in the bacterial genomes of the three main species (*E. coli, K. pneumoniae* and *S. enterica*) that encoded at least one P4 and/or one P2, using the same approaches as above. The defence gene families in bacterial hosts were computed by collecting, for each species, the genes of the defence systems (either in DefenseFinder or PADLOC) and clustering all the corresponding protein sequences using MMSeqs2 (again, the clustering was performed per each of the three species, and for each dataset, DefenseFinder or PADLOC individually). A defence gene family was considered to be associated with P4 or with P2 if at least one of the genes of the family is within an *att*-defined P4 or P2 genome of the three main species.

Calculation of P4 and P2 pangenome accumulation curves

We first clustered at 40% identity all proteins of the dereplicated P4 satellites, and all proteins of the dereplicated P2 prophages, using the MMSeq2 cluster function, with the parameters cluster_mode 1 and the default coverage. Then, for each category of elements (P4 or P2), we randomized the order of the elements and calculated the number of new gene families added by each consecutive element, annotating whether their representative sequences correspond to a defence-associated gene or a core gene of the respective element.

Calculation of genomic similarity between P4-like satellites and between P2-like prophages

752
753

We searched for sequence similarity between all proteins of satellites using MMseqs2
with the sensitivity parameter set at 7.5 to align all versus all proteins. For specific
datasets (core or defence genes), the datasets were limited to the genes that were
annotated with the specific core markers or identified as defence-associated. We then
performed the combinations between different datasets as well (e.g., defence vs core or
defence vs whole proteome). The outputs of MMSeqs2 were converted to the blast format
and we kept for analysis the hits respecting the following thresholds: e-value lower than
0.0001, at least 35% identity, and a coverage of at least 50% of the proteins (since
MMSeqs2 searches for local similarity). The hits were used to retrieve the bi-directional
best hits between pair of genomes, which were in turn used to compute a score of gene
repertoire relatedness weighted by sequence identity:

766
$$wGRR = \frac{\sum_i^p id(A_i, B_i)}{\min(A, B)}$$

where A_i and B_i is the pair *I* of homologous proteins present in *A* and *B*, id(A_i,B_i) is their
sequence identity in the local alignment, and min(*A*,*B*) is the number of proteins of the
genome encoding the fewest proteins (*A* or *B*). wGRR is the fraction of bi-directional best
hits between two genomes weighted by the sequence identity of the homologs. It varies
between zero (no bi-directional best hits) and one (all genes of the smallest genome have
an identical homolog in the largest genome). Hence, a wGRR=1 means that the two
elements are identical (or one is entirely contained within the other), whereas wGRR=0
means the elements have no homologous protein-coding genes. wGRR integrates
information on the frequency of homologs and sequence identity. For example, when the
smallest genome has 10 proteins, a wGRR of 0.2 can result from two identical homologs
or five homologs each with a lower sequence similarity (40%). All wGRR calculations in
the manuscript were obtained through GRIS version 2
(https://gitlab.pasteur.fr/jgugliel/gris).

Calculation of the pairwise similarity of core genes of P4 or P2

To calculate the pairwise similarity between each core gene, we first collected and concatenated in a fasta file all the individual core genes of P4-like satellites (7 core genes) and P2-like prophages (7 core genes). Each of these files with concatenated proteins was simultaneously used as both query and target database in Diamond BlastP, in order to retrieve the best pairwise matches. The parameters used were the following: --query-cover 50, --ultra-sensitive and --max-target-seqs 0. We then retrieved the best matches for each core gene between each pair of elements.

Identification of pseudogenes

Pseudogenes were detected using PseudoFinder version 1.1.0[25], using the default parameters. We first generated the GenBank files from the sequences to be analysed (P4-like satellites, P2-like prophages, or uncharacteristic genomic regions of bacterial genomes). We used as a reference database the complete set of bacterial proteins from the RefSeq bacterial collection used in this study (see Bacterial genomic and metagenomic datasets). We then analysed the putative pseudogenes reported by PseudoFinder either as an existing ORFs or as an intergenic region predicted to be a pseudogene. To functionally annotate the latter, we used the ORF from the RefSeq bacterial dataset that better matches each intergenic sequenced (as reported by PseudoFinder, using BlastX). The putative pseudogenes (ORFs and intergenic) were then functionally annotated with the HMM profiles corresponding to the P4 and P2 core genes and analysed with DefenseFinder and PADLOC to detect defence-associated genes. Additionally, we annotated the pseudogenes using EggNoggMapper version 2.1.12[62], with the -m diamond parameter and with all the other parameters left as default.

Cloning of candidate defence systems

Putative defence systems, as well as unknown P4 genes, were synthesized with their native promoter and cloned into the low-copy plasmid pFR66, which carries a pSC101 origin of replication, a kanamycin resistance cassette, and a superfolder GFP gene under

the control of a pTet promoter[10]. Thus, the GFP was replaced by the candidate defence

systems, placing them under the control of the anhydrotetracycline (aTc) inducible pTet

promoter. Fifty µl of *E. coli* K-12 MG1655 chemically competent cells prepared by

rubidium chloride method were transformed by heat shock at 42°C with one µl of plasmid

carrying the candidate defence system. After 1 h of recovery at 37°C in LB (Lysogeny

broth) medium, transformants were selected on LB plates supplemented with

kananamycin (Kan) at 50 µg/ml. All constructions were verified by Nanopore sequencing.

The fragments of candidate defence genes are shown in Dataset E1, whilst the complete

plasmid sequences are shown in Dataset E2.

Phage amplification

The panel of 28 phages tested, including both classical model phages and a

representative selection of phages from the BASEL collection[63], as well as the $T5_{Mos}$

phage[64], was amplified on *E. coli* K-12 MG1655. For amplification of phage stocks from -

80°C, a frozen piece of phage stock mixed with 200 µl of an overnight culture of *E. coli* and

20 mL of warm (56°C) LB + CaCl2 5 mM + 0.5% agar was poured onto square plates (12 x

12 cm) containing LB + CaCl2 5 mM + 1% agar and incubated overnight at 37°C except for

the T7-like phages that were incubated at room temperature. The top-agar layer was

recovered with 1 mL of PBS 1X from plates on which confluent lysis was observed and

transferred into a 50 mL conical tube. The top agar was then disrupted by vortexing until

broken into small pieces. Tubes were then left to incubate 10 min at room temperature

before centrifugation at 4,000 g for 5 min. Finally, the supernatant was recovered and

filtered with a 0.45 µM filter. The phage lysates were stored at 4°C. To determine the titer

of the phage lysates, the lysates were serially diluted and spotted onto lawns of *E. coli*

MG1655.

Phage plaque assays

E. coli K-12 MG1655 strains carrying each putative defence system or the control plasmid

pFR66 were grown overnight in LB supplemented with Kan 50 µg/ml. Bacterial lawns were

prepared by mixing 200 µL of a stationary culture with 100 µl of 1 mM CaCl2 and 20 mL of

LB containing 0.5% agar, and the mixture was poured onto square plates (12 x 12 cm) of

LB supplemented with Kan 50 μg/ml with or without 0.5 μg/ml aTc. Serial dilutions of high-

titre (>10^8 pfu/mL) phage stocks were spotted onto each plate and incubated overnight

at 37°C or at room temperature for T7-like phages. The next day, plaques were counted,

and fold resistance was calculated as the number of plaques in the presence of each

system divided by the number of plaques on the control plate. When plaques were too

small to be counted individually, the highest-concentration dilution showing no visible

plaques was conservatively scored as containing a single plaque.

Machine learning detection of antiviral domains in P4-like satellites and annotation of

novel defence systems

To annotate the candidate defence systems and characterize potential antiviral domains,

sequence and structural homology analyses were performed. Sequence homology was

first assessed using Pfam-A annotation using PyHMMER[65] (v0.11), a Pyhton library

binding to HMMER[66] (v3.4). To annotate protein regions without Pfam annotations,

HHpred[67] was used with 3 iterations on Pfam-A (online version) to identify conserved

domains via profile-to-profile Hidden Markov Model (HMM) comparisons. Structural

predictions were used for both homology detection and representation. For homology

detection, 3D protein models were generated using ESM-Fold, and the resulting PDB files

were processed through the Foldseek web server[68] to identify structural homologs within

the PDB and AlphaFold databases. In the second approach, AlphaFold 3.0.1 was

employed to generate structural predictions, evaluating monomeric or dimeric

configurations (when the system has 2 genes). When possible, we annotated whether the

novel systems were predicted as potentially defensive[69] (GeneCLR hit in File S3). These

integrated structural data were used to infer the presence of specialized antiviral effector

or sensor domains by comparing the predicted folds to known defence-related protein

architectures.

Genes of unknown functions were scored by both the ESM-650M$_{DF}$ and the

geneCLR$_{DF}$ scores as defined in[69]. The methods were run on representative sequences

when clustering the database at 95% of identity. Thus, we mapped the genes of unknown

function to their representative 95 and retrieved the corresponding scores.

Detection of homologs of P4/P2 defence genes in bacterial and bacteriophage genomes

We collected all defence-associated proteins in P4-like satellites and P2-like genomes,

and then used this set with Diamond BlastP to search for homologs in the complete

RefSeq bacterial genomes, including their plasmids. We used Diamond BlastP with the

parameters --query-cover 50, --ultra-sensitive and --max-target-seqs 0. We then

retrieved all matches with at least 85% identity with the query sequence and consider

these as homologs of the matched defence genes. We did a similar analysis to search for

homologs of P4 or P2 defence-associated proteins in the dataset of 27022 bacteriophage

genomes described above.

The genomic context of the homologs detected in bacterial genomes is assigned as a

plasmid (if in a secondary replicon of the bacterial genome), a prophage (if in a region

detected as such by geNomad), a satellite (if in a region detected as such by

SatelliteFinder), a phage-plasmid (if the plasmid replicon correspond to known phage-

plasmids[37] or if geNomad detects a secondary replicon as a prophage) or an integrative

mobile element (i.e., Integrative Conjugative Elements – ICEs –, Integrative Mobilizable

Elements – IMEs – or iOriTs, if in a region detected as described in[34]). If the defence

homolog was detected in a bacterial genomic region that does not correspond to any of

the categories above, we assigned it as an "uncharacteristic genomic region".

To account for the possibility that homology of defence systems is caused by vertical

descent within the element, i.e., cases where an homolog of a P4/P2 defence gene was

found in a bacterial region that corresponds to that same P4/P2, we calculated the core

(minus the integrase) wGRR (see section above) between all P4 and P4*, and all P2 and

P2*, and discarded homologous matches with elements that had a wGRR of at least 0.95.

Phylogenetic analyses

To compute the phylogenies of P4 and P2 elements, we extracted all their respective core

proteins (except for the integrase), and aligned each set of proteins individually, using

using mafft-linsi[70] (v. 7.490, default parameters). The individual alignments of core

proteins were then concatenated in two alignments (one with the P4 core proteins,

another with the P2 core proteins). For elements where some core genes were missing, a

sequence of gaps ("-") was added. The resulting alignment was trimmed with clipkit[71] (v.

1.3.0, default parameters). We then used IQ-Tree[72] (v. 1.6.12) to build the phylogenetic

trees, with the options –bb 1000 to run the ultrafast bootstrap option with 1000 replicates

and -m TEST option for automatic model choice . The resulting tree files were visualized

and edited using the v5 webserver of iTOL[73]

The phylogenies of individual defence genes were inferred by aggregating in single fasta

files all the protein sequences corresponding to homologous matches of specific

defence genes (PD-T7-2, Lit and Avs_V). Systems with more than one gene (i.e., PD-T7-2)

were aggregated in separate files, and each gene was processed separately. The

alignments and phylogenies were assembled and visualized as described above.

Functional annotation of uncharacteristic genomic regions

Homologous matches of defence genes of P4/P2 in bacterial chromosomic regions that

do not correspond to known MGEs were defined as "uncharacteristic genomic regions".

To analyse the genomic context of these homolog genes, we extracted 20 genes upstream

and 20 genes downstream. These regions were then further annotated with

DefenseFinder[19] and PADLOC[20], P4 and P2-specific HMM markers, and EggNogMapper[62],

all with the same parameters described above in each section. We also used

PseudoFinder[25], as described above, to detect pseudogenes in these uncharacteristic

genomic regions, and the intergenic pseudogenes detected were further classified with

P4 and P2 specific HMM profiles. To maintain specificity, we excluded from the search for

core markers the integrase of P4/P2, as well as Delta and AlpA of P4, since homologs to

these components can be found in other MGEs.

Visualization of genomic regions

Genome visualization across the manuscript uses Clinker version 0.0.3[74] with the -i parameter set at 0.3 at the other parameters were left as default. The JSON output files were edited to better represent the colors of the gene clusters.

Acknowledgements

We thank all the Microbial Evolutionary Genomics Unit members for the regular discussions throughout the project. The authors also wish to acknowledge the students of the 2025 M2 Microbiology course at Institut Pasteur (Pierric Chuppa, Hugo Cibla, Léa Dunand-Sauthier, Esteban Francois, Thibault Frisch, Theana Gardais, Chiara Gouviaux-Deletraz, Dorian Joffres, Ekaterina Korotaeva, Kevin Koule, Jeane Le Floch, Nicolas Le Jalle, Laly Meyer, Laura Ochsenbein, Orane Pion, Maël Rochefort, Corentin Rohel, Alexandre Roubaud, Inès Santos and Sofia Tagkalidou), who collaborated in the preliminary assays of infection the candidate defence systems. This work used the computational and storage services (MAESTRO cluster) provided by the IT department at Institut Pasteur, Paris. We acknowledge support from the Laboratoire d'Excellence IBEID Integrative Biology of Emerging Infectious Diseases [ANR-10-LABX-62-IBEID], the Agence Nationale de Recherche [ANR 23 CE20 0046 01 TRIADE] and [ANR-20-CE12-0004 TransfoConflict] and the HORIZON programme [HORIZON-MSCA-2024-PF-01 TriSH]. This work used the computational and storage services (MAESTRO cluster) provided by the IT department at Institut Pasteur, Paris.

REFERENCES

1. Clokie, M. R. J., Millard, A. D., Letarov, A. V. & Heaphy, S. Phages in nature. *Bacteriophage* **1**, 31–45 (2011).

2. Touchon, M., Moura de Sousa, J. A. & Rocha, E. P. Embracing the enemy: the diversification of microbial gene repertoires by phage-mediated horizontal gene transfer. *Curr. Opin. Microbiol.* **38**, 66–73 (2017).

3. Shkoporov, A. N., Turkington, C. J. & Hill, C. Mutualistic interplay between bacteriophages and bacteria in the human gut. *Nat. Rev. Microbiol.* https://doi.org/10.1038/s41579-022-00755-4 (2022) doi:10.1038/s41579-022-00755-4.

4. Dedrick, R. M. *et al.* Prophage-mediated defence against viral attack and viral counter-defence. *Nat. Microbiol.* **2**, 16251 (2017).

5. Brenes, L. R. & Laub, M. T. E. coli prophages encode an arsenal of defense systems to protect against temperate phages. *Cell Host Microbe* **33**, 1004-1018.e5 (2025).

6. Beamud, B., Benz, F. & Bikard, D. Going viral: The role of mobile genetic elements in bacterial immunity. *Cell Host Microbe* **32**, 804–819 (2024).

7. Clabby, T., Tesson, F., Gaborieau, B. & Bernheim, A. Why do bacteria accumulate antiphage defence systems? *Philos. Trans. R. Soc. B Biol. Sci.* **380**, 20240082 (2025).

8. Ares-Arroyo, M., Coluzzi, C., Moura De Sousa, J. A. & Rocha, E. P. C. Hijackers, hitchhikers, or co-drivers? The mysteries of mobilizable genetic elements. *PLOS Biol.* **22**, e3002796 (2024).

9. Penadés, J. R., Seed, K. D., Chen, J., Bikard, D. & Rocha, E. P. C. Genetics, ecology and evolution of phage satellites. *Nat. Rev. Microbiol.* **23**, 410–422 (2025).

10. Rousset, F. *et al*. Phages and their satellites encode hotspots of antiviral

systems. *Cell Host Microbe* **30**, 740-753.e5 (2022).

11. Puigbò, P., Makarova, K. S., Kristensen, D. M., Wolf, Y. I. & Koonin, E. V.

Reconstruction of the evolution of microbial defense systems. *BMC Evol. Biol*. **17**, 94

(2017).

12. Hussain, F. A. *et al*. Rapid evolutionary turnover of mobile genetic elements

drives bacterial resistance to phages. *Science* **374**, 488–492 (2021).

13. Hatfull, G. F. Bacteriophage genomics. *Curr. Opin. Microbiol*. **11**, 447–453 (2008).

14. Dion, M. B., Oechslin, F. & Moineau, S. Phage diversity, genomics and phylogeny.

Nat. Rev. Microbiol. **18**, 125–138 (2020).

15. Kindt, J., Tzlil, S., Ben-Shaul, A. & Gelbart, W. M. DNA packaging and ejection

forces in bacteriophage. *Proc. Natl. Acad. Sci*. **98**, 13671–13674 (2001).

16. Leclercq, S. & Cordaux, R. Do phages efficiently shuttle transposable elements

among prokaryotes? *Evolution* **65**, 3327–3331 (2011).

17. de Sousa, J. A. M., Fillol-Salom, A., Penadés, J. R. & Rocha, E. P. C. Identification

and characterization of thousands of bacteriophage satellites across bacteria.

Nucleic Acids Res. gkad123 (2023) doi:10.1093/nar/gkad123.

18. Chikhi, R. *et al*. Logan: Planetary-Scale Genome Assembly Surveys Life's

Diversity. Preprint at https://doi.org/10.1101/2024.07.30.605881 (2024).

19. Tesson, F. *et al*. Systematic and quantitative view of the antiviral arsenal of

prokaryotes. *Nat. Commun*. **13**, 2561 (2022).

20. Payne, L. J. *et al*. PADLOC: a web server for the identification of antiviral defence

systems in microbial genomes. *Nucleic Acids Res*. **50**, W541–W550 (2022).

21.   Fillol-Salom, A. *et al*. Bacteriophages benefit from mobilizing pathogenicity

islands encoding immune systems against competitors. *Cell* **185**, 3248-3262.e20

(2022).

22.   Christie, G. E. & Calendar, R. Bacteriophage P2. *Bacteriophage* **6**, e1145782

(2016).

23.   Nilsson, A. S. & Haggård-Ljungquist, E. Evolution of P2-like phages and their

impact on bacterial evolution. *Res. Microbiol*. **158**, 311–317 (2007).

24.   Egorov, A. A., Hauryliuk, V. & Atkinson, G. C. Systematic annotation of hyper-

variability hotspots in phage genomes and plasmids. Preprint at

https://doi.org/10.1101/2024.10.15.618418 (2024).

25.   Syberg-Olsen, M. J., Garber, A. I., Keeling, P. J., McCutcheon, J. P. & Husnik, F.

Pseudofinder: Detection of Pseudogenes in Prokaryotic Genomes. *Mol. Biol. Evol*. **39**,

msac153 (2022).

26.   Mariano, G. & Blower, T. R. Conserved domains can be found across distinct

phage defence systems. *Mol. Microbiol*. **120**, 45–53 (2023).

27.   Georjon, H. & Bernheim, A. The highly diverse antiphage defence systems of

bacteria. *Nat. Rev. Microbiol*. **21**, 686–700 (2023).

28.   Haudiquet, M. *et al*. Structural basis for Lamassu-based antiviral immunity and

its evolution from DNA repair machinery. *Proc. Natl. Acad. Sci*. **122**, e2519643122

(2025).

29.   Gorbalenya, A. E. & Koonin, E. V. Endonuclease (R) subunits of type-I and type-III

restriction-modification enzymes contain a helicase-like domain. *FEBS Lett*. **291**,

277–281 (1991).

30.   Gao, L. *et al*. Diverse enzymatic activities mediate antiviral immunity in

prokaryotes. *Science* **369**, 1077–1084 (2020).

31.   Liu, H. W., Roisné-Hamelin, F. & Gruber, S. SMC-based immunity against

extrachromosomal DNA elements. *Biochem. Soc. Trans.* **51**, 1571–1583 (2023).

32.   Andersen, S. E. *et al*. Serine recombinases are conserved genetic markers of

antiphage defense systems. Preprint at https://doi.org/10.1101/2025.10.07.681051

(2025).

33.   Tan, J. M. J. *et al*. A DNA-gated molecular guard controls bacterial Hailong anti-

phage defence. *Nature* **643**, 794–800 (2025).

34.   Ares-Arroyo, M., Junker, R., Nucci, A., Touchon, M. & Rocha, E. P. C. Genomic

Islands as minimal hitchers of conjugative elements. Preprint at

https://doi.org/10.64898/2026.01.13.699239 (2026).

35.   Johnson, C. M., Harden, M. M. & Grossman, A. D. Interactions between mobile

genetic elements: An anti-phage gene in an integrative and conjugative element

protects host cells from predation by a temperate bacteriophage. *PLOS Genet.* **18**,

e1010065 (2022).

36.   Camargo, A. P. *et al*. Identification of mobile genetic elements with geNomad.

Nat. Biotechnol. **42**, 1303–1312 (2024).

37.   Pfeifer, E., Moura de Sousa, J. A., Touchon, M. & Rocha, E. P. C. Bacteria have

numerous distinctive groups of phage–plasmids with conserved phage and variable

plasmid gene repertoires. *Nucleic Acids Res.* **49**, 2655–2673 (2021).

38.   Reiter, W.-D., Palm, P. & Yeats, S. Transfer RNA genes frequently serve as

integration sites for prokaryotic genetic elements. *Nucleic Acids Res.* **17**, 1907–1914

(1989).

39.   Aframian, N., Omer Bendori, S., Hen, T., Guler, P. & Eldar, A. Expression level of

anti-phage defence systems controls a trade-off between protection range and

autoimmunity. *Nat. Microbiol.* **10**, 1954–1962 (2025).

40.   Frumkin, I., Vassallo, C. N., Chen, Y. H. & Laub, M. T. Emergence of antiphage

functions from random sequence libraries reveals mechanisms of gene birth. *Proc.*

Natl. Acad. Sci. **122**, e2513255122 (2025).

41.   Christie, G. E. & Calendar, R. INTERACTIONS BETWEEN SATELLITE

BACTERIOPHAGE P4 AND ITS HELPERS. *Annu. Rev. Genet.* **24**, 465–490 (1990).

42.   Lindqvist, B. H., Dehò, G. & Calendar, R. Mechanisms of genome propagation

and helper exploitation by satellite phage P4. *Microbiol. Rev.* **57**, 683–702 (1993).

43.   Smith, G. R. Chi hotspots of generalized recombination. *Cell* **34**, 709–710 (1983).

44.   Mosig, G. Recombination and recombination-dependent DNA replication in

bacteriophage T4. *Annu. Rev. Genet.* **32**, 379–413 (1998).

45.   Vulić, M., Dionisio, F., Taddei, F. & Radman, M. Molecular keys to speciation: DNA

polymorphism and the control of genetic exchange in enterobacteria. *Proc. Natl.*

Acad. Sci. **94**, 9763–9767 (1997).

46.   Moura de Sousa, J. A., Pfeifer, E., Touchon, M. & Rocha, E. P. C. Causes and

Consequences of Bacteriophage Diversification via Genetic Exchanges across

Lifestyles and Bacterial Taxa. *Mol. Biol. Evol.* **38**, 2497–2512 (2021).

47.   Koonin, E. V., Makarova, K. S., Wolf, Y. I. & Krupovic, M. Evolutionary

entanglement of mobile genetic elements and host defence systems: guns for hire.

Nat. Rev. Genet. **21**, 119–131 (2020).

48.   Sheinman, M. *et al.* Identical sequences found in distant genomes reveal

frequent horizontal transfer across the bacterial domain. *eLife* **10**, e62719 (2021).

49.   Oliveira, P. H., Touchon, M., Cury, J. & Rocha, E. P. C. The chromosomal

organization of horizontal gene transfer in bacteria. *Nat. Commun.* **8**, 841 (2017).

50.   Baker, J., Limberger, R., Schneider, S. J. & Campbell, A. Recombination and

modular exchange in the genesis of new lambdoid phages. *New Biol.* **3**, 297–308

(1991).

51.   Dragoš, A. *et al.* Pervasive prophage recombination occurs during evolution of

spore-forming *Bacilli*. *ISME J.* **15**, 1344–1358 (2021).

52.   Touchon, M. *et al.* Phylogenetic background and habitat drive the genetic

diversification of Escherichia coli. *PLOS Genet.* **16**, e1008866 (2020).

53.   Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life

scale using DIAMOND. *Nat. Methods* **18**, 366–368 (2021).

54.   Hockenberry, A. J. & Wilke, C. O. BACPHLIP: predicting bacteriophage lifestyle

from conserved protein domains. *PeerJ* **9**, e11396 (2021).

55.   Néron, B. *et al.* MacSyFinder v2: Improved modelling and search engine to

identify molecular systems in genomes. *Peer Community J.* **3**, e28 (2023).

56.   Casjens, S. R. & Grose, J. H. Contributions of P2- and P22-like prophages to

understanding the enormous diversity and abundance of tailed bacteriophages.

Virology **496**, 255–276 (2016).

57.   Perrin, A. & Rocha, E. P. C. PanACoTA: a modular tool for massive microbial

comparative genomics. *NAR Genomics Bioinforma.* **3**, lqaa106 (2021).

58.   Rajeev, L., Malanowska, K. & Gardner, J. F. Challenging a paradigm: the role of

DNA homology in tyrosine recombinase reactions. *Microbiol. Mol. Biol. Rev. MMBR*

73, 300–309 (2009).

59.   Camacho, C. *et al*. BLAST+: architecture and applications. *BMC Bioinformatics*

10, 421 (2009).

60.   Mageeney, C. M. *et al*. New candidates for regulated gene integrity revealed

through precise mapping of integrative genetic elements. *Nucleic Acids Res*. **48**,

4052–4065 (2020).

61.   Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence

searching for the analysis of massive data sets. *Nat. Biotechnol*. **35**, 1026–1028

(2017).

62.   Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J.

eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain

Prediction at the Metagenomic Scale. *Mol. Biol. Evol*. **38**, 5825–5829 (2021).

63.   Maffei, E. *et al*. Systematic exploration of Escherichia coli phage–host

interactions with the BASEL phage collection. *PLOS Biol*. **19**, e3001424 (2021).

64.   Burman, N. *et al*. A virally encoded tRNA neutralizes the PARIS antiviral defence

system. *Nature* **634**, 424–431 (2024).

65.   Larralde, M. & Zeller, G. PyHMMER: a Python library binding to HMMER for

efficient sequence analysis. *Bioinformatics* **39**, btad214 (2023).

66.   Eddy, S. R. Accelerated Profile HMM Searches. *PLoS Comput. Biol*. **7**, e1002195

(2011).

67.   Gabler, F. *et al*. Protein Sequence Analysis Using the MPI Bioinformatics Toolkit.

Curr. Protoc. Bioinforma. **72**, e108 (2020).

68.   Van Kempen, M. *et al*. Fast and accurate protein structure search with Foldseek.

Nat. Biotechnol. **42**, 243–246 (2024).

69.   Mordret, E. *et al.* Protein and genomic language models chart a vast landscape

of antiphage defenses. Preprint at https://doi.org/10.1101/2025.01.08.631966 (2025).

70.   Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software

Version 7: Improvements in Performance and Usability. *Mol. Biol. Evol.* **30**, 772–780

(2013).

71.   Steenwyk, J. L., Buida, T. J., Li, Y., Shen, X.-X. & Rokas, A. ClipKIT: A multiple

sequence alignment trimming software for accurate phylogenomic inference. *PLOS*

Biol. **18**, e3001007 (2020).

72.   Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and

Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol.*

Biol. Evol. **32**, 268–274 (2015).

73.   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v5: an online tool for

phylogenetic tree display and annotation. *Nucleic Acids Res.* **49**, W293–W296 (2021).

74.   Gilchrist, C. L. M. & Chooi, Y.-H. clinker & clustermap.js: automatic generation of

gene cluster comparison figures. *Bioinformatics* **37**, 2473–2475 (2021).

1142