Likely role of promoter reconstitution in Mpr-mediated D29 resistance by *Mycobacterium smegmatis*

Buhari Yusuf, [1,2,3,4] Yanan Ju, [1,2,3,5] Biao Zhou, [6] Abdul Malik, [1,2,3,4] Md Shah Alam, [1,2,3,4] Lijie Li, [1,2,3,5] Haftay Abraha Tadesse, [1,2,3,4] Belachew Aweke Mulu, [1,2,3,4] Cuiting Fang, [1,2,3,7,8,9] Xirong Tian, [1,2,3,7,9] Hongyi Chen[10], Li Wan,[1,2,3] , Liqiang Feng,[1,2,3] Xiaoli Xiong[1,2,3,8,9], Shuai Wang, [1,2,3] Tianyu Zhang [1,2,3,7,8,9]

[1] State Key Laboratory of Respiratory Disease, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

[2] China-New Zealand Joint Laboratory on Biomedicine and Health, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

[3] Guangdong-Hong Kong-Macao Joint Laboratory of Respiratory Infectious Diseases, Guangzhou Institutes of Biomedicine and Health, Chinese Academy of Sciences, Guangzhou 510530, China

[4] University of Chinese Academy of Sciences, Beijing 100049, China

[5] School of Life Sciences, University of Science and Technology of China, Hefei 230027, China

[6] State Key Laboratory of Respiratory Disease, Guangzhou Chest Hospital, Institute of Tuberculosis, Guangzhou Medical University, Guangzhou 510095, China

[7] Guangzhou Medical University-Guangzhou Institutes of Biomedicine and Health Joint School of Life Sciences, Guangzhou Medical University, Guangzhou 511436, China

[8] State Key Laboratory of Respiratory Disease, Guangzhou Chest Hospital, Guangzhou, China

[9] Guangzhou National Laboratory, Guangzhou 510005, China

[10] West China School of Public Health, Sichuan University, Chengdu 610041, China

Correspondence: Tianyu Zhang, zhang_tianyu@gibh.ac.cn

Abstract

The <u>m</u>ulti-copy <u>p</u>hage <u>r</u>esistance gene (*mpr*) of *M. smegmatis* is a major factor in resistance to the lytic mycobacteriophage D29. Mpr is a membrane-bound exonuclease that cleaves phage DNA post injection, hence blocking downstream stages in the phage infection cycle. The mechanism of resistance allows for adsorption, is non-abortive and independent of any mutation in the gene. Rather, it depends on overexpression of a wild type copy of the gene. However, the underlying factor behind_*mpr* overexpression in spontaneous D29-resistant mutants of *M. smegmatis* remained elusive. Here, we report that D29 infection triggers insertion sequence (IS) rearrangements, including the transposition and integration of IS6120 directly upstream of *mpr*. Mutants with IS6120 integration upstream of *mpr* show highly elevated Mpr expression. Whole genome sequence analysis reveals that IS6120 introduced a putative transcription factor-binding site and a canonical -35 promoter element at the integration site, hence reconstituting a fuller promoter (*rcp*) than the original promoter (*wtp*) at the site. Promoter reporter assays suggest that *rcp* is a far stronger promoter than *wtp*, implying that elevated *mpr* expression in D29-resistant mutants with this transposition event could be due to promoter reconstitution. While strains with this transposition event appear to grow normally, *rcp*-driven, vector-borne Mpr overexpression appears to be toxic as it barely allows for colony formation on agar plates. This study reports a previously unknown factor likely behind *mpr* regulation in *M. smegmatis*, adding to the existing knowledge of mycobacterial anti-phage defense mechanisms and guiding rational phage engineering efforts for therapeutic applications.

Keywords: *Mycobacterium smegmatis*, mycobacteriophage D29, promoter reconstitution, Mpr, phage resistance.

Introduction

Antimicrobial resistance (AMR) remains a serious public health threat, and despite efforts to develop novel therapeutics or vaccines, the threat remains ever existent. This is because the pace at which bacterial pathogens develop resistance to antimicrobials surpasses the rate at which new therapeutics are introduced into clinical use. Conventional antibiotic treatments occasionally fail due to extensive drug resistance, hence complicating disease management. In fact, it is estimated that global death toll due to bacterial AMR may surpass 39 million between 2025 and 2050 (1). For this reason, interest in phage therapy as an alternative to conventional antibiotic therapy is gaining significant traction.

Phage therapy is the use of bacteria-eating viruses (or bacteriophages) to treat bacterial infections. Bacteriophages are highly specific, relatively safe and hold significant potential as alternatives to redundant, toxic antibiotic therapy. Currently, phage therapy is only available to patients under the framework of compassionate use (2). Compassionate use of phage therapy against drug-resistant mycobacterial infections have further highlighted the significant potential of phage therapy even in complicated cases of failed antibiotic therapy (3–9). However, bacterial pathogens can develop resistance to bacteriophages, which represents a potent threat to sustainable utility of phage therapy.

Bacterial pathogens have evolved an extensive arsenal of anti-phage defense systems, including clinical bacterial isolates (10–13). The lytic mycobacteriophage D29 has demonstrated therapeutic potential in compassionate use against BCG and *Mycobacterium abscessus* infections (5,7). However, a resistance mechanism against D29 and its parent L5 already exists in the model bacterium *Mycobacterium smegmatis* (14). The multicopy phage resistance gene (*mpr*) of *M. smegmatis* is a major factor in D29 resistance. Mpr is a membrane-bound exonuclease that mediates resistance via cleavage of phage DNA post injection, which blocks downstream stages of the phage lifecycle and inhibit phage propagation (15). Mpr contains a domain of unknown function (DUF4352) that is believed to be of viral origin, suggesting that *M. smegmatis* obtained this protein during the course of its evolution with viral predators and repurposed it for anti-

phage defense (16). The protein has also been reported to aid non-homologous end joining

(NHEJ)-mediated anti-phage defense in *M. smegmatis* (17), further highlighting its relevance to

antiviral defense. However, the role of Mpr in D29 resistance is independent of any genetic

change in the gene, but rather depends on overexpression of a wild type copy of the gene. The

exact factor behind elevated Mpr expression in spontaneous D29-resistant mutants remained

unknown.

We report here that infection with D29 triggers IS rearrangements in *M. smegmatis*, including the

transposition and integration of IS6120 directly upstream of *mpr*. IS6120 introduces a canonical -

35 promoter element as well as a putative transcription factor-binding site (TFBS) for leucine-

responsive regulatory protein (Lrp), hence reconstituting a fuller promoter than the original

promoter at the integration site. Promoter reporter assays show that the reconstituted promoter

(*rcp*) is far stronger than the wild type promoter (*wtp*), which could represent a previously

unknown regulatory mechanism for the toxic protein Mpr in *M. smegmatis*.

Materials and Methods

Bacterial strains, phage and growth conditions

Cloning experiments were carried out in *E. coli* Trans1 (TransGen Biotech., China). *E. coli* was maintained in liquid or on solid Luria-Bertani medium. Selective growth of *E. coli* was carried out in the presence of kanamycin (kan, 50 µg/mL), ampicillin (amp, 100 µg/mL) or zeocin (zeo, 30 µg/mL). *M. smegmatis* mc^2 155 was maintained in Middlebrook 7H9 (Difco) medium supplemented 10% OADC, 0.2 % glycerol and 0.05% Tween 80. *M. smegmatis* was selectively grown in liquid or on solid medium in the presence of the following concentrations of antibiotics: kan (liquid, 20 µg/mL; solid, 50 µg/mL); zeo (30 µg/mL). All liquid cultures were maintained at 37 $^{\circ}$C with shaking at 200 revolutions per minute (rpm). The phage used in this study is the lytic mycobacteriophage D29. All D29 infection experiments were carried out at 37 $^{\circ}$C in the presence of 2 mM CaCl$_2$.

Phage amplification

Wild type *M. smegmatis* (Ms^{Wt}) was grown in 7H9 medium to OD$_{600}$ 0.8-1 at 37 $^{\circ}$C with shaking. The cells were pelleted and washed thrice in MP buffer, then re-suspended in 1.5 mL of the same buffer. Ten-fold serial dilutions of the phage stock were prepared in MP buffer until 10^{-4} dilution factor. A four-component mixture was prepared by adding top agar, washed Ms^{Wt} cells, D29 at 10^{-4} dilution and CaCl$_2$ at a final concentration of 2 mM. The mixture was mixed by pipetting and poured into plain 7H10 plates, spread evenly, allowed to dry and incubated at 37 $^{\circ}$C for 12-24 hours. This was followed by addition of 2.5 mL of sterile MP buffer into each plate and rocking on an orbital shaker at room temperature for 4-5 hours. The MP buffer was then recovered from the plates by pipetting, centrifuged at 3,500 ×g for 10 minutes, then passing the supernatant through a 0.22 µm filter. The filtrate containing the amplified phage was stored at 4 $^{\circ}$C.

Generation of spontaneous D29-resistant strains

Spontaneous D29-resistant strains were generated by subjecting Ms^{Wt} to four sequential rounds

of extended exposures to mycobacteriophage D29 to increase the chance of retaining resistant

clones and eliminating susceptible ones (18,19). In the first exposure cycle, bacterial cells were

grown to OD_{600} 1.0, washed three times in MP buffer and re-suspended in 1 mL of MP buffer,

infected with D29 and incubated at 37 ^{o}C for 7 hours with regular shaking. Cells from the 7-hour

incubation were collected by centrifugation at 13,000 $\times$g for 3 minutes, washed three times in

PBS-Tween 80, and re-suspended in 1 mL of the same medium. 50 uL of the washed cells were

sub-cultured in 7H9 medium containing 0.05 % Tween 80 until OD_{600} 1.0. This resulting culture

was washed three times in MP buffer, infected with D29 and incubated at 37 ^{o}C for 14 hours

with regular shaking. This continued until the fourth exposure cycle, with the incubation period

extended by 7 hours after the previous exposure cycle (21 and 28 hours for third and fourth

exposure cycles, respectively). For every infection cycle, MP buffer was used as the medium of

infection, bacterial cells were infected at MOI of 5, and it was ensured that $CaCl_2$ was added to a

final concentration of 2 mM. After the final 28-hour exposure, the cells were collected by

centrifugation, washed in PBS-Tween 80, diluted and plated to collect single colonies.

Phage susceptibility testing (PST)

Phage susceptibility testing was carried out by plaque and spot-kill assays as well as on D29-

seeded 7H10 plates. Bacterial cells were grown in 7H9 medium to OD_{600} 0.8-1. The grown cells

were pelleted by centrifugation at 3,500 $\times$g for 10 minutes. The supernatant was discarded and

the pellet was washed three times in MP buffer and re-suspended in 0.5-1 mL of the same buffer.

Ten-fold serial dilutions of D29 were prepared. For plaque assay, a soft overlay of top agar,

phage, bacterial cells and 2 mM $CaCl_2$ was prepared and poured into plain 7H10 plates in

triplicates. The plates were dried and incubated at 37 ^{o}C for 12-24 hours before enumeration of

plaque forming units. For spot-kill assay, a soft overlay of top agar, bacterial cells and $CaCl_2$ was

poured into a 7H10 plate, spread evenly and allowed to dry. Two microliters of ten-fold serial

dilutions of the phage were spotted on the dried soft overlay, allowed to dry, and incubated at 37

^{o}C for 12-24 hours. For PST on phage-seeded plates, 2 μL of ten-fold serial dilutions of the

bacteria was spotted on 7H10 plate seeded with $10^7 - 10^9$ plaque forming units per milliliter

(PFU/mL) of D29, allowed to dry, and incubated at 37 ^{o}C for 3-5 days. Serial dilution of the

bacteria was carried out in MP buffer containing 2 mM $CaCl_2$, and 7H10 plate containing no

phage was used as a negative control.

Phage adsorption assay

Phage adsorption assay was carried out as described previously (19,20). Bacterial cells were

grown to mid log phase, washed three times and re-suspended in MP buffer. The cells were

diluted in MP buffer to OD_{600} 0.3. Ten-fold serial dilutions of D29 were prepared in MP buffer

until 10^{-3}. To determine terminal phage adsorption, 30 µL of 10^{-3} dilution of the phage was added

to 200 µL aliquot of diluted bacteria, and $CaCl_2$ was added to a final concentration of 2 mM. The

mixture was incubated at 37 $^{\circ}$C with shaking for 60 minutes. Simultaneously, same quantity of

same phage inoculum was added to 200 µL of sterile MP buffer, 2 mM $CaCl_2$ was added, mixed,

and immediately centrifuged at 5000 ×g for 3 minutes. The supernatant was collected, diluted

and used for plaque assay on MP buffer-washed Ms^{Wt}. After the 60 minutes shaking of phage-

bacteria mixture, the mixture was also centrifuged at 5000 ×g for 3 minutes, the supernatant was

collected, diluted and used for plaque assay on MP buffer-washed Ms^{Wt}. Plates were allowed to

dry, incubated at 37 $^{\circ}$C for 24 hours before enumeration of plaque forming units. Plates were

prepared in triplicates, and data was presented as percentage of adsorbed phage. For

determination of phage adsorption rates, same phage-bacteria mixtures and incubation conditions

were prepared for plaque assay at 0-, 30- and 60-minute time points. Collected data was

presented as percentage of unadsorbed phage.

Colony morphology and spot inhibition by kanamycin and zeocin

186   All strains were grown in 7H9 medium to OD_{600} 0.8-1 and diluted to OD_{600} 0.6. Two microliters

187   from diluted culture of each strain was spotted on plain 7H10 plate and allowed to dry. For

188   susceptibility testing to kan (50 µg/mL) or zeo (30 µg/mL), the cells were spotted on 7H10 plates

189   containing indicated concentration of either drug. Dried plates were incubated at 37 $^{\circ}$C for 3-4

190   days.

Whole genome sequencing (WGS)

192   Genomic DNA (gDNA) extraction and sequencing were carried out by Shanghai Jingnuo

193   Biotechnology Co., Ltd. (Shanghai, China). A modified Cetyltrimethylammonium bromide

194   (CTAB) method was used for extraction of gDNA (21). Extracted gDNA was quantified using

Qubit 2.0 Fluorometer (ThermoFisher, USA), followed by library construction using Hieff NGS® OnePot Flash DNA Library Prep Kit (Yeasen, China) by enzymatic fragmentation of gDNA, end repair, tail adenylation, adapter ligation, purification and PCR amplification. The library was initially quantified using Qubit 2.0 and diluted. Fragment sizes were verified using Agilent 2100 bioanalyzer (Agilent, USA) and effective library concentration was quantified again by qPCR. Libraries were pooled onto a flow cell, clustered using cBOT and sequenced using the NovaSeq 6000 high-throughput sequencing platform (Illumina, Inc., USA).

Bioinformatics

Next generation sequencing (NGS) data was analyzed on the GalaxyEU bioinformatics server using default settings (22). Raw sequence reads were checked for quality using FASTQC (23), and undesirable reads/reads of low quality were trimmed using FASTP (24). Mapping to the reference genome (*M. smegmatis* mc^2 155, NCBI accession number CP000480.1) and variant calling were achieved using Snippy (25), and the output data was exported as VCF and tab-separated files. All detected variants were verified by Sanger sequencing.

RNA extraction and RT-qPCR

RNA extraction was carried out according to manufacturer instructions using Magen HiPure Bacterial RNA Mini Kit (Magen, China). For bacterial growth to detect mRNA levels in the presence and absence of D29 infection, the strains were first washed thrice in 7H9 medium without Tween 80, sub-cultured in 7H9 (without Tween 80) containing 2 mM CaCl$_2$ with/without D29 at MOI 5. The strains were then washed thrice in PBS-Tween 80 prior to RNA extraction. Extracted RNA was reverse transcribed using ExonScript All-in-One RT SuperMix with DNase (EXONGEN, China) according to manufacturer instructions. The RT-qPCR reaction was prepared and run according to manufacturer instructions using Taq Pro Universal SYBR qPCR Master Mix (Vazyme Biotech., China). All reactions were run in C1000 Touch^{TM} Thermal Cycler (Bio-Rad, USA). SigA was used as control, and all wild type/untreated group expressions were normalized to 1, and the treated/test group expression was presented as fold change. Expression levels (fold) were represented as bar graphs using GraphPad Prism version 9.0.0 (GraphPad, USA).

IS transposition analysis

Sequences of the different IS elements encoded by *M. smegmatis* genome were obtained from

ISFinder database (26). ISMapper (27) was used to map out the different IS integration sites in

raw sequence reads of each sequenced D29-resistant mutant. Briefly, paired end Illumina reads

of a D29-resistant mutant, a FASTA file of the IS query sequence and a genbank file of the

reference genome (*M. smegmatis* mc^2 155, NCBI accession number CP000480.1) were used as

input files. The paired end Illumina reads were mapped to the IS query sequence using BWA-

MEM (28), which generates a SAM file as output. Sequences directly flanking the IS query (i.e.

pairs of unmapped reads that mapped to the end of the IS query sequence) as well as soft clipped

regions (reads containing sequences that mapped to the IS query and extend into neighboring

sequence) were extracted from the SAM file using SAMtools (29) and SAMBLASTER (30)

respectively. The extracted sequences were converted to FASTQ format using BEDTools (31)

and filtered using BioPython to extract soft clipped read portions that are 5-30 bp in length. The

resulting sequences were sorted into left and right flanking sequences and separately mapped to

the reference genome using BWA-MEM, and BEDTools was used to determine the exact

position of IS integration in the D29-resistant mutant. All detected IS integrations were verified

by PCR.

Promoter reporter assay

Lawn appearance

Strains were grown in 7H9 medium to OD_{600} 0.8-1, then diluted to OD_{600} 0.6. Ten microliters

from diluted culture of each strain was streaked on a plain 7H10 plate. The plate was dried and

incubated at 37 ^{o}C until appearance of clear bacterial lawn.

Live fluorescence

Strains were grown in 7H9 medium to OD_{600} 0.8-1, then diluted to OD_{600} 0.6. Two-fold serial

dilutions of the diluted cultures were prepared in triplicates for each strain in a 96-well

microplate to obtain OD_{600} 0.6, 0.3, 0.15, 0.075, 0.0375 and 0.01875. Fluorescence was

quantified in FlexStation 3 Multi-Mode Microplate Reader (Molecular Devices, USA) at

excitation and emission wavelengths of 488 nm and 520 nm respectively. Quantification data for

fluorescent strains were normalized by subtracting Ms^{Wt} fluorescence from that of each strain,

and the results were plotted as bar graphs in GraphPad Prism.

Flow cytometry

Flow cytometry was carried out as previously described (32) with modification. The strains were grown in 7H9 medium to OD_{600} 0.8-1. Two milliliters of each strain was pelleted, washed twice in PBS-Tween 80, and fixed in 1 mL of 4 % glutaraldehyde in PBS-Tween 80 at room temperature for 30 minutes. The fixed cells were pelleted, re-suspended in 2 mL of sterile PBS, and passed through a 40 µm filter prior to flow cytometry using a CytoFLEX flow cytometer (Beckman Coulter, USA). A minimum of 10,000 cells were selected to count in focus GFP events for each strain. Flow cytometry data was analyzed using FlowJo version 10.8.1.

Confocal microscopy

Strains were grown in 7H9 medium to OD_{600} 0.8-1. One milliliter of each strain was pelleted and washed twice in PBS-Tween 80. The supernatant was discarded, the cells were re-suspended in 1 mL of 4 % glutaraldehyde in PBS-Tween 80 and allowed to stand at room temperature for 30 minutes. The fixed cells were pelleted and re-suspended in 200 µL of PBS-Tween 80. For each strain, 5 µL of re-suspended fixed cells as well as a 10-fold serial dilution were mounted on a microscope slide and covered with a coverslip, then observed under a Zeiss LSM 800 Confocal Laser Scanning Microscope (Zeiss, Germany).

Drug susceptibility testing (DST)

Drug susceptibility testing was carried out by broth microdilution. Bacterial strains were cultured in 7H9 medium at 37 $^{\circ}$C with shaking until OD_{600} 0.8-1, then diluted to a cell density of ~10^5 colony forming units per milliliter (CFU/mL) in 7H9 without Tween 80. Bacterial cultures in two-fold serial dilutions of each drug as well as a plain control were prepared in 96-well microplates. The plates were incubated at 37 $^{\circ}$C for 3 – 5 days. Minimum inhibitory concentration (MIC) was defined as the concentration at which visible bacterial growth was inhibited.

Results

Spontaneous D29 resistance in *M. smegmatis*

We screened 137 colonies of *M. smegmatis* for D29 resistance after multiple cycles of exposure, and a vast majority of them (82.5 %, n = 113) were classified as D29-resistant isolates. We then selected and purified 24 colonies, re-screened them for D29 resistance and sequenced their genomes. Of these 24 isolates, two strains, A72.1 and 2.6.1, particularly stand out for their high level of resistance to D29. Both strains appear to manifest no obvious change in surface morphology, but exhibit high level of resistance when screened by spot-kill and plaque assays (Figure 1A, B), which suggests impaired D29 infection efficiency against A72.1 and 2.6.1. Notably, greater susceptibility is seen at higher D29 concentration (Figure 1A), possibly suggesting dose-dependent resistance. Phage adsorption assay indicates little to no effect on D29 adsorption to both strains, suggesting that the mechanism of resistance in both strains occurs at stage(s) beyond phage adsorption (Figure 1C). The D29-resistant mutants A72.1 and 2.6.1 sustained point mutations and two IS transposition events each, including IS6120 integration directly upstream of *mpr* in both mutants.

D29 infection triggers IS rearrangements in *M. smegmatis*

It is likely that other genetic factors alone or in combination with the identified mutations may be responsible for D29 resistance phenotypes of the mutants (20). For this reason, we carried out IS transposition analysis on raw FASTQ data obtained for each of the mutants as well as Ms[Wt]. We first collected the sequences of all known IS elements encoded by the *M. smegmatis* genome

from ISFinder database (26) and determined their prevalence across the genome (Table S1,

Figure S1). We then identified all IS rearrangements that were present in the mutants but not in

Ms^{Wt}, then verified them by PCR alongside the Ms^{Wt} control (Figure 2).

A total of 10 IS transposition events were detected in seven D29-resistant mutants, where 4

mutants had single event each and 3 had double. There had been six IS6120, three ISMsm1 and

one IS1096 transposition events in the mutants, in which the IS integrations took place in six

coding and three non-coding regions. A72.1 and 2.6.1 – the two strains that manifested

incredibly higher level of D29 resistance compared to the other mutants – appear to have

sustained two IS transposition events each, making them strains of particular interest (Figure 3).

Details about the integration position and protein products of the affected genes are outlined in

Table S2.

A72.1 and 2.6.1 show elevated Mpr expression

The two mutants A72.1 and 2.6.1 sustained two transposition events each. ISMsm1 integrated in

the intergenic region between *MSMEG_2105* and *MSMEG_2104* in A72.1, but integrated in

MSMEG_5393 in 2.6.1. However, in both strains, IS6120 integrated within the 168 bp intergenic

region between *mpr* and *MSMEG_1237*, just upstream of *mpr*. Mpr-MSMEG_1237 interaction

map on STRING database suggests possible co-expression (Figure 4A). We therefore quantified

the expression of *mpr* and *MSMEG_1237* to determine if any of the genes had been affected by

IS6120 integration, using Ms^{Wt} and B6.1 as control strains. Both A72.1 and 2.6.1 appear to

manifest highly elevated Mpr expression, whereas *MSMEG_1237* expression remained similar

among the strains (Figure 4B). We also simultaneously quantified mRNA levels at the 168 bp

intergenic region (*MSGEG_1236-1237IR*) in both A72.1 and 2.6.1 as well as the control strains

to detect any variations. The mRNA levels at this region don't seem to differ much, and identical

DNA sizes were seen on agarose gel for the four strains after qPCR amplification (data not

shown).

IS6120 integration upstream of *mpr* introduces a putative transcription factor-binding site
(TFBS) and a canonical -35 promoter element

337   There is a 168 bp region between *mpr* and *MSMEG_1237*, and integration of IS6120 at this

338   region appears to primarily affect the expression of *mpr*, not *MSMEG_1237* (Figure 4). In

339 addition, IS6120 has the same orientation as *mpr* at the integration site, which raises the question

340 of whether IS6120 integration has had some regulatory influence over *mpr*. One way this could

341 happen is by promoter reconstitution, which inspired us to analyze the intergenic sequence prior

342 to IS6120 integration (or <u>wild</u> <u>type</u> <u>seq</u>uence, *wtseq*) and that after integration (or <u>rec</u>onstituted

343 <u>seq</u>uence, *rcseq*) for possible presence of new promoter/regulatory element(s). We used four

344 promoter prediction programs, and each predicted at least one complete promoter in *rcseq*. One

345 program, Sima70Pred, predicted three promoters in *wtseq* and fourteen in *rcseq*, but the

346 remaining programs predicted no promoters in *wtseq* (Table S3). All promoters predicted by all

347 programs in *rcseq* have the sequence "**<u>TTGACAA</u>T**TCCTCgtcccgag<u>ttgtacgtt</u>" in common,

348 which contains a putative TFBS for leucine-responsive regulatory protein (Lrp) as well as -35

349 and -10 promoter elements (Figure 5). The TFBS and the canonical -35 promoter element were

350 introduced at the site by IS6120 integration, implying that integration of IS6120 at the site

351 brought in these two elements to reconstitute a more complete promoter than that in *wtseq*.

352 **IS6120 integration reconstitutes a stronger promoter upstream of *mpr***

353 To verify the activity of the <u>rec</u>onstituted <u>p</u>romoter (*rcp*), we carried out GFP reporter assays by

354 constructing GFP-overexpressing strains using *rcp* or *wtp* (<u>w</u>ild <u>t</u>ype <u>p</u>romoter). We constructed

355 Ms$^{wtp\text{-}GFP}$ by transforming Ms^{Wt} with pMV-*wtp*-GFP, a GFP overexpression vector in which

356 *hsp60* promoter was replaced by *wtp* (the entire 168 bp region between *mpr* and *MSMEG_1237*,

357 amplified from Ms^{Wt}). Similarly, we constructed Ms$^{rcp\text{-}GFP}$ in which the *hsp60* promoter was

358 replaced by *rcp* (a region of equal length directly upstream of *mpr*, amplified from A72.1).

359 Ms^{GFP}, a strain overexpressing GFP from the *hsp60* promoter, was used as a control.

360 The first evidence of promoter activity manifested at colony level after transformation of bacteria

361 with the respective vectors. Ms^{GFP} and Ms$^{rcp\text{-}GFP}$ colonies manifested the characteristic green

362 color of GFP. Lawn of Ms$^{rcp\text{-}GFP}$ also shows the characteristic green coloration on plain 7H10

363 plates (Figure 6A). Though Ms$^{rcp\text{-}GFP}$ produces lower live fluorescence than Ms^{GFP}, it produces

364 far greater fluorescence than Ms$^{wtp\text{-}GFP}$, suggesting that introduction of the new elements by

365 IS6120 significantly increased the strength of the promoter at the integration site (Figure 6B).

366 We also counted GFP events for each of the strains by flow cytometry, and the number of GFP

367 events detected for Ms$^{rcp\text{-}GFP}$ was clearly far greater than that for Ms$^{wtp\text{-}GFP}$, which constitutes

368 further evidence of superior promoter activity (Figure 7). Finally, confocal micrographs

corroborate the flow cytometry data, showing that *wtp* drives significantly lower GFP expression

than *rcp*, and that the GFP signals produced by Ms$^{rcp\text{-GFP}}$ and Ms^{GFP} are somewhat identical

(Figure 8). Collectively, these imply that IS6120 integration upstream of *mpr* brings about

replacement of *wtp* by *rcp*, which raises the question of whether *M. smegmatis* regulates the

expression of Mpr by promoter reconstitution. In addition, we attempted to overexpress Mpr in

Ms^{Wt} by constructing Ms$^{wtp\text{-Mpr}}$ and Ms$^{rcp\text{-Mpr}}$ using the vectors pMV-*wtp-mpr* or pMV-*rcp-mpr*

respectively. However, strains transformed with pMV-*rcp-mpr* rarely formed colonies on the

plate (Figure S2). Due to low efficiency of Ms$^{rcp\text{-Mpr}}$ to form colonies, we are unable to provide

any definitive inference *vis-à-vis* the relationship between *rcp* and Mpr expression or D29

resistance. However, it does appear like expressing Mpr using *rcp* exerts toxicity on the bacterial

cells.

D29 resistance had no effect on *M. smegmatis* resistance to selected drugs

To assess potential fitness tradeoffs associated with D29 resistance, we performed drug

susceptibility testing (DST) on Ms^{Wt} and three resistant mutants (B6.1, A72.1, 2.6.1) against

seven frontline anti-tubercular agents: isoniazid, rifampicin, linezolid, clofazimine, bedaquiline,

levofloxacin, and ethambutol. Remarkably, all strains—including the resistant mutants—

exhibited identical susceptibility profiles to these antibiotics (Table S4). This lack of divergence

suggests that D29 resistance does not impinge upon core cellular processes targeted by these

drugs, nor does it incur detectable collateral costs in tolerance to these drugs. These findings

imply that mycobacterial adaptation to phage pressure may occur independently of classical

antibiotic resistance pathways.

Discussion

AMR triggered a renewed interest in phage therapy as a potential alternative to antibiotic therapy (3,33,34). However, phage resistance represents a potent threat to sustainable usability of phage therapy, addressing which would require dissecting the mechanisms of phage interactions with their bacterial hosts. Efforts towards dissecting phage-bacteria interactions revealed invaluable details regarding receptor requirements (19) and adaptive responses to phage pressure, including the roles played by IS elements (20,42,43) as well as Mpr in M. smegmatis (15). Studies on Mpr role in D29 resistance have relied on vector-borne overexpression of the gene. Here, we report two D29-resistant mutants with elevated Mpr expression that likely stemmed from promoter reconstitution by IS6120 integration directly upstream of *mpr*.

ISs are mobile genetic elements that are widespread in bacterial genomes and play important roles in bacterial evolution and adaptation to stress. It is becoming increasingly evident that phage infection triggers IS transposition, which causes these elements to integrate into certain genes in bacteria and cause phage resistance via different mechanisms (20,42,43). For example, TM4 infection of *M. smegmatis* triggers IS1096 transposition and integration into *lsr2*, hence activating the LOS cluster, causing PIMs accumulation and reduced phage adsorption (20). Such transposition events often cause large insertions that, coupled with widespread endogenous presence of IS elements in the chromosome, may often be missed by mere SNP calling. This

explains why they were not detected during the initial whole-genome sequence analysis.

Therefore, we analyzed the raw sequence data for all sequenced D29-resistant mutants using

ISMapper (27), and we detected several IS transposition events (Figure 3A). We investigated one

particular transposition event in A72.1 and 2.6.1, which resulted in integration of IS6120 within

the 168 bp region between *MSMEG_1236* (multi-copy phage resistance gene, *mpr*) and

MSMEG_1237 (Figure 3B). Mpr is a membrane-bound DNA exonuclease that, upon

overexpression, defends *M. smegmatis* against D29 by cleaving phage DNA following its

injection into the cytoplasm (14–16).

Mpr is required for the emergence of spontaneous D29-resistant mutants, and such role does not

require any mutation in the gene but overexpression of the wild type gene. However, the genetic

basis for emergence of D29-resistant mutants with elevated Mpr expression remained elusive.

Here, we uncovered that both A72.1 and 2.6.1 have IS6120 integration directly upstream of *mpr*,

and both show profoundly elevated expression of Mpr, not MSMEG_1237 (Figure 4B). In

addition, the integrated IS6120 element runs in the same orientation (5'-3') as *mpr*, not

MSMEG_1237, which could partly explain the IS transposition effect on Mpr rather than

MSMEG_1237. Further analysis of the sequence at the integration site revealed that IS6120

introduced a -35 promoter element and a TFBS for Lrp, which were not originally present at the

site, hence reconstituting a new, fuller promoter (Figure 5). Promoter reporter assays reveal that

the reconstituted promoter (*rcp*) has activity that is far superior than that of the original promoter

(*wtp*) and is nearly as good as the BCG promoter *hsp60* (Figures 7 and 8). This certainly begs the

question of whether elevated Mpr expression in spontaneous D29-resistant mutants has always

been due to reconstitution of a fuller, more powerful promoter upstream of *mpr*, which may also

explain why emergence of such mutants require the presence of an intact *mpr* (15). Failure to

isolate recombinant strains overexpressing *rcp-mpr* suggest extreme toxicity, and this is

consistent with previous effort to express *mpr* in *M. smegmatis* using the *hsp60* promoter (14).

This mechanism of gene regulation by promoter reconstitution is common in bacteria (44–46),

and the extreme toxicity may well be reminiscent of such situations where inherent protein

toxicity often makes it difficult to study (10,45). Mechanism of Mpr-mediated defense is

believed to be non-abortive in nature (16), and it remains to be seen whether A72.1 and 2.6.1

were able to grow normally in the face of elevated Mpr expression because the observed

expression levels were tolerable, or possibly due to neutralization of toxicity by other factors,

likely the second transposition event or other mutations detected in each of these strains. DEMs

of D29 have been reported for vector-borne Mpr-overexpressing *M. smegmatis* (16), but we were

unable to isolate a DEM for A72.1 and 2.6.1 in this study. It is not clear whether this could be

due to chromosome-borne, *rcp*-driven Mpr overexpression or absolute resistance in these strains

(13).

Mpr contains two major domains - a transmembrane domain (amino acids 33-50) and the domain

of unknown function DUF4352 (amino acids 83-211). DUF4352 is believed to be of viral origin,

suggesting that the bacteria likely acquired it during the course of evolution with viral predators,

then kept it for anti-viral defense (16). The fact that Ms^{Wt} is susceptible to D29 is evidence that

wild type level of Mpr expression does not confer resistance. However, regulation of Mpr

expression by promoter reconstitution could suggest that the gene is most useful to the bacteria

under D29-induced stress, not normal conditions. From our observations, Mpr appears to be the

primary anti-D29 resistance tool in *M. smegmatis*.

Events such as gene amplification-duplication are transient adaptive responses to increase

bacterial fitness, and are usually reversible due to the high fitness costs they exert (47,48). IS

transposition could also be reversible (43), but we haven't seen any evidence of such reversion in

A72.1 and 2.6.1 despite high expression of the toxic protein Mpr, and it remains to be seen

whether this might happen in later generations of the bacteria. Tolerance of A72.1 and 2.6.1 to

drugs appears to be unaffected, meaning that the strains had not traded off tolerance to the drugs

(at least those tested in this study) for D29 resistance. They also fare very well under normal

growth conditions, indicating lack of growth-related fitness cost associated with D29 resistance

in the strains.

This study reveals a form of gene regulation by promoter reconstitution in *M. smegmatis*. We

report that exposure of *M. smegmatis* to the lytic mycobacteriophage D29 triggers the

transposition and integration of IS6120 directly upstream of *mpr*, resulting in reconstitution of a

fuller, stronger promoter. This likely represents a previously elusive factor behind emergence of

D29-resistant mutants with elevated Mpr expression.

Acknowledgments

This study was supported by the National Key R&D Program of China (2021YFA1300900), the

Major Project of Guangzhou National Laboratory (GZNL2024A01009, GZNL2025C01003), the

National Natural Science Foundation of China (82502762).

Data availability

The next generation sequencing reads for Ms^{Wt}, 2.6.1 and A72.1 have been deposited at the

NCBI Sequence Read Archive, and can be accessed with SAMN55349604, SAMN55349605

and SAMN55349614 respectively.

References

1. GBD 2021 Antimicrobial Resistance Collaborators. Global burden of bacterial

antimicrobial resistance 1990-2021: a systematic analysis with forecasts to 2050. *Lancet.*

2024;404(10459):1199-1226. doi: 10.1016/S0140-6736(24)01867-1.

2. Expanded Access | FDA https://www.fda.gov/news-events/public-health-

focus/expanded-access

3. Dedrick RM, Guerrero-Bustamante CA, Garlena RA, Russell DA, Ford K, Harris K,

Gilmour KC, Soothill J, Jacobs-Sera D, Schooley RT, Hatfull GF, Spencer H. Engineered

bacteriophages for treatment of a patient with a disseminated drug-resistant

Mycobacterium abscessus. *Nat Med.* 2019; 25(5):730-733. doi: 10.1038/s41591-019-

0437-z.

4. Cristinziano M, Shashkina E, Chen L, Xiao J, Miller MB, Doligalski C, Coakley R, Lobo

LJ, Footer B, Bartelt L, Abad L, Russell DA, Garlena R, Lauer MJ, Viland M,

Kaganovsky A, Mowry E, Jacobs-Sera D, van Duin D, Kreiswirth BN, Hatfull GF,

Friedland A. Use of epigenetically modified bacteriophage and dual beta-lactams to treat

a *Mycobacterium abscessus* sternal wound infection. *Nat Commun.* 2024; 15(1):10360. doi: 10.1038/s41467-024-54666-4.

5. Dedrick RM, Smith BE, Cristinziano M, Freeman KG, Jacobs-Sera D, Belessis Y, Whitney Brown A, Cohen KA, Davidson RM, van Duin D, Gainey A, Garcia CB, Robert George CR, Haidar G, Ip W, Iredell J, Khatami A, Little JS, Malmivaara K, McMullan BJ, Michalik DE, Moscatelli A, Nick JA, Tupayachi Ortiz MG, Polenakovik HM, Robinson PD, Skurnik M, Solomon DA, Soothill J, Spencer H, Wark P, Worth A, Schooley RT, Benson CA, Hatfull GF. Phage Therapy of *Mycobacterium* Infections: Compassionate Use of Phages in 20 Patients With Drug-Resistant Mycobacterial Disease. *Clin Infect Dis.* 2023; 76(1):103-112. doi: 10.1093/cid/ciac453.

6. Dedrick RM, Freeman KG, Nguyen JA, Bahadirli-Talbott A, Smith BE, Wu AE, Ong AS, Lin CT, Ruppel LC, Parrish NM, Hatfull GF, Cohen KA. Potent antibody-mediated neutralization limits bacteriophage treatment of a pulmonary *Mycobacterium abscessus* infection. *Nat Med.* 2021; 27(8):1357-1361. doi: 10.1038/s41591-021-01403-9.

7. Nick JA, Dedrick RM, Gray AL, Vladar EK, Smith BE, Freeman KG, Malcolm KC, Epperson LE, Hasan NA, Hendrix J, Callahan K, Walton K, Vestal B, Wheeler E, Rysavy NM, Poch K, Caceres S, Lovell VK, Hisert KB, de Moura VC, Chatterjee D, De P, Weakly N, Martiniano SL, Lynch DA, Daley CL, Strong M, Jia F, Hatfull GF, Davidson RM. Host and pathogen response to bacteriophage engineered against *Mycobacterium abscessus* lung infection. *Cell.* 2022; 185(11):1860-1874.e12. doi: 10.1016/j.cell.2022.04.024.

8. Little JS, Dedrick RM, Freeman KG, Cristinziano M, Smith BE, Benson CA, Jhaveri TA, Baden LR, Solomon DA, Hatfull GF. Bacteriophage treatment of disseminated cutaneous *Mycobacterium chelonae* infection. *Nat Commun.* 2022; 13(1):2313. doi: 10.1038/s41467-022-29689-4.

9. Dedrick RM, Freeman KG, Nguyen JA, Bahadirli-Talbott A, Cardin ME, Cristinziano M, Smith BE, Jeong S, Ignatius EH, Lin CT, Cohen KA, Hatfull GF. Nebulized Bacteriophage in a Patient With Refractory *Mycobacterium abscessus* Lung Disease. *Open Forum Infect Dis.* 2022; 9(7):ofac194. doi: 10.1093/ofid/ofac194.

10. Dedrick RM, Jacobs-Sera D, Bustamante CA, Garlena RA, Mavrich TN, Pope WH, Reyes JC, Russell DA, Adair T, Alvey R, Bonilla JA, Bricker JS, Brown BR, Byrnes D,

Cresawn SG, Davis WB, Dickson LA, Edgington NP, Findley AM, Golebiewska U, Grose JH, Hayes CF, Hughes LE, Hutchison KW, Isern S, Johnson AA, Kenna MA, Klyczek KK, Mageeney CM, Michael SF, Molloy SD, Montgomery MT, Neitzel J, Page ST, Pizzorno MC, Poxleitner MK, Rinehart CA, Robinson CJ, Rubin MR, Teyim JN, Vazquez E, Ware VC, Washington J, Hatfull GF. Prophage-mediated defence against viral attack and viral counter-defence. *Nat Microbiol.* 2017; 2:16251. doi: 10.1038/nmicrobiol.2016.251.

11. Mageeney CM, Mohammed HT, Dies M, Anbari S, Cudkevich N, Chen Y, Buceta J, Ware VC. *Mycobacterium* Phage Butters-Encoded Proteins Contribute to Host Defense against Viral Attack. *mSystems.* 2020; 5(5):e00534-20. doi: 10.1128/mSystems.00534-20.

12. Gentile GM, Wetzel KS, Dedrick RM, Montgomery MT, Garlena RA, Jacobs-Sera D, Hatfull GF. More Evidence of Collusion: a New Prophage-Mediated Viral Defense System Encoded by Mycobacteriophage Sbash. *mBio.* 2019; 10(2):e00196-19. doi: 10.1128/mBio.00196-19.

13. Mohammed HT, Mageeney C, Ware VC. Identification of a new antiphage system in *Mycobacterium* phage Butters. *bioRxiv* 2023.01.03.522681; doi: https://doi.org/10.1101/2023.01.03.522681

14. Barsom EK, Hatfull GF. Characterization of *Mycobacterium smegmatis* gene that confers resistance to phages L5 and D29 when overexpressed. *Mol Microbiol.* 1996; 21(1):159-70. doi: 10.1046/j.1365-2958.1996.6291342.x.

15. Seniya SP, Jain V. Decoding phage resistance by *mpr* and its role in survivability of *Mycobacterium smegmatis*. *Nucleic Acids Res.* 2022; 50(12):6938-6952. doi: 10.1093/nar/gkac505.

16. Sharma S, Jain V. Structural and functional relevance of the DUF4352 domain in antiviral immunity and physiology of mycobacteria. *Nucleic Acids Res.* 2025;53(20):gkaf1253. doi: 10.1093/nar/gkaf1253.

17. Huang Y, Xu H, Zhang T, Liao Y, Hu J, Huang Z, Hu H, Li P, Fan L, Xie J. Mycobacterial non-homologous end joining is required for antiphage defense. *Nucleic Acids Res.* 2026;54(3):gkag066. doi: 10.1093/nar/gkag066.

18. Sørensen PE, Baig S, Stegger M, Ingmer H, Garmyn A, Butaye P. Spontaneous Phage Resistance in Avian Pathogenic *Escherichia coli*. *Front Microbiol.* 2021;12:782757. doi:

10.3389/fmicb.2021.782757.

19. McKitterick AC, Bernhardt TG. Phage resistance profiling identifies new genes required

for biogenesis and modification of the corynebacterial cell envelope. *Elife.*

2022;11:e79981. doi: 10.7554/eLife.79981.

20. Li Y, Wei Y, Guo X, Li X, Lu L, Hu L, He ZG. Insertion sequence transposition activates

antimycobacteriophage immunity through an *lsr2*-silenced lipid metabolism gene island.

mLife. 2024; 3(1):87-100. doi: 10.1002/mlf2.12106.

21. Wilson K. Preparation of genomic DNA from bacteria. *Curr Protoc Mol Biol.*

2001;Chapter 2:Unit 2.4. doi: 10.1002/0471142727.mb0204s56.

22. Afgan E, Baker D, Batut B, van den Beek M, Bouvier D, Cech M, Chilton J, Clements D,

Coraor N, Grüning BA, Guerler A, Hillman-Jackson J, Hiltemann S, Jalili V, Rasche H,

Soranzo N, Goecks J, Taylor J, Nekrutenko A, Blankenberg D. The Galaxy platform for

accessible, reproducible and collaborative biomedical analyses: 2018 update. *Nucleic*

Acids Res. 2018;46(W1):W537-W544. doi: 10.1093/nar/gky379.

23. Andrews S. (2010). FastQC: A quality control tool for high throughput sequence data.

Available online at https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

24. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 2018;34(17):i884-i890. doi: 10.1093/bioinformatics/bty560.

25. Seemann T (2015) snippy: fast bacterial variant calling from NGS reads Available online

at https://github.com/tseemann/snippy

26. Siguier P, Perochon J, Lestrade L, Mahillon J, Chandler M. ISfinder: the reference centre

for bacterial insertion sequences. *Nucleic Acids Res.* 2006;34(Database issue):D32-6. doi:

10.1093/nar/gkj014.

27. Hawkey J, Hamidian M, Wick RR, Edwards DJ, Billman-Jacobe H, Hall RM, Holt KE.

ISMapper: identifying transposase insertion sites in bacterial genomes from short read

sequence data. *BMC Genomics.* 2015;16(1):667. doi: 10.1186/s12864-015-1860-2.

28. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics. 2009;25(14):1754-60. doi: 10.1093/bioinformatics/btp324.

29. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,

Durbin R; 1000 Genome Project Data Processing Subgroup. The Sequence

Alignment/Map format and SAMtools. *Bioinformatics.* 2009;25(16):2078-9. doi:

10.1093/bioinformatics/btp352.

30. Faust GG, Hall IM. SAMBLASTER: fast duplicate marking and structural variant read extraction. *Bioinformatics.* 2014;30(17):2503-5. doi: 10.1093/bioinformatics/btu314.

31. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics.* 2010;26(6):841-2. doi: 10.1093/bioinformatics/btq033.

32. Kriel NL, Newton-Foot M, Bennion OT, Aldridge BB, Mehaffy C, Belisle JT, Walzl G, Warren RM, Sampson SL, Gey van Pittius NC. Localization of EccA$_3$ at the growing pole in *Mycobacterium smegmatis. BMC Microbiol.* 2022;22(1):140. doi: 10.1186/s12866-022-02554-6.

33. Schooley RT, Biswas B, Gill JJ, Hernandez-Morales A, Lancaster J, Lessor L, Barr JJ, Reed SL, Rohwer F, Benler S, Segall AM, Taplitz R, Smith DM, Kerr K, Kumaraswamy M, Nizet V, Lin L, McCauley MD, Strathdee SA, Benson CA, Pope RK, Leroux BM, Picel AC, Mateczun AJ, Cilwa KE, Regeimbal JM, Estrella LA, Wolfe DM, Henry MS, Quinones J, Salka S, Bishop-Lilly KA, Young R, Hamilton T. Development and Use of Personalized Bacteriophage-Based Therapeutic Cocktails To Treat a Patient with a Disseminated Resistant *Acinetobacter baumannii* Infection. *Antimicrob Agents Chemother.* 2017;61(10):e00954-17. doi: 10.1128/AAC.00954-17.

34. Aslam S, Courtwright AM, Koval C, Lehman SM, Morales S, Furr CL, Rosas F, Brownstein MJ, Fackler JR, Sisson BM, Biswas B, Henry M, Luu T, Bivens BN, Hamilton T, Duplessis C, Logan C, Law N, Yung G, Turowski J, Anesi J, Strathdee SA, Schooley RT. Early clinical experience of bacteriophage therapy in 3 lung transplant recipients. *Am J Transplant.* 2019;19(9):2631-2639. doi: 10.1111/ajt.15503.

35. Besra GS, Khoo KH, Belisle JT, McNeil MR, Morris HR, Dell A, Brennan PJ. New pyruvylated, glycosylated acyltrehaloses from *Mycobacterium smegmatis* strains, and their implications for phage resistance in mycobacteria. *Carbohydr Res.* 1994;251:99-114. doi: 10.1016/0008-6215(94)84279-5.

36. Dhariwal KR, Liav A, Vatter AE, Dhariwal G, Goren MB. Haptenic oligosaccharides in antigenic variants of mycobacterial C-mycosides antagonize lipid receptor activity for mycobacteriophage D4 by masking a methylated rhamnose. *J Bacteriol.* 1986;168(1):283-93. doi: 10.1128/jb.168.1.283-293.1986.

37. Furuchi A, Tokunaga T. Nature of the receptor substance of *Mycobacterium smegmatis*

for D4 bacteriophage adsorption. *J Bacteriol.* 1972;111(2):404-11. doi: 10.1128/jb.111.2.404-411.1972.

38. Imaeda T, Blas FS. Adsorption of Mycobacteriophage on Cell-wall Components. *J Gen Virol.* 1969;5(4):493–8.

39. Wetzel KS, Illouz M, Abad L, Aull HG, Russell DA, Garlena RA, Cristinziano M, Malmsheimer S, Chalut C, Hatfull GF, Kremer L. Therapeutically useful mycobacteriophages BPs and Muddy require trehalose polyphleates. *Nat Microbiol.* 2023;8(9):1717-1731. doi: 10.1038/s41564-023-01451-6.

40. Li X, Long X, Chen L, Guo X, Lu L, Hu L, He ZG. Mycobacterial phage TM4 requires a eukaryotic-like Ser/Thr protein kinase to silence and escape anti-phage immunity. *Cell Host Microbe.* 2023;31(9):1469-1480.e4. doi: 10.1016/j.chom.2023.07.005.

41. Chen J, Kriakov J, Singh A, Jacobs WR, Besra GS, Bhatt A. Defects in glycopeptidolipid biosynthesis confer phage I3 resistance in *Mycobacterium smegmatis. Microbiology (Reading).* 2009;155(Pt 12):4050-4057. doi: 10.1099/mic.0.033209-0.

42. Fujiki J, Nakamura T, Kreimeyer H, Llorente C, Fouts DE, Schnabl B. Insertion sequence-mediated phage resistance contributes to attenuated colonization of cytolytic *Enterococcus faecalis* variants in the gut. *Microbiol Spectr.* 2025;13(5):e0330324. doi: 10.1128/spectrum.03303-24.

43. Styles KM, Locke RK, Cowley LA, Brown AT, Sagona AP. Transposable Element Insertions into the *Escherichia coli* Polysialic Acid Gene Cluster Result in Resistance to the K1F Bacteriophage. *Microbiol Spectr.* 2022;10(3):e0211221. doi: 10.1128/spectrum.02112-21.

44. Durrant MG, Perry NT, Pai JJ, Jangid AR, Athukoralage JS, Hiraizumi M, McSpedon JP, Pawluk A, Nishimasu H, Konermann S, Hsu PD. Bridge RNAs direct programmable recombination of target and donor DNA. *Nature.* 2024;630(8018):984-993. doi: 10.1038/s41586-024-07552-4.

45. Wilkinson ME, Li D, Gao A, Macrae RK, Zhang F. Phage-triggered reverse transcription assembles a toxic repetitive gene from a noncoding RNA. *Science.* 2024;386(6717):eadq3977. doi: 10.1126/science.adq3977.

46. Ton-Hoang B, Bétermier M, Polard P, Chandler M. Assembly of a strong promoter following IS911 circularization and the role of circles in transposition. *EMBO J.*

1997;16(11):3357-71. doi: 10.1093/emboj/16.11.3357.

47. Andersson DI, Hughes D. Gene amplification and adaptive evolution in bacteria. *Annu*

Rev Genet. 2009;43:167-95. doi: 10.1146/annurev-genet-102108-134805.

48. Kugelberg E, Kofoid E, Reams AB, Andersson DI, Roth JR. Multiple pathways of

selected gene amplification during adaptive mutation. *Proc Natl Acad Sci U S A.*

2006;103(46):17319-24. doi: 10.1073/pnas.0608309103.

681   **Tables and Figure Legends**

683   **Figure 1 Spontaneous**
684   **D29 resistance in *M. smegmatis*.** A, A72.1 and 2.6.1 were tested for D29 resistance by spot-kill assay, and
685   both appear to be particularly highly resistant to D29 compared to Ms^{Wt}. B, both strains were also tested for
686   D29 resistance by plaque assay, which appears to be consistent with the spot-kill assay. C, terminal adsorption
687   suggests that D29 adsorption is not inhibited in the two strains, suggesting that the resistance mechanism is at a
688   stage downstream of phage adsorption.

Panel		A	B	C	D	E	F	G	H	I
IS element		IS6120	IS6120	IS6120	IS6120	IS6120	IS1096	ISMsm1	ISMsm1	ISMsm1
Size (bp)	before	154	486	153	166	72	3169	134	79	141
	after	1631	1962	1629	1641	1547	5419	1475	1419	1482

691 **Figure 2 PCR verification of IS rearrangements.** A total of 7 (29.2 %) of the sequenced D29-resistant
692 mutants sustained at least a single IS transposition event. PCR verification of integration at each site as well as
693 affected gene/region are shown in the upper panels A-I, whereas sizes at the site before and after IS integration
694 are outlined in the lower table in the figure. Integration at intergenic region is indicated by a "black diamond
695 icon" between the two genes.

	A33.1	A72.1	B10.1	B28.2	B35.1	2.6.1	5.2.1
MSMEG_1236♦MSMEG_1237		IS6120				IS6120	
MSMEG_1730				IS6120			
MSMEG_2105♦MSMEG_2104		ISMsm1					
MSMEG_3275♦MSMEG_3276				ISMsm1			
MSMEG_3324			IS6120				
MSMEG_3756 (rrlA)					IS1096		
MSMEG_4724							IS6120
MSMEG_5091	IS6120						
MSMEG_5393 (kdpB)						ISMsm1	

Figure 3 IS rearrangements in the D29-resistant mutants. A, the different strains/regions affected by IS rearrangements. IS integration at intergenic region is indicated by a "black diamond" between the two genes. B, both A72.1 and 2.6.1 – the D29-resistant mutants manifesting the highest level of resistance – sustained IS6120 integration between nucleotides G-70 and G-69 upstream of *MSMEG_1236* (*mpr*). This integration resulted in two direct repeats (indicated by black diamonds) at either sides of IS6120 at the integration site as well as introduction of the -35 promoter element "TTGACA" and a transcription factor-binding site (TFBS) for Lrp (leucine-responsive regulatory protein). Figure is drawn in a reverse orientation. Upper case letters, nucleotides introduced by IS6120 integration; underlined letters, TFBs; lower case letters, original sequence at the site; blue-font letters, direct repeats; TSS, transcription start site; atg, *mpr* start codon.

Figure 4 Elevated Mpr expression in A72.1 and 2.6.1. A, STRING protein-protein interaction network for Mpr. Mpr and MSMEG_1237 are adjacent to each other, and a possible co-expression is predicted in their interaction profile. B, RT-qPCR of Mpr, MSMEG_1237 as well as their intergenic region (IR) reveals that Mpr expression is elevated in A72.1 and 2.6.1.

Number of predicted promoters in:

Prediction program	*wtseq*	*rcseq*
BPROM	0	1
BGDP	0	1
Sigma70Pred	3	14
PePPER	0	1

-35   -10
TTGACAATTCCTCgtcccgagttgtacgtt
TFBS

Upper case, IS6120 sequence
Lower case, original sequence

Figure 5 Promoter prediction in *rcseq*. Four promoter prediction programs detected the presence of a new promoter in *rcseq*. All these promoters share one sequence in common, which contains a putative TFBS, -35 and -10 promoter elements.

Figure 6 Quantification of *wtp*- and *rcp*-driven green fluorescence. A, Lawn appearance of GFP-expressing strains using *hsp60*, *wtp* and *rcp*. Ms$^{rcp\text{-GFP}}$ shows the characteristic green coloration of GFP. B, Live fluorescence of each strain as quantified using a microplate reader at different optical densities indicates that *rcp* drives higher fluorescence than *wtp*.

Figure 7 Quantification of *wtp*- and *rcp*-driven GFP events by flow cytometry. Quantification of GFP
events places *hsp60* and *rcp* on par with each other, whereas *wtp* falls far behind in comparison. Quantification
was carried out in a 10,000-cell population for each strain. Populations expressing GFP using *hsp60* (Ms^{GFP}),
rcp (Ms$^{rcp\text{-}GFP}$) and *wtp* (Ms$^{wtp\text{-}GFP}$) had 99.9, 99.8 and 19.5 % GFP-positive cells respectively. In contrast, the
negative control (Ms^{Wt} population) had a negligible 0.91 % GFP-false positive cells.

Ms^{GFP}   Ms^{rcp}-GFP   Ms^{wtp}-GFP

Figure 8 Quantification of fluorescence signal. Fluorescence signal driven by *hsp60*, *rcp* and *wtp* were quantified by confocal laser scanning microscopy. Strength of signal produced by *hsp60* and *rcp* appear to be similar, whereas *wtp* drives far weaker fluorescence signal under the microscope.