Vorinostat Rescues SQSTM1 Palmitoylation and Restores Dysfunctional Autophagy in Huntington Disease

Yasmeen Alshehabi, Fatima Abrar, Milena Rabu, Anthony Dang, Firyal Ramzan, and Dale D. O. Martin

NeurdyPhagy Lab, Department of Biology, Faculty of Science, University of Waterloo, Waterloo, ON, Canada

Running title: Vorinostat restores autophagy in HD

Correspondance: Dale D. O. Martin, dale.martin@uwaterloo.ca

Keywords: neurodegeneration, autophagy, SQSTM1, p62, Vorinostat, Huntington Disease, S-acylation, palmitoylation

Abstract

Protein mislocalization, impaired autophagy, and protein aggregation are central features of Huntington disease (HD), yet therapeutic strategies that simply stimulate autophagy risk exacerbating dysfunction in an already compromised pathway. We previously demonstrated that palmitoylation of the autophagy receptor SQSTM1/p62 is reduced in HD patient brains and YAC128 mice, contributing to defective cargo loading and accumulation of mutant huntingtin (mHTT) aggregates. Here, we identify the FDA-approved histone deacetylase inhibitor Vorinostat (also known as suberoylanilide hydroxamic acid; SAHA) as a small molecule that restores SQSTM1 palmitoylation and autophagic function in HD models. Using acyl-biotin exchange and click chemistry assays, we show that Vorinostat selectively increases SQSTM1 palmitoylation without altering total protein palmitoylation. In turn, Vorinostat enhances autophagic flux, decreases SQSTM1 levels, increases LC3-II, and promotes colocalization of SQSTM1, mHTT, and lysosomes, indicating improved cargo delivery and degradation. Importantly, systemic administration of Vorinostat crosses the blood-brain barrier and restores SQSTM1 palmitoylation in YAC128 mouse cortices. Mechanistically, Vorinostat may function through dual pathways: partial inhibition of depalmitoylating enzymes and transcriptional regulation of palmitoylation machinery. These findings build on our identification of SQSTM1 palmitoylation as a therapeutic target for HD and repositions Vorinostat as a candidate modifier of autophagy dysfunction and palmitoylation impairment in HD.

1. Introduction

Protein aggregation is a hallmark of several neurodegenerative diseases, including Huntington disease (HD), amyotrophic lateral sclerosis (ALS), Alzheimer disease (AD), and Parkinson disease (PD)[1–3]. The intracellular pathway responsible for clearing aggregates, known as autophagy, is often impaired in these devastating conditions[1]. As a result, autophagy is an attractive therapeutic target for clearing the toxic aggregates that exacerbate disease progression. However, only stimulating autophagy is insufficient, as it places additional stress on a system that is already compromised. Therefore, addressing the underlying defects, in conjunction with increasing autophagy, is essential for effective therapeutic intervention.

Macroautophagy, henceforth referred to as autophagy, involves the loading of cargo, such as toxic or aggregated proteins and damaged organelles, into a double-membrane structure known as a phagophore, which seals to form an autophagosome[4] (Figure 1B). This autophagosome is then transported and fused with a lysosome to form the ultimate structure known as an autophagolysosome where the cargo is degraded with hydrolases[4]. The cargo-loading step is crucial, wherein sequestosome 1 (SQSTM1; commonly known as p62) binds ubiquitinated cargo destined for degradation and delivers it to lipidated microtubule-associated protein 1 light chain 3 (LC3-II), located on both sides of the phagophore membrane[4]. In HD, autophagy is dysfunctional at several steps (Figure 1C)[5]. Critically, there is a cargo-loading failure that leads to an accumulation of empty autophagosomes and protein aggregates[6,7].

Figure 1 – SQSTM1 and HTT palmitoylation may support healthy autophagy, and its deficiency may contribute to autophagic dysfunction in HD. (A) Palmitoylation involves the formation of a thioester bond between palmitoyl-CoA and a free cysteine residue in a protein via palmitoyl acyltransferases (ZDHHCs). Depalmitoylation involves the cleavage of a thioester bond via acyl protein thioesterases (APTs) or α/β-hydrolase domain enzymes (ABHDs). **(B)** LC3-I undergoes lipidation to form LC3-II, located on both sides of the phagophore membrane. LC3-II then recruits SQSTM1, which is bound to the ubiquitinated cargo destined for degradation. HTT acts as a scaffold to support the function of SQSTM1. We predict that the palmitoylation of SQSTM1 and HTT plays an important role in their delivery of cargo to the phagophore membrane. Next, the phagophore expands around the cargo and closes to form a sealed autophagosome, which then fuses with a lysosome to form an autophagolysosome. All the contents inside this vesicle are degraded by hydrolytic enzymes, and LC3-II bound to the outside of the double-membrane is released back to the cytosol. **(C)** In HD, there is a defect at the cargo-loading step leading to an accumulation of mHTT aggregates and empty autophagosomes. We predict that this cargo-loading failure is at least partially the result of a reduction in SQSTM1 and HTT palmitoylation. Figure created with Biorender.

HD is a neurodegenerative disease with devastating affects on cognition and motor function[8]. HD is an

autosomal dominant, monogenic disease caused by a mutation in exon 1 of the huntingtin gene (*HTT*)

that encodes a cytosine-adenine-guanine (CAG) repeat expansion (*i.e.,* 36 to 39 CAG repeats for partial

HD penetrance; 40 or more repeats for full HD penetrance) resulting in a polyglutamine expansion in the

huntingtin protein (HTT)[9,10]. In healthy conditions, HTT is ubiquitous and involved in many cellular

processes[11], including autophagy, in which HTT functions as a scaffold, binding to SQSTM1 and

ubiquitinated cargo to support the function of SQSTM1 in cargo-loading[12–14]. However, cargo-loading is

impaired in HD[6,7], which may be at least partly due to reduced palmitoylation of SQSTM1 and mHTT. We

have shown that SQSTM1 palmitoylation plays an important role in cargo-loading and is required for

SQSTM1 localization to autophagosomes[15,16]. Notably, SQSTM1 palmitoylation is significantly reduced in

the brains of HD patients and YAC128 mice[15,16]. In turn, increasing SQSTM1 palmitoylation using broad

palmitoylation inhibitor Palmostatin B improved HTT and SQSTM1 localization to autophagosomes[15,16].

Moreover, the Hayden group has shown that mHTT palmitoylation is reduced in HD mouse models and

further decreases with increasing age and CAG repeat length[17,18]. Subsequently, Palmostatin B restored

mHTT palmitoylation and thus mHTT aggregation and cytotoxicity in HD mouse model primary

neurons[18].

Palmitoylation, the most common form of S-acylation, is the reversible post-translational addition of a

16-carbon fatty acid to a cysteine residue via a thioester bond[19] (Figure 1A). Due to its reversibility and

hydrophobicity, palmitoylation allows selective and dynamic targeting of soluble proteins to membranes

and thus plays a key role in regulating protein trafficking[19]. Palmitoylation is mediated by 23 palmitoyl

acyltransferases (ZDHHCs) in humans, whereas the reverse reaction is mediated by acyl protein

thioesterase (APT), palmitoylprotein thioesterase (PPT), and α/β-hydrolase domain (ABHD) enzymes[20].

The Hayden group has shown that HTT is palmitoylated by ZDHHC13 and ZDHHC17, and depalmitoylated

by APT1 and APT2[17,18,21,22]. Importantly, HD is linked to decreased ZDHHC17 activity and increased APT1

activity[23]. Furthermore, the Yang group has shown that SQSTM1 is depalmitoylated by APT1[15,24]. In turn,

the Saudou group demonstrated that APT1 activity is increased in HD and specific inhibition of APT1

using ML348 is protective in an HD mouse model[25]. Consequently, APT1 may be a therapeutic target in

HD. Therefore, the identification of a small molecule which can increase SQSTM1 palmitoylation holds

therapeutic promise for HD, rescuing cargo-loading in autophagy, and in turn, reducing mHTT

aggregation and toxicity. Through a high-throughput screening of FDA-approved drugs, we identified

Vorinostat, also known as suberoylanilide hydroxamic acid (SAHA), as a potential candidate for

increasing SQSTM1 palmitoylation[26]. Vorinostat is primarily known as a blood-brain barrier (BBB)

permeable inhibitor of lysine deacetylases (KDACs), more commonly known as histone deacetylases

(HDACs), which catalyze the removal of acetyl groups (*i.e.,* deacetylation) from lysine residues on

histone and/or non-histone proteins[27,28]. Herein, we report that Vorinostat significantly increases

SQSTM1 palmitoylation, increases and enhances autophagic flux, and may function as a

depalmitoylating enzyme inhibitor and a transcriptional regulator. Importantly, we found that

Vorinostat crosses the blood-brain barrier and restores SQSTM1 palmitoylation in YAC128 HD mouse

model brains.

2. Materials and Methods

2.1 Cell Culture

HeLa cells were grown at 37^{o}C in 5% CO_2 in Dulbecco's Modified Eagle Medium (DMEM; Wisent #319-015 CL) supplemented with 10% fetal bovine serum (FBS), 2 mM L-glutamine, 1 mM sodium pyruvate, 1X penicillin-streptomycin. Cells were seeded at 170,000 cells per 6 cm plate for biochemistry experiments and 75,000 cells per well onto 0.01 mg/mL poly-D-lysine coated coverslips (#1.5; VWR) in 6-well plates for microscopy experiments. The next day, cells were transfected with 10 µg of plasmid DNA using calcium phosphate precipitation and incubated for 2 hours. The transfection mix contained 1X HEPES-buffered saline (HBS) and 122 mM $CaCl_2$. The media was replaced 2 hours post transfection. Where indicated, drugs were added to transfected cells immediately following the media change and cells were processed 18 hours post transfection (*i.e.,* 16-hour drug incubation). For non-transfected cells, indicated drugs were added for 16 hours prior to processing. Drugs were added at the listed concentrations: 10 µM Vorinostat (Sigma #SML0061), 10 µM ML348 (Cayman #18523), 10 µM Palmostatin B (PalmB; Millipore #178501), 50 µM 2-bromopalmitate (2BP; Sigma #238422), and/or 100 nM Bafilomycin A1 (BafA1; Millipore #196000). All drugs were dissolved in dimethyl sulfoxide (DMSO), except for 2BP which was dissolved in ethanol.

2.2 Click Chemistry Assay

Cell labeling and click chemistry were performed as previously described[29]. Briefly, the alkynyl palmitate fatty acid analog (Click Chemistry Tools #CCT-1165) was saponified in potassium hydroxide and conjugated to fatty-acid free bovine serum albumin (BSA). Cells were then deprived of lipids in DMEM supplemented with 5% charcoal-stripped FBS (Wisent #080-710), 2 mM L-glutamine, 1 mM sodium pyruvate, 1X penicillin-streptomycin, and incubated with the alkynyl palmitate for 4 hours, after which cells were lysed in modified RIPA buffer (50 mM HEPES pH 7.4, 150 mM NaCl, 2 mM $MgCl_2$, 1% Igepal CA-630, 0.5% sodium deoxycholate, 0.1% SDS). Iodoacetamide was added to each sample to block free

cysteine residues. Finally, lysates were subjected to click chemistry with biotin-azide (Click Chemistry

Tools #CCT-1488), tris-hydroxypropyl triazolyl methylamine (THPTA; Click Chemistry Tools #CCT-1010),

copper sulfate, and sodium ascorbate for 1 hour. The reaction was then ended by acetone precipitation.

Since we are looking at total palmitoylated protein, the affinity purification step was skipped, and the

protein was denatured at 95°C for 5 minutes in 1X sample loading buffer (SLB) with 20 mM dithiothreitol

and stored at -20°C.

2.3 Acyl Biotin Exchange (ABE) Assay

The acyl biotin exchange (ABE) was performed as previously described[30]. Briefly, ground cortical brain

tissue (approximately 30 mg) was sonicated (3x6 seconds at 20%) in ABE homogenization buffer (10 mM

10X phosphate buffer pH 7.4, 0.32 M sucrose, 1 mM EDTA, 6 M urea, 2.5% SDS) with 1X PIC and 20 mM

methyl methanthiosulfate (MMTS). Alternatively, cells were sonicated (2x6 seconds at 20%) in lysis

buffer (50 mM HEPES pH 7.4, 1 mM EDTA, 2% SDS) with 1X PIC and 20 mM MMTS. Lysates were

incubated for 30 minutes at 50°C to block free cysteine residues with MMTS. Excess MMTS was removed

by acetone precipitation and resuspended in 4SB (50 mM Tris pH 7.5, 5 mM EDTA, 4% SDS). Samples

were then incubated with 0.7 M hydroxylamine and 1 mM HPDP-Biotin (APExBIO #A8008) for 1 hour

rotating at room temperature to cleave thioester bonds and label newly freed cysteine residues with

biotin. The reaction was ended by acetone precipitation and resuspended in lysis buffer without MMTS.

Following protein quantification, 0.5-1 mg of protein was incubated in 150 mM NaCl dilution buffer (50

mM HEPES pH 7.0, 1% TX-100, 1 mM EDTA, 1 mM EGTA) with High-Capacity NeutrAvidin Agarose beads

(Thermo #29202) for 3 hours rotating at 4°C to bind biotinylated proteins. Following two washes with

0.5 M NaCl dilution buffer and one wash with dilution buffer without NaCl, the protein was eluted from

beads with elution buffer (dilution buffer, 250 mM NaCl, 0.2% SDS, 1% β-mercaptoethanol) over a 10-

minute incubation at 37°C, then denatured at 95°C for 5 minutes in 1X SLB with 1% β-mercaptoethanol

and stored at -20°C.

2.4 SDS-PAGE and Western Blotting

12% Bis-Tris polyacrylamide gels were run in 1X XT MOPS buffer (Bio-Rad #1610788) and transferred onto nitrocellulose (to detect SQSTM1) or a polyvinylidene fluoride (to detect LC3-II) membranes via semi-dry transfer using transfer buffer (192 mM glycine, 25 mM tris base, 20% methanol, 0.05% SDS) and the Transblot Turbo Transfer System. Primary antibodies rabbit anti-SQSTM1 (1:1000; Cell Signaling #D1Q5S) or rabbit anti-SQSTM1 (1:1000; Invitrogen #PA5-20839), rabbit anti-LC3A/B (1:1000; Cell Signaling #D3U4C), and mouse anti-VCP (1:1000; Proteintech #60316-1-Ig) were used to detect SQSTM1, LC3-II, and VCP, respectively. Secondary antibodies goat anti-rabbit Starbright 520 (1:2500; Bio-Rad #12005870), donkey anti-rabbit Alexa Fluor 790 (1:2500; Jackson ImmunoResearch #711-655-152), and goat anti-mouse Alexa Fluor 488 (1:2500; Jackson ImmunoResearch #115-545-003) were used. Tubulin-rhodamine (1:5000; Bio-Rad #12004166) was used to detect tubulin as the loading control. Alexa Fluor 680 Streptavidin (1:5000; Jackson ImmunoResearch #016-620-084) was used to detect biotinylated proteins.

2.5 Fluorescence Microscopy

For fixed cell imaging, cells were treated with 50 nM LysoTracker Deep Red (Invitrogen #L12492) for 1 hour prior to fixation. Cells were washed twice with 1X phosphate buffer saline (PBS), then fixed in 4% paraformaldehyde for 20 minutes at room temperature. Again, cells were washed twice with 1X PBS, then permeabilized in 1X PBS containing 0.1% TX-100, 1 µM CaCl$_2$, 1 µM MgCl$_2$ for 1 minute. Again, cells were washed twice with 1X PBS, then blocked in 0.2% gelatin for 30 minutes. Next, cells were incubated with the primary antibody for 1 hour. Rabbit anti-SQSTM1 (1:500; Invitrogen #PA5-20839) and mouse anti-HTT (1:500; Millipore #MAB2166) were used to detect SQSTM1 and HTT, respectively. Cells were washed twice with 0.2% gelatin, then the last step was repeated with the secondary antibody. Secondary antibodies goat anti-rabbit Alexa Fluor 488 (1:500; Invitrogen) and donkey anti-mouse Alexa Fluor 555 (1:500; Invitrogen) were used. Again, cells were washed twice with 0.2% gelatin, then

incubated in 1 μg/mL DAPI (Sigma #D9542) for 20 minutes. Finally, cells were washed twice with 0.2%

gelatin. The coverslips were mounted onto microscope slides (Fisher) using ProLong Gold Antifade

Mountant (Invitrogen #P36934) and sealed with nail polish. Using a Nikon AXR laser scanning confocal

microscope and its NIS-Elements software, Z-stack images were captured at 60X magnification. Next, 3D

Deconvolution was applied. Protein colocalization was determined by measuring Pearson Correlation

Coefficient (PCC) between channels. For HeLa cells, PCC was determined from regions of interest

manually drawn around single cells. For neurons, PCC was determined from regions of interest using the

automated General Analysis 3 (GA3).

2.6 Molecular Docking

Molecular docking prediction was performed using the AutoDock Vina program[31]. Briefly, the protein

PDBQT file was opened in AutoDockTools software to define the grid box dimensions. Molecular docking

was then performed using AutoDock Vina via the command prompt, specifying the file path to the

program and a folder containing the protein PDBQT file, the ligand PDBQT file, and a text file listing the

PDBQT file names, grid box dimension values, energy range, and exhaustiveness. Energy range was set

to 4 and exhaustiveness to 64. Running this program produced an "output.pdbqt" file containing the top

ligand binding conformations, along with a "log.txt" file listing the corresponding binding affinities.

2.7 qPCR

Total RNA from cells was extracted using RNAZol RT (Bioshop #RNN190). Then, 500 ng of total RNA was

used to prepare cDNA using High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems

#4368813). Next, cDNA was diluted 1:5, and qPCR was performed using primers and the PowerTrack

SYBR Green Master Mix (Applied Biosystems #A46109). Each sample was run in triplicate. qPCR cycling

conditions were 2 minutes at 95°C, followed by 40 cycles of 5 seconds at 95°C and 30 seconds at 60°C,

then finally 15 seconds at 95°C, 1 minute at 60°C, and 15 seconds at 95°C. The forward and reverse

primer sequences were synthesized by IDT and are listed in Table 1. *ACTB* and *GAPDH* were used as

housekeeping genes. Primers for the ZDHHCs were previously published[32]. Primers for *SQSTM1*, *HTT*,

VCP, *MAP1LC3B*, the histone deacetylases, the serine hydrolases, and the housekeeping genes were

designed for this study.

Table 1 List of Human Primers for qPCR

Gene	Forward Sequence (5' to 3')	Reverse Sequence (5' to 3')
ACTB	GCCGCCAGCTCACCA	CTCGTCGCCCACATAGGAAT
GAPDH	TCGGAGTCAACGGATTTGGT	TTCCCGTTCTCAGCCTTGAC
ZDHHC1	CTCACCCGCTCCAGATTGTG	TGAGGCAGGAGGGGAACAAG
ZDHHC2	TGGAGAGAGAGCCAAGAGGAG	TTATAAGTTGGCATCTGTCACAGTATC
ZDHHC3	CTTAACTTCAGTTCGACAGAGTGC	GAAGAGCAGGCCCTCAAAGC
ZDHHC4	GATTGTCTTCATGCTGGGCTTTG	CTCTGTACCACTCGTTAGTAGTCTG
ZDHHC5	GAAGACTGAAGAAAGATAAGAGACATTG	GACACTTCAAAAGTTTACTGTGGATG
ZDHHC6	TCTGCTTGTCCTGGGCTGAG	TGTCTTGGTCTGAAACCCAACC
ZDHHC7	AGTCCATGCTCTGATCCTTTGTG	GGAAGATCAACAGGATTACAGTTATCG
ZDHHC8	TCTGTGACAACTGTGTAGAGGAC	AGCAGGAACAGGAAGAAGTAGC
ZDHHC9	TAGCAATTGGACTTTTGATGATGTTTG	GGGTTTTGCGATTACACGAGAG
ZDHHC11	TCCGTCACCCCAGAAGCCATAC	CGTGAGGCAGGAAGGGAATGAAG
ZDHHC12	TGCTGACCTGGGGAATCACG	CATGAGTGACACAGCGAGGTAG
ZDHHC13	GCATATCTCATCTCAAAGGGACAG	CAACCACATTGAGAGAAGGATTAAAC
ZDHHC14	TCGCTGGCATCCTGTTCTTC	TGCCGTTTGCGATATCTATTTGC
ZDHHC15	TATGAATGAGTCACAGAACCCACTGCTAG	TAATCTTGGTTGTCATCCTCGTTGTCTTCC
ZDHHC16	ATTGTAGCTATCGCCTACCTGTG	AGTAGTGGAAGACAATCAGGATCAG
ZDHHC17	ATGGTTGTGCAACTAATGAAATATGG	GCTATGAGATAAGCAACAATTGAGG
ZDHHC18	CCACCACCGGCCTCTTCTTC	GAGGATGGCAGCGATGATGG
ZDHHC19	GTCCCTGGTTCCTCCCTAGC	AATGCGAAGAAGAGGCCACTG
ZDHHC20	ATCAACCTTTTCCTATCAAACCACTTAG	GCCTTCTTCAGCTCCATTCTCC
ZDHHC21	TCTGGGAATTATGTAACAAGTGTAATTTG	AGAGCCAATGATTATCTTCACCAAC
ZDHHC23	AAAAGTGAACATCAGCATCATCCC	CCAGGAGGAAATGCCAGGAAG
ZDHHC24	CCCCTTCCGAGCTTCTATTGG	CTAGGCCGCTTCCGTATTGG
SQSTM1	GCACTACCGCGATGAGGAC	GCCATCGCAGATCACATTGG
HTT	GCATGGGATCCTGTCCCTTC	GGAGTGTATGGACACCTGGC
VCP	TCTGGAGCCGATTCAAAAGGT	TTCATCCATCTTGGGCTGGG
LYPLA1	CGCTCTTCCTTCCGCTTGC	GCAGGAAAATCACCGCAGC
LYPLA2	AGGCAGCAGAGAACATCAAGG	CAGGTGAGGGCCGTGTAGAG
LYPLAL1	CCTGCATGGCTCAGGTGATT	TCAAGGTGTTCTGGGCAGTC
PPT1	GAAGACCCTGATGGAGGACG	GATCATGGGAGGTGAAGGGC
PPT2	AGACCATCCCAATGCCACAG	CCCGCAGATAAACCAGTTGC
ABHD10	AGCTGATGGGCCACAGATTC	GCATGCTCCACACACCTTTC
ABHD16A	CTCCTGCAGCATCGGTATCC	CGTGTTCTGCCTGGTAGGAG
ABHD17B	TTGCTCTTAGGACAAGATATGGC	ATCAGGAAAGGCAACTCGC

ABHD17C	TTCTACATTGGCCTCGGCTC	GACTCACGCCATACCGGG
HDAC1	TCTATCGCCCTCACAAAGCC	CTCACCAACGTTGAATCTCTGC
HDAC2	GTGAGATTCCCAATGAGTTGCC	GGTGCATGAGGTAACATGCG
HDAC3	ACTACTGTCTGAACGTGCCC	ATTCAACGCATTCCCCATGC
HDAC6	CTATGTCCACCACCAGGCTC	CCCTCATCCAAGGTACAGGC
MAP1LC3B	GCTCATCAAGATAATTAGAAGGCG	GTCTCCTGGGAGGCATAGAC

2.8 Mice

YAC128 transgenic mice on an FVB/N genetic background were obtained from in-house breeding

colonies and were originally purchased from the Jackson Laboratory (Strain ID: 004938). Mice were

housed in standard laboratory cages under a 12-hour light/dark cycle at 23 ± 1^{o}C, with *ad libitum* access

to food and water, and toys for enrichment. All mouse experiments were conducted in accordance with

Animal Use Protocol #44651, approved by the UW Animal Care Committee. Mice were administered

Vorinostat once every second day between 10 am and 12 pm, at a dose of 50 mg/kg through

intraperitoneal (IP) injection (alternating right and left side) for four consecutive weeks for a total of 15

treatments[33]. Vorinostat was dissolved in DMSO at a concentration of 40 mg/mL and stored at -20^{o}C. On

the day of IP injection, the stock was diluted in corn oil to yield a working solution containing 18.75%

DMSO and 81.25% corn oil, with a final Vorinostat concentration of 7.5 μg/μL. Control injections

contained the same DMSO-to-corn oil ratio. Solutions were warmed to 37^{o}C for 5 minutes and vortexed

for 15 seconds prior to IP injection.

2.9 Mouse-Derived Primary Cortical Neurons

Primary cortical neurons were derived from YAC128 mouse embryos at embryonic day 17. Cortices were

dissociated using papain and seeded on 1 mg/mL poly-L-lysine coated wells or glass coverslips (#1.5;

Warner) in 6-well plates at 280,000 and 180,000 neurons per well for biochemistry and microscopy

experiments, respectively. Neurons were plated in plating media [neurobasal media (Gibco #21103049)

supplemented with 2.5% FBS, 2.5% horse serum, 2 mM L-glutamine, 1X penicillin-streptomycin, 2%

B27]. The next day, plating media was replaced with neuronal media (neurobasal media supplemented

with 2 mM L-glutamine, 1X penicillin-streptomycin, 2% B27). Neurons were processed at 17 days *in*

vitro.

2.10 Statistics

234 Statistical analyses and biological repeats are indicated in figure legends. Significant outliers identified

235 using the Grubbs' test were removed and are indicated in figure legends. GraphPad Prism 10 was used

236 for all statistical analyses and graph preparation. Graphs are presented as the standard error of the

237 mean.

3. Results

3.1 Vorinostat increases SQSTM1 palmitoylation but does not affect total protein
palmitoylation

An acyl-biotin exchange (ABE) assay was performed on HeLa cells overexpressing SQSTM1-HA and

treated with 10 µM Vorinostat or 10 µM PalmB. PalmB is a non-specific inhibitor of depalmitoylating

enzymes[34], whereas 2BP is a non-specific inhibitor for palmitoylation[35]. The absence of signal in the

hydroxylamine (HAM-) lane confirms that without thioester bond cleavage, biotin-HPDP cannot be

incorporated on free cysteine residues. Vorinostat treatment led to a significant increase in SQSTM1

palmitoylation (*$p < 0.05$) (Figure 2A,B), while PalmB treatment led to no changes.

Next, we performed a click chemistry assay on HeLa cells treated with 10 µM Vorinostat, 10 µM ML348,

10 µM PalmB, or 50 µM 2BP. ML348 is a specific inhibitor for APT1[34], the enzyme that depalmitoylates

SQSTM1[15,24]. Vorinostat treatment led to no changes in total protein palmitoylation (Figure 2C,D).

ML348 and 2BP treatments led to a slight decrease in total protein palmitoylation, whereas PalmB led to

a slight increase. The absence of a signal in the cold palmitate (palm) lane confirms that without an

alkyne group, biotin-azide cannot be incorporated on the palmitate.

**Figure 2 – Vorinostat increases SQSTM1 palmitoylation but does not affect total protein
palmitoylation. (A)** SQSTM1 palmitoylation was detected via acyl biotin exchange and is sensitive to
Vorinostat treatment in HeLa cells. Composite of the same gel. **(B)** Quantification of (top) SQSTM1
palmitoylation and (bottom) SQSTM1 levels from five biological replicates. Kruskal-Wallis test followed
by Dunn's post-hoc analysis. **(C)** Total palmitoylated protein was detected via click chemistry and is not
sensitive to Vorinostat treatment. Composite of the same gel. **(D)** Quantification of total palmitoylated
protein from three biological replicates. Kruskal-Wallis test followed by Dunn's post-hoc analysis.

22-month-old YAC128 mice were administered 50 mg/kg Vorinostat via intraperitoneal injection once

every second day for four consecutive weeks. Subsequently, an ABE was performed on approximately 30

mg of ground cortices. Vorinostat treatment in YAC128 mice led to a significant increase in SQSTM1 and

VCP palmitoylation (*p<0.05) compared to YAC128 DMSO-treated and WT Vorinostat-treated mice

268   (Figure 3A,B). Vorinostat treatment in WT mice led to no changes in SQSTM1 and VCP palmitoylation.

269   Vorinostat treatment led to no changes in total protein palmitoylation in both WT and YAC128 mice.

Figure 3 – Vorinostat crosses the blood-brain barrier and increases SQSTM1 palmitoylation in YAC128 mice. (A) SQSTM1 palmitoylation in cortices was detected via acyl biotin exchange and is sensitive to Vorinostat treatment in YAC128 mice. Composite of the same gel. **(B)** Quantification of (top) SQSTM1 palmitoylation, (second) VCP palmitoylation, (third) SQSTM1 levels, (fourth) VCP levels, and (bottom) total palmitoylated protein from three biological replicates. Two-way ANOVA test followed by Fisher's LSD post-hoc analysis (*p<0.05).

3.2 Vorinostat enhances autophagy and loading of mHTT aggregates into autophagosomes

In autophagy, SQSTM1 is degraded in autophagolysosomes, and while some of LC3-II located on the inner surface is also degraded, the majority located on the outer surface is recycled back to LC3-I, the inactive delipidated form of LC3[36]. This makes SQSTM1 and LC3-II excellent markers of autophagy, such that increased autophagic flux is indicated by reduced SQSTM1 levels and elevated LC3-II levels[36]. BafA1 is a late-stage autophagy inhibitor that prevents autophagosome-lysosome fusion leading to an increase in both SQSTM1 and LC3-II levels[36]. BafA1 is used to confirm that the effect of the drug treatment on autophagic flux is indeed a result of increased autophagy[36]. For example, if SQSTM1 levels decrease and LC3-II levels increase with drug treatment, but do not substantially increase with drug plus BafA1 treatment, this suggests that the drug treatment blocks autophagic flux, rather than increases it[36].

HeLa cells were treated with 10 μM Vorinostat, 10 μM ML348, or 10 μM PalmB, with or without BafA1. Vorinostat treatment led to a significant decrease in SQSTM1 levels ($*p < 0.05$) (Figure 4A,B). In addition, Vorinostat treatment significantly increased LC3-II levels ($*p < 0.05$). ML348 and PalmB treatment led to no changes in SQSTM1 or LC3-II levels. All BafA1 treatment conditions led to a significant increase in LC3-II ($**p < 0.01$) and SQSTM1 ($***p < 0.001$).

Furthermore, a key characteristic of autophagy is the colocalization of SQSTM1, HTT, and lysosomes, which suggests that SQSTM1 is successfully delivering cargo (*e.g.,* mHTT) with the support of HTT to autophagosomes, which fuse with lysosomes to form autophagolysosomes. Therefore, increased colocalization between these proteins and organelles, as measured by the PCC, indicates enhanced autophagic flux, as the presence of SQSTM1 and HTT in lysosomes signifies their successful delivery. HeLa cells were treated with 10 μM Vorinostat, 10 μM ML348, or 10 μM PalmB and visualized by fixed

cell confocal microscopy. When treated with Vorinostat, ML348, or PalmB, SQSTM1 and HTT

colocalization significantly increased, with a more pronounced increase elicited by Vorinostat

(****$p<0.0001$) (Figure 4C,D). When treated with Vorinostat or PalmB, SQSTM1 and lysosome

colocalization significantly increased, with a more pronounced increase elicited by Vorinostat

(****$p<0.0001$). Finally, when treated with Vorinostat or PalmB, HTT and lysosome colocalization

significantly increased, with a more pronounced increase elicited by Vorinostat (****$p<0.0001$). In

another experiment, primary cortical neurons derived from WT and YAC128 mice were treated with 10

μM Vorinostat, 10 μM ML348, or 10 μM PalmB and visualized by fixed cell confocal microscopy. In WT

cortical neurons treated with Vorinostat, SQSTM1 and HTT colocalization was not affected (Figure 5A,C).

However, in YAC128 cortical neurons treated with Vorinostat, SQSTM1 and mHTT colocalization

significantly increased (***$p<0.001$). Similarly, SQSTM1 and lysosome colocalization was not affected

with Vorinostat treatment in WT cortical neurons but significantly increased (**$p<0.01$) in YAC128

cortical neurons. In addition, HTT and lysosome colocalization, which includes both HTT and mHTT, was

not affected with Vorinostat treatment in WT cortical neurons but significantly increased

(****$p<0.0001$) in YAC128 cortical neurons, which is especially relevant in this context where

colocalization suggests that mHTT aggregates are in autophagolysosomes. When treated with ML348,

WT cortical neurons showed significantly increased SQSTM1 and mHTT colocalization (**$p<0.01$),

whereas YAC128 cortical neurons showed significantly increased SQSTM1 and mHTT colocalization, as

well as HTT and lysosome colocalization (***$p<0.001$ and **$p<0.01$, respectively). When treated with

PalmB, WT cortical neurons showed significantly increased SQSTM1 and mHTT, as well as HTT and

lysosome colocalization (**$p<0.01$ and ****$p<0.0001$, respectively), whereas YAC128 cortical neurons

showed significantly increased HTT and lysosome colocalization (***$p<0.001$).

Figure 4 – Vorinostat increases autophagic flux. (A) SQSTM1 and LC3-II levels were detected and are sensitive to Vorinostat treatment in HeLa cells. **(B)** Quantification of (left) SQSTM1 levels and (right) LC3-II levels from three biological replicates. Kruskal-Wallis test followed by Dunn's post-hoc analysis (*p<0.05) within +/- BafA1 conditions. Two-way ANOVA test followed by Fisher's LSD post-hoc analysis (**p<0.01, ***p<0.001) across BafA1 conditions. **(C)** Vorinostat affects SQSTM1, HTT, and lysosome colocalization levels in HeLa cells. **(D)** Quantification of (left) PCC between SQSTM1 and HTT, (middle) PCC between SQSTM1 and lysosome, and (right) PCC between HTT and lysosome, 15 cells per condition from three biological replicates. One-way ANOVA test followed by Dunnett's post-hoc analysis (*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001). Box and whiskers (min to max).

Figure 5 – Vorinostat redirects mHTT and SQSTM1 to lysosomes in HD neurons. (A) SQSTM1, HTT, and lysosome colocalization in primary cortical neurons derived from WT mice. **(B)** Vorinostat significantly increases SQSTM1 and mHTT co-localization, and lysosome colocalization with SQSTM1 and mHTT in primary cortical neurons derived from YAC128 HD mouse embryos. **(C)** Quantification of (top) PCC between SQSTM1 and HTT, (middle) PCC between SQSTM1 and lysosome, and (bottom) PCC between HTT and lysosome, on an average of 15 neurons per condition from three biological replicates. Two-way ANOVA test followed by Dunnett's post-hoc analysis (**$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$). Box and whiskers (min to max).

3.3 Vorinostat may function as a depalmitoylation inhibitor and transcriptional regulator

SQSTM1 has been shown to be depalmitoylated by APT1[15,24]. Thus, our initial theory was Vorinostat may function via inhibition of APT1, thereby increasing SQSTM1 palmitoylation. To further investigate, molecular docking experiments were performed using AutoDock Vina. These results indicate that ML348, an inhibitor specific to APT1[34], and Vorinostat have a deep fit into the binding pocket of APT1, with ML348 exhibiting a stronger binding affinity (-10.6 kcal/mol) compared to Vorinostat (-8.0 kcal/mol) (Figure 6A). We also looked at APT2 for comparison and found that ML349, an inhibitor specific to APT2[34], and Vorinostat have a deep fit into the binding pocket of APT2, with ML349 exhibiting a stronger binding affinity (-12.9 kcal/mol) compared to Vorinostat (-8.0 kcal/mol). For context, the more negative the value, the stronger the binding affinity.

Alternatively, Vorinostat is already known as an HDAC inhibitor, so its function could be through increasing the transcription of palmitoylating enzymes or decreasing the transcription of depalmitoylating enzymes. To further investigate, qPCR was performed on HeLa cells treated with 10 μM Vorinostat to quantify the relative transcript levels of *ZDHHC1-24*, *SQSTM1*, *HTT*, *VCP*, *LYPLA1* (APT1), *LYPLA2* (APT2), *PPTs*, *ABHDs*, *HDACs*, and *MAP1LC3s* (Figure 6B,C). Compared to DMSO, treatment with Vorinostat led to significantly increased *ZDHHC11*, *ZDHHC23*, and *SQSTM1* transcript levels, as well as significantly decreased transcript levels of *ZDHHC2*, *ZDHHC4*, *ZDHHC5*, *ZDHHC9*, *ZDHHC16*, *ZDHHC17*, *ZDHHC18*, *ZDHHC20*, *ZDHHC21*, *VCP*, *LYPLA2*, *PPT1*, *PPT2*, *ABHD16A*, and *ABHD17C*. Notably, *HTT*, *LYPLA1*, and *ZDHHC19* transcript levels were unaffected by Vorinostat treatment. ZDHHC22 had amplification in the Vorinostat condition, but not DMSO condition, whereas ABHD17A and MAP1LC3A showed no amplification in either condition.

Figure 6 – Vorinostat may function as a depalmitoylation inhibitor and transcriptional regulator. (A) Vorinostat and ML348 molecular docking and binding affinity on APT1 and APT2 were predicted using AutoDock Vina. (B) *ZDHHC1-24* transcript levels were detected via qPCR in HeLa cells. *ZDHHC2,4,5,9,11,16,17,18,20,21,23* transcript levels are sensitive to Vorinostat treatment. One significant outlier (SO) removed from ZDHHC11 and ZDHHC23. Two-tailed unpaired *t*-test (*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001). (C) *SQSTM1*; *HTT*; *VCP*; *LYPLA1,2*; *LYPLAL1*; *PPT1,2*; *ABHD10,16A,17A,17B,17C*; *HDAC1,2,3,6*; and *MAP1LC3A,B* transcript levels were detected via qPCR in HeLa cells. *SQSTM1, VCP, LYPLA2, PPT1, PPT2, ABHD16A,* and *ABHD17C* transcript levels are sensitive to Vorinostat treatment. One SO removed from LYPLA2. Two-tailed unpaired *t*-test (*p<0.05, ***p<0.001).

4. Discussion

A hallmark of HD is the cytotoxic accumulation and aggregation of mHTT, which is partly caused and subsequently exacerbated by dysfunctional autophagy. Our focus is failure at the cargo-loading step, wherein SQSTM1 is unable to deliver cargo, mainly, mHTT aggregates, to the phagophore membrane for degradation. Since palmitoylation plays an important role in protein localization to membranes, we predict that the autophagic dysfunction seen in HD may be a result of reduced SQSTM1 palmitoylation, which our group has confirmed previously in the brains of HD patients and YAC128 mice. Consequently, we aim to promote mHTT intracellular degradation, thereby preventing cytotoxicity and neuronal death, by restoring SQSTM1 palmitoylation. Through a high-throughput screening of FDA-approved drugs, we identified blood-brain permeable Vorinostat as a candidate to increase SQSTM1 palmitoylation.

We found that Vorinostat treatment significantly increases SQSTM1 palmitoylation does not affect total protein palmitoylation both *in vitro* (Figure 2) and *in vivo* (Figure 3). Since the biochemical assays for our *in vivo* work were conducted on cortical tissue, this confirms that Vorinostat successfully crosses the blood-brain barrier. Notably, PalmB did not increase SQSTM1 palmitoylation *in vitro*. This may be because PalmB is a non-selective inhibitor of depalmitoylating enzymes. Since many palmitoylating enzymes undergo regulatory palmitoylation (*e.g.,* ZDHHC6 is palmitoylated by ZDHHC16[37] and ZDHHC5 is palmitoylated by ZDHHC20[38]), broad inhibition of depalmitoylating enzymes could disrupt the function of palmitoylating enzymes, potentially explaining the lack of effect of PalmB on SQSTM1 palmitoylation.

With an increase in SQSTM1 palmitoylation induced by Vorinostat treatment, we then observed a corresponding increase of autophagic flux. We found that Vorinostat treatment significantly decreases SQSTM1 levels and increases LC3-II levels (Figure 4). SQSTM1 and LC3-II are autophagy markers, and their respective decrease and increase in levels suggests increased autophagic flux, since SQSTM1 is degraded in autophagolysosomes, whereas the majority of LC3-II is recycled back into the inactive

delipidated form of LC3. Increased autophagic flux was confirmed with Vorinostat plus BafA1 treatment,

a late-stage autophagy inhibitor[36], which significantly increased SQSTM1 and LC3-II levels.

Furthermore, with an increase in SQSTM1 palmitoylation induced by Vorinostat treatment, we then

observed a corresponding enhancement of autophagy. We found that Vorinostat treatment significantly

increased SQSTM1, mHTT, and lysosome colocalization in HeLa cells (Figure 4), as well as primary

cortical neurons derived from YAC128 mice, but not WT mice (Figure 5). In HeLa cells, this suggests

enhanced SQSTM1 and HTT delivery to autophagosomes. In YAC128 primary cortical neurons, increased

SQSTM1, mHTT, and lysosome colocalization suggests improved delivery of cytotoxic mHTT aggregates

to autophagosomes for subsequent degradation in autophagolysosomes, repairing the hallmark

neuropathological feature of HD. Notably, the observation that Vorinostat had an effect in YAC128 but

not in WT primary cortical neurons is important as it suggests that Vorinostat may exert its effects

selectively under pathological conditions, reducing the risk of potential off-target effects. The effects of

ML348 and PalmB on SQSTM1, mHTT, and lysosome colocalization were neither as strong, consistent,

nor specific as those of Vorinostat, suggesting that Vorinostat may be a more effective drug in HD.

419
Initially, we predicted that Vorinostat increases SQSTM1 palmitoylation by acting as a depalmitoylation

inhibitor, specifically of APT1, the depalmitoylating enzyme of SQSTM1. Consequently, we aimed to

better understand Vorinostat's ability to inhibit depalmitoylating enzymes through molecular docking

experiments to quantify Vorinostat's binding affinity to APT1 and APT2 (Figure 6). Although Vorinostat

had a deep fit into the binding pockets of both APT1 and APT2, its binding affinity to either was weaker

than that of the specific inhibitors, ML348 and ML349. However, prior experiments indicated that

Vorinostat was more effective than ML348 at increasing SQSTM1 palmitoylation, thus prompting us to

reassess our previous thinking. Vorinostat is also a known HDAC inhibitor, promoting histone acetylation

or transcription factor acetylation, thereby increasing transcription. Either way, Vorinostat could

increase the transcription of *ZDHHC* genes, leading to increased SQSTM1 palmitoylation through an

alternative pathway. Therefore, Vorinostat may be acting via a dual mechanism, as summarized in

Figure 7.

Figure 7 – Predicted Vorinostat mechanism. Vorinostat may be working through two concurrent
mechanisms to increase autophagy protein palmitoylation: inhibition of depalmitoylating (depalm)
enzymes and inhibition of acetylation enzymes (HDAC). Inhibition of depalmitoylating enzymes prevents
the depalmitoylation of SQSTM1, thus increasing SQSTM1 palmitoylation. In contrast, inhibition of HDAC
prevents deacetylation of histones, thus increasing their acetylation, promoting active transcription by
opening chromatin. Alternatively, inhibition of HDAC prevents deacetylation of transcription factors (TF),
thus increasing their acetylation, also promoting active transcription. K represents a lysine residue
protruding from the histone, and Ac represents an acetyl group. This may increase transcription of
palmitoylating (palm) enzymes or decrease the transcription of depalmitoylating enzymes, overall
increasing autophagy protein palmitoylation. Alternatively, this may increase or decrease transcription
of relevant proteins, such as SQSTM1, which may also enhance autophagy. Figure generated on
BioRender.

The next step was to confirm whether Vorinostat increases transcription of relevant proteins by

quantifying transcript levels via qPCR. We found that Vorinostat significantly increases *SQSTM1*

transcript levels compared to DMSO, suggesting that Vorinostat may promote autophagy. However,

Vorinostat plus BafA1 did not lead to an increase in SQSTM1 levels (Figure 4). It could be that Vorinostat

plus BafA1 leads to the accumulation of detergent-insoluble SQSTM1, as described by the Yang group[24],

which may limit detection of SQSTM1 via western blot. In addition, we found that Vorinostat

significantly increases *ZDHHC11* and *ZDHHC23* transcript levels compared to DMSO. Work by the Liu

group suggests that ATG2A is palmitoylated by ZDHHC11 and its *depalmitoylation* is required for proper

autophagosome formation[39]. In contrast, studies by the Kim group suggest that MCOLN3/TRPML3 is

palmitoylated by ZDHHC11 and its *palmitoylation* is required for proper autophagosome formation[40].

Minimal connections between ZDHHC23 and autophagy have been established. Therefore, it remains

unclear how Vorinostat-induced increases in the *ZDHHC11* and *ZDHHC23* transcript levels might have a

role in the increase and enhanced autophagy we observed following Vorinostat treatment. Furthermore,

we found that Vorinostat decreases *LYPLA2* transcript levels which encode APT2. APT2 depalmitoylates

HTT, and HTT palmitoylation is reduced in HD mouse models[18]. Therefore, reducing APT2 may increase

HTT palmitoylation, thereby lowering mHTT aggregation and cytotoxicity. Although increased transcript

levels do not necessarily equate to increased protein levels, they serve as a good representation of

potential regulatory changes at the transcriptional level and can provide insight into upstream effects of

Vorinostat treatment.

4.1 Vorinostat and Huntington disease

Vorinostat has been tested in HD mouse models previously, but in the context of its HDAC inhibiting

properties and fixing transcriptional dysregulation found in HD. The Bates group treated R6/2 HD mouse

model and found improved motor impairment, reduced mHTT aggregates in the cortex, and restored

Bdnf transcript levels[33,41]. BDNF is implicated in HD pathogenesis[42]. In R6/2 mice, *Bdnf* transcript levels

decline during disease progression[43]; accordingly, BDNF over-expression was found to improve HD

phenotypes[44]. The Bates group also found that Vorinostat decreases HDAC2 and HDAC4 levels *in vivo*[41].

In R6/2 mice, the group reported that HDAC4 associates with mHTT in a polyQ-dependent manner and

co-localizes with cytoplasmic inclusions, and knockdown of HDAC4 reduced cytoplasmic aggregate

formation, restored *Bdnf* transcript levels, rescued membrane properties of MSNs and of corticostriatal

synaptic transmission, improved motor and neurological impairment, and extended the lifespan of R6/2

mice[45]. To further understand the mechanism of HDAC4, the Cristea group studied the interaction of

HDAC4 in the context of HD through multiomics and propose a potential model of the role for HDAC4 in

HD[46]. One notable finding from their study is the interaction between HDAC4 and VCP, which they also

found has minimal interaction in the Q20 WT mice and maximal interaction in the Q140 HD mice, an

interaction which increases with age. Combining this with our findings on VCP, it is possible that the

increased interaction between HDAC and VCP, and likely VCP acetylation[47], interferes with VCP

palmitoylation, thereby disrupting its normal function. By deceasing HDAC4 levels with Vorinostat, VCP

function may be restored.

4.2 VCP and Huntington disease

Our group has recently shown that VCP palmitoylation is significantly reduced in the brains of HD mouse

models. It is still unclear the effect of decreasing VCP palmitoylation in the context of HD; however, we

looked at whether Vorinostat also rescues VCP palmitoylation in addition to SQSTM1 palmitoylation. We

found that *in vivo* Vorinostat treatment significantly increases VCP palmitoylation. VCP is also involved in

autophagy initiation, and since autophagy membrane-dependent process, this suggests that VCP

palmitoylation may modulate its function in this pathway. Links between VCP and HD have already been

established. The Qi group identified VCP as a mHTT-binding protein on mitochondria, where it drives

excessing mitophagy, or the autophagic degradation of mitochondria[48]. Because palmitoylation is a key

regulator of protein localization, restoring VCP palmitoylation could potentially correct its

mislocalization, reducing VCP accumulation on mitochondria in HD mouse models.

4.3 Future Directions

Although studies have demonstrated positive outcomes of Vorinostat treatment in HD mouse models,

the broader consequences of HDAC inhibition remain an important consideration. Determining the

specific mechanism by which Vorinostat mediates these positive effects should be a priority, as this

knowledge could guide the design of more targeted small molecules. However, it is also possible that

Vorinostat's multi-functional nature contributes to its broad impact on various HD phenotypes, whether

through restoring protein palmitoylation, correcting transcriptional dysregulation, or both. This raises an

important question: in a devastating disease like HD, at what point do the positive effects of Vorinostat

outweigh the risks of potential off-target effects? If nothing else, Vorinostat may provide insight into HD

pathology and guide the identification of novel therapeutic targets, including protein palmitoylation and

autophagy. In this study, we reported that Vorinostat crossed the blood-brain barrier and increased

SQSTM1 palmitoylation *in vivo*, as well as increased and enhanced autophagy *in vitro*, facilitating mHTT

loading into autophagolysosomes for degradation.

Author Contributions:

YA prepared the first draft. YA prepared Figures 1-7. YA completed the experiments in Figures 2-6. FA contributed to the SQSTM1 palmitoylation experiment in Figure 2. MR contributed to the qPCR experiments in Figure 6. AD contributed to the docking experiments in Figure 6. FR helped with animal studies including preparation of primary neuronal cultures. YA and DDOM edited the final draft.

Funding:

DDOM is supported by a Natural Sciences and Engineering Research Council Discovery Grant (RGPIN-2019-04617) and Huntington Society of Canada Navigator Research Award. YA is supported by a Canadian Institutes of Health Research Canada Graduate Scholarship Master's Award, University of Waterloo President's Graduate Scholarship, Brain Canada Rising Star in ALS Research Award, and John E. Thompson Biology Graduate Scholarship. AD is supported by a Huntington Society of Canada Undergraduate Summer Fellowship. FR is supported by a Mitacs Accelerate Grant in collaboration with Circumvent Pharmaceuticals Inc., Canadian Institutes of Health Research Postdoctoral Research Award, and an Uplifting Athletes Young Investigator Draft Award in collaboration with Cure VCP Disease Inc.

Conflicts of Interest:

The authors declare no conflicts of interest.

Acknowledgements:

We would like to acknowledge that the University of Waterloo resides on the traditional territory of five Indigenous communities, including the Ho-de-no-sau-nee-ga (Haudenosaunee), Mississaugas of the Credit First Nation, Anishinabewaki, Attiwonderonk (Neutral), and Mississauga peoples, which is situated on the Haldimand Tract, a land granted to the Six Nations, encompassing six miles on either side of the Grand River. This land is home to many past, present, and future Indigenous peoples.

References

1. Fujikake, N., Shin, M. & Shimizu, S. Association Between Autophagy and Neurodegenerative Diseases. *Front Neurosci* **12**, 255 (2018).

2. Ramzan, F., Abrar, F., Mishra, G. G., Liao, L. M. Q. & Martin, D. D. O. Lost in traffic: consequences of altered palmitoylation in neurodegeneration. *Front. Physiol.* **14**, 1166125 (2023).

3. Wlodarczyk, J. *et al.* Altered Protein Palmitoylation as Disease Mechanism in Neurodegenerative Disorders. *J. Neurosci.* **44**, e1225242024 (2024).

4. Parzych, K. R. & Klionsky, D. J. An Overview of Autophagy: Morphology, Mechanism, and Regulation. *Antioxid Redox Signal* **20**, 460–473 (2014).

5. Pircs, K. *et al.* Distinct subcellular autophagy impairments in induced neurons from patients with Huntington's disease. *Brain* **145**, 3035–3057 (2022).

6. Martinez-Vicente, M. *et al.* Cargo recognition failure is responsible for inefficient autophagy in Huntington's disease. *Nat Neurosci* **13**, 567–576 (2010).

7. Walter, C. *et al.* Activation of AMPK-induced autophagy ameliorates Huntington disease pathology in vitro. *Neuropharmacology* **108**, 24–38 (2016).

8. Ghosh, R. & Tabrizi, S. J. Clinical Features of Huntington's Disease. in *Polyglutamine Disorders* (eds Nóbrega, C. & Pereira de Almeida, L.) 1–28 (Springer International Publishing, Cham, 2018). doi:10.1007/978-3-319-71779-1_1.

9. The Huntington's Disease Collaborative Research Group. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell* **72**, 971–983 (1993).

10. Kay, C. *et al.* Huntington disease reduced penetrance alleles occur at high frequency in the general population. *Neurology* **87**, 282–288 (2016).

11. Saudou, F. & Humbert, S. The Biology of Huntingtin. *Neuron* **89**, 910–926 (2016).

12. Ochaba, J. *et al.* Potential function for the Huntingtin protein as a scaffold for selective autophagy.

Proc Natl Acad Sci U S A **111**, 16889–16894 (2014).

13. Rui, Y.-N. *et al.* Huntingtin Functions as a Scaffold for Selective Macroautophagy. *Nat Cell Biol* **17**,

262–275 (2015).

14. Martin, D. D. O., Ladha, S., Ehrnhoefer, D. E. & Hayden, M. R. Autophagy in Huntington disease and

huntingtin in autophagy. *Trends in Neurosciences* **38**, 26–35 (2015).

15. Abrar, F. *et al.* Reduced Palmitoylation of SQSTM1/p62 in Huntington Disease Is Associated With

Impaired Autophagy. *FASEB J* **39**, e70549 (2025).

16. Alshehabi, Y., Abrar, F. & Martin, D. D. O. Sequestering sequestosome 1 via S-acylation in

autophagy, Huntington disease, and beyond. *Autophagy Rep* **4**, 2547975 (2025).

17. Yanai, A. *et al.* Palmitoylation of huntingtin by HIP14is essential for its trafficking and function. *Nat*

Neurosci **9**, 824–831 (2006).

18. Lemarié, F. L. *et al.* Rescue of aberrant huntingtin palmitoylation ameliorates mutant huntingtin-

induced toxicity. *Neurobiology of Disease* **158**, 105479 (2021).

19. Chamberlain, L. H. & Shipston, M. J. The physiology of protein S-acylation. *Physiol Rev* **95**, 341–376

(2015).

20. Anwar, M. U. & van der Goot, F. G. Refining S-acylation: Structure, regulation, dynamics, and

therapeutic implications. *Journal of Cell Biology* **222**, e202307103 (2023).

21. Lemarié, F. L., Sanders, S. S., Nguyen, Y., Martin, D. D. O. & Hayden, M. R. Full-length huntingtin is

palmitoylated at multiple sites and post-translationally myristoylated following caspase-cleavage.

Front Physiol **14**, 1086112 (2023).

22. Martin, D. D. O. & Sanders, S. S. Let's get fat: emergence of *S*-acylation as a therapeutic target in

Huntington disease. *Biochemical Society Transactions* BST20231290 (2024)

doi:10.1042/BST20231290.

23. Martin, D. D. O. & Sanders, S. S. Let's get fat: emergence of S-acylation as a therapeutic target in

Huntington disease. *Biochemical Society Transactions* **52**, 1385–1392 (2024).

24. Huang, X. *et al.* S-acylation of p62 promotes p62 droplet recruitment into autophagosomes in

mammalian autophagy. *Mol Cell* **83**, 3485-3501.e11 (2023).

25. Virlogeux, A. *et al.* Increasing brain palmitoylation rescues behavior and neuropathology in

Huntington disease mice. *Science Advances* **7**, (2021).

26. Martin, D. D. O. *et al.* Identification of Novel Inhibitors of DLK Palmitoylation and Signaling by High

Content Screening. *Sci Rep* **9**, 3632 (2019).

27. Athira, K. V., Sadanandan, P. & Chakravarty, S. Repurposing Vorinostat for the Treatment of

Disorders Affecting Brain. *Neuromol Med* **23**, 449–465 (2021).

28. Verrillo, L., Di Palma, R., de Bellis, A., Drongitis, D. & Miano, M. G. Suberoylanilide Hydroxamic Acid

(SAHA) Is a Driver Molecule of Neuroplasticity: Implication for Neurological Diseases. *Biomolecules*

13, 1301 (2023).

29. Liao, L. M. Q., Gray, R. A. V. & Martin, D. D. O. Optimized Incorporation of Alkynyl Fatty Acid Analogs

for the Detection of Fatty Acylated Proteins using Click Chemistry. *J Vis Exp*

https://doi.org/10.3791/62107 (2021) doi:10.3791/62107.

30. Hayashi, T., Thomas, G. M. & Huganir, R. L. Dual Palmitoylation of NR2 Subunits Regulates NMDA

Receptor Trafficking. *Neuron* **64**, 213–226 (2009).

31. Trott, O. & Olson, A. J. AutoDock Vina: improving the speed and accuracy of docking with a new

scoring function, efficient optimization and multithreading. *J Comput Chem* **31**, 455–461 (2010).

32. McClafferty, H. & Shipston, M. J. siRNA Knockdown of Mammalian zDHHCs and Validation of mRNA

Expression by RT-qPCR. in *Protein Lipidation* (ed. Linder, M. E.) vol. 2009 151–168 (Springer New

York, New York, NY, 2019).

33. Hockly, E. *et al.* Suberoylanilide hydroxamic acid, a histone deacetylase inhibitor, ameliorates motor

deficits in a mouse model of Huntington's disease. *Proc Natl Acad Sci U S A* **100**, 2041–2046 (2003).

34. Virlogeux, A. *et al.* Increasing brain palmitoylation rescues behavior and neuropathology in

Huntington disease mice. *Sci Adv* **7**, eabb0799 (2021).

35. Davda, D. *et al.* Profiling targets of the irreversible palmitoylation inhibitor 2-bromopalmitate. *ACS*

Chem Biol **8**, 1912–1917 (2013).

36. Klionsky, D. J. *et al.* Guidelines for the use and interpretation of assays for monitoring autophagy

(4th edition)1. *Autophagy* **17**, 1–382 (2021).

37. Abrami, L. *et al.* Identification and dynamics of the human ZDHHC16-ZDHHC6 palmitoylation

cascade. *eLife* **6**, e27826 (2017).

38. Plain, F. *et al.* Control of protein palmitoylation by regulating substrate recruitment to a zDHHC-

protein acyltransferase. *Commun Biol* **3**, 411 (2020).

39. Zheng, W. *et al.* S-palmitoylation modulates ATG2-dependent non-vesicular lipid transport during

starvation-induced autophagy. *EMBO J* **44**, 2596–2619 (2025).

40. Kim, S. W. *et al.* Palmitoylation controls trafficking of the intracellular Ca2+ channel

MCOLN3/TRPML3 to regulate autophagy. *Autophagy* **15**, 327–340 (2018).

41. Mielcarek, M. *et al.* SAHA decreases HDAC 2 and 4 levels in vivo and improves molecular

phenotypes in the R6/2 mouse model of Huntington's disease. *PLoS One* **6**, e27746 (2011).

42. Zuccato, C. & Cattaneo, E. Role of brain-derived neurotrophic factor in Huntington's disease. *Prog*

Neurobiol **81**, 294–330 (2007).

43. Benn, C. L., Fox, H. & Bates, G. P. Optimisation of region-specific reference gene selection and

relative gene expression analysis methods for pre-clinical trials of Huntington's disease. *Mol*

Neurodegener **3**, 17 (2008).

44. Gharami, K., Xie, Y., An, J. J., Tonegawa, S. & Xu, B. Brain-derived neurotrophic factor over-

expression in the forebrain ameliorates Huntington's disease phenotypes in mice. *J Neurochem* **105**,

369–379 (2008).

45. Mielcarek, M. *et al.* HDAC4 reduction: a novel therapeutic strategy to target cytoplasmic huntingtin

and ameliorate neurodegeneration. *PLoS Biol* **11**, e1001717 (2013).

46. Federspiel, J. D., Greco, T. M., Lum, K. K. & Cristea, I. M. Hdac4 Interactions in Huntington's Disease

Viewed Through the Prism of Multiomics. *Mol Cell Proteomics* **18**, S92–S113 (2019).

47. Mori-Konya, C. *et al.* p97/valosin-containing protein (VCP) is highly modulated by phosphorylation

and acetylation. *Genes Cells* **14**, 483–497 (2009).

48. Guo, X. *et al.* VCP recruitment to mitochondria causes mitophagy impairment and

neurodegeneration in models of Huntington's disease. *Nat Commun* **7**, 12646 (2016).