1   Beyond Fixation: Persistent Genetic Variation Under Intense Selection

3   **Authors**: Kenneth R. Arnold[1*], Zachary S. Greenspan[2], Ryan D. Robinson[1], Anastasia Pupo[3],

4   Valeria V. Chavarin[4], Kevin S. Chang[2], Corrina O. Cannell[2], Miao Qi[2], Laurence D. Mueller[2],

5   Michael R. Rose[2], and Mark A. Phillips[1*]

7   [1]Department of Integrative Biology, Oregon State University

9   [2]Department of Ecology and Evolutionary Biology, University of California, Irvine

11   [3] Khoury College of Computer Sciences, Northeastern University

13   [4]Department of Biological Science, California State University, Fullerton

15   [*]Corresponding authors: Kenneth Arnold (kenneth.arnold@oregonstate.edu) and Mark

16   Phillips (philmark@oregonstate.edu)

Keywords: Experimental evolution, Polygenic adaptation, Balancing selection, Antagonistic

pleiotropy, Evolutionary reversibility, Standing genetic variation, Life-history evolution, *Drosophila*

melanogaster

Abstract

Understanding how and why genetic variation is maintained under sustained selection remains a

central question in evolutionary genetics. Experimental evolution shows that adaptation in

sexually reproducing populations is often highly polygenic, proceeding through coordinated,

genome-wide allele frequency shifts from standing variation rather than classic hard sweeps.

Recent explanations emphasize highly polygenic architectures, optimizing-selection, and genetic

redundancy, which can slow fixation by distributing selection across many loci during adaptation.

However, observations from long-term selection experiments reveal a pattern these frameworks

do not fully explain: substantial genetic variation persists after hundreds of generations of intense

directional selection in constant environments. Here, we use long-term experimental evolution in

Drosophila melanogaster to test whether balancing-selection actively maintains genetic variation

under strong life-history selection and preserves evolutionary reversibility. Longstanding

populations selected for accelerated or delayed reproduction were shifted to the opposing regime,

imposing age-structured fitness trade-offs. Notably, selection for early reproduction is associated

with substantial loss of genetic variation, providing a stringent test of whether standing variation

is truly depleted. Following reciprocal shifts, populations showed rapid phenotypic convergence

toward the target regime. At the genomic level, allele-frequency trajectories were strongly

antiparallel and highly repeatable across replicates, revealing coordinated polygenic responses.

Relaxing long-standing early-life selection produced a pronounced rebound in genome-wide

heterozygosity. Deep sequencing uncovered ultra-rare alleles at sites appearing fixed under

standard coverage, indicating low-frequency functional variation persists below detection

thresholds. These results suggest that substantial genetic variation can persist under intense

directional selection and be rapidly redeployed when selection reverses, consistent with

widespread balancing-selection.

Introduction

Understanding how genetic variation is shaped and maintained in populations has long been a central question within evolutionary genetics (Lewontin 1974). A major puzzle is why substantial genetic variation often persists within sexually reproducing populations despite long-term, intense selection, under which classical models predict rapid fixation and loss of diversity. Evolution experiments in sexually reproducing organisms have repeatedly shown that responses to selection are often highly polygenic, fueled by standing genetic variation, and characterized by modest coordinated shifts in allele frequencies rather than classic hard sweeps in which selected alleles and linked sites fix (Burke 2012; Barghi and Schlötterer 2020). In mesocosm experiments mimicking natural conditions, the absence of sweeps and persistence of genetic variation is often attributed to fluctuating selection and adaptive tracking (Rudman et al. 2022, Bitter et al. 2024). However, we know that this explanation cannot be applied to laboratory based experimental evolution studies where environments and selection pressures are typically constant.

Experimental evolution combined with whole-genome sequencing ("E&R") is an established approach for studying the genetic basis of adaptation and complex trait variation (Long et al. 2015; Schlötterer et al. 2015). Across the *Drosophila* studies that comprise most of this work, the shift versus sweep dynamic is widespread (e.g. Burke et al. 2010; Tobler et al. 2014; Graves et al. 2017; Barghi et al. 2019, 2020). It is even observed in outcrossing yeast studies, where large population sizes and longer selection durations allow for a greater possibility of adaptation from de novo mutations (Burke et al. 2014; Phillips et al. 2020; Linder et al. 2022). Recent explanations for the "shifts not sweeps" pattern emphasize highly polygenic architectures, optimizing selection, and genetic redundancy (Chevin and Hospital 2008; Barghi et al. 2019; Hayward and Sella 2022). However, while this framework is compelling, it does not resolve a puzzling pattern that emerges in the case of long-term selection experiments, where populations experience sustained selection for many hundreds of generations. Although genetic redundancy can slow fixation by distributing selection across many loci during ongoing adaptation, it does not provide a mechanism for actively maintaining polymorphisms once populations approach a phenotypic optimum, at which point drift is expected to erode variation.

Evidence for this phenomenon comes from our own long-term *Drosophila* experimental evolution system, in which populations maintained under intense selection for over 1,000 generations retain substantially more genetic variation than expected under neutral drift and under forward simulations incorporating both directional and optimizing selection (Phillips et al. 2016). Why so much genetic variation persists after hundreds of generations of intense selection remains an

open question. Here we propose that pervasive antagonistic pleiotropy may play an important
role in maintaining genetic variation by constraining fixation and slowing the erosion of
polymorphism under sustained selection.

In previous work, we raised the possibility that the long-term persistence of genetic variation in
these populations reflects pervasive balancing selection acting within populations rather than
passive retention (Graves et al. 2017). This idea was motivated by a key result from that study:
despite many generations of selection on reproductive timing, populations recently derived from
the same ancestral stock rapidly converged phenotypically and genotypically with long-standing
selected lines. This repeatability implies that adaptive alleles are retained in the base populations
far longer than expected, given that hundreds of generations of selection and drift should
otherwise erode standing variation. Antagonistic pleiotropy provides a plausible mechanism:
alleles that enhance fitness through one life-history character may impose costs upon other
characters, constraining fixation, and in some cases, sustaining genetic polymorphism over time
(Rose 1982; Rose 1985).

Here we aim to empirically test whether widespread balancing selection contributes to the long-
term maintenance of genetic variation in Drosophila experimental evolution. We focus on
populations that have been under intense selection for early or late reproduction for hundreds of
generations and assess whether they retain the capacity to evolve back toward ancestral
phenotypic and genomic states when selection is reversed.

The early-reproduction populations, termed the A-types, are maintained on a 10-day generation
cycle, whereas the late-reproductive populations, the C-types, are maintained on a 28-day cycle
(Table S1). Both sets were ultimately descended from the same ancestral Ives base population
(Ives 1970; Rose et al. 2004). A-type selection produces accelerated development, reduced adult
lifespan, reduced stress resistance, increased early-life fecundity, and distinct metabolic profiles
(Burke et al. 2016; Kezos et al. 2023; Hubert et al. 2025). Genomic characterization further
indicates that these phenotypes are underlain by widespread genomic responses to selection,
including the apparent fixation of some polymorphic sites and regions of the genome where
variation is almost entirely purged. (At present, the longest-standing A-type populations have
experienced ~1,200 generations of selection for early reproduction, and the more recently derived
A-types ~650). Given that population sizes are on the order of a thousand, beneficial *de novo*
mutations are unlikely to be a factor driving adaptation and populations should have reached
mutation-selection-drift equilibrium under their respective selection regimes. Under standard
expectations when antagonistic pleiotropy is absent, most segregating variation should be

selectively neutral or weakly deleterious, slowly drifting toward fixation or loss. Thus, the ability of
A-type populations to move rapidly back toward C-type states when selection is altered would
imply that genetic variation been actively maintained at equilibrium frequencies, rather than
"passively" retained during incomplete selection.

To assess the possible conclusion described above, we use antiparallel selection shifts between
the A- and C-type regimes and follow two trajectories: A→C (A-types placed under C-type
conditions) and C→A (C-types placed under A-type conditions) illustrated in Figure 1. In doing so,
we both relax the intense early-reproduction selection experienced by A-types and impose this
same regime on C-types. Starting from these contrasting genetic backgrounds, we track
phenotypic and genomic changes over several dozen generations to evaluate how populations
respond to their new life-history cycles. If widespread balancing selection is maintaining functional
variation in this system, we expect that: (1) evolutionary history will not impose strong, persistent
constraints, and populations will converge genomically and phenotypically on long-standing
counterparts, where under our design, this should translate into inverted genomic trajectories for
the A→C and C→A given they are ultimately derived from the same ancestral population; (2)
shifting A-types to C-type conditions will result in a widespread rebound in genetic variation,
revealing low-frequency alleles that persisted despite the apparent loss of polymorphisms in past
work, and (3) replicate populations within each trajectory will show repeatable genomic
responses, consistent with access to shared reservoirs of selectively maintained variation.

Results

Adult Life-History Evolution: Mortality Trajectories

Founder populations exhibited the expected, long-established divergence in age-specific mortality between A-type and C-type selection regimes (Fig. S1, S2; Tables S2-10), providing internal baselines for evaluating phenotypic change along reciprocal trajectories. At the time of the first mortality assay, C→A populations had experienced approximately 33 generations of early-reproduction selection, whereas A→C populations had experienced approximately 12 generations of delayed-reproduction selection. At this stage, both trajectory types differed significantly from their respective founders but exhibited asymmetric convergence toward the target selection regime consistent with differences in total number of generations under selection (Fig. S3A). C→A populations showed strong convergence with A-type founders across much of the adult lifespan, particularly at early adult ages, whereas A→C populations remained substantially divergent from C-type founders, showing early phase mortality profiles (Tables S11-14).

At the next timepoint in the evolutionary trajectory, C→A populations had experienced approximately 143 generations of early-reproduction selection, whereas A→C populations had experienced approximately 33 generations of delayed-reproduction selection. Here we see both trajectory types have converged on the adult mortality phenotypes characteristic of long-term populations maintained under their respective selection regimes (Fig. S3B). Neither A→C nor C→A populations differed significantly from long-established populations of the same regime across adult life-history intervals, and both remained distinguishable from their original founders (Table S15-18).

To visualize systematic changes in adult mortality following reciprocal shifts in selection regime, we summarized mortality trajectories as the difference in mean age-specific mortality between the evolved populations and their regime founder counterparts, plotted as a function of age from egg. In both reciprocal selection treatments, differences in age-specific mortality relative to the founder populations exhibit a smooth, wave-like pattern across the adult lifespan (Fig. 2). For A→C populations, there is convergence on mortality at early ages until a breakpoint day, at which point, mortality differences begin to increase gradually at successive ages. This breakpoint is pushed towards later ages with more selection, indicating age-structured convergence toward C-type mortality profiles rather than uniform shifts across the lifespan.

Developmental Timing Responses

To determine whether developmental timing responded concordantly to reciprocal life-history selection, we assayed pupation timing along the same evolutionary trajectories. For the first assay, A→C populations (~6 generations of delayed-reproduction selection) and C→A populations (~15 generations of early-reproduction selection) exhibited pupation timing intermediate relative to their respective founders (Figs. S3, S4; Table S19). At the next timepoint, A→C (~64 generations) C→A (~182 generations), pupation timing in both trajectory types converged on the phenotypes characteristic of long-term populations maintained under their current selection regimes (Fig.S5; Table S20). Thus, developmental timing shows a consistent and reversible response to reciprocal life-history selection, corroborating patterns observed for adult mortality.

Antiparallel Genomic Trajectories

To characterize genome-wide shifts associated with reciprocal selection, we performed a principal component analysis (PCA) on SNP frequencies from samples taken over the course of the experiment (Fig. 3). The first two components captured most of the variance, with PC1 explaining 60.4% and PC2 explaining 11.0% of genome-wide variation; all higher-order PCs each explained <2%.

Projection onto the first two PCs revealed strongly antiparallel genomic trajectories. A→C populations shifted toward the cluster of long-established C-type populations, whereas C→A populations moved toward A-type populations along opposing directions. Dispersion among replicates was minimal at the final timepoints collected, whereas greater spread was observed among A→C populations at earlier stages relative to C→A populations, consistent with fewer generations having elapsed under delayed-reproduction selection.

To identify genomic axes that differ in their temporal response to selection, we analyzed the first ten principal components (PCs) using linear mixed-effects models with selection treatment (A→C vs. C→A) and generation as fixed effects and population as a random effect. The treatment × generation interaction tests whether PC scores change through time differently between treatments. Significant interactions were detected for PC1 (Table S21: estimate = 7.74, Bonferroni-adjusted $p = 6.26 \times 10^{-5}$) and PC2 (Table S21: estimate = −12.47, Bonferroni-adjusted $p = 2.0 \times 10^{-5}$), indicating opposing, generation-dependent shifts in these components between A→C and C→A populations. No significant interactions were detected for higher-order PCs, each

of which explained <2% of the variance. Thus, treatment-specific temporal genomic divergence
is concentrated along the first two principal components.

To focus on locus-specific responses, we quantified allele-frequency dynamics at individual SNPs
using beta-binomial generalized linear mixed models applied independently to 1,389,963 loci. In
these models, allele-frequency change was modeled as a function of treatment (A→C vs. C→A),
generation, and their interaction, with population included as a random effect. Under this
framework, significant treatment × generation effects identify SNPs exhibiting reciprocal
evolutionary responses (i.e. allele frequencies shifting in opposite directions between the groups).

After Bonferroni correction, we detected a widespread and highly polygenic genomic response to
reciprocal selection (Fig. 4). The majority of significant signal was attributable to treatment ×
generation interactions, with 134,709 SNPs exhibiting divergent allele-frequency trajectories
between A→C and C→A populations. By comparison, 39,666 SNPs showed a significant main
effect of generation with interaction, and only 1,164 SNPs exhibited a significant main effect of
generation alone. Similarly, only 387 SNPs were significant for just treatment. Overlap among
significant SNP sets was limited (Fig. 5), with only 204 SNPs significant for all three terms. Most
SNPs exhibiting significant interaction effects lacked independent treatment effects, indicating
that genomic differentiation between trajectories arises primarily through opposing evolutionary
change over time rather than static differences between selection regimes. Together, these results
demonstrate that reciprocal selection regimes produce extensive, genome-wide, and largely non-
overlapping allele frequency changes, consistent with anti-parallel evolutionary trajectories
shaped by selection history.

Relaxing Early-Life Selection Recovers Hidden Genetic Variation

Mean genome-wide heterozygosity changed in opposite directions along the two reciprocal
trajectories (Fig. 6). A→C populations, which originate from low-heterozygosity A-type founders,
showed a significant increase in heterozygosity over evolutionary time, whereas C→A
populations, derived from high-heterozygosity C-type founders, exhibited a significant decline.
These trends were consistent across replicate populations (n = 10 per trajectory) and were
already evident at earlier generations. Comparisons between founder and endpoint populations
confirmed highly significant changes in both trajectories (Welch two-sample t-tests, $p < 10^{-11}$).
Together, these results demonstrate divergent genome-wide responses to reciprocal selection,
with relaxation of early-life selection permitting recovery of genetic variation, while renewed early-
life selection reduces heterozygosity.

Many sites in the initial generation of the A→C trajectory appeared fixed under standard
sequencing depth (90×), yet the genome-wide rebound observed following selection reversal
suggested that this apparent fixation might instead reflect alleles present at frequencies too low
to be reliably detected. To test whether these populations harbored hidden variation, we compared
allele frequencies from a single founding population (A-type1, Timepoint 1) sequenced at higher
depth (800× coverage) against sites appearing fixed in the standard-depth dataset. Specifically,
we focused on sites that were classified as significant in the beta-binomial GLM analysis and that
showed either zero or full minor-allele counts under standard sequencing depth (Bonferroni-
corrected GLM, $p < 0.005$). Of the SNPs identified as statistically significant following Bonferroni
correction, 524,873 sites appeared fixed in the A-type1 population. Deep sequencing data were
available for 151,329 of these sites, and subsequent analysis was performed on this subset.

Using the deep-sequencing subset, we examined the distribution of minor allele frequencies at
these sites and compared it to a random subset of equal size drawn from the same deep-
sequencing dataset (Fig. 7). While the random subset exhibited a broad range of allele
frequencies (mean minor allele frequency = 0.174), sites that appeared fixed under standard
coverage were strongly skewed toward very low-frequency alleles, with a mean minor allele
frequency of 0.0063. A permutation test (10,000 iterations) confirmed that this mean was
significantly lower than expected under random sampling ($Z = -450.79$, $p < 1 \times 10^{-4}$). Together,
these results demonstrate that many sites appearing fixed at the start of the A→C trajectory are
in fact segregating at low frequencies, indicating that apparent fixation largely reflects limited
sequencing depth rather than true loss of genetic variation.

Reciprocal Trajectories Exhibit Highly Repeatable Genomic Responses

To evaluate the repeatability of genome-wide evolutionary outcomes at the last collected
timepoint, we performed principal component analysis (PCA) on SNP frequency data from A→C
and C→A populations alongside their corresponding extant founder populations (A and C) (Fig.
8). The first two PCs explained 62.8% (PC1) and 13.2% (PC2) of the total variance and clearly
separated populations by trajectory group.

Replicate populations clustered tightly within each selection treatment, indicating highly
repeatable genomic outcomes under sustained selection. C→A populations, which experienced
182 generations of selection, closely overlapped with the A-founder cluster, whereas A→C
populations, after 65 generations of selection, showed slightly greater dispersion and incomplete
convergence toward C-type founders, consistent with their shorter selection history. Despite this

difference in extent, the two trajectories formed distinct, non-overlapping clusters along both PC1

and PC2, demonstrating strong repeatability of populations sharing the same selection group and

clear differentiation between reciprocal evolutionary treatments.

Linear models testing the effect of trajectory group on the first ten PCs supported these patterns.

Group identity was strongly associated with PC1 and PC2 (Table S22 PC1: estimate ≈ 507,

Bonferroni-adjusted $p < 1.5 \times 10^{-40}$; PC2: estimate ≈ −765, Bonferroni-adjusted $p < 1 \times 10^{-41}$),

whereas no significant effects were detected for PCs 3–10 (adjusted $p > 0.05$). Together, these

results indicate that reciprocal selection produces highly repeatable and selection-specific

genomic states.

To further evaluate the repeatability of genome-wide evolutionary outcomes after selection, we

compared allele frequencies in evolved populations to their corresponding founder populations

using beta-binomial generalized linear mixed models. Specifically, C→A populations were

compared to A-type founders (Fig. 9A), and A→C populations were compared to C-type founders

(Fig. 9B), testing whether replicate populations evolving under the same selection regime

converge on similar genomic states relative to extant founder populations.

The comparison of C→A populations to A-type founders identified only two significant SNPs after

Bonferroni corrections, whereas the comparison of A→C populations to C-type founders yielded

64 significant SNPs spread across the major chromosome arms. This quantitative asymmetry

likely reflects unequal generational exposure to selection: C→A populations experienced

approximately 182 generations of early-reproduction selection, whereas A→C populations

experienced only 65 generations of delayed-reproduction selection, despite both being sampled

at the same chronological time point. Despite this difference, both reciprocal trajectories exhibited

limited genomic differentiation relative to their founder counterparts. Together, these results

indicate that reciprocal trajectories converge on largely parallel and repeatable genomic states

under shared selection regimes, with only modest residual differentiation associated with unequal

generational exposure.

Reciprocal Trajectories Exhibit Largely Parallel Genomic Responses

We quantified genome-wide responses to selection separately for the A→C and C→A trajectories

using quasibinomial generalized linear models that tested for consistent allele frequency change

with generation while accounting for replicate population effects (Fig. S7). At liberal significance

thresholds (FDR < 0.05 without effect size filtering), a very large fraction of the genome showed

evidence of change in both trajectories (843,317 SNPs in A→C and 723,486 SNPs in C→A), with

substantial overlap between trajectories (502,987 shared SNPs; ~60% of A→C and ~70% of
C→A). Tightening the statistical threshold reduced this overlap substantially. At FDR < 0.01, A→C
retained 616,981 significant SNPs whereas C→A was reduced to 241,371, with overlap falling to
133,507 sites (~22% of A→C and ~55% of C→A). Introducing an additional effect size filter
(|estimate| ≥ 0.02) further sharpened this contrast: under FDR < 0.05 with the effect size cutoff,
A→C retained 344,437 SNPs while C→A retained only 65,097, with 30,764 overlapping sites
(~9% of A→C and ~47% of C→A). These global patterns indicate that A→C is associated with a
broad genomic response involving many loci of modest effect, whereas C→A with a more
restricted set of loci exhibiting larger allele frequency shifts. These patterns are consistent with
the substantially greater number of generations of selection in the C→A trajectory relative to the
A→C trajectory. This asymmetry is consistent with classical antagonistic pleiotropy models for
discrete, non-overlapping generations (Rose 1982), where under prolonged directional selection,
such loci may exhibit large frequency shifts but are not expected to sweep universally to fixation.
However, trajectories with shorter selection histories may retain a larger number of loci exhibiting
modest frequency changes.

To assess whether these globally significant SNPs reflect replicate-independent responses to
selection rather than signals driven by individual populations, we implemented a leave-one-out
(LOO) parallelism analysis. For each trajectory, we iteratively identified "target" SNPs using nine
replicate populations under the chosen global threshold and then measured allele frequency
change at these same SNPs in the held-out replicate. Parallelism was summarized as the median
allele frequency shift across target SNPs and compared to an equal-sized set of randomly
sampled, non-significant SNPs ("matched controls") drawn from the same trajectory. This
procedure was repeated for all replicates and summarized genome-wide and by chromosome
arm.

Based on an FDR < 0.05 and ≥ 0.02 allele frequency change threshold, target SNPs showed
consistently larger median allele frequency shifts than matched controls across all replicates in
both trajectories (Figure 10). At the genome-wide level, A→C targets exhibited median shifts of
approximately 0.105 compared to 0.015 for controls, while C→A targets shifted by approximately
−0.098 compared to −0.054 for controls, with these differences strongly supported by two-tailed
t-tests across replicate medians (A→C: $p = 6.99 \times 10^{-8}$; C→A: $p = 2.34 \times 10^{-13}$). This pattern was
consistent across all major chromosome arms (2L, 2R, 3L, 3R, and X), where target SNPs showed
larger-magnitude shifts than matched controls in every case. Although control SNPs also
exhibited non-zero median shifts, reflecting genome-wide allele frequency change associated

with each trajectory, target SNPs showed stronger and more directionally consistent changes

across replicates, indicating parallel responses to selection. Together, the LOO results mirror the

global patterns, with A→C characterized by many small and broadly distributed shifts and C→A

by a smaller subset of loci showing larger and more consistent shifts across replicates. The

direction and magnitude of the shifts also reflects the anti-parallel responses we have observed

between the trajectories in our other analyses.

Discussion

Long term experimental evolution studies have repeatedly shown that adaptation in sexually reproducing populations is dominated by modest, coordinated allele frequency shifts across many loci rather than fixation at single large-effect variants (Burke 2012; Barghi et al. 2019; Barghi and Schlötterer 2020). While this "shifts not sweeps" pattern is now well established, deeper questions emerge regarding the mechanisms responsible for maintaining such extensive variation, especially in cases with sustained and intense selection. In the absence of frequent de novo mutations or environmental fluctuations, classical theories predict that drift and directional selection should steadily erode standing genetic variation as populations converge on a new optimum. In this study we demonstrate that genetic variation persists and remains evolutionarily accessible even after hundreds of generations of intense life-history selection, even from populations associated with major loss of variation (Graves et al. 2017). These results provide strong evidence for the active maintenance of polymorphisms through balancing selection.

Reciprocal selection reveals active maintenance of genetic variation

A central result of this study is the rapid phenotypic reversibility of populations with opposing selection regimes. Adult mortality and developmental timing assays demonstrated clear responses to selection, with populations evolving toward the characteristic phenotypes of long-established A- and C-type populations. Crucially, this reversibility was observed even in A-type populations, which are characterized by substantial genome-wide reductions in genetic variation. These results suggest that alleles contributing to delayed reproduction, extended lifespan, and slower development were actively maintained at low frequencies in the A-type populations in some manner.

Importantly, phenotypic reversibility was mirrored at the genomic level. Principal component analysis reveals a global anti-parallel response with A→C and C→A populations moving in opposing directions toward their respective selection state. When we examined locus-specific responses using beta-binomial generalized linear mixed models, the interaction term accounted for the largest number of significant SNPs suggesting global antiparallel trajectories were not diffuse or idiosyncratic. Rather, they appear to be driven by repeatable and parallel shifts in target sites and linked regions. In total, these results are consistent with widespread balancing selection ensuring populations consistently had access to the same subsets of standing variation when selection was reversed. These results build on those of Graves et al. (2017), showing that in this

system recent selection regime is the primary determinant of a population's genetic state, even
when prior selection regimes involved hundreds of generations of very intense selection.

Relaxation of early-life selection reveals hidden variation

Because A-type populations are characterized by substantial genome-wide depletion of genetic
variation, the wholesale rebound in heterozygosity observed in the A→C populations suggest
these apparent losses are not as complete as they otherwise seem. Instead, evidence again
points to wide-spread balancing selection maintaining polymorphisms at very low frequencies
rather than true fixation. This conclusion is supported by our deep sequencing of an A-type
population. Here we see that many of these seemingly "fixed" sites instead harbor minor alleles
below our usual detection thresholds. Thus, when selection is relaxed, these alleles are able to
respond, producing a genome-wide rebound in heterozygosity.

Although this inference is based on deep sequencing of a single replicate population, the
consistency of the rebound across independent lines suggests that these low-frequency
reservoirs are a general feature of A-type populations, providing a plausible explanation for their
rapid genomic and phenotypic recovery. The critical question, then, is not whether genetic
variation persists, but how it is maintained in each population replicate. The population sizes are
too small for mutations to generate coordinated allele-frequency shifts across thousands of loci
within only a few generations. Accidental migration is another possibility, but we also find this to
be unlikely.

First, the nature of the regimes effectively precludes systematic gene flow between the groups:
A-type flies reproduce before C-type adults even emerge, and at late ages when C-types
reproduce, A-types are competitively inferior. Second, for migration to generate the highly parallel
rebound we observe, identical alleles would have to enter multiple replicate populations at roughly
the same time, persist at low frequency despite being disfavored, and then rise in concert when
selection shifts. Prior simulation work in Phillips et al. (2016) demonstrated that it would require
unrealistically high and persistent migration rates to produce the levels of similarity observed
between replicate populations in this system. Such rates are inconsistent with independently
maintained laboratory stocks where migration is accidental by definition.

Highly repeatable and parallel genomic responses argue against historical contingency

A striking feature of the reciprocal trajectories is the tight clustering of replicate populations at our
final collection points. Despite originating from similar yet distinct founders and evolving along

opposite trajectories, populations subjected to the same selection regime converge on highly
reproducible genomic states within their respective regimes. The limited number of loci that
remain significantly differentiated between reciprocally selected populations and their
corresponding founders further underscores the repeatability of these evolutionary outcomes.
Importantly, the small asymmetry observed between the A→C and C→A comparisons is readily
explained by differences in the number of generations experienced under selection, rather than
by intrinsic limits on reversibility. These results argue strongly against a dominant role for historical
contingency or idiosyncratic paths in shaping long-term genomic outcomes. Instead, they indicate
that selection repeatedly draws on the same reservoirs of standing variation and drives
populations toward predictable genomic outcomes.

Next, the largely parallel genomic responses within each reciprocal trajectory clarify how selection
acts on standing variation in this system. Even under stringent significance thresholds, a
substantial fraction of the genome shifts in both A→C and C→A trajectories, highlighting the
pervasive impact of altered life-history selection. Leave-one-out analyses further demonstrate that
this pattern is underlain by a high degree of parallelism in the genomic response to selection
within a group.

Current trajectories support robust inference despite ongoing adaptation

Collectively, our data suggest that the A→C populations are still adapting, and our conclusions
therefore reflect the current direction and consistency of the response rather than a completed
evolutionary endpoint. Nevertheless, the concordance of phenotypic reversibility, genome-wide
antiparallel allele-frequency shifts, rebounds in heterozygosity, and agreement across multiple
genomic analyses indicates that the central features of the response are already well established.
While additional timepoints may refine the magnitude of these changes, they are unlikely to
qualitatively alter the interpretation supported by the patterns already observed.

Antagonistic pleiotropy as a plausible mechanism

Taken together, these results are consistent with the antagonistic pleiotropy models developed
for discrete-generation populations in Rose (1982). Importantly, our interpretations do not require
that individual fitness-components exhibit classical overdominance or frequency-dependent
selection. Rather, antagonistic pleiotropy distributed across many fitness-components can
collectively generate balancing dynamics at the level of net fitness. In such a system, fixation is
prevented because homozygous genotypes have reduced overall fitness compared to
heterozygous genotypes. This population-level resolution of trade-offs naturally leads to the

maintenance of extensive genetic variation, even under constant environmental conditions.

Because the Rose laboratory's A and C regimes impose selection directly on reproductive timing,

they inherently act on age-structured fitness, a core context in which antagonistic pleiotropy is

predicted to generate life-history trade-offs (Rose 1985).

Conclusion

Broadly, these findings contribute to a growing body of evidence that adaptive evolution in sexually reproducing populations frequently proceeds through coordinated, genome-wide shifts in allele frequencies rather than classic hard sweeps (Burke 2012; Barghi et al. 2019; Barghi and Schlötterer 2020). In our reciprocal selection framework, the largely antiparallel responses across replicate populations and chromosome arms, together with the capacity for directional reversal under altered age-specific selection, indicate that substantial genetic variation is maintained over long evolutionary timescales. Rather than being exhausted by prolonged selection, this standing variation remains available to fuel rapid and repeatable responses when the selective regime changes. Such patterns serve as strong signatures of balancing processes operating across many loci, maintaining polymorphism through time and enabling predictable yet reversible genomic trajectories. Viewed in this light, our results contribute to the growing body of literature on the importance of considering balancing selection in evolutionary genetics, not as a marginal phenomenon, but as a major driver of observed patterns of genetic variation (Kawecki et al. 2021, Zwoinska et al. 2026).

Materials and Methods

Experimental Populations

All *Drosophila melanogaster* populations used in this study derive and branch from the outbred population established by P. T. Ives in 1970 and reared on standard banana agar media. From this founding population multiple experimental evolution lines have been derived, typically with five independently replicated populations and maintained and subjected to distinct selection regimes. Of these treatments, we have cultured ten early-reproducing A-types and ten late-reproducing C-types, achieving hundreds of generations of sustained selection (Rose et al. 1992; Chippindale et al. 1997; Burke et al. 2016), demonstrating strong convergence across life-history characteristics and genomics (Graves et al. 2017). Given this strong convergence within each regime, we will refer to these two sets of 10 populations as the founder A-types and the founder C-types respectively.

Building upon this work, we derived 20 new populations from the progenitor A-types and C-types. This was accomplished by imposing an A-type selection on 10 populations derived from the progenitor C-type populations to create the C→A-types. In tandem, C-type selection was imposed on 10 lines derived from the progenitor A-type populations to create the A→C-types.

C→A-type populations were created by collecting eggs from the 10 founder C-types and only permitting the first ~20% developing flies to contribute to the next generation. This was accomplished by rearing eggs from each generation in carousels, which feature food caps housed in a dome to contain the earliest developing flies within until enough individual flies have eclosed to create a new outbred population. The earliest derived flies contained in the carousel were then gassed with CO_2 and pooled into Plexiglass cages to lay eggs for the next generation. Multiple carousels were utilized per population to yield large cohort sizes to prevent inbreeding depression from the intense selection regime. With each vial containing 80+ eggs, 80 vials each in each carousel, and ~3 carousels per population, ~19,200 total eggs were obtained each early generation, with approximately ~3840 young adults constituting the top 20% of early developers permitted to create the next generation. With each successive generation of directional selection, average development time decreased, reducing the need for additional carousels and thereby minimizing the risk of large population bottlenecks and cohorts necessary for each population as the experiment continued. Once the C→A-type populations were reliably reproducing on Day 10 without complications, they were maintained like their progenitor A-type counterparts with only vials and cages, and no carousels.

A→C-type populations were created by collecting eggs from 10 progenitor A-types populations

allowing only those females who were still alive and fecund by day 26 to contribute to the next

generation. Because A-type populations have substantially shorter adult lifespans than C-type

populations, additional cohort cages were required to ensure that a sufficient number of flies

survived to day 26 to propagate the next generation. This involved collecting ~7,500 eggs for each

population to be transferred on day 14 into 5 separate cages containing ~1,500 individuals.

Standard banana media was fed to the cohorts every other day until Day 26 when cages were

condensed to ensure enough living and breeding individuals were pooled to have at least ~1,500

individuals to produce the next generation for a given population. Intense forward selection

continued with multiple cohorts collected from separate cages, and then pooled within a

population cage to elevate census sizes until populations can reliably be maintained with a single

cage.

Data Collection

At several intervals, phenotypic assays and genomic samples were collected to track the two

opposing trajectories as their constituent populations adapted to their new selection regimes. By

measuring these newly derived populations at multi-generation intervals, we were able to map

the initial trajectories of phenotypic and genotypic change in response to selection. Although

phenotypic assays and genomic sampling were conducted at the same chronological time points,

A→C and C→A trajectories accumulated different numbers of generations due to differences in

generation time and are therefore treated as distinct experiments.

Adult Mortality

Adult mortality assays followed the protocols described in Burke et al. (2016), with minor

modifications. Progenitor stock populations were sampled on a 14-day culture cycle for two

generations, staggered by one day to align treatments on a shared calendar and minimize

environmental variation. Each population was represented by two cohorts (e.g., A-type1α and A-

type1β) of approximately 1,500 flies housed in Plexiglas cages and supplied daily with fresh

banana–molasses medium (Phillips et al. 2018) supplemented with yeast to induce oviposition

after transfer on day 14 from egg. At the same time each day, all dead flies were removed, sexed,

and recorded for each cohort until no individuals remained. Mortality data were pooled across the
two cohorts to obtain daily death counts for each population.

The Burke et al. (2016) protocol was modified to eliminate the use of carbon dioxide for
condensing cohorts when census densities were low. Instead, flies remained in their original
cages for the duration of the assay, with internal cleaning performed as needed to remove waste
accumulation from the cage walls. Adult mortality assays were conducted on founder populations
and on the selected populations at multiple points along the adaptive trajectory, corresponding to
approximately 12 and 51 generations for A→C populations and approximately 33 and 141
generations for C→A populations.

Larval Development Analysis

Following two generations of standardized rearing, ancestral founder populations (A-type
and C-type) and trajectory populations from the C→A and A→C treatments were assayed for
pupation timing. For each population, eggs were collected on nutrient-free agar plates, and 60
first-instar larvae were individually transferred to food vials, with three vials established per
population. Vials were maintained in a sealed incubator at 25 °C under constant light conditions.
Larval development was monitored every four hours for 14 days, and the time of pupation was
recorded for each individual upon detection of newly formed pupal casings. To minimize temporal
bias and blocking effects among treatments, one replicate population was initiated per day;
however, once established, all replicates were monitored at identical time intervals throughout the
assay.

Pupation data were analyzed using linear mixed-effects models to evaluate the effects of
selection regime on developmental timing. For each vial, the proportion of individuals pupated
was modeled as a function of selection regime, time, and their interaction, with vial nested within
replicate as a random effect to account for non-independence among larvae sharing the same
vial. Replicate identity corresponded to the day of assay initiation and was included to account for
blocking effects. For each vial we have assayed the percentage of developed flies f_{hijk}, from
selection regime *(h=1,2,..,8)*, time *(i=4,8,…,T)*, replicate *(j=1,2,…5)*, and vial *(k=1,2,3)*. The model
can be summarized as:

539
$$f_{hijk} = \mu + \alpha_h \delta_h + \beta_i \delta_i + \gamma_{hi} \delta_h \delta_i + \theta_{jk} + \varepsilon_{hijk}$$

540   where the Kronecker deltas are defined as:

$$\delta_h = \begin{cases} 0 & if \ h \ = 1 \\ 1 & otherwise \end{cases} \quad \delta_i = \begin{cases} 0 & if \ i \ = 4 \\ 1 & otherwise \end{cases}$$

The random component θ_{jk} refers to the random variation of vials nested within replicate. In short,
Percent Developed = μ+Selection Regime*Time+Replicate/Vial+ε. Replicate refers to reps 1, 2,
3, 4, or 5 and corresponds to day of experiment start because of blocking effects. Pupation assays
were conducted at multiple points along the adaptive trajectory. The first pupation rate assays
corresponded to approximately 15 generations of early-reproduction selection for C→A
populations and approximately 6 generations of delayed-reproduction selection for A→C
populations. The final pupation rate assays corresponded to approximately 150 generations of
early-reproduction selection for C→A populations and approximately 50 generations of delayed-
reproduction selection for A→C populations.

Genomic Sampling

Samples for genomic analysis were collected following the protocols described in Graves et al.
(2017). For each population, approximately 1,500 adults per population were maintained in
separate Plexiglas cohort-cages with banana–molasses medium. Prior to sampling, populations
underwent two lead-in generations on a standardized 14-day culture cycle, with collections
staggered by one day among cages to reduce gene-by-environment effects. Flies were
anesthetized with CO_2 before sampling, and each sample consisted of a pool of ~200 randomly
collected females per population. The ten A-type and ten C-type founder populations were
sequenced at time points 1 and 4, and the ten A→C and ten C→A reciprocal populations were
sampled at time points 2, 3, and 4.

DNA Extraction & Sequencing

Pooled samples of ~200 female flies were immediately flash-frozen in liquid nitrogen and stored
at −80°C prior to genomic extraction. Samples were homogenized in liquid nitrogen prior to
extraction using Qiagen Puregene kits, and DNA concentration was assessed using a NanoDrop
spectrophotometer. Extracted DNA pools were submitted to the UC Irvine Genomics Research
and Technology Hub (GRTH) for gDNA library construction and sequencing. Samples were
mechanically sheared to a target size of ~150 bp, and an AMPure XP bead cleanup (1.8x) was

performed on a subset of samples exhibiting smaller fragment profiles indicative of partial
degradation; all samples were verified by Bioanalyzer prior to library construction. Each replicate
population received a unique dual-index (UDI) barcode (IDT UDI indices), and libraries were
sequenced on an Illumina NovaSeq X Plus using 150 bp paired-end chemistry (PE150). To avoid
index collision, samples from different collection years were multiplexed on separate lanes.
Samples from were initially sequenced at a target depth of ~30 million read pairs (~50x average
genome coverage), re-sequenced again at a target depth of ~60 million read pairs (~100x average
genome coverage). One population (A-type1, Timepoint 1) was additionally subjected to deep
sequencing at a target depth of 500 million read pairs; library concentration was re-verified by
Qubit prior to this run.

SNP Data Processing

Raw sequencing data were received from the sequencing center as ~150 bp paired-end FASTQ
files. Each sample was sequenced twice across two lanes. Reads were mapped to the *Drosophila*
melanogaster reference genome (dm6, including the ISO1 mitochondrial genome) using Novalign
v4.04.01 (Novocraft Technologies 2025). Alignment was performed in paired-end mode (-r
RANDOM, -I PE 250, 50) with default settings. A homopolymer filter was used to exclude low-
complexity reads, and reads failing to meet quality thresholds were removed prior to alignment.
The "Novosort" function was then use to sort the resulting bam files and identify PCR duplicates.
The two BAM files corresponding to a given population were merged using Bamtools (Barnett et
al. 2011). Bam files for individual samples were then combined in the mpileup format using
SAMtools (Li and Durbin 2009). *Popoolation2* (Kofler et al. 2011) was then used to convert this
mpileup to a "sync" file for ease of processing. Lastly, a python script was used to convert the
sync file into SNP table with minor allele counts and total coverage for all polymorphic sites for
each population in the dataset. Only biallelic sites were considered and the minor allele was
defined as the less common nucleotide looking across all samples. We only considered sites that
had a minimum 20X coverage in each sample, and polymorphic sites were defined as those with
a minimum minor allele frequency of 0.02 across all samples. This resulted in a data set with
1,389,963 SNPs. Average SNP coverage per sample ranged from 62X to 113X, with an average
mean coverage of 89 looking across samples.

Principal Component Analysis

To summarize genome-wide allele frequency variation, principal component analysis (PCA) was conducted on SNP frequency matrices using the prcomp function in R (R Core Team 2024). Data were scaled to unit variance but not centered. SNPs with zero minor allele counts, a pseudocount of 1 was added to prevent division by zero in frequency calculations. To examine the evolutionary trajectories of populations over time, each PC (PC1–PC10) was modeled as a function of trajectory and generation using linear mixed-effects models:

$$PC \sim trajectory \times generation + (1 \mid population)$$

where *trajectory* represents the evolutionary direction (A→C or C→A), generation is a continuous variable representing generations since the start of selection, and population is a random intercept accounting for replicate-specific effects. Models were fit using *lmer* from the lme4 package, and significance of fixed effects was assessed using the lmerTest package (Kuznetsova et al. 2017). Bonferroni-adjusted p-values were calculated across PCs and model terms with α = 0.005.

Modeling Genome-Wide Allele-Frequency Dynamics with Beta-Binomial GLMMs

We used a beta-binomial generalized linear mixed model to account for overdispersion in pooled-sequencing allele-count data, which commonly arises from sampling variance, linkage, and unmodeled heterogeneity among loci and replicates (Wiberg et al. 2017). To identify SNPs exhibiting reciprocal evolutionary responses between the A→C and C→A trajectories, we modeled allele counts at each locus with reciprocal responses captured by the trajectory × generation interaction term. Let Y_{ijk} denote the minor allele count at SNP i in population replicate j at generation k, with n_{ijk} total reads. Then:

$$Y_{ijk} \sim BetaBinomial(n_{ijk}, p_{ijk}, \phi),$$

where p_{ijk} is the underlying allele frequency, and ϕ is the overdispersion parameter. The linear predictor for the logit of the allele frequency is:

$$logit(p_{ijk}) = \beta_0 + \beta_1 T_j + \beta_2 G_k + \beta_3 (T_j \cdot G_k) + u_j$$

β_0 is the intercept (baseline allele frequency), T_j is the treatment / trajectory indicator (0 = A→C,

1 = C→A), G_k is the continuous generation variable, β_1, β_2, β_3 are fixed-effect coefficient for

treatment, generation, and their interaction $u_j \sim N(0, \sigma_u^2)$ = random effect for population replicate

j. The interaction term β_3 captures trajectory-specific allele-frequency shifts over generations,

while the random intercept u_j accounts for replicate-level variation. To correct for multiple

comparisons across the genome, p-values for each model term were adjusted using a Bonferroni

correction and SNPs were considered statistically significant at a genome-wide threshold of α =

0.05.

Genome-Wide Heterozygosity Analysis

To assess genome-wide heterozygosity dynamics across the A→C and C→A

experimental evolution trajectories, SNP allele counts were obtained from the trajectory dataset.

Sample identifiers were standardized to reflect replicate populations and treatment direction

(A→C or C→A), and total coverage columns were converted to major allele counts by subtracting

minor allele counts from total coverage.

$$H = 2 \cdot f \cdot (1 - f)$$

where f is the minor allele frequency.

Changes in mean heterozygosity over time were visualized using boxplots stratified by treatment.

Statistical significance between initial and final generations for each trajectory was evaluated

using two-sided t-tests, and p-values were converted to conventional significance labels (ns, *, **,

***) for visualization.

To further investigate the apparent loss of variation observed in the A→C trajectory, we examined

whether sites that appeared fixed at generation 0 truly lacked standing variation. This analysis

was motivated by the observation that many SNPs in the A→C populations appeared fixed based

on standard-coverage sequencing, yet these same populations exhibited rapid phenotypic and

genomic rebound when selection was reversed, an outcome inconsistent with widespread true

fixation.

To characterize highly differentiated sites in the A→C trajectory, SNP data were obtained from the

trajectory SNP table and corresponding deep sequencing counts, from the generation 0 A-type1

population. Generalized linear models (GLMs) with a beta-binomial error structure had previously

been applied to estimate allele-frequency changes across the trajectory; resulting p-values were

Bonferroni-corrected, and SNPs with corrected p-values < 0.005 were retained.

Significant SNPs were then filtered to identify pseudo-fixed sites in A-type1, defined as positions

where the minor allele appeared absent or fixed based on standard coverage (minor allele count

= 0 or equal to total coverage). These sites were merged with deep sequencing data to compute

minor allele frequencies (MAFs), allowing detection of low-frequency alleles that would otherwise

be missed.

The distribution of MAFs at pseudo-fixed sites was visualized using histograms and compared

against a randomly sampled set of genome-wide SNPs of equal size to provide a null background.

To formally test whether pseudo-fixed sites exhibited reduced but nonzero variation, we performed

a permutation test in which the mean MAF of pseudo-fixed sites was compared to a null

distribution generated by repeatedly sampling random SNP sets from the deep sequencing data

(n = 10,000 permutations). Empirical p-values were calculated as the proportion of permuted

means less than or equal to the observed mean, and standardized z-scores were used to quantify

deviation from the null expectation.

Generalized Linear Mixed Model Analysis of 'newly derived' versus founder populations

To test the repeatability of genomic evolution under reversed selection, we asked whether

populations selected for a given treatment (A or C) converged on the same allele-frequency states

as the original founder populations adapted to that regime. Specifically, we tested whether C→A

populations resemble founder A populations, and whether A→C populations resemble founder C

populations. Under repeatable evolution, populations that independently experience the same

selection pressure are expected to arrive at similar allele-frequency distributions, despite distinct

evolutionary histories.

All analyses were conducted using SNP count data from the trajectory dataset. For each

comparison, we retained SNPs that were polymorphic among populations, excluding sites that

were monomorphic across all replicates or fixed in all individuals. Chromosome 4 was excluded

due to its atypical recombination properties and high heterochromatin content.

Repeatability was assessed on a per-SNP basis using generalized linear mixed models that test

whether newly derived populations differ from their corresponding founders. For each SNP, minor-

allele counts were modeled with a quasibinomial error distribution, with total sequencing depth

used as the denominator. Models included population type (newly derived vs. founder) as a fixed

effect and replicate population as a random intercept to account for shared ancestry and non-

independence among samples within regimes.

For each SNP, evidence for non-repeatable evolution was evaluated by comparing the full model

to a null model lacking the population-type effect using likelihood ratio tests. SNPs showing no

significant differentiation between newly derived and founder populations were interpreted as

consistent with repeatable convergence on a shared allele-frequency state, whereas significant

differentiation indicates historical contingency or incomplete reversibility at that locus.

The quasibinomial framework was used to accommodate overdispersion inherent to Pool-seq

allele-count data, following Wiberg et al. 2017. Multiple testing was controlled using a Bonferroni

correction, and SNPs with corrected p-values below 0.05 significance threshold were considered

significantly differentiated between newly derived and founder populations. At a genome-wide

level, the proportion and genomic distribution of differentiated versus non-differentiated SNPs

provide a quantitative measure of repeatability, capturing the extent to which reversed populations

recapitulate the genomic architecture of adaptation to a given selective regime.

706   Leave-One-Out (LOO) parallelism analysis

To assess and compare parallel evolutionary change between the A→C and C→A selection

trajectories, we conducted (i) genome-wide trajectory-specific scans to summarize baseline

patterns of allele-frequency change and (ii) a leave-one-out (LOO) framework to test replicate-

independent parallelism based on methods used by Bitter et al. (2024). All analyses were

performed separately for the A→C and C→A trajectories.

Baseline genome-wide scans were used to describe trajectory-level patterns and to visualize the

distribution of signals across the genome. For each SNP within each trajectory, minor and major

allele counts were modeled with a quasibinomial GLM:

$$(Minor\ count, Major\ Count) \sim generation + replicate$$

where generation was treated as a numeric predictor and *replicate* accounted for baseline

differences among replicate populations. For each SNP, the estimate and p-value for the

generation term were extracted, and p-values were adjusted using the Benjamini–Hochberg false

discovery rate (FDR) procedure. These baseline scans were used to summarize broad

differences between trajectories, but were not used for the replicate-independent parallelism test

described below.

Parallelism was assessed using a leave-one-out (LOO) framework in which each replicate

population was treated as an independent test case. For each iteration, one replicate was

excluded and the remaining replicates were used as a training set to re-fit per-SNP quasibinomial

models of the same form as above. Candidate SNPs ("targets") were defined within each training

set using a false discovery rate (FDR) threshold of 0.05 and a minimum effect-size criterion of

$|\Delta p| \geq 0.02$. Here, Δp represents the net change in minor allele frequency between the first and

last sampled generations, with allele frequencies calculated by pooling allele counts across all

training replicates within each generation.

Parallelism was evaluated by quantifying allele-frequency change in the held-out replicate alone.

For each LOO iteration, Δp was recalculated for the held-out replicate as the difference in minor

allele frequency between the final and initial sampled generations. Parallelism was assessed by

comparing Δp at target SNPs to control SNPs, which were randomly sampled from sites that did

not meet the target criteria in the corresponding training set and matched in number to targets.

Control SNPs capture background genome-wide allele-frequency change due to drift, linkage,

and hitchhiking. Parallel evolutionary response within each replicate was summarized as the

median Δp across target SNPs and across control SNPs, calculated both genome-wide and for

individual chromosome arms. Replicate-level median Δp values for targets and controls were

compared using two-sample, two-tailed t-tests, treating replicate populations as independent

units.

Data Availability

Raw genomic and phenotypic data are available through Dryad (DOI: 10.5061/dryad.tdz08kqdc) and scripts used to process and analyze data are available through Github (https://github.com/krarnold/A-C_Genomic_Trajectory_Project). The DNA sequence data supporting the conclusions of this article are available in the NCBI SRA repository (PRJNA1426013).

Acknowledgments

This work was supported by the National Institutes of Health Maximizing Investigators' Research Award to MAP (NIH R35GM155286). We thank the UCI Genomics High-Throughput Facility for their assistance with library preparation, sequencing, and technical support that made this work possible. We thank the developers of Novoalign for providing the alignment software used in processing and mapping our sequencing reads. Most importantly, we are grateful to the dozens of undergraduate researchers who assisted with the experiments, without whose help this work could not have been done, especially S. Gagucas, G. Medrano, R. Odina, J. Sue, D. Youn, N. Ly, N. Nguyen, T. Nguyen, E. Choi, K. Hoang-Yanis, C. Banoub, M. Ishizuka, A. Attalla, C. Wang, E. Brown, B. Khan, J. Hana, A. Vo, K. Co, J. Anup, J. Mai, J. Lee, K. Bryan, R. Chhetri, and B. Ung.

Author Contributions

M.A.P., M.R.R., and L.D.M. conceptualized and oversaw the project. M.A.P., M.R.R., L.D.M., and K.R.A. designed the experiment. A.P., V.V.C., R.D.R., K.C., C.O.C., and M.Q. collected all fly samples for genomic analyses and carried out mortality assays and larval development experiments under the supervision of K.R.A. K.R.A. extracted samples for genomic sequencing. Z.S.G. and L.D.M analyzed phenotypic data. K.R.A., Z.S.G., and M.A.P. analyzed the genomic data. K.R.A. and M.A.P. wrote the manuscript.

References

Barghi, Neda, Joachim Hermisson, and Christian Schlötterer. 2020. "Polygenic Adaptation: A
Unifying Framework to Understand Positive Selection." *Nature Reviews Genetics* 21 (12):
769–81. https://doi.org/10.1038/s41576-020-0250-z.

Barghi, Neda, and Christian Schlötterer. 2020. "Distinct Patterns of Selective Sweep and
Polygenic Adaptation in Evolve and Resequence Studies." *Genome Biology and Evolution*
12 (6): 890–904. https://doi.org/10.1093/gbe/evaa073.

Barghi, Neda, Raymond Tobler, Viola Nolte, et al. 2019. "Genetic Redundancy Fuels Polygenic
Adaptation in Drosophila." *PLOS Biology* 17 (2): e3000128.
https://doi.org/10.1371/journal.pbio.3000128.

Barnett, Derek W., Erik K. Garrison, Aaron R. Quinlan, Michael P. Strömberg, and Gabor T. Marth.
2011. "BamTools: A C++ API and Toolkit for Analyzing and Managing BAM Files."
Bioinformatics 27 (12): 1691–92. https://doi.org/10.1093/bioinformatics/btr174.

Bitter, M. C., S. Berardi, H. Oken, et al. 2024. "Continuously Fluctuating Selection Reveals Fine
Granularity of Adaptation." *Nature* 634 (8033): 389–96. https://doi.org/10.1038/s41586-
024-07834-x.

Burke, Molly K. 2012. "How Does Adaptation Sweep through the Genome? Insights from Long-
Term Selection Experiments." *Proceedings of the Royal Society B: Biological Sciences*
279 (1749): 5029–38. https://doi.org/10.1098/rspb.2012.0799.

Burke, Molly K., Thomas T. Barter, Larry G. Cabral, et al. 2016. "Rapid Divergence and
Convergence of Life-history in Experimentally Evolved *Drosophila Melanogaster*."
Evolution 70 (9): 2085–98. https://doi.org/10.1111/evo.13006.

Burke, Molly K., Joseph P. Dunham, Parvin Shahrestani, Kevin R. Thornton, Michael R. Rose,
and Anthony D. Long. 2010. "Genome-Wide Analysis of a Long-Term Evolution
Experiment with Drosophila." *Nature* 467 (7315): 587–90.
https://doi.org/10.1038/nature09352.

Burke, Molly K., Gianni Liti, and Anthony D. Long. 2014. "Standing Genetic Variation Drives
Repeatable Experimental Evolution in Outcrossing Populations of Saccharomyces

Cerevisiae." *Molecular Biology and Evolution* 31 (12): 3228–39.
https://doi.org/10.1093/molbev/msu256.

Chevin, Luis-Miguel, and Frédéric Hospital. 2008. "Selective Sweep at a Quantitative Trait Locus
in the Presence of Background Genetic Variation." *Genetics* 180 (3): 1645–60.
https://doi.org/10.1534/genetics.108.093351.

Chippindale, A. K., A. M. Leroi, H. Saing, D. J. Borash, and M. R. Rose. 1997. "Phenotypic
Plasticity and Selection in Drosophila Life History Evolution. 2. Diet, Mates and the Cost
of Reproduction." *Journal of Evolutionary Biology* 10 (3): 269–93.
https://doi.org/10.1046/j.1420-9101.1997.10030269.x.

Graves, J.L., K.L. Hertweck, M.A. Phillips, et al. 2017. "Genomics of Parallel Experimental
Evolution in *Drosophila*." *Molecular Biology and Evolution*, January 12, msw282.
https://doi.org/10.1093/molbev/msw282.

Hayward, Laura Katharine, and Guy Sella. 2022. "Polygenic Adaptation after a Sudden Change
in Environment." *eLife* 11 (September): e66697. https://doi.org/10.7554/eLife.66697.

Hubert, David L, Kenneth R Arnold, Zachary S Greenspan, et al. 2025. "Selection for Early
Reproduction Leads to Accelerated Aging and Extensive Metabolic Remodeling in
Drosophila Melanogaster." *Genome Biology and Evolution* 17 (5): evaf082.
https://doi.org/10.1093/gbe/evaf082.

Ives, Philip T. 1970. "FURTHER GENETIC STUDIES OF THE SOUTH AMHERST POPULATION
OF DROSOPHILA MELANOGASTER." *Evolution; International Journal of Organic
Evolution* 24 (3): 507–18. https://doi.org/10.1111/j.1558-5646.1970.tb01785.x.

Kawecki, Tadeusz J., Berra Erkosar, Cindy Dupuis, Brian Hollis, R. Craig Stillwell, and Martin
Kapun. 2021. "The Genomic Architecture of Adaptation to Larval Malnutrition Points to a
Trade-off with Adult Starvation Resistance in *Drosophila*." *Molecular Biology and Evolution*
38 (7): 2732–49. https://doi.org/10.1093/molbev/msab061.

Kezos, James N., Thomas T. Barter, Mark A. Phillips, et al. 2023. "Building Bridges from Genome
to Physiology Using Machine Learning and *Drosophila* Experimental Evolution."
Physiological and Biochemical Zoology 96 (3): 192–205. https://doi.org/10.1086/724827.

Kofler, Robert, Ram Vinay Pandey, and Christian Schlötterer. 2011. "PoPoolation2: Identifying
Differentiation between Populations Using Sequencing of Pooled DNA Samples (Pool-
Seq)." *Bioinformatics* 27 (24): 3435–36. https://doi.org/10.1093/bioinformatics/btr589.

Kuznetsova, Alexandra, Per B. Brockhoff, and Rune H. B. Christensen. 2017. "**lmerTest** Package:
Tests in Linear Mixed Effects Models." *Journal of Statistical Software* 82 (13).
https://doi.org/10.18637/jss.v082.i13.

Lewontin, Richard C. 1974. *The Genetic Basis of Evolutionary Change*. Columbia Biological
Series 25. Columbia Univ. Pr.

Li, Heng, and Richard Durbin. 2009. "Fast and Accurate Short Read Alignment with Burrows–
Wheeler Transform." *Bioinformatics* 25 (14): 1754–60.
https://doi.org/10.1093/bioinformatics/btp324.

Linder, Robert A, Behzad Zabanavar, Arundhati Majumder, et al. 2022. "Adaptation in Outbred
Sexual Yeast Is Repeatable, Polygenic and Favors Rare Haplotypes." *Molecular Biology*
and Evolution 39 (12): msac248. https://doi.org/10.1093/molbev/msac248.

Long, Anthony, Gianni Liti, Andrej Luptak, and Olivier Tenaillon. 2015. "Elucidating the Molecular
Architecture of Adaptation via Evolve and Resequence Experiments." *Nature Reviews*
Genetics 16 (10): 567–82. https://doi.org/10.1038/nrg3937.

Novocraft Technologies. 2025. *Novoalign*. V. 4.04.01. Released.
http://www.novocraft.com/products/novoalign/.

Phillips, Mark A., Ian C. Kutch, Anthony D. Long, and Molly K. Burke. 2020. "Increased Time
Sampling in an Evolve-and-resequence Experiment with Outcrossing *Saccharomyces*
Cerevisiae Reveals Multiple Paths of Adaptive Change." *Molecular Ecology* 29 (24): 4898–
912. https://doi.org/10.1111/mec.15687.

Phillips, Mark A., Anthony D. Long, Zachary S. Greenspan, et al. 2016. "Genome-Wide Analysis
of Long-Term Evolutionary Domestication in Drosophila Melanogaster." *Scientific Reports*
6 (1): 39281. https://doi.org/10.1038/srep39281.

R Core Team. 2024. *R: A Language and Environment for Statistical Computing*. R Foundation for
Statistical Computing. https://www.R-project.org/.

Rose, Michael R. 1982. "Antagonistic Pleiotropy, Dominance, and Genetic Variation." *Heredity* 48
(1): 63–78. https://doi.org/10.1038/hdy.1982.7.

Rose, Michael R. 1985. "Life History Evolution with Antagonistic Pleiotropy and Overlapping
Generations." *Theoretical Population Biology* 28 (3): 342–58.
https://doi.org/10.1016/0040-5809(85)90034-6.

Rose, Michael R., Hardip B. Passananti, and Margarida Matos, eds. 2004. *Methuselah Flies*.
WORLD SCIENTIFIC. https://doi.org/10.1142/5457.

Rose, Michael R., Linh N. Vu, Sung U. Park, and Joseph L. Graves. 1992. "Selection on Stress
Resistance Increases Longevity in Drosophila Melanogaster." *Experimental Gerontology*
27 (2): 241–50. https://doi.org/10.1016/0531-5565(92)90048-5.

Rudman, Seth M., Sharon I. Greenblum, Subhash Rajpurohit, et al. 2022. "Direct Observation of
Adaptive Tracking on Ecological Time Scales in *Drosophila*." *Science* 375 (6586):
eabj7484. https://doi.org/10.1126/science.abj7484.

Schlötterer, C, R Kofler, E Versace, R Tobler, and S U Franssen. 2015. "Combining Experimental
Evolution with Next-Generation Sequencing: A Powerful Tool to Study Adaptation from
Standing Genetic Variation." *Heredity* 114 (5): 431–40.
https://doi.org/10.1038/hdy.2014.86.

Tobler, Ray, Susanne U. Franssen, Robert Kofler, et al. 2014. "Massive Habitat-Specific Genomic
Response in D. Melanogaster Populations during Experimental Evolution in Hot and Cold
Environments." *Molecular Biology and Evolution* 31 (2): 364–75.
https://doi.org/10.1093/molbev/mst205.

Wiberg, R. Axel W., Oscar E. Gaggiotti, Michael B. Morrissey, and Michael G. Ritchie. 2017.
"Identifying Consistent Allele Frequency Differences in Studies of Stratified Populations."
Methods in Ecology and Evolution 8 (12): 1899–909. https://doi.org/10.1111/2041-
210X.12810.

Zwoinska, Martyna K., R. Axel W. Wiberg, Milena Trabert, Philipp Kaufmann, and Elina Immonen.
2026. "From Constraint to Opportunity: Relaxing Sexual Antagonism Reveals Adaptive
Potential Maintained by Balancing Selection." Preprint, Evolutionary Biology, February 6.
https://doi.org/10.64898/2026.02.04.703537.

Figures

Figure 1. Experimental design of reciprocal A→C and C→A selection trajectories.
Schematic illustrating the derivation of 20 trajectory populations from 20 long-established founder
populations under antiparallel life-history selection. The x-axis represents generational time, and
the y-axis represents shifts in the reproductive window imposed by directional selection. Ten
founder C-type populations were subjected to A-type selection, favoring progressively earlier
reproduction, generating C→A trajectories that ultimately reached the 10-day generation cycle
characteristic of A-type populations. Conversely, ten founder A-type populations were subjected
to C-type selection, favoring delayed reproduction and increased longevity, generating A→C
trajectories that ultimately reached the 28-day generation cycle characteristic of C-type
populations. Phenotypic assays and genomic sampling were conducted at defined points along
each trajectory to quantify evolutionary responses across generations.

Figure 2. Waves of Convergence selection. Differences in mean age-specific adult mortality plotted as a function of age from egg. (A) C→A trajectory: differences between C→A populations and A-type founder populations after ~33 (magenta) and ~143 (red) generations of early-reproduction selection, shown relative to the baseline difference between C- and A-type founders (purple). (B) A→C trajectory: differences between A→C populations and C-type founder populations after ~12 (light blue) and ~51 (dark blue) generations of delayed-reproduction selection, shown relative to the baseline difference between A- and C-type founders (purple). Points indicate mean mortality differences at each age; lines show smoothed trends.

905 **Figure 3. Antiparallel genomic trajectories under reciprocal selection.** Lines indicate allele-

906 frequency trajectories of replicate populations for A→C (blue) and C→A (red) across PC1 and

907 PC2. Facet labels indicate the percentage of variance explained by each PC. Each line

908 represents a replicate population, with points corresponding to successive generations (x-axis

909 labels show generation numbers for each trajectory).

**Figure 4. Genome-wide distribution of SNPs with Bonferroni-significant selection ×
generation interactions**. The Manhattan plot shows $-\log_{10}$-transformed Bonferroni-adjusted p-
values for the interaction term across all major chromosome arms (2L, 2R, 3L, 3R, X). SNP
positions are shown in megabases (Mb) along the x-axis, and chromosomes are colored
individually for clarity. The dashed horizontal line indicates the $\alpha = 0.05$ significance threshold
applied to Bonferroni-adjusted p-values. Significant SNPs are broadly distributed across the
genome, with no single chromosome dominating the signal, consistent with a highly polygenic
response and anti-parallel evolutionary trajectories across selection regimes.

Figure 5. Genome-wide SNP overlap among Bonferroni-significant effects. The UpSet plot displays the number of SNPs with Bonferroni-significant effects for generation, selection, and the selection × generation interaction. Notably, 134,709 SNPs exhibit significant selection × generation interactions, highlighting pervasive, genome-wide divergence in allele frequency trajectories between reciprocal selection regimes. The plot illustrates that while some loci also show main effects of generation or treatment alone, the interaction effect dominates, consistent with highly polygenic, anti-parallel evolutionary responses.

Figure 6. Genome-wide heterozygosity across generations for reciprocal selection trajectories. Mean genome-wide heterozygosity was measured in replicate populations (n = 10 per trajectory) evolving under A→C (early-to-late reproduction) and C→A (late-to-early reproduction) selection regimes. Boxplots show the distribution of mean heterozygosity across replicates at each sampled generation, including earlier time points. A→C populations exhibit increasing heterozygosity over time, whereas C→A populations show a progressive decline. Stars above boxes indicate significant differences between founder and midpoint populations based on statistical tests.

Figure 7. Deep sequencing reveals allele frequency variation at sites previously classified as fixed. Histograms depict the minor allele frequency (MAF) of SNPs in the A-type1 population sequenced at ~800× coverage. Gray bars represent a random subset of SNPs equal in number to the pseudo-fixed sites identified in standard sequencing data, while purple bars indicate sites classified as pseudo-fixed under standard coverage (Bonferroni-corrected GLM, $p < 0.005$). Pseudo-fixed sites are overwhelmingly skewed toward low-frequency alleles, indicating that these loci appear fixed under standard coverage but retain hidden variation at higher sequencing depth. Permutation testing (10,000 replicates) confirmed that the observed mean MAF of pseudo-fixed sites (0.0063) is dramatically lower than expected under random sampling (mean null = 0.1745, SD = 0.00037; Z = –450.79, $P < 10^{-4}$), demonstrating that these pseudo-fixed sites represent a significant depletion of minor alleles relative to genome-wide expectations.

Figure 8. PCA of SNP frequencies after reciprocal selection shifts. Principal component analysis of SNP frequencies at the last collected timepoints for populations of the A→C (light blue) and C→A (light red) trajectories, alongside founder A (dark red) and C (dark blue) populations. Ellipses represent 95% confidence intervals around group centroids. C→A populations, which experienced 182 generations of selection, cluster tightly with the A founders, whereas A→C populations, with 65 generations of selection, show greater spread and incomplete convergence with C founders.

Figure 9. Genome-wide differentiation of SNP frequencies after reciprocal selection shifts.
Manhattan plots show beta-binomial GLMM results comparing allele frequencies at the final
sampled time point in reciprocally selected populations to long-established populations
maintained under the same selection regime: (A) A→C populations relative to C-type founders
and (B) C→A populations relative to A-type founders. Each point represents a SNP plotted by
chromosomal position (Mb) on the x-axis and −$\log_{10}$(Bonferroni-adjusted p-value) on the y-axis.
Dashed horizontal lines denote a significance threshold of α = 0.05 on the Bonferroni-adjusted p-
values. Chromosomes are color-coded as 2L (light blue), 2R (green), 3L (red), 3R (blue), and X
(orange).

Figure 10. Leave-one-out analysis comparing allele frequency changes at target SNPs identified from all but one replicate population to matched control SNPs. This analysis was carried out independently for the A→C and C→A trajectories. For each LOO iteration, the median allele frequency shift of target and matched control SNPs in the held-out replicate is shown, genome-wide and separately for each major chromosome arm for the A→C and C→A trajectories. Target SNPs were defined using FDR < 0.05 and ≥ 0.02 total change in allele frequency in the training set, with controls drawn from non-significant SNPs. Points represent medians from individual LOO iterations; the dashed line indicates no net allele frequency change.