Striatal and frontal signatures of social context and cost-benefit decision making in developmental stuttering

Nicole E. Neef[1,2]*, Elena Winter[1], Hellmuth Obrig[3,4], Andreas Neef[5], Toralf Mildner[6], Shaza Haj Mohamad[1], Christian H. Riedel[1], Katharina Scholze[2,7]

[1]Department of Diagnostic and Interventional Neuroradiology, University Medical Center Göttingen, Göttingen, Germany

[2]Department of Neuropsychology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[3]Clinic for Cognitive Neurology, University Hospital Leipzig, Leipzig, Germany

[4]Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[5]Göttingen Campus Institute for Dynamics of Biological Networks, University of Göttingen, Göttingen, Germany

[6]NMR Group, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[7]Research Center of the German Foundation for Depression and Suicide Prevention, Department of Psychiatry, Psychosomatics, and Psychotherapy, Goethe University, Frankfurt am Main, Germany

*Correspondence: nicole.neef@med.uni-goettingen.de

Short title: Stuttering and social speech

Keywords: nucleus accumbens, BA 47, stuttering anticipation, motivational salience, social cognition, context-dependent regulation

Abstract

Developmental stuttering is usually framed as a sensorimotor disorder, yet it manifests in communicative situations that engage motivational, self-referential, and regulatory processes. In this case-control study, we combined a socio-economic decision task outside the scanner with a socially modulated speech task during fMRI to test how listener presence and self-referential speech shape neural activity in 34 adults who stutter and 32 controls. Both groups valued talking with another person about themselves more than talking with another person about someone else or talking to themselves. On the neural level, listener-directed (vs. private) speech and self-disclosure (vs. guessing the preferences of a famous other) elicited stronger responses in the ventral striatum and medial prefrontal cortex, extending social valuation effects previously reported in fluent speakers to adults who stutter. Within the stuttering group, individual differences revealed a systematic reweighting of socially modulated activation as a function of symptom burden: higher anticipation of stuttering and greater overall impact were associated with stronger engagement of motivational circuitry, greater recruitment of frontal evaluative-control regions, and reduced contextual differentiation within speech-language cortex. Stuttering anticipation and lived experience gradually shift the balance between control, language, and motivational salience-processing systems, contributing to the disorder's marked heterogeneity and context sensitivity. These findings indicate system-level signatures of the interaction between social context and symptom severity, rather than isolated motor deficits, in developmental stuttering. More generally, they reveal how recurrent experiences shape brain activity through the interplay between language, motivational and control systems that governs human social interactions.

Introduction

Human speech is inherently social. Speaking to a listener carries motivational relevance (1) because it serves motivationally salient interpersonal goals such as affiliation (2,3), self-disclosure (4–6), and social evaluation (7,8). Listener presence fundamentally changes the incentives and consequences of speaking, as directed speech involves reputational and interpersonal stakes. By contrast, speech produced without a listener lacks these social consequences. Thus, social context modulates the value of speaking, and directed speech may recruit social-motivational control processes that shape speech production beyond lower-level sensorimotor mechanisms.

This social-context-dependent valuation of speaking is particularly relevant for developmental stuttering, a common speech disorder characterized by repetitions of sounds and syllables, prolongations, and blocks (9), and by pronounced sensitivity to social-evaluative context (10–12). Stuttering is highly situationally variable (13,14) and is markedly reduced during private speech (15,16) when utterances are not directed to a listener or embedded in a communicative goal. By contrast, adults who stutter (AWS) frequently report increased difficulty when speaking under conditions of audience presence or communicative pressure (17). On the neural level, perceived discomfort is associated with right amygdala activity during stuttering occurrence (18). Neuroimaging research has implicated alterations within cortico-basal ganglia-thalamo-cortical circuits involved in speech timing and motor control (19–21) yet it remains unclear how social context interacts with reward-related systems during speaking. Given the context sensitivity of stuttering, altered valuation of directed speech may represent a mechanism contributing to symptom expression (22,23).

A reward-related circuitry, including the nucleus accumbens and ventral striatum, supports the valuation of actions (24) and is reliably engaged by social rewards (25,26). Beyond responses to primary reward, ventral striatum is sensitive to being observed, receiving social feedback (27,28), and communicating self-relevant information (29,30). Consistent with a role in assigning motivational salience to social acts, both adults and children who stutter show altered grey-matter volume of the ventral striatum/nucleus accumbens, when compared to fluent speakers and typically developing peers (22,31,32). Interestingly, adults show increased right ventral striatum/nucleus accumbens morphology compared with fluent speakers, while children show reduced nucleus accumbens volume compared to their peers. This pattern points to an atypical developmental trajectory of striatal circuitry in stuttering. Rather than a uniform increase or decrease, it suggests that the motivational salience system for social communication is reorganized over time when language development is accompanied by a communication disorder such as stuttering. In this context, speech directed toward a listener is likely to recruit motivational circuits differently in people who stutter than in fluent speakers, with downstream effects on speech-motor and control systems in a listener-dependent manner.

Given that ventral striatum encodes the motivational salience of social communication and shows atypical development in stuttering, we aimed to test whether listener presence and self-referential content specifically modulate speech-related activity within this circuitry and its interaction with language and control networks. We investigated how social context modulates neural responses during speech, and whether this modulation differs between adults who do or do not stutter (AWS/AWNS). Participants completed a monetary decision task in which they either disclosed their own preferences or inferred the preferences of a well-known person, under share conditions (speech audible to the experimenter) or private conditions (speech inaudible) (29,30). This design enabled us to isolate the effect of directed speech (share) versus non-directed speech (private) while controlling for speech content. From a computational perspective, our paradigm requires participants to integrate social value, internal costs (e.g. effort, anticipated dysfluency), and monetary reward, engaging multiple valuation and control systems rather than a unitary 'speech' module. This follows proposals that ecologically valid paradigms combining several value dimensions are particularly informative for understanding neurodevelopmental and mental health-related differences in social behavior (33). We hypothesized that directed speech would increase activity in reward-related circuitry, particularly the nucleus accumbens, and that AWS would show altered social-context-dependent modulation relative to AWNS. To relate neural responses to subjective valuation, we additionally quantified the relative monetary value of speaking in these conditions in an independent behavioral examination (29).

Materials and methods

Participants

Thirty-four adults who stutter (AWS; 29 males, five females; aged 19-46 years; one left-hander) and 32 age- and sex-matched adults who do not stutter (AWNS; 28 males, four females; aged 21-46 years; one left-hander) took part in a larger project on persistent developmental stuttering. Handedness was assessed with the Edinburgh Inventory (34). AWS were recruited via advertisements and social media of the German Stuttering Association; AWNS were recruited from the Max Planck Institute for Human Cognitive and Brain Sciences volunteer database. Stuttering in AWS had been previously confirmed by a speech-language pathologist, and all self-identified as people who stutter. AWNS reported no personal or family history of stuttering or other speech or language disorders. All participants had normal or corrected-to-normal vision, met MRI safety criteria, and provided written informed consent. This study was approved by the University of Leipzig ethics committee [123/17-ek].

Stuttering severity of AWS was scored using the Stuttering Severity Instrument-4 (SSI-4) (35) and the German Version of the Overall Assessment of the Speaker's Experience of Stuttering (OASES) (36,37). SSI-4 outcomes ranged from very mild (n = 11), mild (n = 8), moderate (n = 6), severe (n = 2), to very severe (n = 7). OASES outcomes ranged from mild (n = 1), mild-to-moderate (n = 6), moderate (n =

16), moderate-to-severe (n = 7), severe (n = 2) to very severe (n = 1). Anticipation of stuttering was measured with the Premonitory Awareness in Stuttering Scale (PAiS) (38), a 12-item self-report questionnaire adapted from the Premonitory Urge for Tics Scale (39). Example items include: "Right before I stutter, I can sense that I am about to stutter," and "Right before I stutter, I feel tense." Trait and state anxiety were assessed with the German version of the State-Trait Anxiety Inventory (40), revealing modestly elevated but non-clinical anxiety levels in AWS relative to AWNS (Table 1).

Behavioural task: subjective value of speaking

The behavioural paradigm was adapted from Tamir and Mitchell (29) to assess social decision-making under monetary trade-offs. Participants chose between answering a question aloud or waiting passively, allowing quantification of the subjective value assigned to speaking in different social contexts. Four conditions were tested: *self share* (verbal report of one's own preference to a listener), *self private* (spoken but unheard), *other share* (verbal report of beliefs about a public figure's preferences to a listener), and *other private* (spoken but unheard). Angela Merkel was selected as the "other" due to her high familiarity at the time of the study, enabling plausible belief formation. Each trial required a two-alternative forced choice between speaking and waiting for 5 seconds. Both options were associated with small monetary outcomes (€0.01 - €0.04), with the relative payoff between speaking and waiting varying randomly from −€0.03 to +€0.03. In waiting trials, participants viewed a fixation cross and performed no task. Participants were seated in a sound-attenuated booth, while the experimenter (listener) was located in a separate room (Fig. 1A). Trial type, options, and payoffs were displayed at trial onset. Choices were made via button press. When speaking was selected, participants read the question aloud and responded "yes" or "no." Speech was recorded in all conditions but transmitted to the listener only during share trials. Practice trials demonstrated the share versus private manipulation. The task modified the original button-press paradigm by requiring overt speech, permitting assessment of speech production alongside socio-economic decision-making. Each condition was presented 60 times (240 trials total), randomly distributed across four runs (7:30 min each). Item assignment and trial order were randomized per participant. The task duration was approximately 45 minutes. Stimulus presentation was controlled using Presentation software (Neurobehavioral Systems). Questions concerned everyday, non-controversial preferences.

To estimate the subjective value of speaking, we computed the proportion of trials in which the speaking option was chosen at seven relative payoff levels (-€0.03 to +€0.03) for each participant and condition. These data were used to estimate the point of subjective equality (PSE), defined as the payoff difference at which speaking and waiting were chosen with equal probability, thereby indexing the subjective value of speaking relative to silence. Choice probabilities were fitted with cumulative normal functions using a Levenberg-Marquardt least-squares algorithm (Igor Pro 9, Wavemetrics, Fig. 1B). Negative PSE values indicate a willingness to incur a monetary cost to speak, whereas positive values indicate that

speaking required financial compensation. In cases where choice behaviour provided insufficient constraint (e.g. near-deterministic responding), weakly informative Gaussian priors were applied to stabilise parameter estimation without materially affecting well-constrained fits.

PSE values were fed into a repeated measures ANOVA with the between-subjects factor Group (AWNS versus AWS) and the within-subjects factors Mode (share versus private) and Person (self versus other). Group statistics were calculated with RStudio 2022.07.1 using the ez library.

Determination of stuttering in experimental recordings

All experimental recordings were assessed for stuttering occurrence with a scoring system (0 for no stuttering, 1 for stuttering). Speech blocks, prolongations, and part-word repetitions were considered stuttering. In addition, dysrhythmic repetitions of monosyllabic words were also classified as stuttering. Speech blocks were considered to occur when an unnatural pause appeared that was longer than 300 ms and did not correspond to the expected prosodic patterns, for example, a syntactically or semantically inappropriate pause in a position where no prosodic boundary would be typically expected. Pauses were also considered stuttering in case an audible interruption (glottis noise) occurred in the otherwise continuous phonation. To ensure conservative and reliable coding, ambiguous instances or unclear audio recordings were not labelled as stuttering. Other types of disfluencies, such as polysyllabic whole-word repetitions and interjections (e.g., "um," "like"), were not considered stuttering. Stuttering response rate was then calculated by dividing the number of stuttered trials by the total number of share trials for each participant. We used paired Wilcoxon Signed-Rank tests to assess whether the condition speaking privately or to someone else influenced the stuttering response rate in individuals who stutter.

To assess inter-rater reliability, a second rater independently evaluated 10% of the trials. Participant selection was stratified to reflect the overall distribution of stuttering severity in the full sample, resulting in a subset that included one participant with mild stuttering, six with moderate stuttering, two with mild-to-moderate stuttering, two with moderate-to-severe stuttering, and one with severe stuttering. For each of these 12 participants, the spoken trials from one run were analysed, with one of the four runs per participant chosen pseudo-randomly. This yielded a total of 549 reanalysed trials.

Cohen's Kappa was calculated to estimate inter-rater reliability, accounting for the binary nature of the data and its imbalanced distribution. The analysis yielded a Cohen's Kappa of 0.81 (95% CI [0.75, 0.87], indicating strong agreement according to McHugh, 2012 (41).

As task familiarity or habituation may influence stuttering behaviour over time, we conducted a Friedman test to examine whether the proportion of stuttered trials varied across runs. The analysis revealed no significant effect of time, $\chi^2(3) = 0.97$, $p = 0.81$, indicating that the proportion of stuttered trials remained stable throughout the experiment.

fMRI task: social modulation of speech

The fMRI paradigm was designed to assess whether social context modulates nucleus accumbens activity. To maximise the number of active trials, the waiting option used in the behavioural task was removed. Participants completed the same four conditions (*self share*, *self private*, *other share*, *other private*) and responded on every trial using a 5-point Likert scale ranging from *extremely unlikely* to *extremely likely*. During share trials, a study assistant outside the scanner listened to participants' responses via headphones (Fig. 2A). Each condition was presented 48 times (192 trials total), randomly distributed across four runs in an event-related design (9:40 min each). Trial order and item assignment were randomized per participant. The task lasted approximately 45 minutes and was implemented using Presentation software (Neurobehavioral Systems). All questions were novel and did not overlap with the behavioural experiment.

MRI data acquisition

Whole brain fMRI data were acquired using a 3 Tesla MAGNETOM Prismafit scanner (Siemens Healthineers, Erlangen, Germany) equipped with a 32-channel receive head coil, using a dual gradient-echo echo planar imaging sequence.(42) Scanning parameters were as follows: repetition time (TR) = 2000 ms; echo times (TEs) = 12/33 ms; partial Fourier factor = 6/8; bandwidth = 1966 Hz/Px; flip angle = 90°; voxel size = $2.5 \times 2.5 \times 2.5$ mm³; interslice gap = 0.25 mm; and field of view (FOV) = 204 mm. Image readout was accelerated by using GeneRalized Autocalibrating Partial Parallel Acquisition (GRAPPA) and multiband (SMS) acquisition, both by a factor of 2. Across 4 runs, a total of 1160 volumes were acquired per participant, each consisting of 60 slices aligned to the AC–PC plane and acquired in interleaved order. Before the start of the stimulation paradigm in each run, 30 auxiliary volumes were taken, which were used to calculate the weighting factors for the echo combination. Additionally, sagittal-oriented T1-weighted images were acquired using a 3D magnetization-prepared 2 rapid gradient echo (MP2RAGE)(43) sequence with the following parameters: inversion times (TIs) = 700/2500 ms; TR = 5000 ms; TE = 1.96 ms; bandwidth = 240 Hz/Px, flip angles = 4/5°, voxel size = $1 \times 1 \times 1$ mm³; acquisition matrix $256 \times 256 \times 176$; GRAPPA factor = 3; and acquisition time = 8:22 min.

MRI data analyses

A hybrid time series consisting of a weighted sum of both individual echo time series was created using Statistical Parameteric Mapping (SPM12, Welcome Trust Centre for Neuroimaging, UCL, London, UK) and an additional script both implemented in MATLAB (The MathWorks, Natick, MA, USA). In the first step, both echo time series (including their auxiliary volumes) were re-sliced using identical alignment parameters derived from the first echo's time series. Subsequently, the auxiliary volumes of

both echoes were used to calculate maps of the respective tSNR and weighting factors. Using these weighting factor maps, the final hybrid time series for the functional volumes was created by weighted summation.(44) Functional hybrid data were further analysed with FSL v6.0.7.16 using the FEAT pipeline (45,46). Preprocessing comprised spatial smoothing with a 5 mm full-width at half-maximum (FWHM) Gaussian kernel, high-pass temporal filtering (cut-off 90 s), co-registration to the individual anatomical image, and normalization to the MNI ICBM152 nonlinear asymmetric template (mni_icbm152_t1_tal_nlin_asym_09a). T1-weighted images were segmented using FastSurfer.(47)

At the first level, a general linear model was specified with the four tasks convolved with the hemodynamic response function; temporal derivatives were included as nuisance regressors. Within each participant, conditions were then combined using a fixed-effects analysis to model a 2×2 ANOVA with the factors Mode (share vs private), Person (self vs other), and their interaction. Contrast estimates (COPE images) were then entered into higher-level random-effects analyses to test main effects of Mode and Person, the Mode $\times$ Person interaction, and group differences (AWNS vs AWS) for each effect. Statistical inference was performed on the whole brain, using cluster-wise correction with a cluster-forming threshold of $Z > 3.1$ and a family-wise error-corrected cluster-extent threshold of $p < 0.05$ (48).

Correlation analyses with clinical measures

We examined whether individual differences in (i) anticipation of stuttering and (ii) overall stuttering impact were associated with the modulation of brain activity by social context. To this end, we performed additional cluster-level correlation analyses testing positive and negative associations between PAiS and OASES scores and neural modulation by Mode (share > private) as well as by Person (self > other). PAiS and OASES scores were not significantly correlated (Spearman $r(31) = 0.27$, $p =$ .131, two-tailed), whereas OASES scores showed a trend toward a positive association with SSI-4 total scores (Spearman $r(31) = 0.31$, $p = .081$, two-tailed). Because OASES scores showed a trend toward a positive association with SSI-4 total scores (Spearman $r(31) = 0.31$, $p = .081$, two-tailed), and to avoid redundancy and inflation of type-I error from including partially overlapping clinical measures, SSI-4 was not entered as an additional predictor in the voxel-wise correlation analyses. Instead, we focused on PAiS and OASES as conceptually distinct dimensions of stuttering experience. Age was included as a nuisance covariate. Statistical inference was based on the spatial extent of clusters in the respective contrasts, using cluster-wise correction with a cluster-forming threshold of $Z > 3.1$ and a family-wise error-corrected cluster-extent threshold of $p < 0.05$. Correlation analyses were implemented in FEAT using a fixed-effects model. Given that developmental stuttering is not a homogeneous disorder (50), together with the marked inter-individual variability in neural activation patterns (51), and the absence of robust group-level differences in the random-effects analyses in the critical contrast (share > private), we focused on within-sample correlations between contrast estimates and clinical scores. A fixed-effects model was therefore chosen to maximize sensitivity to these individual-difference associations, with the

understanding that inference is confined to this sample. We included age as a covariate of no interest to account for age-related variance in brain activation and clinical scores, ensuring that observed associations primarily reflect stuttering-related individual differences.

Results

Effects of group and social context on the subjective value of speaking

Statistics of the PSE values indicated a significantly higher relative monetary value of speaking in AWS compared with AWNS (Fig. 1C). A repeated-measures ANOVA revealed a main effect of Group ($F(1,57) = 5.1$, $p = .027$), with higher PSE values in AWS (mean = 1.47 cents, 95% CI [1.23, 1.70]) than in AWNS (mean = 0.83 cents, 95% CI [0.64, 1.03]). In addition, the ANOVA showed a Group × Mode × Person interaction ($F(1,57) = 4.1$, $p = 0.046$), as well as main effects of Mode ($F(1,57) = 9.1$, $p = 0.004$) and Person ($F(1,57) = 4.3$, $p = 0.042$). Across groups, participants were willing to sacrifice monetary reward to share information rather than keep it private (shared: mean = 1.36 cents, 95% CI [1.13, 1.60]; private: mean = 0.99 cents, 95% CI [0.78, 1.20]; Fig. 1D), and for disclose information about themselves rather than judge the preferences of another person (self: mean = 1.23 cents, 95% CI [0.53, 1.46]: other: mean = 1.12 cents, 95% CI [0.90, 1.34]; Fig. 1D).

To further characterize the three-way interaction, we conducted post hoc unpaired *t*-tests. These analyses suggested that AWS assigned higher value than AWNS to sharing information about themselves and to speaking about others in private; however, neither comparison survived correction for multiple testing. No group differences were observed for the remaining conditions (*other share* and *self private*; Supplementary Table 1).

No effect of listener presence on stuttering rate

We next tested whether stuttering rate was modulated by listener presence. In the *share* conditions, participants were aware that the experimenter could hear their speech. Conversely, in the *private* conditions they knew that their utterances were not audible to the experimenter. Stuttering rates were highly similar across share and private conditions (Supplementary Figure 1A) and showed a strong positive relationship, such that participants who stuttered more in the share condition also stuttered more in the private condition. Stuttering rates varied markedly across individuals, consistent with the broad range of symptom severity captured by the SSI-4 (Supplementary Figure 1B). In AWS, Wilcoxon signed-rank tests indicated no significant differences in stuttering rate between the *self share* and *self private* conditions ($V = 308$, $p = 0.052$), nor between the *other share* and *other private* conditions ($V = 200$, $p = 0.153$). Descriptive stuttering rate data for both AWS and AWNS are provided in Supplementary Table 2.

Effects of listener presence and self-relatedness on brain activity

Whole-brain analysis revealed significant effects of listener presence and self-relatedness of speech content. Speech produced in the presence of a listener (share > private) engaged widespread cortical and subcortical brain regions spanning medial and lateral prefrontal, premotor, posteromedial parietal, posterior cingulate, and temporo-parietal cortex, together with the caudate nucleus extending into the nucleus accumbens and cerebellar Crus I (Fig. 2B; Supplementary Table 3). Speech referring to the self rather than another person (self > other) recruited a distributed set of frontal and parietal association regions, including anterior cingulate, inferior frontal, and ventrolateral/orbitofrontal prefrontal cortex extending into the caudate nucleus, nucleus accumbens and thalamus, as well as precuneus and inferior parietal cortex, with additional involvement of lateral temporal and occipital regions, cerebellar Crus II, and hippocampus (Fig. 2B; Supplementary Table 3).

Listener-related modulation of brain activity by group

Listener-related modulation of brain activity (share > private) was observed in both groups and showed a largely overlapping spatial pattern. In AWNS and AWS, directed speech engaged medial prefrontal and premotor cortex, posterior cingulate and posteromedial parietal regions, as well as subcortical structures including the caudate nucleus, thalamus, and cerebellum. Additional frontal and temporal regions and extended activity of the nucleus accumbens were evident in the AWS map, whereas medial frontal and parietal components were more prominent in AWNS; however, the direct between-group comparison was not significant (Fig. 3; Supplementary Table 4).

Group differences in self-related speech responses

Self-related speech (self > other) engaged frontal and parietal association cortex in both groups, with prominent involvement of anterior cingulate and ventral frontal regions. In AWNS, self-related responses additionally extended into orbitofrontal and inferior frontal cortex, inferior temporal cortex, hippocampus, and brainstem regions encompassing the ventral tegmental area and dorsal raphe nucleus. In AWS, effects were more circumscribed, centered on anterior cingulate and frontal pole regions, with additional cerebellar involvement (Fig. 4; Supplementary Table 5).

Direct group comparison showed stronger self-related responses in AWNS than AWS in anterior cingulate cortex, whereas AWS showed greater responses than AWNS in medial inferior parietal cortex.

Correlates of stuttering anticipation in listener-related modulation

Within AWS, stuttering anticipation (PAiS) was associated with listener-related modulation of brain activity (share > private). Higher PAiS scores correlated positively with share-related responses in the nucleus accumbens (Fig. 5A; Supplementary Table 6). In contrast, negative correlations were observed in left inferior frontal cortex, encompassing pars opercularis, triangularis, and orbitalis, as well as posterior superior temporal regions including ventral posterior superior temporal sulcus and adjacent temporal cortex (Fig. 5B; Supplementary Table 6).

Correlates with stuttering-related burden in listener-related modulation

Within AWS, greater stuttering-related burden (OASES) was positively associated with listener-related modulation of brain activity (share > private). Higher OASES scores correlated with increased responses in medial frontal and cingulate regions, including medial prefrontal cortex, anterior cingulate cortex, and medial superior frontal cortex, as well as inferior frontal and frontal opercular regions (Fig. 6; Supplementary Table 7).

Discussion

Speaking is a motor act embedded in socially motivated behavior that engages valuation and control systems. In a socio-economic decision task where participants chose between speaking and remaining silent, both listener presence and self-related speech carried clear subjective value for AWS and AWNS extending prior findings from fluent speakers (29) to AWS. Using fMRI, we further found that listener-directed and self-related speech engage reward- and evaluation-related brain regions, including ventral striatum and medial frontal cortex, in both groups. Although listener presence did not produce group differences, individual variability in neural modulation was systematically related to stuttering anticipation and overall impact, indicating that social context shapes brain activity in AWS along clinically relevant dimensions.

In the discussion, we focus on aspects of the data that are specific to stuttering, namely, how inter-individual differences in anticipation of stuttering and overall stuttering impact relate to modulation of speech-related brain activity by social context. Moreover, the observed brain-behaviour associations were estimated using fixed-effects analyses and therefore formally apply only to the present sample, rather than supporting population-level inference. These exploratory correlations need to be interpreted as characterizing heterogeneity within the cohort of AWS, and as hypothesis-generating with respect to how clinically relevant dimensions of the stuttering phenotype may shape processing of speech in social contexts.

Prior structural and resting-state studies already indicate that cortico-basal ganglia and frontotemporal networks are altered in developmental stuttering, including anomalous resting-state network

architecture, atypical connectivity of speech-motor and auditory regions, and morphology of the nucleus accumbens that differs between children and adults who stutter (22,32,49,50). Our results extend this work by showing how these large-scale systems are differentially engaged by social context and stuttering load during active speech. Together with developmental MRI data showing distinct trajectories of striatal and cortical networks in children who stutter (23), our findings support the view that social communicative experience contributes to experience-dependent reorganization of cortico-basal ganglia loops over time.

Behavioral effects of listener presence

Despite reports that AWS avoid challenging communicative situations, we did not observe systematic avoidance in this low-stakes setting: AWS did not forgo monetary reward to avoid speaking, and stuttering frequency did not differ reliably between shared and private trials. This suggest that, under these conditions, the subjective value of speaking and self-disclosure remains high in AWS and can be dissociated from overt avoidance behavior. It also implies that the neural effects we observed are unlikely to be driven simply by increased dysfluency under social exposure, but instead reflect changes in the motivational and regulatory state in which speech is produced.

Premonitory awareness, motivational valuation and speech-language hubs

Within AWS, listener-related modulation of the nucleus accumbens scaled with individual differences in stuttering anticipation: higher PAiS scores were associated with stronger differentiation between directed and private speech in this ventral striatal region. Given the established role of the nucleus accumbens in encoding motivational relevance and subjective value in socially meaningful contexts (51–57), this pattern suggest that socially directed speech acquires greater motivational salience as anticipatory load increases, rather than simply becoming more rewarding. In multi-system terms, anticipatory load appears to amplify automatic motivational signals in ventral striatum, consistent with how affective cues bias approach-avoid decisions via striatal circuitry (33). Structural work has identified altered morphology of the nucleus accumbens in persistent developmental stuttering, with reduced volume in children but increased right-lateralized volume in adults (22,32), suggesting a non-linear developmental trajectory of ventral striatal circuitry. Our finding that listener-related modulation in this region scales with anticipatory load is consistent with the idea that atypical ventral striatal development is functionally expressed as heightened motivational salience of socially directed speech.

Higher stuttering anticipation was also associated with reduced listener-related modulation in cortical speech-language regions, including inferior frontal and posterior temporal areas that support speech

planning and perception (58–60), as well as the lateral orbitofrontal cortex (BA 47), implicated in controlled semantic processing and in integration of value and context into language and decision-making (61,62). These regions showed less differentiation between directed and private speech as anticipation increased, suggesting that increasing anticipatory load alters how socially directed speech is processed within cortical speech-language systems. One possibility is that, with higher anticipatory load, these regions are already strongly engaged during private speech, leaving less capacity for further modulation by listener presence. Alternatively, processing may shift away from fine-grained speech-language operations toward motivational-salience systems when social context is perceived as challenging.

Recent work indicates that the core language network, although not itself computing others' mental states (63,64), can be rapidly informed by feedback from theory-of-mind regions during naturalistic comprehension, with reliable functional coupling between these networks over time (65). Within broader prefrontal circuitry, lateral orbitofrontal and frontopolar regions have been linked to credit assignment and to counteracting biased value signals (33,66,67) complementing striatal mechanisms of value updating (68). In this context, concurrently reduced modulation in ventrolateral frontal regions, together with stronger listener-related responses in nucleus accumbens, is consistent with a shift in the relative engagement of ventral striatal motivational circuitry and cortical speech-language systems as stuttering anticipation increases.

Stuttering-related burden is associated with self-referential and regulatory regions

Listener-related increases in activity for share > private speech were observed in a medial prefrontal-cingulate network associated with social evaluation, self-referential processing, and context-dependent regulation (69–73). Their greater engagement for directed relative to private speech suggests that speaking under observation places additional demands on evaluative and monitoring processes. Within AWS, higher OASES scores were associated with stronger listener-related modulation in medial prefrontal and ventrolateral frontal regions, indicating that individuals who experience greater stuttering-related burden recruit this evaluative-regulatory circuitry more strongly under listener presence. OASES and PAiS were not correlated (Spearman $r(31) = 0.27$, $P = 0.13$, two-tailed), supporting the view that these neural associations track distinct aspects of the stuttering experience and phenotype. Within a multi-system framework (33), dorsal anterior cingulate cortex and medial prefrontal cortex may act as a hub, where learned negative consequences of speaking and current contextual cues jointly shape the value of speaking under observation.

The self > other contrast revealed group differences in regions implicated in self-referential and evaluative processing. Relative to AWS, AWNS showed greater engagement of the subcallosal cortex, often linked to affective valuation and self-related appraisal (74,75), whereas AWS exhibited stronger

activity in the medial inferior parietal cortex, including angular gyrus, associated with perspective-taking and integration of self-related information within a broader social context (76,77). Although these differences require replication, the pattern suggests that self-related speech may rely on partially distinct self-referential circuits in AWS and AWNS, potentially reflecting differences in how self-relevant information is evaluated or contextualized during speaking. On current accounts, ventromedial prefrontal cortex/subcallosal regions encode cognitive 'maps' for self-relevant value, whereas parietal regions support perspective-taking and contextual integration (33). Within this view, self-related speech in AWS may depend relatively more on distributed perspective-taking networks and relatively less on ventromedial prefrontal-centered self-valuation schemas, possibly reflecting experience-dependent adaptations to chronic social evaluative contexts. Together with the OASES correlations, these group differences indicate that both the evaluation of self-related speech and its modulation by listener presence vary systematically with the perceived impact of stuttering.

This dovetails with evidence that, after fluency-shaping therapy, greater relief from the social-emotional burden of stuttering is linked to lower fractional anisotropy in the right frontal aslant tract, suggesting that both frontal gray- and white-matter systems contribute to how stuttering burden is experienced and reduced. Consistent reports of altered frontotemporal and cerebellar-orbitofrontal connectivity that scale with stuttering severity (49) further suggest that individuals with higher burden recruit evaluative and regulatory hubs within a broader fronto-cerebellar and temporo-parietal network. In our task, this is reflected in stronger context-dependent engagement of medial prefrontal and ventrolateral frontal regions when speaking under observation.

Prior work using naturalistic interpersonal communication paradigms has shown that disfluent speech in adults who stutter is accompanied by right amygdala activity and reduced prefrontal engagement (18), highlighting a prominent role for limbic circuits during socially demanding speech. Our results complement these findings by showing that, even in a relatively low-stakes setting, higher stuttering-related burden shifts listener-related modulation toward medial prefrontal und lateral orbitofrontal valuation circuits. Together, these studies indicate that limbic, cognitive and sensorimotor systems jointly shape how social context is mapped onto brain networks in developmental stuttering.

Overall, our findings argue against a uniform neural deficit in developmental stuttering and instead support a shift in the balance between salience, control, and language processing systems that tracks anticipation and burden. This system-level perspective helps explain why stuttering is highly context-sensitive and heterogeneous and suggests that targeting motivational and evaluative processes, alongside speech motor mechanisms, may be important for understanding and treating the disorder.

Limitations

The paradigm isolates listener presence and self-relatedness but does not directly manipulate social evaluation or communicative consequences, and the reported associations are correlational rather than causal. More naturalistic interaction paradigms and longitudinal designs will be needed to clarify how motivational and regulatory systems interact with speech-motor processes over time (78). In addition, the sample size limits statistical power and the generalizability of the correlation findings, particularly given the marked heterogeneity of the stuttering phenotype. Nonetheless, the present results provide a principled starting point and concrete effect-size estimates for future, larger-scale studies that aim to further map how social-motivational and speech-related neural systems interact in people who stutter.

Conclusion

Overall, our findings argue against a uniform neural deficit in developmental stuttering and instead point to a shift in the balance between salience, control, and speech and language processing systems that tracks anticipation and burden. This multi-system perspective may help account for the context sensitivity and heterogeneity of stuttering and underscores the contribution of social and emotional factors to the neural organization of speech processing. Our findings do not directly speak to treatment efficacy but suggest that clinical assessment may benefit from explicitly considering how social exposure and stuttering anticipation modulate activity in salience, control, and speech-language systems, alongside traditional measures of overt fluency. This may help explain why the same individual can show very different fluency profiles across situations and support approaches that focus on reducing social and emotional load in challenging contexts rather than targeting speech motor behavior in isolation.

Supplementary material

Supplementary material is available at *bioRxiv* online.

Data availability

Data availability: The brain imaging data of the group averages in the standard space are available at: https://owncloud.gwdg.de/index.php/s/Jpd63stMeZZmBPW/download

Acknowledgements

We are grateful for the generous support of Angela D. Friederici. Additionally, we wish to thank all of our participants for volunteering their time.

Funding

This research was supported by the Max Planck Society and the University Medical Center Göttingen.

Competing interest

All authors report no competing interest.

CRediT statements

Nicole E. Neef: Conceptualization, Investigation, Data curation, Formal analysis, Project administration, Visualization, Supervision, Validation, Writing - original draft, Writing - review & editing. Elena Winter: Formal analysis, Writing - review & editing. Andreas Neef: Methodology, Formal analysis, Writing - review & editing. Toralf Mildner: Methodology, Writing - review & editing. Shaza Haj Mohamad: Formal analysis, Writing - review & editing. Hellmuth Obrig: Investigation, Writing - review & editing. Christian H. Riedel: Resources, Writing - review & editing. Katharina Scholze: Investigation, Data curation, Formal analysis, Writing - review & editing.

Disclosures

The authors have nothing to disclose.

References

1. Kawamichi H, Yoshihara K, Sasaki AT, Sugawara SK, Tanabe HC, Shinohara R, *et al.* (2015): Perceiving active listening activates the reward system and improves the impression of relevant experiences. *Soc Neurosci* 10: 16–26.

2. Hüttner N, Müller F, Cañal-Bruland R (2025): Dyadic Motive Fit: How Individual Differences in Dyads' Motives Predict Performance and Communication in Joint Action. *Motiv Sci* 11: 470–484.

3. Popliger M, Talwar V, Crossman A (2011): Predictors of children's prosocial lie-telling: Motivation, socialization variables, and moral understanding. *J Exp Child Psychol* 110: 373–392.

4. Rimé B, Bouchat P, Paquot L, Giglio L (2020): Intrapersonal, interpersonal, and social outcomes of the social sharing of emotion. *Curr Opin Psychol* 31: 127–134.

5. Willems YE, Finkenauer C, Kerkhof P (2020): The role of disclosure in relationships. *Curr Opin Psychol* 31: 33–37.

6. Ignatius E, Kokkonen M (2007): Factors contributing to verbal self-disclosure. *Nord Psychol* 59: 362–391.

7. Diel K, Grelle S, Hofmann W (2021): A Motivational Framework of Social Comparison. *J Pers Soc Psychol* 120: 1415–1430.

8. Scholl W (2013): The socio-emotional basis of human interaction and communication: How we construct our social world. *Soc Sci Inf* 52: 3–33.

9. Bloodstein O, Ratner NB (2008): *A Handbook on Stuttering*. Clifton Park, NY: Thomson/Delmar Learning.

10. Luckman C, Jackson ES (2025): A qualitative investigation of how stutterers perceive social interactions. *J Fluen Disord* 85: 106137.

11. Boyle MP, Rosen AL (2025): More than meets the eye: Self-rated covert stuttering is linked to reduced psychosocial and communicative outcomes. *J Fluen Disord* 86: 106162.

12. Ortiz-Alvarez A, Arenas R (2025): A phenomenological exploration of the contextual variability of stuttering. *J Fluen Disord* 84: 106120.

13. Tichenor SE, Yaruss JS (2021): Variability of Stuttering: Behavior and Impact. *Am J Speech-lang Pat* 30: 75–88.

14. Constantino CD, Leslie P, Quesal RW, Yaruss JS (2016): A preliminary investigation of daily variability of stuttering in adults. *J Commun Disord* 60: 39–50.

15. Jackson ES, Miller LR, Warner HJ, Yaruss JS (2021): Adults who stutter do not stutter during private speech. *J Fluen Disord* 70: 105878.

16. Andrews G, Howie PM, Dozsa M, Guitar BE (1982): StutteringSpeech Pattern Characteristics Under Fluency-Inducing Conditions. *J Speech, Lang, Hear Res* 25: 208–216.

17. Sheehan J, Hadley R, Gould E (1967): Impact of authority on stuttering. *J Abnorm Psychol* 72: 290–293.

18. Toyomura A, Fujii T, Yokosawa K, Kuriki S (2018): Speech Disfluency-dependent Amygdala Activity in Adults Who Stutter: Neuroimaging of Interpersonal Communication in MRI Scanner Environment. *Neuroscience* 374: 144–154.

19. Chang S-E, Guenther FH (2020): Involvement of the Cortico-Basal Ganglia-Thalamocortical Loop in Developmental Stuttering. *Front Psychol* 10: 3088.

20. Alm PA (2014): Stuttering in relation to anxiety, temperament, and personality: Review and analysis with focus on causality. *J Fluen Disord* 40: 5–21.

21. Cler GJ, Krishnan S, Papp D, Wiltshire CEE, Chesters J, Watkins KE (2021): Elevated Iron Concentration in Putamen and Cortical Speech Motor Network in Developmental Stuttering. *Brain* 144: 2979–2984.

22. Neef NE, Bütfering C, Auer T, Metzger FL, Euler HA, Frahm J, *et al.* (2018): Altered morphology of the nucleus accumbens in persistent developmental stuttering. *J Fluency Disord* 55: 84–93.

23. Neef NE, Chang S-E (2024): Knowns and unknowns about the neurobiology of stuttering. *PLOS Biol* 22: e3002492.

24. Levy DJ, Glimcher PW (2012): The root of all value: a neural common currency for choice. *Curr Opin Neurobiol* 22: 1027–1038.

25. Delgado MR, Fareri DS, Chang LJ (2023): Characterizing the Mechanisms of Social Connection. *Neuron* 111: 3911–3925.

26. Rilling JK, Gutman DA, Zeh TR, Pagnoni G, Berns GS, Kilts CD (2002): A Neural Basis for Social Cooperation. *Neuron* 35: 395–405.

27. Sobczak A, Yousuf M, Bunzeck N (2023): Anticipating Social Feedback Involves Basal Forebrain and Mesolimbic Functional Connectivity. *NeuroImage* 274: 120131.

28. Izuma K, Saito DN, Sadato N (2008): Processing of Social and Monetary Rewards in the Human Striatum. *Neuron* 58: 284–294.

29. Tamir DI, Mitchell JP (2012): Disclosing Information about the Self Is Intrinsically Rewarding. *Proc Natl Acad Sci* 109: 8038–8043.

30. Tamir DI, Zaki J, Mitchell JP (2015): Informing Others Is Associated With Behavioral and Neural Signatures of Value. *J Exp Psychol: Gen* 144: 1114–1123.

31. Montag C, Bleek B, Reuter M, Müller T, Weber B, Faber J, Markett S (2019): Ventral Striatum and Stuttering: Robust Evidence from a Case-Control Study Applying DAR[TEL]. *Neuroimage Clin* 23: 101890.

32. Chow HM, Garnett EO, Koenraads SPC, Chang S-E (2023): Brain developmental trajectories associated with childhood stuttering persistence and recovery. *Dev Cogn Neurosci* 60: 101224.

33. Scholl J, Klein-Flügge M (2018): Understanding psychiatric disorder by capturing ecologically relevant features of learning and decision-making. *Behav Brain Res* 355: 56–75.

34. Oldfield RC (1971): The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia* 9: 97–113.

35. Riley GD (2009): *SSI-4: Stuttering Severity Instrument*, Fourth Edition.

36. Yaruss JS, Quesal RW (2014): OASES: Overall Assessment of the Speaker's Experience of Stuttering (((Deutsche Übersetzung: H. A. Euler & & A. Alpermann), translators)).

37. Euler H, Kohmäscher A, Cook S, Metten C, Miele K (2016): *OASES-A: Erfassung Der Erfahrungen von Stotternden Erwachsenen [Deutsche Übersetzung Des OASES-A] [Overall Assessment of the Speaker' Experience of Stuttering (German Translation of the OASES-A)]*. Retrieved from https://stutteringtherapyresources.com/products/oases-german?variant=33149202530444

38. Cholin J, Heiler S, Whillier A, Sommer M (2016): Premonitory Awareness in Stuttering Scale (PAiS). *J Fluency Disord* 49: 40–50.

39. WOODS DW, PIACENTINI J, HIMLE MB, CHANG S (2005): Premonitory Urge for Tics Scale (PUTS). *J Dev Behav Pediatr* 26: 397–403.

40. Laux L, Glanzmann P, Schaffner P, Spielberger C (1981): *Das State-Trait-Angstinventar (STAI): Theoretische Grundlagen Und Handanweisungen.*

41. McHugh ML (2012): Interrater reliability: the kappa statistic. *Biochem Med* 22: 276–282.

42. Moeller S, Yacoub E, Olman CA, Auerbach E, Strupp J, Harel N, Uğurbil K (2010): Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial parallel imaging with application to high spatial and temporal whole-brain fMRI. *Magn Reson Med* 63: 1144–1153.

43. Marques JP, Kober T, Krueger G, Zwaag W van der, Moortele P-FV de, Gruetter R (2010): MP2RAGE, a self bias-field corrected sequence for improved segmentation and T1-mapping at high field. *Neuroimage* 49: 1271–1281.

44. Poser BA, Versluis MJ, Hoogduin JM, Norris DG (2006): BOLD contrast sensitivity enhancement and artifact reduction with multiecho EPI: Parallel-acquired inhomogeneity-desensitized fMRI. *Magn Reson Med* 55: 1227–1235.

45. Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM (2012): FSL. *NeuroImage* 62: 782–790.

46. Woolrich MW, Ripley BD, Brady M, Smith SM (2001): Temporal Autocorrelation in Univariate Linear Modeling of FMRI Data. *NeuroImage* 14: 1370–1386.

47. Henschel L, Conjeti S, Estrada S, Diers K, Fischl B, Reuter M (2020): FastSurfer - A fast and accurate deep learning based neuroimaging pipeline. *NeuroImage* 219: 117012.

48. Worsley KJ (2001): Statistical analysis of activation images. In: Jezzard P, Matthews PM, Smith S, editors. *Functional MRI: An Introduction to Methods*. OUP.

49. Sitek KR, Cai S, Beal DS, Perkell JS, Guenther FH, Ghosh SS (2016): Decreased Cerebellar-Orbitofrontal Connectivity Correlates with Stuttering Severity: Whole-Brain Functional and Structural Connectivity Associations with Persistent Developmental Stuttering. *Front Hum Neurosci* 10: 190.

50. Chang S-E, Angstadt M, Chow HM, Etchell AC, Garnett EO, Choo AL, *et al.* (2018): Anomalous network architecture of the resting brain in children who stutter. *J Fluency Disord* 55: 46–67.

51. Berridge KC, Robinson TE (1998): What is the role of dopamine in reward: hedonic impact, reward learning, or incentive salience? *Brain Res Rev* 28: 309–369.

52. Berridge KC, Kringelbach ML (2015): Pleasure Systems in the Brain. *Neuron* 86: 646–664.

53. Salamone JD, Correa M (2012): The Mysterious Motivational Functions of Mesolimbic Dopamine. *Neuron* 76: 470–485.

54. Knutson B, Greer SM (2008): Anticipatory affect: neural correlates and consequences for choice. *Philos Trans R Soc B: Biol Sci* 363: 3771–3786.

55. Smith DV, Clithero JA, Boltuck SE, Huettel SA (2014): Functional connectivity with ventromedial prefrontal cortex reflects subjective value for social rewards. *Soc Cogn Affect Neurosci* 9: 2017–2025.

56. Zink CF, Pagnoni G, Martin ME, Dhamala M, Berns GS (2003): Human Striatal Response to Salient Nonrewarding Stimuli. *J Neurosci* 23: 8092–8097.

57. Enriquez-Traba J, Arenivar M, Yarur-Castillo HE, Noh C, Flores RJ, Weil T, *et al.* (2025): Dissociable control of motivation and reinforcement by distinct ventral striatal dopamine receptors. *Nat Neurosci* 28: 105–121.

58. Bohland JW, Bullock D, Guenther FH (2010): Neural Representations and Mechanisms for the Performance of Simple Speech Sequences. *J Cogn Neurosci* 22: 1504–1529.

59. Guenther FH, Ghosh SS, Tourville JA (2006): Neural modeling and imaging of the cortical interactions underlying syllable production. *Brain Lang* 96: 280–301.

60. Guenther FH, Hickok G (2015): Chapter 9 Role of the auditory system in speech production. *Handb Clin Neurology* 129: 161–175.

61. Rudebeck PH, Murray EA (2014): The Orbitofrontal Oracle: Cortical Mechanisms for the Prediction and Evaluation of Specific Behavioral Outcomes. *Neuron* 84: 1143–1156.

62. Beer JS, Knight RT, D'Esposito M (2005): Controlling the Integration of Emotion and Cognition. *Psychol Sci* 17: 448–453.

63. Shain C, Paunov A, Chen X, Lipkin B, Fedorenko E (2022): No evidence of theory of mind reasoning in the human language network. *Cereb Cortex* 33: 6299–6319.

64. Paunov AM, Blank IA, Jouravlev O, Mineroff Z, Gallée J, Fedorenko E (2022): Differential Tracking of Linguistic vs. Mental State Content in Naturalistic Stimuli by Language and Theory of Mind (ToM) Brain Networks. *Neurobiol Lang* 3: 413–440.

65. Paunov AM, Blank IA, Fedorenko E (2019): Functionally distinct language and Theory of Mind networks are synchronized at rest and during language comprehension. *J Neurophysiol* 121: 1244–1265.

66. Lamba A, Nassar MR, FeldmanHall O (2023): Prefrontal cortex state representations shape human credit assignment. *eLife* 12: e84888.

67. Witkowski PP, Rondot LJ, Kurth-Nelson Z, Garvert MM, Dolan RJ, Behrens TE, Boorman E (2025): Neural mechanisms of credit assignment for delayed outcomes during contingent learning. *eLife* 13: RP101841.

68. Algermissen J, Swart JC, Scheeringa R, Cools R, Ouden HEM den (2024): Prefrontal signals precede striatal signals for biased credit assignment in motivational learning biases. *Nat Commun* 15: 19.

69. Chen Y, Huang Y, Yen N (2022): Role of anterior midcingulate cortex in self-reward representation and reward allocation judgments within social context. *Hum Brain Mapp* 43: 2377–2390.

70. Mitchell JP, Banaji MR, Macrae CN (2005): The Link between Social Cognition and Self-referential Thought in the Medial Prefrontal Cortex. *J Cogn Neurosci* 17: 1306–1315.

71. Northoff G, Bermpohl F (2004): Cortical midline structures and the self. *Trends Cogn Sci* 8: 102–107.

72. Somerville LH, Kelley WM, Heatherton TF (2010): Self-esteem Modulates Medial Prefrontal Cortical Responses to Evaluative Social Feedback. *Cereb Cortex* 20: 3005–3013.

73. Somerville LH, Jones RM, Ruberry EJ, Dyke JP, Glover G, Casey BJ (2012): The Medial Prefrontal Cortex and the Emergence of Self-Conscious Emotion in Adolescence. *Psychol Sci* 24: 1554–1562.

74. Palomero-Gallagher N, Eickhoff SB, Hoffstaedter F, Schleicher A, Mohlberg H, Vogt BA, *et al.* (2015): Functional Organization of Human Subgenual Cortical Areas: Relationship between Architectonical Segregation and Connectional Heterogeneity. *NeuroImage* 115: 177–190.

75. Mathiak KA, Alawi EM, Koush Y, Dyck M, Cordes JS, Gaber TJ, *et al.* (2015): Social Reward Improves the Voluntary Control over Localized Brain Activity in fMRI-based Neurofeedback Training. *Front Behav Neurosci* 9: 136.

76. Seghier ML (2012): The angular gyrus: multiple functions and multiple subdivisions. *Neurosci : a Rev J bringing Neurobiol, Neurol psychiatry* 19: 43–61.

77. Schurz M, Radua J, Aichhorn M, Richlan F, Perner J (2014): Fractionating theory of mind: A meta-analysis of functional brain imaging studies. *Neurosci Biobehav Rev* 42: 9–34.

78. Castellucci GA, Guenther FH, Long MA (2022): A Theoretical Framework for Human and Nonhuman Vocal Interaction. *Annu Rev Neurosci* 45. https://doi.org/10.1146/annurev-neuro-111020-094807

Tables

Table 1 Demographic details of participants, clinical features of patients

	AWNS (n=32)	AWS (n=34)	Difference, (P)
Age in years, mean (SD)	32.1 (6.7)	31.2 (6.4)	0.57 (0.574) [b]
Sex, male (%)	28 (88)	29 (85)	0.83 (1) [c]
Handedness, *Mdn* LQ (IQR)	100 (12.1)	90 (21.5)	619.5 (0.300) [d]
School education, *Mdn* (IQR)	4 (0)	4 (1)	**679.0 (0.033)** [d]
Professional education, *Mdn* (IQR)	7 (2)	6 (3)	645.0 (0.176) [d]
Trait anxiety percentile rank, *Mdn* (IQR)	27 (40)	65 (49) [a]	**253.5 (< 0.001)** [d]
State anxiety percentile rank, *Mdn* (IQR)	16 (43)	51 (38) [a]	**318.0 (0.005)** [d]
SSI-4 overall score, *Mdn* (IQR)	0 (0)	22 (14.5)	**0 (< 0.001)** [d]
% stuttered syllables, *Mdn* (IQR)	0 (0.3)	5.5 (8.8)	**0 (< 0.001)** [d]
OASES total score, *Mdn* (IQR)	n/a	2.8 (0.8) [a]	n/a
PAiS, *Mdn* (IQR)	n/a	18 (7.8) [a]	n/a

LQ = laterality quotient; IQR = interquartile range; SSI = Stuttering Severity Index;(35) OASES = Overall Assessment of the Speaker´s Experience of Stuttering; (37,41) Premonitory awareness in stuttering;(38); school and professional education were coded as ordinal variables. Highest school qualification was classified as: (0) no qualification, (1) general education school-leaving certificate after grade 9, (2) general education school-leaving certificate after grade 10, (3) university of applied sciences entrance qualification, and (4) higher education entrance qualification. Highest professional qualification was classified as: (0) no qualification; (1) vocational training <1 year; (2) vocational training <2 years; (3) vocational training <3 years; (4) completed vocational qualification; (5) currently enrolled at a university or university of applied sciences; (6) bachelor's degree or master craftsman qualification; (7) master's degree or equivalent; and (8) doctoral degree or higher. an = 33; differences were tested with unpaired, two-tailed t-test (R command: t.test), cFisher's Exact Test (R command: fisher.test), and d Wilcoxon Signed-Rank Test (R command: wilcox.exact).

Figures

Figure 1 Task outside the MRI scanner and results. (A) Experimental setup. Participants were seated in a soundproof booth and viewed the stimuli on a screen. The investigator, located outside the booth, viewed the same stimuli and listened to participants' responses during share trials only. (B) Representative individual data showing PSEs from one AWNS (left) and one AWS (right). The AWNS example shows the most negative PSE in the self-share condition, indicating a willingness to forgo 2 cents to share self-related information. The AWS example shows negative PSEs for both self- and other-related conditions, indicating a willingness to forgo approximately 1.5 cents to speak. (C) Group-level PSEs by condition, shown separately for each group (left) and pooled (right). (D) PSEs pooled across all participants and conditions.

Figure 2 Task inside the MRI scanner and results pooled across groups. (A) Experimental setup. Participants lay in the MRI scanner and viewed the stimuli and responded to each trial. The investigator, located outside the scanner room, viewed the same stimuli and listened to participants' responses during share trials only. (B) Whole-brain analysis revealed a main effect of Mode, with greater nucleus accumbens (NAc) responses during share compared with private trials, and a main effect of Person, with greater NAc responses during self compared with other trials (results pooled across groups). (C) For visualization purposes, we performed a whole-brain analysis across all conditions to illustrate task-positive speech-related activity. Abbreviations: aINS, anterior insula; IFG, inferior frontal gyrus; NAc, Nucleus accumbens; PPN, pedunculopontine nucleus; postCG, postcentral gyrus; post STSd, dorsal posterior superior temporal gyrus; preCG, precentral gyrus; preSMA, presupplementary motor area; SMA, supplementary motor area; VTA, ventral tegmental area.

Figure 3 Listener presence. For the share > private contrast, both AWNS (A) and AWS (B) showed increased BOLD responses in the nucleus accumbens (NAc) and caudate nucleus (Caud), together with bilateral anterior cingulate cortex (area 9m) and thalamic activity. In AWS, additional effects were observed in the left inferior frontal gyrus, pars orbitalis (area 47l), the left ventral posterior superior temporal sulcus (STSvp) extending into middle temporal gyrus, and posterior cingulate cortex (area 31pd).

Figure 4 Self-related speech. For the self > other contrast, both AWNS (A) and AWS (B) showed increased BOLD responses in anterior cingulate cortex (a24) and frontal pole. In AWNS, effects extended into orbitofrontal cortex (47m) and nucleus accumbens (NAc), with additional clusters in inferior frontal gyrus (44), inferior frontal junction (6r), and posterior inferior temporal gyrus (pITG). Group differences (C) were observed in right subcallosal cortex (s32) and left medial inferior parietal cortex (PFm).

Figure 5 Correlates of stuttering anticipation. In AWS, within the cluster mask of the share > private contrast (dark grey overlay), higher PAiS scores were positively associated with listener-related modulation in the right nucleus accumbens (NAc) (A), and negatively associated with modulation in left frontal operculum (FOP4), inferior frontal gyrus pars orbitalis (IFG), and ventral posterior superior temporal sulcus (STSvp) (B).

Figure 6 Correlates of overall stuttering experience. In AWS, within the cluster mask of the share > private contrast (dark grey overlay), higher OASES scores were positively associated with listener-related modulation in medial prefrontal and medial superior frontal cortex, anterior cingulate cortex (9m), and inferior frontal gyrus (pars orbitalis and triangularis) and frontal operculum (FOP4, 45).