Neuronal degeneracy reflects context-specific trade-offs between energy and information

Philip Sommer[a,b], Fleur Zeldenrust[c], Peter Jedlicka[d,e,f,+], Alexander D. Bird[d,e,g,+], *Jochen Triesch[a,+]

[a]Frankfurt Institute for Advanced Studies, Frankfurt am Main, Germany
[b]Department of Physiology, Anatomy and Genetics, University of Oxford, Oxford, United Kingdom
[c]Donders Institute for Brain, Behaviour and Cognition, Radboud University, Nijmegen, The Netherlands
[d]Computer-Based Modelling in the field of 3R Animal Protection, ICAR3R, Faculty of Medicine, Justus Liebig
University, Giessen, Germany
[e]Translational Neuroscience Network Giessen, Germany
[f]Institute of Clinical Neuroanatomy, Neuroscience Center, Goethe University, Frankfurt
[g]Ernst Strüngmann Institute (ESI) for Neuroscience in cooperation with the Max Planck Society, Frankfurt am Main, Germany

[+]Joint senior authors

*triesch@fias.uni-frankfurt.de

Keywords

degeneracy, information, energy-efficiency, neuronal firing rates, Pareto-optimality

In brief

We show that degeneracy of neuronal properties reflects structured trade-offs between energy efficiency and information transmission.

Highlights

- Neuronal variability is structured along a low-dimensional manifold reflecting degenerate solutions to an energy-information trade-off.

- We derive an updated neuronal energy budget, enabling instantaneous energy estimates from integrate-and-fire models.

- Our analysis predicts the trajectories of neuronal adaptation to energy deprivation.

- Optimal energy-information trade-offs are achieved for intermediate-low firing rates (2 – 5 Hz) as observed in neocortex.

- Synaptic efficacies are scaled non-multiplicatively during food deprivation. Modeling indicates that increased downscaling of strong synapses is energetically favorable.

Abstract

Neurons exhibit an impressive diversity. Even within the same cell type, firing rates can vary by several orders of magnitude, and key parameters — including resting membrane potential, membrane resistance, and synaptic inputs — differ substantially across neurons. It is presently unclear if this diversity reflects independent sources of variation such as a noisy expression of different ion channels in a neuron's membrane, or if there is any hidden structure in this large diversity of neuronal parameters and the neuron's resulting behavior. One potential source of such structure is evolutionary pressure that has sculpted brains to work in an energy-efficient manner.

Here, we use computational modeling to identify a manifold within a neuron's parameter space permitting energy-efficient signaling. Neurons whose parameters lie on this manifold represent degenerate solutions to the problem of energy-efficient signaling. We confirm that neuronal features in different sensory brain areas indeed fall on such a manifold and that food restriction induces systematic shifts of neuronal parameters on this manifold. Our results reveal that the large diversity of neuronal properties is tightly regulated to ensure energy-efficient signaling in the presence of different computational tasks and context.

Introduction

Neurons exhibit a striking variability in their intrinsic biophysical properties. Several studies (Marder and Goaillard, 2006; Goaillard and Marder, 2021) have investigated the vast space of neural parameters and shown that features such as ion channel densities can vary by two to three orders of magnitude across neurons of the same type. Work by Prinz *et al.* (2003, 2004) and Deistler *et al.* (2022) demonstrated that disparate, but well-balanced, combinations of synaptic and intrinsic parameters can give rise to similar functional outcomes (realistic neuronal and network behavior) (Marder and Goaillard, 2006), a phenomenon known as degeneracy. Degenerate solutions likely provide an evolutionary advantage by reducing the need for tight (homeostatic) regulation of individual ion channel levels while maintaining overall neuronal function (Marder and Prinz, 2002; Mishra and Narayanan,

2021; Stöber *et al.*, 2023). Furthermore, heterogeneous response properties can enhance the quality of stimulus representation and coding range (Padmanabhan and Urban, 2010; Mejias and Longtin, 2012; Perez-Nieves *et al.*, 2021; Gast *et al.*, 2024; Dahmen *et al.*, 2025). This flexibility also enables neurons to compensate for perturbations, such as ion channel knockouts (O'Leary *et al.*, 2013), increasing robustness, adaptability, and potentially evolvability (Schneider *et al.*, 2023). The question remains, however, how much of this variability comes from unavoidable, or unimportant, stochasticity in ion channel expression (Yang and Prescott, 2023) and how much is constrained by functional objectives.

Brains are energy-hungry, consuming 20% of the body's total energy despite accounting for only 2% of human body mass (Attwell and Laughlin, 2001) and evolution has driven them to work in an energy-efficient manner (Niven and Laughlin, 2008; Hyder *et al.*, 2013). Energy availability is a constraint on brain size (Isler and Schaik, 2009; Navarrete *et al.*, 2011), wiring patterns (Chen *et al.*, 2006), synaptic transmission (Harris *et al.*, 2015, 2019), neural coding (Levy and Baxter, 1996), and information processing (Schreiber *et al.*, 2002) with multiple factors contributing to energy efficiency (Yu and Yu, 2017; Padamsey and Rochefort, 2023). In food restriction experiments, Padamsey *et al.* (2022) have recently observed that individual neurons in mouse visual cortex can reduce their energy consumption at the cost of information transmission. Specifically, these neurons appear to reduce energy consumption by changing their integration properties through the adaptation of membrane resistance, resting potential, and synaptic efficacies.

Trade-offs between functional performance, energy efficiency, robustness, and flexibility are ubiquitous in neural systems and impose fundamental constraints on neuronal design (Giudice and Crespi, 2018). A powerful framework for analyzing such multi-objective trade-offs is Pareto-optimality (Shoval *et al.*, 2012; Remme *et al.*, 2018; Pallasdies *et al.*, 2021; Jedlicka *et al.*, 2022), which describes optimal solutions balancing multiple competing objectives. A neuron is said to be Pareto optimal if its performance in one task, for example information coding, cannot be improved without reducing its performance in at least one other task, such as energy consumption. In this framework, alternative neuronal parameter sets should form a Pareto front with different points representing different trade-offs between the competing objectives.

We propose that a large part of experimentally observed neuronal heterogeneity can be

understood as a manifestation of an information-energy trade-off, where neuronal pheno- types evolve towards and along a Pareto front maximizing information transmission while minimizing metabolic costs. Depending on the type of information encoded, availability of energy, and demanded precision, the energy-information trade-off might vary. To test this hypothesis, we analyze experimental data from two different primary sensory areas in rodents, each employing distinct coding strategies. We predict that the energy-information trade-off is a fundamental organizing principle of neural activity, shaping the observed diversity in neuronal properties.

Here we investigate energy-information trade-offs theoretically by combining data-driven simulations of an integrate-and-fire neuron model (Brette and Gerstner, 2005) and a detailed compartmental model (Goetz *et al.*, 2021) with information measures (Shannon, 1948; Schreiber, 2000) and updated energy calculations (Attwell and Laughlin, 2001; Howarth *et al.*, 2012; Yu *et al.*, 2017). Our energy estimates take a new approach, being based on both first principles and ion-counting in biophysical models. Our work elucidates how the interplay between energy constraints and information transmission shapes neuronal diversity. We provide novel evidence that neuronal variability is not random, but resides in a constrained, low-dimensional subspace; a subspace that differs depending on coding task and energy availability. By demonstrating that neurons operate on a structured manifold of energy-efficient solutions, we provide a new perspective on degeneracy in neural systems, with neuronal parameter variability emerging from fundamental trade-offs rather than arbitrary fluctuations (Edelman and Gally, 2001). This framework offers a novel explanation for the observed variability in neuronal parameters and suggests that metabolic constraints play a key role in sculpting cortical firing rate distributions and coding strategies.

Results

Degenerate neuronal parameters reflect an optimally balanced trade-off between energy efficiency and information transmission.

Neuronal parameters vary widely across brain regions and cell types in experimental recordings (Padamsey *et al.*, 2022; Zeldenrust *et al.*, 2024; Joshi *et al.*, 2025) (Fig. 1A and 2A,B). We searched for structure in this variability by calculating the energy consumption of an adaptive exponential integrate-and-fire model (AdExp) (Brette and Gerstner, 2005) using our updated energy budget (see Methods). For visual cortex (Fig. 1) we systematically varied membrane resistance, resting potential, and synaptic weights — parameters experimentally shown to change under metabolic stress (Padamsey *et al.*, 2022) — and evaluated neuronal performance in an orientation selectivity task (Fig. S1 — see Methods). For somatosensory cortex (Fig. 2) we explored the full parameter space extracted from experimental data (V_{rest}, R_m, C_m, V_{th}, V_{reset}, Δ_T, τ_{ad}, a_{ad}, b_{ad}) (Zeldenrust *et al.*, 2024) and assessed coding in a binary switching task (Fig. S11 — see Methods). We show that, rather than being arbitrary, the variability of neuronal parameters reflects the information-energy trade-offs made by different neurons in different situations. Using our newly derived estimates of neuronal energy consumption (see Methods), we identify a distinct subregion in the space of neuronal parameters where neurons transmit information in the most energy-efficient manner; this region lies close to the border where cells fall silent (Fig. 1B, shown as transparent circles). Experimental data from Padamsey *et al.* (2022) populate the predicted manifold (Fig. 1C), meaning that the recorded neurons lie along a structured Pareto-front representing optimal trade-offs between the two competing objectives of minimising energy consumption and maximising information transmission (proxied by orientation selectivity index, OSI). The Pareto front represents degenerate solutions to the biologically relevant tasks of the neuron.

When under metabolic stress from food restriction, neuronal parameters in the visual cortical dataset alter in a predictable manner. Neurons from energy-deprived animals exhibit an increased membrane resistance and a more depolarized resting potential compared to control animals (Padamsey *et al.*, 2022) (Fig. 1A). These changes are consistent with

a shift toward energy conservation (Fig. 3), but the neuronal parameters remain on the   112
predicted manifold. The neurons appear to dynamically adjust their parameters depending   113
on energy availability whilst moving along the Pareto front.   114

Sensory neurons operate on their Pareto-optimal frontier.   115

Sensory neurons must balance two fundamental objectives: energy efficiency and infor-   116
mation transmission. Using a Pareto-optimality framework, we show that neurons from   117
distinct cortical areas populate distinct Pareto fronts that each reflect this fundamental   118
trade-off (Fig. 1B and Fig. 2A,B). In the visual cortex, neurons lie at the Pareto-front spanned   119
by the energy efficiency and information selectivity (OSI) objectives (Fig. 1D), while in   120
the somatosensory cortex, neuronal feature combinations are distributed along a front   121
spanned by energy efficiency and mutual information (MI) (Fig. 2C,D). However the pa-   122
rameters of excitatory and inhibitory cells in somatosensory cortex lie on distinct manifolds.   123
These observations suggest that both area-specific demands (visual vs somatosensory) and   124
cell-type-specific properties (excitatory vs inhibitory) shape distinct energy–information   125
trade-offs.   126

Depending on functional requirements, sensory neurons may prioritize information ac-   127
curacy or reduced metabolic cost. For example, neurons processing rapidly changing   128
inputs may optimize for information fidelity, whereas neurons in energetically constrained   129
environments may shift toward more energy-efficient configurations (Remme *et al.*, 2018).   130
In line with this, neurons from visual cortex of food-restricted (FR) animals shift away   131
from the high-information region occupied by neurons from control (CTR) animals and   132
toward the energy-efficient regime (Fig. 1D). These energy-saving changes increase the   133
membrane time constant (by an increased membrane resistance), which strongly reduces   134
the ability of the neurons to encode changing stimuli but only slightly impairs orientation   135
selectivity. In contrast, somatosensory neurons — which did not experience metabolic   136
stress — do not populate the energy-conserving parameter space explored by neurons   137
from FR animals (Fig. 2G). Simulations with binary switching tasks (see Methods) reveal   138
that shifting parameters in this direction (as observed by (Padamsey *et al.*, 2022): increased   139
membrane resistance, depolarized resting potential, and reduced synaptic drive) does   140

reduce energy demand but simultaneously decreases the information transmission (MI) 141
substantially (Fig. 2H). 142

Together, these results indicate that, despite a diversity in parameters, neurons lie along an 143
energy–information Pareto frontier. The precise frontiers spanned by different neuronal 144
types in different contexts reflect their functional adaptation to the competing constraints 145
they experience biologically. 146

Firing rates are tuned to maximize energy-efficient coding. 147

The coding strategies employed by different neurons also reflect their energetic constraints 148
(Lennie, 2003). Sparse representations reduce the number of spikes required for informa- 149
tion transmission, maximizing computational efficiency (Olshausen and Field, 1996) and 150
energy efficiency (Levy and Baxter, 1996; Sengupta *et al.*, 2010). Levy and Baxter (1996) 151
and Laughlin and Sejnowski (2003) made the assumption that neurons maximize their 152
representational capacity per unit of energy expended to predict that firing frequencies 153
should be restricted to optimal ranges of around 6 Hz. Across cortical areas and cell types, 154
we identify a consistent range of optimal firing rates that maximize coding-efficiency 155
(Fig. 1E, Fig. 2E) prioritizing information transmission over energy-efficiency (reducing 156
their MI/E_{tot} ratio compared to simulation results — Fig. 1F). This range corresponds to 157
the low-to-medium firing rates commonly observed in the mammalian neocortex. 158

Notably, the exact position and shape of the coding-efficiency curves (Fig. 1E, Fig. 2E) 159
show distinct differences. This suggests that neurons are tuned to area- and cell-type- 160
specific coding demands, reflecting divergent computational roles. Moreover, visual and 161
somatosensory neurons display variations in the steepness and width of their optimal 162
efficiency curves, suggesting different tolerances to firing rate deviations, thus, indicating 163
differences in coding robustness and flexibility. Within barrel cortex, inhibitory neurons 164
appear suited to encode information at higher rates than excitatory neurons (Fig. 2E). 165
Furthermore, barrel cortex neurons do not exhibit combinations of high excitability and 166
slow adaptation — reflected in the absence of cells with both a high membrane resistance 167
(R_m) and long adaptation time constants (τ_{ad}) (Fig. 2F). This constrained distribution likely 168
reflects functional specialization, limiting biophysical variability to support task-specific 169

demands.   170

We propose that neurons homeostatically regulate their intrinsic excitability and post-   171
synaptic weights to maintain firing rates within their optimal range — consistent with   172
mechanisms of activity-dependent adaptation (Turrigiano *et al.*, 1994; Desai *et al.*, 1999;   173
Turrigiano and Nelson, 2004). Padamsey *et al.* (2022) observed that the food-restricted   174
visual cortical neurons did not have significantly different firing rates to the control case.   175
Self-tuning not only maximizes information efficiency but also contributes to population   176
sparsity, a hallmark of efficient coding at the network level.   177

Figure 1: **Visual cortex neurons balance energy efficiency and information transmission along a structured degeneracy manifold.**
A, Experimental data from food-restricted (FR, red) and control (CTR, black) animals (Padamsey *et al.*, 2022) show both a high variability in neuronal parameters and systematic shifts under metabolic stress. FR neurons exhibit an increased membrane resistance, depolarized resting potentials, and reduced synaptic currents. Ellipsoids represent two standard deviations. **B**, Simulated parameter space (AdExp model with an orientation selectivity task (Fig. S1) — see Methods). Silent neurons are shown as transparent circles, while spiking neurons are color-coded by orientation selectivity index (OSI; colorbar in C). A structured subspace of high information transmission per unit energy emerges, forming a Pareto-optimal manifold located near the boundary between active and silent regimes. **C**, Experimental data align with this predicted manifold, suggesting neurons maintain near-optimal trade-offs energy expenditure and information transmission. **D**, Visual cortex neurons (CTR and FR) populate a Pareto front between energy efficiency and orientation selectivity (OSI). Color indicates OSI per expended energy. Total energy consumption (E_{tot}) is computed using our updated energy budget (see Methods). **E**, Coding efficiency (mutual information per spike) is maximal at low-to-medium firing rates. Experimental neurons operate within this optimal range. Color indicates coding efficiency per unit energy. **F**, Mutual information per energy also peaks at low firing rates, offering an explanation for sparse cortical activity. Experimental cells prioritize information transmission over energy-efficiency thus populating the small MI/E_{tot} ratios left of the peak. Color indicates total energy expenditure.

Figure 2: **Distinct cortical areas and neuron types prioritize different energy–information trade-offs.**

A, Excitatory and **B**, inhibitory barrel cortex neurons (Zeldenrust *et al.*, 2024) shown as colored dots across three out of nine electrophysiological dimensions (V_{rest}, R_m, C_m, V_{th}, V_{reset}, Δ_T, τ_{ad}, a_{ad}, b_{ad}). Simulated neurons (AdExp model with binary switching task (Fig. S11 — see Methods — parameters interpolated from 9D principal component space) shown as a semi-transparent volume colored by normalized mutual information (MI). **C**, Excitatory (orange) and **D**, inhibitory (light blue) neurons lie along distinct Pareto fronts spanning high energy efficiency and high MI (normalized to 1), reflecting cell-type-specific optimization strategies. **E**, Coding efficiency (MI/spike) differs in peak and shape between cell types, indicating different optimal firing regimes and cost structures. **F**, No experimental neurons exhibit both high excitability (high R_m) and slow adaptation (τ_{ad}), suggesting functional specialization limits parameter combinations. **G**, Barrel cortex neurons do span the same manifold as visual cortex neurons and not enter the parameter region corresponding to food-restricted animals (red-shaded area, increased R_m, depolarized V_{rest}). **H**, Simulated transitions into the FR regime show a decline in mutual information (top) and energy consumption (bottom) in binary switching tasks (Fig. S11 — see Methods), demonstrating a trade-off where metabolic adaptations save energy at the cost of coding performance.

Refined neuronal energy budget with rate-dependence allows elevated firing rates in food-restriction being more energy-efficient than controls.

To understand neuronal energy expenditure more precisely, we introduce a refined energy budget model (for cortical pyramidal neurons) that builds upon previous estimates (Attwell and Laughlin, 2001; Howarth *et al.*, 2012; Yu *et al.*, 2017). Unlike these previous models, which assumed constant firing rates (4 Hz), our approach accounts for rate-dependent energy consumption, adaptation currents, non-spiking regimes, and additional ionic conductances (I_h). We demonstrate that including an additional conductance for hyperpolarization-activated cyclic nucleotide–gated (HCN) channels allows a neuron to increase its resting potential while reducing the ATP consumption by an additional $\sim 25\%$ (Fig. S8). Using a detailed multi-compartment neuronal model (Goetz *et al.*, 2021), we derive an improved approximation of the energetic cost of non-specific excitatory synaptic currents, enabling us to estimate the instantaneous energy state of integrate-and-fire neuron models (Fig. 3 — see Methods). This refined energy budget reveals that even the modestly (though not statistically significantly) elevated firing rates observed in food-restricted neurons (Padamsey *et al.*, 2022) are in fact energetically favorable. Although food-restricted neurons fire slightly more frequently compared to control, their substantially reduced excitatory synaptic currents lead to an overall decrease in total energy consumption. At the same time, the increase in firing rate partially compensates for reduced synaptic drive by preserving information transmission. Consequently, a small upward shift in firing rate represents an efficient trade-off: it maximizes information transfer for only a marginal energetic cost, allowing neurons to operate near an optimal information–energy balance under metabolic constraints.

Figure 3: **New refined neuronal energy budget estimate with synaptic and non-synaptic contributions and rate-dependence.**

A, Updated energy budget (for derivation see Methods) of cells from control (CTR), left, and food-restricted (FR), right, animals with six energy contributors. Cells from food-restricted animals show smaller total energy demand for all firing rates. **B**, CTR and **C**, FR energy components ratios projected out for the "optimal" firing rates of 4 Hz as indicated in the pie charts. They show different ratios of energy components between cells from CTR and FR animals.

Neurons follow predictable trajectories in response to energy deprivation.

To understand how neurons might respond dynamically to metabolic stress, we analyzed food-restricted neuronal cell data (Padamsey *et al.*, 2022) and simulated parameter adjustments under energy constraints (see Methods). We assume that neurons during energy shortages try to reduce their energy use, while still maintaining their coding abilities as well as possible. Maintaining the highest possible OSI among the neighbouring parameter combinations that allow lower energy expenditure predicts systematic trajectories in parameter space that:

- reduce excitatory synaptic input currents to conserve energy,

- increase membrane resistance and depolarize the resting potential to enhance excitability or hyperpolarize for small resting energies,

- maintain low firing rates to minimize the energetic cost of action potential generation and neurotransmitter release.

From any combination in the three dimensional parameter space (Fig. 1B) the trajectory moves into the optimal information-per-energy manifold, minimizing the total energy consumption and maintaining a near-optimal energy-information trade-off (Fig. 4). Once in this manifold the trajectories move along this manifold to combinations of high input resistances and depolarized resting potentials maintaining nearly constant firing rates as observed experimentally (Padamsey *et al.*, 2022). This confirms that biophysical parameters are strongly constrained by functional needs, while energy is minimized as long as function is maintained (Remme *et al.*, 2018).

The adaptive changes of membrane resistance, resting potential, and synaptic weights align with prior findings that neurons across multiple systems optimize energy efficiency (transmit maximal information per energy used) in axonal transmission (Levy and Baxter, 1996) as well as synaptic transmission (Lezmy *et al.*, 2021; Harris *et al.*, 2012) with presynaptic transmitter release (Levy and Baxter, 2002; Savtchenko *et al.*, 2013; Yuan *et al.*, 2018) and the postsynaptic conductance (EPSCs) (Harris *et al.*, 2015, 2019). This supports the

notion that energy constraints drive the evolution and development of neuronal biophysics (Niven and Laughlin, 2008), with neurons self-regulating their metabolic expenditure while continuously preserving functional output.

Figure 4: **Energy-deprived neurons follow a constrained adaptation trajectory, minimizing their energy expenditure, while maintaining their coding.**

A, Model neurons (Fig. 1B, AdExp model with an orientation selectivity task (Fig. S1) — see Methods) facing energy scarcity adjust their parameters to maintain an optimal energy-information trade-off. Trajectories in the three-dimensional parameter space (membrane resistance, resting potential, total synaptic weight) found altered in food-deprivation experiments (Padamsey *et al.*, 2022) first converge onto the predicted high-efficiency manifold, then shift within this manifold towards configurations that reduce energy consumption while preserving information transmission. **B**, Blue and purple traces illustrate example voltage responses before and after adaptation; the cells have preserved spiking activity (0.09 Hz vs 0.11 Hz) despite metabolic constraints. **C**, Normalized histogram of membrane voltages reveals a narrower distribution for depolarized resting potentials, while hyperpolarized resting potentials exhibit broader variability.

Non-multiplicative synaptic scaling, membrane noise, and adaptation currents jointly contribute to tuning curve broadening during energy scarcity.

We have seen that food restriction results in broader neuronal tuning curves (Padamsey *et al.*, 2022) — but there are multiple potential mechanisms that could drive this effect. To investigate its cause, we first inspected the observed synaptic scaling behavior. We examined whether synaptic weight scaling in energy-deprived neurons is multiplicative or non-multiplicative. Fitting the experimental mEPSC distributions (Padamsey *et al.*, 2022) with log-normal functions (Turrigiano *et al.*, 1998; Buzsaki and Mizuseki, 2014) revealed non-multiplicative weight scaling (Kolmogorov-Smirnov-test, $D = 0.44, p = 0.003^{**}$) (Fig. 5A). The non-multiplicative scaling disproportionately weakens strong synapses, reducing overall excitatory drive while preserving firing rates at lower metabolic cost. This is energetically favorable: neurons with non-multiplicative scaling reduce ATP consumption significantly because only a few changes in strong synapses are needed to control the mean firing rate: $1.75 \cdot 10^9$ ATP/s in multiplicative scaling (Fig. 5B) vs. $1.69 \cdot 10^9$ ATP/s in non-multiplicative scaling (Fig. 5C). However, downscaling stronger synapses disproportionately reduces the variance of the synaptic inputs resulting in a reduced dynamic range (Fig. S1E). This also decreases the dynamic range of the membrane potential, which degrades stimulus selectivity, as seen in the broader tuning curves of energy-deprived neurons (Fig. 5B–C) and overall affects the coding abilities. Extrapolating non-multiplicative synaptic weight scaling based on two experimental conditions (CTR and FR), however, comes with high uncertainties. Thus, as a conservative and well-established assumption, we applied multiplicative weight scaling in our other models (Turrigiano *et al.*, 1998); in reality, stronger synapses are downscaled disproportionately more than predicted by a strictly multiplicative model. Consequently, we expect the actual effects to be more pronounced.

A second factor contributing to tuning curve broadening is intrinsic membrane noise, with neurons from food-restricted (FR) animals consistently showing lower orientation selectivity than control (CTR) across all tested noise levels (Fig. 5E) when presented different oriented gratings (Fig. S1A–D). This reduction in selectivity is reflected in broader

tuning curves (Fig. 5D–F) and becomes more pronounced as noise increases. The resulting 263
heightened sensitivity of FR neurons to intrinsic noise likely stems from their reduced 264
voltage gap between resting and threshold potentials, making them more susceptible to 265
random fluctuations. In addition, experimental *ex vivo* data show rather constant levels of 266
membrane noise in FR neurons (0.9 mV/ms) and controls (0.8 mV/ms) (Fig. S9A-C). 267

Finally, we find that adaptation currents play a crucial role in tuning curve broaden- 268
ing. Simulations with an oriented grating inputs (Fig. S1 — see Methods) using a leaky 269
integrate-and-fire (LIF) model show minimal broadening, whereas the adaptive exponen- 270
tial integrate-and-fire (AdExp) (Brette and Gerstner, 2005; Zerlaut and Destexhe, 2017) 271
model captures the experimentally observed effects (Fig. S3). We propose that subthreshold 272
and spike-triggered adaptation currents suppress strong responses to preferred stimuli 273
while maintaining overall activity levels. Overall, this reduces orientation selectivity. 274

Figure 5: **Tuning curve broadening under energy deprivation is amplified by non-multiplicative synaptic weight scaling and membrane noise.**

A, Experimental *ex vivo* mEPSC distributions of neurons from control (CTR, black) and food-restricted (FR, dark red) animals (Padamsey *et al.*, 2022). A multiplicative downscaling of CTR weights (red) (Turrigiano *et al.*, 1998) fails to reproduce the experimental FR distribution, indicating a non-multiplicative scaling mechanism (dark red). CTR neurons show a distribution with mean $\mu_{CTR} = 9.70$ and standard deviation $\sigma_{CTR} = 6.92$, while FR neurons have a lower mean $\mu_{FR} = 5.26$ and narrower spread $\sigma_{FR} = 2.75$. Multiplicatively scaling the CTR distribution to match the FR mean yields $\mu_{CTR,scaled} = 5.26$ and $\sigma_{CTR,scaled} = 3.75$, indicating that the FR distribution cannot be explained by multiplicative scaling of the CTR distribution. **B**, Simulated tuning curves (AdEx model with an orientation selectivity task (Fig. S1) — see Methods) using multiplicative weight scaling (red) show broadening. **C**, In contrast, non-multiplicative downscaling of synaptic weights (dark red) shows more pronounced broadening, suggesting that strong synapses are preferentially weakened — reducing selectivity while improving energy efficiency. **D**, Simulated tuning curves without intrinsic membrane noise show smaller broadening compared to high membrane noise levels as in **F**. CTR neurons retain higher orientation selectivity (OSI, solid lines) and narrower tuning (lower full width at half maximum (FWHM), dashed lines) than FR neurons. **E**, Orientation selectivity index (OSI, solid) and full width at half maximum (FWHM, dashed) as a function of membrane noise amplitude. Solid lines show a monotonic decline in OSI for CTR as well as FR but with consistently lower values in the FR case. Dashed lines show an increased decline in FWHM for FR compared to CTR representing a loss in selectivity of FR neurons. Errorbars are within width of line. **F**, Tuning curve at high membrane noise (4 mV/ms) illustrates strong broadening and loss of selectivity in FR neurons, with markedly increased FWHM (dashed lines) compared to CTR neurons.

Resting potential adaptation enhances coding efficiency and is optimized *in vivo*.

Why do neurons adjust their resting membrane potential rather than their spiking threshold in response to metabolic stress, such as during food restriction (Padamsey *et al.*, 2022)? To address this, we compared scenarios in which the voltage gap between resting and threshold potential was modulated either by depolarizing the resting potential or by lowering the spiking threshold (AdExp model with an orientation selectivity task (Fig. S1) — see Methods). These two options are not symmetrical, as both excitatory and inhibitory synaptic reversal potentials remain fixed. Altering the range of voltages a neuron can experience below threshold will therefore change the strength and integration of synaptic inputs. Across a range of gap values the orientation selectivity index (OSI) was consistently higher when the resting potential was adapted (Fig. 6A – Top). By contrast, threshold adaptation led to a rapid decline in OSI and an increase in firing rate (Fig. S10A), reducing coding efficiency (Fig. S10E) and increasing metabolic cost (Fig. 6A – Bottom). These results indicate that shifting the resting potential maintains selectivity at lower energetic expense.

To further investigate this mechanism, we systematically varied the resting potential while maintaining a fixed voltage gap to threshold, spanning the range between inhibitory and excitatory synaptic reversal potentials. This revealed a distinct coding optimum — reflected in the orientation selectivity index (OSI) — for specific resting potentials (Fig. 6B – Top) with a higher overall selectivity in the control case. Strikingly, these optima coincide with the experimentally observed resting potentials in control and food-restricted animals (CTR: –72 mV, FR: –66 mV; dashed lines in Fig. 6B). Moreover, energy consumption tracked coding performance: control neurons were most efficient near their mean experimental resting potential, and food-restricted neurons around their mean resting potential (Fig. 6B – Bottom). Notably, transitioning from CTR to FR parameters shifts neurons toward lower overall coding capacity (indicated by dotted lines), but yields an improved energy efficiency. These findings suggest that cortical neurons dynamically tune their resting membrane potential to operate within a biophysically optimal regime that is coding-efficient as well as metabolically efficient.

Experimental data (Padamsey *et al.*, 2022) show variability in spiking thresholds, but mean

threshold values remain unchanged under food restriction. Instead, neurons adjust their 305
resting potentials to optimize energy efficiency. This suggests that resting potential modula- 306
tion is a primary mechanism for energy adaptation, while threshold adjustments may serve 307
different functional roles. In previous work we have found that sub-threshold adaptation 308
of the h-current increases information transfer when inputs vary in time (Zeldenrust *et al.*, 309
2024). Depression of the h-current leads to more false alarms and decreases information 310
transfer rates. On the other hand, threshold adaptation increases the working range of 311
neurons. This was not tested for in the food-restriction experiments (Padamsey *et al.*, 312
2022), but might represent another important difference in how neurons adapt to optimally 313
represent the stimuli they receive *in vivo*. 314

Figure 6: **Resting potential adaptation enhances coding performance and energy efficiency.**
A, Comparison between two mechanisms of increasing excitability — raising the resting potential
(turquoise) versus lowering the threshold (purple) — for a range of voltage gaps in orientation
selectivity tasks (Fig. S1 — see Methods). Errorbars are within width of line. **A – Top**, Orientation
selectivity index (OSI) is consistently higher when excitability is increased via resting potential
depolarization. **A – Bottom**, Total energy expenditure (E_{tot}) is markedly higher when reducing the
threshold, driven by elevated firing rates (Fig. S10A). **B**, Coding performance and energy cost as a
function of resting membrane potential, while keeping the rest–threshold voltage gap constant (35%
synaptic weight reduction in FR relative to CTR, based on experimental data, using an experimentally
measured membrane noise level of 0.9 mV/ms). CTR neurons have a 6 mV wider gap than FR
neurons. **B – Top**, Orientation selectivity index (OSI) across a range of resting potentials. Coding
performance peaks near the experimentally observed resting potentials for cells of control (–72 mV)
and food-restricted (–66 mV) animals (dashed lines) (Padamsey *et al.*, 2022). CTR neurons achieve
higher overall selectivity. Shaded areas indicate optimal coding regimes for CTR (black) and FR
(red) neurons. **B – Bottom**, Total energy consumption over the same resting potential range. Energy
efficiency closely follows coding performance: CTR and FR neurons are most efficient near their
respective *in vivo* resting potentials.

Discussion   315

We have demonstrated that there is hidden structure in the diversity of neuronal properties   316
reflecting an energy-information trade-off. We have developed an energy budget model   317
that accounts for neuronal energy expenditure based on firing rate and intrinsic neuronal   318
properties (see Methods). Using this model we have found that neurons control their   319
intrinsic properties to populate a neuron type- and context-dependent Pareto-optimal   320
space, balancing energy efficiency and information transmission. This trade-off predicts a   321
low to medium firing rate regime as observed in neocortex as particularly efficient. It also   322
predicts systematic adaptations under energy scarcity, favouring more energy-efficient   323
configurations at the expense of a relatively small information loss. While we have revealed   324
a hidden structure in the space of neuronal properties, reflecting neuronal degeneracy, it is   325
important to ask why there is so much diversity in neuronal parameters in the first place.   326

Benefits of neuronal diversity   327

Heterogeneity of intrinsic neuronal properties is ubiquitous in the brain and cells of the   328
same cell type show diverse structures and response properties (Altschuler and Wu, 2010;   329
Joshi *et al.*, 2025). It is conceivable that this diversity simply reflects randomness during   330
neuronal development. However, computational work has shown that such heterogeneous   331
populations of neurons improve the performance of neural networks by increasing the   332
variability in the representation of stimulus features (Duarte and Morrison, 2019; Shen *et*   333
al., 2023) resulting in optimized information transmission (Haggard and Chacron, 2023),   334
higher efficiency, and more robust learning relative to networks with homogenous units   335
(Zeldenrust *et al.*, 2021; Dahmen *et al.*, 2025). Furthermore, heterogeneity can serve as an   336
additional degree of freedom to reduce time and energy spent on hyperparameter tuning   337
when training artificial neural networks, which is one of the largest costs in practical   338
engineering applications (Perez-Nieves *et al.*, 2021).   339

Biological evidence for benefits of neuronal diversity comes from a study showing that   340
the loss of heterogeneity (difference of resting membrane potential to spiking threshold)   341
is linked to synchronization and epileptic behavior (Rich *et al.*, 2022), highlighting the   342

functional importance of intrinsic variability in biological brains. Heterogeneity of neuronal properties can also be spatially organized. Along the dorsal–to–ventral axis of the hippocampus, CA1 pyramidal neurons (Dougherty *et al.*, 2012) and dentate granule cells (Kumari and Narayanan, 2026) (with similar gradients reported in medial entorhinal cortex stellate cells) exhibit a systematic increase in input resistance and a concomitant depolarization of resting membrane potential. It is currently unclear what the functional benefit of this may be. Based on our findings we propose that these gradients reflect energy-efficient solutions to gradually changing information processing needs along this axis.

Energy efficiency has long been identified as a principle of nervous system organization (Barlow, 1961). This principle has been applied at various levels ranging from synaptic transmission (Goldman, 2004; Harris *et al.*, 2012) to information coding at the population level Barlow (1961). In this study, we have focused on the question of how coordinated changes in neuronal parameters affect information coding in single neurons response to sensory stimuli. Our work complements earlier studies showing that balanced excitatory and inhibitory currents give rise to more energy-efficient (bits/ATP) and informative (bits/spike) coding (Sengupta *et al.*, 2013; Yu *et al.*, 2018).

Managing changes in energy supply

Biological brains have to deal with diverse changes in energy supply. Food scarcity induces systematic changes in resting potential and membrane resistance, affecting neuronal integration properties (Padamsey *et al.*, 2022). An increased membrane time constant alters information integration. This might have big effects in tasks such as coincidence detection where high temporal precision is needed (Remme *et al.*, 2018). Changes in the distribution of time constants could impact network-level task performance, its learning and robustness (Perez-Nieves *et al.*, 2021; Shen *et al.*, 2023).

While such adaptations to food scarcity take weeks, sensory cortices may be able to modify their energy-information trade-off on much faster time sacles. Recent findings suggest that inhibitory circuits dynamically regulate coding precision and perception (Gauld *et al.*, 2024). The authors show that circuit properties and cell intrinsic properties work together to form a "latent" pool of neurons that is normally silenced by sensory-evoked inhibition,

but that can be artificially recruited via optogenetic stimulation to enhance perception. When activated, these neurons increase the encoding quality of a stimulus. These findings suggest that sensory cortices may be able to rapidly control their energy-information trade-off by altering the level of inhibition.

Interest in the energetic supply and demand of neurons has grown substantially in recent years due to the recognition that metabolic dysfunction is tightly linked to neurological disease (Bazzigaluppi *et al.*, 2017), e.g., in the case of mitochondrial dysfunctions (Klemmensen *et al.*, 2024). Although neurons are astonishingly flexible when handling transient energetic mismatches, chronic metabolic perturbations have detrimental effects on neuronal function, ultimately contributing to the development of neurodegeneration (Köhler-Solís and Schirmeier, 2025). Potentially, a long-term energy depletion results in a chronic depolarization and drifting reversal potentials making a network prone to seizures (Le Masson *et al.*, 2014). Our framework could provide insights into how neurons adapt to chronic energy deficiencies, such as those observed in Alzheimer's disease (Engl and Attwell, 2015).

Energy efficiency: from brains to engineering applications

Evolution has resulted in many ways of improving the energy efficiency of nervous systems. An improved understanding of the various mechanisms can also inspire engineering efforts to build more energy-efficient artificial computing systems. This has long been a major motivation behind the field of neuromorphic engineering (Indiveri, 2025). For example, recent approaches have considered mechanisms for reducing the energy consumption of encoding information (Zhang *et al.*, 2025; Dimakou *et al.*, 2025; N'dri *et al.*, 2025) or learning (Li and van Rossum, 2020; Pache and van Rossum, 2023). We believe that engineers still have much to learn from how biological nervous systems flexibly manage their energy budgets in the face of fluctuating supply and demand.

Conclusion

By integrating experimental data with computational modeling, we have identified degenerate solutions to the problem of neuronal parameter selection revealing an optimal energy-information trade-off. Our findings suggest that neurons operate within a constrained subspace of parameter combinations that ensure close to optimal information transmission. Our model predicts a high but well-orchestrated variability of neuronal parameters (reflecting degeneracy), which is context and cell-type-specific. Under energy limitation, neurons systematically shift their parameters along a Pareto-optimal frontier, reducing metabolic costs while preserving functional output as much as possible. These results provide a theoretical framework for understanding neuronal degeneracy, metabolic adaptation, and functional diversity across cortical areas. Our work highlights energy efficiency as a fundamental principle of neural computation, offering new insights into how neurons balance metabolic demands with information processing needs. Overall, efficient energy management presents itself as a promising research area spanning biological adaptation, disease mechanisms and technological design.

Acknowledgments

We want to thank Zahid Padamsey (Weill Cornell Medicine, Qatar) for his provision and explanation of the experimental data, as well as his feedback on our results. We are grateful to Arnd Roth (University College London) for his advice and expertise on detailed compartmental models and energy estimation. We are thankful to Richard Gao (Goethe University Frankfurt) for his advice on the SBI pipeline.

Author contributions

P.S., A.D.B., P.J., J.T. designed the study.

P.S. and A.D.B. performed the simulations and analyzed the data.

P.S., F.Z., A.D.B., P.J., J.T. wrote the paper. 421

F.Z. supervised the experiments. 422

We reanalyzed the experimental data from a previous study (Zeldenrust *et al.*, 2017). 423
In this study Fleur Zeldenrust and Tansu Celikel supervised the experiments. Fleur 424
Zeldenrust designed the experimental protocol. Fleur Zeldenrust and Niccolo Calcini 425
did the experimental data analysis. Niccolo Calcini, Koen Kole, Ate Bijlsma, Xuan Yang 426
performed the experiments. 427

During the preparation of this work the authors used ChatGPT-5 for inspirational language 428
work, improvements to human-generated texts for readability and style, and to ensure 429
that the texts are free of errors in grammar, spelling, punctuation, and tone as well as the 430
readability and debugging of code. After using this tool, the authors reviewed and edited 431
the content as needed and take full responsibility for the content of the published article. 432

Funding: 433
P.S. was supported by Studienstiftung des Deutschen Volkes. 434
F.Z. acknowledges funding from NWO (a Vidi grant VI.Vidi.213. 137 and "DBI2" (024.005.022), 435
a Gravitation program of the Dutch Ministry of Science, Education and Culture). 436
J.T. was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foun- 437
dation) through the Excellence Cluster EXC3066 "The Adaptive Mind" and the priority pro- 438
gram SPP 2041, project numbers TR 881/7-2; TR 881/8-2, priority program SPP 2411, project 439
numbers 520617944; TR 881/11-1 and Research Unit FOR 5368, DYNAMIC center, funded 440
by the LOEWE program of the HMWK (grant number: LOEWE1/16/519/03/09.001(0009)/98) 441
J.T. acknowledges support from the Johanna Quandt foundation. 442

The funders had no role in the study design, data collection, analysis, or decision to publish. 444

Declaration of interests 445

The authors declare no competing interests. 446

Material and methods

Experimental data

The raw experimental data is accessible by the original publications (Padamsey *et al.*, 2022; Zeldenrust *et al.*, 2024). Extracted data and example data is available with the analysis code on github (`https://github.com/trieschlab/energy-information-trade-off`).

Energy-deprivation data from Padamsey *et al.* (2022)

Padamsey *et al.* (2022) investigated the effects on food deprivation on layer 2/3 pyramidal neurons in visual cortex.

In vivo response to oriented gratings

Padamsey *et al.* (2022) presented drifting gratings (12 different angles from $0°$ to $330°$, $30°$ increments) with a temporal frequency of 1 Hz for 1.5 s in random order followed by a 1 s grey screen. The authors measured following neuronal parameters via *in vivo* electrophysiology:

- Current clamp recordings: 33 control and 26 food-restricted animals

- Voltage clamp recordings: 13 control and 10 food-restricted animals.

The key findings include:

- an 18 % mean increase in membrane resistance R_m,

- a 10 % mean increase in resting membrane potential V_{rest},

- a 42 % mean reduction of synaptic currents (excitatory and inhibitory),

- a constant spiking threshold V_{th},

- a constant spiking rate r_{post},   467

- a constant E-I ratio as quantified by the synaptic currents.   468

The *in vivo* electrophysiological measurements of a single cell contains its membrane resis-   469
tance R_m, its resting membrane potential V_{rest} and its spiking rate r_{post}. For comparison   470
with our simulations, the synaptic scaling parameter w_{scale} was estimated for each experi-   471
mental condition by fitting experimental firing rates to simulated firing rates integrating   472
5000 homogeneous Poisson inputs. The resulting synaptic weights were then scaled by a   473
factor to match experimental and simulated tuning curves of the non-homogeneous firing   474
of oriented grating presentation.   475

Ex vivo measurements of synaptic weight distributions   476

In *ex vivo* electrophysiology measurements of visual cortex neurons (Padamsey *et al.*, 2022)   477
recorded AMPAR-mediated mean miniature excitatory postsynaptic currents (mEPSC) at   478
-70 mV in 20 control and 20 food-restricted cells in the presence of 1 mM TTX (Padamsey *et*   479
al., 2022). Results showed: 480

- a decreased mean mEPSC amplitude,   481

- a constant mEPSC frequency.   482

The mEPSC distributions were fitted with log-normal functions as proxies for synaptic   483
weights (Buzsaki and Mizuseki, 2014) (see Fig. 5).   484

We assume that information stored in synaptic weights can be identified by the average   485
excitatory postsynaptic currents.   486

Estimation of membrane noise   487

Fluctuations in membrane voltage were estimated from existing *in vivo* blind patch clamp   488
recordings in control and food restricted mice (Padamsey *et al.*, 2022). It was first necessary   489
to remove experimental artefacts from the raw traces (Fig. S9A). These artefacts came in   490

three forms: 1) Sudden sharp deviations in voltage. 2) Low-frequency oscillations. 3) Slow baseline drift. 1) Sharp deviations were removed by cutting out $0.2\,\mathrm{ms}$ either side of voltages outside the windows $[-70\,\mathrm{mV}, -50\,\mathrm{mV}]$ (CTR) and $[-90\,\mathrm{mV}, -65\,\mathrm{mV}]$ (FR). 2) Slow oscillations were removed with Notch filters (SciPy Signal Processing package) centered around the first spectral peaks in each case (quality 0.1) (Fig. S9B). 3) Baseline drift was removed by subtracting the moving average (over 0.5 ms) from the voltage. Removing these artefacts gave cleaner estimates of Gaussian membrane voltage noise (Fig. S9C).

Simulation-based inference of neuronal parameters of switching task data of Zeldenrust *et al.* (2024)

Zeldenrust *et al.* (2024) measured the intrinsic information encoding properties of excitatory and inhibitory neurons in L2/3 of the mouse barrel (somatosensory) cortex. We found:

- excitatory neurons: high thresholds and strong adaptation resulting in sparse firing and a strong compression of information.

- inhibitory neurons: firing at high rates, resulting in a higher transfer of information.

To infer neuronal parameters from experimental voltage traces and injected currents from Zeldenrust *et al.* (2024), we employed simulation-based inference (SBI) using the `sbi` Python package (Tejero-Cantero *et al.*, 2020). Parameter inference was performed using a two-stage sequential neural posterior estimation (SNPE) framework (Greenberg *et al.*, 2019; Lueckmann *et al.*, 2021).

Pre-processing and fixed parameters. The resting potential (V_{rest}) was estimated as the y-intercept of a linear fit between the membrane voltage and the injected current trace. The spiking threshold (V_{th}) was taken directly from the experimental estimates reported in Zeldenrust *et al.* (2024). These parameters were fixed throughout the inference procedure.

Stage 1: Passive and spike-generation parameters. In the first stage, we inferred parameters governing passive membrane properties and spike initiation: membrane resistance (R_m), membrane capacitance (C_m), and the spike slope factor (Δ_T). Summary statistics

included the mean and standard deviation of the voltage trace, selected voltage percentiles   517
(5, 10, 25, 50, 75, 95%), spike count, and spike-triggered averages.   518

Priors were defined as: 519

- $R_m \in [25, 500]$ MΩ 520

- $C_m \in [2, 120]$ pF   521

- $\Delta_T \in [0.5, 5]$ mV   522

Posterior distributions were estimated using SNPE with direct sampling. If direct sampling   523
exceeded a predefined runtime threshold, posterior sampling was performed using Markov   524
chain Monte Carlo (MCMC) with slice sampling.   525

Stage 2: Adaptation-related parameters. In the second stage, we inferred parameters   526
governing post-spike and adaptive dynamics: reset potential (V_{reset}), adaptation time   527
constant (τ_{ad}), subthreshold adaptation strength (a_{ad}), and spike-triggered adaptation   528
increment (b_{ad}). Summary statistics included inter-spike interval (ISI) features (mean,   529
standard deviation, coefficient of variation), spike count, first and last spike times, and   530
after-spike averages.   531

Priors were defined as: 532

- $V_{\text{reset}} \in [V_{\text{rest}} - 10, V_{\text{th}} - 1]$ mV   533

- $\tau_{\text{ad}} \in [5, 500]$ ms   534

- $a_{\text{ad}} \in [1, 10]$ nS   535

- $b_{\text{ad}} \in [0.001, 0.1]$ nA   536

Posterior validation and dimensionality reduction. Posterior mean parameter estimates   537
were used to simulate voltage traces. Model quality was assessed by comparing experi-   538
mental and simulated spike trains using the Earth Mover's Distance (EMD) (Sihn and Kim,   539
2019), with data retained for further analysis only if EMD was below 270 ms.   540

One- and two-dimensional posterior marginals are shown in Fig. S12. Correlations between 541
inferred parameters across neurons are shown in Fig. S13. The quality-checked parameter 542
sets were projected into principal component space (Fig. S14) and used as basis for further 543
simulations. 544

Neuron model 545

We combine simulations of a generic leaky-integrate-and-fire (LIF) neuron model and an 546
adaptive exponential integrate-and-fire (AdExp) neuron model in conjunction with our 547
updated version of the energy calculations based on Attwell and Laughlin (2001); Howarth 548
et al. (2012); Yu *et al.* (2017) and information-theoretic measures. 549

Our simulations employ the leaky-integrate-and-fire (LIF) neuron model (Brunel and van 550
Rossum, 2007), which is a simplified neuron model capturing the essential dynamics of 551
neuronal behavior combined with computational simplicity: 552

$$C_m \cdot \frac{dV_m(t)}{dt} = g_L \cdot (V_{\text{rest}} - V_m(t)) + I_{\text{syn}}(t),$$

where C_m is the membrane capacitance, $V_m(t)$ the membrane potential, g_L the leak con- 553
ductance, V_{rest} the leak reversal potential, and $I_{\text{syn}}(t)$ the synaptic input current which is 554
modeled by conductance-based excitatory (AMPA) and inhibitory (GABA A) synapses. 555
The neuron spikes if the membrane voltage exceeds the threshold voltage V_{th} ($V_m > V_{\text{th}}$) 556
and is reset to V_{reset} after a spike was fired (for some of the simulations we kept it constant 557
at -60 mV but also set it to V_{rest} for others to check consistency — see Fig. S4). No qualitative 558
nor significant quantitative changes we observed except for the coefficient variation — see 559
Fig. S4.). The membrane time constant τ_m determines the temporal charge and discharge 560
properties of the membrane and is defined as: 561

$$\tau_m = C_m \cdot R_m = \frac{C_m}{g_L} \quad \text{with} \quad R_m = \frac{1}{g_L}.$$

To model biological spiking behavior more realistically the LIF can be extended by an adaptation current and an exponential term for spike initiation, which captures both subthreshold adaptation and spike-triggered adaptation. The adaptive exponential integrate-and-fire (AdEx) neuron model (Brette and Gerstner, 2005) can be represented by the following set of differential equations with the membrane potential dynamics:

$$C_m \cdot \frac{dV_m(t)}{dt} = g_L \cdot (V_{\text{rest}} - V_m(t)) + g_L \cdot \Delta_T \cdot \exp\left(\frac{V_m(t) - V_T}{\Delta_T}\right) - w_{\text{ad}}(t) + I_{\text{syn}}(t),$$

and the adaptation current dynamics:

$$\tau_{\text{ad}} \cdot \frac{dw_{\text{ad}}(t)}{dt} = a \cdot (V_m(t) - V_{\text{rest}}) - w_{\text{ad}}(t),$$

where Δ_T is the slope factor for the exponential term, V_T the exponential onset voltage, $w_{\text{ad}}(t)$ the adaptation current, τ_{ad} the adaptation time constant, a_{ad} the subthreshold adaptation, and b_{ad} the spike-triggered adaptation for which $w_{\text{ad}}(t)$ is incremented for each time $V_m(t)$ reaching the spiking threshold. These differential equations are numerically integrated with built-in functions of Brian2 based on Fourth-order Runge-Kutta algorithms.

Stochastic membrane noise

To simulate intrinsic membrane fluctuations, we added a noise term I_{noise} to the membrane equation. This current was modeled as uncorrelated Gaussian white noise with zero mean and a standard deviation σ_{noise} derived from the experimental *ex vivo* mEPSC recordings. The noise was scaled by the square root of the integration timestep Δt to ensure correct physical units and variance:

$$I_{\text{noise}}(t) = \sigma_{\text{noise}} \cdot \sqrt{\Delta t} \cdot \mathcal{N}(0, 1),$$

where $\mathcal{N}(0, 1)$ denotes a standard normal random variable. Due to numerical stability constraints in the presence of stochastic input, simulations including I_{noise} were performed using the forward Euler integration scheme instead of the default Runge–Kutta solver in

Brian2.   582

Model oriented gratings as synaptic input   583

The synaptic input current is the sum of all excitatory and all inhibitory synaptic inputs: 584

$$I_{\text{syn}}(t) = I_{\text{syn}_e}(t) + I_{\text{syn}_i}(t).$$

The synaptic inputs are modeled by conductance-based excitatory and inhibitory synapses: 585

$$I_{\text{syn}_e}(t) = g_e(t) \cdot (E_e - V_m(t)),$$

$$I_{\text{syn}_i}(t) = g_i(t) \cdot (E_i - V_m(t)),$$

where $g_e(t)$ is the excitatory synaptic conductance, E_e the reversal potential of excita-   586
tory synapses, $g_i(t)$ the inhibitory synaptic conductance, and E_i the reversal potential of   587
inhibitory synapses. The dynamics of the conductances is described by: 588

$$\frac{dg_e(t)}{dt} = -\frac{g_e(t)}{\tau_e},$$

$$\frac{dg_i(t)}{dt} = -\frac{g_i(t)}{\tau_i},$$

where τ_e is the postsynaptic potential (PSP) time constant for excitatory synapses and τ_i for   589
inhibitory synapses. For every presynaptic spike the respective conductance is increased by   590
the size of the synaptic weight w_e or w_i. The excitatory synaptic weights are approximated   591
as log-normally distributed with the *ex vivo* measured mEPSC distribution of the control   592
(CTR) animals with $\mu_{w_e} = 2.07$ and $\sigma_{w_e} = 0.64$ (Padamsey *et al.*, 2022), where the mESPCs   593
are assumed as a proxy for the synaptic weights (Song *et al.*, 2005). The same weight   594
distribution is assumed for inhibitory synaptic weights but their total weight is scaled to   595
25 % of the total excitatory synaptic weight to result in an overall balance of excitation   596
and inhibition (Deneve and Machens, 2016). The synaptic inputs are excitation-inhibition   597

balanced as experimentally measured ([Padamsey *et al.*, 2022](); [Haider *et al.*, 2013](); [Barral and Reyes, 2016]()). Thus, we maintain in our simulations a constant ratio between excitation and inhibition with the conductance ratio of $g_e/g_i \approx 0.52$ and the synaptic current ratio of $I_{\mathrm{syn}_e}/I_{\mathrm{syn}_i} \approx 1.5$ consistent with experimental measurement. The constancy of the excitation-inhibition balance is achieved by scaling excitatory and inhibitory synapses by the same factor w_{scale} proportionately (coupling them). For $w_{\mathrm{scale}} = 100$ the mean excitatory synaptic weight is $w_e = 0.52$ nS and the mean excitatory synaptic weight is $w_i = 0.4$ nS. The excitatory presynaptic firing rates arriving at the synapses are normally distributed with $\mu_{r_e} = 1.37$ Hz and $\sigma_{r_e} = 0.23$ Hz ([Goetz *et al.*, 2021]()). The inhibitory presynaptic firing rates are assumed to be uniform with a firing rate of 6 Hz. We included 5000 synapses in total from which 75 % (3750) are excitatory and 25% (1250) are inhibitory (the latter being driven by a homogeneous Poisson process). Of the excitatory synapses 50 % code for a presented stimulus (driven by an inhomogeneous Poisson process) and 50 % are assumed noise synapses (driven by a homogeneous Poisson process) (see Fig. S1C, Fig. S2).

To model *in vivo* like synaptic input of oriented gratings for layer 2/3 pyramidal neurons we assume that stimuli are solely coded in firing rate (rate-based code) and match the firing rates and synaptic weights of the coding synapses. This is motivated by the findings that the connection between neurons with similar receptive fields are stronger ([Goetz *et al.*, 2021]()) compared to unrelated receptive field properties. The matching logic is performed similarly to [Goetz *et al.* (2021)]() (see Fig. S1A–C):

- For the preferred orientation, large synapses are assigned to high firing rates, small synapses with low firing rates.

- For the non-preferred orientation, large synapses fire with low firing rates, small synapses with high firing rates.

- For the intermediate orientations, we interpolate between preferred and non-preferred and draw the firing rates from two Gaussian distributions with same mean and variance.

For simulation purposes different oriented gratings are presented for 1.5 s with 1 s intermittent background firing and with the same increments as in the food-deprivation

experiments (Padamsey *et al.*, 2022) following the scheme 0°, 30°, 60°, 90°, 120°, 150°, 180° 627
with 90° as the preferred stimulus and 0° and 180° as the non-preferred stimulus (Fig. S1A). 628
For more references check Goetz *et al.* (2021). 629

Parameter	Value	Unit	Reference
C_m	100	pF	Padamsey *et al.* (2022)
V_{th}	-50	mV	Zerlaut and Destexhe (2017)
V_{reset}	V_{rest}	mV	
E_e	0.0	mV	Zerlaut and Destexhe (2017)
E_i	-80.0	mV	Zerlaut and Destexhe (2017)
τ_e	3.0	ms	Gerstner *et al.* (2014)
τ_i	5.0	ms	Gerstner *et al.* (2014)
Δ_T	2	mV	Zerlaut and Destexhe (2017)
V_T	$V_{\text{thresh}} - 2.5 \cdot \Delta_T$	mV	Zerlaut and Destexhe (2017)
τ_{ad}	500	ms	Zerlaut and Destexhe (2017)
a	4	nS	Zerlaut and Destexhe (2017)
b	0.02	nA	Zerlaut and Destexhe (2017)
N_{syn}	5000	-	Goetz *et al.* (2021)
N_e	3750	-	Goetz *et al.* (2021)
$N_{e,\text{signal}}$	1875	-	-
N_i	1250	-	Goetz *et al.* (2021)
μ_{w_e}	2.07	-	Padamsey *et al.* (2022)
σ_{w_e}	1.14	-	Padamsey *et al.* (2022)
μ_{w_i}	2.07	-	Padamsey *et al.* (2022)
σ_{w_i}	1.14	-	Padamsey *et al.* (2022)
μ_{r_e}	1.37	Hz	Goetz *et al.* (2021)
σ_{r_e}	0.23	Hz	Goetz *et al.* (2021)
μ_{r_i}	6	Hz	Goetz *et al.* (2021)
σ_{r_i}	0	Hz	Goetz *et al.* (2021)

Table 1: Parameters for the LIF and AdExp neuron model and oriented gratings synaptic inputs.

Model switching ON-OFF states as synaptic input

Following the approach of Zeldenrust *et al.* (2017), we modelled the stimulus as a hidden binary state (Markov process) switching stochastically between preferred (ON) and non-preferred (OFF) conditions. The transition from OFF to ON occurred at a rate r_{ON}, and from ON to OFF at r_{OFF}. The resulting switching time constant of the hidden state is given by:

$$\tau_{\text{switch}} = \frac{1}{r_{\text{ON}} + r_{\text{OFF}}}.$$

In line with experimental observations (Zeldenrust *et al.*, 2024), excitatory and inhibitory neurons exhibit distinct operational timescales — ranging from a few Hz in excitatory cells to over 20 Hz in inhibitory neurons. Accordingly, the switching dynamics of the inputs were adapted: we used $\tau_{\text{switch}} = 50$ ms for excitatory and 250 ms for inhibitory neurons to match their respective membrane integration properties.

Synaptic inputs were modelled analogously to earlier sections: excitatory and inhibitory synaptic weights were drawn from a log-normal distribution (Fig. S11A), excitatory presynaptic firing rates from a normal distribution (Fig. S11B), and inhibitory rates were set to a uniform distribution. Stimulus encoding was implemented via a matching scheme: during ON epochs, strong synapses were paired with high firing rates, whereas during OFF epochs, this pairing was inverted (Fig. S11C,D):

- For the preferred stimulus (ON) large synapses are assigned to high firing rates, small synapses with low firing rates.

- For the non-preferred stimulus (OFF) large synapses fire with low firing rates, small synapses with high firing rates.

The corresponding firing rates were translated into spiking input with an inhomogeneous Poisson process (a time-varying rate code).

For subsequent analyses, parameter grids spanning ± 3 standard deviations in the nine-dimensional principal component space (see Fig. S14) were simulated separately for excita-

tory and inhibitory neurons.

Parameter	Value	Unit	Reference
R_m	100	mΩ	Zeldenrust *et al.* (2024)
C_m	50	pF	Zeldenrust *et al.* (2024)
V_{rest}	-70	mV	Zeldenrust *et al.* (2024)
V_{th}	-63	mV	Zeldenrust *et al.* (2024)
V_{reset}	V_{rest}	mV	
E_e	0.0	mV	Zerlaut and Destexhe (2017)
E_i	-80.0	mV	Zerlaut and Destexhe (2017)
τ_e	3.0	ms	Gerstner *et al.* (2014)
τ_i	5.0	ms	Gerstner *et al.* (2014)
Δ_T	1	mV	Zeldenrust *et al.* (2024)
V_T	$V_{\text{thresh}} - 2.5 \cdot \Delta_T$	mV	Zerlaut and Destexhe (2017)
τ_{ad}	10	ms	Zeldenrust *et al.* (2024)
a	4	nS	Zeldenrust *et al.* (2024)
b	0.0805	nA	Zeldenrust *et al.* (2024)
N_{syn}	1333	-	Goetz *et al.* (2021)
N_e	1000	-	Zeldenrust *et al.* (2024)
$N_{e,\text{signal}}$	500	-	-
N_i	333	-	-
r_{ON_e}	1.3	-	Zeldenrust *et al.* (2024)
r_{OFF_e}	2.7	-	Zeldenrust *et al.* (2024)
r_{ON_i}	6.7	-	Zeldenrust *et al.* (2024)
r_{OFF_i}	13.3	-	Zeldenrust *et al.* (2024)
μ_{w_e}	2.07	-	Padamsey *et al.* (2022)
σ_{w_e}	1.14	-	Padamsey *et al.* (2022)
μ_{w_i}	2.07	-	Padamsey *et al.* (2022)
σ_{w_i}	1.14	-	Padamsey *et al.* (2022)
μ_{r_e}	1.37	Hz	Goetz *et al.* (2021)
σ_{r_e}	0.23	Hz	Goetz *et al.* (2021)
μ_{r_i}	6	Hz	Goetz *et al.* (2021)
σ_{r_i}	0	Hz	Goetz *et al.* (2021)

Table 2: Parameters for the LIF and AdExp neuron model and switching ON-OFF synaptic inputs.

Information measures

Many information measures have the caveats that they can only be applied to specific
situations with binary sensory input. We calculate four different information measures:

- mutual information MI between in- & output,

- transfer entropy TE between in- & output with one time bin as lag,

- coding efficiency of a spike train CE,

- tuning curve,

- orientation selectivity index OSI,

- coefficient of variation of neuronal activity CV_{ISI},

- coefficient of variation of subthreshold membrane variability CV_{V_m},

and relate them to the energy consumption.

The mutual information (MI) measures the overlap between the information content of
two (sub)systems Shannon (1948). The mutual information between a given stimulus s
and the neuronal response/firing rate r is denoted as $MI(r|s)$ (for continous time series
(Zeldenrust *et al.*, 2017)) and given by:

$$MI(r|s) = \sum_{r,s} p(s,r) \cdot \log\left(\frac{p(s,r)}{p(s) \cdot p(r)}\right),$$

where $p(s,r)$ is the joint probability describing the likelihood of observing a particular
combination of s and r concomitantly, $p(s)$ the stimulus probability of observing particular
stimulus s, and $p(r)$ the neuronal response probability of observing particular neuronal
response r. It is a symmetric measure where no distinction between cause and effect can
be made.

By introducing transition probabilities into mutual information dynamical and directional
information transport can be quantified. The resulting measures is called transfer entropy

(TE). This asymmetrical measure estimates causally transmitted information with a directed exchange of information to distinguish effectively driving and responding signal with specific time lag (Schreiber, 2000). The information flow from a given stimulus s to the neuronal response/firing rate r considering the the past stimulus s' with time lag τ is denoted as the transfer entropy $TE(r|s, \tau)$ and is given by:

$$TE(r|s, \tau) = \sum_{s, r, s'(\tau)} p(s, r, s'(\tau)) \cdot \log\left(\frac{p(s, r, s'(\tau)) \cdot p(s)}{p(s, r) \cdot p(s, s'(\tau))}\right),$$

where $p(s, r, s'(\tau))$ is the joint probability describing the likelihood of observing a particular combination of current stimulus s, neuronal response r, and past stimulus s' delayed by a time lag τ concomitantly, $p(s)$ is the stimulus probability of observing particular stimulus s, $p(s, r)$ is the joint probability describing the likelihood of observing a particular combination of s and r concomitantly without considering past stimuli, and $p(s, s'(\tau))$ is the joint probability of the current stimulus s and the past stimulus s' describing the relationship between the current and past states of a stimulus.

To estimate the mutual information and transfer entropy the spike trains were discretized in time and frequency space. The bin sizes were optimized by maximizing the information content for different combinations of time and frequency bins (optimal combinations: bin time$_{MI}$ = 1200 ms and bins rate$_{MI}$ = 150 and bin time$_{TE}$ = 294 ms and bins rate$_{TE}$ = 58. The time lag to estimate the transfer entropy was one bin size ($\tau = 294$ ms), see Fig. S5).

The coding efficiency of a spike train CE describes how much information is transmitted per spike. It is calculated by dividing the information content of a spike train (either measured by MI denoted as CE_{MI} or by TE denoted as CE_{TE}) by the number of spikes in the train represented by the firing rate r_{post}:

$$CE_{MI} = \frac{MI}{r_{\text{post}}}$$

$$CE_{TE} = \frac{TE}{r_{\text{post}}}$$

The tuning curve describes the response of the neuron to presented stimuli. The neuronal response is estimated by the firing rate of the neuron during each stimulus condition presentation.

The orientation selectivity index describes how selective a neuron responds to a stimulus. It compares the response of a neuron to its preferred orientation r_{pref} to its non-preferred (equivalently orthogonal, $90°$ shifted grating relative to preferred orientation) orientation response r_{ortho}:

$$OSI = \frac{r_{\text{pref}} - r_{\text{ortho}}}{r_{\text{pref}} + r_{\text{ortho}}}.$$

An OSI of 1 describes a selective neuron which has exclusive/strong response for preferred and no response to non-preferred stimuli. Conversely, an OSI of 0 describes a neuron responding to all stimuli equally making it non-selective for this group of stimuli.

The coefficient of variation of neuronal activity CV_{ISI} is a measure for variability of the inter-spike intervals of a spike train:

$$CV_{\text{ISI}} = \frac{\sigma_{\text{ISI}}}{\mu_{\text{ISI}}},$$

where σ_{ISI} is the standard deviation of inter-spike intervals and μ_{ISI} is the mean inters-spike interval duration.

The coefficient of variation of subthreshold membrane variability CV_{V_m} describes how variable the subthreshold membrane potential V_m is:

$$CV_{V_m} = \frac{\sigma_{V_m}}{\mu_{V_m}}.$$

Zeldenrust fraction of transferred information

To quantify the information transferred by the neuron for switching ON-OFF stimulation, we use the fraction of transferred information (FI) the ratio between the mutual infor-

mation between postsynaptic spike train ($MI_{\text{spike train}}$) and hidden state and the mutual information between synaptic input and hidden state (MI_{input}):

$$FI = \frac{MI_{\text{spike train}}}{MI_{\text{input}}}.$$

This quantifies the amount of information kept during the spike generation process.

The used information measures of our model assume rate-based coding, whereas real neurons likely employ a combination of rate, phase, and temporal coding strategies.

Energy budget

With their landmark study Attwell and Laughlin (2001) gave a simple estimate of the total energy of grey matter neurons as the sum of different subcellular processes. It was later updated by Howarth *et al.* (2012) and biologically validated by Yu *et al.* (2017) who compared expected and measured glucose consumption. Following the aforementioned studies, we assume the hydrolysis of ATP as the main energy provider and thus ATP as the main energy carrier ("energy currency of the cell"). For all energy budgets we calculate how much ATP a neuron consumes per second.

The total energy demand of neurons can be split into a non-signalling (house-keeping and maintaining a resting potential) and signalling component being activity/frequency dependent (action potential generation and conductance, synaptic transmission process, glutamate/GABA recycling, presynaptic Ca^{2+} entry). The total energy demand splits into six main subprocesses:

$$E_{tot} = E_{\text{HK}} + E_{\text{RP}} + E_{\text{AP}} + E_{\text{ST}} + E_{\text{glu}} + E_{Ca^{2+}}.$$

The different components can be estimated separately and vary due to the type and amount of molecules and ions involved and also depend on the cell morphology.

House keeping energy budget

Cells have a basic demand of energy to maintain their fundamental functions and shape such as synthesis and turnover of proteins, phospholipids and other macromolecules, microtubule cytoskeletal rearrangements and turnover, axoplasmic transport by actin tread milling, and mitochondrial proton leak (Engl and Attwell, 2015) (synaptogenesis (Balasubramanian, 2021)). These energy demands do not dependent on the activity of the neuron and thus are also consumed when the neuron is silent. All these house-keeping processes are summed in the term E_{HK} and represent how many hydrolyses of ATP molecules per second a neuron demands.

We chose to keep the house-keeping energy demand constant for all activity regimes following Yu *et al.* (2017). Attwell and Laughlin (2001) and Howarth *et al.* (2012) calculated it as 1/3 of the signalling energy and others report 1/5 (Hyder *et al.*, 2013). The effect of this assumption increases with firing rate. Because our results suggest medium-low firing-rates being the energetically most efficient ones, we argue that the effect of this assumption has only minor effect on the results. But for investigating neurons with high firing rates the activity-dependence of the house-keeping energy demand should be considered further. Furthermore, Balasubramanian (2021) report that 10 % of total energy remains unused under normal operation and serves as energy reserviour to buffer periods of high energetic demands.

Resting potential energy demand

To maintain the resting potential, a state maintained far from thermodynamic equilibrium, a neuron continuously needs to pump sodium out of the cell (and potassium into the cell) due to the leakiness of the cellular membrane to ions. Pumping sodium against the electrical (charge difference) and chemical (concentration difference) gradient involves the sodium-potassium ion pump which extrudes 3 Na^+ and imports 2 K^+ under the consumption of 1 molecule of ATP. Assuming that the cell membrane at rest is (only) permeable to Na^+ and K^+ and all charge fluxes through these leaky conductances are compensated by the sodium-potassium pump with given stoichiometry and ATP demand (in opposite direction) (no net charge flux, balanced sodium and potassium conductances), Attwell and Laughlin (2001) derive Formula (4) in their paper to estimate the energy demand E_{RP} to maintain the membrane at a specific resting potential V_{RP}:

$$E_{\mathrm{RP}}(V_{\mathrm{RP}}) = \frac{(V_{\mathrm{Na}^+} - V_{\mathrm{RP}})(V_{\mathrm{RP}} - V_{\mathrm{K}^+})}{eR_m(V_{\mathrm{RP}} + 2V_{\mathrm{Na}^+} - 3V_{\mathrm{K}^+})},$$

where V_{Na^+} is the sodium reversal/Nernst potential, V_{K^+} the potassium reversal/Nernst potential, e the elementary charge, and R_m the membrane resistance. The somatic estimate of input resistance derived from our experimental data slightly overestimates the true membrane resistance relevant for calculating the resting potential energy, by approximately 9%. To assess the overestimation of membrane resistance from somatic input resistance measurements, we performed simulations using a detailed compartmental model of a layer 2/3 pyramidal neuron (Goetz *et al.*, 2021). A current of 0.05 nA was injected into the soma, resulting in a voltage deflection of 7.331 mV and an input resistance (R_{in}) of 146.62 MΩ. Given the model's total surface area ($A_{\mathrm{neuron}} = 0.000111$ cm^2), this corresponded to an estimated membrane resistance ($R_m = R_{\mathrm{in}} \cdot A_{\mathrm{neuron}}$) of 16.3 kΩ·cm^2. In contrast, the model's true specific membrane resistance was 15 kΩ·cm^2, indicating that somatic input resistance overestimates the true membrane resistance by approximately 9%. This bias is likely even larger for the larger neurons considered by Attwell and Laughlin (2001). We therefore applied a correction factor based on this estimate when calculating energy budgets from experimental measurements.

We are aware, that the above equation to estimate the resting energy represents a conservative estimate of energy savings as drastically changing the resting potential violates some of the original assumptions made by Attwell and Laughlin (2001). In particular, this equation assumes that the resting potential is governed by the ratio of just one conductance for each of Na$^+$ and K$^+$. A higher resting potential therefore implies a higher Na$^+$ conductance which actually increases ATP cost (Fig S8). In reality, the membrane potential is maintained by a larger number of ionic conductances. Hyperpolarization-activated cyclic nucleotide-gated (HCN) channels depolarize the resting potential, contribute to resting potential stability and dendritic integration (Wahl-Schott and Biel, 2008). By incorporating HCN-mediated currents into the energy budget, we refine estimates of resting potential maintenance. The somato-dendritic gradient of HCN channels also influences temporal integration, potentially tuning synaptic efficacies and response selectivity (Zhuchkova *et al.*, 2013). The I_h might also be adapted during food-restriction helping to save energy (more degrees of freedom). Thus, for our energy calculations we extent the original formula

derived by (Attwell and Laughlin, 2001) by an additional I_h resulting in

$$E_{\text{RP}} = \frac{g_{\text{Na}} \cdot (V_{\text{Na}} - V_{\text{RP}})/3 + g_{\text{h}} \cdot (V_{\text{h}} - V_{\text{RP}})/4}{e}.$$

The full derivation with the basic assumptions is presented in the supplements.

Action potential energy budget

Firing an action potential and propagating it to along the whole axon (of a neuron to the dendritic terminals) involves a depolarization (sodium influx, rising phase) followed by a repolarization or even a hyperpolarization (potassium efflux, falling phase) of the neuronal membrane. The re- and hyperpolarization do not happen spontaneously because intracellular sodium has to be extruded against the electrochemical gradient. Thus, the energy demand of an action potential can be calculated by the total sodium charge influx into the cell during depolarization (due to an action potential being fired), which has to be extruded by the sodium-potassium pump (with stoichiometry 3 Na$^+$ per ATP) during repolarization under the consumption of ATP (Attwell and Laughlin, 2001):

$$E_{\text{AP}}(r_{\text{post}}) = \frac{\Delta Q_{\text{Na}^+}}{3 \cdot e} \cdot r_{\text{post}} \quad \text{with} \quad \Delta Q_{\text{Na}^+} = A_{\text{overlap}} \cdot S_m \cdot c_m \cdot \Delta V,$$

where E_{AP} is the energy demand for action potential firing at a mean firing rate r_{post}, ΔQ_{Na^+} is the sodium charge influx which has to be extruded by the sodium-potassium pump, e is the elementary charge, A_{overlap} is the overlap factor between sodium and potassium currents, S_m is the surface area of the neuron, c_m is the specific membrane capacitance, and ΔV being the difference between the spike peak voltage V_{peak} and the threshold voltage V_{th}:

$$\Delta V = (V_{\text{peak}} - V_{\text{th}}).$$

The surface area of a neuron is estimated as the sum of the surface area of the axon, soma, and the dendritic tree (Attwell and Laughlin, 2001):

$$S_m = S_{\text{axon}} + S_{\text{soma}} + S_{\text{dendrite}}.$$

The surface area of the axons is calculated by: 818

$$S_{\text{axon}} = \pi \cdot d_{\text{axon}} \cdot l_{\text{axon}}.$$

The surface area of the soma is calculated by: 819

$$S_{\text{soma}} = \pi \cdot d_{\text{soma}}^2.$$

The surface area of the dendritic tree is calculated by: 820

$$S_{\text{dendrite}} = \pi \cdot d_{\text{dendrite}} \cdot l_{\text{dendrite}}.$$

The theoretical minimum charge needed to depolarize the membrane from V_{th} to V_{peak} 821
(amplitude of the action potential) is $\Delta Q_{\text{Na}^+,\text{min}} = S_m \cdot c_m \cdot \Delta V$. In biological neurons, 822
however, sodium influx and potassium efflux overlap temporally due to channel kinetics, 823
increasing the effective sodium charge influx (Sengupta *et al.*, 2010). This excess cost by 824
increased sodium charge influx is commonly described by the overlap ratio A_{overlap} and 825
describes the energy-efficiency of action potentials. 826
Attwell and Laughlin (2001) calculated the contribution of action potentials to their 827
energy budget based on measurements from the squid giant axon, where temporal 828
sodium–potassium current overlap results in an energy expenditure approximately four 829
times the theoretical minimum (Hodgkin and Huxley, 1952; Hodgkin, 1975) However, the 830
squid giant axon is an energetically inefficient reflex neuron compared to most mammalian 831
neurons. Alle *et al.* (2009) (for rat hippocampal non-myelinated mossy fibers) and Carter 832
and Bean (2009) (for mouse pyramidal cells in hippocampal CA1 and neocortex) showed 833
that action potentials in mammalian neurons are much more energy-efficient by measuring 834
the overlap between the rapidly decaying sodium current and delayed-onset potassium 835
current waveforms. The authors showed that action potentials in mammalian excitatory 836

axons and somata cost only 1.3 instead of 4 times the theoretical minimum. These findings led to a revised and lower estimate of spike-related energy consumption in mammalian cortex (Howarth *et al.*, 2012).

Carter and Bean (2009) also provided experimental evidence that the amount of sodium entry during action potential generation varies considerably among different species and cell types (Carter and Bean (2009), Fig. 4). An increased sodium influx due to an incomplete sodium inactivation decreases energy efficiency but enables fast spiking (narrower APs) as a functional benefit (Carter and Bean, 2009). Thus, it is important to consider the type of neurons because they trade-off energy efficiency and function differently.

Alternative approaches for estimating the energetic cost of action potential generation have also been proposed (Yi *et al.*, 2016). Rather than quantifying ion flux directly, these methods infer energy demand from the electrical power dissipated by ionic conductances during spiking, offering a complementary perspective to ion-count-based estimates.

It is important to note that the membrane capacitance C_m reported in Padamsey *et al.* (2022) reflects the somatic capacitance only. While this value is appropriate for simulating integrative properties in single-compartment neuron models (LIF or AdExp), it underestimates the total energy cost of action potential generation, which also involves the axon and dendritic arbor. To accurately capture the full energetic demand, we compute the total membrane capacitance by multiplying the specific membrane capacitance (c_m) with the complete surface area (S_m) of a pyramidal neuron.

We additionally assessed the energetic contribution of adaptation currents. The mean adaptation current increases with firing rate and with the adaptation time constant (Fig. S7A). Within the firing-rate range relevant for optimal coding in our study (up to ~5 Hz), adaptation currents contribute no more than 13 % of the mean excitatory synaptic current (Fig. S7B). To examine the ionic composition and energetic cost of adaptation currents during spike generation, we stimulated a detailed compartmental model (Goetz *et al.*, 2021) with brief step current injections (15 ms duration), once with adaptation currents (I_M and $I_{K,\mathrm{Ca}}$) enabled and once with these currents disabled (Fig. S7C). Tracking ionic fluxes revealed that the presence of adaptation currents slightly reduces the total ionic load per action potential, resulting in a lower energetic cost per spike (Fig.S7D). Given their

minor contribution in the low firing-rate regime studied here, adaptation currents were 867
not explicitly included in the energy budget. However, their energetic impact may become 868
relevant at higher firing rates and under stronger adaptation. 869

Synaptic transmission energy budget 870
The release of neurotransmitter from the presynaptic terminal and their binding to post- 871
synaptic (ionotropic) receptors results in an influx of Na^+ and Ca^{2+}. We calculated the 872
energy demand of the postsynaptic cell to re-establish the resting ion gradients due to 873
presynaptic activity as: 874

$$E_{\text{ST}} = \frac{\Delta Q_{\text{Na}^+}}{3 \cdot e} \quad \text{with} \quad \Delta Q_{\text{Na}^+} = I_{e,tot} \cdot \Delta t \cdot A_{\text{syn}},$$

where E_{ST} is the energy demand to restore the resting membrane potential due to the 875
"effective" (post)synaptic sodium charge influx ΔQ_{Na^+} which has to be extruded by the 876
sodium-potassium pump. The effective sodium charge influx ΔQ_{Na^+} is the product of 877
the total excitatory synaptic current $I_{e,tot}$ entering the cell in time interval Δt (energy 878
measures per second demands $\Delta t = 1$ s), and the proportionality factor A_{syn} which 879
accounts for the different stoichiometry of sodium and calcium (potassium) influx as well 880
as inhibitory chloride influxes (which have a much smaller electrochemical gradient at 881
inhibitory synapses and thus consume way less energy than sodium at excitatory synapses 882
(Attwell and Laughlin, 2001; Howarth *et al.*, 2012). Additionally, Attwell and Laughlin 883
(2001) treat all neurons as glutamatergic since they outnumber inhibitory neurons by 9:1 884
and 90 % of synapses release glutamate (Braitenberg and Schüz, 1998). 885

We derive A_{syn} as follows: 886

Assuming that the (measured/simulated) excitatory synaptic current consists only of 887
sodium ions the energy demand to reverse the influx of sodium charge $\Delta Q_{\text{Na}^+} = I_{e,tot} \cdot \Delta t$ 888
(per second $\Delta t = 1$ s) is: 889

$$E_{ST,\text{Na}^+\text{approx}} = \frac{I_{e,tot}}{3 \cdot e}.$$

To obtain biophysically grounded estimates of synaptic ionic currents, we leveraged a   890
detailed multi-compartment model of a layer 2/3 pyramidal neuron (Goetz *et al.*, 2021).   891
Although this model was originally parameterized for rat cortex, comparative analyses   892
across mammalian species indicate that key electrophysiological and metabolic scaling   893
relationships are largely conserved, with humans representing the primary outlier in   894
allometric scaling (Beaulieu-Laroche *et al.*, 2021). We therefore use this model as a rep-   895
resentative mammalian reference to estimate the relative contributions of ionic currents   896
during synaptic transmission. Using this framework, we quantified the mean dendritic   897
ionic currents associated with excitatory synaptic input, yielding the following average   898
synaptic current components: 899

$$< I_{\text{Na}^+,\text{dend}} >= -0.0577 \, \text{nA},$$

$$< I_{\text{K}^+,\text{dend}} >= +0.0591 \, \text{nA},$$

$$< I_{\text{Ca}^{2+},\text{dend}} >= -0.0096 \, \text{nA}.$$

The ratio of Na^+- to Ca^{2+}-currents is approximately 6:1. Thus, the Ca^{2+}-current accounts   900
for $1/7$ of the total current ($I_{e,tot}$). Splitting the excitatory synaptic current into its relative   901
contributions $I_{e,tot} = I_{\text{Na}^+} + I_{\text{Ca}^{2+}}$:   902

$$I_{\text{Na}^+} = \frac{6}{7} \cdot I_{e,tot},$$

$$I_{\text{Ca}^{2+}} = \frac{1}{7} \cdot I_{e,tot}.$$

The ATP-demand to extrude Ca^{2+} is much different to extrude Na^+. One molecule of   903
ATP extrudes 3 Na^+ or 1 Ca^{2+}. Accounting for the double charge of Ca^{2+} and the ATP-   904
stoichiometry the energy consumption of the contributions are: 905

$$E_{ST,\text{Na}^+} = \frac{\frac{6}{7} \cdot I_{e,tot}}{3 \cdot e} = \frac{6 \cdot I_{e,tot}}{21 \cdot e},$$

$$E_{ST,\text{Ca}^{2+}} = \frac{\frac{1}{7} \cdot I_{e,tot}}{2 \cdot e} = \frac{1 \cdot I_{e,tot}}{14 \cdot e}.$$

In sum this results in:

$$E_{ST,\text{Na}^+\text{Ca}^{2+}} = E_{ST,\text{Na}^+} + E_{ST,\text{Ca}^{2+}} = \frac{6 \cdot I_{e,tot}}{21 \cdot e} + \frac{1 \cdot I_{e,tot}}{14 \cdot e} = \frac{5}{14} \cdot \frac{I_{e,tot}}{e}.$$

Setting this measure in relation to $E_{ST,\text{Na}^+\text{approx}}$ we receive a factor A_{syn} which corrects the energy estimate of the non-specific excitatory synaptic current approximation (of the excitatory synaptic currents only consisting of Na^+) by accounting for different stoichiometries of Na^+ and Ca^{2+}:

$$A_{\text{syn}} = \frac{E_{ST,\text{Na}^+\text{Ca}^{2+}}}{E_{ST,\text{Na}^+\text{approx}}} = \frac{\frac{5}{14}}{\frac{1}{3}} \approx 1.07.$$

The mean ionic currents of the compartmental model vary for different stimuli presentations in a range from 2:11 for preferred till 1:8 for non-preferred. Calculating A_{syn} analogously to before these ratios results in a range of A_{syn} of 1.055 for non-preferred until 1.08 for preferred stimuli, thus, not varying a lot. To account also for energetic demands of inhibitory synapses we set this correction factor to $A_{\text{syn}} \approx 1.1$.

Attwell and Laughlin (2001) already mention the estimation of synaptic transmission energy from postsynaptic ion fluxes but also that few experimental data are available. But the postsynaptic ionic currents are easily accessible (inate output variable) of our neuronal modeling. Attwell and Laughlin (2001) calculated the energy budget of the synaptic transmission process by the presynaptic activity:

$$E_{\text{ST}}(r_{\text{post}}) = r_{\text{post}} \cdot p_{\text{ves}} \cdot N_{boutons} \cdot E_{\text{ves}},$$

where p_{ves} is the release probability for glutamate vesicles, $N_{boutons}$ the number of downstream synaptic contacts to following cells, and E_{ves} is the energy needed to reverse the sodium ion influx through NMDA and non-NMDA ionotropic receptors and metabotropic receptors (G-protein signalling) due to one vesicle of glutamate of around 137 000 ATP molecules per vesicle (Attwell and Laughlin, 2001). This calculation assumes a linear increase of synaptic transmission energy demand with the firing rate and that that the

pre- and postsynaptic firing rates always behave equally. This assumption is based on the   927
idea that the release of presynaptic glutamate results in the opening of an equal amount of   928
postsynaptic receptors and thus constant postsynaptic influx of ions and metabolic con-   929
sequences. But this assumption of symmetry in pre- and postsynaptic firing rate/energy   930
use neglects the postsynaptic non-linear integration process and does not capture synaptic   931
scaling. These estimations of ATP consumption due to synaptic transmission make the   932
simplifying assumption used in Attwell and Laughlin (2001) that neurons in a population   933
are recurrently connected so that pre- and post-synaptic firing rates are equal. For sensory   934
systems with a more hierarchical organisation (Van Essen and Maunsell, 1983), it is likely   935
that these two rates will differ, so we relax this assumption here.   936

Neurotransmitter recycling energy budget   937

Excitatory synaptic signalling involves mainly the presynaptic release of glutamate and its   938
binding to glutamate receptors at the postsynaptic cell. The released glutamate is resumed   939
by the presynaptic cell, metabolically recycled/reprocessed, repacked in vesicles, and   940
transported to the dendritic terminals for further signalling (Li and Sheng, 2022). We   941
calculated the energy demand of the cell for neurotransmitter recycling (mainly glutamate)   942
E_{glu} as: 943

$$E_{\text{glu}}(r_{\text{post}}) = r_{\text{post}} \cdot p_{\text{ves}}(r_{\text{post}}) \cdot N_{boutons} \cdot E_{\text{glu,rec}} \quad \text{with} \quad p_{\text{ves}}(r_{\text{post}}) = 0.6 \cdot e^{\frac{-r_{\text{post}}}{5.0}},$$

where p_{ves} is the release probability for glutamate vesicles depending on the neuronal firing   944
rate r_{post}, $N_{boutons}$ the number of downstream synaptic contacts/boutons to neurons, and   945
$E_{\text{glu,rec}}$ the energy to recycle one vesicle of glutamate. One vesicle contains approximately   946
4000 glutamate molecules (Riveros *et al.*, 1986) and to recycle one glutamate molecule   947
3.67 ATP molecules are needed (Yu *et al.*, 2017). The presynaptic release probability   948
p_{ves} is assumed to decay exponentially as a function of presynaptic firing rate (r_{post})   949
Attwell and Laughlin (2001); Howarth *et al.* (2012); Yu *et al.* (2017) due to the depletion of   950
neurotransmitter during prolonged stimulation (Del Castillo and Katz, 1954). The steady-   951
state release probability is an exponentially decreasing function of rate (Tsodyks and   952
Markram, 1997) for a broad class of presynaptic firing patterns (Bird and Richardson, 2018).   953
Attwell and Laughlin (2001); Howarth *et al.* (2012) used the simplification of a constant   954
value of $p_{\text{ves}} = 0.25$ to estimate the release probability for a mean activity of $r_{\text{post}} = 4$ 955

Hz. But our simulations involve a broad range of different firing rates. Thus, we account for the (quick) exponential decay of presynaptic release probability with a time constant of the synaptic depression of 200 ms (Tsodyks *et al.*, 1998). Moreover, the presynaptic release probability has been found to adapt to energy levels by regulating the speed of synaptic vesicle cycle (Yuan *et al.*, 2018). This is in line with findings of activity-enhanced presynaptic energy production (Li and Sheng, 2022).

Presynaptic calcium extrusion energy budget

The synaptic transmission process includes an influx of calcium which triggers, if enough calcium accumulates in a bouton, the release of a neurotransmitter vesicle. This intracellular calcium has to be extruded against the electrochemical gradient, which consumes energy. Another consumer of energy is the vesicular release (endocytosis) in form of a fusion of vesicle and cell membrane. Similar to the glutamate release energy budget, also the cost for calcium extrusion is activity-dependent. Analogously, we calculated the energy demand of the cell for calcium extrusion $E_{\text{Ca}^{2+}}$ as:

$$E_{\text{Ca}^{2+}}(r_{\text{post}}) = r_{\text{post}} \cdot p_{\text{ves}}(r_{\text{post}}) \cdot N_{boutons} \cdot E_{\text{Ca}^{2+},\text{ex}} \quad \text{with} \quad p_{\text{ves}}(r_{\text{post}}) = 0.6 \cdot e^{\frac{-r_{\text{post}}}{5.0}},$$

where $E_{\text{Ca}^{2+},\text{ex}}$ is the energy needed to extrude the calcium influx of a single presynaptic action potential to release one neurotransmitter vesicle. For the release of one vesicle $12\,000$ Ca^{2+} ions are needed and it takes 1 ATP molecule to extrude 1 Ca^{2+} ion (Attwell and Laughlin, 2001).

Additional minor presynaptic energy contributors are synapse assembly, their main-tainance by cargo transport (Li and Sheng, 2022).

Minor energy consumption of inhibitory currents

Neuronal energy consumption depends on morphology (size of membrane area), synaptic connectivity, the precise combination and densities of ionic conductances, their type, function and neurotransmitter. While we focused on Na^+, K^+, and Ca^{2+} gradients, other ions (H^+, Cl^-, and HCO_3^-) also contribute to energy costs (Attwell and Laughlin, 2001). Inhibition, while metabolically less demanding and often neglected (around 1 % of the total energy demand (Attwell and Laughlin, 2001; Howarth *et al.*, 2012)), plays a critical role in

network function (Buzsaki *et al.*, 2007; Raimondo *et al.*, 2017) and was incorporated into 983
our model through a synaptic proportionality factor A_{syn} derived from a compartmental 984
model (Goetz *et al.*, 2021). 985

Minor energy consumption of adaptation currents 986

In addition, the energy cost of adaptation currents may merit consideration. As shown 987
by Yi *et al.* (2016), spike-frequency adaptation mediated by Ca^{2+}-activated (I_{AHP}) and 988
K^{+}-activated (I_{M}) currents nearly doubles the energy cost of action potentials under strong 989
adaptation. These slow hyperpolarizing K^{+} currents increase the sodium influx required 990
to reach threshold, thereby enhancing the overlap between inward sodium and outward 991
potassium currents and reducing action potential efficiency. To account for this additional 992
cost, it may be necessary to introduce either a frequency-dependent adjustment of the 993
overlap factor (A_{overlap}) or an explicit energy term linked to the adaptation current (w_{ad}), 994
which remains to be quantified. In our simulations, the mean adaptation current increased 995
with firing rate (Fig. S7A) but remained below 14 % of the mean excitatory current across 996
physiologically relevant firing rates (Fig. S7B). Interestingly, disabling adaptation currents 997
(I_{M} and $I_{\text{K,Ca}}$) in a detailed compartmental model (Goetz *et al.*, 2021) increased the energetic 998
cost of a single spike by 5%, due to greater sodium and calcium influx(Fig. S7C-D). 999

In total, the reversal of ion-fluxes consumes the largest amount of energy with the synaptic 1000
transmission processes being the main contributor. 1001

Parameter	Value	Unit	Reference
E_{HK}	$5.38 \cdot 10^8$	ATP/s	Yu *et al.* (2017)
V_{Na^+}	50	mV	Attwell and Laughlin (2001)
V_{K^+}	-100	mV	Attwell and Laughlin (2001)
e	$1.6 \cdot 10^{-19}$	C	-
A_{overlap}	1.24	-	Carter and Bean (2009)
d_{axon}	0.3	μm	Braitenberg and Schüz (1998)
l_{axon}	4	cm	Braitenberg and Schüz (1998)
d_{soma}	25	μm	Attwell and Laughlin (2001)
d_{dendrite}	$3 \cdot d_{\text{axon}}$	μm	Braitenberg and Schüz (1998)
l_{dendrite}	$l_{\text{axon}}/9$	cm	Braitenberg and Schüz (1998)
c_m	0.01	F/m^2	Attwell and Laughlin (2001)
V_{peak}	25	mV	Carter and Bean (2009)
V_{th}	-50	mV	Zerlaut and Destexhe (2017)
A_{syn}	1.1	-	
$N_{boutons}$	8000	-	Braitenberg and Schüz (1998)
$E_{\text{glu,rec}}$	$1.468 \cdot 10^4$	ATP/s	Yu *et al.* (2017)
$E_{\text{Ca}^{2+},\text{ex}}$	$1.2 \cdot 10^4$	ATP/s	Attwell and Laughlin (2001)

Table 3: Neuronal parameters to calculate the energy budget of a layer 2/3 pyramidal excitatory neuron of rats with cellular anatomies, specific electrical properties, and natural constants. To calculate energy demands of human neurons simply multiply l_{axon} and $N_{boutons}$ by 1.89 to account for their larger size and different synaptic densities (higher metabolism) (Yu *et al.*, 2017).

References

Alle H, Roth A, Geiger JRP (2009) Energy-Efficient Action Potentials in Hippocampal Mossy Fibers. *Science* 325:1405–1408.

Altschuler SJ, Wu LF (2010) Cellular Heterogeneity: Do Differences Make a Difference? *Cell* 141:559–563.

Attwell D, Laughlin SB (2001) An Energy Budget for Signaling in the Grey Matter of the Brain. *Journal of Cerebral Blood Flow & Metabolism* 21:1133–1145.

Balasubramanian V (2021) Brain power. *Proceedings of the National Academy of Sciences* 118:e2107022118.

Barlow HB (1961) Possible principles underlying the transformations of sensory messages. *Sensory communication* 1:217–233.

Barral J, Reyes AD (2016) Synaptic scaling rule preserves excitatory–inhibitory balance and salient neuronal network dynamics. *Nature Neuroscience* 19:1690–1696.

Bazzigaluppi P, Amini AE, Weisspapir I, Stefanovic B, Carlen P (2017) Hungry Neurons: Metabolic Insights on Seizure Dynamics. *International Journal of Molecular Sciences* 18:2269.

Beaulieu-Laroche L, Brown NJ, Hansen M, Toloza EHS, Sharma J, Williams ZM, Frosch MP, Cosgrove GR, Cash SS, Harnett MT (2021) Allometric rules for mammalian cortical layer 5 neuron biophysics. *Nature* 600:274–278.

Bird AD, Richardson MJ (2018) Transmission of temporally correlated spike trains through synapses with short-term depression. *PLOS Computational Biology* 14:e1006232.

Braitenberg V, Schüz A (1998) *Cortex: statistics and geometry of neuronal connectivity* Springer Science & Business Media.

Brette R, Gerstner W (2005) Adaptive Exponential Integrate-and-Fire Model as an Effective Description of Neuronal Activity. *Journal of Neurophysiology* 94:3637–3642.

Brunel N, van Rossum MC (2007) Lapicque's 1907 paper: from frogs to integrate-and-fire. *Biological cybernetics* 97:337–339.

Buzsaki G, Kaila K, Raichle M (2007) Inhibition and Brain Work. *Neuron* 56:771–783.

Buzsaki G, Mizuseki K (2014) The log-dynamic brain: how skewed distributions affect network operations. *Nature Reviews Neuroscience* 15:264–278.

Carter BC, Bean BP (2009) Sodium Entry during Action Potentials of Mammalian Neurons: Incomplete Inactivation and Reduced Metabolic Efficiency in Fast-Spiking Neurons. *Neuron* 64:898–909. [1032] [1033] [1034]

Chen BL, Hall DH, Chklovskii DB (2006) Wiring optimization can relate neuronal structure and function. *Proceedings of the National Academy of Sciences* 103:4723–4728. [1035] [1036]

Chrysafides S M B SJ, S S (2024) Physiology, resting potential. *StatPearls* . [1037]

Dahmen D, Hutt A, Indiveri G, Kennedy A, Lefebvre J, Mazzucato L, Motter AE, Narayanan R, Payvand M, Planert H, Gast R (2025) How heterogeneity shapes dynamics and computation in the brain. *Neuron* . [1038] [1039] [1040]

Deistler M, Macke JH, Gonçalves PJ (2022) Energy-efficient network activity from disparate circuit parameters. *Proceedings of the National Academy of Sciences* 119:e2207632119. [1041] [1042]

Del Castillo J, Katz B (1954) Statistical factors involved in neuromuscular facilitation and depression. *The Journal of Physiology* 124:574. [1043] [1044]

Deneve S, Machens CK (2016) Efficient codes and balanced networks. *Nature Neuroscience* 19:375–382. [1045] [1046]

Desai NS, Rutherford LC, Turrigiano GG (1999) Plasticity in the intrinsic excitability of cortical pyramidal neurons. *Nature Neuroscience* 2:515–520. [1047] [1048]

Dimakou A, Pezzulo G, Zangrossi A, Corbetta M (2025) The predictive nature of spontaneous brain activity across scales and species. *Neuron* 113:1310–1332. [1049] [1050]

Dougherty KA, Islam T, Johnston D (2012) Intrinsic excitability of CA1 pyramidal neurones from the rat dorsal and ventral hippocampus. *The Journal of Physiology* 590:5707–5722. [1051] [1052]

Duarte R, Morrison A (2019) Leveraging heterogeneity for neural computation with fading memory in layer 2/3 cortical microcircuits. *PLoS Computational Biology* 15:e1006781. [1053] [1054]

Edelman GM, Gally JA (2001) Degeneracy and complexity in biological systems. *Proceedings of the National Academy of Sciences of the United States of America* 98:13763–8. [1055] [1056]

Engl E, Attwell D (2015) Non signalling energy use in the brain. *The Journal of Physiology* 593:3417–3429. [1057] [1058]

Gast R, Solla SA, Kennedy A (2024) Neural heterogeneity controls computations in spiking neural networks. *Proceedings of the National Academy of Sciences* 121:e2311885121.

Gauld OM, Packer AM, Russell LE, Dalgleish HW, Iuga M, Sacadura F, Roth A, Clark BA, Häusser M (2024) A latent pool of neurons silenced by sensory-evoked inhibition can be recruited to enhance perception. *Neuron* 112:2386–2403.e6.

Gerstner W, Kistler WM, Naud R, Paninski L (2014) *Neuronal Dynamics: From single neurons to networks and models of cognition* Cambridge University Press, Cambridge.

Giudice MD, Crespi BJ (2018) Basic functional trade-offs in cognition: An integrative framework. *Cognition* 179:56–70.

Goaillard JM, Marder E (2021) Ion Channel Degeneracy, Variability, and Covariation in Neuron and Circuit Resilience. *Annual Review of Neuroscience* 44:1–23.

Goetz L, Roth A, Häusser M (2021) Active dendrites enable strong but sparse inputs to determine orientation selectivity. *Proceedings of the National Academy of Sciences* 118:e2017339118.

Goldman M (2004) Enhancement of information transmission efficiency by synaptic failures. *Neural Computation* 16:1137–1162.

Greenberg DS, Nonnenmacher M, Macke JH (2019) Automatic posterior transformation for likelihood-free inference. *Proceedings of the 36th International Conference on Machine Learning (ICML)* pp. 2404–2414.

Haggard M, Chacron MJ (2023) Coding of object location by heterogeneous neural populations with spatially dependent correlations in weakly electric fish. *PLOS Computational Biology* 19:e1010938.

Haider B, Häusser M, Carandini M (2013) Inhibition dominates sensory responses in the awake cortex. *Nature* 493:97–100.

Harris JJ, Jolivet R, Engl E, Attwell D (2015) Energy-Efficient Information Transfer by Visual Pathway Synapses. *Current Biology* 25:3151–3160.

Harris JJ, Engl E, Attwell D, Jolivet RB (2019) Energy-efficient information transfer at thalamocortical synapses. *PLoS Computational Biology* 15:e1007226.

Harris J, Jolivet R, Attwell D (2012) Synaptic Energy Use and Supply. *Neuron* 75:762–777. 1087

Hilgert S, Hannah S, Niemeyer N, Huthmacher L, Hürkey S, Schleimer J, Schreiber S, Duch 1088
C, Ryglewski S (2025) Robustness through variability: ion channel isoform diversity 1089
safeguards neuronal excitability. *bioRxiv* p. 2025.09.16.676508. 1090

Hodgkin AL, Huxley AF (1952) A quantitative description of membrane current and its 1091
application to conduction and excitation in nerve. *The Journal of Physiology* 117:500–544. 1092

Hodgkin JA (1975) The optimum density of sodium channels in an unmyelinated nerve. 1093
Philosophical Transactions of the Royal Society of London. B, Biological Sciences 270:297–300. 1094

Howarth C, Gleeson P, Attwell D (2012) Updated Energy Budgets for Neural 1095
Computation in the Neocortex and Cerebellum. *Journal of Cerebral Blood Flow &* 1096
Metabolism 32:1222–1232. 1097

Hyder F, Rothman DL, Bennett MR (2013) Cortical energy demands of signaling and 1098
nonsignaling components in brain are conserved across mammalian species and activity 1099
levels. *Proceedings of the National Academy of Sciences* 110:3549–3554. 1100

Indiveri G (2025) Neuromorphic is dead. long live neuromorphic. *Neuron* 113:3311–3314. 1101

Isler K, Schaik CPv (2009) The Expensive Brain: A framework for explaining evolutionary 1102
changes in brain size. *Journal of Human Evolution* 57:392–400. 1103

Jedlicka P, Bird AD, Cuntz H (2022) Pareto optimality, economy–effectiveness trade-offs 1104
and ion channel degeneracy: improving population modelling for single neurons. *Open* 1105
Biology 12:220073. 1106

Joshi N, Burg SvD, Celikel T, Zeldenrust F (2025) Neuronal identity is not static: An 1107
input-driven perspective. *PLOS Computational Biology* 21:e1013821. 1108

Klemmensen MM, Borrowman SH, Pearce C, Pyles B, Chandra B (2024) Mitochondrial 1109
dysfunction in neurodegenerative disorders. *Neurotherapeutics* 21:e00292. 1110

Kumari S, Narayanan R (2026) Intrinsic excitability of rat hippocampal granule cells 1111
increases along the dorsal-to-ventral axis. *bioRxiv* p. 2026.01.01.697271. 1112

Köhler-Solís A, Schirmeier S (2025) Neuronal metabolism: Surprisingly flexible. *Current* 1113
Biology 35:R1098–R1101. 1114

Laughlin SB, Sejnowski TJ (2003)   Communication in Neuronal Networks.   *Sci-* 1115
ence 301:1870–1874. 1116

Le Masson G, Przedborski S, Abbott L (2014) A Computational Model of Motor Neuron 1117
Degeneration. *Neuron* 83:975–988. 1118

Lennie P (2003) The Cost of Cortical Computation. *Current Biology* 13:493–497. 1119

Levy WB, Baxter RA (1996) Energy Efficient Neural Codes. *Neural Computation* 8:531–543. 1120

Levy WB, Baxter RA (2002) Energy-Efficient Neuronal Computation via Quantal Synaptic 1121
Failures. *The Journal of Neuroscience* 22:4746–4755. 1122

Lezmy J, Harris JJ, Attwell D (2021) Optimising the energetic cost of the glutamatergic 1123
synapse. *Neuropharmacology* 197:108727. 1124

Li HL, van Rossum MC (2020) Energy efficient synaptic plasticity. *eLife* 9:e50804. 1125

Li S, Sheng ZH (2022) Energy matters: presynaptic metabolism and the maintenance of 1126
synaptic transmission. *Nature Reviews Neuroscience* 23:4–22. 1127

Lueckmann JM, Boelts J, Greenberg DS, Gonçalves PJ, Macke JH (2021)   Bench- 1128
marking simulation-based inference.   *Proceedings of the National Academy of Sci-* 1129
ences 118:e2103070118. 1130

Marder E, Goaillard JM (2006) Variability, compensation and homeostasis in neuron and 1131
network function. *Nature Reviews Neuroscience* 7:563–574. 1132

Marder E, Prinz AA (2002) Modeling stability in neuron and network function: the role of 1133
activity in homeostasis. *BioEssays* 24:1145–1154. 1134

McCormick DA, Pape HC (1990) Properties of a hyperpolarization-activated cation current 1135
and its role in rhythmic oscillation in thalamic relay neurones. *The Journal of Physiol-* 1136
ogy 431:291–318. 1137

Mejias JF, Longtin A (2012) Optimal heterogeneity for coding in spiking neural networks. 1138
Physical Review Letters 108:228102. 1139

Mishra P, Narayanan R (2021) Ion-channel degeneracy: Multiple ion channels heteroge- 1140
neously regulate intrinsic physiology of rat hippocampal granule cells. *Physiological* 1141
Reports 9:e14963. 1142

Navarrete A, Schaik CPv, Isler K (2011) Energetics and the evolution of human brain size. 1143
Nature 480:91–93. 1144

Niven JE, Laughlin SB (2008) Energy limitation as a selective pressure on the evolution of 1145
sensory systems. *Journal of Experimental Biology* 211:1792–1804. 1146

N'dri AW, Barbier T, Teulière C, Triesch J (2025) Predictive coding light. *Nature Communica-* 1147
tions 16:8880. 1148

O'Leary T, Williams AH, Caplan JS, Marder E (2013) Correlations in ion channel ex- 1149
pression emerge from homeostatic tuning rules. *Proceedings of the National Academy of* 1150
Sciences 110:E2645–E2654. 1151

Olshausen BA, Field DJ (1996) Emergence of simple-cell receptive field properties by 1152
learning a sparse code for natural images. *Nature* 381:607–609. 1153

Pache A, van Rossum MC (2023) Energetically efficient learning in neuronal networks. 1154
Current Opinion in Neurobiology 83:102779. 1155

Padamsey Z, Katsanevaki D, Dupuy N, Rochefort NL (2022) Neocortex saves energy by 1156
reducing coding precision during food scarcity. *Neuron* 110:280–296.e10. 1157

Padamsey Z, Rochefort NL (2023) Paying the brain's energy bill. *Current Opinion in* 1158
Neurobiology 78:102668. 1159

Padmanabhan K, Urban NN (2010) Intrinsic biophysical diversity decorrelates neuronal 1160
firing while increasing information content. *Nature Neuroscience* 13:1276–1282. 1161

Pallasdies F, Norton P, Schleimer JH, Schreiber S (2021) Neural optimization: Understand- 1162
ing trade-offs with Pareto theory. *Current Opinion in Neurobiology* 71:84–91. 1163

Perez-Nieves N, Leung VCH, Dragotti PL, Goodman DFM (2021) Neural heterogeneity promotes robust learning. *Nature Communications* 12:5791.

Prinz AA, Billimoria CP, Marder E (2003) Alternative to Hand-Tuning Conductance-Based Models: Construction and Analysis of Databases of Model Neurons. *Journal of Neurophysiology* 90:3998–4015.

Prinz AA, Bucher D, Marder E (2004) Similar network activity from disparate circuit parameters. *Nature Neuroscience* 7:1345–1352.

Raimondo JV, Richards BA, Woodin MA (2017) Neuronal chloride and excitability — the big impact of small changes. *Current Opinion in Neurobiology* 43:35–42.

Remme MWH, Rinzel J, Schreiber S (2018) Function and energy consumption constrain neuronal biophysics in a canonical computation: Coincidence detection. *PLoS Computational Biology* 14:e1006612.

Rich S, Chameh HM, Lefebvre J, Valiante TA (2022) Loss of neuronal heterogeneity in epileptogenic human tissue impairs network resilience to sudden changes in synchrony. *Cell Reports* 39:110863.

Riveros N, Fiedler J, Lagos N, Mun~oz C, Orrego F (1986) Glutamate in rat brain cortex synaptic vesicles: influence of the vesicle isolation procedure. *Brain Research* 386:405–408.

Savtchenko LP, Sylantyev S, Rusakov DA (2013) Central synapses release a resource-efficient amount of glutamate. *Nature Neuroscience* 16:10–12.

Schneider M, Bird AD, Gidon A, Triesch J, Jedlicka P, Cuntz H (2023) Biological complexity facilitates tuning of the neuronal parameter space. *PLoS computational biology* 19:e1011212.

Schreiber S, Machens CK, Herz AVM, Laughlin SB (2002) Energy-Efficient Coding with Discrete Stochastic Events. *Neural Computation* 14:1323–1346.

Schreiber T (2000) Measuring Information Transfer. *Physical Review Letters* 85:461–464.

Sengupta B, Laughlin SB, Niven JE (2013) Balanced Excitatory and Inhibitory Synaptic Currents Promote Efficient Coding and Metabolic Efficiency. *PLoS Computational Biology* 9:e1003263.

Sengupta B, Stemmler M, Laughlin SB, Niven JE (2010) Action Potential Energy Efficiency Varies Among Neuron Types in Vertebrates and Invertebrates. *PLoS Computational Biology* 6:e1000840.

Shannon CE (1948) A Mathematical Theory of Communication. *Bell System Technical Journal* 27:379–423.

Shen G, Zhao D, Dong Y, Li Y, Zeng Y (2023) Dive into the Power of Neuronal Heterogeneity. *arXiv* .

Shoval O, Sheftel H, Shinar G, Hart Y, Ramote O, Mayo A, Dekel E, Kavanagh K, Alon U (2012) Evolutionary trade-offs, pareto optimality, and the geometry of phenotype space. *Science* 336:1157–1160.

Sihn D, Kim SP (2019) A spike train distance robust to firing rate changes based on the earth mover's distance. *Frontiers in Computational Neuroscience* 13:82.

Song S, Sjöström PJ, Reigl M, Nelson S, Chklovskii DB (2005) Highly Nonrandom Features of Synaptic Connectivity in Local Cortical Circuits. *PLoS Biology* 3:e68.

Stöber TM, Batulin D, Triesch J, Narayanan R, Jedlicka P (2023) Degeneracy in epilepsy: multiple routes to hyperexcitable brain circuits and their repair. *Communications Biology* 6:479.

Tejero-Cantero A, Boelts J, Deistler M, Lueckmann JM, Durkan C, Gonçalves P, Greenberg D, Macke J (2020) sbi: A toolkit for simulation-based inference. *Journal of Open Source Software* 5:2505.

Tsodyks M, Pawelzik K, Markram H (1998) Neural Networks with Dynamic Synapses. *Neural Computation* 10:821–835.

Tsodyks MV, Markram H (1997) The neural code between neocortical pyramidal neurons depends on neurotransmitter release probability. *Proceedings of the National Academy of Sciences* 94:719–723.

Turrigiano G, Abbott LF, Marder E (1994) Activity-dependent changes in the intrinsic properties of cultured neurons. *Science* 264:974–977.

Turrigiano GG, Leslie KR, Desai NS, Rutherford LC, Nelson SB (1998) Activity-dependent scaling of quantal amplitude in neocortical neurons. *Nature* 391:892–896. 1219 1220

Turrigiano GG, Nelson SB (2004) Homeostatic plasticity in the developing nervous system. *Nature Reviews Neuroscience* 5:97–107. 1221 1222

Van Essen DC, Maunsell JH (1983) Hierarchical organization and functional streams in the visual cortex. *Trends in Neurosciences* 6:370–375. 1223 1224

Wahl-Schott C, Biel M (2008) HCN channels: Structure, cellular regulation and physiological function. *Cellular and Molecular Life Sciences* 66:470. 1225 1226

Yang J, Prescott SA (2023) Homeostatic regulation of neuronal function: importance of degeneracy and pleiotropy. *Frontiers in Cellular Neuroscience* 17:1184563. 1227 1228

Yi GS, Wang J, Li HY, Wei XL, Deng B (2016) Metabolic Energy of Action Potentials Modulated by Spike Frequency Adaptation. *Frontiers in Neuroscience* 10:534. 1229 1230

Yu L, Shen Z, Wang C, Yu Y (2018) Efficient Coding and Energy Efficiency Are Promoted by Balanced Excitatory and Inhibitory Synaptic Currents in Neuronal Network. *Frontiers in Cellular Neuroscience* 12:123. 1231 1232 1233

Yu L, Yu Y (2017) Energy efficient neural information processing in individual neurons and neuronal networks. *Journal of Neuroscience Research* 95:2253–2266. 1234 1235

Yu Y, Herman P, Rothman DL, Agarwal D, Hyder F (2017) Evaluating the gray and white matter energy budgets of human brain function. *Journal of Cerebral Blood Flow & Metabolism* 38:1339–1353. 1236 1237 1238

Yuan Y, Huo H, Fang T (2018) Effects of Metabolic Energy on Synaptic Transmission and Dendritic Integration in Pyramidal Neurons. *Frontiers in Computational Neuroscience* 12:79. 1239 1240

Zeldenrust F, Calcini N, Yan X, Bijlsma A, Celikel T (2024) The tuning of tuning: How adaptation influences single cell information transfer. *PLOS Computational Biology* 20:e1012043. 1241 1242 1243

Zeldenrust F, Gutkin B, Denéve S (2021) Efficient and robust coding in heterogeneous recurrent networks. *PLoS Computational Biology* 17:e1008673. 1244 1245

Zeldenrust F, Knecht Sd, Wadman WJ, Denève S, Gutkin B (2017) Estimating the Information Extracted by a Single Spiking Neuron from a Continuous Input Time Series. *Frontiers in Computational Neuroscience* 11:49.   1246 1247 1248

Zerlaut Y, Destexhe A (2017) A mean-field model for conductance-based networks of adaptive exponential integrate-and-fire neurons. *arXiv* .   1249 1250

Zhang M, Chitic R, Bohté SM (2025) Energy optimization induces predictive-coding properties in a multicompartment spiking neural network model. *PLoS Computational Biology* 21:e1013112.   1251 1252 1253

Zhuchkova E, Remme MWH, Schreiber S (2013) Somatic versus Dendritic Resonance: Differential Filtering of Inputs through Non-Uniform Distributions of Active Conductances. *PLoS ONE* 8:e78908.   1254 1255 1256

Supporting information

Frequently used abbreviations

- AdExp — Adaptive exponential integrate-and-fire neuron model

- ATP — Adenosine triphosphate

- CE — Coding efficiency

- CTR — Control condition; normally fed mice

- CV — Coefficient of variation

- FR — Food-restricted condition; mice subjected to 2 weeks of dietary restriction

- FWHM — Full width at half maximum

- HCN — Hyperpolarization-activated cyclic nucleotide-gated (HCN) channels

- ISI — Inter-spike interval

- LIF — Leaky integrate-and-fire neuron model

- mEPSC — Miniature excitatory postsynaptic current

- MI — Mutual information

- OSI — Orientation selectivity index

- SBI — simulation based inference

- TE — Transfer entropy

Figure S1: **Design of *in vivo*-like excitatory and inhibitory synaptic input to a layer 2/3 pyramidal neuron in primary visual cortex with orientation variation stimulus.**

A, Oriented gratings are present for 1.5 s with 1 s intermittent background firing (BG) and the orientation values 0°, 30°, 60°, 90°, 120°, 150°, 180° as done by Padamsey *et al.* (2022). **B – Top**, Excitatory synaptic weights follow a log-normal distribution ($\mu_{w_e} = 2.07, \sigma_{w_e} = 1.14$). **B – Bottom**, Excitatory presynaptic firing rates are normally distributed ($\mu_{r_e} = 1.47$ Hz, $\sigma_{r_e} = 0.23$ Hz). **C**, The seven orientations are assigned synaptic inputs based on the matching scheme, visualized as cityscape histograms: presynaptic firing rates (dark red) as a function of synaptic strength (green). The preferred stimulus (90°) elicits the strongest total synaptic drive, whereas non-preferred orientations (0°, 180°) exhibit the weakest. Intermediate orientations fall between these extremes. **D**, Raster plot of synaptic input spike times, distinguishing inhibitory (light blue), excitatory signal and excitatory noise (red as a function of the synaptic strength). **E – Top**, Cumulative density function of the experimentally observed mEPSC distribution as a proxy for synaptic strength in control cells (Song *et al.*, 2005). **E – Bottom**, The same for food-restricted cells, showing a disproportionate reduction in the strongest synapses under food restriction reduces the effect of the strongest synapses.

Broadness of tuning curve

The strong broadening of tuning curves under food restriction suggests that synaptic weights undergo non-multiplicative downscaling. Strong synapses are preferentially weakened, reducing overall energy expenditure but reducing the selectivity of the neuron. But, it is unclear if the *ex vivo* measured synaptic weight distribution represent the *in vivo* scaling behavior. The broadness of weight distribution affects the neuronal response and the width of tuning curve in opposite way: narrow weight distribution lead to broad tuning curves because only few synapses are coding for the preferred stimulus (Fig. 5A–C). The number of signaling synapses significantly influences the sharpness of tuning curves. In Fig. S2A, we show that a higher proportion of signaling synapses leads to a more pronounced tuning curve. This effect arises from the log-normal distribution of synaptic strengths in our input model. For the preferred orientation ($90°$), strong synapses are paired with high firing rates, resulting in a strong synaptic drive. When the fraction of signaling synapses increases, the contribution of strong synapses becomes more dominant relative to weaker ones, further enhancing the response to the preferred stimulus. The resulting tuning curves of the 50% signaling synapse ratio matched the experimental data best.

Figure S2: **A higher signaling synapse ratio increases tuning curve sharpness.**

Using an AdExp model with orientation selectivity task (Fig. S1) shows that tuning curves for different ratios of signaling synapses (**A**, 10% to **I**, 90%) show increased peakiness with a higher proportion of signaling synapses. This effect arises from the log-normal distribution of synaptic strengths: for the preferred orientation (90°), strong synapses are paired with high firing rates, leading to a stronger synaptic drive. As the proportion of signaling synapses increases, the relative contribution of strong synapses becomes more dominant, enhancing the selectivity of the neuronal response. The tuning curves with a excitatory signaling synapse ratio of 50% matched the experimental tuning curves (Padamsey *et al.*, 2022) best.

Adaptation and spike-generation non-linearities shape tuning curve broadening

At experimentally measured levels of intrinsic membrane noise, a generic leaky integrate-and-fire (LIF) neuron fails to reproduce the tuning curve broadening observed under food restriction (Fig. S3A) in simulations with oriented grating input (Fig. S1). Introducing an adaptation current does not qualitatively alter this behavior (Fig. S3B), indicating that adaptation alone is insufficient to account for the experimentally observed loss of selectivity.

In contrast, incorporating an exponential spike-initiation non-linearity produces a modest broadening of tuning curves (Fig. S3C). This effect is substantially amplified when adaptation is combined with non-linear spike initiation, as in the adaptive exponential integrate-and-fire (AdExp) model (Fig. S3D), which reproduces the pronounced broadening observed in food-restricted neurons well.

This behavior can be understood by phase-plane analysis. The exponential term introduces a non-linear voltage nullcline, reshaping the local dynamical landscape near threshold. In this regime, the separation between the voltage (V) and adaptation (w) nullclines provides a measure of dynamical robustness (Hilgert *et al.*, 2025): larger separations imply greater resistance to perturbations from membrane noise, synaptic fluctuations, or parameter variability. Control neurons exhibit a larger nullcline separation, conferring greater robustness and preserving selectivity, whereas food-restricted parameter configurations reduce this separation, rendering the system more susceptible to noise-driven transitions and thereby amplifying tuning curve broadening.

Figure S3: **Spike-generation non-linearities and adaptation amplify tuning curve broadening.**
A, A leaky integrate-and-fire (LIF) model with an orientation selectivity task (Fig. S1) does not exhibit tuning curve broadening at experimentally measured membrane noise levels. The corresponding phase plane contains only a linear voltage (V) nullcline. **B**, Adding a linear adaptation current to the LIF model does not induce tuning curve broadening; both the V- and w-nullclines remain linear. **C**, Introducing an exponential spike-initiation term produces a modest broadening of tuning curves. The phase plane now contains a non-linear V-nullcline, increasing sensitivity to fluctuations near threshold. **D**, The adaptive exponential integrate-and-fire (AdExp) model reproduces robust tuning curve broadening under experimental noise levels. The combination of a non-linear V-nullcline and a linear w-nullcline reduces the nullcline separation in food-restricted (FR) neurons relative to control (CTR) neurons, indicating decreased robustness to perturbations and explaining the stronger loss of selectivity in the FR condition.

Figure S4: **Correlation between firing rate and information measures.**

Correlation between mean firing rate and **A** coefficient of variation of subthreshold membrane voltage, **B** coefficient of variation of inter spike intervals, **C** mutual information, **D** transfer entropy, and **E** the mutual information based on the tuning curve only for AdExp model simulations with an orientation selectivity task (Fig. S1). **A – E** showing a similar relationship as previously reported by (Harris *et al.*, 2015). **F**, Orientation selectivity index increases, has a peak at small firing rates and slowly decays as firing rates increase.

Optimal binsize to maximize MI and TE

To estimate the mutual information and transfer entropy the spike trains were discretized in time and frequency space. The bin sizes were optimized by maximizing the information content for different combinations of time and frequency bins (optimal combinations: bin time$_{MI}$ = 1200 ms and bins rate$_{MI}$ = 150 and bin time$_{TE}$ = 294 ms and bins rate$_{TE}$ = 58. The time lag to estimate the transfer entropy was one bin size (τ = 294 ms), see Fig. S5).

Figure S5: **Optimization of the binsize for to calculate mutual information (MI) and transfer entropy (TE) in orientation selectivity tasks (Fig. S1)**
A, Different bin size combinations in time and frequency space were evaluated for mutual information calculation. The white dot marks the optimal discretization parameters: bin time$_{MI}$ = 1200 ms and bins rate$_{MI}$ = 150. **B**, The same procedure was applied for transfer entropy calculation, yielding optimal parameters of bin time$_{TE}$ = 782 ms and bins rate$_{TE}$ = 39, with the time lag of τ = 294 ms.

Derivation and rationale behind the energy budget calculation

Calculating proportionality factor A$_{syn}$

To refine the estimate of synaptic transmission energy costs, we computed the contributions of different ionic currents using a detailed compartmental model (Goetz *et al.*, 2021)

presented in Fig. S6. While previous models assumed that excitatory synaptic currents were dominated by sodium influx, our analysis revealed that calcium currents contribute significantly to the overall energy demand. By quantifying the ratio of sodium to calcium currents and incorporating their distinct ATP requirements for extrusion, we derived a correction factor, A_{syn}, to adjust the energy estimate. Across different stimulus conditions, A_{syn} remained relatively stable, ranging from 1.055 for non-preferred stimuli to 1.08 for preferred stimuli. To account for inhibitory synaptic energy demands, we set $A_{syn} \approx 1.1$, providing a more accurate estimate of the metabolic cost associated with synaptic transmission.

Figure S6: **Subthreshold ionic currents of the detailed compartmental model (Goetz et al., 2021).** **A**, Total synaptic ionic currents summed across all compartments of the detailed model (Goetz et al., 2021) over a 10 ms interval. The dominant contributors are the sodium current (I_{Na}) and potassium current (I_K), while the calcium current (I_{Ca}) provides a non-negligible contribution to the synaptic energy budget. **B**, Morphology of the detailed compartmental model (Goetz et al., 2021).

Figure S7: **Estimating the cost of adaptation currents.**

A, Mean adaptation current of the AdExp model with given parameters (Tab. 1) as a function of firing rate for four adaptation time constants (1, 50, 100, 500 ms). Higher adaptation time constants lead to larger mean adaptation currents across all firing rates. **B**, Ratio of mean adaptation current to mean excitatory synaptic current as a function of firing rate. For short adaptation time constants (1–100 ms), this ratio decreases with firing rate, while for a long time constant (500 ms), the ratio increases (indicating greater metabolic cost at higher activity levels). **C**, Membrane voltage traces from the detailed compartmental model (Goetz *et al.*, 2021) in response to a brief step current (100 ms onset, 15 ms duration). Simulations with and without adaptation currents (I_M and $I_{K,Ca}$) are compared. Without adaptation, the post-spike depolarization is elevated. **D**, Ionic currents (I_{Na^+}, I_{K^+}, $I_{Ca^{2+}}$) recorded following spike initiation (from 1.5 ms post-spike). Disabling adaptation currents increases net I_{Na^+} and $I_{Ca^{2+}}$ (also I_{K^+}) influx, thereby raising the energetic cost per spike.

Derivation of resting potential energy demand

In the following we perform the derivation of formula formula (4) in Attwell and Laughlin (2001) for the consumed energy E_{RP} (ATP per second) of a neuron as a function of the resting membrane potential V_{RP}, the reverse potentials V_{Na} and V_{K}, the Faraday constant F, and the input resistance R_m. The given relationships are:

I. Ionic currents at rest: $I_{pump} = g_{\text{Na}}(V_{\text{Na}} - V_{RP}) + g_{\text{K}}(V_{\text{K}} - V_{RP})$

II. Total consumption of ATP/time by Na^+/K^+-pump:
$I_{pump} = \frac{g_{\text{Na}}(V_{\text{Na}} - V_{RP})}{3}$

III. $\frac{1}{R_m} = g_{\text{K}} + g_{\text{Na}}$

IV. $E_{RP} = \frac{I_{pump}}{F}$

The steps of derivation are:

Step 1: Set the equations equal to each other
Equate the two expressions for I_{pump}:

$$g_{\text{Na}}(V_{\text{Na}} - V_{RP}) + g_{\text{K}}(V_{\text{K}} - V_{RP}) = \frac{g_{\text{Na}}(V_{\text{Na}} - V_{RP})}{3}$$

Step 2: Simplify the equation
Multiply both sides by 3 to eliminate the fraction:

$$3g_{\text{Na}}(V_{\text{Na}} - V_{RP}) + 3g_{\text{K}}(V_{\text{K}} - V_{RP}) = g_{\text{Na}}(V_{\text{Na}} - V_{RP})$$

Subtract $g_{\text{Na}}(V_{\text{Na}} - V_{RP})$ from both sides:

$$3g_{\text{K}}(V_{\text{K}} - V_{RP}) + 2g_{\text{Na}}(V_{\text{Na}} - V_{RP}) = 0$$

Step 3: Rearrange to isolate V_{RP}

Distribute the terms:

$$3g_K V_K - 3g_K V_{RP} + 2g_{Na} V_{Na} - 2g_{Na} V_{RP} = 0$$

Combine like terms:

$$3g_K V_K + 2g_{Na} V_{Na} = 3g_K V_{RP} + 2g_{Na} V_{RP}$$

Factor out V_{RP} on the right side:

$$3g_K V_K + 2g_{Na} V_{Na} = (3g_K + 2g_{Na}) V_{RP}$$

Solve for V_{RP}:

$$V_{RP} = \frac{3g_K V_K + 2g_{Na} V_{Na}}{3g_K + 2g_{Na}}$$

Thus, the resting membrane potential V_{RP} is given by this equation.

Step 4: Rearrange to isolate g_{Na} and g_K

Distribute the terms:

$$3g_K V_K - 3g_K V_{RP} + 2g_{Na} V_{Na} - 2g_{Na} V_{RP} = 0$$

Combine like terms:

$$3g_K V_K + 2g_{Na} V_{Na} = 3g_K V_{RP} + 2g_{Na} V_{RP}$$

Rearrange to separate g_{Na} and g_K:

$$2g_{Na}(V_{Na} - V_{RP}) = 3g_K(V_{RP} - V_K)$$

Solve for g_{Na}:

$$g_{Na} = \frac{3g_K(V_{RP} - V_K)}{2(V_{Na} - V_{RP})}$$

Step 5: Use the input resistance relationship to solve for g_K (III)

Using the relationship $\frac{1}{R_m} = g_{\text{Na}} + g_{\text{K}}$:

$$g_{\text{Na}} + g_{\text{K}} = \frac{1}{R_m}$$

Substitute the expression for g_{Na}:

$$\frac{3g_{\text{K}}(V_{RP} - V_{\text{K}})}{2(V_{\text{Na}} - V_{RP})} + g_{\text{K}} = \frac{1}{R_m}$$

Factor out g_{K}:

$$g_{\text{K}} \left(\frac{3(V_{RP} - V_{\text{K}})}{2(V_{\text{Na}} - V_{RP})} + 1 \right) = \frac{1}{R_m}$$

Combine terms inside the parentheses:

$$g_{\text{K}} \left(\frac{3(V_{RP} - V_{\text{K}}) + 2(V_{\text{Na}} - V_{RP})}{2(V_{\text{Na}} - V_{RP})} \right) = \frac{1}{R_m}$$

Simplify the numerator:

$$g_{\text{K}} \left(\frac{3V_{RP} - 3V_{\text{K}} + 2V_{\text{Na}} - 2V_{RP}}{2(V_{\text{Na}} - V_{RP})} \right) = \frac{1}{R_m}$$

$$g_{\text{K}} \left(\frac{V_{RP} + 2V_{\text{Na}} - 3V_{\text{K}}}{2(V_{\text{Na}} - V_{RP})} \right) = \frac{1}{R_m}$$

Solve for g_{K}:

$$g_{\text{K}} = \frac{2(V_{\text{Na}} - V_{RP})}{R_m(V_{RP} + 2V_{\text{Na}} - 3V_{\text{K}})}$$

Step 6: Substitute g_{K} back into the expression for g_{Na}

$$g_{\text{Na}} = \frac{3 \cdot \frac{2(V_{\text{Na}} - V_{RP})}{R_m(V_{RP} + 2V_{\text{Na}} - 3V_{\text{K}})} \cdot (V_{RP} - V_{\text{K}})}{2(V_{\text{Na}} - V_{RP})}$$

$$g_{\text{Na}} = \frac{3(V_{RP} - V_{\text{K}})}{R_m(V_{RP} + 2V_{\text{Na}} - 3V_{\text{K}})}$$

Step 7: Substitute I_{pump} into E_{RP} (II in IV)

From the given relationships:

$$E_{RP} = \frac{I_{pump}}{F} = \frac{\frac{g_{Na}(V_{Na}-V_{RP})}{3}}{F} = \frac{g_{Na}(V_{Na}-V_{RP})}{3F}$$

Step 8: Substitute g_{Na} in the energy consumption equation E_{RP}

$$E_{RP} = \frac{I_{pump}}{F} = \frac{g_{Na}(V_{Na}-V_{RP})}{3F} = \frac{\frac{3(V_{RP}-V_{K})}{R_m(V_{RP}+2V_{Na}-3V_{K})}(V_{Na}-V_{RP})}{3F}$$

Simplify:

$$E_{RP} = \frac{(V_{Na}-V_{RP})(V_{RP}-V_{K})}{FR_m(V_{RP}+2V_{Na}-3V_{K})}$$

and arrive at the final formula (4) given by (Attwell and Laughlin, 2001).

Alternative formulation for better understanding of the derivation:

$$E_{RP} = -\frac{(V_{Na}-V_{RP})(V_{K}-V_{RP})}{FR_m(2(V_{Na}-V_{RP})-3(V_{K}-V_{RP}))}$$

Estimating the cost of the resting potential with additional ionic conductances

Here we provide an informative derivation of the effect of including an inward-hyperpolarising (I_h) conductance, mediated by hyperpolarization-activated cyclic nucleotide–gated (HCN) channels, on the ATP cost of maintaining a given membrane resting potential. As neuronal membranes are maintained by a large number of conductances (Goaillard and Marder, 2021), this derivation could be extended further. The above assumptions are modified as followed:

1. Ionic currents at rest. The pump currents now consist of an additional term to account for the I_h, so that:

$I_{pump} = g_{Na}(V_{Na}-V_{RP}) + g_K(V_K-V_{RP}) + g_h(V_h-V_{RP})$

The reversal potential of the I_h current, V_h, is taken to be -43mV (McCormick and Pape, 1990).

2. As the I_h current is a mixed cation current typically assumed to consist of $\sim 3/4\,\text{Na}^+$ (McCormick and Pape, 1990), the total consumption of ATP/time by the Na^+/K^+-pump is:

$I_{pump} = \frac{g_{\text{Na}}(V_{\text{Na}} - V_{RP})}{3} + \frac{g_h(V_h - V_{RP})}{4}$

3. The membrane resistance now depends on the sum of the three conductances:

$\frac{1}{R_m} = g_{\text{K}} + g_{\text{Na}} + g_h$

4. As before:

$E_{RP} = \frac{I_{pump}}{F}$

Now, the above system is underconstrained in terms of ionic conductances. We introduce a parameter α to describe the ratio of g_{Na} to g_{K} so that $g_{\text{Na}} = \alpha g_{\text{K}}$. Typical estimates of α are in the region of 0.05 (Chrysafides S M and S, 2024). This relationship could change with the resting potential, but the Na^+ conductance will typically remain far lower than the K^+ conductance at rest. It is then possible to use the first two relationships above to write g_{K} in terms of g_h for a given resting potential:

$$g_{\text{K}} = \frac{9g_h(V_{RP} - V_{\text{Na}})}{12(V_{\text{K}} - V_{RP}) + 8\alpha(V_{\text{Na}} - V_{RP})}.$$

The membrane resistance relationship can then be used to write the three ionic conductances

$$g_{\text{Na}} = \frac{9\alpha}{R_m}\left(\frac{V_{RP} - V_h}{9(1+\alpha)(V_{RP} - V_h) + 12(V_{\text{K}} - V_{RP}) + 8\alpha(V_{\text{Na}} - V_{RP})}\right),$$

$$g_{\text{K}} = \frac{9}{R_m}\left(\frac{V_{RP} - V_h}{9(1+\alpha)(V_{RP} - V_h) + 12(V_{\text{K}} - V_{RP}) + 8\alpha(V_{\text{Na}} - V_{RP})}\right),$$

$$g_h = \frac{4}{R_m}\left(\frac{3(V_{\text{K}} - V_{RP}) + 2\alpha(V_{\text{Na}} - V_{RP})}{9(1+\alpha)(V_{RP} - V_h) + 12(V_{\text{K}} - V_{RP}) + 8\alpha(V_{\text{Na}} - V_{RP})}\right).$$

This results in the energy to maintain the resting potential:

$$E_{RP} = \frac{g_{\text{Na}} \cdot (V_{\text{Na}} - V_{RP})/3 + g_h \cdot (V_h - V_{RP})/4}{F}.$$

These conductances can be plotted individually (Fig. S8A) and show how, in contrast to the model with only two conductances, it is possible for sodium channels to close as the resting potential increases. This further implies that both increasing the membrane resistance R_m and the resting potential V_{RP} can save energy (Fig. S8B). The joint effect of the experimentally observed changes could therefore lead to a more substantial ATP saving given the extra degree of freedom provided by the HCN channels (Fig. S8C). In reality, many other channels are involved in maintaining the resting potential and we hypothesise that energy savings here could be even greater. Clarifying the exact mechanisms involved in these changes would require further experiments.

Figure S8: **Effect of considering additional ionic conductances of estimates of ATP consumption.**
A, Individual ionic conductances as a function of resting potential with (solid lines) and without (dashed lines) considering I_h. Input resistance is 92.3MΩ. **B**, ATP consumption necessary to maintain membrane properties as a function of resting potential (left, $R_m = 92.3$MΩ) and input resistance (right, $V_{RP} = -72.3$mV) with (solid lines) and without (dashed lines) considering I_h. **C**, ATP consumption necessary to maintain membrane properties as a function of both resting potential and input resistance with (left) and without (right) considering I_h. Mean values for the control (black) and food restricted (red) cells are highlighted.

Figure S9: **Experimental membrane noise amplitude extraction and effects of different membrane noise levels on neuronal behavior of cells from control (CTR) and food-restricted (FR) animals.**

A, Preprocessing pipeline for membrane noise analysis: large deviations are excluded by windowing, followed by Notch-filtering to remove low-frequency oscillations, and normalization using a moving average to correct for slow baseline drifts. **B**, Power spectral density of membrane potential fluctuations, illustrating the frequency components of the signal. **C**, Probability density distribution of membrane potential fluctuations, used to estimate the standard deviation of intrinsic noise. **D**, Mean firing rate rises exponentially with increasing membrane noise for simulations of AdExp models with an orientation selectivity task (Fig. S1). **E**, Total energy expenditure mirrors the increase of firing rate but CTR energy demand is constantly higher than FR. **F**, Orientation selectivity per energy as a function of membrane noise, including non-spiking trials. **G**, Coding efficiency (mutual information per spike) mirrors the orientation selectivity index (OSI), showing an initial increase at low membrane noise levels due to stochastic resonance, followed by a monotonic decline at higher noise.

Figure S10: **Resting potential and synaptic reversal values shape firing rate, energy use, and information transmission (AdExp model with an orientation selectivity task (Fig. S1)).**

A, Postsynaptic firing rate, **B**, mutual information, **C**, and mutual information per spike (coding efficiency) as functions of the voltage gap between threshold and resting potential, modulated either by depolarizing the resting potential (turquoise) or lowering the threshold (purple). Lower thresholds produce higher firing rates and, consequently, higher mutual information, though at increased energetic cost. Energy consumption scales approximately linearly with firing rate at high rates. But, the coding-efficiency (mutual information per spike) is higher for a decreased resting potential. **D–F**, Sliding a fixed resting–threshold gap across the membrane potential range reveals that food-restricted cells (red) maintain **D**, higher firing rates, **E**, higher information content. However, at more depolarized resting potentials ($>$ –70 mV), control cells become more energy-efficient. **F**, Similarly to the higher energy efficiency, the coding-efficiency is higher for low resting potentials in control cells and higher for depolarized resting potentials in food-restricted cells.

Figure S11: **Design of Markov-process switching ON-OFF stimulus excitatory and inhibitory synaptic input similar to the stimulation protocol of Zeldenrust *et al.* (2024).**
A – Top, Excitatory synaptic weights follow a log-normal distribution ($\mu_{w_e} = 2.07, \sigma_{w_e} = 1.14$). **B – Bottom**, Excitatory presynaptic firing rates are normally distributed ($\mu_{r_e} = 1.47$ Hz, $\sigma_{r_e} = 0.23$ Hz). **B**, The preferred ON and non-preferred OFF stimuli assigned synaptic inputs based on the matching scheme, visualized as cityscape histograms: presynaptic firing rates (dark red) as a function of synaptic strength (green). The preferred stimulus (ON) elicits the strongest total synaptic drive, whereas non-preferred stimulus (OFF) exhibit the weakest. **C**, Hidden state representing the ON-OFF stimulus switching task. **D**, Raster plot of synaptic input times, distinguishing inhibitory (light blue), excitatory signal (dark red), and excitatory noise (dark red).

Figure S12: **SBI posterior distributions and validation of inferred parameters.**
Top row, Experimental voltage trace (Zeldenrust *et al.*, 2024) compared to a simulated trace generated using the posterior mean parameter estimates. **Right**, Voltage traces simulated using parameters sampled at the posterior mean ±1 standard deviation, illustrating uncertainty in the inferred dynamics. **Left**, One- and two-dimensional marginal posterior densities for inferred parameters, with posterior means (blue) and maximum a posteriori (MAP — purple) estimates indicated.

Figure S13: **Correlation structure of excitatory and inhibitory cell parameters and functional properties.**

Pairwise correlations among electrophysiological parameters and derived properties such as mutual information and total energy consumption from Zeldenrust *et al.* (2024) are shown separately for excitatory and inhibitory neurons.

Figure S14: **Principal component loadings (PC1–3) for all, excitatory, and inhibitory neurons.**
A, Loadings of the first three principal components (PC1–3) across all neurons, capturing the dominant axes of electrophysiological variability in our parameter inference of experimental data from Zeldenrust *et al.* (2024). **B**, PC1–3 loadings restricted to excitatory neurons, highlighting excitatory-specific variance structure. **C**, PC1–3 loadings restricted to inhibitory neurons, revealing distinct variance contributions relative to excitatory cells.