ER Stress-Induced β-Cell Apoptosis is Linked to Novel Select Lipid Signaling at the Transcriptional Level: Implications in T1D Development

[1,2]Xiaoyong Lei, [1,2]Anil K. Challa, [3]Susan E. Nozell, [1,2]Tomader Ali, [4]Daniel J. Stephenson, [4]Andrew Nafzinger, [2,5,6]Chad S. Hunter, [2,7]Adam R. Wende, [8]Ernesto S. Nakayasu, [1,2]Ying Gai-Tusing, [4,9,10]Charles E. Chalfant, [1,2]Sasanka Ramanadham**

[1]Department of Cell, Developmental, and Integrative Biology, University of Birmingham at Birmingham, AL 35294, USA

[2]Comprehensive Diabetes Center University of Birmingham at Birmingham, AL 35294, USA

[3]Nozell

[4]Department of Medicine, Division of Hematology & Oncology, University of Virginia, Charlottesville, VA, 22903

[5]Department of Medicine, University of Birmingham, Birmingham, AL 35294, USA

[6]KU Medical Center, The University of Kansas, Kansas City, Kansa, 66160

[7]Department of Molecular and Cellular Pathology, University of Birmingham, Birmingham, AL 35294, USA

[8]Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA 99354, USA

[9]esearch Service, Central Virginia Veterans Affairs Health Care System, Richmond, VA, 23249

[10]Program in Cancer Biology, University of Virginia NCI Comprehensive Cancer Center, Charlottesville, VA, 22903

**<u>*Corresponding Author</u>

Sasanka Ramanadham, PhD
Professor, Department of Cell, Developmental, and Integrative Biology *(CDIB)*
Senior Scientist, Comprehensive Diabetes Center
University of Alabama at Birmingham *(UAB)*
Shelby Biomedical Research Building, Rm. 1205
Birmingham, AL 35294-2182, USA
Office: 205-996-5973; 205-996-5220 (fax)
Email: sramvem@uab.edu

Running Title: Lipid signaling, transcription, NFκB, and β-cell death

Key words: ER stress, iPLA$_2$β-derived lipids, NFκB, transcriptional regulation, β-cell death

Abstract

Type 1 diabetes (T1D) is a consequence of β-cell death. ER stress precedes T1D onset and prolonged ER stress in β-cells can lead to β-cell apoptosis. We reported that lipid signaling generated by the Ca^{2+}-independent phospholipase $A_2\beta$ (iPLA$_2\beta$), encoded by *Pla2g6*, participates in ER stress-mediated β-cell apoptosis. β-Cell membranes are enriched in arachidonic acid containing glycerophospholipids and the iPLA$_2\beta$ catalyzes the hydrolysis of arachidonic acid in ER stressed β-cells. Metabolism of arachidonic acid leads to the generation of various proinflammatory lipids, raising the possibility that they contribute to ER stress and β-cell death leading to T1D. However, molecular mechanisms by which such β-cell-iPLA$_2\beta$-derived lipid (iDL) signaling contributes to β-cell apoptosis are not understood. It is well known that ER stress-mediated β-cell apoptosis is associated with induction of transcription factors, NFκB and STAT1. We report here that both induce *Pla2g6* and, unexpectedly, we find that iPLA$_2\beta$, which lacks DNA-binding motifs, associates with *NFkB*, *Stat1*, and *Pla2g6* promoter regions. Consistently, p65-NFκB and pSTAT1 induction is reduced with select inhibition or knockdown of iPLA$_2\beta$. Surprisingly, iPLA$_2\beta$ expression is also reduced by select inhibition of iPLA$_2\beta$, raising the possibility of feedback regulation by iDLs. In support, we find that select iDLs, recognized to be proinflammatory, enhance association of iPLA$_2\beta$ with *Pla2g6*, *Nfkb*, and *Stat1* promoter regions leading to induction of all three gene products and β-cell apoptosis. Our findings reveal previously unrecognized transcriptional regulation by iDL signaling and, iPLA$_2\beta$ itself, that leads to gene products that promote β-cell apoptosis. Analogous findings in human islets validate this mechanism raising the possibility that targeting select lipid signaling can reduce ER stress in β-cells and ameliorate T1D development.

Introduction

Type 1 diabetes (T1D) is a consequence of autoimmune destruction of β-cells and the involvement of immune cells is extensively-studied[1]. The β-cells also generate signals that support immune responses by mechanisms that are not well-understood. Our on-going studies highlight roles of select lipid signaling in β-cell apoptosis[2-9] and T1D development[10-13]. While several factors have been identified as inducers of β-cell apoptosis, there are no studies linking it with lipid signaling generated by β-cells themselves.

Accumulating evidence suggests that β-cell apoptosis due to ER stress is a critical contributor to β-cell death leading to T1D[14-27]. Importantly, ER stress in β-cells precedes T1D onset in both NOD, a spontaneous autoimmune mouse model of T1D, and humans[28,29]. Despite the mounting evidence linking ER stress with β-cell apoptosis leading to T1D, underlying molecular mechanisms triggered in the β-cells themselves that contribute to their death are not well-understood.

Our work reveals that the group VIA phospholipase A_2 (iPLA$_2$β) participates in β-cell apoptosis[2,4,5,30]. The iPLA$_2$β hydrolyzes the *sn*-2 substituent from membrane glycerophospholipids to release fatty acids[31]. In islets, iPLA$_2$β is predominantly localized in β-cells[2,4,32], and is induced in ER-stressed β-cells[2,4,32]. Various strategies (inhibitors, siRNA, and genetic-modulation) indicate a critical role for iPLA$_2$β in β-cell apoptosis due to ER stress[2,30,33-37]. We reported that inhibition or reduction of iPLA$_2$β in the NOD preserves β-cell mass and glucose tolerance, and reduces T1D incidence[10,13], supporting a critical role for iPLA$_2$β-derived lipids (iDLs) in T1D development.

The β-cell membranes are enriched in membrane glycerophospholipids containing arachidonic acid in the *sn*-2 position. ER stress in β-cells is associated with induction of iPLA$_2$β at the transcription level[30]. This can lead to increased hydrolysis of arachidonic acid, which can be metabolized by various enzymes to generate oxidized lipids, many of which are proinflammatory[38]. These include prostaglandins (PGs) that are generated via cyclooxygenases and are linked to stress responses[39-43]. They are also recognized to serve signaling/co-factor

functions to promote transcriptional activity[44-47]. However, the underlying molecular mechanisms these iDLs trigger in β-cells to promote their own death have not yet been identified.

Multiple transcription factors are activated in stressed β-cells leading to induction of genes encoding factors which contribute to β-cell apoptosis[48-50]. Among these are NFκB and STAT1 that are induced in β-cells during ER stress[48-50]. Herein, we report that in ER stressed β-cells, NFκB and STAT1 induce *Pla2g6* at the transcription level and intriguingly, that iPLA$_2$β, which does not contain DNA-binding motifs, associates with *Nfκb* and *Pla2g6* promoter regions. Our findings suggest a novel function of iDLs as co-factors that facilitate transcriptional regulation by iPLA$_2$β to trigger generation of gene products that promote ER stress and β-cell apoptosis. Hence, our study identifies potential candidate lipids that could be targeted to reduce ER stress and β-cell death and consequentially, ameliorate T1D development.

Methods

General materials. Coomassie reagents, SDS-PAGE reagents, kaleidoscope pre-stained molecular mass standards, and Triton X-100 (161-0324, 161-0407; BioRad, Hercules, CA); S-BEL ($iPLA_2\beta$ inhibitor), PGE_2 (10006801, 14010; Cayman Chemicals, Ann Arbor, MI); acetyl histone H4 (AcH4antibody, normal mouse IgG, protein A beads (06-911, 12-371, 16-125; EMD Millipore, Billerica, MA); Collagenase 2, fetal bovine serum (FBS) and common cell culture reagents (17101-015; Gibco, Carlsbad, CA); Immobilin-P PVDF membrane (IPVH00010; Millipore Corp., Bedford, MA; protease inhibitor cocktail (D4693), alpha-MEM media (P8340), and common reagents (Sigma Chemical Co., St. Louis, MO); and enhanced chemiluminescence reagent, (34095; Thermo- Fisher, Waltham, MA).

Cell culturing. Rat-derived INS-1 β-cells were cultured in Roswell Park Memorial Institute (RPMI, Cat. 11875093, Thermo Fisher Scientific Inc. Waltham, MA) 1640 medium, containing 11 mM glucose, 10% fetal calf serum, 10 mM HEPES buffer, 2 mM glutamine, 1 mM sodium pyruvate, 50 mM β-mercaptoethanol (BME), and 0.1% (w/v) each of penicillin and streptomycin in cell culture conditions (37°C, 5%CO_2/95% air). Mouse-derived MIN6 β-cells were cultured in Dulbecco's modified Eagle's medium (DMEM, Life Technologies Corporation, Grand Island, NY) containing 15% fetal calf serum, β-mercaptoethanol (150 µM), and 0.1% (w/v) each of penicillin and streptomycin in cell culture conditions (37°C, 5%CO_2/95% air). For any given experiment, except where indicated, the cells were seeded in 6-well plates and used at ~70% confluency. Data (Figs. 1 and 3) generated in INS-1 cells were used to guide all the remaining studies in MIN6 cells.

***Pla2g6* knockdown by siRNA**. Cells were seeded in 6-well tissue culture plates () in 2 ml antibiotic-free normal growth medium supplemented with FBS. After incubating the cells (37°C, 5%CO_2/95% air) until the cells 60-80% confluency was reached. The group VI $iPLA_2$ siRNA (m) (sc-43820, Santa Cruz, Dallas, TX) duplex solution was then added directly to the dilute Transfection Reagent (sc-36868, Santa Cruz Biotechnology, Dallas, Texas) using a pipette. After gentle mixing by pipetting, the mixture was incubated for 20 min at room temperature in cell culture hood and then transferred to an incubator (37°C, 5%CO_2/95% air). After 5-7h, 1 ml of normal growth medium containing 2X the normal serum and antibiotics concentration (2x normal growth medium) was added to the transfection mixture. After an additional 18-24h, the medium was aspirated and replaced with fresh 1x normal growth medium prior experimentation.

***Pla2g6* knockdown by CRISPR-Cas9 protocol**. Constructs that encoded guide 0 (control) or guide1 and 2 were used to generate control cell line and *Pla2g6* knockdown cells by using Lipofectamine™ 3000 Transfection

Reagent (L3000015, Thermo Fisher Sci). Transfected cells were selected with 0.25 μg/mL puromycin, and single-cell clones expanded and used for treatments.

Treatments. Cells were treated with DMSO or cytokines (50 U/ml IL-1β + 150 U/ml IFNγ) (401-ML and 485-MI; R & D Systems, Minneapolis, MN) and processed for immunoblotting, TUNEL, ChIP, and lipidomics analyses. To assess the impact of select pathways, cells were pretreated (1h) with inhibitors of iPLA₂β (*S*-BEL, 1 μM, Cayman Chemicals, Ann Arbor, MI, #10006801 or FKGK18 (5×10^{-8}M, generously provided by Dr. George Kokotos, University of Athens, Greece); iPLA₂γ (*R*-BEL, 1 μM, #10006800), cPLA₂ (Cay10502, 10 μM, #10008657), STAT1 activation (AZD 1480, 10 μM, #10702) from Cayman Chemicals, Ann Arbor, MI,); and NFκB activation (Bay11-7082, 10 μM, Sigma-Aldrich Inc., St. Louis, MO, #196870). To assess the role of select pathways, cells were treated with 10 nM PGE₂ (#14010), 10 μM PGF₂α (#16010), 10 μM 8-Iso PGF₂α (#16350, Cayman Chemical, Ann Arbor, Michigan)

Protein analyses. The 1° antibodies for targeted proteins included: iPLA₂β (sc-166616, 1:1,000), p65-NFκB (#sc-372, 1:1,000), CHOP (#sc-575, 1:1,000), IKBα (sc-371, 1:1000), tubulin (#sc-8035, 1:1,000), and Oct 1 (#sc-8024, 1:1,000) from Santa Cruz Biotechnology, Dallas, Texas; and pPERK (Cell Signaling Danvers, MA, #3179, 1:500). As quantitation of p65-NFkB, relative to tubulin and Oct-1 (nuclear marker), was found to be similar, only tubulin was used as loading control in subsequent experiments. The 2° antibodies concentrations were 1:10,000 (Goat anti mouse, sc-2031, Goat anti rabbit, sc-2030, Santa Cruz Biotechnology, Dallas, Texas). Immunoreactive bands were visualized by enhanced chemiluminescence. In the text, unless otherwise noted, representative immunoblots and corresponding densitometry quantitations are presented. The cumulative quantitations ± SEMs are presented as Supplemental figures.

***In situ* detection of DNA cleavage by TUNEL staining.** Cells were harvested and washed twice with ice-cold PBS and processed for TUNEL analyses using In Situ Cell Death Detection Kit (Cat. 1684795, Roche Diagnostic Corp., Indianapolis, IN) The cells were then immobilized on slides by cytospin and fixed with 4% paraformaldehyde in PBS (pH 7.4, 1h, room temperature). The cells were then washed with PBS and incubated in permeabilization solution (0.1% Triton X-100 in 0.1% sodium citrate in phosphate-buffered saline for 30 min at room temperature). The permeabilization solution was then removed and TUNEL reaction mixture (50 μL) added, and the cells incubated (1h at 37°C) in a humidified chamber. The cells were tjen washed again with PBS and counterstained with 1 μg/mL DAPI (4', 6'-diamidino-2-phenylindole, DAPI Cat. 62248, Thermo Fisher Scientific Inc., Waltham, MA in phosphate-buffered saline for 10 min to identify cellular nuclei. The incidence of apoptosis was assessed under a fluorescence microscope using a FITC filter. Cells with TUNEL-positive nuclei

considered apoptotic. DAPI staining was used to determine the total number of cells in a field. A minimum of four fields per slide was used to calculate the percent of cells that were apoptotic.

ChIP protocol. The ChIP analyses were performed by using Magna ChIP® A – a chromatin immunoprecipitation kit (Sigma-Aldrich Inc., St. Louis, MO, #17-610 Magna ChIP®), as previously described[51,52]. Briefly, MIN6 β-cells (3 x T225 flasks) were cultures in the absence (DMSO only) or presence of treatments for different durations, as described in appropriate results sections. The nuclei from cross-linked cells were resuspended in Tris/EDTA. The soluble chromatin was adjusted in RIPA buffer and pre-cleared with salmon sperm blocked protein A beads (Cat. 16-125; EMD Millipore, Billerica, MA). Immunoprecipitation was performed with 5 μg of antibodies directed against acetylated histone (AcH4), iPLA$_2$β or IgG. After pre-clearing and before immunoprecipitation, equal amounts of sonicated DNA (10 % volume of each sample) were reserved for qPCR analysis. The *Pla2g6, Nfkb, Stat1, Gcg, Pepck*, and *Il-6* promoter regions were probed with specific primers (**Table 1**) against the immunoprecipitated DNA by qPCR. Real-time qPCR was performed using Fast SYBR® Green PCR Master Mix (Thermo Fisher Scientific, Auburn, AL, #A25742) and carried out in a plate-based LightCycler® 480 System (Roche Life Sciences), as we described[5,10,13]. AcH4 was used to confirm transcriptional activity and IgG as a negative control.

Human islets. Islets from healthy donors were obtained through Integrated Islet Distribution Program (IIDP) Alberta Islet Distribution Program (AIDP)[53,54], Clinical islet Laboratory, Univ. Alberta Hospital. The islets upon receipt were immediately cleaned of non-islet material under a microscope. The islets were counted and cultured in RPMI 1640 medium (containing 10% FCS, 200 mM glutamine, 1% of 100 × penicillin-streptomycin, 25 mM HEPES) for 24h at 37°C under an atmosphere of 5%CO_2-95% air. The islets were then distributed into dishes for treatment and different analyses.

Lipidomics. Islets were then treated with cytokines (100 U/ml IL-1β + 300 U/ml IFNγ) from 2 to 36 h. In some experiments the islets were cultured with cytokines ± *S*-BEL at peak response time. The media were collected for analysis of eicosanoids by UPLC ESI-MS/MS analyses, as described[10,12,13]. The media was combined with an IS mixture comprised of 10% methanol (400 μL), glacial acetic acid (20 μL) and internal standard (20 μL) containing the following deuterated eicosanoids (1.5 pmol/μL, 30 pmol total). (All standards purchased from Cayman Chemicals): (d4) 6keto-prostaglandin F$_1$α, (d4) prostaglandin F$_2$α, (d4) prostaglandin E$_2$, (d4) prostaglandin D$_2$, (d8) 5-hydroxyeicosatetranoic acid (5-HETE), (d8) 12-hydroxyeicosa-tetranoic acid (12-HETE) (d8) 15-hydroxyeicosa-tetranoic acid (15-HETE), (d6) 20-hydroxyeicosa-tetranoic acid (20-HETE), (d11) 8,9-epoxyeicosatrienoic acid, (d8) 14,15 hpoxyeicosa-trienoic acid, (d8) arachidonic acid, (d5) eicosapentaenoic acid (EET), (d5) docosahexaenoic acid, (d4) prostaglandin A$_2$, (d4) leukotriene B$_4$, (d4) leukotriene C$_4$, (d4) leukotriene D$_4$, (d4) leukotriene E$_4$, (d5) 5(*S*),6(*R*)-lipoxin A$_4$, (d11) 5-iPF$_2$α-VI, (d4) 8-iso

prostaglandin $F_2\alpha$, (d11) (±)14,15-dihydroxyeicosatrienoic acid (DHET), (d11) (±)8,9-DHET, (d11) (±)11,12-DHET, (d4) prostaglandin E_1, (d4) thromboxane B_2, (d6) dihomo-gamma linoleic acid, (d5) resolvin D2, (d5) resolvin D1 (RvD1), (d5) maresin2, (d7) 5-OxoETE, and (d5) resolvin D3. Samples and vial rinses (5% MeOH; 2 mL) were applied to Strata-X SPE columns (Phenomenex), previously washed with methanol (2 mL) and then dH_2O (2 mL). Eicosanoids eluted with isopropanol (2 mL), were dried *in vacuuo* and reconstituted in EtOH:dH_2O (50:50;100 µL) prior to ultra-high performance liquid chromatography electrospray ionization-MS/MS (UPLC ESI-MS/MS) analysis.

The eicosanoids were separated using a Shimadzu Nexera X2 LC-30AD coupled to a SIL-30AC auto injector, coupled to a DGU-20A5R degassing unit in the following way. A 14 min, reversed phase LC method utilizing an Acentis Express C18 column (150 mm x 2.1 mm, 2.7 µm) was used to separate the eicosanoids at a 0.5 mL/min flow rate at 40°C. The column was equilibrated with 100% Solvent A [acetonitrile:water:formic acid (20:80:0.02, v/v/v)] for 5 min and then 10 µL of sample was injected. 100% Solvent A was used for the first two min of elution. Solvent B [acetonitrile:isopropanol:formic acid (20:80:0.02, v/v/v)] was increased in a linear gradient to 25% solvent B at 3 min, to 30% at 6 min, to 55% at 6.1 min, to 70% at 10 min, to 100% at 10.10 min, 100% solvent B was held constant until 13.0 min, where it was decreased to 0% solvent B and 100% Solvent A from 13.0 min to 13.1 min. From 13.1 min to 14.0 min. Solvent A was held constant at 100%. The eicosanoids were analyzed via mass spec using an AB Sciex Triple Quad 5500 Mass Spectrometer. Q1 and Q3 were set to detect distinctive precursor and product ion pairs. Ions were fragmented in Q2 using N2 gas for collisionally-induced dissociation. Analysis used multiple-reaction monitoring in negative-ion mode. Eicosanoids were monitored using precursor → product MRM pairs. The mass spectrometer parameters used were: Curtain Gas: 20 psi; CAD: Medium; Ion Spray Voltage: -4500 V; Temperature: 300 °C; Gas 1: 40 psi; Gas 2: 60 psi; Declustering Potential, Collision Energy, and Cell Exit Potential vary per transition.

TUNEL analyses. Human islets (250/condition) were treated with cytokines or PGs, as described above, for 24h and then processed for TUNEL analyses

ChIP Analyses. Human islets (3500-5000/condition) were treated with cytokines for 2h or PGs for 4h, as described above, and processed for ChIP using antibodies select for p65 or iPLA2β and subsequent qPCR analyses for *PLA2G6*, *p65-NFκB*, and *GcG* promoter regions.

Statistical Analysis. All statistical analyses were done using GraphPad Prism (version 10). A "p" value < 0.05 was defined as significant. Student's t-test was used for comparisons between 2 groups. For lipidomics analyses, "p" values were determined by two-way ANOVA, followed by Dunnett's multiple comparisons test.

Results

iPLA$_2$β induction is mitigated by inhibition of cytokine-induced ER Stress. β-cell ER stress precedes T1D development and inflammatory cytokines play critical roles in this process. We reported that ER stress induces iPLA$_2$β expression increased in ER-stressed β-cells[8] and that targeting iPLA$_2$β reduces T1D incidence[11]. In the present study, we sought to examine the molecular mechanisms that lead it increases in iPLA$_2$β expression. To address this, we first examined whether mitigation of ER stress impacts iPLA$_2$β expression. We find that proinflammatory cytokines-induced INS-1 β-cell apoptosis is prominent after 24h (**Fig. 1A**) and that this was associated with induction of iPLA$_2$β and ER stress factors (pPERK and CHOP) (**Figs. 1B-E**, cumulative densitometries are presented in **Suppl. Fig. 1**). As expected, the chemical chaperon PBA reduced induction of pPERK and CHOP, and β-cell apoptosis. Moreover, it reduced iPLA$_2$β expression, suggesting a causative link between ER stress and iPLA$_2$β expression in β-cells. The participation of NFκB in β-cell death leading to T1D is well established and we find a time-dependent nuclear accumulation of p65-NFκB following exposure to cytokines that peaked at 16h (**Fig. 1F-G**). However, in the presence of PBA, there was reduction in p65-NFκB (**Fig. 1H**). Given the established roles of both NFκB and STAT1 as transcriptional inducers in β-cells during T1D development, we considered the possibility that their activity in stressed β-cells is linked to iPLA$_2$β expression.

ER-stressed-mediated activation of NFκB and STAT1 in β-cells is linked to iPLA$_2$β expression. Given the transcriptional activity of p65-NFκB and STAT1, we assessed whether ER-stressed induces iPLA$_2$β through an NFκB- and/or STAT1-dependent mechanism. β-cells were pre-treated with Bay11-7082 or AZD 1480 (inhibitors of NFκB and STAT1 activation, respectively) prior to addition of cytokines. We find that both Bay11 (**Figs. 2A,B-F**) and AZD (**Figs. 2G, H-I**) reduced p65-NFκB and pSTAT1 expression, as well as pPERK and CHOP (cumulative densitometries are presented in **Suppl. Fig. 1**) and these outcomes were associated with reduced β-cell apoptosis (**Fig. 1J**). Further, cytokine-induced iPLA$_2$β expression was also reduced in the presence of the two inhibitors. These findings suggest that iPLA$_2$β induction in ER-stressed β-cells occurs via NFκB and STAT1-dependent pathways.

In their roles as transcription factors, p65-NFκB and pSTAT1 induce genes associated with inflammation and apoptosis. We therefore considered the possibility that iPLA$_2$β expression during ER stress is through p65-NFκB and/or pSTAT1 at the transcriptional level. Survey of the *Pla2g6* gene identified putative promoter regions containing binding motifs that are recognized by p65-NFκB and pSTAT1. This raised the possibility that p65-NFκB and pSTAT1 can induce transcription of *Pla2g6*. To address this, INS-1 cells were treated with cytokines and then processed for CHIP protocol using 1° antibodies targeting p65, STAT1, acetylated histone (AcH4), or IgG. The IPed DNA was then processed for qPCR analyses for *IL-1* (positive control) and *Pla2g6* promoter

regions. Preliminary studies suggested peak effects with 2h cytokine treatment (**Suppl. Fig. 2**). We find exposure to cytokines promotes enrichment of p65-NFκB and pSTAT1 at the *IL-6* promoter region in β-cells (**Fig. 3A**) confirming triggering of transcriptional activation by cytokines and verified by AcH4 enrichment at the *IL-6* promoter region. Under the same conditions, p65-NFκB and pSTAT1, and AcH4 were also found to be enriched at the *Pla2g6* promoter region (**Fig. 3B**). Specificity of p65-NFκB, pSTAT1, and AcH4 enrichment at both *IL-6* and *Pla2g6* promoter regions was reflected by the absence of IgG (negative control) enrichment at the two promoter regions. These findings suggest that iPLA$_2$β expression in ER-stressed β-cells is modulated of p65-NFκB and STAT1, in part, at the transcriptional level.

Impact of iPLA$_2$β on NFκB and STAT1 Activation in ER Stressed β-Cells. As targeting iPLA$_2$β reduced T1D incidence, we considered the impact of iPLA$_2$β itself on ER stress at the molecular level. To address this, we compared the effects of selective inhibition of iPLA$_2$β, iPLA$_2$γ, and cPLA$_2$α. We find that inhibition of iPLA$_2$β with either *S*-BEL[55] (**Fig. 4A**) or FKGK18 (**Fig. 4B**)[56,57] was effective in reducing p65-NFκB, pSTAT1, pPERK, and CHOP. However, inhibition of iPLA$_2$γ with *R*-BEL[55] (**Fig. 4C**) or cPLA$_2$α with Cay10502 (**Fig. 4D**) did not lead to reductions in p65-NFκB, pSTAT1, pPERK, or CHOP. (Cumulative densitometries are presented in **Suppl. Fig. 3**). These findings suggest a select impact of iPLA$_2$β in ER-stress β-cells.

To verify that such an impact is specifically iPLA$_2$β-dependent, siRNA was used to knock down *Pla2g6* mRNA and the cells were treated with vehicle or cytokines. While the cytokines induced iPLA$_2$β, p65, pSTAT1, pPERK, and CHOP in cells transfected with scrambled RNA (scrRNA), they were not in cells transfected with siRNA targeting *Pla2g6* (**Fig. 5**). Further, p65-NFκB induction evident with scrRNA was also reduced with siRNA treatment (**Figs. 5A,C**), in the absence of reductions in IκBα (**Figs. 5D**). (Cumulative densitometries are presented in **Suppl. Fig. 3**). Consistently, only inhibition of iPLA$_2$β (with *S*-BEL or FKGK18) or iPLA$_2$β knockdown (with siRNA), but not inhibition of iPLA$_2$γ or cPLA$_2$α, mitigated cytokine-induced β-cell apoptosis (**Fig. 6**). These findings suggest a select impact of iPLA$_2$β on ER stress outcomes in β-cells, including activation of NFκB and STAT1.

CRISPR-Cas9 knockdown of iPLA$_2$β reduces its association with promoters. We previously reported that during ER stress, iPLA$_2$β translocates to the nucleus[8] and in view of the above findings, we considered the possibility iPLA$_2$β modulates NFκB at the gene level. To address this question, we established a CRISPR-Cas9 approach to knockdown iPLA$_2$β. Using Benchling (www.benchling.com), we designed two CRISPR guides (G1 and G2) targeting the coding region in exon 2 (**Fig. 7A**). Following the method described by Ran et al[58], we generated plasmid constructs that express the guide RNAs along with Cas9 protein fused either to EGFP or puromycin to facilitate screening and selection. We then transfected MIN6 β-cells with either control plasmids

(G0) or plasmids encoding gRNAs (G1 and G2) targeting *Pla2g6*; in addition to indels at each of the two target sites, simultaneous cutting of both target sites followed by NHEJ repair would result in a predicted deletion. We performed genotyping by PCR followed by TBE-PAGE analyses to confirm the presence of the deletion allele (**Fig. 7B**). Whereas only the expected wildtype band (298 bp) is predominantly expressed in G0-treated (control) cells, the cells transfected with both the CRISPR plasmids (G1 and G2) had additional truncated bands, in particular one at 161bp, which is expected from the 137 bp deletion by synchronous activity of both CRISPR guides. Additional minor bands that are uniquely found in the CRISPR targeted samples are likely to be alleles with different deletions arising from NHEJ, which is a relatively common phenomenon[59]. The 137 bp deletion leads to a frameshift in the coding region resulting in an early truncation. The mutant allele encodes a 30 aa peptide with only 8 native amino acids at the N-terminal and 22 new amino acids resulting from the frameshift (**Fig. 7C**). Immunoblotting analyses verified the knockdown of basal and cytokine-induced iPLA$_2$β with the CRISPR constructs (G1 and G2), but not with the control plasmids (G0) (**Fig. 7D,E**). Consistently, cytokine-induced β-cell apoptosis was unaffected by G0 but reduced with the CRISPR targeting (**Fig. 7F**). As both gRNAs appeared to work similarly, they were used in combination in subsequent experiments. Such studies demonstrated reductions in cytokine-induced iPLA$_2$β, p65-NFκB, pSTAT1, and pPERK with CRISPR targeting, relative to controls (**Figs. 7G-I**). These findings provide evidence of successful application of the CRISPR-Cas9 protocol to knockdown iPLA$_2$β in β-cells, resulting in the mitigation of ER stress outcomes.

Evidence for iPLA$_2$β impact at the transcription level. To determine if iPLA$_2$β modulates NFκB or STAT1 at the transcription level, CRISPR-Cas9-modified MIN6 cells were treated with DMSO vehicle or cytokines for 2h, as determined in pilot studies (**Suppl. Fig. 2**). After 2h, a ChIP protocol was performed using antibodies directed against iPLA$_2$β, AcH4, and IgG. The IPed DNA was then used for qPCR analyses of *Pla2g6* and *Nfκb* promoter regions. Intriguingly, we find enrichment of iPLA$_2$β at both *Pla2g6 and Nfκb* promoter regions under both basal and cytokine-exposed conditions in both the G0 and gRNAs-treated, relative to IgG enrichment (**Figs. 8A,B**). Under the same conditions, enrichment of AcH4 at the *Pla2g6* promoter region was also evident, indicative of triggering of *Pla2g6* transcription. Interestingly, enrichment of iPLA$_2$β at the *Pla2g6* and *Nfκb* promoter regions or of Ach4 at the *Pla2g6* promoter region were enhanced by cytokines, relative to DMSO, in the G0 group but not in the gRNA-treated group (**Figs. 8C**). These findings suggest a correlation between iPLA$_2$β abundance and its association with promoters and ability to modulate transcriptional activity.

Because iPLA$_2$β is known to not contain any DNA binding motifs, the above findings were unexpected. However, as iDL signaling participates in ER stress-induced β-cell death, we anticipated that inhibition of iPLA$_2$β may impair its association with promoters. We therefore performed ChIP analyses as above, but this time compared the MIN6 cells response in the absence and presence of *S*-BEL. As above, we find enrichment of iPLA$_2$β at the *Pla2g6*, *Nfκb*, and of AcH4 at the *Pla2g6* promoter regions under both basal (**Fig. 8D**) and cytokine-

exposed (**Fig. 8E**) conditions in the absence and presence of *S*-BEL, relative to IgG. Surprisingly, in contrast to the above findings in iPLA$_2$β-reduced MIN6 cells, enrichment of iPLA$_2$β at the *Pla2g6* and *Nfκb* promoter regions was similar in the absence and presence of *S*-BEL (**Fig. 8F**). However, Ach4 enrichment at the *Pla2g6* promoter region was decreased, suggesting mitigation of transcriptional activity. These findings raise the possibility of a unique separation of association with a promoter region from activation of transcription, involving lipid signaling.

ER stress-induced production of prostaglandins is reduced by inhibition of iPLA$_2$β. We previously reported dramatic changes in proinflammatory eicosanoids profile in immune cells during T1D development[12]. A potential explanation for the above findings may be related to the abundances of IDLs in the β-cells, whose production would be limited by iPLA$_2$β activation. To address this possibility, we performed lipidomics analyses to identify changes in lipid abundances associated with ER stress. As insulinoma cells generated weak lipid intensity signals and the required temporal analyses necessitated large batches of islets, we utilized human islets for these analyses. We previously reported that cytokines induce ER stress and human β-cells apoptosis that are mitigated by *S*-BEL[5]. Here, we report that under the same conditions, islets production of several proinflammatory eicosanoids increases in a temporal fashion that peaks at 24h. Among those, the production of PGE$_2$, PGF$_2\alpha$, and 8-Iso-PGF$_2\alpha$ is blunted by the addition of *S*-BEL (**Figs. 9A-C**). These PGs are noted for their potent pro-inflammatory effects[38] raising the possibility of that their signaling contributes to ER stress-induced β-cell apoptosis.

Select prostaglandins contribute to ER stress and apoptosis. To address this possibility, MIN6 cells were treated with PGE$_2$, PGF$_2\alpha$, and 8-Iso-PGF$_2\alpha$ alone or in combination for 24h and then processed for TUNEL analyses. We find that each PG alone induced MIN6 β-cell apoptosis, relative to DMSO (**Supplemental Fig. 4**) that was ~50% of that caused by cytokines. And, in combination they induced ~67% of that caused by cytokines (**Fig. 10A**). Further, the combination of the three PGs induced p65-NFκB, pSTAT1, pPERK, and CHOP, relative to DMSO-treated cells (**Fig. 10B-F**). These findings suggest that PGE$_2$, PGF$_2\alpha$, and 8-Iso-PGF$_2\alpha$ signaling participates in ER stress-mediated β-cell apoptosis.

Feedback regulation of iPLA$_2$β expression. While regulation of enzymes by PGs has been reported[60], their potential impact on β-cell iPLA$_2$β has not. This possibility surfaced by the findings that while select inhibition of iPLA$_2$β with *S*-BEL (**Fig. 11**) or FKGK18 (**Fig. 11B**) had a minor effect alone in MIN6 cells, they markedly reduced cytokine-induced iPLA$_2$β expression. A select nature of iPLA$_2$β effect was reflected by the absence of a similar reduction with inhibition of iPLA$_2$γ or cPLA$_2$α (**Fig. 11C,D**). Analogous to cytokine effects, the combination PG treatment also increased iPLA$_2$β expression (**Fig. 11E**). (Cumulative densitometries are

presented in **Suppl. Figs. 3 and 6**) These findings suggest that PGs production by stressed cells can induce iPLA$_2$β expression.

Select prostaglandins promote iPLA$_2$β association with promoters. Translocation of iPLA$_2$β to the nucleus amplifies the activity of truncated iPLA$_2$β resulting in accelerated hydrolysis of the *sn*-2 fatty acid from membrane glycerophospholipids and this would be expected to increase the abundances of iDLs in the nucleus. To assess whether the select PGs participate in transcriptional regulation to upregulate iPLA$_2$β expression, we performed ChIP analyses with MIN6 cells treated with cytokines or the combination of PGE$_2$, PGF$_2$α, and 8-Iso-PGF$_2$α in the absence or presence of *S*-BEL. Pilot studies indicated that 4h treatment produced peak effects (**Suppl. Fig. 7**) and subsequent analyses revealed that the combination PGs increased the association of iPLA$_2$β with *Pla2g6*, *Nfκb*, and *Stat1* promoter regions, relative to IgG (**Fig. 12A**). Further, the association of iPLA$_2$β with *Pla2g6* (**Fig. 1B**), *Nfκb* (**Fig. 1C**), and *Stat1* (**Fig. 1D**) following exposure to cytokines was significantly increased, relative to DMSO. Specificity of such enrichment was evidenced by the lack of PGs-induced enrichment of iPLA$_2$β with *Gcg* (**Fig. 1E**) or *Pepck* (**Fig. 1F**) promoter regions. Addition of *S*-BEL did not have an effect alone or significantly impact stimulated association of iPLA$_2$β with any promoter regions. These findings suggest a mechanism by which the PGs may confer upon iPLA$_2$β the ability to activate transcription to not only promote gene product NFκB but, also iPLA$_2$β. These findings suggest that PGs produced by stressed β-cells contribute to β-cell death by inducing iPLA$_2$β, NFκB, and STAT1 expression at the transcription level.

Transcriptional regulation by select iDLs in human islets. To assess whether transcriptional regulation of and by iPLA$_2$β translates to islet β-cells, ChIP analyses were performed in human islets. First, the association of p65 with *Pla2g6* promoter was confirmed by treating the islets with cytokines followed by a ChIP protocol using antibodies directed against p65-NFkB or IgG. Subsequent qPCR analyses revealed cytokine-induced enrichment of p65-NFκB at the *PLA2G6*, but not at the *GCG*, promoter region, relative to IgG (**Fig. 13A, Suppl. Fig. 7**). Also evident were cytokine-induced enrichment of iPLA$_2$β at the *PLA2G6* and *NFκB*, but not at the *GCG*, promoter regions, relative to IgG (**Fig. 13B**). Further, the select PGs also promoted enrichment of iPLA$_2$β at the *PLA2G6* and *NFκB*, but not at the *GCG*, promoter regions, relative to IgG (**Fig. 13C**). Consistently, the PGs combination induced a 2-fold increase in β-cell apoptosis, relative to DMSO (**Fig. 13D**). In comparison, the cytokines indued a 3-fold increase in β-cell apoptosis. These findings validate a link between iPLA$_2$β/iDLs and transcriptional regulation of relevant apoptotic factors in human islet β-cells.

Discussion

T1D is a consequence β-cell destruction resulting from concerted actions of immune cells and β-cells. While the immunologic components have been heavily dissected, the role of β-cells in causing its own demise are not well-understood. We recently reported that select reduction of iPLA$_2$β in macrophages or T-cells reduces their inflammatory phenotype and lowers T1D incidence[10,13]. However, the nature of signals derived from β-cells to drive this process are not completely understood.

Our work has implicated iPLA$_2$β activation in β-cells as a critical contributor to ER stress-induced β-cell death[2-5,8]. Reports suggesting that ER stress in β-cells precedes T1D onset[29,61] raises the possibility that iDL signaling is an early trigger leading to β-cell death and T1D. Activation of PLA$_2$s leads to hydrolysis of the *sn*-2 substituent of membrane glycerophospholipids and islet β-cell membranes are enriched in *sn*-2 arachidonic acid[62]. Subsequent metabolism of arachidonic acid leads to generation of oxidized bioactive lipids including prostaglandins, several of which are potently inflammatory[38]. Here we sought to identify molecular mechanisms through which iDL signaling participates in ER stress-induced β-cell death. Utilizing insulinoma cells and human islets; employing chemical, siRNA, and CRISPR-Cas9 approaches; and immunoblotting, TUNEL, qPCR, ChIP, and lipidomics analyses, we report (a) a select role of iPLA$_2$β in ER stress and β-cell apoptosis, (b) regulation of iPLA$_2$β expression by NFκB and STAT1 at the transcriptional level, (c) regulation of NFκB, STAT1, and iPLA$_2$β expression by iPLA$_2$β at the transcriptional level, (d) potential co-factor roles of select inflammatory prostaglandins to facilitate transcriptional activation by iPLA$_2$β, and (e) feedback regulation of iPLA$_2$β expression.

ER stress elicits adaptive unfolded protein response (UPR) initially, however, prolonged ER stress can lead to terminal UPR, where hyperactivation of the UPR arms (PERK, IRE1α, ATF6) triggers apoptotic targets[63,64]. The nuclear transcription factors NFκB and STAT1, by virtue of being able to trigger pro-apoptotic genes expression are key players in cytokine-mediated β-cell death and T1D development[65-74]. Our finding that inhibiting NFκB or STAT1 activation decreases iPLA$_2$β expression, in association with a decrease in β-cell apoptosis, led us to further explore the link between iPLA$_2$β and the two transcription factors in ER stressed β-cells. Initial ChIP analyses with antibodies directed against NFκB and STAT1 revealed that cytokine exposure promotes association of NFκB and STAT1 at *Pla2g6* promoter region. While gene analyses identified evolutionarily conserved region in the *Pla2g6* promoter that harbors putative response elements for transcription factors that respond to ER stress, including NFκB and STAT1[75], our findings for the first time provide evidence for a potential regulation of iPLA$_2$β by NFκB and STAT1 at the transcriptional level.

Inhibition or genetic knockdown of iPLA$_2$β decreases while overexpression of iPLA$_2$β increases the susceptibility of islet β-cells to ER stress and apoptosis and reduces T1D incidence[4,11,12]. Here, we present evidence that both outcomes are impacted selectively by iPLA$_2$β, but not iPLA$_2$γ or cPLA$_2$α. In view of this, we considered the impact of modulating iPLA$_2$β on NFκB. We find that select chemical inhibition, siRNA-mediated downregulation, or CRISPR-Cas9-mediated

reduction of iPLA$_2\beta$ reduces NFκB and STAT1 activation, suggesting their potential modulation by iPLA$_2\beta$ in ER stressed β-cells. When ChIP analyses were repeated with antibodies directed against iPLA$_2\beta$, we find enrichment of iPLA$_2\beta$ at *Nfκb* and *Stat1* promoter regions in ER-stressed β-cells, suggesting potential transcriptional regulation of *Nfκb* and *Stat1* by iPLA$_2\beta$. However, as iPLA$_2\beta$ does not contain DNA binding motifs, we considered the possibility that co-factors, previously not identified, facilitate such regulation.

Lipids are recognized to promote gene transcription[76-78] and our lipidomics analyses reveal increased production of proinflammatory eicosanoids by β-cells with exposure to cytokines. Among the more prominent were PGE$_2$, PGF$_2\alpha$, and 8-iso-PGF$_2\alpha$. These lipids induced β-cell apoptosis, that was associated with induction of NFκb, STAT1, and ER stress, suggesting that their signaling participates in ER stress-induced β-cell apoptosis. Interestingly, they also induced iPLA$_2\beta$. This is not unprecedented as PGE$_2$ has been reported to feedback regulate COX2[77], which metabolizes arachidonic acid to generate PGs. In view of our findings, we assessed whether the PGs confer activity at the transcriptional level in ER stressed β-cells. Intriguingly, we find that the PGs promotes iPLA$_2\beta$ association with *Pla2g6* and *Nfkb* promoter regions, that were not significantly impacted in the presence of the iPLA$_2\beta$ inhibitor *S*-BEL, providing a likely mechanism by which the PGs serve as co-factors to promote association of iPLA$_2\beta$ with *Pla2g6* and *Nfkb* promoters.

This likely mechanism, however, was not congruent with the findings that chemical inhibition of iPLA$_2\beta$, which would be expected to reduce eicosanoid production[79], reduced NFκB and STAT1 activation, ER stress, and β-cell apoptosis, though it did not impair iPLA$_2\beta$ enrichment at *Pla2g6* or *Nfκb* promoter regions. Further assessments with chemical inhibitors revealed that only inhibition of iPLA$_2\beta$, but not of iPLA$_2\beta$ or cPLA$_2\alpha$, reduced iPLA$_2\beta$. Taken together, we posit that there are two separate but required processes in play: one, is the cytokine-induced enrichment of iPLA$_2\beta$ with promoter regions, which does not require co-factors. Second and subsequent to iPLA$_2\beta$ association with promoters, triggering of its transcriptional "activity" is facilitated by the tested PGs, serving as co-factors/co-activators. This is consistent with the mitigated ER stress outcomes with chemical inhibition of iPLA$_2\beta$, though its association with promoters remains intact. Such a mechanism may also explain why under basal conditions, overexpression of iPLA$_2\beta$ does not induce deleterious outcomes, but under stress they are unmasked[4,80]. This is consistent with a prerequisite need for *de novo* generated lipid signaling to facilitate iPLA$_2\beta$ activity at the promoter regions and warrants further investigation.

Our findings provide a rational novel sequence of feedback events (**Fig.13**), where ER stress in β-cells (a) induces NFκB and STAT1, which in turn induce *Pla2g6* to increase expression iPLA$_2\beta$ and (b) induces iPLA$_2\beta$ localization to nucleus and its association with *Pla2g6*, *Nfκb,* and *Stat1* promoter regions, and (c) leads to increased generation of iPLA$_2\beta$-derived PGs which facilitate iPLA$_2\beta$ transcriptional activity. This sequence of events links select lipid signaling with triggering of transcription of NFκB and STAT1 genes, leading to generation of apoptosis-related factors that contribute to β-cell death. Importantly, both cytokines and the select iDLs promoted association of iPLA$_2\beta$ with *Pla2g6* and *Nfκb* promoter regions suggesting that ER stress elicits the same sequence of events, in the context of human islet β-cell apoptosis leading to T1D development.

Figure 1. NFκB Activation During ER Stress-Induced β-Cell Apoptosis is Associated with iPLA$_2$β Induction. INS-1 β-cells were treated with DMSO or Cytokines (50 U/ml IL-1β + 150 U/ml IFNγ) for 16h or 24h (**A**), 16h (**B-E, H**), or 2-48h (**F-G**). **A**. Cells were processed for TUNEL analyses and mean ± SEMs of %apoptotic cells, relative to total cells, are presented. ([*,†]Significantly different from corresponding DMSO group, $p < 0.05$; [†]significantly different from corresponding DMSO and cytokine groups, $p < 0.05$, n=3/group.) **B-E, H**. Representative blot from cells treated without or with PBA for iPLA$_2$β, p65-NFκB, pPERK, and CHOP expression and corresponding densitometries. Tubulin and Oct-1 (nuclear marker) were uses as loading controls in n=3 independent analyses. Cumulative densitometries are presented in **Suppl. Fig. 2**. **F, G**. Representative blot of iPLA$_2$β expression in cytosolic and nuclear fractions prepared, as described[7], and cumulative densitometry.

Figure 2. <u>**Inhibition of NFκB and STAT1 Reduces ER Stress-Mediated iPLA$_2$β Induction and β-Cell Apoptosis**</u>. **A**. MIN-6 cells were pre-treated with NFκB inhibitor Bay11-7082 (10 µM) or STAT1 inhibitor AZD 1480 (10 µM) for 1h. The cells were then treated with DMSO or cytokines (50 U/ml IL-1β + 150 U/ml IFNγ) for 16h and processed for p65-NFκB, SpTAT1, iPLA$_2$β, pPERK, CHOP, and loading control tubulin immunoblotting analyses. Representative blots after BAY11 treatment (**A**) and corresponding densitometries (**B-F**)). Representative blots after AZD treatment (**G**) and corresponding densitometries (**H-I**). Cumulative densitometries are presented in **Suppl. Fig. 1**. **J**. Cells treated for 24h were processed for TUNEL analyses. ([a]Significantly different from DMSO, $p < 0.0001$; [b]Significantly different from CTK+Bay11, $p = 0.0007$; [c]Significantly different from CTK+AZD, $p = 0.0028$.)

Figure 3. <u>**Transcriptional Induction of *Pla2g6* by NFκB and STAT1**</u>. INS-1 Cells were treated with 50 U/ml IL-1β + 150 U/ml IFNγ for 2h and processed for ChIP analyses. Nuclear preparations were immunoprecipitated with antibodies directed at p65, STAT1, AcH4, or IgG control and probed for association with *IL-6* (**A**) and *Pla2g6* (**B**) promoter regions by qPCR.

Figure 4. Selective Impact of iPLA$_2$β Inhibition on NFκB and STAT1 Activation in ER Stressed β-Cells. MIN6 β-cells were pre-treated for 1h with selective inhibitors for iPLA$_2$β (**A**) <u>S-B</u>EL (1 µM) or (**B**) <u>FK</u>GK18 (5 x 10^{-8} M); iPLA$_2$γ (**C**) <u>R-B</u>EL (1 µM); or cPLA$_2$α (**D**) Cay10502, (10 µM) and then treated for 16h with DMSO, or Cytokines (50 U/ml IL-1β + 150 U/ml IFNγ). The cells were then processed for p65-NFκB, pPERK, CHOP, pSTAT1, and loading control tubulin immunoblotting. Representative blots and densitometries are presented. Cumulative densitometries are presented in **Suppl. Fig. 3**.

Figure 5. Knockdown of iPLA$_2$β Reduces ER Stress and NFκB and STAT1 Activation in ER Stressed β-Cells. MIN6 β-cells were transfected with s̲crambled RNA or s̲iRNA targeting *Pla2g6* and treated with D̲MSO or C̲ytokines (50 U/ml IL-1β + 150 U/ml IFNγ) and processed for iPLA$_2$β, p65-NFκB, IKBα, pSTAT1, pPERK, CHOP, and loading control tubulin immunoblotting. Representative blots (**A**) and corresponding densitometries (**B-G**) are presented. Cumulative densitometries are presented in **Suppl. Fig. 3**.

Figure 6. Select Targeting of iPLA$_2$β Inhibits ER Stress-Induced β-Cell Apoptosis. **A.** MIN6 β-cells were pretreated with *S*-B̲EL (1 μM), F̲KGK18 (5 x 10^{-8} M), *R*-B̲EL (1 μM), or Cay10502 (10 μM) for 1h or transfected with s̲crambled or siRNA targeting *Pla2g6* and then treated with D̲MSO or c̲ytokines (50 U/ml IL-1β + 150 U/ml IFNγ) for 24h and processed for TUNEL analyses. Data are means±SEMs of % apoptotic cells, relative to total cells. ([a]Cytokine group significantly different from DMSO group, $p<0.0001$; [b,c,d,e,f,g,h]Cytokine group significantly different from cytokines + inhibitor or siRNA groups: $p=0.0001$, $p=0.0002$, $p=0.0228$, $p=0.0007$, $p=0.0010$, $p=0.0004$, $p=0.0001$, respectively.

Figure 7. CRISPR-Cas9 Approach to Examine iPLA$_2$β Role in Transcription. MIN6 β-cells were transfected with control (G0) or guide RNA constructs (G1 and G2) targeting *Pla2g6*. **A.** *Pla2g6* CRIPR-Cas9 gRNAs. **B**. Genotyping analyses by PCR. **C**. Predicted deletion region. **D, E**. Cells were treated with D̲MSO or C̲ytokines (50 U/ml IL-1β + 150 U/ml IFNγ) for 16h and then processed for iPLA$_2$β and loading control tubulin immunoblotting. Representative blots and densitometries are presented. **F**. Cells were treated with D̲MSO or C̲ytokines (50 U/ml IL-1β + 150 U/ml IFNγ) for 24h and then processed for TUNEL analyses. Data are means±SEMs of percent apoptotic cells, relative to total cells. ([†,*]Significantly different from G0-DMSO, $p<0.00001$, $p<0.005$; [#]different from G0-CTK $p<0.0005$, n=3/group). **G-I**. Cells were treated with D̲MSO or C̲ytokines (50 U/ml IL-1β + 150 U/ml IFNγ) for (16h) and then processed for pPERK, p65-NFκB, and loading control tubulin immunoblotting and representative blots (**G,** n=3-4) and cumulative densitometries presented (**H-I**). [a,b,c]G0 CTK group significantly from GO DMSO group, $p<0.0001$, $p=0.05$, $p=0.0046$, respectively; [d]G0 CTK group significantly different from G1+G2 CTK group, $p<0.0057$.)

Figure 8. iPLA$_2$β Associates with *Pla2g6* and *p65* Promoters in ER Stressed β-Cells. MIN6 β-cells were transfected with control (G0) or guide RNA constructs (G1 and G2) targeting *Pla2g6* (**Top panels**) or were pretreated with *S*-B̲EL (10 μM, 1h) (**Bottom panels**) and then treated with D̲MSO or C̲ytokines (50 U/ml IL-1β + 150 U/ml IFNγ) for (2h). Subsequently, ChIP was performed using antibody directed against iPLA$_2$β and acetylated histone (AcH4) followed by qPCR analyses of *Pla2g6* and *Nfκb* promoter regions. AcH4 was used to confirm transcriptional activation and IgG as a negative control. **A, D.** Fold enrichment, relative to IgG, under

basal (DMSO) conditions. **B, E.** Cytokine-induced fold enrichment, relative to IgG. **C, F.** Cytokine-induced fold enrichment, relative to DMSO: **C.** [a,b]G0 C groups significantly different from other groups, $p<0.01$, $p<0.05$, respectively; n=3.); **F.** [a,b]Cytokine groups significantly different from corresponding DMSO groups, $p<0.005$, $p<0.05$, respectively; [c]C+B significantly different from corresponding C $p<0.05$; n=3.)

Figure 9. Lipidomics analyses. Human islets (500/condition) were treated with cytokines (100 U/ml IL-1β + 300 U/ml IFNγ) for 2-36h. In some replicates, at peak response times, cytokines + *S*-BEL groups were added. A. PGE_2. B. $PGF_2\alpha$. C. 8-Iso-$PGF_2\alpha$. (Insets,[*,†]significantly different from DMSO group, $p < 0.05$ and $p < 0.01$, respectively, n=10-18. *CTK+*S*-BEL group significantly different from CTK group, $p < 0.05$, n values indicated above the bars.)

Figure 10. Impact of Select PGs β-Cell Apoptosis and ER Stress. A. MIN6 β-cells were treated with <u>D</u>MSO, <u>C</u>ytokines as a positive control (50 U/ml IL-1β + 150 U/ml IFNγ) or a PGs cocktail (10 nM PGE_2 + 10 μM $PGF_2\alpha$ + 10 μM 8-Iso-$PGF_2\alpha$) and processed for TUNEL analyses after 24h. Data are means ± SEMs of % apoptotic cells, relative to total cells. ([a]Cytokine group significantly different from DMSO group, $p < 0.0001$; [b]PGs group significantly different from DMSO group, $p = 0.004$, n=3/group). **B**. Representative immunoblotting analyses blots for p65-NFκB, pSTAT1, pPERK, CHOP, and loading control tubulin after 16h. **C-F**. Corresponding cumulative densitometries are presented. ([a,b,c]Cytokine or PGs group significantly different from DMSO groups, p, 0.05, $p = 0.0013$, and $p = 0.009$, respectively, n=3-4/group. ND=not determined.)

Figure 11. iDLs-Dependent Induction of iPLA₂β. MIN6 β-cells were treated with <u>C</u>ytokines ± inhibitors as in Fig. 4, or PGs, as in Fig. 9, and processed for iPLA₂β immunoblotting analyses. Representative blots and corresponding densitometries for CTK ± *S*- BEL (A), CTK ± FKGK18, CTK ± *R*- BEL, CTK ± *S*- BEL, and cytokines and PGs (E) are presented. Cumulate densitometries are presented in Supplemental Figures 3 and 6.

Figure 12. Select iDLs-Indued iPLA₂β Enrichment at Promoter Regions in β-Cells. MIN6 β-cells were treated with a PGs cocktail, as in Fig. 10, for 4h and processed for ChIP and qPCR analyses. **A**. Fold enrichment of iPLA₂β at promoter regions, relative to IgG. **B-F**. Fold enrichment of iPLA₂β at promoter regions, relative to DMSO: **B**, *Pla2g6*; **C**, *NfkB*; **D**, *Stat1*; **E**. *Gcg*; **F**. *Pepck*. ([a,c]PGs groups significantly different from corresponding DMSO groups, $p < 0.001$ and $p < 0.01$, respectively. [b,d]PGs + *S*-BEL groups significantly different from corresponding DMSO groups, $p < 0.01$ and $p < 0.05$, respectively.)

Figure 13. Validation of p65 or iPLA$_2$β Enrichment at Promoter Regions in Human Islets. Human islets (3500-5000/condition) from healthy donors were treated with cytokines (100 U/ml IL-1β + 300 U/ml IFNγ for 2h (**A, B**) or a PGs cocktail (**C**) as in Fig. 10 for 4h. The islets were then processed for ChIP analyses using antibodies directed against p65-NFκB or iPLA$_2$β and subsequent qPCR analyses. **D**. Human islets (250/condition) from healthy donors were treated with cytokines (100 U/ml IL-1β + 300 U/ml IFNγ or a PGs cocktail for 24h and processed for TUNEL analyses. Data are mean ± SEMs of %apoptotic cells, relative to total cells. ([a,b]Significantly different from DMSO group, p < 0.0001 and p = 0.004, respectively.)

Figure 14. Proposed Model of iDLs Contribution to T1D Development.

Author's Contributions

XL was involved with the design, experimentation, data analyses and manuscript preparation; **AKC** generated the CRISPR-Cas9 guides, **TA** contributed to NFκB immunoblotting; **SEN** contributed to the design of the ChIP experiments; **DS**, **AF**, and **CEC** performed the lipidomics mass spectrometry and data analyses; **CSH**, **ARW**, and **ESN** contributed to the transcriptional approaches; **YGT** assisted with the cell culturing and human islets experiments; and **SR** oversaw all experimental design, data analyses and preparation of the manuscript. All co-authors contributed to editing of the manuscript.

Acknowledgements

This work was supported in part by funding (to SR) from NIH/NIDDK (R01DK069455, R01DK110292), NIH/HIRN (U01 DK127786), Breakthrough T1D (3-SRA-2023-1362-S-B), Beatson Foundation (#2023-20), UAB-Department of Cell Developmental, and Integrative Biology, UAB-Comprehensive Diabetes Center, and UAB Diabetes Research Center (P30 DK079626). The contents of this manuscript do not represent the views of the Department of Veterans Affairs or the United States Government. This work was mainly supported by the National Institutes of Health via R01DK126444 (to CEC, SR), the Veteran's Administration (VA Merit Review, BX001792 (to CEC), VA Merit Review award, BX006063 (to CEC), VA Merit Review award, RD001334 (to CEC), and a Senior Research Career Scientist Award, IK6RD002739 (to CEC)). This work was also peripherally supported by way of research cores, methodology and technology development, and software/bioinformatic development by the National Institutes of Health via P01 CA171983 (to CEC), P01 CA302570 (to CEC), NIH/NCI Cancer Center Support Grant P30 CA044579, R01 AI139072 (to CEC), and R01 GM155691 (to CEC). We thank Dr. George Kokotos, University of Athens for the generous supply of FKGK18. Activity of the AIDP is approved by the Health Research Ethics Board at the University of Alberta (Approval # Pro-00001620).

Table 1. ChIP Primers

Name	Sequence (5' to 3')		DNA detected	Product Size
Mm Pla2g6 F	GGGACGAGCATATGCAAACC		*Pla2g6*	222 bp
Mm Pla2g6 R	GCCCTGGCTATCCTGGAACT			
Mm Nfκb F	TCCTGTTCGAGTCTCCATGCAG		*Rela*	513 bp
Mm Nfκb R	CATCAGCTTGCGAAAAGGAGCC			
Mm Stat1 F	GCCTCTCATTGTCACCGAAGAAC		*Stat1*	
Mm Stat1R	TGGCTGACGTTGGAGATCACCA			
Mm Pepck F	GAGTGACACCTCACAGCTGTGG		*Pepck*	
Mm Pepck R	GGCAGGCCTTTGGATCATAGCC			
Mm Gcg F	CGTAAAAAGCAGATGAGCAAAGTG			
Mm Gcg R	GAACAGGTGTAGACAGAGGGAGTCC		*Gcg*	

TGAACCGAAACTCTGGCAGCTG/CATCAGCTTGCGAAAAGGAGCC

Name	Sequence (5' to 3')		DNA detected	Product Size
HsPLA2G6 F	CTCTAGCAGCCACCCCTGAA		*PLA2G6*	391 bp
HsPLA2G6 R	CTCTAGCAGCCACCCCTGAA			
HsNFkB F	TGAACCGAAACTCTGGCAGCTG		*Rela*	513 bp
HsNFkB R	CATCAGCTTGCGAAAAGGAGCC			

References

1. Anderson, M. S., and Bluestone, J. A. (2005) The NOD mouse: a model of immune dysregulation. *Annual review of immunology* **23**, 447-485

2. Lei, X., Zhang, S., Barbour, S. E., Bohrer, A., Ford, E. L., Koizumi, A., Papa, F. R., and Ramanadham, S. (2010) Spontaneous development of endoplasmic reticulum stress that can lead to diabetes mellitus is associated with higher calcium-independent phospholipase A2 expression: a role for regulation by SREBP-1. *The Journal of biological chemistry* **285**, 6693-6705

3. Lei, X., Zhang, S., Bohrer, A., Bao, S., Song, H., and Ramanadham, S. (2007) The group VIA calcium-independent phospholipase A2 participates in ER stress-induced INS-1 insulinoma cell apoptosis by promoting ceramide generation via hydrolysis of sphingomyelins by neutral sphingomyelinase. *Biochemistry* **46**, 10170-10185

4. Lei, X., Bone, R. N., Ali, T., Wohltmann, M., Gai, Y., Goodwin, K. J., Bohrer, A. E., Turk, J., and Ramanadham, S. (2013) Genetic modulation of islet beta-cell iPLA2beta expression provides evidence for its impact on beta-cell apoptosis and autophagy. *Islets* **5**, 29-44

5. Lei, X., Bone, R. N., Ali, T., Zhang, S., Bohrer, A., Tse, H. M., Bidasee, K. R., and Ramanadham, S. (2014) Evidence of contribution of iPLA2beta-mediated events during islet beta-cell apoptosis due to proinflammatory cytokines suggests a role for iPLA2beta in T1D development. *Endocrinology* **155**, 3352-3364

6. Lei, X., Zhang, S., Bohrer, A., Barbour, S. E., and Ramanadham, S. (2012) Role of calcium-independent phospholipase A2beta in human pancreatic islet beta-cell apoptosis. *American journal of physiology. Endocrinology and metabolism* **303**, E1386-1395

7. Lei, X., Zhang, S., Bohrer, A., and Ramanadham, S. (2008) Calcium-independent phospholipase A2 (iPLA2b)-mediated ceramide generation plays a key role in the cross-talk between the endoplasmic reticulum (ER) and mitochondria during ER stress-induced insulin-secreting cell apoptosis. *J. Biol. Chem.* **283**, 34819-34832

8. Ramanadham, S., Hsu, F. F., Zhang, S., Jin, C., Bohrer, A., Song, H., Bao, S., Ma, Z., and Turk, J. (2004) Apoptosis of insulin-secreting cells induced by endoplasmic reticulum stress is amplified by overexpression of group VIA calcium-independent phospholipase A2 (iPLA2b) and suppressed by inhibition of iPLA2b. *Biochemistry* **43**, 918-930

9. Ali, T., Lei, X., Barbour, S. E., Koizumi, A., Chalfant, C. E., and Ramanadham, S. (2021) Alterations in beta-Cell Sphingolipid Profile Associated with ER Stress and iPLA(2)beta: Another Contributor to beta-Cell Apoptosis in Type 1 Diabetes. *Molecules* **26**

10. Almutairi, A., White, T. D., Stephenson, D. J., Stephenson, B. D., Gai-Tusing, Y., Goel, P., Phillips, D. W., Welner, R. S., Lei, X., Hammock, B. D., Chalfant, C. E., and Ramanadham, S. (2024) Selective Reduction of Ca2+-Independent Phospholipase A2beta (iPLA2beta)-Derived Lipid Signaling from Macrophages Mitigates Type 1 Diabetes Development. *Diabetes*

11. Bone, R. N., Gai, Y., Magrioti, V., Kokotou, M. G., Ali, T., Lei, X., Tse, H. M., Kokotos, G., and Ramanadham, S. (2015) Inhibition of Ca2+-independent phospholipase A2beta (iPLA2beta) ameliorates islet infiltration and incidence of diabetes in NOD mice. *Diabetes* **64**, 541-554

12. Nelson, A. J., Stephenson, D. J., Bone, R. N., Cardona, C. L., Park, M. A., Tusing, Y. G., Lei, X., Kokotos, G., Graves, C. L., Mathews, C. E., Kramer, J., Hessner, M. J., Chalfant, C. E., and Ramanadham, S. (2020) Lipid mediators and biomarkers associated with type 1 diabetes development. *JCI Insight* **5**

13. White, T. D., Almutairi, A., Gai-Tusing, Y., Stephenson, D. J., Stephenson, B. D., Lei, X., Lu, B., Hammock, B. D., DiLorenzo, T. P., Chalfant, C. E., and Ramanadham, S. (2024) Differential Lipid Signaling from CD4+ and CD8+ T-Cells Contributes to Type 1 Diabetes Developmen. *Frontiers Immunology* **doi.org/10.3389/fimmu.2024.1444639**

14. Araki, E., Oyadomari, S., and Mori, M. (2003) Impact of endoplasmic reticulum stress pathway on pancreatic β-cells and diabetes mellitus. *Experimental Biology and Medicine* **228**, 1213-1217

15. Delepine, M., Nicolino, M., Barrett, T., Golamaully, M., Lathrop, G. M., and Julier, C. (2000) EIF2AK3, encoding translation initiation factor 2-alpha kinase 3, is mutated in patients with Wolcott-Rallison syndrome. *Nat Genet* **25**, 406

16. Iwawaki, T., and Oikawa, D. (2013) The role of the unfolded protein response in diabetes mellitus. *Seminars in immunopathology* **35**, 333-350

17. Butler, A. E., Janson, J., Bonner-Weir, S., Ritzel, R., Rizza, R. A., and Butler, P. C. (2003) Beta-cell deficit and increased beta-cell apoptosis in humans with type 2 diabetes. *Diabetes* **52**, 102-110

18. Butler, A. E., Janson, J., Soeller, W. C., and Butler, P. C. (2003) Increased beta-cell apoptosis prevents adaptive increase in beta-cell mass in mouse model of type 2 diabetes: evidence for role of islet amyloid formation rather than direct action of amyloid. *Diabetes* **52**, 2304-2314

19. Kayo, T., and Koizumi, A. (1988) Mapping of murine diabetogenic gene mody on chromosome 7 at D7Mit258 and its involvement in pancreatic islet and beta cell development during the perinatal period. *J Clin Invest* **101**, 2112-2118

20. Lei, X., Barbour, S. E., and Ramanadham, S. (2010) Group VIA Ca2+-independent phospholipase A2 (iPLA2β) and its role in β-cell programmed cell death. *Biochimie* **92**, 627-637

21. Mandrup-Poulsen, T. (2001) Beta-cell apoptosis: stimuli and signaling. *Diabetes* **50**, S58-63

22. Oyadomari, S., Koizumi, A., Takeda, K., Gotoh, T., Akira, S., Araki, E., and Mori, M. (2002) Targeted disruption of the Chop gene delays endoplasmic reticulum stress-mediated diabetes. *J. Clin. Invest.* **109**, 525-532

23. Yamada, T., Ishihara, H., Tamura, A., Takahashi, R., Yamaguchi, S., Takei, D., Tokita, A., Satake, C., Tashiro, F., Katagiri, H., Aburatani, H., Miyazaki, J.-i., and Oka, Y. (2006) WFS1-deficiency increases endoplasmic reticulum stress, impairs cell cycle progression and triggers the apoptotic pathway specifically in pancreatic {beta}-cells. *Hum. Mol. Genet.* **15**, 1600-1609

24. Yoshioka, M., Kayo, T., Ikeda, T., and Koizumi, A. (1997) A novel locus, Mody4, distal to D7Mit189 on chromosome 7 determines early-onset NIDDM in nonobese C57BL/6 (Akita) mutant mice. *Diabetes* **46**, 887-894

25. Rabinovitch, A., and Suarez-Pinzon, W. L. (1998) Cytokines and their roles in pancreatic islet beta-cell destruction and insulin-dependent diabetes mellitus. *Biochem Pharmacol* **55**, 1139-1149

26. Sesti, G. (2002) Apoptosis in the beta cells: cause or consequence of insulin secretion defect in diabetes? *Ann Med* **34**, 444-450

27. Socha, L., Silva, D., Lesage, S., Goodnow, C., and Petrovsky, N. (2003) The role of endoplasmic reticulum stress in nonimmune diabetes: NOD.k iHEL, a novel model of beta cell death. *Annals of the New York Academy of Sciences* **1005**, 178-183

28. Oyadomari, S., Araki, E., and Mori, M. (2002) Endoplasmic reticulum stress-mediated apoptosis in pancreatic beta-cells. *Apoptosis : an international journal on programmed cell death* **7**, 335-345

29. Tersey, S. A., Nishiki, Y., Templin, A. T., Cabrera, S. M., Stull, N. D., Colvin, S. C., Evans-Molina, C., Rickus, J. L., Maier, B., and Mirmira, R. G. (2012) Islet beta-cell endoplasmic reticulum stress precedes the onset of type 1 diabetes in the nonobese diabetic mouse model. *Diabetes* **61**, 818-827

30. Lei, X., Zhang, S., Bohrer, A., Barbour, S. E., and Ramanadham, S. (2012) Role of calcium-independent phospholipase A(2)beta in human pancreatic islet beta-cell apoptosis. *American journal of physiology. Endocrinology and metabolism* **303**, E1386-1395

31. Gijon, M. A., and Leslie, C. C. (1997) Phospholipases A2. *Semin Cell Dev Biol* **8**, 297-303

32. Gross, R. W., Ramanadham, S., Kruszka, K. K., Han, X., and Turk, J. (1993) Rat and human pancreatic islet cells contain a calcium ion independent phospholipase A2 activity selective for hydrolysis of arachidonate which is stimulated by adenosine triphosphate and is specifically localized to islet beta-cells. *Biochemistry* **32**, 327-336

33. Lei, X., Bone, R. N., Ali, T., Wohltmann, M., Gai, Y., Goodwin, K. J., Bohrer, A. E., Turk, J., and Ramanadham, S. (2013) Genetic modulation of islet beta-cell iPLA(2)beta expression provides evidence for its impact on beta-cell apoptosis and autophagy. *Islets* **5**, 29-44

34. Lei, X., Bone, R. N., Ali, T., Zhang, S., Bohrer, A., Tse, H. M., Bidasee, K. R., and Ramanadham, S. (2014) Evidence of Contribution of iPLAbeta-Mediated Events During Islet beta-Cell Apoptosis Due to

Proinflammatory Cytokines Suggests a Role for iPLAbeta in T1D Development. *Endocrinology*, en20132134

35. Lei, X., Zhang, S., Bohrer, A., Bao, S., Song, H., and Ramanadham, S. (2007) The group VIA calcium-independent phospholipase A2 participates in ER stress-induced INS-1 insulinoma cell apoptosis by promoting ceramide generation via hydrolysis of sphingomyelins by neutral sphingomyelinase. *Biochemistry* **46**, 10170-10185

36. Lei, X., Zhang, S., Bohrer, A., and Ramanadham, S. (2008) Calcium-independent phospholipase A2 (iPLA2 beta)-mediated ceramide generation plays a key role in the cross-talk between the endoplasmic reticulum (ER) and mitochondria during ER stress-induced insulin-secreting cell apoptosis. *The Journal of biological chemistry* **283**, 34819-34832

37. Ramanadham, S., Hsu, F. F., Zhang, S., Jin, C., Bohrer, A., Song, H., Bao, S., Ma, Z., and Turk, J. (2004) Apoptosis of Insulin-Secreting Cells Induced by Endoplasmic Reticulum Stress Is Amplified by Overexpression of Group VIA Calcium-Independent Phospholipase A2 (iPLA2β) and Suppressed by Inhibition of iPLA2β. *Biochemistry* **43**, 918-930

38. Luo, P., and Wang, M. H. (2011) Eicosanoids, beta-cell function, and diabetes. *Prostaglandins & other lipid mediators* **95**, 1-10

39. Inceoglu, B., Bettaieb, A., Haj, F. G., Gomes, A. V., and Hammock, B. D. (2017) Modulation of mitochondrial dysfunction and endoplasmic reticulum stress are key mechanisms for the wide-ranging actions of epoxy fatty acids and soluble epoxide hydrolase inhibitors. *Prostaglandins & other lipid mediators* **133**, 68-78

40. Schwemer, T. F., Deutscher, N., Diermann, N., Boger, R., Schwedhelm, E., Blankenberg, S., and Friedrich, F. W. (2019) Effect of ranolazine on plasma arginine derivatives and urinary isoprostane 8-iso-PGF(2alpha) in patients with myocardial infarction in the randomized RIMINI-Trial. *Sci Rep* **9**, 5708

41. Chen, M., Yang, Z. D., Smith, K. M., Carter, J. D., and Nadler, J. L. (2005) Activation of 12-lipoxygenase in proinflammatory cytokine-mediated beta cell toxicity. *Diabetologia* **48**, 486-495

42. Chopra, S., Giovanelli, P., Alvarado-Vazquez, P. A., Alonso, S., Song, M., Sandoval, T. A., Chae, C. S., Tan, C., Fonseca, M. M., Gutierrez, S., Jimenez, L., Subbaramaiah, K., Iwawaki, T., Kingsley, P. J., Marnett, L. J., Kossenkov, A. V., Crespo, M. S., Dannenberg, A. J., Glimcher, L. H., Romero-Sandoval, E. A., and Cubillos-Ruiz, J. R. (2019) IRE1alpha-XBP1 signaling in leukocytes controls prostaglandin biosynthesis and pain. *Science* **365**

43. Wagner, K. M., McReynolds, C. B., Schmidt, W. K., and Hammock, B. D. (2017) Soluble epoxide hydrolase as a therapeutic target for pain, inflammatory and neurodegenerative diseases. *Pharmacol Ther* **180**, 62-76

44. Cho, K. N., Choi, J. Y., Kim, C. H., Baek, S. J., Chung, K. C., Moon, U. Y., Kim, K. S., Lee, W. J., Koo, J. S., and Yoon, J. H. (2005) Prostaglandin E2 induces MUC8 gene expression via a mechanism involving ERK MAPK/RSK1/cAMP response element binding protein activation in human airway epithelial cells. *The Journal of biological chemistry* **280**, 6676-6681

45. Gao, Y., Zhao, C., Wang, W., Jin, R., Li, Q., Ge, Q., Guan, Y., and Zhang, Y. (2016) Prostaglandins E2 signal mediated by receptor subtype EP2 promotes IgE production in vivo and contributes to asthma development. *Sci Rep* **6**, 20505

46. Faour, W. H., Alaaeddine, N., Mancini, A., He, Q. W., Jovanovic, D., and Di Battista, J. A. (2005) Early growth response factor-1 mediates prostaglandin E2-dependent transcriptional suppression of cytokine-induced tumor necrosis factor-alpha gene expression in human macrophages and rheumatoid arthritis-affected synovial fibroblasts. *The Journal of biological chemistry* **280**, 9536-9546

47. Aoki, T., Frosen, J., Fukuda, M., Bando, K., Shioi, G., Tsuji, K., Ollikainen, E., Nozaki, K., Laakkonen, J., and Narumiya, S. (2017) Prostaglandin E2-EP2-NF-kappaB signaling in macrophages as a potential therapeutic target for intracranial aneurysms. *Sci Signal* **10**

48. Chen, X., Shi, C., He, M., Xiong, S., and Xia, X. (2023) Endoplasmic reticulum stress: molecular mechanism and therapeutic targets. *Signal Transduct Target Ther* **8**, 352

49. Schmitz, M. L., Shaban, M. S., Albert, B. V., Gokcen, A., and Kracht, M. (2018) The Crosstalk of Endoplasmic Reticulum (ER) Stress Pathways with NF-kappaB: Complex Mechanisms Relevant for Cancer, Inflammation and Infection. *Biomedicines* **6**

50. Sahin, G. S., Lee, H., and Engin, F. (2021) An accomplice more than a mere victim: The impact of beta-cell ER stress on type 1 diabetes pathogenesis. *Mol Metab* **54**, 101365

51. Nozell, S., Laver, T., Moseley, D., Nowoslawski, L., De Vos, M., Atkinson, G. P., Harrison, K., Nabors, L. B., and Benveniste, E. N. (2008) The ING4 tumor suppressor attenuates NF-kappaB activity at the promoters of target genes. *Molecular and cellular biology* **28**, 6632-6645

52. Nozell, S., Laver, T., Patel, K., and Benveniste, E. N. (2006) Mechanism of IFN-beta-mediated inhibition of IL-8 gene expression in astroglioma cells. *J Immunol* **177**, 822-830

53. Kin, T., O'Gorman, D., Schroeder, A., Onderka, C., Richer, B., Rosichuk, S., Zhai, X., and Shapiro, A. M. (2011) Human islet distribution program for basic research at a single center. *Transplant Proc* **43**, 3195-3197

54. Kin, T., O'Gorman, D., Zhai, W., Moriarty, J., Park, K., Ganguly, A., Rosichuk, S., and Shapiro, A. M. J. (2024) Contribution of a Single Islet Transplant Program to Basic Researchers in North America, Europe, and Asia through Distributing Human Islets. *OBM Transplantation* **08**, 212

55. Jenkins, C. M., Han, X., Mancuso, D. J., and Gross, R. W. (2002) Identification of calcium-independent phospholipase A2 (iPLA2)beta, and not iPLA2gamma, as the mediator of arginine vasopressin-induced arachidonic acid release in A-10 smooth muscle cells. Enantioselective mechanism-based discrimination of mammalian iPLA2s. *J. Biol. Chem.* **277**, 32807-32814

56. Ali, T., Kokotos, G., Magrioti, V., Bone, R. N., Mobley, J. A., Hancock, W., and Ramanadham, S. (2013) Characterization of FKGK18 as inhibitor of group VIA Ca2+-independent phospholipase A2 (iPLA2beta): candidate drug for preventing beta-cell apoptosis and diabetes. *PloS one* **8**, e71748

57. Kokotos, G., Hsu, Y. H., Burke, J. E., Baskakis, C., Kokotos, C. G., Magrioti, V., and Dennis, E. A. (2010) Potent and selective fluoroketone inhibitors of group VIA calcium-independent phospholipase A2. *Journal of medicinal chemistry* **53**, 3602-3610

58. Ran, F. A., Hsu, P. D., Wright, J., Agarwala, V., Scott, D. A., and Zhang, F. (2013) Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* **8**, 2281-2308

59. Chang, H. H. Y., Pannunzio, N. R., Adachi, N., and Lieber, M. R. (2017) Non-homologous DNA end joining and alternative pathways to double-strand break repair. *Nat Rev Mol Cell Biol* **18**, 495-506

60. Bravo-Ruiz, I., Medina, M. A., and Martinez-Poveda, B. (2021) From Food to Genes: Transcriptional Regulation of Metabolism by Lipids and Carbohydrates. *Nutrients* **13**

61. Marhfour, I., Lopez, X. M., Lefkaditis, D., Salmon, I., Allagnat, F., Richardson, S. J., Morgan, N. G., and Eizirik, D. L. (2012) Expression of endoplasmic reticulum stress markers in the islets of patients with type 1 diabetes. *Diabetologia* **55**, 2417-2420

62. Ramanadham, S., Hsu, F. F., Bohrer, A., Nowatzke, W., Ma, Z., and Turk, J. (1998) Electrospray ionization mass spectrometric analyses of phospholipids from rat and human pancreatic islets and subcellular membranes: comparison to other tissues and implications for membrane fusion in insulin exocytosis. *Biochemistry* **37**, 4553-4567

63. Oakes, S. A., and Papa, F. R. (2015) The role of endoplasmic reticulum stress in human pathology. *Annual review of pathology* **10**, 173-194

64. Papa, F. R. (2012) Endoplasmic reticulum stress, pancreatic beta-cell degeneration, and diabetes. *Cold Spring Harbor perspectives in medicine* **2**, a007666

65. Cardozo, A. K., Heimberg, H., Heremans, Y., Leeman, R., Kutlu, B., Kruhoffer, M., Orntoft, T., and Eizirik, D. L. (2001) A comprehensive analysis of cytokine-induced and nuclear factor-kappa B-dependent genes in primary rat pancreatic beta-cells. *J Biol Chem* **276**, 48879-48886

66. Cnop, M., Welsh, N., Jonas, J. C., Jorns, A., Lenzen, S., and Eizirik, D. L. (2005) Mechanisms of pancreatic beta-cell death in type 1 and type 2 diabetes: many differences, few similarities. *Diabetes* **54 Suppl 2**, S97-107

67. Eldor, R., Yeffet, A., Baum, K., Doviner, V., Amar, D., Ben-Neriah, Y., Christofori, G., Peled, A., Carel, J. C., Boitard, C., Klein, T., Serup, P., Eizirik, D. L., and Melloul, D. (2006) Conditional and specific NF-

kappaB blockade protects pancreatic beta cells from diabetogenic agents. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 5072-5077

68. Karin, M., and Lin, A. (2002) NF-kappaB at the crossroads of life and death. *Nature immunology* **3**, 221-227

69. Carrero, J. A., Calderon, B., Towfic, F., Artyomov, M. N., and Unanue, E. R. (2013) Defining the transcriptional and cellular landscape of type 1 diabetes in the NOD mouse. *PloS one* **8**, e59701

70. Lamhamedi-Cherradi, S. E., Zheng, S., Hilliard, B. A., Xu, L., Sun, J., Alsheadat, S., Liou, H. C., and Chen, Y. H. (2003) Transcriptional regulation of type I diabetes by NF-kappa B. *J Immunol* **171**, 4886-4892

71. Ortis, F., Pirot, P., Naamane, N., Kreins, A. Y., Rasschaert, J., Moore, F., Theatre, E., Verhaeghe, C., Magnusson, N. E., Chariot, A., Orntoft, T. F., and Eizirik, D. L. (2008) Induction of nuclear factor-kappaB and its downstream genes by TNF-alpha and IL-1beta has a pro-apoptotic role in pancreatic beta cells. *Diabetologia* **51**, 1213-1225

72. Papaccio, G., Graziano, A., D'Aquino, R., Valiante, S., and Naro, F. (2005) A biphasic role of nuclear transcription factor (NF)-kappaB in the islet beta-cell apoptosis induced by interleukin (IL)-1beta. *Journal of cellular physiology* **204**, 124-130

73. Salem, H. H., Trojanowski, B., Fiedler, K., Maier, H. J., Schirmbeck, R., Wagner, M., Boehm, B. O., Wirth, T., and Baumann, B. (2014) Long-term IKK2/NF-kappaB signaling in pancreatic beta-cells induces immune-mediated diabetes. *Diabetes* **63**, 960-975

74. Zhao, Y., Krishnamurthy, B., Mollah, Z. U., Kay, T. W., and Thomas, H. E. (2011) NF-kappaB in type 1 diabetes. *Inflammation & allergy drug targets* **10**, 208-217

75. Ovcharenko, I., Nobrega, M. A., Loots, G. G., and Stubbs, L. (2004) ECR Browser: a tool for visualizing and accessing data from comparisons of multiple vertebrate genomes. *Nucleic acids research* **32**, W280-286

76. Kliewer, S. A., Sundseth, S. S., Jones, S. A., Brown, P. J., Wisely, G. B., Koble, C. S., Devchand, P., Wahli, W., Willson, T. M., Lenhard, J. M., and Lehmann, J. M. (1997) Fatty acids and eicosanoids regulate gene expression through direct interactions with peroxisome proliferator-activated receptors alpha and gamma. *Proceedings of the National Academy of Sciences of the United States of America* **94**, 4318-4323

77. Sheppe, A. E. F., and Edelmann, M. J. (2021) Roles of Eicosanoids in Regulating Inflammation and Neutrophil Migration as an Innate Host Response to Bacterial Infections. *Infect Immun* **89**, e0009521

78. Wang, X., Han, X., and Powell, C. A. (2022) Lipids and genes: Regulatory roles of lipids in RNA expression. *Clin Transl Med* **12**, e863

79. Ramanadham, S., Hsu, F. F., Bohrer, A., Ma, Z., and Turk, J. (1999) Studies of the role of group VI phospholipase A2 in fatty acid incorporation, phospholipid remodeling, lysophosphatidylcholine generation, and secretagogue-induced arachidonic acid release in pancreatic islets and insulinoma cells. *The Journal of biological chemistry* **274**, 13915-13927

80. Ramanadham, S., Song, H., Bao, S., Hsu, F. F., Zhang, S., Ma, Z., Jin, C., and Turk, J. (2004) Islet complex lipids: involvement in the actions of group VIA calcium-independent phospholipase A(2) in beta-cells. *Diabetes* **53 Suppl 1**, S179-185

Supplemental Figures

Supplemental Figure 1. Impact of ER Stress, NFκB, and STAT1 on iPLA₂β and Stress Factors. MIN-6 β-cells were treated with cytokines, as in **Figs. 1 and 2** in the absence or presence of PBA, BAY 11, or AZD for 16h. Cytosols were then prepared and processed for immunoblotting analyses for p65-NFκb, iPLA₂β, pSTAT1, pPERK, and CHOP. (Cytokine groups significantly different from corresponding DMSO groups, [a]$p < 0.0001$ and [b]$p = 0.0003$; [c,d,*,e]Cytokine + intervention groups significantly different from corresponding cytokine groups, $p < 0.05$, $p < 0.01$, $p < 0.05$ (1-tailed), and $p < 0.005$, respectively.

Supplemental Figure 2. Time-Course of iPLA₂β Association with *Pla2g6* Promoter Region. INS-1 β-cells were treated with DMSO or Cytokines, as in **Suppl. Fig. 1**, for 2-8h and then processed for ChIP analyses using select antibodies for iPLA₂β and AcH4 and subsequent qPCR analyses for *Pla2g6* promoter region.

Supplemental Figure 3. Select Targeting of iPLA₂β Reduces Stress Responses. MIN-6 β-cells were treated with DMSO or cytokines for 24h, as in **Suppl. Fig. 1**, in the absence or presence of DMSO, PBA, *S*-BEL, FKGK18, *R*-BEL, Cay10502, scrambled RNA, or siRNA targeting *Pla2g6*. Cytosols were then prepared and processed for immunoblotting analyses for p65-NFκB, iPLA₂β, pSTAT1, pPERK, and CHOP, and IκBα immunoblotting analyses. (Significance differences are presented as inserts.)

Supplemental Figure 4. Concentration-Dependent Induction of β-cells by Select Prostaglandins. MIN-6 β-cells were treated with DMSO, Cytokines, or PGs (10 nM PGE₂, 10 μM PGF₂α, 10 μM 8-Iso-PGF₂α) for 24h and processed for TUNEL. The mean ± SEMs of %apoptotic cells, relative to total cells, are presented. ([a-e]Significantly different from DMSO group, $p < 0.05$ - $p < 0.001$.)

Supplemental Figure 5. Time-Course of Select PGs-Induced Enrichment of iPLA₂β at Promoter Regions. MIN6 β-cells were treated with DMSO or PGs cocktail (10 nM PGE₂ + 10 μM PGF₂α + 10 μM 8-Iso-PGF₂α) for 2-8h. The cells were then processed for ChIP analyses using select antibodies for iPLA₂β, and subsequently for qPCR analyses of *Pla2g6, Nfkb, Stat1, Gcg,* and *Pepck* promoter regions.

Supplemental Figure 6. Impact of Select PGs on iPLA₂β Expression. MIN6 β-cells were treated with DMSO or PGs cocktail for 16h and then processed for iPLA₂β immunoblotting analyses. ([†]Significantly different from D, $p < 0.05$, 1-tailed test, n=4/group.)

Supplemental Figure 7. Time-Course of Cytokines-Induced Enrichment of p65-NFKB at Promoter Regions in Human Islets. Human islets from health donors (3500-5000) were treated with <u>D</u>MSO or <u>C</u>ytokines (100 U/ml IL-1β + 300 U/ml IFNγ) for 4 and 8h. The islets were then processed for ChIP analyses using select antibodies for p65-NFkB and subsequent qPCR analyses for *PLA2G6* and *GcG* promoter regions. Enrichment relative to IgG (**A**) and DMSO (**B**) are presented.

...s-Induced β-Cell Apoptosis is Linked to Novel Lipid Signaling at the Transcriptional Level: Implications...
Development

...ong Lei, [1,2]Anil K. Challa, [3]Susan E. Nozell, [1,2]Tomader Ali, [4]Daniel J. Stephenson, [4]Andrew Nafzinger, [2,5,6]...
[2,7]Adam R. Wende, [8]Ernesto S. Nakayasu, [1,2]Ying Gai-Tusing, [4,9,10]Charles E. Chalfant, [1,2]Sasanka Ramana...

Main Text Figures 1-14

Figure 1. NFκB Activation During ER Stress-Induced β-Cell Apoptosis is Associated with iPLA$_2$β Induc

A. TUNEL

B. NFκB, iPLA$_2$β, and ER Stress Factors

iPLA$_2$β
pPERK
CHOP
Tub
D P C C+P
p65
Oct-1
D P C C+P

24 16 24 16 24 16 24 (h)

C. iPLA$_2$β
Fold Rel. to Tub
D P C C+P

D. pPERK
Fold Rel. to Tub
D P C C+P

E. CHOP
Fold Rel. to Tub
D P C C+P

F. NFκB Induction by Cytokines
NUC
CYTO
Tub
0 2 8 16 24 48h

G. NFκB ± Cytokines
Fold-Ch. (Rel. to 0h)
☐ Nuc
☐ Cyto
0 0 2 2 8 8 16 16 24 24 48 48
IL-1β + IFNγ

H. p
Fold Rel. to Tub
D

Figure 2. Inhibition of NFκB and STAT1 Reduces ER Stress-Mediated iPLA₂β Expression and β-Cell Apoptosis

Figure 3. <u>Transcriptional Induction of *Pla2g6* by NFκB and STAT1</u>

A. <u>*IL-6* Promoter</u>

B. <u>*Pla2g6* Promoter</u>

Figure 4. <u>Selective</u> Impact of iPLA₂β Inhibition on NFκB Activaton in ER Stressed β-Cells

Figure 5. <u>Knockdown of iPLA$_2$β Reduces ER Stress and NFκB and pSTAT Activation in β-Cells</u>

B-G. <u>Densitometries</u>

A. <u>iPLA$_2$β siRNA-Induced Effects</u>

iPLA$_2$β

p65

pPERK

CHOP

Tub

pSTAT1

Tub

IKBα

Tub

D   scr   C   C+si

B. <u>iPLA$_2$β</u>

C. <u>p65</u>

D. IκBα

F. <u>pSTAT1</u>

F. <u>pPERK</u>

G. <u>CHOP</u>

Figure 6. Select Targeting iPLA$_2$β Inhibits ER Stress-Induced β-Cell Apoptosis

Figure 7. CRISPR-Cas9 Approach to Examine iPLA$_2$β Role in Transcription

A. *Pla2g6* CRISPR-Cas9 Guide RNAs

1 (minus strand)	CGCTACTGAGGGTGTTGACG AGG
2 (plus strand)	GAATGTCTCCAATCGCACCT GGG

B. Genotyping by PCR-PAGE

C. Predicted Deletion Region

137 bp deletion

D. iPLA$_2$β Expression

G0 G0 G1 G1 G2 G2 G1+2 G1+2
- + - + - + - +

E. iPLA$_2$β

F. β-Cell Apoptosis

CRISPR-Cas9 Reduction of *Pla2g6* Mitgates ER Stress in β-Cells

H. p65

I. pPERK

Fig. 8. iPLA$_2$β Enrichment at *Pla2g6* and *p65* in ER-Stresed β-cells: CRISPR vs. *S*-BEL Approache

iPLA$_2$β Enrichment at Promoter Regions in ER-Stresed β-Cells ± gRNAs

sal iPLA$_2$β Enrichment Relative to IgG   **B. Induced iPLA$_2$β Enrichment Relative to IgG**   **C. Fold iPLA$_2$β Enrichment Relative to G0 D**

iPLA$_2$β Enrichment at Promoter Regions in ER-Stresed β-Cells ± *S*-BEL

sal iPLA$_2$β Enrichment Relative to IgG   **E. Induced iPLA$_2$β Enrichment Relative to IgG**   **F. Fold iPLA$_2$β Enrichment Relative to G0 DM**

Figure 9. Cytokines-Induced Select Prostaglandins Production in Human Islets

Figure 10. Impact of Select PGs on β-Cell Apoptosis and ER Stress

A. PGs-Induced β-Cell Apoptosis

Induce ER Stress

Figure 11. <u>iDLs-Dependent Induction of iPLA$_2$β</u>

A. <u>CTK ± S-BEL</u>

LA$_2$β

D   S-B   C   C+S-B

B. <u>CTK ± FKGK18</u>

D   FK   C   C+FK

C. <u>CTK ± R-BEL</u>

D   SR-B   C   C+R-B

D. <u>CTK ± Cay</u>

D   Cay   C   C+Cay

E. <u>PGs Induce iPLA$_2$β</u>

iPLA$_2$β

Tub

D   C   PGs

Figure 12. Select iDLs-Indued iPLA₂β Enrichment at Promoter Regions in β-cells

Figure 13. Validation of p65 or iPLA₂β Enrichment at Promoter Regions in Human Islets

A. Enrichment of p65 at Promoter Regions

B. Cytokines-Induced Enrichment of iPLA₂β at Promoter Regions

C. Select Prostgandins-Induced Enrichment of iPLA₂β at Promoter Regions and Induce β-cell Apoptosis

D. Apoptosis

Figure 14. Select iDLs Confer an iPLA$_2$β Transcriptional Role in Stressed β-cells

...ong Lei, [1,2]Anil K. Challa, [3]Susan E. Nozell, [1,2]Tomader Ali, [4]Daniel J. Stephenson, [4]Andrew Nafzinger, [2,5,6]...

[2,7]Adam R. Wende, [8]Ernesto S. Nakayasu, [1,2]Ying Gai-Tusing, [4,9,10]Charles E. Chalfant, [1,2]Sasanka Ramana...

Supplemental Figures 1-7

Supplemental Figure 1. <u>Impact of ER Stress, NFκB, and STAT1 on iPLA₂β and Stress Factors</u>

Supplemental Figure 2. Time-Course of iPLA$_2\beta$ Association with *Pla2g6* and *AcH4* Promoter Regions

A. *Pla2g6*

B. *AcH4*

Supplemental Figure 3. <u>Select Targeting of iPLA$_2$β Reduces Stress Responses</u>

Supplemental Figure 4. <u>Concentration-Dependent Induction of β-cells by Select Prostaglandins</u>

pplemental Figure 5. <u>Time-Course of Select PGs-Induced Enrichment of iPLA$_2$β at Promoter Region</u>

A. *Pla2g6*

B. *Nfκb*

C. *Stat1*

D. *Gcg*

E. *Pepck*

Supplemental Figure 6. <u>Impact of Select PGs on iPLA$_2$β Expression</u>

Supplemental Figure 7. Time-Course of Select Cytokines-Induced Enrichment of p65-NFκB at Promoter Regions in Human Islets