Preclinical validation of AAV9-TECPR2 gene therapy in a novel knock-in model of TECPR2-related disorder

Bruna Lenfers Turnes[1], Peter Casey-Caplan[1], Leo Mejia[1], Tiffany Berry[1], Julie Zhao[1], Francesco Villa[1], Emma Cropper[1], Maryam Arab[2], Biyao Zhang[1], Darred Surin[3], Darius Ebrahimi-Fakhari[1,2], Silmara de Lima[4], Alan S. Kopin[6], Nick Andrews[5], Nathaniel Hodgson[1], Michela Fagiolini*[1]

Affiliations:

[1] F.M. Kirby Neurobiology Center & Department of Neurology, Boston Children's Hospital, Harvard Medical School, Boston, MA, USA

[2]Harvard Medical School, Boston, MA, USA

[3]Department of Biology, Boston University, Boston, MA, USA

[4]Department of Ophthalmology, University of Pittsburg, Pittsburgh, PA, USA

[5]In Vivo Scientific Services, Salk Institute for Biological Studies, La Jolla, CA, USA.

[6]Tufts University School of Medicine, Boston, MA, USA

*Corresponding author: Michela Fagiolini, Michela.Fagiolini@childrens.harvard.edu

One-Sentence Key Message

TECPR2 deficiency disrupts brainstem sensory–motor circuits, impairing autophagy and tactile, gait, and startle function and is prevented by neonatal AAV9.

Abstract

TECPR2-related disorder is a rare, autosomal recessive neurodevelopmental and neurodegenerative disease characterized by early-onset motor dysfunction, sensory- and autonomic neuropathy, and progressive neurological decline with early mortality. Currently, there are no effective treatments for individuals affected by this debilitating condition. To advance our understanding of disease mechanisms and explore therapeutic strategies, we developed and then characterized a knock-in (KI) mouse model carrying the human *TECPR2* c.1319delC frameshift mutation.

TECPR2-KI mice exhibit a subset of disease-relevant phenotypes, most prominently abnormal
gait, along with reduced body weight and altered tactile sensitivity. We additionally observe a
reduction in acoustic startle responses, consistent with dysfunction of brainstem-associated
sensorimotor pathways. Histopathological analyses reveal progressive accumulation of axonal
spheroids in the dorsal column nuclei, together with abnormalities in autophagy-related markers,
features previously reported in individuals with *TECPR2*-related disorder.

To assess the therapeutic potential of gene replacement, we delivered *TECPR2* via intracisternal
infusion of AAV9/*TECPR2* in neonatal KI mice. Gene therapy restored mechanosensory function,
normalized gait and startle responses, maintain autophagic homeostasis, and partially reduced
axonal pathology. These findings demonstrate that *TECPR2*-associated deficits are not only
replicable in this new mouse model but are also amenable to postnatal intervention.

Our study introduces a genetically accurate murine model of TECPR2 deficiency, identifies
brainstem-associated phenotypes, and provides preliminary evidence supporting the feasibility of
AAV9-mediated TECPR2 gene delivery, establishing a foundation for future translational research
in a currently untreatable disease.

46
47

INTRODUCTION

TECPR2-related disorder is a devastating, ultra-rare neurodevelopmental and neurodegenerative
condition for which no disease-modifying therapies currently exist. It is classified along the
spectrum of the hereditary sensory and autonomic neuropathies (specifically HSAN9, OMIM
#615031) and the complex hereditary spastic paraplegias (HSPs). Since its first identification in
2012, fewer than 40 individuals have been reported worldwide, underscoring both the rarity and
the limited clinical awareness of the disorder (1-5).

TECPR2-related disorder is caused by biallelic loss-of-function mutations in the *TECPR2* gene,
which encodes the tectonin beta-propeller repeat-containing protein 2, a multifunctional scaffold
protein with broad expression across the central nervous system (CNS). TECPR2 has been
implicated as a key regulator of the autophagy–lysosome pathway, particularly in mediating the
targeting of autophagosomes to lysosomes and promoting autophagic flux (6). Efficient autophagic
clearance is essential for maintaining neuronal health, especially in long-projecting neurons that
depend on the constant turnover of organelles and proteins to preserve axonal integrity (7,8).

Accordingly, dysfunction of autophagy-related genes has been increasingly associated with a wide

range of neurodevelopmental and neurodegenerative disorders (9-12). Loss of TECPR2 function

results in impaired autolysosomal fusion, accumulation of protein aggregates, and formation of

neuroaxonal spheroids, ultimately leading to axonal dystrophy and neuronal loss (6).

Pathological spheroids have been documented not only in affected human patients but also in other

species, highlighting the conserved nature of TECPR2-deficiency related axonopathy. For

instance, Spanish Water Dogs with a spontaneous *TECPR2* loss-of-function mutations display

progressive axonal degeneration, and a CRISPR-Cas9-generated *Tecpr2* knockout mouse model

exhibits comparable neuropathological features (11,5). These findings reinforce the importance of

developing robust and genetically faithful animal models to study disease mechanisms and

evaluate potential interventions.

Clinically, *TECPR2*-related disorder manifests with global developmental delay and later

intellectual disability, early-onset hypotonia, sensory and autonomic neuropathy, progressive

spasticity, and progressive bulbar dysfunction with dysphagia and central apneas. Dysfunction of

brainstem-mediated circuits, contributing to dysphagia, impaired reflexes, and central apneas,

accounts for significant morbidity and early mortality. Most affected individuals succumb to

disease in the first or second decade of life, and survivors experience profound, lifelong disability

(1,3,5). Despite the recognition of these hallmark features, the precise temporal progression of

neuropathological changes and their mechanistic underpinnings remain poorly defined,

representing a major barrier to therapeutic development.

Efforts to identify therapeutic strategies for *TECPR2*-related disorders have been hindered by the

rarity of the condition and the lack of suitable *in vivo* models that capture disease-relevant

mutations and phenotypes. Although a recent study demonstrated that antisense oligonucleotides

(ASOs) targeting exon 8 of TECPR2 produced promising effects *in vitro* (13), the development of

candidate therapeutics for *TECPR2*-related disorders, genetic approaches (e.g. ASO, gene therapy)

or otherwise, will require *in vivo* models to test the efficacy of the approach.

Given these limitations, gene replacement therapy using adeno-associated virus serotype 9 (AAV9)

has emerged as a potential alternative. AAV9 vectors are well characterized for their ability to

transduce both neurons and glia across the CNS following intracisternal, intrathecal or intravenous

delivery, and they have been successfully used in the treatment of other pediatric neurological

disorders, including spinal muscular atrophy (14). Importantly, this approach allows for full-length

TECPR2 gene delivery irrespective of the specific mutation, offering a mutation-agnostic strategy

for therapeutic intervention.

To address the critical gap in preclinical models, we generated a knock-in (KI) mouse line carrying

the human *TECPR2* c.1319delC frameshift mutation. This model enables the assessment of

behavioral, sensory, and neuropathological alterations associated with TECPR2 deficiency and

was used to evaluate intracisternal AAV9-mediated *TECPR2* gene delivery at neonatal stages.

Together, this work provides (i) a genetically accurate *in vivo* model for investigating *TECPR2*-

related pathophysiology, (ii) insight into previously unrecognized sensorimotor deficits involving

brainstem circuits, and (iii) initial preclinical evidence supporting the feasibility of *TECPR2* gene

replacement.

RESULTS

Behavioral alterations in *TECPR2*-KI mice recapitulate aspects of the clinical phenotype

observed in *TECPR2*-related disorder

To model human *TECPR2*-associated disease, we generated a homozygous *Tecpr2* c.1319delC

knock-in (KI) mouse line. This mutation introduces a premature stop codon at the same position

as the pathogenic c.1319delT founder variant identified in a large cohort of patients (5) thereby

recapitulating the genetic basis of disease. We performed longitudinal behavioral phenotyping at

postnatal days 30, 60, and 90 to assess developmental and progressive manifestations.

At P90, *TECPR2*-KI mice exhibited a small significant reduction in body weight compared with

WT littermates (Fig. 1A), consistent with reports of growth impairment in some individuals with

TECPR2-related disorder (5). We examined their self-grooming behavior over a 10-minute video

recording with automatic classification of behaviors and observed that TECPR2-KI mice show

increased grooming behavior at P90 (Fig. 1B). Given reports of sensory abnormalities in patients,

we assessed tactile sensitivity in *TECPR2*-KI mice using von Frey filaments. At >P90, *TECPR2*-

KI mice exhibited a small increase in mechanical thresholds compared with WT controls (Fig. 1C),

while responses to pin prick and noxious heat were preserved (data not shown). Together, these

findings suggest subtle and age-dependent alterations in tactile sensitivity in *TECPR2*-KI mice,

though the magnitude and consistency of the effect warrant cautious interpretation.

Given the motor impairment present in patients with *TECPR2*-related disorder, we also analyzed

gait dynamics using a novel bottom-up imaging and data acquisition technology and analysis

platform (19). *TECPR2*-KI mice displayed progressive gait abnormalities, characterized by a

reduced hindlimb base width (increased intoeing), starting as early as P30 and worsening with age

(>P90) (Fig. 1D,E), supporting a relatively early behavioral onset of defects. Brainstem nuclei

implicated in tactile processing also contribute to sensorimotor integration of auditory inputs.

Therefore, we evaluated the acoustic startle reflex and observed that *TECPR2*-KI mice exhibited

a markedly reduced maximum startle amplitude at >P90 compared with WT littermates (Fig. 1F).

This represents a previously unrecognized phenotype in *TECPR2*-related disease and suggests an

additional brainstem-mediated sensory–motor impairment. Notably, grip strength and locomotion

did not differ between genotypes (Supplementary Fig. 1A,B), excluding generalized muscle

weakness as a cause of gait or startle abnormalities.

Collectively, these findings indicate that *TECPR2*-KI mice exhibit a subset of disease-relevant

phenotypes, most prominently brainstem acoustic startle disfunction, progressive gait

abnormalities, along with additional mechanosensory deficits.

Figure 1: *TECPR2*-KI mutation [NM_001081057.2; c.1319delC, p.Ser440Serfs19*] leads to
changes in behavior phenotype. **A,** Relative body weight of *TECPR2*-KI and WT littermate male
and female mice at P90 ($n = 20$-29 mice per group, data are presented as mean values ± s.e.m., *P*
values from unpaired t test). **B,** Grooming time of *TECPR2*-KI and WT littermate male and female
mice at >P90 ($n = 23$-41 mice per group; data are presented as mean values ± s.e.m., *P* values from
unpaired t test). **C,** Mechanical withdrawal thresholds of *TECPR2*-KI and WT littermate male and
female mice using an up–down von Frey method at >P90 ($n = 21$-33 mice per group; data are
presented as mean values ± s.e.m., *P* values from unpaired t test). **D,** Distance between hindpaws
used as a measure of intoeing in male and female *TECPR2*-KI and WT littermates over time
($n = 14$-62 mice per group, data are presented as mean values ± s.e.m., *P* values from two-way
ANOVA followed by Šídák's multiple comparisons test). **E,** Representative images of intoeing in
TECPR2-KI and WT littermate control. **F,** Maximum acoustic startle response in male and female
TECPR2-KI and WT littermates at P90 ($n = 16$-18 mice per group, data are presented as mean

values ± s.e.m., *P < 0.03, **P < 0.002, ***P < 0.001* from two-way ANOVA followed by Šídák's

multiple comparisons test).

TECPR2-KI mice develop progressive axonal pathology and defective autophagy in the

brainstem

Histopathological examination revealed that _TECPR2_-KI mice, but not WT littermates, developed

progressive axonal spheroid accumulation in the gracile and cuneate nuclei of the medulla

oblongata (Fig. 2A). These spheroids, representing focal axonal swellings, were also present in the

cervical–thoracic spinal cord (data not shown) but not in cortical layers, indicating selective

vulnerability of brainstem and spinal sensory tracts. Quantification demonstrated that spheroid

accumulation increased with age, first appearing at P60 (~100 spheroids per mm^2) and becoming

more pronounced at P90 (~300 spheroids per mm^2) (Fig. 2B). This progressive axonal pathology

closely mirrors previous reports in _Tecpr2_ knockout mice (6) and parallels the neurodegeneration

observed in patients. Ultrastructural analysis by transmission electron microscopy revealed

prominent structural abnormalities in _TECPR2_-KI mouse brainstem, showing atypical

autophagosome morphology and robust accumulations of single- and double-membrane-limited

autophagosome vacuoles with electron-dense content, consistent with impaired autophagosome

clearance (Supplementary Fig. 3A–D). Supporting this, RT–qPCR confirmed reduced endogenous

Tecpr2 expression in KI brainstem tissue (Fig. 2C). Immunostaining showed significant

accumulation of the autophagy substrate SQSTM1/p62 in mutant mice, while levels of

neurofilament heavy chain (NF-H), a biomarker of axonal integrity, were unchanged (Fig. 2D–F).

These findings indicate that loss of TECPR2 function causes selective autophagic defects in

brainstem nuclei, leading to progressive spheroid formation and axonal pathology.

Figure 2. *TECPR2*-KI mutation induces accumulation of spheroids and SQSMT1/p62 in the brainstem. A, Representative H&E staining of coronal brainstem sections (cuneate and gracile nuclei) of *TECPR2*-KI mice showing (indicated by arrows) progressive appearance of spheroids at P30, P60 and P90. Scale bars, 50 μm. **B**, Quantification of spheroids in brainstem of *TECPR2*-KI mice (*n* = 3-6 mice per group). **C**, RT–qPCR quantification of endogenous TECPR2 in brainstem of *TECPR2*-KI and WT littermates male and female mice (*n* = 3-7 mice per group, data are presented as mean values ± s.e.m., *P* values from two-way ANOVA followed by Šídák's multiple comparisons test). **D,** Representative SQSTM1/p62 and neurofilament heavy chain (NF-H) staining of brainstem of TECPR2-KI and WT littermates at P90, showing accumulation of SQSMT1/p62, general marker of autophagic flux, in TECPR2-KI mice. Scale bars, 100 μm. **E,** Quantification of SQSTM1/p62 and **F,** NF-H (*n* = 3-5 mice per group; data are presented as mean values ± s.e.m., *P* values from unpaired t test).

AAV9-mediated TECPR2 gene replacement is associated with improvements in select motor and sensory measures

To determine whether gene replacement could mitigate disease phenotypes, we delivered a single intracisternal injection of AAV9.pU1a-TECPR2_WT (VCAV-04485, 5.00E+12 GC/mL; 1uL) to *TECPR2*-KI mice and WT littermates at P1–3 (Fig. 3A). We chose the cisterna magna as the route of administration because of its proximity to the dorsal column nuclei, the earliest site of pathology, and its established efficacy in neonatal CNS gene therapy (14). The treatment was well tolerated in WT mice, supporting safety and long-term tolerability. In KI mice, AAV9-based gene placement prevented only a subset of the disorder manifestations. Although transient prevention of body weight loss was noted in males at P30 (Supplementary Fig. 2A), treated KI mice continued to exhibit reduced body weight at >P90 (Fig. 3B) and excessive grooming behavior (Fig. 3C). Improvements were observed across several functional measures. AAV9/TECPR2-treated KI mice showed normalized mechanical sensitivity at >P90 (Fig. 3D), improved hindlimb base width (paw intoeing) at P30 (Fig. 3E), and complete recovery of acoustic startle response amplitudes to WT levels (Fig. 3F). Together, these findings indicate that AAV9-mediated TECPR2 expression can partially ameliorate mechanosensory and brainstem-associated sensorimotor deficits, while growth and grooming abnormalities remain largely unaffected.

Figure 3. AAV9/TECPR2 mediated gene replacement in *TECPR2*-KI mice is associated with improvements in select motor and sensory measures. A, Illustration of experimental regimen. **B,** Relative body weight of TECPR2-KI, WT littermate, TECPR2-KI+AAV and WT littermate + AAV male and female mice at P90 (*n* = 17-29 mice per group, data are presented as mean values ± s.e.m., *P* values from ANOVA followed by Tukey's multiple comparisons test). **C,** Grooming time of TECPR2-KI, WT littermate, TECPR2-KI+AAV and WT littermate + AAV male and female mice at >P90 (*n* = 12-50 mice per group; data are presented as mean values ± s.e.m., *P*

values from ANOVA followed by Tukey's multiple comparisons test). **D**, Mechanical withdrawal thresholds of *TECPR2*-KI, WT littermate, *TECPR2*-KI+AAV and WT littermate + AAV male and female mice using an up–down von Frey method (n = 5-33 mice per group; data are presented as mean values ± s.e.m., *P* values from one-way ANOVA followed by Dunnett's T3 multiple comparisons test). **E,** Distance between hindpaws used as a measure of intoeing in male and female *TECPR2*-KI, WT littermate, *TECPR2*-KI+AAV and WT littermate + AAV over time (n = 5-62 mice per group, data are presented as mean values ± s.e.m., *P* values from two-way ANOVA followed by Šídák's multiple comparisons test). **F,** Maximum acoustic startle response in male and female *TECPR2*-KI, WT littermate, *TECPR2*-KI+AAV and WT littermate + AAV (n = 7-30 mice per group, data are presented as mean values ± s.e.m., *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$* from two-way ANOVA followed by Tukey's multiple comparisons test).

AAV9-mediated TECPR2 gene placement reduces spheroid pathology and improves autophagic flux

We next examined whether AAV9-TECPR2 treatment could mitigate axonal and autophagic defects in *TECPR2*-KI mice. Histological analysis revealed that intracisternal AAV9 delivery at P1–3 partially prevented spheroid accumulation in the gracile and cuneate nuclei, reducing their density by approximately 30% at P90 compared with untreated mutants (Fig. 4A,B). Robust viral-mediated expression of TECPR2 was confirmed in the brainstem of treated animals (Fig. 4C). Importantly, treated WT littermates displayed normal histoarchitecture and physiological TECPR2 expression, indicating that the therapy was safe and well tolerated. Immunohistochemical analyses further showed improvement of autophagy-related markers in AAV9-treated KI mice. Accumulation of the autophagy substrate SQSTM1/p62, prominent in untreated mutants, was similar to WT in the brainstem of treated animals (Fig. 4D,E), while levels of neurofilament heavy chain (NF-H) were comparable across groups (Fig. 4F).

Figure 4. AAV9/TECPR2 mediated gene replacement in *TECPR2*-KI mice reduces spheroid accumulation and ameliorates autophagic flux at P90. A, Representative H&E staining of coronal brainstem sections of *TECPR2*-KI, WT littermates and *TECPR2*-KI+AAV mice showing (indicated by arrows) spheroid accumulation at P90. Scale bars, 50 and 100 μm. **B,** Quantification of spheroids in brainstem of *TECPR2*-KI, WT littermate, *TECPR2*-KI+AAV and WT littermate + AAV male and female mice (*n* = 4-7 mice per group, data are presented as mean values ± s.e.m., *P* values from one-way ANOVA followed by Dunnett's multiple comparisons test). **C,** RT–qPCR quantification of endogenous and viral TECPR2 in brainstem of *TECPR2*-KI and WT littermates (endogenous), and *TECPR2*-KI+AAV and WT littermate + AAV (viral) male and female mice (*n* = 7-30 mice per group, data are presented as mean values ± s.e.m., *P* values from one-way ANOVA followed by Dunnett's multiple comparisons test). **D,** Representative SQSTM1/p62 staining of brainstem of *TECPR2*-KI, WT littermates and TECPR2-KI+AAV at P90. Scale bars, 100 μm. **E,** Quantification of SQSTM1/p62 and **F,** NF-H in brainstem of TECPR2-KI, WT littermates and

TECPR2-KI+AAV male and female mice (n = 3-7 mice per group, data are presented as mean

values ± s.e.m., P values from one-way ANOVA followed by Turkey's multiple comparison test).

DISCUSSION

TECPR2-related disorder is a rare neurodevelopmental and neurodegenerative disease for which

no disease-modifying therapies currently exist. In this study, we describe a knock-in mouse model

carrying the recurrent *TECPR2* c.1319delC founder mutation and characterize behavioral, sensory,

and neuropathological abnormalities associated with TECPR2 deficiency. Our data suggest an

association between impaired autophagic flux and selective involvement of brainstem sensory–

motor circuits and identify an acoustic startle deficit as a functional readout relevant to circuit

dysfunction. We further evaluate early postnatal AAV9-mediated *TECPR2* gene delivery and

observe partial improvements in autophagy-related markers, axonal pathology, and select

functional outcomes. Together, these results provide a framework for investigating disease

mechanisms and support further evaluation of gene replacement approaches in *TECPR2*-related

disorder, particularly to treat brain-stem related disease manifestations which account for early

mortality.

Neurodevelopmental and neurodegenerative disorders caused by impaired autophagy present a

major translational challenge (11,12) as they often combine early circuit dysfunction with

progressive neurodegeneration but lack validated disease models and actionable outcome

measures. *TECPR2*-related disorder exemplifies this gap: despite clear genetic causation and

severe clinical manifestations, therapeutic development has been limited by disease rarity and the

absence of *in vivo* systems that faithfully capture patient mutations and circuit-level pathology. In

this study, we address these limitations by generating a knock-in mouse model harboring the

recurrent *TECPR2* c.1319delC mutation and using it to define disease mechanisms, identify

translatable functional readouts, and evaluate early gene replacement therapy.

By introducing the patient-derived frameshift mutation into the endogenous mouse locus, we

generated a model that reproduces the predominant loss-of-function mechanism observed in

affected individuals. Unlike prior knockout or spontaneous mutation models, this knock-in

approach preserves the native genomic context while recapitulating the precise pathogenic allele

found in patients, increasing its relevance for both mechanistic and translational studies.

Longitudinal phenotyping revealed that homozygous *TECPR2*-KI mice develop reduced body
weight, increased grooming behavior, progressive gait abnormalities, and selective tactile sensory
impairment, some of which overlap with clinical features described in *TECPR2*-related disorder
(5,15). These phenotypes evolve over time and can be quantified reproducibly, establishing the
TECPR2-KI mouse as a robust platform for tracking disease progression and therapeutic response.
A key insight enabled by this model is the identification of selective brainstem circuit vulnerability
as a prominent feature of TECPR2 deficiency. Despite broad TECPR2 expression, neuropathology
in our model was largely restricted to sensory–motor pathways of the brainstem and spinal cord,
with progressive accumulation of axonal spheroids in the gracile and cuneate nuclei and relative
sparing of cortical neurons. These findings are consistent with recent work using a neuropathy-
associated *Tecpr2* knock-in mouse, which revealed endolysosomal loss-of-function phenotypes in
neurons and microglia, supporting a central role for disrupted vesicular and degradative pathways
in *TECPR2*-related disease (16). Together, these studies support a model in which TECPR2 loss
leads to selective vulnerability of long-range projection neurons and associated circuits, rather than
global neurodegeneration.
A key insight enabled by this model is the identification of selective brainstem circuit vulnerability
as a central feature of TECPR2 deficiency. Despite broad TECPR2 expression, neuropathology
was strikingly restricted to sensory–motor pathways of the brainstem and spinal cord, with
progressive accumulation of axonal spheroids in the gracile and cuneate nuclei and relative sparing
of cortical neurons. This pattern mirrors neuropathological findings in patients and other TECPR2-
deficient species and supports a circuit-selective disease mechanism rather than widespread
neurodegeneration. Long-range projection neurons in these pathways likely exhibit heightened
vulnerability due to their exceptional reliance on efficient autophagy to maintain axonal
homeostasis.
Consistent with this interpretation, ultrastructural and molecular analyses revealed marked defects
in autophagic processing within affected brainstem nuclei. Accumulation of abnormal autophagic
vacuoles and increased SQSTM1/p62 indicate impaired autophagosome maturation and clearance,
directly linking TECPR2 loss to defective autophagy and axonal degeneration. These findings
clarify how disruption of a core cellular homeostatic pathway translates into focal circuit
dysfunction and highlight autophagy as a mechanistically grounded therapeutic target.

Beyond previously recognized phenotypes, *TECPR2*-KI mice exhibited a pronounced reduction in
acoustic startle responses, a behavioral deficit not previously reported in *TECPR2*-related disorder.
The acoustic startle reflex depends on well-defined auditory–motor brainstem circuits, and its
impairment provides functional evidence of brainstem sensorimotor dysfunction. This finding is
particularly relevant given the prominence of brainstem-related clinical features in patients,
including impaired reflexes, dysphagia, and central apnea. Importantly, the acoustic startle
response represents a non-invasive, quantifiable, and cross-species functional readout, making it a
promising biomarker for disease progression and therapeutic efficacy in future clinical studies.
Using this knock-in mouse model, we next evaluated the effects of early gene replacement.
Neonatal intracisternal delivery of AAV9-*TECPR2* resulted in widespread transduction of
brainstem nuclei and was associated with improvements in autophagy-related markers, a partial
reduction in axonal spheroid burden, and improvements across multiple functional outcomes.
Mechanosensory thresholds, gait abnormalities, and acoustic startle responses similarly showed
improvement, indicating that early restoration of TECPR2 expression can ameliorate aspects of
brainstem-associated dysfunction. These results provide *in vivo* evidence that aspects of TECPR2-
related pathology can be modified by postnatal gene replacement. Not all phenotypes were fully
prevented. Reduced body weight and excessive grooming persisted in treated mice at later ages,
suggesting contributions from early developmental alterations, incomplete vector distribution, or
non–cell-autonomous effects involving peripheral or systemic tissues. These findings underscore
the importance of therapeutic timing and raise the possibility that even earlier intervention may be
required for full disease modification. They also suggest that combination strategies, such as gene
replacement paired with pharmacological modulation of lysosomal or autophagy-related
pathways, may further enhance therapeutic efficacy.
More broadly, this work informs understanding of selective vulnerability in neurodevelopmental
and neurodegenerative disorders linked to autophagy dysfunction. The relative sparing of cortical
neurons despite widespread TECPR2 expression highlights how circuit architecture and metabolic
demand shape disease expression. By linking impaired autophagic flux to degeneration of specific
brainstem sensory–motor pathways, our findings suggest convergent mechanisms that may extend
to other disorders characterized by axonal spheroids and brainstem involvement.
Few limitations should be considered. Our analyses focused primarily on the brainstem and spinal
cord, and future studies should evaluate peripheral nerves, forebrain circuits, and systemic

metabolic contributions. Only a single AAV dose and delivery route were tested, and comparative studies using alternative delivery strategies will be important for optimizing translational approaches. Finally, while the acoustic startle deficit emerges as a promising biomarker, its relevance in patients remains to be established through targeted clinical evaluation.

In summary, we developed a knock-in mouse model carrying the recurrent c.1319delC founder mutation that reproduces the underlying genetic alteration and exhibits circuit-selective pathology and functional impairments associated with TECPR2 deficiency. This model enabled the identification of brainstem sensory–motor circuits as a prominent site of vulnerability and revealed an acoustic startle phenotype that may serve as a translatable functional readout. In addition, early AAV9-mediated TECPR2 delivery was associated with improvement of autophagy-related markers and partial amelioration of select behavioral outcomes. By combining a genetically defined mouse model with quantifiable functional endpoints and a clinically established gene delivery strategy, this study provides a preclinical framework to evaluate TECPR2 gene replacement and to inform future investigations of therapeutic timing, target engagement, and outcome measures.

MATERIALS AND METHODS

Generation of the TECPR2 Knock-In Mouse Model

TECPR2 knock-in (KI) mice were generated at the Harvard Genome Modification Facility. CRISPR/Cas9/guide RNA and donor oligo were injected into fertilized C57BL/6 eggs to generate the equivalent HSP50 exon 8 mutation in the mice. This mutation is a deletion of a single C nucleotide in mouse exon 8: NM_001081057.2; c.1319delC, p.Ser440Serfs19*. Male F1 heterozygous mice were transferred to Charles River Laboratories, where they used these mice for IVF of oocytes from female C57BL/6 to generate heterozygous breeders and used to set up a breeding colony. Mice were genotyped using PCR and sequencing by the Molecular Genetics Core Facility at Boston Children's Hospital to confirm the presence of the mutation. TECPR2 KI mice were housed in standard clear plastic cages with no more than 5 animals per cage under controlled conditions (lights on 07:00-19:00; humidity 30%-50%; temperature 22-23°C) with ad libitum access to food and water. All experiments were performed between 9:00 and 17:00 in a room

maintained at a temperature of $21 \pm 1°C$. All experimental protocols comply with relevant ethical

regulations and were approved by Boston Children's Hospital Institutional Animal Care and Use

Committee (protocol 00002125). Animals were randomized to treatment groups.

AAV production

Through the Luke Heller TECPR2 Foundation, we have obtained WT human TECPR2and GFP-

AAV9 constructs, generated at the Horae Gene Therapy Center, UMass Chan Medical School

(Worcester, MA, USA) under the direction of Dr. Guangping Gao. The AAV genome is composed

of U1a promoter, TECPR2 CDS, a synthetic short polyA and AAV2 ITRs at both ends. The genome

was packaged into the AAV9 capsid by triple-transfection in HEK293 cells.

In vivo injection of viral vectors

For AAV9 delivery into the CSF via cisterna magna, mice at postnatal day 1–3 (P1–P3) were

anesthetized by rapid induction of hypothermia for 2–4 min on ice water until loss of

consciousness. The injection site was disinfected with 70% ethanol. A 33-gauge needle attached

to a Hamilton syringe was lowered approximately 2 mm into the cisterna magna at a 45° angle,

and 1 µL of viral solution (VCAV-04485, 5.00E+12 GC/mL) was slowly injected. Body

temperature was gradually restored and maintained on a 37 °C warming pad for 10 min after

injection before returning pups to the parental cage. Injections were performed prior to genotyping

and blinded to genotype; all pups within each litter received the same viral vector.

Body Weight and Growth Curve Analysis

Body weights of the TECPR2 KI mice and their WT littermates were recorded with an electronic

scale at postnatal day 30, 60 and 90 (P30, P60 and P90, respectively). Growth curves were plotted

to compare the development of mutant and WT mice.

Mechanical Sensitivity Testing

Mechanical sensitivity was assessed using the von Frey filament test. Mice were placed in

individual chambers on a mesh platform, and calibrated filaments were applied to the plantar

surface of the left hind paw, with a positive response consisting of a paw lifting or flinching

response to the fiber. The response patterns were collected and converted into corresponding 50%

withdrawal thresholds using the Up–Down Reader software and associated protocol (17). All tests
were performed by investigators blinded to the genotype and treatment.

Acoustic startle reflexes.

Mice were tested for startle reflexes in response to broadband auditory stimulation at varying
intensities as previously described (18). The animals were tested in a sound shielded startle booth
(Kinder Scientific) and the force generated by foot movement was sensed by piezoelectric motion
sensor fixed beneath an elevated platform. Mice were placed in a smaller sub chamber anchored
to the topside of the platform, which restricted them from rearing on their limbs but freely
permitted penetration of the sound stimulus. Sound stimuli were calibrated with the door closed
using a sound pressure level meter (Allied Electronics) with the microphone mounted in the
position normally occupied by the animal holder. The generation of sound stimuli and recording
of force amplitude signals was performed by startle monitor software (Kinder Scientific).
Broadband white noise was presented at a background level of 60 dB throughout the experiment
and auditory test stimuli consisted of 50-ms broadband white noise pulses in 10-dB steps from 60
to 120 dB. Different intensities of the test stimulus were presented in a pseudo-random order at
randomized inter trial intervals (ITI) that varied between 8 and 22 s, with no ITI repeated more
than three times. Five repetitions were averaged for each of the intensities for one test subject and
responses were normalized to the weight of the mouse tested. Startle response measurements were
performed by investigators blinded to the genotype.

Intoeing and grooming behaviors

Intoeing was quantified using a novel bottom-up imaging and data acquisition technology and
analysis platform that provides automated, quantitative, and objective measures of naturalistic
rodent behavior in an observer-independent and unbiased fashion as previously described (19).
The location of the paws was extracted from body frames using the DeepLabCut and the distance
between hindpaws (measure of intoeing) in the unit of pixel distance was calculated. Grooming
behaviors were calculated by the time the animal spent grooming the face and body in seconds for
10 minutes recording using the ARBEL algorithm, a machine learning tool with light-based image
analysis for automatic classification of 3D behaviors (20).

Tissue processing and histopathological analysis

Animals were anesthetized with pentobarbital and perfused with a 4% paraformaldehyde (PFA) solution in PBS. The brain was dissected and fixed in 4% PFA solution overnight. Samples were then transferred to a 30% sucrose solution until the tissue sank. Samples were immediately frozen in OCT embedding medium and kept in -80oC until further processing. Coronal cryosections (50 µm) were collected through the brainstem and spinal cord for immunostaining and histopathological analysis. Slices were then stained with hematoxylin and eosin (H&E) to identify spheroids or probed with antibodies for neurofilament and autophagic flux. Quantification of spheroids was done measuring the density of spheroids per mm2 in the cuneate and gracile nuclei areas bilaterally in 10-12 sections per animal, then an average density was calculated using QuPath software (21).

Immunofluorescence staining

Sections were washed in PBS, permeabilized with PBS + 0.3% Triton X-100 (PBST-0.3) for 15 min and blocked in PBST-0.3 containing 5% normal goat serum (NGS) for 1 h at room temperature. Samples were then incubated overnight at 4 °C with primary antibodies in blocking buffer (Mouse anti-SQSTM1/p62; 1:500; Cell Signaling Technologies #88588; rabbit anti-neurofilament heavy chain; 1:1,000; Sigma Aldrich #N4142). After washes in PBST-0.3, sections were incubated with secondary antibodies for 2 h at room temperature (1:500; Alexa Fluor–conjugated goat anti-rabbit or goat anti-mouse, Invitrogen). Nuclei were counterstained with DAPI (0.5 µg/mL). Sections were mounted with ProLong Glass antifade reagent and imaged on a Zeiss LSM880 confocal microscope. Neurofilament (NF-H) intensity and SQSTM1/p62+ puncta were measured from epifluorescent images. NF-H intensity was measured using sections from WT tissues to set a threshold and the mean pixel intensity from WT and TECPR2 KI tissues was measured using Zeiss Zen 3.8 Software. The quantification of SQSTM1/p62 puncta was done by measuring the area of the cuneate and gracile from epifluorescent images and manually detecting SQSTM1/p62+ puncta in those regions using Zeiss Zen 3.8 Software. Quantitative analysis was performed by an investigator who was blind to the experimental conditions.

Electron microscopy

Animals were anesthetized with pentobarbital and perfused with 2.5% PFA, 2.5% glutaraldehyde, in 0.1 M cacodylate buffer pH 7.4. Brain samples were fixed with 2.5% PFA, 2.5% glutaraldehyde, in 0.1 M cacodylate buffer containing 5 mM CaCl2 (pH 7.4), post-fixed in 1% osmium tetroxide supplemented with 0.5% potassium hexacyanoferrate trihydrate and potassium dichromate in 0.1 M cacodylate for 1 h, stained with 2% uranyl acetate in double-distilled water for 1 h, dehydrated in graded ethanol solutions, and embedded in epoxy resin (Agar Scientific, AGR1030). Ultrathin sections (70–90 nm) of brainstem, obtained with a LeicaUltracut UCT microtome, were stained with lead citrate and then examined using either a Phillips CM-12 transmission electron microscope (TEM) equipped with a Gatan One View camera or an FEI Tecnai SPIRIT TEM equipped with a bottom-mounted 2 k × 2 k FEI Eagle CCD camera.

Statistical Analysis

Data are expressed as the mean ± SEM. Significant differences between groups were assessed with one or two-way ANOVA; once the significance of the group differences ($p \leq 0.05$) was established, appropriate post-hoc comparisons between pairs of groups were used. Statistical analyses were done using GraphPad Prism Software.

List of Supplementary Materials

Materials and Methods

Fig. S1 to S3.

References

1.	D. Oz-Levi, B. Ben-Zeev, E. K. Ruzzo, Y. Hitomi, A. Gelman, K. Pelak, Y. Anikster, H. Reznik-Wolf, I. Bar-Joseph, T. Olender, A. Alkelai, M. Weiss, E. Ben-Asher, D. Ge, K. V. Shianna, Z. Elazar, D. B. Goldstein, E. Pras, D. Lancet, Mutation in TECPR2 reveals a role for autophagy in hereditary spastic paraparesis. Am. J. Hum. Genet. 91, 1065–1072 (2012).

2. P. P. Zhu, K. R. Denton, T. M. Pierson, X. J. Li, C. Blackstone, Mutations in TECPR2,

encoding a positive regulator of autophagy, cause a new syndrome of childhood-onset spasticity

and axonal neuropathy. Am. J. Hum. Genet. 97, 671–682 (2015).

3. G. Heimer, D. Oz-Levi, E. Eyal, S. Edvardson, A. Nissenkorn, E. K. Ruzzo, A.

Szeinberg, C. Maayan, M. Mai-Zahav, O. Efrati, E. Pras, H. Reznik-Wolf, D. Lancet, D. B.

Goldstein, Y. Anikster, S. A. Shalev, O. Elpeleg, B. Ben Zeev, TECPR2 mutations cause a new

syndrome of progressive neurodegeneration with developmental delay. Brain 139, 4034–4044

(2016).

4. S. Anazi, S. Maddirevula, V. Salpietro, Y. T. Asi, S. Alsahli, A. Alhashem, H. E.

Shamseldin, F. AlZahrani, N. Patel, N. Ibrahim, F. M. Abdulwahab, M. Hashem, N. Alhashmi,

F. Al Murshedi, A. Al Kindy, A. Alshaer, A. Rumayyan, S. Al Tala, W. Kurdi, A. Alsaman, A.

Alasmari, S. Banu, T. Sultan, M. M. Saleh, H. Alkuraya, M. A. Salih, H. Aldhalaan, T. Ben-

Omran, F. Al Musafri, R. Ali, J. Suleiman, B. Tabarki, A. W. El-Hattab, C. Bupp, M. Alfadhel,

N. Al Tassan, D. Monies, S. T. Arold, M. Abouelhoda, T. Lashley, H. Houlden, E. Faqeih, F. S.

Alkuraya, Expanding the genetic heterogeneity of intellectual disability. Hum. Genet. 136, 1419–

1429 (2017).

5. S. Neuser, B. Brechmann, G. Heimer, I. Brösse, S. Schubert, L. O'Grady, M. Zech, S.

Srivastava, D. A. Sweetser, Y. Dincer, V. Mall, J. Winkelmann, C. Behrends, B. T. Darras, R. J.

Graham, P. Jayakar, B. Byrne, B. E. Bar-Aluma, Y. Haberman, A. Szeinberg, H. M. Aldhalaan,

M. Hashem, A. Al Tenaiji, O. Ismayl, A. E. Al Nuaimi, K. Maher, S. Ibrahim, F. Khan, H.

Houlden, V. S. Ramakumaran, A. T. Pagnamenta, J. E. Posey, J. R. Lupski, W. H. Tan, G.

ElGhazali, I. Herman, T. Muñoz, G. M. Repetto, A. Seitz, M. Krumbiegel, M. Poli, U. Kini, S.

Efthymiou, J. Meiler, R. Maroofian, F. S. Alkuraya, R. Abou Jamra, B. Popp, B. Ben-Zeev, D.

Ebrahimi-Fakhari, Clinical, neuroimaging, and molecular spectrum of TECPR2-associated

hereditary sensory and autonomic neuropathy with intellectual disability. Hum. Mutat.42, 762–

776 (2021).

6. M. Fraiberg, B. C. Tamim-Yecheskel, K. Kokabi, N. Subic, G. Heimer, F. Eck, K.

Nalbach, C. Behrends, B. Ben-Zeev, O. Shatz, Z. Elazar, Lysosomal targeting of

autophagosomes by the TECPR domain of TECPR2. Autophagy 17, 3096–3108 (2021).

7.   S. Maday, E. L. J. Holzbaur, Autophagosome biogenesis in primary neurons follows an
ordered and spatially regulated pathway. Dev. Cell 30, 71–85 (2014).

8.   J. H. Hurley, L. N. Young, Mechanisms of autophagy initiation. Annu. Rev. Biochem.
86, 225–244 (2017).

9.   M. Komatsu, S. Waguri, T. Chiba, S. Murata, J. I. Iwata, I. Tanida, T. Ueno, M. Koike,
Y. Uchiyama, E. Kominami, K. Tanaka, Loss of autophagy in the central nervous system causes
neurodegeneration in mice. Nature 441, 880–884 (2006).

10.   D. Stadel, V. Millarte, K. D. Tillmann, J. Huber, B. C. Tamim-Yecheskel, M. Akutsu, A.
Demishtein, B. Ben-Zeev, Y. Anikster, F. Perez, V. Dötsch, Z. Elazar, V. Rogov, H. Farhan, C.
Behrends, TECPR2 cooperates with LC3C to regulate COPII-dependent ER export. Mol. Cell
60, 89–104 (2015).

11.   D. Ebrahimi-Fakhari, A. Saffari, L. Wahlster, J. Lu, S. Byrne, G. F. Hoffmann, H.
Jungbluth, M. Sahin, Congenital disorders of autophagy: An emerging novel class of inborn
errors of neuro-metabolism. Brain 139, 317–337 (2016).

12.   H. S. Dafsari, D. Martinelli, A. Saffari, D. Ebrahimi-Fakhari, M. Fanto, C. Dionisi-Vici,
H. Jungbluth, An update on autophagy disorders. J. Inherit. Metab. Dis. 48, e12798 (2025).

13.   L. A. Williams, D. J. Gerber, A. Elder, W. C. Tseng, V. Baru, N. Delaney-Busch, C.
Ambrosi, G. Mahimkar, V. Joshi, H. Shah, K. Harikrishnan, H. Upadhyay, S. H. Rajendran, A.
Dhandapani, J. Meier, S. J. Ryan, C. Lewarch, L. Black, J. Douville, S. Cinquino, H. Legakis, K.
Nalbach, C. Behrends, A. Sato, L. Galluzzi, T. W. Yu, D. Brown, S. Agrawal, D. Margulies, A.
Kopin, G. Dempsey, Developing antisense oligonucleotides for a TECPR2 mutation-induced,
ultra-rare neurological disorder using patient-derived cellular models. Mol. Ther. Nucleic Acids
29, 189–203 (2022).

14.   R. M. Bailey, D. Armao, S. Nagabhushan Kalburgi, S. J. Gray, Development of
intrathecal AAV9 gene therapy for giant axonal neuropathy. Mol. Ther. Methods Clin. Dev. 9,
160–171 (2018).

15. G. Heimer, S. Neuser, B. Ben-Zeev, D. Ebrahimi-Fakhari, TECPR2-related hereditary
sensory and autonomic neuropathy with intellectual disability. In: Adam MP, Bick S, Mirzaa
GM, Pagon RA, Wallace SE, Amemiya A, editors. GeneReviews® [Internet]. Seattle (WA):
University of Washington; 1993–2026 (updated 2022).

16. D. Bhattacharya, P. da Silva-Buttkus, K. Nalbach, L. Cheng, L. Garrett, M. Irmler, G.
Kislinger, G. Werner, R. Rodde, C. Lengger, J. Beckers, A. Zimprich, S. M. Hölter, V. Gailus-
Durner, H. Fuchs, M. Hrabe de Angelis, B. Wefers, W. Wurst, M. S. Brill, M. Schifferer, S. F.
Lichtenthaler, C. Behrends, Neuropathy-associated Tecpr2 mutation knock-in mice reveal
endolysosomal loss of function phenotypes in neurons and microglia. Cell Death Dis. 16, 775
(2025).

17. R. Gonzalez-Cano, B. Boivin, D. Bullock, L. Cornelissen, N. Andrews, M. Costigan, Up-
Down Reader: An Open Source Program for Efficiently Processing 50% von Frey Thresholds.
Front. Pharmacol. 9, 433 (2018).

18. C. A. Nist-Lund, B. Pan, A. Patterson, Y. Asai, T. Chen, W. Zhou, H. Zhu, S. Romero, J.
Resnik, D. B. Polley, G. S. Géléoc, J. R. Holt, Improved TMC1 gene therapy restores hearing
and balance in mice with genetic inner ear disorders. Nat. Commun. 10, 236 (2019).

19. Z. Zhang, D. P. Roberson, M. Kotoda, B. Boivin, J. P. Bohnslav, R. González-Cano, D.
A. Yarmolinsky, B. L. Turnes, N. K. Wimalasena, S. Q. Neufeld, L. B. Barrett, N. L. M.
Quintão, V. Fattori, D. G. Taub, A. B. Wiltschko, N. A. Andrews, C. D. Harvey, S. R. Datta, C.
J. Woolf, Automated preclinical detection of mechanical pain hypersensitivity and analgesia.
Pain 163, 2326–2336 (2022).

20. O. Barkai, B. Zhang, B. L. Turnes, M. Arab, D. A. Yarmolinsky, Z. Zhang, L. B. Barrett,
C. J. Woolf, A machine learning tool with light-based image analysis for automatic classification
of 3D pain behaviors. Cell Rep. Methods 5, 101145 (2025).

21. P. Bankhead, M. B. Loughrey, J. A. Fernández, Y. Dombrowski, D. G. McArt, P. D.
Dunne, S. McQuaid, R. T. Gray, L. J. Murray, H. G. Coleman, J. A. James, M. Salto-Tellez, P.
W. Hamilton, QuPath: Open source software for digital pathology image analysis. Sci. Rep. 7,
16878 (2017).

Acknowledgments

Luke Heller TECPR2 Foundation

Jordan Avi Ogman Foundation

Quiver Bioscience [formerly Q-State Biosciences]

Animal Behavior and Physiology Core at Boston Children's Hospital

Molecular Genetics Core Facility at Boston Children's Hospital

Cellular Imaging Core at Boston Children's Hospital

Neurobiology Imaging Facility at Harvard Medical School

Electron Microscopy Core at Harvard Medical School

Funding

National Institute of Health grant P50HD105351 (MF)

United States - Israel Binational Science Foundation (MF)

Author contributions

Conceptualization: BLT, NH, NA, MF

Methodology: BLT, SL, ASK, NH, NA, MF

Investigation: BLT, PCC, LM, TB, JZ, FV, EC, MA, BZ, SL, NH, NA

Funding acquisition: NH, NA, MF

Project administration: BLT, NH, NA, MF

Supervision: BLT, NA, NH, MF

Writing – original draft: BLT

Writing – review & editing: BLT, NH, ASK, NA, DEF, MF

 Supplementary

Supplementary Fig. 1: TECPR2-KI mutation [NM_001081057.2; c.1319delC, p.Ser440Serfs19*] does not compromise locomotion and grip strength. A, Locomoting time of TECPR2-KI and WT littermate male and female mice at P30, P60 and P90 (n = 5-59 mice per group, data are presented as mean values ± s.e.m., P values from two-way ANOVA followed by Šídák's multiple comparisons test). **B**, Grip strength of TECPR2-KI and WT littermate male and female mice at >P90 (n = 23-41 mice per group; data are presented as mean values ± s.e.m., P values from unpaired t test). C, Mechanical withdrawal thresholds of TECPR2-KI and WT littermate male and female mice using an up–down von Frey method at >P90 (n = 9-25 mice per group; data are presented as mean values ± s.e.m., P values from unpaired t test).

Supplementary Fig. 2: TECPR2-KI mutation [NM_001081057.2; c.1319delC, p.Ser440Serfs19*] induces changes in body weight in male mice. A, Body weight of TECPR2-

KI and WT littermate male and **B**, female mice at P30, P60 and P90 (n = 4-33 mice per group, data

are presented as mean values ± s.e.m., P values from two-way ANOVA followed by Šídák's

multiple comparisons test).

Supplementary Fig. 3: TECPR2-KI mutation induces structural abnormalities in TECPR2-

KI mouse brainstem. A-D, Ultrastructural analysis by transmission electron microscopy revealed

prominent structural abnormalities in TECPR2-KI mouse brainstem.