1 **Preclinical validation of AAV9-TECPR2 gene therapy in a novel knock-in model of**
2 **TECPR2-related disorder**
3

4 Bruna Lenfers Turnes[1], Peter Casey-Caplan[1], Leo Mejia[1], Tiffany Berry[1], Julie Zhao[1],
5 Francesco Villa[1], Emma Cropper[1], Maryam Arab[2], Biyao Zhang[1], Darred Surin[3], Darius
6 Ebrahimi-Fakhari[1,2], Silmara de Lima[4], Alan S. Kopin[6], Nick Andrews[5], Nathaniel
7 Hodgson[1], Michela Fagiolini[1]
8

Affiliations:
[1] F.M. Kirby Neurobiology Center & Department of Neurology, Boston Children's
Hospital, Harvard Medical School, Boston, MA, USA
[2]Harvard Medical School, Boston, MA, USA
[3]Department of Biology, Boston University, Boston, MA, USA
[4]Department of Ophthalmology, University of Pittsburg, Pittsburgh, PA, USA
[5]In Vivo Scientific Services, Salk Institute for Biological Studies, La Jolla, CA, USA.
[6]Tufts University School of Medicine, Boston, MA, USA
17

18 Corresponding author: Michela Fagiolini, Michela.Fagiolini@childrens.harvard.edu
19

20 **One-Sentence Key Message**
21 TECPR2 deficiency disrupts brainstem sensory–motor circuits, impairing autophagy and
22 tactile, gait, and startle function and is prevented by neonatal AAV9.
23

Abstract

TECPR2-related disorder is a rare, autosomal recessive neurodevelopmental and

neurodegenerative disease characterized by early-onset motor dysfunction, sensory-

and autonomic neuropathy, and progressive neurological decline with early mortality.

Currently, there are no effective treatments for individuals affected by this debilitating

condition. To advance our understanding of disease mechanisms and explore

therapeutic strategies, we developed and then characterized a knock-in (KI) mouse

model carrying the human *TECPR2* c.1319delC frameshift mutation.

TECPR2-KI mice exhibit a subset of disease-relevant phenotypes, most prominently

abnormal gait, along with reduced body weight and altered tactile sensitivity. We

additionally observe a reduction in acoustic startle responses, consistent with

dysfunction of brainstem-associated sensorimotor pathways. Histopathological analyses

reveal progressive accumulation of axonal spheroids in the dorsal column nuclei,

together with abnormalities in autophagy-related markers, features previously reported

in individuals with *TECPR2*-related disorder.

To assess the therapeutic potential of gene replacement, we delivered *TECPR2* via

intracisternal infusion of AAV9/*TECPR2* in neonatal KI mice. Gene therapy restored

mechanosensory function, normalized gait and startle responses, maintain autophagic

homeostasis, and partially reduced axonal pathology. These findings demonstrate that

TECPR2-associated deficits are not only replicable in this new mouse model but are

also amenable to postnatal intervention.

Our study introduces a genetically accurate murine model of TECPR2 deficiency,

identifies brainstem-associated phenotypes, and provides preliminary evidence

supporting the feasibility of AAV9-mediated TECPR2 gene delivery, establishing a

foundation for future translational research in a currently untreatable disease.

Introduction

TECPR2-related disorder is a devastating, ultra-rare neurodevelopmental and

neurodegenerative condition for which no disease-modifying therapies currently exist. It

is classified along the spectrum of the hereditary sensory and autonomic neuropathies

(specifically HSAN9, OMIM #615031) and the complex hereditary spastic paraplegias

(HSPs). Since its first identification in 2012, fewer than 40 individuals have been

reported worldwide, underscoring both the rarity and the limited clinical awareness of

the disorder (1-5).

TECPR2-related disorder is caused by biallelic loss-of-function mutations in the

TECPR2 gene, which encodes the tectonin beta-propeller repeat-containing protein 2, a

multifunctional scaffold protein with broad expression across the central nervous system

(CNS). TECPR2 has been implicated as a key regulator of the autophagy–lysosome

pathway, particularly in mediating the targeting of autophagosomes to lysosomes and

promoting autophagic flux (6). Efficient autophagic clearance is essential for maintaining

neuronal health, especially in long-projecting neurons that depend on the constant

turnover of organelles and proteins to preserve axonal integrity (7,8). Accordingly,

dysfunction of autophagy-related genes has been increasingly associated with a wide

range of neurodevelopmental and neurodegenerative disorders (9-12). Loss of TECPR2

function results in impaired autolysosomal fusion, accumulation of protein aggregates,

and formation of neuroaxonal spheroids, ultimately leading to axonal dystrophy and

neuronal loss (6).

Pathological spheroids have been documented not only in affected human patients but

also in other species, highlighting the conserved nature of TECPR2-deficiency related

axonopathy. For instance, Spanish Water Dogs with a spontaneous *TECPR2* loss-of-

function mutations display progressive axonal degeneration, and a CRISPR-Cas9-

generated *Tecpr2* knockout mouse model exhibits comparable neuropathological

features (11,5). These findings reinforce the importance of developing robust and

genetically faithful animal models to study disease mechanisms and evaluate potential

interventions.

Clinically, *TECPR2*-related disorder manifests with global developmental delay and later

intellectual disability, early-onset hypotonia, sensory and autonomic neuropathy,

progressive spasticity, and progressive bulbar dysfunction with dysphagia and central

apneas. Dysfunction of brainstem-mediated circuits, contributing to dysphagia, impaired
reflexes, and central apneas, accounts for significant morbidity and early mortality. Most
affected individuals succumb to disease in the first or second decade of life, and
survivors experience profound, lifelong disability (1,3,5). Despite the recognition of
these hallmark features, the precise temporal progression of neuropathological changes
and their mechanistic underpinnings remain poorly defined, representing a major barrier
to therapeutic development.

Efforts to identify therapeutic strategies for *TECPR2*-related disorders have been
hindered by the rarity of the condition and the lack of suitable *in vivo* models that
capture disease-relevant mutations and phenotypes. Although a recent study
demonstrated that antisense oligonucleotides (ASOs) targeting exon 8 of TECPR2
produced promising effects *in vitro* (13), the development of candidate therapeutics for
TECPR2-related disorders, genetic approaches (e.g. ASO, gene therapy) or otherwise,
will require *in vivo* models to test the efficacy of the approach.

Given these limitations, gene replacement therapy using adeno-associated virus
serotype 9 (AAV9) has emerged as a potential alternative. AAV9 vectors are well
characterized for their ability to transduce both neurons and glia across the CNS
following intracisternal, intrathecal or intravenous delivery, and they have been
successfully used in the treatment of other pediatric neurological disorders, including
spinal muscular atrophy (14). Importantly, this approach allows for full-length *TECPR2*
gene delivery irrespective of the specific mutation, offering a mutation-agnostic strategy
for therapeutic intervention.

To address the critical gap in preclinical models, we generated a knock-in (KI) mouse
line carrying the human *TECPR2* c.1319delC frameshift mutation. This model enables
the assessment of behavioral, sensory, and neuropathological alterations associated
with TECPR2 deficiency and was used to evaluate intracisternal AAV9-mediated
TECPR2 gene delivery at neonatal stages.

Together, this work provides (i) a genetically accurate *in vivo* model for investigating
TECPR2-related pathophysiology, (ii) insight into previously unrecognized sensorimotor
deficits involving brainstem circuits, and (iii) initial preclinical evidence supporting the
feasibility of *TECPR2* gene replacement.

Results

Behavioral alterations in *TECPR2*-KI mice recapitulate aspects of the clinical phenotype observed in *TECPR2*-related disorder

To model human *TECPR2*-associated disease, we generated a homozygous *Tecpr2* c.1319delC knock-in (KI) mouse line. This mutation introduces a premature stop codon at the same position as the pathogenic c.1319delT founder variant identified in a large cohort of patients (5) thereby recapitulating the genetic basis of disease. We performed longitudinal behavioral phenotyping at postnatal days 30, 60, and 90 to assess developmental and progressive manifestations.

At P90, *TECPR2*-KI mice exhibited a small significant reduction in body weight compared with WT littermates (Fig. 1A), consistent with reports of growth impairment in some individuals with *TECPR2*-related disorder (5). We examined their self-grooming behavior over a 10-minute video recording with automatic classification of behaviors and observed that TECPR2-KI mice show increased grooming behavior at P90 (Fig. 1B). Given reports of sensory abnormalities in patients, we assessed tactile sensitivity in *TECPR2*-KI mice using von Frey filaments. At >P90, *TECPR2*-KI mice exhibited a small increase in mechanical thresholds compared with WT controls (Fig. 1C), while responses to pin prick and noxious heat were preserved (data not shown). Together, these findings suggest subtle and age-dependent alterations in tactile sensitivity in *TECPR2*-KI mice, though the magnitude and consistency of the effect warrant cautious interpretation.

Given the motor impairment present in patients with *TECPR2*-related disorder, we also analyzed gait dynamics using a novel bottom-up imaging and data acquisition technology and analysis platform (19). *TECPR2*-KI mice displayed progressive gait abnormalities, characterized by a reduced hindlimb base width (increased intoeing), starting as early as P30 and worsening with age (>P90) (Fig. 1D,E), supporting a relatively early behavioral onset of defects. Brainstem nuclei implicated in tactile processing also contribute to sensorimotor integration of auditory inputs. Therefore, we

evaluated the acoustic startle reflex and observed that *TECPR2*-KI mice exhibited a
markedly reduced maximum startle amplitude at >P90 compared with WT littermates
(Fig. 1F). This represents a previously unrecognized phenotype in *TECPR2*-related
disease and suggests an additional brainstem-mediated sensory–motor impairment.
Notably, grip strength and locomotion did not differ between genotypes (Supplementary
Fig. 1A,B), excluding generalized muscle weakness as a cause of gait or startle
abnormalities.
Collectively, these findings indicate that *TECPR2*-KI mice exhibit a subset of disease-
relevant phenotypes, most prominently brainstem acoustic startle disfunction,
progressive gait abnormalities, along with additional mechanosensory deficits.

Figure 1: ***TECPR2*-KI mutation [NM_001081057.2; c.1319delC, p.Ser440Serfs19*]**
leads to changes in behavior phenotype. A, Relative body weight of *TECPR2*-KI and
WT littermate male and female mice at P90 ($n = 20$-29 mice per group, data are

presented as mean values $\pm$ s.e.m., *P* values from unpaired t test). **B,** Grooming time
of *TECPR2*-KI and WT littermate male and female mice at >P90 ($n = $ 23-41 mice per
group; data are presented as mean values $\pm$ s.e.m., *P* values from unpaired t test). **C,**
Mechanical withdrawal thresholds of *TECPR2*-KI and WT littermate male and female
mice using an up–down von Frey method at >P90 ($n = $ 21-33 mice per group; data
are presented as mean values $\pm$ s.e.m., *P* values from unpaired t test). **D,** Distance
between hindpaws used as a measure of intoeing in male and female *TECPR2*-KI and
WT littermates over time ($n = $ 14-62 mice per group, data are presented as mean
values $\pm$ s.e.m., *P* values from two-way ANOVA followed by Šídák's multiple
comparisons test). **E,** Representative images of intoeing in *TECPR2*-KI and WT
littermate control. **F,** Maximum acoustic startle response in male and female *TECPR2*-KI
and WT littermates at P90 ($n = $ 16-18 mice per group, data are presented as mean
values $\pm$ s.e.m., **P < 0.03, **P < 0.002, ***P < 0.001* from two-way ANOVA followed
by Šídák's multiple comparisons test).

TECPR2-KI mice develop progressive axonal pathology and defective autophagy in the brainstem

Histopathological examination revealed that *TECPR2*-KI mice, but not WT littermates,
developed progressive axonal spheroid accumulation in the gracile and cuneate nuclei
of the medulla oblongata (Fig. 2A). These spheroids, representing focal axonal
swellings, were also present in the cervical–thoracic spinal cord (data not shown) but
not in cortical layers, indicating selective vulnerability of brainstem and spinal sensory
tracts. Quantification demonstrated that spheroid accumulation increased with age, first
appearing at P60 (~100 spheroids per mm^2) and becoming more pronounced at P90
(~300 spheroids per mm^2) (Fig. 2B). This progressive axonal pathology closely mirrors
previous reports in *Tecpr2* knockout mice (6) and parallels the neurodegeneration
observed in patients. Ultrastructural analysis by transmission electron microscopy
revealed prominent structural abnormalities in *TECPR2*-KI mouse brainstem, showing
atypical autophagosome morphology and robust accumulations of single- and double-
membrane-limited autophagosome vacuoles with electron-dense content, consistent
with impaired autophagosome clearance (Supplementary Fig. 3A–D). Supporting this,

RT–qPCR confirmed reduced endogenous Tecpr2 expression in KI brainstem tissue (Fig. 2C). Immunostaining showed significant accumulation of the autophagy substrate SQSTM1/p62 in mutant mice, while levels of neurofilament heavy chain (NF-H), a biomarker of axonal integrity, were unchanged (Fig. 2D–F). These findings indicate that loss of TECPR2 function causes selective autophagic defects in brainstem nuclei, leading to progressive spheroid formation and axonal pathology.

Figure 2. *TECPR2*-**KI mutation induces accumulation of spheroids and SQSMT1/p62 in the brainstem. A**, Representative H&E staining of coronal brainstem sections (cuneate and gracile nuclei) of *TECPR2*-KI mice showing (indicated by arrows) progressive appearance of spheroids at P30, P60 and P90. Scale bars, 50 µm. **B**, Quantification of spheroids in brainstem of *TECPR2*-KI mice (*n* = 3-6 mice per group). **C**, RT–qPCR quantification of endogenous TECPR2 in brainstem of *TECPR2*-KI and WT littermates male and female mice (*n* = 3-7 mice per group, data are presented as mean values ± s.e.m., *P* values from two-way ANOVA followed by Šídák's multiple comparisons test). **D,** Representative SQSTM1/p62 and neurofilament heavy chain (NF-H) staining of brainstem of TECPR2-KI and WT littermates at P90, showing

accumulation of SQSMT1/p62, general marker of autophagic flux, in TECPR2-KI mice.

Scale bars, 100 µm. **E,** Quantification of SQSTM1/p62 and **F,** NF-H (n = 3-5 mice per

group; data are presented as mean values ± s.e.m., P values from unpaired t test).

AAV9-mediated TECPR2 gene replacement is associated with improvements in

select motor and sensory measures

To determine whether gene replacement could mitigate disease phenotypes, we

delivered a single intracisternal injection of AAV9.pU1a-TECPR2_WT (VCAV-04485,

5.00E+12 GC/mL; 1uL) to *TECPR2*-KI mice and WT littermates at P1–3 (Fig. 3A). We

chose the cisterna magna as the route of administration because of its proximity to the

dorsal column nuclei, the earliest site of pathology, and its established efficacy in

neonatal CNS gene therapy (14). The treatment was well tolerated in WT mice,

supporting safety and long-term tolerability. In KI mice, AAV9-based gene placement

prevented only a subset of the disorder manifestations. Although transient prevention of

body weight loss was noted in males at P30 (Supplementary Fig. 2A), treated KI mice

continued to exhibit reduced body weight at >P90 (Fig. 3B) and excessive grooming

behavior (Fig. 3C). Improvements were observed across several functional measures.

AAV9/TECPR2-treated KI mice showed normalized mechanical sensitivity at >P90 (Fig.

3D), improved hindlimb base width (paw intoeing) at P30 (Fig. 3E), and complete

recovery of acoustic startle response amplitudes to WT levels (Fig. 3F). Together, these

findings indicate that AAV9-mediated TECPR2 expression can partially ameliorate

mechanosensory and brainstem-associated sensorimotor deficits, while growth and

grooming abnormalities remain largely unaffected.

Figure 3. AAV9/TECPR2 mediated gene replacement in *TECPR2*-KI mice is
associated with improvements in select motor and sensory measures. A,
Illustration of experimental regimen. **B,** Relative body weight of TECPR2-KI, WT
littermate, TECPR2-KI+AAV and WT littermate + AAV male and female mice at P90
(n = 17-29 mice per group, data are presented as mean values ± s.e.m., *P* values
from ANOVA followed by Tukey's multiple comparisons test). **C**, Grooming time of
TECPR2-KI, WT littermate, TECPR2-KI+AAV and WT littermate + AAV male and female

mice at >P90 (n = 12-50 mice per group; data are presented as mean
values ± s.e.m., *P* values from ANOVA followed by Tukey's multiple comparisons
test). **D**, Mechanical withdrawal thresholds of *TECPR2*-KI, WT littermate, *TECPR2*-
KI+AAV and WT littermate + AAV male and female mice using an up–down von Frey
method (n = 5-33 mice per group; data are presented as mean values ± s.e.m., *P*
values from one-way ANOVA followed by Dunnett's T3 multiple comparisons test). **E,**
Distance between hindpaws used as a measure of intoeing in male and female
TECPR2-KI, WT littermate, *TECPR2*-KI+AAV and WT littermate + AAV over time
(n = 5-62 mice per group, data are presented as mean values ± s.e.m., *P* values
from two-way ANOVA followed by Šídák's multiple comparisons test). **F,** Maximum
acoustic startle response in male and female *TECPR2*-KI, WT littermate, *TECPR2*-
KI+AAV and WT littermate + AAV (n = 7-30 mice per group, data are presented as
mean values ± s.e.m., **P < 0.05, **P < 0.01, ***P < 0.001* from two-way ANOVA
followed by Tukey's multiple comparisons test).

AAV9-mediated TECPR2 gene placement reduces spheroid pathology and
improves autophagic flux

We next examined whether AAV9-TECPR2 treatment could mitigate axonal and
autophagic defects in *TECPR2*-KI mice. Histological analysis revealed that intracisternal
AAV9 delivery at P1–3 partially prevented spheroid accumulation in the gracile and
cuneate nuclei, reducing their density by approximately 30% at P90 compared with
untreated mutants (Fig. 4A,B). Robust viral-mediated expression of TECPR2 was
confirmed in the brainstem of treated animals (Fig. 4C). Importantly, treated WT
littermates displayed normal histoarchitecture and physiological TECPR2 expression,
indicating that the therapy was safe and well tolerated. Immunohistochemical analyses
further showed improvement of autophagy-related markers in AAV9-treated KI mice.
Accumulation of the autophagy substrate SQSTM1/p62, prominent in untreated
mutants, was similar to WT in the brainstem of treated animals (Fig. 4D,E), while levels
of neurofilament heavy chain (NF-H) were comparable across groups (Fig. 4F).

Figure 4. AAV9/TECPR2 mediated gene replacement in *TECPR2*-KI mice reduces spheroid accumulation and ameliorates autophagic flux at P90. A, Representative H&E staining of coronal brainstem sections of *TECPR2*-KI, WT littermates and *TECPR2*-KI+AAV mice showing (indicated by arrows) spheroid accumulation at P90. Scale bars, 50 and 100 µm. **B,** Quantification of spheroids in brainstem of *TECPR2*-KI, WT littermate, *TECPR2*-KI+AAV and WT littermate + AAV male and female mice (*n* = 4-7 mice per group, data are presented as mean values ± s.e.m., *P* values from one-way ANOVA followed by Dunnett's multiple comparisons test). **C,** RT–qPCR quantification of endogenous and viral TECPR2 in brainstem of *TECPR2*-KI and WT littermates (endogenous), and *TECPR2*-KI+AAV and WT littermate + AAV (viral) male and female mice (*n* = 7-30 mice per group, data are presented as mean values ± s.e.m., *P* values from one-way ANOVA followed by Dunnett's multiple comparisons test). **D,** Representative SQSTM1/p62 staining of brainstem of *TECPR2*-KI, WT littermates and TECPR2-KI+AAV at P90. Scale bars, 100 µm. **E,** Quantification

of SQSTM1/p62 and **F,** NF-H in brainstem of TECPR2-KI, WT littermates and TECPR2-

KI+AAV male and female mice (n = 3-7 mice per group, data are presented as mean

values□±□s.e.m., P values from one-way ANOVA followed by Turkey's multiple

comparison test).

Discussion

TECPR2-related disorder is a rare neurodevelopmental and neurodegenerative disease

for which no disease-modifying therapies currently exist. In this study, we describe a

knock-in mouse model carrying the recurrent *TECPR2* c.1319delC founder mutation

and characterize behavioral, sensory, and neuropathological abnormalities associated

with TECPR2 deficiency. Our data suggest an association between impaired autophagic

flux and selective involvement of brainstem sensory–motor circuits and identify an

acoustic startle deficit as a functional readout relevant to circuit dysfunction. We further

evaluate early postnatal AAV9-mediated *TECPR2* gene delivery and observe partial

improvements in autophagy-related markers, axonal pathology, and select functional

outcomes. Together, these results provide a framework for investigating disease

mechanisms and support further evaluation of gene replacement approaches in

TECPR2-related disorder, particularly to treat brain-stem related disease manifestations

which account for early mortality.

Neurodevelopmental and neurodegenerative disorders caused by impaired autophagy

present a major translational challenge (11,12) as they often combine early circuit

dysfunction with progressive neurodegeneration but lack validated disease models and

actionable outcome measures. *TECPR2*-related disorder exemplifies this gap: despite

clear genetic causation and severe clinical manifestations, therapeutic development has

been limited by disease rarity and the absence of *in vivo* systems that faithfully capture

patient mutations and circuit-level pathology. In this study, we address these limitations

by generating a knock-in mouse model harboring the recurrent *TECPR2* c.1319delC

mutation and using it to define disease mechanisms, identify translatable functional

readouts, and evaluate early gene replacement therapy.

By introducing the patient-derived frameshift mutation into the endogenous mouse
locus, we generated a model that reproduces the predominant loss-of-function
mechanism observed in affected individuals. Unlike prior knockout or spontaneous
mutation models, this knock-in approach preserves the native genomic context while
recapitulating the precise pathogenic allele found in patients, increasing its relevance for
both mechanistic and translational studies. Longitudinal phenotyping revealed that
homozygous *TECPR2*-KI mice develop reduced body weight, increased grooming
behavior, progressive gait abnormalities, and selective tactile sensory impairment, some
of which overlap with clinical features described in *TECPR2*-related disorder (5,15).
These phenotypes evolve over time and can be quantified reproducibly, establishing the
TECPR2-KI mouse as a robust platform for tracking disease progression and
therapeutic response.

A key insight enabled by this model is the identification of selective brainstem circuit
vulnerability as a prominent feature of TECPR2 deficiency. Despite broad TECPR2
expression, neuropathology in our model was largely restricted to sensory–motor
pathways of the brainstem and spinal cord, with progressive accumulation of axonal
spheroids in the gracile and cuneate nuclei and relative sparing of cortical neurons.
These findings are consistent with recent work using a neuropathy-associated *Tecpr2*
knock-in mouse, which revealed endolysosomal loss-of-function phenotypes in neurons
and microglia, supporting a central role for disrupted vesicular and degradative
pathways in *TECPR2*-related disease (16). Together, these studies support a model in
which TECPR2 loss leads to selective vulnerability of long-range projection neurons and
associated circuits, rather than global neurodegeneration.

A key insight enabled by this model is the identification of selective brainstem circuit
vulnerability as a central feature of TECPR2 deficiency. Despite broad TECPR2
expression, neuropathology was strikingly restricted to sensory–motor pathways of the
brainstem and spinal cord, with progressive accumulation of axonal spheroids in the
gracile and cuneate nuclei and relative sparing of cortical neurons. This pattern mirrors
neuropathological findings in patients and other TECPR2-deficient species and supports
a circuit-selective disease mechanism rather than widespread neurodegeneration.

Long-range projection neurons in these pathways likely exhibit heightened vulnerability
due to their exceptional reliance on efficient autophagy to maintain axonal homeostasis.
Consistent with this interpretation, ultrastructural and molecular analyses revealed
marked defects in autophagic processing within affected brainstem nuclei. Accumulation
of abnormal autophagic vacuoles and increased SQSTM1/p62 indicate impaired
autophagosome maturation and clearance, directly linking TECPR2 loss to defective
autophagy and axonal degeneration. These findings clarify how disruption of a core
cellular homeostatic pathway translates into focal circuit dysfunction and highlight
autophagy as a mechanistically grounded therapeutic target.
Beyond previously recognized phenotypes, *TECPR2*-KI mice exhibited a pronounced
reduction in acoustic startle responses, a behavioral deficit not previously reported in
TECPR2-related disorder. The acoustic startle reflex depends on well-defined auditory–
motor brainstem circuits, and its impairment provides functional evidence of brainstem
sensorimotor dysfunction. This finding is particularly relevant given the prominence of
brainstem-related clinical features in patients, including impaired reflexes, dysphagia,
and central apnea. Importantly, the acoustic startle response represents a non-invasive,
quantifiable, and cross-species functional readout, making it a promising biomarker for
disease progression and therapeutic efficacy in future clinical studies.
Using this knock-in mouse model, we next evaluated the effects of early gene
replacement. Neonatal intracisternal delivery of AAV9-*TECPR2* resulted in widespread
transduction of brainstem nuclei and was associated with improvements in autophagy-
related markers, a partial reduction in axonal spheroid burden, and improvements
across multiple functional outcomes.
Mechanosensory thresholds, gait abnormalities, and acoustic startle responses similarly
showed improvement, indicating that early restoration of TECPR2 expression can
ameliorate aspects of brainstem-associated dysfunction. These results provide *in vivo*
evidence that aspects of TECPR2-related pathology can be modified by postnatal gene
replacement. Not all phenotypes were fully prevented. Reduced body weight and
excessive grooming persisted in treated mice at later ages, suggesting contributions
from early developmental alterations, incomplete vector distribution, or non–cell-
autonomous effects involving peripheral or systemic tissues. These findings underscore

the importance of therapeutic timing and raise the possibility that even earlier intervention may be required for full disease modification. They also suggest that combination strategies, such as gene replacement paired with pharmacological modulation of lysosomal or autophagy-related pathways, may further enhance therapeutic efficacy.

More broadly, this work informs understanding of selective vulnerability in neurodevelopmental and neurodegenerative disorders linked to autophagy dysfunction. The relative sparing of cortical neurons despite widespread TECPR2 expression highlights how circuit architecture and metabolic demand shape disease expression. By linking impaired autophagic flux to degeneration of specific brainstem sensory–motor pathways, our findings suggest convergent mechanisms that may extend to other disorders characterized by axonal spheroids and brainstem involvement.

Few limitations should be considered. Our analyses focused primarily on the brainstem and spinal cord, and future studies should evaluate peripheral nerves, forebrain circuits, and systemic metabolic contributions. Only a single AAV dose and delivery route were tested, and comparative studies using alternative delivery strategies will be important for optimizing translational approaches. Finally, while the acoustic startle deficit emerges as a promising biomarker, its relevance in patients remains to be established through targeted clinical evaluation.

In summary, we developed a knock-in mouse model carrying the recurrent c.1319delC founder mutation that reproduces the underlying genetic alteration and exhibits circuit-selective pathology and functional impairments associated with TECPR2 deficiency. This model enabled the identification of brainstem sensory–motor circuits as a prominent site of vulnerability and revealed an acoustic startle phenotype that may serve as a translatable functional readout. In addition, early AAV9-mediated TECPR2 delivery was associated with improvement of autophagy-related markers and partial amelioration of select behavioral outcomes. By combining a genetically defined mouse model with quantifiable functional endpoints and a clinically established gene delivery strategy, this study provides a preclinical framework to evaluate TECPR2 gene replacement and to inform future investigations of therapeutic timing, target engagement, and outcome measures.

Materials and Methods

Generation of the TECPR2 Knock-In Mouse Model

TECPR2 knock-in (KI) mice were generated at the Harvard Genome Modification Facility. CRISPR/Cas9/guide RNA and donor oligo were injected into fertilized C57BL/6 eggs to generate the equivalent HSP50 exon 8 mutation in the mice. This mutation is a deletion of a single C nucleotide in mouse exon 8: NM_001081057.2; c.1319delC, p.Ser440Serfs19*. Male F1 heterozygous mice were transferred to Charles River Laboratories, where they used these mice for IVF of oocytes from female C57BL/6 to generate heterozygous breeders and used to set up a breeding colony. Mice were genotyped using PCR and sequencing by the Molecular Genetics Core Facility at Boston Children's Hospital to confirm the presence of the mutation. TECPR2 KI mice were housed in standard clear plastic cages with no more than 5 animals per cage under controlled conditions (lights on 07:00-19:00; humidity 30%-50%; temperature 22-23°C) with ad libitum access to food and water. All experiments were performed between 9:00 and 17:00 in a room maintained at a temperature of 21 ± 1°C. All experimental protocols comply with relevant ethical regulations and were approved by Boston Children's Hospital Institutional Animal Care and Use Committee (protocol 00002125). Animals were randomized to treatment groups.

AAV production

Through the Luke Heller TECPR2 Foundation, we have obtained WT human TECPR2and GFP-AAV9 constructs, generated at the Horae Gene Therapy Center, UMass Chan Medical School (Worcester, MA, USA) under the direction of Dr. Guangping Gao. The AAV genome is composed of U1a promoter, TECPR2 CDS, a synthetic short polyA and AAV2 ITRs at both ends. The genome was packaged into the AAV9 capsid by triple-transfection in HEK293 cells.

In vivo injection of viral vectors

For AAV9 delivery into the CSF via cisterna magna, mice at postnatal day 1–3 (P1–P3)
were anesthetized by rapid induction of hypothermia for 2–4 min on ice water until loss
of consciousness. The injection site was disinfected with 70% ethanol. A 33-gauge
needle attached to a Hamilton syringe was lowered approximately 2 mm into the
cisterna magna at a 45° angle, and 1 µL of viral solution (VCAV-04485, 5.00E+12
GC/mL) was slowly injected. Body temperature was gradually restored and maintained
on a 37 °C warming pad for 10 min after injection before returning pups to the parental
cage. Injections were performed prior to genotyping and blinded to genotype; all pups
within each litter received the same viral vector.

Body Weight and Growth Curve Analysis
Body weights of the TECPR2 KI mice and their WT littermates were recorded with an
electronic scale at postnatal day 30, 60 and 90 (P30, P60 and P90, respectively).
Growth curves were plotted to compare the development of mutant and WT mice.

Mechanical Sensitivity Testing
Mechanical sensitivity was assessed using the von Frey filament test. Mice were placed
in individual chambers on a mesh platform, and calibrated filaments were applied to the
plantar surface of the left hind paw, with a positive response consisting of a paw lifting
or flinching response to the fiber. The response patterns were collected and converted
into corresponding 50% withdrawal thresholds using the Up–Down Reader software and
associated protocol (17). All tests were performed by investigators blinded to the
genotype and treatment.

Acoustic startle reflexes.
Mice were tested for startle reflexes in response to broadband auditory stimulation at
varying intensities as previously described (18). The animals were tested in a sound
shielded startle booth (Kinder Scientific) and the force generated by foot movement was
sensed by piezoelectric motion sensor fixed beneath an elevated platform. Mice were
placed in a smaller sub chamber anchored to the topside of the platform, which
restricted them from rearing on their limbs but freely permitted penetration of the sound

stimulus. Sound stimuli were calibrated with the door closed using a sound pressure
level meter (Allied Electronics) with the microphone mounted in the position normally
occupied by the animal holder. The generation of sound stimuli and recording of force
amplitude signals was performed by startle monitor software (Kinder Scientific).
Broadband white noise was presented at a background level of 60 dB throughout the
experiment and auditory test stimuli consisted of 50-ms broadband white noise pulses in
10-dB steps from 60 to 120 dB. Different intensities of the test stimulus were presented
in a pseudo-random order at randomized inter trial intervals (ITI) that varied between 8
and 22 s, with no ITI repeated more than three times. Five repetitions were averaged for
each of the intensities for one test subject and responses were normalized to the weight
of the mouse tested. Startle response measurements were performed by investigators
blinded to the genotype.
481

Intoeing and grooming behaviors
Intoeing was quantified using a novel bottom-up imaging and data acquisition
technology and analysis platform that provides automated, quantitative, and objective
measures of naturalistic rodent behavior in an observer-independent and unbiased
fashion as previously described (19). The location of the paws was extracted from body
frames using the DeepLabCut and the distance between hindpaws (measure of
intoeing) in the unit of pixel distance was calculated. Grooming behaviors were
calculated by the time the animal spent grooming the face and body in seconds for 10
minutes recording using the ARBEL algorithm, a machine learning tool with light-based
image analysis for automatic classification of 3D behaviors (20).
492

Tissue processing and histopathological analysis
Animals were anesthetized with pentobarbital and perfused with a 4%
paraformaldehyde (PFA) solution in PBS. The brain was dissected and fixed in 4% PFA
solution overnight. Samples were then transferred to a 30% sucrose solution until the
tissue sank. Samples were immediately frozen in OCT embedding medium and kept in -
80oC until further processing. Coronal cryosections (50 μm) were collected through the
brainstem and spinal cord for immunostaining and histopathological analysis. Slices

were then stained with hematoxylin and eosin (H&E) to identify spheroids or probed with
antibodies for neurofilament and autophagic flux. Quantification of spheroids was done
measuring the density of spheroids per mm2 in the cuneate and gracile nuclei areas
bilaterally in 10-12 sections per animal, then an average density was calculated using
QuPath software (21).
505

Immunofluorescence staining

Sections were washed in PBS, permeabilized with PBS + 0.3% Triton X-100 (PBST-0.3)
for 15 min and blocked in PBST-0.3 containing 5% normal goat serum (NGS) for 1 h at
room temperature. Samples were then incubated overnight at 4 °C with primary
antibodies in blocking buffer (Mouse anti-SQSTM1/p62; 1:500; Cell Signaling
Technologies #88588; rabbit anti-neurofilament heavy chain; 1:1,000; Sigma Aldrich
#N4142). After washes in PBST-0.3, sections were incubated with secondary antibodies
for 2 h at room temperature (1:500; Alexa Fluor–conjugated goat anti-rabbit or goat anti-
mouse, Invitrogen). Nuclei were counterstained with DAPI (0.5 μg/mL). Sections were
mounted with ProLong Glass antifade reagent and imaged on a Zeiss LSM880 confocal
microscope. Neurofilament (NF-H) intensity and SQSTM1/p62+ puncta were measured
from epifluorescent images. NF-H intensity was measured using sections from WT
tissues to set a threshold and the mean pixel intensity from WT and TECPR2 KI tissues
was measured using Zeiss Zen 3.8 Software. The quantification of SQSTM1/p62 puncta
was done by measuring the area of the cuneate and gracile from epifluorescent images
and manually detecting SQSTM1/p62+ puncta in those regions using Zeiss Zen 3.8
Software. Quantitative analysis was performed by an investigator who was blind to the
experimental conditions.
524

Electron microscopy

526 Animals were anesthetized with pentobarbital and perfused with 2.5% PFA, 2.5%
527 glutaraldehyde, in 0.1 M cacodylate buffer pH 7.4. Brain samples were fixed with 2.5%
528 PFA, 2.5% glutaraldehyde, in 0.1 M cacodylate buffer containing 5 mM $CaCl_2$ (pH 7.4),
529 post-fixed in 1% osmium tetroxide supplemented with 0.5% potassium
530 hexacyanoferrate trihydrate and potassium dichromate in 0.1 M cacodylate for 1 h,

stained with 2% uranyl acetate in double-distilled water for 1 h, dehydrated in graded ethanol solutions, and embedded in epoxy resin (Agar Scientific, AGR1030). Ultrathin sections (70–90 nm) of brainstem, obtained with a LeicaUltracut UCT microtome, were stained with lead citrate and then examined using either a Phillips CM-12 transmission electron microscope (TEM) equipped with a Gatan One View camera or an FEI Tecnai SPIRIT TEM equipped with a bottom-mounted 2 k × 2 k FEI Eagle CCD camera.

Statistical Analysis

Data are expressed as the mean ± SEM. Significant differences between groups were assessed with one or two-way ANOVA; once the significance of the group differences ($p \leq 0.05$) was established, appropriate post-hoc comparisons between pairs of groups were used. Statistical analyses were done using GraphPad Prism Software.

List of Supplementary Materials

Materials and Methods

Fig. S1 to S3.

Acknowledgments

Luke Heller TECPR2 Foundation

Jordan Avi Ogman Foundation

Quiver Bioscience [formerly Q-State Biosciences]

Animal Behavior and Physiology Core at Boston Children's Hospital

Molecular Genetics Core Facility at Boston Children's Hospital

Cellular Imaging Core at Boston Children's Hospital

Neurobiology Imaging Facility at Harvard Medical School

Electron Microscopy Core at Harvard Medical School

References

1.   D. Oz-Levi, B. Ben-Zeev, E. K. Ruzzo, Y. Hitomi, A. Gelman, K. Pelak, Y.
Anikster, H. Reznik-Wolf, I. Bar-Joseph, T. Olender, A. Alkelai, M. Weiss, E. Ben-Asher,
D. Ge, K. V. Shianna, Z. Elazar, D. B. Goldstein, E. Pras, D. Lancet, Mutation in
TECPR2 reveals a role for autophagy in hereditary spastic paraparesis. Am. J. Hum.
Genet. 91, 1065–1072 (2012).

2.   P. P. Zhu, K. R. Denton, T. M. Pierson, X. J. Li, C. Blackstone, Mutations in
TECPR2, encoding a positive regulator of autophagy, cause a new syndrome of
childhood-onset spasticity and axonal neuropathy. Am. J. Hum. Genet. 97, 671–682
(2015).

3.   G. Heimer, D. Oz-Levi, E. Eyal, S. Edvardson, A. Nissenkorn, E. K. Ruzzo, A.
Szeinberg, C. Maayan, M. Mai-Zahav, O. Efrati, E. Pras, H. Reznik-Wolf, D. Lancet, D.
B. Goldstein, Y. Anikster, S. A. Shalev, O. Elpeleg, B. Ben Zeev, TECPR2 mutations
cause a new syndrome of progressive neurodegeneration with developmental delay.
Brain 139, 4034–4044 (2016).

4.   S. Anazi, S. Maddirevula, V. Salpietro, Y. T. Asi, S. Alsahli, A. Alhashem, H. E.
Shamseldin, F. AlZahrani, N. Patel, N. Ibrahim, F. M. Abdulwahab, M. Hashem, N.
Alhashmi, F. Al Murshedi, A. Al Kindy, A. Alshaer, A. Rumayyan, S. Al Tala, W. Kurdi,
A. Alsaman, A. Alasmari, S. Banu, T. Sultan, M. M. Saleh, H. Alkuraya, M. A. Salih, H.
Aldhalaan, T. Ben-Omran, F. Al Musafri, R. Ali, J. Suleiman, B. Tabarki, A. W. El-
Hattab, C. Bupp, M. Alfadhel, N. Al Tassan, D. Monies, S. T. Arold, M. Abouelhoda, T.
Lashley, H. Houlden, E. Faqeih, F. S. Alkuraya, Expanding the genetic heterogeneity of
intellectual disability. Hum. Genet. 136, 1419–1429 (2017).

5.   S. Neuser, B. Brechmann, G. Heimer, I. Brösse, S. Schubert, L. O'Grady, M.
Zech, S. Srivastava, D. A. Sweetser, Y. Dincer, V. Mall, J. Winkelmann, C. Behrends, B.
T. Darras, R. J. Graham, P. Jayakar, B. Byrne, B. E. Bar-Aluma, Y. Haberman, A.
Szeinberg, H. M. Aldhalaan, M. Hashem, A. Al Tenaiji, O. Ismayl, A. E. Al Nuaimi, K.
Maher, S. Ibrahim, F. Khan, H. Houlden, V. S. Ramakumaran, A. T. Pagnamenta, J. E.
Posey, J. R. Lupski, W. H. Tan, G. ElGhazali, I. Herman, T. Muñoz, G. M. Repetto, A.

Seitz, M. Krumbiegel, M. Poli, U. Kini, S. Efthymiou, J. Meiler, R. Maroofian, F. S. Alkuraya, R. Abou Jamra, B. Popp, B. Ben-Zeev, D. Ebrahimi-Fakhari, Clinical, neuroimaging, and molecular spectrum of TECPR2-associated hereditary sensory and autonomic neuropathy with intellectual disability. Hum. Mutat.42, 762–776 (2021).

6. M. Fraiberg, B. C. Tamim-Yecheskel, K. Kokabi, N. Subic, G. Heimer, F. Eck, K. Nalbach, C. Behrends, B. Ben-Zeev, O. Shatz, Z. Elazar, Lysosomal targeting of autophagosomes by the TECPR domain of TECPR2. Autophagy 17, 3096–3108 (2021).

7. S. Maday, E. L. J. Holzbaur, Autophagosome biogenesis in primary neurons follows an ordered and spatially regulated pathway. Dev. Cell 30, 71–85 (2014).

8. J. H. Hurley, L. N. Young, Mechanisms of autophagy initiation. Annu. Rev. Biochem. 86, 225–244 (2017).

9. M. Komatsu, S. Waguri, T. Chiba, S. Murata, J. I. Iwata, I. Tanida, T. Ueno, M. Koike, Y. Uchiyama, E. Kominami, K. Tanaka, Loss of autophagy in the central nervous system causes neurodegeneration in mice. Nature 441, 880–884 (2006).

10. D. Stadel, V. Millarte, K. D. Tillmann, J. Huber, B. C. Tamim-Yecheskel, M. Akutsu, A. Demishtein, B. Ben-Zeev, Y. Anikster, F. Perez, V. Dötsch, Z. Elazar, V. Rogov, H. Farhan, C. Behrends, TECPR2 cooperates with LC3C to regulate COPII-dependent ER export. Mol. Cell 60, 89–104 (2015).

11. D. Ebrahimi-Fakhari, A. Saffari, L. Wahlster, J. Lu, S. Byrne, G. F. Hoffmann, H. Jungbluth, M. Sahin, Congenital disorders of autophagy: An emerging novel class of inborn errors of neuro-metabolism. Brain 139, 317–337 (2016).

12. H. S. Dafsari, D. Martinelli, A. Saffari, D. Ebrahimi-Fakhari, M. Fanto, C. Dionisi-Vici, H. Jungbluth, An update on autophagy disorders. J. Inherit. Metab. Dis. 48, e12798 (2025).

13. L. A. Williams, D. J. Gerber, A. Elder, W. C. Tseng, V. Baru, N. Delaney-Busch, C. Ambrosi, G. Mahimkar, V. Joshi, H. Shah, K. Harikrishnan, H. Upadhyay, S. H.

Rajendran, A. Dhandapani, J. Meier, S. J. Ryan, C. Lewarch, L. Black, J. Douville, S. Cinquino, H. Legakis, K. Nalbach, C. Behrends, A. Sato, L. Galluzzi, T. W. Yu, D. Brown, S. Agrawal, D. Margulies, A. Kopin, G. Dempsey, Developing antisense oligonucleotides for a TECPR2 mutation-induced, ultra-rare neurological disorder using patient-derived cellular models. Mol. Ther. Nucleic Acids 29, 189–203 (2022).

14.   R. M. Bailey, D. Armao, S. Nagabhushan Kalburgi, S. J. Gray, Development of intrathecal AAV9 gene therapy for giant axonal neuropathy. Mol. Ther. Methods Clin. Dev. 9, 160–171 (2018).

15.   G. Heimer, S. Neuser, B. Ben-Zeev, D. Ebrahimi-Fakhari, TECPR2-related hereditary sensory and autonomic neuropathy with intellectual disability. In: Adam MP, Bick S, Mirzaa GM, Pagon RA, Wallace SE, Amemiya A, editors. GeneReviews® [Internet]. Seattle (WA): University of Washington; 1993–2026 (updated 2022).

16.   D. Bhattacharya, P. da Silva-Buttkus, K. Nalbach, L. Cheng, L. Garrett, M. Irmler, G. Kislinger, G. Werner, R. Rodde, C. Lengger, J. Beckers, A. Zimprich, S. M. Hölter, V. Gailus-Durner, H. Fuchs, M. Hrabe de Angelis, B. Wefers, W. Wurst, M. S. Brill, M. Schifferer, S. F. Lichtenthaler, C. Behrends, Neuropathy-associated Tecpr2 mutation knock-in mice reveal endolysosomal loss of function phenotypes in neurons and microglia. Cell Death Dis. 16, 775 (2025).

17.   R. Gonzalez-Cano, B. Boivin, D. Bullock, L. Cornelissen, N. Andrews, M. Costigan, Up-Down Reader: An Open Source Program for Efficiently Processing 50% von Frey Thresholds. Front. Pharmacol. 9, 433 (2018).

18.   C. A. Nist-Lund, B. Pan, A. Patterson, Y. Asai, T. Chen, W. Zhou, H. Zhu, S. Romero, J. Resnik, D. B. Polley, G. S. Géléoc, J. R. Holt, Improved TMC1 gene therapy restores hearing and balance in mice with genetic inner ear disorders. Nat. Commun. 10, 236 (2019).

19.   Z. Zhang, D. P. Roberson, M. Kotoda, B. Boivin, J. P. Bohnslav, R. González-Cano, D. A. Yarmolinsky, B. L. Turnes, N. K. Wimalasena, S. Q. Neufeld, L. B. Barrett,

N. L. M. Quintão, V. Fattori, D. G. Taub, A. B. Wiltschko, N. A. Andrews, C. D. Harvey,

S. R. Datta, C. J. Woolf, Automated preclinical detection of mechanical pain

hypersensitivity and analgesia. Pain 163, 2326–2336 (2022).

20.   O. Barkai, B. Zhang, B. L. Turnes, M. Arab, D. A. Yarmolinsky, Z. Zhang, L. B.

Barrett, C. J. Woolf, A machine learning tool with light-based image analysis for

automatic classification of 3D pain behaviors. Cell Rep. Methods 5, 101145 (2025).

21.   P. Bankhead, M. B. Loughrey, J. A. Fernández, Y. Dombrowski, D. G. McArt, P.

D. Dunne, S. McQuaid, R. T. Gray, L. J. Murray, H. G. Coleman, J. A. James, M. Salto-

Tellez, P. W. Hamilton, QuPath: Open source software for digital pathology image

analysis. Sci. Rep. 7, 16878 (2017).