Main Manuscript for

Myosin binding protein-C limits strain induced cross-bridge detachment in

response to rapid stretch in cardiac and skeletal muscle

Nichlas M. Engels[1], Rachel L. Sadler[2], Michel N. Kuehn[3,4], Devin L. Nissen[5], Dylan

Reichert[6], Marius Meinhold[7,8],Wolfgang A. Linke[3], Weikang Ma[5,9,10], Anthony L.

Hessel[3,4], and Samantha P. Harris[2]*

[1]Department of Cellular and Molecular Medicine, University of Arizona, Tucson, AZ

[2]Department of Physiology, University of Arizona, Tucson, AZ

[3]Institute of Physiology II, University of Muenster, Muenster, Germany

[4]Accelerated Muscle Biotechnologies, Mansfield, MA, USA

[5]BioCAT, Department of Biology, Illinois Institute of Technology, Chicago, IL, USA

[6]Department of Biomedical Engineering, University of Arizona, Tucson, AZ, USA

[7]Department of Comparative Biosciences, University of Wisconsin–Madison, Madison,

WI, USA.

[8]School of Veterinary Medicine, University of Wisconsin–Madison, Madison, WI, USA.

[9]Center for Synchrotron Radiation Research and Instrumentation, Illinois Institute of

Technology, Chicago, IL, USA

[10]Pritzker Institute of Biomedical Science and Engineering, Illinois Institute of

Technology, Chicago, IL, USA

* Samantha P. Harris

Email: samharris@arizona.edu

Author Contributions: N.M.E, A.L.H, and S.P.H conceptualized the project, , N.M.E conducted the investigation, N.M.E visualized the datasets, S.P.H was the project administrator, A.L.H, and S.P.H supervised the study, N.M.E wrote the original draft, all authors reviewed, edited, and agreed to the final draft.

Competing Interest Statement: W.M. consults for Edgewise Therapeutics, Cytokinetics Inc. and Kardigan Bio, but these activities have no relation to the current work. A.L.H. and M.N.K. are founders and owners of Accelerated Muscle Biotechnologies, but that work is not related to the current project. All other authors declare no competing interests.

Classification:

Biological Sciences (Major)

Minor; Physiology

Keywords: myosin binding protein-C, stretch activation, cross-bridge kinetics, mouse models

This PDF file includes:

       Main Text
       Figures 1 to 6

Abstract

Myosin binding protein-C (MyBP-C) consists of a family of regulatory proteins expressed in sarcomeres of cardiac, fast and slow twitch skeletal muscles. The 3 MyBP-C paralogs expressed in each muscle type are encoded by separate genes but maintain a similar structure. Given the overall similarity in structure and localization of each of paralog, it is assumed that MyBP-C expressed in different muscles have similar functional effects. Here we directly tested this assumption by making use of our cut and paste approach to remove and replace N-terminal regions of MyBP-C in sarcomeres of different muscle types. We found that the different MyBP-C paralogs similarly slowed cross-bridge cycling kinetics, increased Ca^{2+} sensitivity of tension, and damped force oscillations. However, responses to a rapid stretch in actively contracting fibers, taken as indices of cross-bridge detachment and attachment kinetics, differed in each muscle type and responses depended on the presence or absence of a given paralog of MyBP-C. Altered responses to stretch were most evident for fast MyBP-C where loss of MyBP-C in psoas muscle resulted in transient responses to stretch that resembled those found in cardiomyocytes. Replacement of cardiac MyBP-C with fast MyBP-C in cardiomyocytes led to responses similar to psoas muscle. In separate X-ray diffraction experiments we also found that loss of MyBP-C in Ca^{2+}-activated psoas muscle increased lattice disorder, reduced the ordering of myosin heads, and decreased thin filament length. Taken together, these results indicate that the different MyBP-C paralogs exert both common and unique effects on myosin cross-bridge kinetics.

Significance Statement

MyBP-C is a family of regulatory proteins found in muscle sarcomeres, where they

regulate contraction and relaxation. Mutations in all MyBP-C paralogs cause disease in

skeletal and cardiac muscles. We used a powerful "cut and paste" strategy to selectively

remove MyBP-C from slow-twitch, fast-twitch, and cardiac muscle to show that each

MyBP-C effects cross-bridge behavior similarly, though to varying degrees. Each MyBP-

C had a notable effect on transient responses to rapid stretch, where MyBP-C was found

to limit strain-induced cross-bridge detachment, especially in fast-twitch muscles. Strain-

induced cross-bridge detachment is critical for rapid filling of the left ventricle in diastole

and for sustained contraction in skeletal muscle. MyBP-C paralogs appear adapted to

meet the mechanical demands of each muscle type.

74
75   **Main Text**
76
77   *Note: Not all manuscripts need to follow the standard section order or headings outlined below.*
78   *Authors are welcome to choose headings and subheadings that best fit their work. Materials and*
79   *Methods should appear after the Results and Discussion in most cases.*
80
81   **Introduction**
82
Myosin binding protein-C (MyBP-C) consists of a family of regulatory proteins located in

sarcomeres of striated muscle. Within this family, there exist three unique paralogs of

MyBP-C, each encoded by a separate gene and expressed in specific fiber types. *Mybpc1*

encodes the slow twitch skeletal paralog (sMyBP-C), *Mybpc2* encodes the fast twitch

skeletal paralog (fMyBP-C), and *Mybpc3* encodes the cardiac paralog (cMyBP-C)(1) The

three paralogs share significant structural homology, as they all consist of a chain of 7

immunoglobulin (Ig)-like domains and 3 fibronectin type III domains termed C1-C10, a

regulatory motif referred to as the M-domain, and a proline/alanine-rich domain near the

N-terminus(1). Cardiac (cMyBP-C) contains an additional 8th Ig-like domain at the N-

terminus termed C0 (Figure 1). Each paralog is located in the C-zone of sarcomeres,

where C-terminal domains C8-C10 anchor MyBP-C to the thick filament backbone and

interact with myosin heads of crown 1 and 3 in relaxed muscle(2, 3). The N-terminal

domains (C0-C2 for cardiac and C1-C2 for skeletal) interact with both thin and thick

filaments(4-7). For example, N-terminal interactions with the thin filament shift the

tropomyosin cable from the closed to the blocked state, thus promoting activation (i.e.

cross-bridge formation) in the presence of Ca^{2+} or delaying relaxation as Ca^{2+} falls(8, 9).

In contrast, N-terminal domain interactions with the thick filament have an inhibitory effect

on myosin heads by stabilizing myosin heads in the OFF state(10, 11). While less is known

regarding the middle domains (C3-C7) of MyBP-C, recent evidence suggests they play a

critical auto-inhibitory role limiting the effects of N-terminal domains to activate the thin

filament(12, 13). However, neither the position of the N-terminal domains nor the middle

domains were resolved in recent cryo-EM reconstructions and so the structural orientation

and interactions of these domains in the sarcomere are still unclear.

As evidence for MyBP-C's structural and regulatory roles in the sarcomere, mutations in

all three MyBP-C genes lead to disease in their respective muscle types. For instance,

mutations in *Mybpc3* are a leading cause of hypertrophic cardiomyopathy (HCM)(14-16),

while mutations in *Mybpc1* and *Mybpc2* are linked to distal arthrogryposis (DA1) and

myogenic muscle tremors(17-19). Although the three proteins share significant structural

homology, sequence homology between the paralogs is low, sharing only ~39% sequence

identity between the three paralogs in human muscles(20). Additionally, sMyBP-C exists

as a subfamily of multiple splice variants which likely have significant functional differences

that have yet to be thoroughly characterized(21).

Here we sought to compare the relative structural and functional effects of the 3 main

MyBP-C paralogs using our cut and paste approach to systematically remove and replace

MyBP-C in each of the different muscle types. The approach used 3 lines of gene-edited

mice, referred to as $SpyC_1$, $SnoopC_2$, and $SpyC_3$, that allow selective removal and

replacement of MyBP-C in slow twitch skeletal, fast twitch skeletal, and cardiac muscles,

respectively (Figure 1)(13, 22-24). The unique strength of these mouse models is that they

allow for a direct investigation of both the structural and mechanical role of MyBP-C within

a single fiber type, where MyBP-C can be selectively removed and replaced in a single,

healthy muscle fiber after it has been isolated from the animal. This is because all 3 mouse

lines contain a tobacco etch virus (TEV) protease recognition site followed by either a

short SpyTag or SnoopTag sequence between domains C7 and C8(22, 23, 25, 26)

(Supplemental Figure 1). Upon treatment of permeabilized muscle fibers with TEV

protease (TEVp), the endogenous domains C1-C7 in skeletal muscle and C0-C7 in

cardiac muscle are selectively cleaved and eliminated from the muscle, while domains

C8-C10 remain anchored to the thick filament. Following cleavage of MyBP-C, added

recombinant proteins containing a cognate SpyCatcher (Sc) or SnoopCatcher(SnC) form

a spontaneous covalent bond with the SpyTag or SnoopTag encoded at the N-terminus

of C8, thus permitting covalent ligation ("paste") of any desired recombinant protein *in situ*

at the position of endogenous MyBP-C in the sarcomere (Figure 1).

Using these gene-edited mouse lines, we performed permeabilized muscle mechanics to

measure changes in isometric force, cross-bridge cycling and stretch activation, as well

as small-angle X-ray diffraction to identify structural changes to the sarcomere during

isometric contraction. Here, we found that each MyBP-C paralog shares certain functional

effects including sensitizing myofilaments to Ca^{2+}, slowing cross-bridge cycling kinetics,

and damping force oscillations at submaximal Ca^{2+} activation. However, we found that

each paralog differentially regulates tension responses following a quick stretch,

apparently by differentially affecting strain-induced cross-bridge detachment. In psoas

muscle these changes correlated with increased thick and thin filament disorder during

half-maximal isometric contraction.

144
145　**Results**
146
MyBP-C enhances Ca^{2+}-sensitivity of tension and limits cross-bridge

redevelopment rates (k_{tr}) in cardiac muscle and fast twitch and slow twitch muscles

To determine the contribution of each MyBP-C paralog to function in the different muscle

fiber types (cardiac, fast skeletal, and slow skeletal), we first investigated the effect of the

different MyBP-C paralogs on steady state Ca^{2+}-sensitivity of tension as well as the rate

of cross-bridge cycling. Both Ca^{2+}sensitivity and crossbridge cycling, the latter of which is

indicated by tension recovery from a slack-restretch maneuver, (k_{tr})(27) are both two

parameters that are frequently altered under disease conditions. For these experiments,

each MyBP-C paralog was selectively cleaved in cardiac left ventricle, fast twitch psoas,

and slow twitch soleus muscles from SpyC$_3$, SnoopC$_2$, and SpyC$_1$ mice, respectively, by

incubating permeabilized muscle fibers with TEV protease (TEVp). Results shown in

Figure 2 demonstrate that loss of the N'-terminal domains caused a rightward shift of the

tension-pCa curve, indicating reduced Ca^{2+}-sensitivity of tension such that higher [Ca^{2+}] is

required to achieve a given amount of force. In SpyC$_3$ cardiac muscle the pCa$_{50}$ for tension

(where pCa50=-log[Ca^{+}]) was shifted from 5.401 ± 0.01 prior to TEVp treatment to

5.32 ± 0.01 after treatment (n=13, $p < 0.001$) (Figure 2a). Similarly, in SnoopC$_2$ fast twitch

psoas muscle, the pCa$_{50}$ was shifted from 5.63 ± 0.01 to 5.57 ± 0.01, (n=10, $p < 0.001$)

(Figure 2b). In SpyC$_1$ slow twitch soleus we did not observe a change in Ca^{2+}-sensitivity.

(5.38 ± 0.02 to 5.36 ± 0.02,n=8, n.s) (Figure 2c). However, since mouse soleus muscle

expresses a near even mixture of slow and fast twitch fiber type populations with variable

MyBP-C paralogs, (28) the lack of a significant difference in Ca^{2+}-sensitivity of tension for

soleus muscle may reflect a relatively low proportion of sMyBP-C relative to total MyBP-

C. However, k_{tr} was faster at sub-maximal levels of Ca^{2+} activation in all 3 muscle types,

indicating an increase in cross-bridge cycling rates at these $[Ca^{2+}]$ (Figure 2d-f). The

increased cycling suggests a change in either cross-bridge attachment or detachment

rates.

Loss of MyBP-C promotes the occurrence of spontaneous oscillatory contractions

(SPOC) in all muscle types.

Another common observation identified in each muscle type following cleavage of MyBP-

C by TEVp was the appearance of spontaneous oscillatory contractions (SPOC) that

became evident during Ca^{2+}-activated contraction(29). The SPOC oscillations are evident

as waves of contraction and relaxation that occur spontaneously, i.e., without cyclic

changes in activating Ca^{2+} (see Supplemental Videos 1 and 2). Recently, the SPOC

phenomenon has been proposed as an underlying mechanism of myogenic tremor in

certain skeletal myopathies linked to mutations in MyBP-C(19). We have also previously

reported SPOC behavior in detergent-permeabilized cardiomyocytes from $SpyC_3$ left

ventricles that had been treated with TEVp to remove the C0-C7 domains of cardiac

MyBP-C. Importantly, the oscillations were specifically due to loss of cMyBP-C because

they were eliminated by ligation of recombinant cMyBP-C(22). Here we investigated

whether SPOC is a common phenomenon following removal of MyBP-C in both slow and

fast twitch muscles in addition to cardiomyocytes. Figure 2g-i shows raw force traces from

submaximal Ca^{2+}-activated force contractions in cardiac tissue from $SpyC_3$, in psoas

muscles from $SnoopC_2$ mice and in soleus muscle from $SpyC_1$ mice before and after

treatment with TEVp. Results show that removal of C1-C7 domains from both $SnoopC_2$

psoas and $SpyC_1$ soleus muscle consistently resulted in robust SPOC behavior which was

evident as force overshoots and instabilities in the raw force traces during Ca^{2+} activation

(see also Supplemental videos S1 and S2). Similar to SPOC observed in cardiomyocytes,

SPOC in skeletal fibers occurred most commonly at submaximal $[Ca^{2+}]$ near the pCa_{50} for

force generation. Therefore, MyBP-C paralogs appear critical in preventing SPOC from

occurring in both cardiac and skeletal muscles where undamped or prolonged SPOC may

contribute to muscle dysfunction in disease.

199   MyBP-C paralogs differentially regulate muscle responses to rapid stretch during

200 Ca^{2+} activation

We next investigated whether MyBP-C affects rapid tension responses following stretch

in Ca^{2+} activated fibers because these responses give insights into cross-bridge

detachment/attachment events as well as the delayed recovery of force referred to as SA.

For these experiments, muscles were subjected to a rapid 2% stretch relative to initial

muscle length followed by a 3 sec hold at the new length before returning to the initial

length. The resulting transient responses to the acute stretch are typically interpreted in

terms of cross-bridge behaviors(30, 31). As shown in Figure 3, the overall shape of the

resulting tension transients in response to quick stretch varied significantly depending on

the muscle type (i.e., responses to rapid stretch differed in cardiac, slow twitch, or fast

twitch skeletal muscles).

Figure 1 shows representative traces from cardiac muscle and psoas muscle that

illustrate the 3 typical phases of the transient force responses to stretch. In cardiac muscle

following rapid stretch there is a rapid increase in force (P_1) that coincides with and is

proportional to the initial stretch/increase in length. P_1 is related to both strain on bound

cross-bridges as well as strain on elastic elements within the sarcomere, such as titin(32).

Following the P_1 peak, the force drops rapidly in such a way that the rate of the decrease

in tension (K_{rel}) is related to the kinetics of cross-bridge detachment. Next, the nadir of the

tension decrease is termed P_2 and typically approaches or drops below the pre-stretch

steady state tension (33). Following P_2, there is a delayed rise in tension where tension

increases to a new steady-state value that is greater than the pre-stretch steady state

force (P_3). This rise to a new steady state force (P_3) termed SA (33-35), is most prominent

in muscles that do oscillatory work such as some insect flight muscles and vertebrate

cardiac muscle, whereas it is typically less prominent and shorter lived in skeletal

muscle(34, 36). In fast twitch skeletal muscle, the rapid increase to P1 is similar to cardiac

muscle. However, there is a notable difference following the initial relaxation between

cardiac and fast twitch skeletal muscle, where P_2 typically remains greater than the steady

state force prior to stretch(34). Next, instead of a delayed rise in tension to a new steady

state (P_3), the force briefly plateaus before slowly decreasing to a steady-state. Since the

fast twitch muscle tension transient typically lacks a delayed regeneration of force, the SA

response is considered minimal in fast twitch skeletal muscle.

Here, we found that the tension responses to rapid stretch were affected following removal

of MyBP-C in each muscle, albeit, to varying degrees. A representative trace for each

muscle type before and after removal of MyBP-C is shown in Figure 3. Results show that

removal of cMyBP-C in cardiomyocytes had overall modest effects on tension transients,

i.e. loss of cMyBP-C slightly reduced the P_2 nadir (by ~4% at pCa 4.5 and 5.2), increased

the P_{df} amplitude (P_3-P_2) by ~4% at pCa_{50}, and increased k_{rel}, the rate of relaxation

following the P_1 peak (97.56 ± 7.37 to 129.37 ± 8.20 at pCa_{50}) (Fig. 3d,g,j). However,

effects of removing fMyBP-C in psoas muscle were much more profound where tension

transients following loss of fMyBP-C (Figure 3b) now closely resembled a typical transient

response seen in cardiac muscle. That is, removal of fMyBP-C significantly decreased the

P_2 nadir below the pre-stretch steady state and increased the P_{df} amplitude. K_{rel} was also

significantly increased at sarcomere length (SL) 3.0 (Supplementary Figure 2), although

K_{rel} remained unchanged at SL 2.4 possibly due to a lower rate of data acquisition for this

data set (Figure 3e,h,k). Most strikingly, the acute removal of fMyBP-C resulted in the

appearance of a new delayed phase of force regeneration (phase 3) that is usually minimal

or absent in fast twitch fibers. This was primarily due to the decrease in P_2 because the

P_3 amplitudes remained unchanged in cardiac and skeletal muscle. However, the P3 does

appear slightly increased in the presence of fMyBP-C (Supplemental Figure 3). In slow

twitch soleus muscle, similar to cardiac muscle, the overall shape of the transient was not

dramatically changed following loss of sMyBP-C, but both the P_2 nadir decreased further

beyond the pre-stretch steady state, P_{df} amplitude increased, and K_{rel} rates were also

increased (Figure 3f,i,l). Taken together, loss of fMyBP-C conferred SA-like properties to

fast skeletal fibers, suggesting that canonical cross-bridge tension transient responses to

stretch are modulated by MyBP-C paralogs in the different fiber types.

Skeletal muscle MyBP-C paralogs alter tension transient properties in cardiac

muscle.

To directly test the extent to which differences in transient tension responses to stretch

are due to MyBP-C paralog expression in different muscle types, we next assessed

whether fMyBP-C and sMyBP-C could confer skeletal-like responses to stretch in

cardiomyocytes. We chose cardiac muscle as the background for paralog exchanges in

this set of experiments because we were able to remove nearly all endogenous cMyBP-

C following TEV protease treatment in cardiac myocytes compared with the more limited

removal of skeletal muscle paralogs from either psoas or soleus muscles that express a

mixture of skeletal MyBP-C paralogs (22). Additionally, because mouse hearts contain

primarily a single myosin isoform (α-myosin)(37), other potentially confounding effects due

to variable myosin isoform expression in different skeletal muscle fiber types are

avoided(38). Therefore, to directly assess effects of the different MyBP-C paralogs,

cardiomyocytes from $SpyC_3$ mice were first treated with TEVp to eliminate endogenous

cMyBP-C and then cMyBP-C was replaced *in situ* with either recombinant cMyBP-C,

fMyBP-C or sMyBP-C via SpyCatcher encoded at their C'-terminal ends. As shown in

Figure 5a, recombinant cMyBP-C was first added as control. Ligation of recombinant

cardiac C0C7Sc did not alter the overall shape of the tension transient compared to before

TEVp treatment, although the P_2 value tended to be higher. This could be due to the lack

of post-translational modifications (e.g., phosphorylation) of the recombinant C0C7Sc

compared to the endogenous cMyBP-C prior to TEVp treatment since phosphorylation of

cMyBP-C reduces P_2(39). Ligation of cardiac C0C7Sc also restored Ca^{2+}-sensitivity of

tension (pCa$_{50}$: Pre: 5.41 ± 0.01, Post: 5.32 ± 0.01, cC0C7Sc: 5.39 ± 0.02) and reduced

k_{tr} rates to values before treatment with TEVp (Figure 4a,d). Similar effects were observed

for ligation of fast C1C7Sc, where Ca^{2+}-sensitivity was restored (pCa$_{50}$: Pre:5.54 ± 0.02,

Post:5.44 ± 0.01, fC1C7Sc: 5.54 ± 0.01) (Fig. 4.b,e). However, after ligation of fC1C7Sc

transient responses to stretch in cardiac muscle were altered such that they more closely

resembled those typically seen in fast twitch skeletal fibers instead of those seen in

cardiomyocytes. That is, fC1C7Sc significantly elevated the P_2 amplitude above the pre-

stretch steady state force and nearly eliminated the rise in P_3 force (SA), leading to

negative P_{df} values (Figure 5e,h). Additionally, fC1C7Sc decreased the phase 2 relaxation

rate K_{rel}, trending to be slightly slower than endogenous cMyBP-C. (Fig. 5k). Taken

together, these experiments demonstrate that removal of fMyBP-C from psoas muscle altered the tension transient to resemble cardiac muscle, while addition of recombinant fC1C7Sc to cardiac muscle altered the tension transient to resemble fast twitch skeletal muscle.

Similar results were seen following replacement with sC1C7Sc, which was able to restore Ca^{2+}-sensitivity of tension (pCa_{50}: Pre 5.55 ± 0.02. Post: 5.49 ± 0.01, sC1C7Sc: 5.54 ± 0.02) and reduce k_{tr} rates to a similar extent as addition of cMyBP-C (Figure 4c,f). Therefore, both recombinant fMyBP-C and sMyBP-C are sufficient to restore typical interactions of endogenous cMyBP-C in cardiac muscle. sC1C7Sc also raised the P_2 nadir to a level above pre-stretch steady state and nearly eliminated P_{df} amplitude of the tension transient, thus significantly blunting the SA response typically seen in cardiac muscle (Fig. 5f,i). Taken together, these results demonstrate that skeletal muscle paralogs of MyBP-C reduce canonical SA responses in cardiac muscle and instead confer skeletal-like responses to rapid stretch.

Structural changes following loss of fMyBP-C determined by X-ray diffraction

The startling result that fMyBP-C cleavage in psoas leads to cardiac-like SA properties encouraged us to identify underlying sarcomere-level structural changes. We therefore used the small-angle X-ray diffraction technique on permeabilized psoas fibers from SnoopC$_2$ mice under Ca^{2+}-activated conditions to determine structural changes in the sarcomere before and after cleavage of fMyBP-C. In particular, we sought to determine whether fMyBP-C contributed to changes in lattice spacing, lattice order, and mass distributions on the thick or thin filaments that could potentially account for the change in transient force behaviors described above. X-ray reflections and their relationship to

sarcomere structures can be visualized in the cross-sectional diagram of the sarcomere

(Fig. 6a,b,c).

fMyBP-C stabilizes lattice order during isometric contraction, reduces myosin

head order, and shortens thin filament length during isometric contraction

Figure 6 shows a representative X-ray diffraction pattern collected from active $SnoopC_2$

psoas muscle. We report on features of equatorial reflections, including the spacing of the

1,0 reflection (d10), the thick-filament only lattice plane spacing, and its width in equatorial

space, which represents the degree of heterogeneity in the lattice and thick filament

disorder (σ_D)(40). Under activating conditions at pCa 6.0 (approximately the pCa_{50} for

force generation), we found that lattice spacing decreased with increasing SL as expected.

However, there were no significant differences in lattice spacing following removal of

fMyBP-C (Fig. 6d). This result differs from the increase in lattice spacing following loss of

fMyBP-C observed under relaxing conditions (23), but may be explained by the relatively

greater influence of strongly bound cross-bridges to act on the thin filament and limit lattice

expansion during Ca^{2+} activation. However, we found that the heterogeneity of lattice

spacing (σ_D) was significantly increased at both SL 2.4 and 3.0 following removal of

fMyBP-C (Fig. 6f). The latter suggests a modest degradation in the lattice integrity during

contraction when fMyBP-C is cleaved.

In order to further investigate the role of fMyBP-C in the orientation and order of myosin

heads as well as the thin filament, we investigated the $I_{1,1}/I_{1,0}$ equatorial intensity ratio and

M3, T3, M6 meridional reflections as well as the 6[th] actin layer line (ALL6) in the X-ray

diffraction patterns. The ratio of the 1,0 and 1,1 reflection intensities $(I_{1,1}/I_{1,0})$ reflects the

distribution of mass between thick and thin filaments, where increases are typically associated with a shift of myosin heads to the ON state and/or binding of myosin heads to the thin filament. We found that a loss of fMyBP-C did not have a significant effect on the $I_{1,1}/I_{1,0}$ intensity ratio during contraction, suggesting that loss of fMyBP-C did not significantly affect the distribution of myosin heads away from the thick filament backbone or proportion of bound crossbridges under Ca^{2+} activated conditions. The $I_{1,1}/I_{1,0}$ intensity ratios at both short and long SL (SL 2.4 and 3.0) also remained unchanged during isometric contraction, as previously reported by others (41, 42). Next, the spacing of the M3 reflection (S_{M3}), a measure of the axial periodicity of myosin heads, did not significantly change at either SL 2.4 or 3.0 following loss of fMyBP-C (Supplementary Figure x). Because S_{M3} is also taken as a measure of whether myosin heads are primarily in the OFF or ON state, results suggest that fMyBP-C does not contribute significantly to the OFF/ON distribution of myosin heads or numbers of bound myosin heads under active conditions, consistent with the $I_{1,1}/I_{1,0}$ intensity ratio. The intensity of the M3 reflection (I_{M3}), however, decreased following removal of fMyBP-C (Figure 7g), consistent with an increase in the angular dispersion of myosin heads(43, 44), suggesting that loss of fMyBP-C promotes disorder of myosin heads. The spacings of other thick filament reflections, such as (S_{M6}), indicative of thick filament length, were unchanged following removal of fMyBP-C (Supplementary Figure 4).

Finally, we sought to determine whether any thin filament structural changes occurred following fMyBP-C removal. We found that the spacing of the third-order troponin reflection (S_{T3}) was significantly decreased following removal of fMyBP-C at both SL 2.4 and SL 3.0, suggesting a reordering of the troponin complex on the thin filament. Additionally, the spacing of actin layer line 6 (S_{ALL6}), reflecting the pitch left-handed helix of actin, was

358   significantly decreased following removal of fMyBP-C at both SL 2.4 and SL 3.0 µm (Fig.

359   6i).

360
361   **Discussion**
362
Myosin binding protein-C paralogs commonly influence Ca^{2+}-sensitivity, cross-

bridge cycling, and oscillatory contractions in striated muscles

In this study, we sought to identify key similarities and differences in the role of the three

major MyBP-C paralogs in regulating muscle contractility, as well as identify key roles for

fMyBP-C in the structural organization of isometrically contracting sarcomeres. Using

three genetically modified mouse lines specific to slow (SpyC$_1$), fast (SnoopC$_2$), and

cardiac-MyBP-C (SpyC$_3$) that allow for selective removal of MyBP-C and replacement with

recombinant MyBP-C in the respective muscle types, we were able to test mechanical

properties of each muscle type in the presence and absence of MyBP-C. We found that

selective cleavage of MyBP-C led to a reduction in Ca^{2+}-sensitivity in both cardiac and fast

twitch myocytes. Effects on Ca^{2+}-sensitivity did not reach significance in soleus, a

predominantly slow twitch muscle, likely due to significant co-expression of fast and slow

MyBP-C (since fMyBP-C would not be cleaved in skeletal muscles from SpyC$_1$ mice).

However, recombinant slow MyBP-C was able to restore Ca^{2+}-sensitivity when added to

SpyC$_3$ cardiac myocytes (Figure 4f), so all 3 paralogs apparently share the common ability

to shift the tension-pCa relationship to the left. Similarly, cleavage of each MyBP-C paralog

increased cross-bridge cycling rates at submaximal Ca^{2+} activation as measured by k_{tr}.

This result is in good agreement with previously reported changes in both cardiac and fast

skeletal MyBP-C KO mice as well as previous work with the SnoopC$_2$ and SpyC$_3$ mouse

lines(22, 45, 46). Moreover, loss of each MyBP-C paralog led to the appearance of

spontaneous oscillatory contractions at sub-maximal Ca^{2+} activations near the pCa$_{50}$ for

force generation. While the precise mechanisms by which oscillatory contractions are

induced and propagated are still unclear, the phenomenon is an intrinsic, emergent

property of sarcomeres that has been likened to a third state of muscle activity between

activation and relaxation(29, 47). Notably, because force oscillations occur at Ca^{2+}

concentrations expected to be encountered in living muscle and are influenced by

metabolites such as ADP/P_i(48), tension instabilities are likely to have physiological

relevance. Consistent with this idea, mutations in slow MyBP-C lead to force oscillations

that are associated with skeletal muscle tremor in humans(17, 49, 50). Results presented

here therefore suggest that a novel functional role of all 3 MyBP-C paralogs is to limit

mechanical force instabilities and promote contractile synchrony. Conversely, loss of

MyBP-C or mutations in MyBP-C underlie contractile dysfunction in different muscle types

(51).

Myosin Binding Protein C paralogs differentially contribute to tension transients in

striated muscle

While we identified several common functional effects of the different MyBP-C

paralogs as discussed above, a notable departure from this trend was evident when

comparing the differential responses to a rapid 2% stretch in each muscle type which we

found were attributable to expression of the different MyBP-C paralogs. For instance,

under control conditions all 3 muscle types (cardiac, fast and slow skeletal) had unique

responses to rapid stretch that were mostly stereotypical for that muscle type as previously

described by others(19, 33, 34, 52, 53). This was most evident when comparing the

responses of cardiac and psoas (fast skeletal) muscles, where cardiac muscle exhibited

a canonical 3 phase response that included a final phase such that after reaching a nadir

(P_2), steady state force rises above the pre-stretch force, P_3. (SA, discussed in more detail

below). By contrast, P_2 was highly elevated in psoas muscle and the phase 3 rise in force

was typically minimal and/or short-lived. Notably, classical SA behavior is generally

considered minimal or absent in fast twitch muscle, although enhancement of SA in

skeletal muscles may be important under conditions of fatigue (e.g., high inorganic P_i) (34,

52, 54). Strikingly, as shown in Figure 3, we found that loss of fMyBP-C after TEVp

cleavage in psoas muscle completely altered the stereotypical response to stretch and

completely converted the response into one resembling that of cardiac muscle, including

a significant decrease in P_2 and the appearance of a prominent phase 3 SA (delayed force

increase) response. Conversely, when fMyBP-C was added to cardiac myocytes (Fig. 5),

the tension transient converted into a response more closely resembling one typically

found in fast skeletal muscle where P_2 was elevated and phase 3 was eliminated. Taken

together, our results provide compelling evidence that the different MyBP-C paralogs

contribute significantly to the stereotypical tension transient responses in their respective

muscle types. Furthermore, to our knowledge, this is the first report showing that loss of

fMyBP-C confers classic SA-like properties to fast twitch skeletal muscle fibers.

Tension transient responses to stretch, including SA responses, are thought to be

mediated largely by changes in cross-bridge cycling kinetics (30). Accordingly, numerous

factors that are known to impact cross-bridge dynamics, including calcium, ADP, and P_i

concentration, the extent of stretch, temperature, and the contribution of other sarcomere

proteins, such as thin filament proteins as well as myosin heavy chain isoforms, have all

have been shown to significantly impact the shape and rate of tension transient responses

to stretch(34, 54-56). In this respect, it may not be surprising that MyBP-C, known to affect

cross-bridge cycling kinetics, also influences tension transients. Indeed, consistent with

results reported here, Stelzer et al (2006) reported increased K_{rel}, reduced P_2, and

increased K_{df} rates in cMyBP-C KO mice, effects that were largely replicated by PKA

phosphorylation of WT cMyBP-C (33, 39). PKA phosphorylation of sMyBP-C in slow twitch

fibers also increased force development rates (57). Our results thus confirm and extend

previous findings by showing that all 3 MyBP-C paralogs limit cross-bridge cycling kinetics

in their respective muscle types. However, to further explore the extent to which each

paralog by itself was sufficient to determine the overall shape of tension transients when

expressed in a similar background, we also utilized the SpyC$_3$ cardiac mouse line that

primarily expresses a single myosin isoform (α-myosin) in cardiac myocytes to

systematically replace endogenous cMyBP-C with recombinant cardiac, slow , and fast

MyBP-C. Again, we found that each MyBP-C paralog conferred a characteristic tension

transient in cardiac muscle, thus allowing us to attribute observed changes in the tension

transients primarily to the different paralogs of MyBP-C rather than to different myosin

isoforms or other confounding variables.

MyBP-C paralogs determine the extent of strain-induced cross-bridge detachment,

P$_2$

A key difference that emerged when comparing the tension transients in the different

muscle types was the extent of the P$_2$ nadir, which presumably reflects the magnitude of

cross-bridge detachment in response to strain in each muscle type. Notably, P$_2$ was

reduced following loss of MyBP-C in all 3 muscle types, suggesting that a normal function

of MyBP-C is to limit strain-induced cross-bridge detachment. In cardiac muscle, strain-

dependent cross-bridge detachment is emerging as a key regulator of the mechanical

control of relaxation where stretch enhances LV relaxation during the transition from late

systole to early diastole(55, 58, 59) . In fact, Stehle et al. concluded that load-dependent

cross-bridge detachment is the only mechanism that explains how cardiac muscle can

relax as quickly as it contracts(60). Consistent with this, small stretches during the slow

phase of myofibril relaxation were shown to accelerate the onset of sarcomere "give"

which initiates the fast exponential phase of relaxation(61). We also recently showed that

loss of cMyBP-C significantly accelerates the fast phase of relaxation suggesting that

cMyBP-C provides an internal load (potentially by cross-linking thick and thin filaments) or

otherwise stabilizes sarcomere dynamics to slow relaxation(24).

In fast skeletal muscle acceleration of strain-dependent cross-bridge detachment after

loss of MyBP-C may contribute to the appearance of SA-like properties. For instance,

Wakefield et al. modeled cardiac tension transients in terms of a model where the shape

of the transient was the sum of 3 main factors: a passive Power-law relaxation term, a

Reversal term that is responsible for the transient drop in force (P_2 nadir), and a Drag term

that represents the initial fast rise in tension and relaxation immediately after the

stretch(59). If the Reversal term in skeletal muscle is small as suggested by Davis and

Epstein(32), and further limited by MyBP-C, then the shape of the tension transient in

skeletal muscle would be expected to be dominated by a combination of the Drag and

Power-law terms. However, if loss of fMyBP-C increases strain-dependent cross-bridge

detachment (g_1 in their model), then the Reversal term would become greater and deepen

the amplitude of the P_2 nadir(59). At the same time, loss of MyBP-C could further reduce

the Drag term as well as leading to an overall increase in the R/D ratio. Together, the two

effects would lead to the appearance of an SA response as observed following the loss of

fMyBP-C.

Through which interactions are the various MyBP-C isoforms able to affect the Drag and

Reversal terms? If MyBP-C links the thick and thin filaments and is able to slow filament

sliding(45, 57), then this interaction would be expected to increase the Drag term. MyBP-

C's presence may also limit the Reversal term, for instance, by preventing P_i-dependent

reversal of the powerstroke. Consistent with this cMyBP-C slows phosphate (P_i)-

dependent myosin cross-bridge detachment and prolongs cross-bridge lifetimes in cardiac

muscle (62, 63). Tanner et al. further concluded that cMyBP-C inhibits P_i-dependent

reversal of the powerstroke or otherwise stabilizes cross-bridge attachment in a way that

increases the probability of myosin completing its powerstroke(64). Notably, to account

for differences in relative rates of P_2 (P_{2-slow}) in different fiber types, Davis and Epstein

argued that P_i-dependent reversal of the powerstroke is largely absent in fast fibers,

whereas it is prominent in slow fibers(32). Straight et al. also concluded that a reverse

stroke of slow skeletal myosin accounts for augmentation of SA in slow skeletal muscle,

but not in fast skeletal muscle(34), although the same group later found that increases in

[P_i] concentration, such as during muscle fatigue, augments SA even in fast twitch

fibers(54). Using optical tweezers and synthetic cardiac myofilaments, Hwang et al.

demonstrated that a reversal stroke is present in cardiac myosin, but absent in fast skeletal

myosin, where it is a key determinant of cardiac contraction and relaxation and is further

necessary to produce stable force even at low levels of Ca^{2+} activation(65). While the

reversal stroke (or absence of) is therefore likely intrinsic to the myosin isoform, our results

suggest that MyBP-C paralogs are key regulators of cross-bridge detachment rates,

potentially via modulation of P_i-dependent reversal of the powerstroke(66).

SA is a phenomenon best described in some insect flight muscles that operate at high

frequencies (100-1000 Hz) where muscle is turned on and off in response to mechanical

signals rather than being driven by slower and more energetically costly chemical and

electrical signals(35, 67, 68). In these so-called asynchronous muscles, a single action

potential elicits a sustained rise in [Ca^{2+}] that is permissive for multiple cycles of muscle

activation and relaxation driven by muscle stretch. The initiating mechanical event, muscle

stretch, itself is the result of contraction of an antagonistic muscle such that asynchronous

muscles act in pairs. By accelerating strain-induced cross-bridge detachment, it is

plausible that stretch synchronizes relaxation of a population of cross-bridges that contract

together resulting in SA. In this respect it is notable that MyBP-C, which appears to limit

strain-induced cross-bridge detachment as we show here, is absent in insect flight

muscles, although other proteins such as projectin and flightin may perform analogous

roles(69-71). While mammalian cardiac and skeletal muscles utilize a synchronous

contractile mechanism (where a single action potential leads to a single twitch), SA occurs

to some extent in both muscle types and may play important physiological roles. Examples

include coordinating proper ejection timing of the ventricle and mechanical control of

relaxation in the heart (55, 58, 72), or during exercise where contraction of one muscle

may lead to stretch in its antagonistic partner, priming it for contraction moments later(52,

54). Our results suggest that by modulating strain-dependent cross-bridge detachment

MyBP-C fine-tunes the contribution of SA in different muscle types and that regulation of

MyBP-C, for instance by phosphorylation within a given muscle type, may thus contribute

to a greater or lesser dependence on mechanical signals for activation and relaxation(33,

57).

Potential mechanisms for MyBP-C regulation of the tension transient response

To gain additional insights into mechanisms by which loss of MyBP-C affects cross-bridge

kinetics, we investigated structural effects of loss of MyBP-C using X-ray diffraction

analyses of Ca^{2+}-activated psoas muscle before and after cleavage of MyBP-C with TEVp.

We chose psoas muscle for X-ray diffraction studies because the effects of loss of MyBP-

C were greatest in psoas muscle (e.g., the reduction in P_2 was greatest) and because we

had previously characterized loss of MyBP-C in psoas muscle under relaxed conditions(23). In that study we found that loss of MyBP-C expanded the sarcomere lattice, reduced the length of the thin (actin) filament, and repositioned myosin heads from their OFF to ON positions on the thick filament. These results were overall consistent with effects of MyBP-C to stabilize the inhibited state of myosin and limit lattice expansion potentially through interactions of MyBP-C with actin. By contrast, here we found under Ca^{2+}-activated conditions that loss of MyBP-C did not significantly affect lattice spacing possibly because strongly bound cross-bridges effectively limit the separation of the thick and thin filaments. However, an indicator of lattice disorder or heterogeneity, σ_D, was significantly increased at both SL 2.4 and SL 3.0 µm following loss of MyBP-C. Palmer et al., also reported greater myofilament disorder in a mouse model of cMyBP-C knockdown(63). σ_D is increased during active conditions compared to passive conditions, likely due to contributions of cross-bridge cycling that tend to move the positions of both thick and thin filament relative to their equilibrium position in the lattice(73). Therefore, results suggest that MyBP-C appears to plays a role in maintaining thick and thin filaments near their equilibrium positions during contraction, possibly by linking thick and thin filaments together. If so, then MyBP-C may promote efficient/strong cross-bridge binding by maintaining alignment of binding sites relative to one another on thick and thin filaments. Conversely, an increase in lattice disorder with loss of MyBP-C may lead to less efficient binding of myosin cross-bridges and/or promote cross-bridge detachment, thereby decreasing myosin duty ratio and increasing cycling rates(74, 75). The idea that MyBP-C links thick and thin filaments together is also consistent with MyBP-C acting as a drag force or internal load that slows cross-bridge cycling kinetics and relaxation(24, 45, 57).

MyBP-C also had significant effects on both the myosin head position and the thin filament.

The proportion of myosin heads in the ON state appeared unchanged following loss of

MyBP-C as indicated by the spacing of the M3 reflection (S_{M3})(Supplementary Fig. X3),

but the ordering of the myosin heads relative to the thick filament backbone was

significantly decreased at both SLs as indicated by a reduction in the M3 intensity (I_{M3})

(Fig. 6). Increased disorder of the myosin heads captured in the M3 intensity may be

indicative of less efficient cross-bridge binding to the thin filament and may explain the

faster cycling rates identified in mechanics experiments. Finally, loss of MyBP-C

decreased both the spacing of the third order troponin reflection (S_{T3}) and the spacing of

actin layer line 6 reflection (S_{ALL6}). The reduced spacing of these two actin-based

reflections indicate an overall reduction of thin filament length, consistent with the ability

of MyBP-C to bind to and exert a strain on the thin filament. In line with this, MyBP-C is

essential for thick filament stability and lattice rigidity(62, 63, 76). Notably, filament

compliance can affect stability of strongly bound cross-bridges and relaxation rates(77-

79). Taken together, the changes in X-ray diffraction reflections are consistent with a role

for MyBP-C in minimizing overall disorder of the sarcomere lattice during contraction and

maintaining myosin heads in ordered positions. Whereas proper alignment of thick and

thin filaments relative to one another may promote strong cross-bridge binding and cross-

bridge stability, loss of thick and thin filament alignment may promote cross-bridge

detachment, especially under strain.

Limitations

A limitation of our X-ray diffraction analysis was that structural changes were investigated

during isometric contraction, making it difficult to draw conclusions about whether dynamic

changes occur during rapid stretch that are not captured during steady state Ca^{2+}

activation. Future studies will include time-resolved X-ray diffraction measurements taken

as the muscle is being stretched in order to obtain structural information during the various

phases of the tension transient. Another limitation was that the soleus muscle contains a

mixed population of fast type and slow type fibers and we did not attempt to differentiate

their different contributions in mechanical force measurements. As a result, the

mechanical contribution of slow MyBP-C was likely underestimated due to the significant

presence of fast MyBP-C which was not subject to cleavage in $SpyC_1$ mice. In addition,

there are at least two splice variants of slow skeletal MyBP-C expressed in mouse and we

did not attempt to determine their relative expression(21).

Conclusions

Major conclusions from this study are that all 3 paralogs of MyBP-C (cardiac, fast skeletal

and slow skeletal) exert robust functional effects on force generation in their respective

muscle types. Using selective cleavage of each paralog we showed that all 3 had common

functional effects to increase Ca^{2+} sensitivity of tension and limit cross-bridge cycling rates

at submaximal levels of activation. In addition, all 3 were effective at damping force

oscillations that appeared following loss of MyBP-C when MyBP-C was cleaved with

TEVp. Finally, all 3 paralogs slowed strain-dependent cross-bridge detachment rates

which was evident as accelerated detachment rates (K_{rel}) and a lower force nadir (P_2)

following loss of MyBP-C when muscles were subjected to a rapid stretch. However, the

overall transient force response to a rapid stretch was unique in each muscle type and

dependent on the MyBP-C paralog expressed. In particular, MyBP-C paralogs uniquely

modulated strain dependence of cross-bridge detachment and thereby influenced SA

properties in each muscle type. This regulation of cross-bridge relaxation may have

significant physiological importance, as strain-induced cross-bridge shortening is key to

rapid relaxation at onset of diastole to allow for proper ventricular filling, and may also

enable rapid bursts in contraction in skeletal muscle.

603
604
605 **Materials and Methods**
606
Animal procedures:

All animal procedures were performed under protocols approved by the Institutional

Animal Care and Use Committee (IACUC) at the University of Arizona. $SpyC_1$, $SnoopC_2$,

and $SpyC_3$ (C57BL6/NJ) mice were created by the Genetically Engineered Mouse Models

(GEMM) core at the University of Arizona. All mice were bred and housed in a pathogen-

free barrier animal facility at the University of Arizona. Animal genotypes were confirmed

using PCR of DNA extracted from toe-clip samples and amplicons visualized on 1%

agarose gels.

Mice used in this study were euthanized via cervical dislocation following anesthesia with

isoflurane. Left ventricles, psoas, and soleus muscles were immediately dissected and

flash frozen in liquid nitrogen for long-term storage at -80°C for force measurements. For

X-ray diffraction measurements, muscles were permeabilized using glyercol (1:1

rigor:glycerol where Rigor solution contained (in mM): KCl (100), $MgCl_2$ (2),

ethyleneglycol-bis(β-aminoethyl)-N,N,N',N'-tetraacetic acid (EGTA, 5), Tris (10),

dithiothreitol (DTT, 1), 1:100 HALT™ Protease Inhibitor Cocktail (ThermoFisher Scientific,

Waltham, MA, USA)) at 4°C for 48 hours. After 48 hours the solution was refreshed and

then muscles were stored up to 60 days at -20°C until use.

Recombinant protein expression and purification:

Protein plasmids with desired DNA sequences were purchased from Genscript

(Piscataway, NJ, USA). BL21 and DE3 competent cells (C2527, New England Biolabs,

Ipswich, MA, USA) were transfected with plasmid DNA and protein purification was

performed as described previously.(13, 22)

Muscle force measurements:

Tissue was thawed in a skinning solution consisting of a Relax Buffer containing 1% Triton

X-100, 0.1% Saponin, and 1:100 HALT™ Protease Inhibitor Cocktail (ThermoFisher

Scientific, Waltham, MA, USA). Relax buffer contained (in mM): 100 KCl, 10 imidazole, 2

EGTA, 5 MgCl$_2$, 4 ATP (A2383, Sigma, St. Louis, MO, USA), and 1:100 HALT™ Protease

Inhibitor Cocktail at pH of 7.0. Multicellular preparations of cardiac cells and skeletal

muscle fibers were then isolated by mechanical disruption using a homogenizer (Omni

International 60CE54. Model: TH115). After homogenization, cardiac and skeletal muscle

cells were incubated with rotation in a skinning solution at 4°C for 15 and 30 min,

respectively. Detergent-permeabilized cardiac and skeletal muscle cells were then

washed thoroughly with fresh Relax buffer to remove residual detergents. Next, cardiac

cells and skeletal muscle fibers (~80-200 mm in length, 50-100 mm in width) were selected

and attached to a high-speed positional motor (Model: 315C-I, Aurora Scientific Inc.,

Aurora, Ontario, Canada) and a force transducer (Model 403 A series, Aurora Scientific

Inc., Aurora, Ontario, Canada) using aquarium sealant (Marineland, 100% clear silicone

rubber or Loctite, Clear Silicone Waterproof Multipurpose Adhesive Sealant). The

positional motor and force transducer were attached over a temperature-adjustable 8-well

platform (Model 803B, Aurora Scientific Inc., Aurora, Ontario, Canada) regulated by a

thermocouple (825A, Aurora Scientific Inc., Aurora, ON, Canada). The 8-well platform was

then mounted on the stage of an inverted light microscope (Model IX-53, Olympus

Instrument Co., Japan). The aquarium sealant was allowed to cure for 30 min before the

experiment was started. SL was then measured using video imaging in passive conditions

(at pCa 9.0) and SL was set to a final length of 2.3 μm and 2.4 μm for cardiac and skeletal

muscle, respectively. All muscle force measurements were obtained at 15°C using a range

of pCa solutions with free Ca^{2+} pCa 9.0 to 4.5, where pCa = -log[Ca^{2+}]. Force

measurements (F) recorded at each submaximal pCa solution were normalized to the

maximal force (F_0) at pCa 4.5 and to cross-sectional area of the muscle fiber assuming

circular dimensions. Data was fitted using a sigmoidal four-parameter logistic curve within

Graphpad Prism. k_{tr} was assessed by performing a slack-restretch test, where the muscle

was shortened to 80% of its reference length for 20 ms, stretched to 105% reference

length for 5 ms and returned to reference length. Tension transient parameters were

assessed by performing a rapid 2% stretch over 5 ms and held for 3 s. Cross-bridge

cycling rates (k_{tr} and shortening induced k_{df}) and cross-bridge relaxation (k_{rel}), were fit

using a single exponential curve and normalized to maximal rate (k_0) at pCa 4.5 and

plotted against isometric force.

N-terminal MyBP-C cleavage and *in situ* paste of recombinant proteins:

Mechanics experiments typically consisted of 3 consecutive sets of measurements: First,

before N-terminal cleavage of MyBP-C (control), second, after N-terminal cleavage of

MyBP-C with TEVp, and third in a subset of experiments after *in situ* paste of recombinant

MyBP-C. Proteolytic cleavage of domains C0-C7 in cardiac cells and C1-C7 in skeletal

muscle cells was performed by incubating myocytes in 20 μM AcTEV™ Protease

(ThermoFisher, 12575015) at 25°C for 30 min while the permeabilized muscle preparation

was attached to the high-speed positional motor and force transducer. After TEVp

incubation, cells were rigorously washed in fresh relax buffer (3 x for 3 min) to remove

TEV protease as well as cleaved endogenous MyBP-C N-terminal domains. Paste of

recombinant protein was done by incubating permeabilized cell preparations for 15 min at

15°C with 20 µM recombinant protein of choice containing a SpyCatcher sequence.

Myocytes were then washed (3 x for 3 min) using fresh relax buffer to remove excess

(non-ligated) recombinant protein. N-terminal MyBP-C cleavage and protein ligation was

assessed using SDS-PAGE gel electrophoresis and western blot as described

previously(23).

Small angle X-ray diffraction:

All X-ray diffraction experiments were performed with the small-angle instrument at the

BioCAT beamline 18ID at the Advanced Photon Source, Argonne National Laboratory.

Muscle experiments were performed both before and after the facility wide upgrade of

the Advanced Photon Source with changes to beam flux (Pre: ~3×10^{12}, Post: ~5×10^{13}

photons per second). The X-ray diffraction beam (0.103 nm wavelength), focusing at the

detector plane (Pre: ~0.06 x 0.15 mm. Post: 0.02 x 0.02 mm), sample to detector

distance (~2m), and detector (Mar 165, Rayonix Inc., USA) are identical to our previous

study in relaxed muscle fibers(23). An in-line camera was present to properly target

beam onto muscle fibers and assess for any irregularities in attachment of the muscle.

Muscle fibers were attached to custom fabricated clamps and connected to a high-speed

length controller (322C, Aurora Scientific, Canada) and a force transducer (402A, Aurora

Scientific, Canada) with a sampling rate of 1,000 Hz. Measurements from the force

transducer to identify activation levels of the fibers were visualized using Real Time

Muscle Data Acquisition and Analysis system (600A, Aurora Scientific, Canada).

Experiments were performed at ~25°C. SL was set at SL 2.4 µm or SL 3.0 µm using a 4-

mW Helium-Neon laser based on width of the first-order diffraction. Muscles were

activated by removal of relaxing solution (in mM: potassium propionate (45.3), N,N-

Bis(2-hydroxyethyl)-2-aminoethanesulfonic acid BES (40); EGTA (10), $MgCl_2$ (6.3), Na-

ATP (6.1), DTT (10), protease inhibitors [Complete], pH 7.0), three 30 second

incubations with pre-activating solution (in mM: potassium propionate (45.3), N,N-Bis(2-

hydroxyethyl)-2-aminoethanesulfonic acid BES (40); $MgCl_2$ (6.3), Na-ATP (6.1), DTT

(10), protease inhibitors [Complete], pH 7.0), and finally by incubation with pCa 6.0

solution (in mM: potassium propionate (170), magnesium acetate (2.5), MOPS (10) and

ATP (2.5), and CaEGTA and K_2EGTA were mixed at different proportions to obtain

different pCa ($-\log[Ca^{2+}]$), pH 7.0) values. Following incubation with pCa 6.0, tension

was monitored until the muscle reached a steady-state level of force, upon which the

bath was lowered with a servo-motor and X-ray images were taken in open air. No part

of the sample was exposed to the X-ray beam more than once. Samples then underwent

TEVp protease incubation as described above, and the process was repeated.

X-ray Diffraction Analysis:

Analysis of x-ray diffraction patterns was performed using MuscleX (version 1.22.0)

open-source data reduction and analysis package (BioCAT, Argonne National

Laboratories) as previously described(23, 80).

Statistics:

Statistical analyses were performed either in Graphpad Prism or JMP Pro (V16, SAS

Institute Inc., Cary, NC, USA). All measurements reported as the mean $\pm$ SEM.

Repeated measures one-way ANOVA tests with main effects pCa, TEVp treatment, and

their interaction, as well as an individual fiber random effect were performed on

measurements before TEV protease cleavage, after TEV protease cleavage, and when

present, after addition of recombinant protein. A mixed effects test was used for any

missing values. A paired t-test was used specifically to compare pCa50 values. The

significance level was $p < 0.05$. For additional statistical details see tables contained in

Supplemental Dataset 1.

728
729 **Acknowledgments**
730
This research was performed on APS beam time award(s) from the Advanced Photon
Source, a U.S. Department of Energy (DOE) Office of Science user facility operated for
the DOE Office of Science by Argonne National Laboratory under Contract No. DE-
AC02-06CH11357. BioCAT is supported by grant P30 GM138395 from the National
Institute of General Medical Sciences of the National Institutes of Health. The content is
solely the responsibility of the authors and does not necessarily reflect the official views
of the National Institute of General Medical Sciences or the National Institutes of Health.
We would like to thank Dr. Thomas C Irving for expertise and help with editing the
manuscript and John Sullivan for generating a video of SPOC in soleus muscle. Funding
for this study was provided by the National Institute of Health (HL080367 [SPH]),
HL140925 [SPH], AR081935 [SPH], T32 HL007249 [NME]), the American Heart
Association (827628 [NME]), and German Research Foundation (DFG) 454867250 to
ALH.
744

745 **References**
746
747

1.   L. Heling, M. A. Geeves, N. M. Kad, MyBP-C: one protein to govern them all. *J Muscle Res*
Cell Motil **41**, 91–101 (2020).
2.   D. Tamborrini *et al.*, Structure of the native myosin filament in the relaxed cardiac
sarcomere. *Nature* **623**, 863–871 (2023).
3.   D. Dutta, V. Nguyen, K. S. Campbell, R. Padron, R. Craig, Cryo-EM structure of the human
cardiac myosin filament. *Nature* **623**, 853–862 (2023).
4.   M. V. Razumova, K. L. Bezold, A. Y. Tu, M. Regnier, S. P. Harris, Contribution of the
myosin binding protein C motif to functional effects in permeabilized rat trabeculae. *J*
Gen Physiol **132**, 575–585 (2008).

5.   A. E. Whitten, C. M. Jeffries, S. P. Harris, J. Trewhella, Cardiac myosin-binding protein C
decorates F-actin: implications for cardiac function. *Proc Natl Acad Sci U S A* **105**, 18360–
18365 (2008).

6.   G. Kunst *et al.*, Myosin binding protein C, a phosphorylation-dependent force regulator
in muscle that controls the attachment of myosin heads by its interaction with myosin
S2. *Circ Res* **86**, 51–58 (2000).

7.   A. Ababou, M. Gautel, M. Pfuhl, Dissecting the N-terminal myosin binding site of human
cardiac myosin-binding protein C. Structure and myosin binding of domain C2. *J Biol*
Chem **282**, 9204–9215 (2007).

8.   C. Risi *et al.*, N-Terminal Domains of Cardiac Myosin Binding Protein C Cooperatively
Activate the Thin Filament. *Structure* **26**, 1604–1611 e1604 (2018).

9.   J. Y. Mun *et al.*, Myosin-binding protein C displaces tropomyosin to activate cardiac thin
filaments and governs their speed by an independent mechanism. *Proc Natl Acad Sci U S*
A **111**, 2170–2175 (2014).

10.   S. Nelson, S. Beck-Previs, S. Sadayappan, C. Tong, D. M. Warshaw, Myosin-binding
protein C stabilizes, but is not the sole determinant of SRX myosin in cardiac muscle. *J*
Gen Physiol **155** (2023).

11.   J. W. McNamara *et al.*, Ablation of cardiac myosin binding protein-C disrupts the super-
relaxed state of myosin in murine cardiomyocytes. *J Mol Cell Cardiol* **94**, 65–71 (2016).

12.   C. Y. Doh *et al.*, Molecular characterization of linker and loop-mediated structural
modulation and hinge motion in the C4-C5 domains of cMyBPC. *J Struct Biol* **214**,
107856 (2022).

13.   A. C. Greenman, R. L. Sadler, S. P. Harris, Autoinhibition of cMyBP-C by its middle
domains. *J Mol Cell Cardiol* **200**, 82–92 (2025).

14.   A. A. Glazier, A. Thompson, S. M. Day, Allelic imbalance and haploinsufficiency in
MYBPC3-linked hypertrophic cardiomyopathy. *Pflugers Arch* **471**, 781–793 (2019).

15.   S. P. Harris, R. G. Lyons, K. L. Bezold, In the thick of it: HCM-causing mutations in myosin
binding proteins of the thick filament. *Circ Res* **108**, 751–764 (2011).

16.   A. S. Helms *et al.*, Spatial and Functional Distribution of MYBPC3 Pathogenic Variants
and Clinical Outcomes in Patients With Hypertrophic Cardiomyopathy. *Circ Genom*
Precis Med **13**, 396–405 (2020).

17.   C. A. Gurnett *et al.*, Myosin binding protein C1: a novel gene for autosomal dominant
distal arthrogryposis type 1. *Hum Mol Genet* **19**, 1165–1173 (2010).

18.   J. W. McNamara, S. Sadayappan, Skeletal myosin binding protein-C: An increasingly
important regulator of striated muscle physiology. *Arch Biochem Biophys* **660**, 121–128
(2018).

19.   J. M. Mariano *et al.*, Pulsatile Myofilament Activity in Myotrem Myopathy Associated
with Myogenic Tremor. *Int J Mol Sci* **26** (2025).

20.   C. E. Oakley, B. D. Hambly, P. M. Curmi, L. J. Brown, Myosin binding protein C: structural
abnormalities in familial hypertrophic cardiomyopathy. *Cell Res* **14**, 95–110 (2004).

21.   M. A. Ackermann, A. Kontrogianni-Konstantopoulos, Myosin binding protein-C slow: a
multifaceted family of proteins with a complex expression profile in fast and slow twitch
skeletal muscles. *Front Physiol* **4**, 391 (2013).

22.   N. C. Napierski *et al.*, A Novel "Cut and Paste" Method for In Situ Replacement of
cMyBP-C Reveals a New Role for cMyBP-C in the Regulation of Contractile Oscillations.
Circ Res **126**, 737–749 (2020).

23. A. L. Hessel *et al.*, Myosin-binding protein C regulates the sarcomere lattice and
stabilizes the OFF states of myosin heads. *Nat Commun* **15**, 2628 (2024).
24. A. V. Dvornikov, S. P. Harris, Myosin-binding protein C slows cardiac myofibril relaxation
kinetics. *J Physiol* **603**, 5351–5368 (2025).
25. B. Zakeri *et al.*, Peptide tag forming a rapid covalent bond to a protein, through
engineering a bacterial adhesin. *Proc Natl Acad Sci U S A* **109**, E690–697 (2012).
26. G. Veggiani *et al.*, Programmable polyproteams built using twin peptide superglues. *Proc*
Natl Acad Sci U S A **113**, 1202–1207 (2016).
27. B. Brenner, E. Eisenberg, Rate of force generation in muscle: correlation with
actomyosin ATPase activity in solution. *Proc Natl Acad Sci U S A* **83**, 3542–3546 (1986).
28. G. K. Dhoot, M. C. Hales, B. M. Grail, S. V. Perry, The isoforms of C protein and their
distribution in mammalian skeletal muscle. *J Muscle Res Cell Motil* **6**, 487–505 (1985).
29. J. E. Wolfe *et al.*, SPontaneous Oscillatory Contraction (SPOC): auto-oscillations observed
in striated muscle at partial activation. *Biophys Rev* **3**, 53–62 (2011).
30. A. F. Huxley, R. M. Simmons, Proposed mechanism of force generation in striated
muscle. *Nature* **233**, 533–538 (1971).
31. L. E. Ford, A. F. Huxley, R. M. Simmons, Tension responses to sudden length change in
stimulated frog muscle fibres near slack length. *J Physiol* **269**, 441–515 (1977).
32. J. S. Davis, N. D. Epstein, Kinetic effects of fiber type on the two subcomponents of the
Huxley-Simmons phase 2 in muscle. *Biophys J* **85**, 390–401 (2003).
33. J. E. Stelzer, R. L. Moss, Contributions of stretch activation to length-dependent
contraction in murine myocardium. *J Gen Physiol* **128**, 461–471 (2006).
34. C. R. Straight, K. M. Bell, J. N. Slosberg, M. S. Miller, D. M. Swank, A myosin-based
mechanism for stretch activation and its possible role revealed by varying phosphate
concentration in fast and slow mouse skeletal muscle fibers. *Am J Physiol Cell Physiol*
317, C1143–C1152 (2019).
35. M. Linari, M. K. Reedy, M. C. Reedy, V. Lombardi, G. Piazzesi, Ca-activation and stretch-
activation in insect flight muscle. *Biophys J* **87**, 1101–1111 (2004).
36. J. O. Vigoreaux, J. Vigoreaux, SpringerLink (2006) Nature's versatile engine : insect flight
muscle inside and out. in *Molecular Biology Intelligence Unit* (Landes
Bioscience/Eurekah.com

Springer Science+Business Media

Springer US, Georgetown, Tex

New York, N.Y

New York, NY).
37. R. Rice *et al.*, Cardiac myosin heavy chain isoform exchange alters the phenotype of
cTnT-related cardiomyopathies in mouse hearts. *J Mol Cell Cardiol* **48**, 979–988 (2010).
38. B. C. Tanner, Y. Wang, J. Robbins, B. M. Palmer, Kinetics of cardiac myosin isoforms in
mouse myocardium are affected differently by presence of myosin binding protein-C. *J*
Muscle Res Cell Motil **35**, 267–278 (2014).
39. R. Mamidi, K. S. Gresham, S. Verma, J. E. Stelzer, Cardiac Myosin Binding Protein-C
Phosphorylation Modulates Myofilament Length-Dependent Activation. *Front Physiol* **7**,
38 (2016).

40. W. Ma, T. C. Irving, Small Angle X-ray Diffraction as a Tool for Structural Characterization
of Muscle Disease. *Int J Mol Sci* **23** (2022).
41. A. L. Hessel *et al.*, The distinctive mechanical and structural signatures of residual force
enhancement in myofibers. *Proc Natl Acad Sci U S A* **121**, e2413883121 (2024).
42. V. Joumaa *et al.*, Effect of Active Lengthening and Shortening on Small-Angle X-ray
Reflections in Skinned Skeletal Muscle Fibres. *Int J Mol Sci* **22** (2021).
43. G. Piazzesi *et al.*, Changes in conformation of myosin heads during the development of
isometric contraction and rapid shortening in single frog muscle fibres. *J Physiol* **514 (Pt**
2), 305–312 (1999).
44. H. E. Huxley, W. Brown, The low-angle x-ray diagram of vertebrate striated muscle and
its behaviour during contraction and rigor. *J Mol Biol* **30**, 383–434 (1967).
45. F. S. Korte, K. S. McDonald, S. P. Harris, R. L. Moss, Loaded shortening, power output,
and rate of force redevelopment are increased with knockout of cardiac myosin binding
protein-C. *Circ Res* **93**, 752–758 (2003).
46. T. Song *et al.*, Fast skeletal myosin-binding protein-C regulates fast skeletal muscle
contraction. *Proc Natl Acad Sci U S A* **118** (2021).
47. K. Sato, M. Ohtaki, Y. Shimamoto, S. Ishiwata, A theory on auto-oscillation and
contraction in striated muscle. *Prog Biophys Mol Biol* **105**, 199–207 (2011).
48. T. Kagemoto, A. Li, C. Dos Remedios, S. Ishiwata, Spontaneous oscillatory contraction
(SPOC) in cardiomyocytes. *Biophys Rev* **7**, 15–24 (2015).
49. X. Li *et al.*, Two novel mutations in myosin binding protein C slow causing distal
arthrogryposis type 2 in two large Han Chinese families may suggest important
functional role of immunoglobulin domain C2. *PLoS One* **10**, e0117158 (2015).
50. K. Ha *et al.*, MYBPC1 mutations impair skeletal muscle function in zebrafish models of
arthrogryposis. *Hum Mol Genet* **22**, 4967–4977 (2013).
51. S. P. Harris, Making waves: A proposed new role for myosin-binding protein C in
regulating oscillatory contractions in vertebrate striated muscle. *J Gen Physiol* **153**
(2021).
52. J. C. Robinett, L. M. Hanft, B. Biesiadecki, K. S. McDonald, Molecular regulation of
stretch activation. *Am J Physiol Cell Physiol* **323**, C1728–C1739 (2022).
53. S. Galler, O. Andruchov, G. M. Stephenson, D. G. Stephenson, Qualitatively different
cross-bridge attachments in fast and slow muscle fiber types. *Biochem Biophys Res*
Commun **385**, 44–48 (2009).
54. P. C. Woods, D. M. Swank, M. S. Miller, Stretch activation combats force loss from
fatigue in fast-contracting mouse skeletal muscle fibers. *J Gen Physiol* **157** (2025).
55. B. M. Palmer *et al.*, Enhancing diastolic function by strain-dependent detachment of
cardiac myosin crossbridges. *J Gen Physiol* **152** (2020).
56. R. H. Abbott, G. J. Steiger, Temperature and amplitude dependence of tension transients
in glycerinated skeletal and insect fibrillar muscle. *J Physiol* **266**, 13–42 (1977).
57. J. C. Robinett, L. M. Hanft, J. Geist, A. Kontrogianni-Konstantopoulos, K. S. McDonald,
Regulation of myofilament force and loaded shortening by skeletal myosin binding
protein C. *J Gen Physiol* **151**, 645–659 (2019).
58. C. S. Chung, C. W. Hoopes, K. S. Campbell, Myocardial relaxation is accelerated by fast
stretch, not reduced afterload. *J Mol Cell Cardiol* **103**, 65–73 (2017).
59. J. I. Wakefield, S. P. Bell, B. M. Palmer, Inorganic phosphate accelerates cardiac
myofilament relaxation in response to lengthening. *Front Physiol* **13**, 980662 (2022).

60.   R. Stehle *et al.*, Mechanical properties of sarcomeres during cardiac myofibrillar
relaxation: stretch-induced cross-bridge detachment contributes to early diastolic filling.
J Muscle Res Cell Motil **27**, 423–434 (2006).
61.   C. Poggesi, C. Tesi, R. Stehle, Sarcomeric determinants of striated muscle relaxation
kinetics. *Pflugers Arch* **449**, 505–517 (2005).
62.   B. M. Palmer *et al.*, Effect of cardiac myosin binding protein-C on mechanoenergetics in
mouse myocardium. *Circ Res* **94**, 1615–1622 (2004).
63.   B. M. Palmer *et al.*, Roles for cardiac MyBP-C in maintaining myofilament lattice rigidity
and prolonging myosin cross-bridge lifetime. *Biophys J* **101**, 1661–1669 (2011).
64.   B. C. W. Tanner, M. J. Previs, Y. Wang, J. Robbins, B. M. Palmer, Cardiac myosin binding
protein-C phosphorylation accelerates beta-cardiac myosin detachment rate in mouse
myocardium. *Am J Physiol Heart Circ Physiol* **320**, H1822–H1835 (2021).
65.   Y. Hwang, T. Washio, T. Hisada, H. Higuchi, M. Kaya, A reverse stroke characterizes the
force generation of cardiac myofilaments, leading to an understanding of heart function.
Proc Natl Acad Sci U S A **118** (2021).
66.   M. S. Woody, D. A. Winkelmann, M. Capitanio, E. M. Ostap, Y. E. Goldman, Single
molecule mechanics resolves the earliest events in force generation by cardiac myosin.
Elife **8** (2019).
67.   J. W. Pringle, The excitation and contraction of the flight muscles of insects. *J Physiol*
108, 226–232 (1949).
68.   G. J. Steiger, J. C. Ruegg, Energetics and "efficiency" in the isolated contractile machinery
of an insect fibrillar muscle at various frequencies of oscillation. *Pflugers Arch* **307**, 1–21
(1969).
69.   T. P. Vu Manh *et al.*, Expression of cardiac myosin-binding protein-C (cMyBP-C) in
Drosophila as a model for the study of human cardiomyopathies. *Hum Mol Genet* **14**, 7–
17 (2005).
70.   M. C. Reedy, B. Bullard, J. O. Vigoreaux, Flightin is essential for thick filament assembly
and sarcomere stability in Drosophila flight muscles. *J Cell Biol* **151**, 1483–1500 (2000).
71.   B. Bullard, A. Pastore, Through thick and thin: dual regulation of insect flight muscle and
cardiac muscle compared. *J Muscle Res Cell Motil* **40**, 99–110 (2019).
72.   K. B. Campbell, M. Chandra, Functions of stretch activation in heart muscle. *J Gen*
Physiol **127**, 89–94 (2006).
73.   L. C. Yu, A. C. Steven, G. R. Naylor, R. C. Gamble, R. J. Podolsky, Distribution of mass in
relaxed frog skeletal muscle and its redistribution upon activation. *Biophys J* **47**, 311–
321 (1985).
74.   T. L. Daniel, A. C. Trimble, P. B. Chase, Compliant realignment of binding sites in muscle:
transient behavior and mechanical tuning. *Biophys J* **74**, 1611–1621 (1998).
75.   I. A. Telley, J. Denoth, E. Stussi, G. Pfitzer, R. Stehle, Half-sarcomere dynamics in
myofibrils during activation and relaxation studied by tracking fluorescent markers.
Biophys J **90**, 514–530 (2006).
76.   L. R. Nyland *et al.*, Cardiac myosin binding protein-C is essential for thick-filament
stability and flexural rigidity. *Biophys J* **96**, 3273–3280 (2009).
77.   K. S. Campbell, Filament compliance effects can explain tension overshoots during force
development. *Biophys J* **91**, 4102–4109 (2006).
78.   B. Kiss *et al.*, Nebulin stiffens the thin filament and augments cross-bridge interaction in
skeletal muscle. *Proc Natl Acad Sci U S A* **115**, 10369–10374 (2018).

79.   K. S. Campbell, Compliance Accelerates Relaxation in Muscle by Allowing Myosin Heads
939   to Move Relative to Actin. *Biophys J* **110**, 661–668 (2016).
940   80.   J. Jiratrakanvong *et al.*, MuscleX: data analysis software for fiber diffraction patterns
941   from muscle. *J Synchrotron Radiat* **31**, 1401–1408 (2024).

954 **Figures and Tables**
955

Figure 1: Sarcomere organization, MyBP-C structure, and utilization of the "cut-and-paste" mouse models to investigate muscle response to rapid stretch. A) Illustration of a muscle sarcomere highlighting Myosin Binding Protein-C location in the C-zone. B) Schematic of slow, fast skeletal, and cardiac Myosin Binding Protein-C paralogs. Slow and cardiac isoforms contain variable phosphorylation sites within the proline/alanine-rich as well as M-domain. Both fast and cardiac contain a linker between domains C4C5 while slow does not. Cardiac MyBP-C also contains an additional N-terminal immunoglobulin domain termed C0 as well as a 28 amino acid loop within the C5 domain. C) Schematic representation of the cut-and-paste system in fast skeletal Myosin Binding Protein-C. D) Representative force transient traces from cardiac and skeletal muscle highlighting shape differences and defining stretch activation terms for different muscle type.

Figure 2: Removal of MyBP-C alters Ca^{2+}-sensitivity, cross-bridge cycling rates, and produces
oscillatory contractions in each muscle type. A-C) Tension-pCa curves for cardiac (A), fast
skeletal (B), and slow skeletal (C) before (blue) and after (red) removal of MyBP-C. D-F) Changes
in cross-bridge cycling rates in cardiac (D), fast (E), and slow skeletal (F) before and after
removal of MyBP-C. G-I) Representative force traces highlighting force overshoots and tension
oscillations at steady-state force in cardiac (G), fast (H), and slow skeletal (I) before and after
removal of MyBP-C. Force traces only in Figure 3i (sMyBP-C) were smoothed for clearer
visualization of oscillations on Graphpad Prism. Data presented as mean ± s.e.m. pCa_{50} values
assessed with paired t-test. **** = $p < 0.0001$ in (a-c). k_{tr} assessed with a two-way ANOVA with
pCa and treatment as main effects also assessing their interaction. Individual fiber was included
as a random effect. Treatment showed significance in (D-F) (D: Treatment: Prob > F = 0.0254, E:
Treatment*pCa Prob > F = < 0.0033, F: Treatment Prob > F = 0.0038) A Student's t-test or Tukey
Honestly Statistically Different (HSD) post-hoc test was performed on significant main effects (p
< 0.05). Dataset from: Cardiac (n=13 cardiomyocytes. N=4 mice). Psoas (n=10 fibers. N=3 mice).
Soleus (n=8. N=3 +sMyBP-C. N=4 -sMyBP-C). Figure 2B, 2E, and 2H are taken from Hessel et, al.
2024 publication in Nature Communications (23).

993
994

995
996
Figure 3: MyBP-C paralogs differentially affect tension transients following rapid stretch in each
muscle type **a-c:** Representative force traces following 2% rapid stretch before (blue) and after
(red)TEVp treatment of cardiac (a) fast skeletal psoas (b), and slow skeletal soleus (c). **d-f:**
Mechanical measurements of the P_2 nadir before and after TEVp treatment in cardiac
(d)(pCa*Treatment Prob > F = <0.0001, fast skeletal psoas (e) (Treatment Prob > F = <0.0001,
and slow skeletal soleus (f) (n.s). **g-i:** Mechanical measurements of P_{df}, the amplitude of delayed
force redevelopment, before and after TEVp treatment of cardiac (g)(Treatment Prob > F =
0.0001), fast skeletal psoas (h)(Treatment*pCa Prob > F = 0.0118), and slow skeletal soleus
(i)(Treatment Prob > F = 0.006). **j-l:** Mechanical measurements of the rate of cross-bridge

1006 relaxation (k_{rel}) following rapid 2% stretch before and after TEVp treatment of cardiac (j)(Prob >
1007 F = <0.0001), fast skeletal psoas (k)(Treatment: Prob > F = 0.0272), and slow skeletal soleus
1008 (l)(n.s). Data presented as mean ± s.e.m. A repeated measures ANOVA design was performed
1009 throughout with pCa, treatment, and their interaction as main effects. A random effect was
1010 included for each individual fiber. A Student's t-test or Tukey's HSD post-hoc test was performed
1011 on significant main effects (p < 0.05). Dataset: Cardiac from (n=13 cardiomyocytes. N=4 mice).
1012 Psoas (n=10 fibers. N=3 mice). Soleus (n=8. N=3 +sMyBP-C. N=4 -sMyBP-C). Further detail in
1013 supplement.

1014

1015

1016

Figure 4: Cut-and-paste experiments performed in detergent permeabilized cardiac muscle to
assess Ca^{2+}-sensitivity and k_{tr} after *in situ* replacement of recombinant cardiac, fast skeletal, and
slow skeletal MyBP-C. **a-c:** Relative tension-pCa with inset containing average pCa$_{50}$ following
each curve for cC0C7Sc(a), fC1C7Sc (b), and sC1C7Sc (c). ** = p = <0.0001, *** p < 0.0005, ****
p = <0.005 **d-f:** Changes in k_{tr} before and after removal of endogenous cMyBP-C and following
paste of cC0C7Sc (d) (pCa*Treatment Prob > F = 0.0046), fC1C7Sc (e) (Treatment Prob > F =
0.0061) and sC1C7Sc (f) (Treatment Prob > F = 0.0046). Statistics throughout are an ANOVA
design with pCa, treatment, and their interaction as main effects, as well as an individual fiber
random effect. Statistically significant differences were assessed using a Tukey's HSD post-hoc
test on statistically significant differences (p < 0.05). Dataset: cC0C7Sc (n=7, N=3), fC1C7Sc (n=6,
N=5), sC1C7Sc (n=8, N=6).

1028

1029
1030

1031

1032

1033

1034

Figure 5: Amplitudes of strain-induced cross-bridge detachment (P_2), stretch activation (P_{df}), and cross-bridge relaxation kinetics (k_{rel}) in cardiac muscle following *in situ* replacement with cC0C7Sc, fC1C7Sc, and sC1C7Sc. **a-c:** Representative force traces of cardiac tension transients before (blue) and after (red) removal of endogenous cMyBP-C, and following replacement (black) with recombinant cC0C7Sc (a) fC1C7Sc (b), and sC1C7Sc (c). **d-f:** Amplitude of strain induced cross-bridge detachment (P_2) before and after removal of endogenous cMyBP-C and after replacement with cC0C7Sc (d) (pCa*Treatment*pCa Prob > F = <0.0001), fC1C7Sc (e) (Treatment Prob > F = <0.0001), and sC1C7Sc (f))(Treatment*pCa Prob > F = 0.0234) **g-i:** Amplitude of stretch activation (P_{df}) before and after removal of endogenous cMyBP-C and after *in situ* replacement with cC0C7Sc (g)(Treatment*pCa Prob > F = 0.0029), fC1C7Sc (h)(Treatment*pCa Prob > F = 0.0076), and sC1C7Sc (i)(Treatment*pCa Prob>F = 0.0181. **j-l:** Rates of stretch induced cross-bridge relaxation (k_{rel}) before and after removal of endogenous

cMyBP-C and after *in situ* replacement with cC0C7Sc (j)(Treatment Prob > F = <0.0001), fC1C7Sc

(k)(Treatment Prob > F = <0.0001), and sC1C7Sc (l)(Treatment*pCa Prob > F = 0.0076). Statistics

throughout are an ANOVA design with pCa, treatment, and their interaction as main effects,

including an individual fiber random effect. A post-hoc Tukey's HSD test was performed on

statistically different main effects (p < 0.05). Dataset: cC0C7Sc (n=7, N=3), fC1C7Sc (n=6, N=5),

sC1C7Sc (n=8, N=6).

1053

 Figure 6: Equatorial and meridional reflections of isometrically contracting psoas Muscle in
presence and absence of fMyBP-C. **a)** Representative X-ray diffraction image at SL 3.0 and pCa
6.0 highlighting meridional reflections (M3, T3, M6, A6) assessed in this study. **b)** Graphic
showing a cross-sectional view of the sarcomere lattice as well as important equatorial
reflections. **c) Left:** Schematic of the quasi-helical arrangement of myosin heads on the thick
filament backbone, where the M3 reflection is highlighted as difference in space between
individual myosin crowns. **Right:** Schematic of the thin filament, highlighting both the left-
handed and right-handed helix of the actin monomers and their spacings. **d)** Lattice spacing of
permeabilized psoas muscle at SL 2.4 and SL 3.0 before and after removal of fMyBP-C
(Treatment: Prob > F 0.1494) **e)** I_{11}/I_{10} intensity ratio of permeabilized psoas muscle at SL 2.4 and

30 before and after removal of fMyBP-C. (Treatment: Prob > F = 0.2284) **f)** Heterogeneity of
lattice spacing (σ_D) of permeabilized psoas muscle at SL 2.4 and 3.0 before and after removal of
fMyBP-C. (Treatment: Prob > F = 0.003) **g)** Intensity of the M3 reflection (I_{M3}) of permeabilized
psoas muscle at SL 2.4 and 3.0 before and after removal of fMyBP-C. (Treatment: Prob > F =
0.0001) **h)** T3 spacing (S_{T3}) of permeabilized psoas muscle at pCa 6.0 at SL 2.4 and 3.0 before and
after removal of fMyBP-C. (Treatment: Prob > F = 0.0377) **i)** Actin layer line 6 spacing (S_{ALL6}) of
permeabilized psoas muscle at SL 2.4 and 3.0 before and after removal of fMyBP-C. (Treatment:
Prob > F = 0.0026) Statistics throughout are a repeated measure ANOVA design with SL,
treatment, and their interaction as main effects, including a random effect on individual fiber. A
Student's t-test or Tukey's HSD Post-Hoc test was performed on statistically significant
differences and presented in figures as connecting letters. Dataset: (n=43 fibers, N=13 animals).