Age-associated B cells (ABCs) develop from a CNS-localized progenitor pool into a pro-inflammatory phenotype after stroke

Annabel M. McAtee[1]*, Mathew Kenwood[2]*, Thomas Ujas[1], Mary K. Colson[1], Julie Watkins[1], Edric D. Winford[3], Katherine M. Cotter[1], Daimen Britsch[1], Dene Betz[2], Jadwiga Turchan-Cholewo[4], Jenny Lutshumba[4], Connor Stuart[1], Gargi Shah[1], Loganann Runice[1], Mark Ebbert[1], Salvatore Cherra[1], Peter T. Nelson[5], Jamie L. Sturgill[6], Nancy L. Monson[7,8,9], Mark P. Goldberg[2], Ann M. Stowe[1,4]

[1]Department of Neuroscience, University of Kentucky, 741 S. Limestone, BBSRB 479, Lexington, Kentucky 40508, USA
[2]Department of Neurology, Long School of Medicine, University of Texas at San Antonio, 7703 Floyd Curl Drive, MSC 7883, San Antonio, Texas 78229, USA
[3]Department of Neurology, Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Gertrude H. Sergievsky Center, Vagelos College of Physicians and Surgeons, Columbia University, New York, New York 10032 USA
[4]Department of Neurology, University of Kentucky, 740 S. Limestone, Kentucky Clinic J-455 Lexington, Kentucky 40536, USA
[5]Sanders-Brown Center on Aging and Department of Pathology, University of Kentucky, 800 S. Limestone, Rm 111B, Lexington, Kentucky 40536, USA
[6]Department of Microbiology, Immunology, and Molecular Genetics, University of Kentucky, 760 Press Ave., HKRB 532, Lexington, Kentucky 40506, USA
[7]Department of Neurology, University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Suite NL9.114, Dallas, Texas 75390-8813, USA
[8]Department of Immunology, University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Suite NL9.114, Dallas, Texas 75390-8813, USA
[9]O'Donnell Brain Institute, University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Suite NL9.114, Dallas, Texas 75390-8813, USA
*equal contribution

Corresponding author:
Ann M. Stowe, PhD, FAHA
Department of Neurology, BBSRB 479
741 S. Limestone St., Lexington, KY USA 40508
Phone: 001-859-323-8420
e-mail: ann.stowe@uky.edu
ORCID iD: 0000-0001-8111-4429

Preprint Server: BioRxiv

Key words: IgM, ischemic stroke, autoreactivity, B cell receptor, progenitor

The authors have no competing interests to declare.

Abstract

Aging and age-related diseases like ischemic stroke induce chronic lymphocyte recruitment into the central nervous system (CNS). Conflicting effects on post-stroke functional recovery, however, are secondary to the differences in responding lymphocyte populations that shift immunophenotype with both ischemic injury and age. To better define CNS-localized B cell subsets, we used flow cytometry, single-cell RNA sequencing, and B cell receptor sequencing on B cells isolated from uninjured and post-stroke brains of aged male and female mice. We identified a novel B1b cell progenitor pool distinct from canonical pleural and peritoneal B1 niches. Trajectory analysis showed B1b progenitors transition into age-associated B cell (ABC) subsets, and clonal expansion of IgM+ ABCs (ABC/B1b) and plasma cells following ischemic stroke. We also confirmed analogous ABCs and developing B cell populations in post-mortem human parenchymal tissue isolated from aged brain donors. These studies reveal unique B cell populations that proliferate within the aging CNS and are associated with impaired post-stroke functional recovery in mice. Identification of inflammatory, CNS-resident ABC/B1b cells that are conserved across species is critical as they have the potential to be sequestered from peripheral immunotherapies and/or contribute to age-related neurodegenerative diseases.

Introduction

Ischemic stroke risk increases with age, and older patients have significantly reduced post-stroke recovery [1, 2]. Rodent models of ischemic stroke replicate the age-related decline in functional recovery attributed, in part, to the aging immune system [3, 4]. Acutely after stroke, there is an influx of innate immune cells in response to damage-associated molecular patterns (DAMPs) originating from CNS cell stress and death [5, 6]. However, during chronic stroke there is a switch to adaptive immune cell infiltration (B and T cells) that persists for months to years after injury [7]. Throughout life, the majority of B cells arise from hematopoietic stem cells in the

bone marrow [8], though B1 progenitors also exist in peripheral compartments (e.g. peritoneal

and pleural cavities) [9]. B cells progress through developmental stages leading to diverse B cell

receptors (BCRs) [9] before leaving the bone marrow and travelling to secondary lymphoid

organs, including the spleen and lymph nodes, where they develop into mature B cells and await

antigen exposure [8]. Aging induces the additional formation of a Tbet$^+$ B cell population [10]

known as age-associated B cells (ABCs). ABCs are found in many peripheral compartments and

comprise 30-50% of splenic B cells in aged animals. ABCs are memory-like, antigen-

experienced, or quiescent B cells that have inflammatory effects depending on disease and

tissue context [11]. Despite thorough characterization of ABC phenotypes outside of the CNS,

few studies have investigated brain-resident ABCs in aging and after stroke [12].

Previous studies illustrate a detrimental effect of post-stroke B cell populations, including

worsened cognitive outcome and CNS-directed autoreactivity [13-15], while others showed a

protective role for anti-inflammatory regulatory B cells [16-18]. Remarkably, chronic depletion of

CD20$^+$ B cell subsets with anti-CD20 antibody after stroke improved cognitive outcomes in young

male mice [14] and aged male mice reconstituted with young-male bone marrow also

demonstrate improved post-stroke deficits [3]. ABCs specifically induce microglia phagocytosis in

male mice, with unclear effects on stroke outcome [12], while previous ABC characterization via

single-cell RNA sequencing (scRNAseq) in the periphery only used aged female mice [19]. Of

note, oligoclonal bands found in the cerebrospinal fluid (CSF) of stroke patients suggest local

antibody synthesis by brain/CSF-resident B cells [20], and patients with elevated CSF antibodies

for myelin are at higher risk of post-stroke cognitive decline [21]. Taken together, larger

preclinical studies are needed to characterize brain-resident B cell development into ABCs

during aging, across sexes, and after ischemic stroke.

To better understand how brain-resident B cells contribute to post-stroke outcomes, we

utilized flow cytometry, scRNAseq, and BCR sequencing (scBCRseq) of brain-localized B cells

from healthy aged and post-stroke mice. We show that ABC populations expand in both sexes,

with novel progenitor B1b cell subsets that directly develop into ABCs with pro-inflammatory

phenotypes and expanded clonality after stroke. This highlights a conserved CNS-resident

population that may not be exposed to peripheral immunotherapies. Finally, we identify ABC

populations in aged post-mortem human brains. Our results reveal that with aging, brain-

localized progenitor B cells develop into a proinflammatory ABC phenotype, shared between

sexes, that may inhibit recovery after ischemic stroke, with conserved populations present in the

aging human brain.

Methods

All experiments were performed in accordance with the University of Kentucky IACUC

under accreditation of AAALAC. Aged (>18 months) male and female C57BL/6JN mice were

obtained from the National Institute on Aging. Mice were socially housed and kept in a 14:10

light-dark cycle room with free access to food and water. Methods for tMCAo, behavioral testing,

brain processing, B cell isolation, flow cytometry, immunohistochemistry, scRNAseq, scBCRseq,

scRNAseq processing & data analysis, high dimensional weighted gene co-expression analysis,

transcription factor regulatory network analysis in hdWGCNA, pseduotime trajectory analysis,

external data processing, label transfer, and multicompartment aged B cell analysis, clonal

analysis, Alpha Fold 3 structural prediction, and human rapid postmortem brain processing are

included in the Supplemental Methods document.

Results

ABCs are present in aged murine brains and increase after stroke

Flow cytometry on aged post-stroke male and female mouse brains 3 weeks following a

30-min transient middle cerebral artery occlusion (tMCAo) identified B and T cell populations

(Supplemental (S)Fig.1), with specific CD11c$^+$ and CD11b$^+$CD11c$^+$ ABCs increasing in all three

brain regions (injured cortex, uninjured cortex, cerebellum) after stroke, but by a much larger

magnitude in males than in females (Fig.1a,b: statistics in STable 1). A CD23$^+$CCR7$^+$ B cell

memory/activated subset expressing the chemokine receptor CCR7 (facilitating tissue

diapedesis) [22], however, exhibited the greatest sex-specific interactions following stroke

(Fig.1c). It has been suspected that ABCs may differentiate into autoantibody-secreting cells and

transition directly into plasma cells after exposure to CNS antigens [23]. Interestingly, CD138$^+$

plasma cells were highest in the aged uninjured female left hemispheres, but also demonstrated

significant effects of injury, brain region, and sex interactions (Fig.1d).

To identify associations of brain-localized B cell subsets with functional recovery in aging

mice, we performed an open field assay to assess anxiety like-behavior, and rotarod assay to

assess motor coordination. Open field showed that aged mice, in general, remain in the

periphery, indicating anxiety, with no effect of stroke on exploration or anxiety-like behaviors

(SFig.1e,f; STable 2). For rotarod, mice were pre-trained for 14 days (SFig.1g) and assessed at

baseline and once weekly for 3 weeks post-stroke. Post-stroke aged male mice showed

sustained deficits in motor coordination, as reported previously in aged animals [3] (Fig.1e). In

both sexes, 3-week motor recovery showed no association with ABC subsets but instead

negatively correlated with the number of plasma cells in the ischemic hemisphere (Fig.1f; males

R^2 =0.978, females R^2 = 0.535). B cells isolated from the post-stroke brain confirmed Tbet$^+$

expression [10] in a subset of B220$^+$ cells, including larger antibody-producing B cells (Fig.1g).

These findings confirm that ABC subsets are present in uninjured aged brains and increase after

stroke, with plasma cells, possibly differentiated from ABCs, contributing negatively to functional

recovery.

<u>Single-cell atlas of aged brain-resident B cells reveals developmental stages within the CNS</u>

To gain an in-depth understanding of the heterogeneity of brain-resident B cells in aging and

after stroke, we next created a multi-omic single-cell atlas using B cell-enriched single cell

suspensions isolated from aged healthy or ischemic brains 3 weeks after tMCAo (Fig.2a). Aged

post-stroke males were not included in this study due to the disproportionately high incidence of

B cell death that did not meet 10x Genomics requirement for minimum number of live cells per

animal. After pre-processing and *in silico* filtering for inclusion of CD45$^+$ non-microglia only,

12,081 total cells were obtained from uninjured aged males (n=4, 6073 cells), uninjured aged

females (n=4, 3919 cells), and 3-week post-stroke aged female mice (n=4, 2089 cells; SFig.2).

CD45$^+$ cells contained hematopoietic stem cells (HSCs), monocytes, plasma cells, and a wide

diversity of B cells (10,966 cells; CD19$^+$/CD20$^+$ (*Ms4a1*); Fig.2b,c) spanning multiple

developmental stages defined by canonical markers for each population (SFig.2b). This includes

Mki67-expressing mitotic progenitor B cells and early pro- and pre-B cells (Fig.2d), in line with

previously coarsely reported early B cells in the brain [24].

To expand on the phenotypic resolution of developing B cells (Fig. 2e) and validate our

scRNAseq data finding that CNS-resident progenitor B cells are present in aged and post-stroke

brains, we identified early developing B cell subsets (Fig.2f; STable 3). To avoid the confounding

nature of immune infiltration resulting from blood-brain barrier breakdown in ischemic regions, we

performed flow cytometry on the cerebellum, a remote uninjured region of the brain. Pre-pro-,

pro-, pre-, and immature B cell populations were present in all groups, with a significant injury

effect in pre-pro- and pro-B cell populations (p<0.05; Fig.2g). Thus, the aging CNS houses early

developmental B cell subsets creating the potential for local expansion and differentiation within

the parenchyma.

<u>Age-associated B cells develop locally within the CNS</u>

We next reclustered the CNS-resident CD19$^+$ population, resulting in 5,835 cells from

aged males, 3,646 cells from aged females, and 1,855 cells from post-stroke aged females

(SFig.2d-j). B cells were individually labeled via an anchor label transfer with previously

published aged mouse B cell annotations (GSE174834) [25] ranging in developmental stage

from mitotic B cells to ABCs (Fig.3a; SFig.3). We confirmed these populations with canonical

marker genes (Fig.3b) [26]. Cell annotation identified similar population composition shared

amongst all three groups, excluding ABCs where post-stroke females had higher mean

proportion of ABCs compared to uninjured females (13.4% in post-stroke vs. 4.4% in uninjured;

Fig.3c) consistent with our flow cytometry data (Fig.1).

To assess changes across the continuum of B cell development and differentiation, we

performed pseudotime trajectory analysis [27] leveraging the known mitotic B cell cluster. This

analysis produced a *single trajectory* from mitotic B cells, traversing pro- and pre-B cells, through

immature and mature stages to finally culminate in ABCs (Fig.3d,e). These results demonstrate a

continuum of B cell development in the aging brain culminating in transcriptionally unique

ABC/B1b cells that are significantly enriched in the 3-week post-stroke period.

To better understand this novel ABC/B1b phenotype, we reclustered these cells and

obtained two resulting populations (Fig.3f). The ABC cluster, enriched for *Tbx21* (Fig.3g),

comprises a greater proportion in post-stroke females compared to uninjured females (Fig.3h).

Cluster markers outline genes that differentiate ABCs from B1b cells (Fig.3i; SFig.4). To confirm

terminal differentiation into an ABC phenotype, we integrated a publicly available dataset with B

cells from aged mouse blood, skull bone marrow, and dura [25] with our own brain data (Fig.3j;

SFig 5a-c). CNS-resident B cell composition mirrors the composition of skull bone marrow and

dura B cells, namely via a large population of pre-B cells (SFig 5d-f) compared to blood.

Pseudotime trajectories of combined compartments show the same population of ABCs at the

end of one trajectory as found in the aged brain (Fig.3k). Furthermore, expression of *Fgl2*

(fibrinogen-like protein 2), a consistent cluster marker of ABCs, represents a novel marker of

mouse ABCs irrespective of compartment (SFig.5g).

<u>ABCs exhibit activated phenotypes</u>

Differential gene expression (DEG) identified gene changes secondary to stroke

(uninjured vs. post-stroke females) or with aging in uninjured male vs. female mice

(Supplemental Data Set 1). Post-stroke females compared to uninjured females showed 519

genes (up: 471; down: 48) to be differentially expressed after stroke (Fig.4a) while aged mice

demonstrated 593 genes (up: 430; down: 163) differentially expressed between uninjured males

versus females (Fig.4b). DEGs after stroke within female mice and sex differences with aging

resulted in an overlap of 228 (33.9%) upregulated genes and 40 (23.4%) downregulated genes,

with 83% downregulated genes after stroke shared with aging males (Fig.4c). Biological process

gene ontology (GO) analysis of upregulated genes after stroke revealed terms enriched for

cytoplasmic translation, mature B cell differentiation, cytokine production involved in

inflammatory response, and positive regulation of B cell proliferation (SFig.6a). Biological

process gene ontology of stroke upregulated genes was very similar to GO terms seen in

between males and females in healthy aging, with terms enriched for increased cytoplasmic

translation, protein metabolism, positive regulation of cytokine production, both suggesting

increased metabolic activity and activation in response to inflammation. Taken together, these

results suggest a link between brain-resident B cell activation responses of stroke-induced

inflammation and sex differences in aging, with high pro-inflammatory markers in males even in

the uninjured state.

We utilized high dimensional Weighted Gene Co-expression Analysis (hdWGCNA) [28] to

create correlated gene expression modules to investigate gene-specific changes across the B

cell developmental trajectory. After dynamic tree splitting, 7 modules were present that

represented both B cell population-localized and diffuse gene expression networks (Fig.4d; SFig.

6b,c). Interrogation of module-specific functionality revealed modules significantly enriched for B

cell activation and proliferation (M3), chromosome reorganization (M1, M5), cellular respiration

and mitochondrial activity (M4), Rho GTPase activity (M2, M6), and ribosomal genes (M7;

Fig.4e,f). Genes encompassed within M3 relating to B cell activity were highly enriched in mature

B cells (*Stap-1, Fcrl1, Gpr183*) and ABCs (*Cd81, Cd22, Themis2*; SFig.6d). We examined CNS-

resident B cell module enrichment across stroke and aging conditions by computing differential

module expression scores, utilizing the top 1000 genes in each module. Module M3, containing

genes associated with B cell activation and BCR signaling and enriched in mature B cells and

ABCs, was significantly upregulated in post-stroke vs. uninjured females ($Log_2FC=0.583$;

$P_{adj}=5.19E-45$) as well as in uninjured males vs. uninjured females ($Log_2FC=0.588$; $P_{adj}=2.68E-$

81). The same pattern of upregulation can be seen in M6 with stroke ($Log_2FC=0.564$; $P_{adj}=2.79E-$

16) and in uninjured males ($Log_2FC=0.969$; $P_{adj}=2.67E-106$; Fig.4g). M6 is also enriched in

mature B cells and ABCs. M7, albeit not B cell lineage-specific, demonstrates a conserved global

increase in GTPase motility and gene expression similarly upregulated with stroke and in

uninjured males. Utilizing Rank-Rank Hygrometric Overlap, we found a significantly strong

correlation (r=0.64; $p=2.2*10^{-16}$) between transcriptomic profiles of post-stroke females vs.

uninjured females and uninjured male vs. uninjured female (Fig.4g; SFig.6e,f). These results

suggest that this transcriptional signature, shared between stroke and aging sex differences and

seen in coarse DEG analysis, is related to B cell activation in mature B cell and ABC/B1b CNS-

localized populations.

Transcription factor regulation network analysis implicates *Cebpb* regulon driving ABC activation

237   To understand if the B cell activation transcriptional signature is driven by specific

transcription factors (TF), we further utilized hdWGCNA [28] to investigate the transcriptional

programs within the identified modules by utilizing machine learning models to infer regulatory

relationships between TFs and their predicted targets [29, 30]. The increase in B cell activation in

uninjured aged males over uninjured aged females (module M3) is driven by a TF regulon

ensemble encompassing *Cebpb, Junb, Fos, Nr4a1*, and *Bcl6.* These TFs are not only

differentially expressed, excluding *Bcl6*, but their positive target genes are upregulated and

differentially expressed in uninjured male B cells compared to females (Fig.4h; SFig.7).

Comparing differential TF regulons between post-stroke females and uninjured females reveals

Cebpb and *Irf8* (Fig.4i; SFig.7f). Their target genes are also differentially expressed in post-

stroke females relative to uninjured females, specifically – and again – in B cell activation module

M3. *Cebpb* was previously implicated in a B cell phagocytic phenotype after injury [31]. These

findings outline a potential TF program that drives sex-specific B cell activation via a shared TF,

Cebpb, that is also activated after stroke in females.

We calculated hdWGCNA module pseudotime trajectories to investigate how the

modules changed within our developmental trajectory. Modules M3, M6, and M7, previously

identified to be rich in B cell activation genes, have the highest module expression scores in the

ABC/B1b population (SFig.8a), suggesting increased B cell activation in ABCs after stroke and in

uninjured males compared to uninjured females. We also investigated if any previously identified

TFs were significantly localized across pseudotime, with a specific focus on mature B cells and

ABC/B1b cells. Investigation of TF network dynamics across differentiation demonstrates that

early B cell populations are Myb^+ & $Bach2^{high}$ and, during transition, are marked by a loss of *Myb*,

previously shown to induce Tbet (*Tbx21*) after its loss while decreasing *Bach2* [32]. Concurrently,

the induction of *Myc* and *Cebpb* occurs in mature and late-stage ABC/B1b populations (Fig.4j),

suggesting the development into a phagocytic phenotype [24]. We further confirmed the previous

observation of B cell development phenotype-specific TF recruitment across compartments *of*

Myb, Bach2, Myc and *Cebpb* (SFig.8c). These findings implicate multiple TFs in the concerted

activation of mature B cells and ABCs in the brain.

<u>Stroke induces clonal expansion of IgM$^+$ B cells in the aged brain</u>

We performed concurrent scBCRseq to investigate the effects of aging and stroke on

brain-resident BCR repertoires. Density plots laid over the initial UMAP with all CD45$^+$ cell

populations confirm BCRs are present in our identified B cell clusters across all groups (Fig.5a).

We defined BCR clones by identical VDJ gene usage in both heavy and light chains, allowing for

amino acid changes and somatic hypermutation. Hyperexpanded clones (defined as comprising

greater than 10% of the total BCR repertoire) were found primarily in the ABC/B1b and plasma

cell clusters (Fig.5b). Clonal network analysis identified BCR sequence similarity amongst cell

populations using either plasma cells or ABC/B1b cells (Fig.5c) as the starting node. Both the

plasma cell cluster and ABC/B1b cell cluster share clonal similarity with other B cell

developmental populations, again indicating the potential for CNS-specific proliferation. Post-

stroke females exhibited the largest proportion of hyperexpanded clones, while uninjured males

had no hyperexpansion (Fig.5d,e). Shannon index scores confirmed BCR clonal diversity [33], in

which a low score indicates decreased diversity presumably due to clonal expansion. Uninjured

females had a Shannon index of 4.12 vs. post-stroke females (4.00; Fig.5f), with the differences

between groups largely driven by one mouse. Interestingly, in all groups, the predominant heavy

chain constant region was IgM (Fig.5g), indicating minimal class switching. Amongst the

hyperexpanded clonotypes, *IGHV7-3* was the most expressed variable gene in both post-stroke

and uninjured females (Fig.5h).

We generated ABodyBuilder2 Fv models for the six representative antibodies (top two

clonotypes per group) and predicted paratope residues with proABC-2 (Fig.5i). The top clone in

uninjured animals of both sexes was in the ABC/B1b cell cluster, while the top post-stroke clone

was in the plasma cell cluster (Fig.5i). Using a conservative cutoff (pt ≥ 0.45), antibodies

contained a median of 6.5 predicted contact residues (range 3–7), with most contacts on the

heavy chain (median 4 residues; range 3–5). At the sensitivity cutoff (pt ≥ 0.40), the predicted

interface expanded to a median of 18.5 residues (range 18–22) while remaining strongly

complimentary-determining region (CDR)-enriched (median 89.2% of residues in CDRs).

Predicted contacts consistently involved CDR-H3 (positions 95–102; present in all six antibodies),

which is the largest portion of the binding surface and is key to determining antibody specificity,

whereas no CDR-H1 residues exceeded pt ≥ 0.45. Residues above each threshold were

mapped onto the Fv structures to visualize candidate binding surfaces for future antigen-focused

experiments.

Identification of ABC populations in human post-mortem brains

The presence of brain-localized ABCs was confirmed in humans who were participants in

the University of Kentucky Alzheimer's Disease Research Center (UK-ADRC) autopsy cohort.

Details of the recruitment, inclusion/exclusion criteria, and follow-up from the UK-ADRC cohort

have been published previously [34, 35]. An emphasis of this cohort is on the rapid post-mortem

(RPM) biobank, with tissue collection near time of death (autopsies usually performed ~2-3 hours

postmortem). Samples of left and right motor cortex and cerebellum (lacking dura) were collected

from 14 RPM donors with an average age of death at 86 years (Fig.6a; STable 4). Among the

included participants, 77% (n=10) had cognitive impairment documented before death, with 69%

having the diagnosis of dementia (n=9). Three study donors (23%) had a documented prior

history of stroke. Results of left and right brain samples were averaged after confirming no cell

population differences between hemispheres. Both CD11b$^+$ and CD11c$^+$ ABCs (Fig. 6b-d) were

broadly present within the CD19$^+$ B cell populations isolated from every brain region, which also

showed high CD23$^+$ expression (Fig.6e). The proportion of CD11b$^+$ ABCs in both the cerebellum

and motor cortex showed no significant sex differences, though males tended to have a lower

proportion than females in the cerebellum (p=0.093; STable 5). CD11c$^+$ ABCs comprised a wide

range, with proportions ranging from 2-75% of B cells. There were no significant differences

detected in ABC proportion between sexes in either location, though males exhibited a significant

increase in cerebellar CD11c$^+$ ABCs with increasing age, a correlation not found within the same

RPM donors for the CD11b$^+$ ABCs (Fig.6f,g). The 3 donors with prior stroke exhibited some of

the lowest representative values for CD11c$^+$ ABCs. We confirmed ABC presence in one of the

RPM brain donors with prior stroke (82-year-old male) using immunohistochemistry to confirm

CD11c$^+$ ABCs in parenchyma and not in unperfused vasculature or leptomeninges (Fig.6h).

Investigation of a scRNAseq dataset of human brain and leptomeningeal B cells (GSE288856)

[36] in Alzheimer's Disease (AD) verified that CD11c$^+$ (*Itgax*) T-Bet$^+$ (*Tbx21*) ABCs are

proportionally enriched in AD patients (40%) compared to case controls (18%), along with

phagocytic and immature B cell populations (Fig.6i).

Discussion

We identified that stroke, in the context of aging, results in an expansion of pro-

inflammatory ABCs, and that the aged brain houses early parenchymal-based B cell progenitors.

Trajectory analysis demonstrated a continuum of B cell development from these early

progenitors through immature and mature subsets to terminally differentiate into the stroke-

expanded *Bach2*^{low} *Cebpb*$^+$ ABC/B1b phenotype. We corroborated the finding of ABCs in human

cortical and cerebellar tissue, identifying populations that are specifically affected by sex and age.

If there is an early developmentally-derived CNS B cell pool maintained by local progenitors as

our study supports, treatments can be developed to specifically target this niche, particularly in

aged individuals with impaired peripheral B cell development.

The *de novo* development of B1 cells is considered to occur from the yolk sac (first wave)

and fetal liver (second wave and predominant source) prior to bone marrow (third wave), though

populations are largely replenished with age through self-renewal driven by Bach2 [37]. Of

critical note, IgM$^+$ B1 cells canonically reside in pleural and peritoneal cavities, and the presence

of Bach2 is a survival factor for B1 cells in peripheral niches [9, 38]. To our knowledge, B1 cells

have not been studied in the brain. Interestingly, B1a and B1b B cells are increased in the

cervical lymph nodes of 3xTgAD Alzheimer phenotype disease mice [39], and innate-like

marginal zone B cells play a role in post-stroke infection prevention [40]. More recent work

identified early proliferating B cells in young brain, leptomeninges, and dura, with brain

expansion in the acute disease phase of experimental autoimmune encephalomyelitis (EAE), a

preclinical model of multiple sclerosis [41]. However, our data are the first to localize the

development of these functionally-relevant innate-like B cells to the CNS in the aged brain (>18

mos. old) and after ischemic stroke.

ABCs are memory-like or antigen-experienced B cells which have been most extensively

studied in models of autoimmune diseases like multiple sclerosis and lupus [11]. Little is known

about their role(s) in the healthy aged brain and after stroke [12], though a previous study

showed that CD11b^{high} ABCs induce changes in microglia that result in higher rates of microglial

phagocytosis 7 days post-stroke [12]. We, however, are the first to show that these B cells may

terminally differentiate locally in the brain as opposed to the periphery. This is significant as it

implicates the brain as a functioning ectopic lymphoid tissue both in normal aging and after

stroke, supporting sustained B cell development. Further studies should determine if enhanced

post-stroke differentiation requires the help of T cells, as prior studies have shown [14], or if this

is a T cell-independent process more akin to B1 maturation. Interestingly, our flow cytometry

data show that these populations increase in both hemispheres after stroke, not only the

ischemic hemisphere, a finding in line with our prior studies that identified increased B cell

populations remote from the infarct that support neurorecovery [16, 42].

Meningeal B cells originate predominantly from the calvaria bone marrow and can enter

the brain directly, bypassing systemic circulation [43]. We are limited to conclude that early

immature B cells divide and transition through maturity *exclusively* within the brain and do not

migrate from the periphery or dura. However, BCR analysis shows expansion of IgM$^+$ brain-

resident mature B cells, ABCs, and plasma cells, suggesting a within-brain transition for all B cell

subsets. Additionally, we found a strong negative correlation between total plasma cells in the

CNS and post-stroke motor performance. Antibody-producing B cells contribute to post-stroke

cognitive impairment in preclinical models [14], with IgM antibodies remaining in the brain of

stroke patients for years [14]. Interestingly, our work did not identify class-switched

hyperexpanded IgG or IgA BCRs, which confirms similarity to the polyreactive, nonspecific IgM$^+$

autoantibodies traditionally formed by B1b cells. Clinical data show a detrimental association of

self-reactive antibodies and cognitive decline [44]. It remains unknown what the antigen targets

are for the expanded clones identified in this study; future studies will be required to elucidate

possible B cell-derived CNS autoantibodies and how they may hinder stroke recovery. But of

note, only B cells from females were hyperexpanded and may contribute to brain-localized

plasma antibody-producing populations though the caveat remains we did not scBCRseq post-

stroke males.

Male post-stroke mice exhibit the greatest increases in ABC populations in flow cytometry

data. However, as mentioned we could not isolate sufficient live male post-stroke B cells at 3

weeks post-stroke to perform scRNAseq in this group, highlighting a limitation of this study.

Advances in single cell sequencing technologies that allow for sequencing of fixed whole cells

may overcome this technical hurdle [45]. Large sequencing efforts in other brain-resident

immune cells, such as microglia, demonstrate similar age-associated sex differences in

activation of male immune cells [46]. This transcriptional signature of activation was also

upregulated after stroke in females in ABCs and, after TF analysis, was inferred to be controlled

by *Cebpb. Cebpb* was previously identified to be increased in peripheral blood mononuclear cells

of elderly patients [47] and has been implicated in inducing post-injury B cells into a myeloid like

phagocytic phenotype [48], consistent with previously described ABC function [49]. Future

studies that directly perturb B cell *Cebpb* in aged rodent models are required to elucidate how

Cebpb and its target genes work to increase B cell activation, specifically in ABCs, and their

functional relevance to health and recovery in the aging brain.

Our postmortem data confirmed the presence of both $CD11b^+$ and $CD11c^+$ ABC populations

in the human brain, with a range of expression up to 80% of total $CD19^+$ B cells. Most B cells

also expressed CD23, which is associated with mature and activated B cells [50]. We validated

these populations in a publicly available human scRNAseq data set to show a significant ABC

enrichment in Alzheimer's disease. We found gender-specific and brain location-specific

differences in ABC populations, with caveats related to sample sizes that limited statistical power.

We also noted that $CD11b^+$ ABCs trended to be higher in females when isolated from the

cerebellum. $CD11b^+$ ABCs in healthy human donor blood exhibit migratory properties [51], and

CD11b facilitates B cell migration by aiding in adhesion to endo- and sub-endothelium in the

vasculature via the intercellular adhesion molecule 1(ICAM1) [52]. These peripheral B cells

express CD11b, regardless of chemokine presence and attraction, and also expressed CD27, a

memory B cell marker [52]. Future studies should include CD27, CCR7, and CXCR5, the latter

receptors instrumental in homing for CNS-specific migration from parenchymal ectopic lymphoid

structures [53], to confirm migratory versus CNS-localized progenitor B cell pools in aged

humans.

In conclusion, this study is the first to identify a B cell resident lineage in the aged brain

that develops from early progenitors to ABCs. We also show that aged male mice have a

substantially increased baseline B cell inflammatory state in the brain over female counterparts.

Stroke induced clonal expansion of IgM^+ ABC/B1b and plasma cells that exhibit a phagocytic

gene signature. We also confirmed developing, mature and ABC B cell subsets in human

parenchyma. These findings have important implications for future work examining sex

differences, brain injuries, and neurodegenerative diseases in the aging brain. Stroke, traumatic

brain injury, and Alzheimer's disease have all had clinical trials with peripheral immunotherapies

that have shown a broad range of results in terms of efficacy and safety. It is possible that

detrimental – or perhaps even some beneficial – immune effects originate from CNS-localized,

self-renewing B cell pools sequestered from peripheral immunotherapies and thus may require

unique CNS targeting for therapeutic efficacy in diseases of aging.

Acknowledgments

Thank you to the brain donors and their supportive families for their invaluable

contribution to this work. These studies are supported by grants from the National Institutes of

Health to AMS: R01NS088555, RF1NS088555; AMM: F30NS141486; MKC: T32AG057461; TU:

T32NS077889, 3RF1NS088555-07A1S1; KMC: T32NS077889; SJC: R01NS129668; UK-ADRC:

P30 AG072946. Services and support for scRNAseq experiments were provided by the College

of Arts & Sciences Imaging Center at the University of Kentucky. This work was supported by the

UK Light Microscopy core facility (RRID: SCR_026405), Flow Cytometry & Immune Monitoring

core (RRID: SCR_026358) and Biospecimen Procurement & Translational Pathology Shared

Resource Facility (P30CA177558).

Data Availability:

Publicly available B cell sequencing data was obtained from GSE174834. Raw and Processed

data from this study will be available upon publication and deposit on a publicly available site

with link to ODC-Stroke.

Figures

Figure 1: B cell subsets demonstrate sex- and brain area-dependent expression patterns following ischemic stroke that impairs long-term motor recovery.

Flow cytometry in immune cells isolated from the left and right hemispheres and cerebellum (designated x axis) in uninjured mice and mice at 3-weeks following a transient middle cerebral artery occlusion (tMCAo; red symbols) show sex-specific interactions for aged females (circles; green bars) vs. males (squares; blue bars). This includes **(a)** CD11c$^+$ and **(b)** CD11b$^+$CD11c$^+$ ABCs, with additional interactions for **(c)** CD23$^+$CCR7$^+$ memory B cell subsets and **(d)** plasma cells**. (e)** tMCAo induced motor deficits in both aged males and females without an effect of sex. Of all B cell subsets, only **(f)** plasma cells showed a negative correlation with functional recovery in both females (top graph) and males (bottom graph). **(g)** To confirm expression of the canonical ABC transcription factor Tbet, immunostaining in B cells isolated from the post-stroke brain show high B220 expression (green; left panel) and more reduced Tbet expression (red, middle, right panels) with very large B cell subsets remaining with high Tbet expression.

Figure 2: scRNAseq of B cells in healthy aging and after ischemic stroke

(a) Experiment design of B cell enriched cell suspension scRNAseq & scBCRseq isolated from healthy aged (Female n = 4; Male n = 4) and 21-day post-stroke mouse brains (Female n = 4; Biorender). **(b)** UMAP of 12,081 total CD45+ cells annotated via canonical marker genes. Hemopoietic stem cells (HSCs; n = 656), monocytes (n = 180), plasma cells (n = 279), all B cells (n = 10,966). **(c)** Gene feature plots of Ms4a1 (Cd20) & Cd19 on all Cd45+ cells confirm B cell lineage. **(d)** Dotplot of immature B cell markers across CD45+ cells show all developmental stages to age-associated B cell (ABC) populations all unique from the monocyte lineage. **(e)** Schematic of B cell development phenotypes from Biorender. **(f)** Number of immature B cell subsets in uninjured aged mouse cerebellum and at 3wks after transient middle cerebral artery occlusion (tMCAo). Blue circle = naïve for both males (left) and females (right); purple squares = tMCAo.

4
466

Figure 3: B cells develop into ABCs in the brain and other compartments

(a) UMAP of 10,966 brain-resident B cells annotated via canonical marker genes and via an anchor-based label transfer from a published multi-compartment B cell scRNAseq dataset. Mitotic B cells (n = 509), Pro-B cells (n = 203), Pre-B cells (n = 2251), Immature B cells (n = 6141), Mature naïve B cells (n = 659), ABCs (n = 710), B1b cells (n = 493). **(b)** Top 5 B cell population gene markers per identified B cell subset ranked by Log2FC. **(c)** B cell UMAP split by treatment condition (left), with B cell population proportions split by treatment condition and

shown in distribution graphs (right). **(d)** UMAP of 10,966 B cells colored Monocole3 pseudotime
with trajectories branch tree plotted. Root node labeled by white circled 1 (in dark purple area),
determined by identifying earliest branch node inside of *KI67*+ mitotic B cells. **(e)** Violin nested
boxplot of pseudotime values across B cell phenotypes ordered increasingly by mean
pseudotime**. (f)** UMAP of 1018 ABC/B1b cells colored by phenotype (ABCs, purple; B1b cells,
pink). **(g)** Gene feature plot of *Tbx21* (Tbet), the canonical ABC transcription factor. **(h)** ABC/B1b
cells split by treatment condition in both UMAP (left) and distribution bar graphs (right). **(i)** Dotplot
of top 10 ABCs & B1b cell marker genes (ABCs, bottom row). **(j)** UMAP of a multicompartment
integrated 29,806 B cells split across 4 compartments (Blood, Skull Bone Marrow, Dura, Brain)
and annotated via identified marker genes to identify B cell subsets. **(k)** Monocole3 pseudotime
trajectories of integrated atlas with root node selected from earliest tree node designated as #1 in
the dark purple mitotic cell population.

Figure 4: Gene and transcription factor regulator networks reveal shared inflammatory
signaling between uninjured male and post-stroke female B cells

(a-b) Volcano plots of differentially expressed genes between treatment conditions (Log2FC >
0.25 or < -0.25; Padj < 0.05, min.pct > 0.1; Top 10 genes by Log2FC labeled). **(c)** Venn diagram
of significant up- and downregulated genes shared between stroke female vs uninjured female
and uninjured male vs uninjured female. **(d)** hdWGCNA Module expression score feature plots of
first 1000 genes in each module on 10,966 brain resident B cells. **(e)** Top 5 significant Gene
Ontology (GO) Biological Process 2025 terms for each module via Enrichr. Dot size is Enrichr
enrichment score and colored by -Log10 adjusted P value. **(f)** Top 5 significant Reactome
Pathway 2024 terms for each module via Enrichr. Dot size is Enrichr enrichment score and
colored by -Log10 adjusted P value. **(g)** hdWGCNA differential module score heatmap (Log2FC
> 0.2 or < -0.2, Padj < 0.05, Wilcox rank sum, pseudocount = 0.01; ** Padj < 0.01, *** Padj, <
0.001). **(h-i)** Transcription Factor (TF) differential regulon scatter plot across treatment conditions
where predicted positive relationship regulons (promoting) plotted on the X-axis and predicted
negative relationship (suppressing) regulons plotted on the Y-axis. TFs are colored by module
association (See hdWGCNA methods for more detail). **(j)** Dotplot of *Myb*, *Bach2*, *Myc*, *Cebpb*
across B cell phenotypes.

Figure 5: Stroke induces hyperexpansion of IgM⁺ ABC/B1b B cells and plasma cells

(a) UMAP of B cell clusters overlayed with BCR density contour, confirming sequenced BCRs were present in the expected populations across groups. **(b)** Clone size across the identified B cell subtypes. **(c)** BCR clonal network analysis showing the network with plasma cells as the primary node (left) and the ABC/B1 cluster as the primary node (right). **(d)** Clonal expansion status across groups. Injured females showed the highest proportion of hyperexpanded cells, followed by uninjured females. **(e)** UMAP of B cells showing expansion status across. **(f)** Shannon diversity indices show minimal difference in diversity across groups. **(g)** IgH constant chain usage is large IgM across groups. **(h)** variable gene usage of hyperexpanded clones in uninjured and post-stroke females. Uninjured males did not have any hyperexpanded clones. **(i)** top expanded clone in each group depicted on UMAP of B cells. post-stroke female top clone is in the plasma cell cluster while uninjured males and females are ABC/B1b cluster. Structural predictions of binding regions are depicted. Red region is 0.45pt confidence, orange region is 0.40pt confidence.

Figure 6: ABCs are present in human rapid post-mortem post-stroke brain

(a) Experimental overview for collection of rapid post-mortem human brain samples. **(b)** Gating strategy for flow cytometry of human brain samples. Samples were first gated for time, single cells, and live cells (not shown). **(c-e)** CD11b⁺ **(c)** and CD11c⁺ **(d)** ABCs and CD23⁺ mature B

cells **(e)** in human motor cortex and cerebellum expressed as percent of live CD45$^+$CD19$^+$ cells.
Females = green circles. Males = blue squares. Red depicts donors with a prior reported history
of stroke. **(f-g)** correlation of age with percentage of CD11b and CD11c ABCs in cerebellar tissue
show male-only increase with age for CD11c$^+$ ABCs. **(h)** Immunohistochemistry staining of post-
mortem human brain for MAP2 (green, neurons), B220 (red), CD11c (purple) identifies B220$^+$
CD11c$^+$ MAP2$^-$ B cells (arrow). **(i)** UMAP of human B cells from GSE288856 of brain and
leptomeninges with proportion of cell types grouped by case diagnosis of Alzheimer's disease
(AD) vs. cognitively normal control (CTRL). Dotplot of B cell and ABC markers across B cell
subsets for the B cell-related genes *Cd19*, *MS4A1*, *Tbx21, ITGAX*, and *ITGAM*.

References

References

1. Winkelmeier, L., et al., *Age and Functional Outcomes in Patients With Large Ischemic Stroke Receiving*
Endovascular Thrombectomy. JAMA Netw Open, 2024. 7(8): p. e2426007. PMID:39133490
2. Knoflach, M., et al., *Functional recovery after ischemic stroke--a matter of age: data from the Austrian*
Stroke Unit Registry. Neurology, 2012. 78(4): p. 279-85. PMID:22238419
3. Ritzel, R.M., et al., *Aging alters the immunological response to ischemic stroke.* Acta Neuropathol, 2018.
136(1): p. 89-110. PMID:29752550
4. Tuohy, M.C., E.M.C. Hillman, R. Marshall, and D. Agalliu, *The age-dependent immune response to*
ischemic stroke. Curr Opin Neurobiol, 2023. 78: p. 102670. PMID:36586305
5. Iadecola, C., M.S. Buckwalter, and J. Anrather, *Immune responses to stroke: mechanisms, modulation,*
and therapeutic potential. The Journal of Clinical Investigation, 2020. 130(6): p. 2777-2788.
6. Iadecola, C. and J. Anrather, *The immunology of stroke and dementia.* Immunity, 2025. 58(1): p. 18-39.
PMID:39813992
7. Malone, M.K., et al., *The immunopathology of B lymphocytes during stroke-induced injury and repair.*
Seminars in Immunopathology, 2022.
8. LeBien, T.W. and T.F. Tedder, *B lymphocytes: how they develop and function.* Blood, 2008. 112(5): p.
1570-80. PMID:18725575
9. Morris, G., et al., *Emerging role of innate B1 cells in the pathophysiology of autoimmune and*
neuroimmune diseases: Association with inflammation, oxidative and nitrosative stress and
autoimmune responses. Pharmacological Research, 2019. 148: p. 104408.
10. Hao, Y., P. O'Neill, M.S. Naradikian, J.L. Scholz, and M.P. Cancro, *A B-cell subset uniquely responsive*
to innate stimuli accumulates in aged mice. Blood, 2011. 118(5): p. 1294-304. PMID:21562046
11. Cancro, M.P., *Age-Associated B Cells.* Annual Review of Immunology, 2020. 38(Volume 38, 2020): p.
315-340.
12. Korf, J.M., et al., *CD11b(high) B Cells Increase after Stroke and Regulate Microglia.* J Immunol, 2022.
209(2): p. 288-300. PMID:35732342
13. Doyle, K.P. and M.S. Buckwalter, *Does B lymphocyte-mediated autoimmunity contribute to post-stroke*
dementia? Brain Behav Immun, 2016. PMID:27531189
14. Doyle, K.P., et al., *B-lymphocyte-mediated delayed cognitive impairment following stroke.* J Neurosci,
2015. 35(5): p. 2133-45. PMID:25653369
15. Ortega, S.B., et al., *Stroke induces a rapid adaptive autoimmune response to novel neuronal antigens.*
Discov Med, 2015. 19(106): p. 381-92. PMID:26105701
16. Ortega, S.B., et al., *B cells migrate into remote brain areas and support neurogenesis and functional*
recovery after focal stroke in mice. Proceedings of the National Academy of Sciences, 2020.
117(9): p. 4983-4993.
17. Offner, H., et al., *Experimental stroke induces massive, rapid activation of the peripheral immune*
system. J Cereb Blood Flow Metab, 2006. 26(5): p. 654-65. PMID:16121126
18. Ren, X., et al., *Regulatory B cells limit CNS inflammation and neurologic deficits in murine*
experimental stroke. J Neurosci, 2011. 31(23): p. 8556-63. PMID:21653859
19. Wu, Y., et al., *Temporal single-cell analysis reveals age-associated delay in immune resolution after*
respiratory viral infection. bioRxiv, 2025. PMID:41446064
20. Prüss, H., et al., *Evidence of Intrathecal Immunoglobulin Synthesis in Stroke: A Cohort Study.* Archives
of Neurology, 2012. 69(6): p. 714-717.
21. Becker, K.J., P. Tanzi, D. Zierath, and M.S. Buckwalter, *Antibodies to myelin basic protein are*
associated with cognitive decline after stroke. Journal of Neuroimmunology, 2016. 295-296: p. 9-
11.
22. Förster, R., et al., *CCR7 Coordinates the Primary Immune Response by Establishing Functional*
Microenvironments in Secondary Lymphoid Organs. Cell, 1999. 99(1): p. 23-33.
23. Xie, G., et al., *Age-Associated B Cells in Autoimmune Diseases: Pathogenesis and Clinical*
Implications. Clin Rev Allergy Immunol, 2025. 68(1): p. 18. PMID:39960645
24. Wang, R., et al., *A novel phenotype of B cells associated with enhanced phagocytic capability and*
chemotactic function after ischemic stroke. Neural Regeneration Research, 2023. 18(11).

25. Brioschi, S., et al., *Heterogeneity of meningeal B cells reveals a lymphopoietic niche at the CNS*
borders. Science, 2021. 373(6553): p. 408-+. PMID:WOS:000679232100025
26. Morgan, D. and V. Tergaonkar, *Unraveling B cell trajectories at single cell resolution.* Trends Immunol,
2022. 43(3): p. 210-229. PMID:35090788
27. Cao, J., et al., *The single-cell transcriptional landscape of mammalian organogenesis.* Nature, 2019.
566(7745): p. 496-502. PMID:30787437
28. Morabito, S., F. Reese, N. Rahimzadeh, E. Miyoshi, and V. Swarup, *hdWGCNA identifies co-*
expression networks in high-dimensional transcriptomics data. Cell Rep Methods, 2023. 3(6): p.
100498. PMID:37426759
29. Childs, J.E., et al., *Relapse to cocaine seeking is regulated by medial habenula NR4A2/NURR1 in*
mice. Cell Rep, 2024. 43(3): p. 113956. PMID:38489267
30. Vorontsov, I.E., et al., *HOCOMOCO in 2024: a rebuild of the curated collection of binding models for*
human and mouse transcription factors. Nucleic Acids Res, 2024. 52(D1): p. D154-d163.
PMID:37971293
31. Wang, R., et al., *A novel phenotype of B cells associated with enhanced phagocytic capability and*
chemotactic function after ischemic stroke. Neural Regen Res, 2023. 18(11): p. 2413-2423.
PMID:37282471
32. Piovesan, D., et al., *c-Myb Regulates the T-Bet-Dependent Differentiation Program in B Cells to*
Coordinate Antibody Responses. Cell Rep, 2017. 19(3): p. 461-470. PMID:28423310
33. Pelissier, A., S. Luo, M. Stratigopoulou, J.E.J. Guikema, and M. Rodríguez Martínez, *Exploring the*
impact of clonal definition on B-cell diversity: implications for the analysis of immune repertoires.
Front Immunol, 2023. 14: p. 1123968. PMID:37138881
34. Schmitt, F.A., et al., *Celebrating 40 years of the University of Kentucky Alzheimer's Disease Research*
Center. Alzheimers Dement, 2025. 21(5): p. e70181. PMID:40365904
35. Schmitt, F.A., et al., *University of Kentucky Sanders-Brown healthy brain aging volunteers: donor*
characteristics, procedures and neuropathology. Curr Alzheimer Res, 2012. 9(6): p. 724-33.
PMID:22471862
36. Hobson, R., et al., *Clonal CD8(+) T☐cells populate the leptomeninges and coordinate with immune*
cells in human degenerative brain diseases. Nat Immunol, 2026. 27(2): p. 323-335.
PMID:41593242
37. Oh, S.G., J. Noh, E. Jang, and J. Youn, *The transcription repressor Bach2 is required for maintaining*
the B-1 cell population by regulating self-renewal. Front Immunol, 2025. 16: p. 1553089.
PMID:40170867
38. Oh, S.-G., J. Noh, E. Jang, and J. Youn, *The transcription repressor Bach2 is required for maintaining*
the B-1 cell population by regulating self-renewal. Frontiers in Immunology, 2025. Volume 16 -
2025.
39. Kim, K., et al., *Therapeutic B-cell depletion reverses progression of Alzheimer's disease.* Nature
Communications, 2021. 12(1): p. 2185.
40. McCulloch, L., C.J. Smith, and B.W. McColl, *Adrenergic-mediated loss of splenic marginal zone B cells*
contributes to infection susceptibility after stroke. Nature Communications, 2017. 8(1): p. 15051.
41. Florescu, A., et al., *Dynamic alterations of dural and bone marrow B cells in an animal model of*
progressive multiple sclerosis. J Exp Med, 2025. 222(12). PMID:40965509
42. Torres, V.O., et al., *B cells upregulate NMDARs, respond to extracellular glutamate, and express*
mature BDNF to protect the brain from ischemic injury. Neurobiology of Disease, 2025. 207: p.
106819.
43. Cugurra, A., et al., *Skull and vertebral bone marrow are myeloid cell reservoirs for the meninges and*
CNS parenchyma. Science, 2021. 373(6553). PMID:34083447
44. Becker, K.J., P. Tanzi, D. Zierath, and M.S. Buckwalter, *Antibodies to myelin basic protein are*
associated with cognitive decline after stroke. J Neuroimmunol, 2016. 295-296: p. 9-11.
PMID:27235342
45. Phan, H.V., et al., *High-throughput RNA sequencing of paraformaldehyde-fixed single cells.* Nat
Commun, 2021. 12(1): p. 5636. PMID:34561439

46. Li, X., et al., *Transcriptional and epigenetic decoding of the microglial aging process.* Nat Aging, 2023.
639   3(10): p. 1288-1311. PMID:37697166

640   47. Wootla, B., A. Denic, A.E. Warrington, and M. Rodriguez, *A monoclonal natural human IgM protects*
641   *axons in the absence of remyelination.* J Neuroinflammation, 2016. 13(1): p. 94. PMID:27126523

642   48. Budeus, B., A. Kibler, and R. Kuppers, *Human IgM-expressing memory B cells.* Front Immunol, 2023.
643   14: p. 1308378. PMID:38143767

644   49. Xie, G., et al., *Age-Associated B Cells in Autoimmune Diseases: Pathogenesis and Clinical*
645   *Implications.* Clinical Reviews in Allergy & Immunology, 2025. 68(1): p. 18.

646   50. Herisson, F., et al., *Direct vascular channels connect skull bone marrow and the brain surface enabling*
647   *myeloid cell migration.* Nat Neurosci, 2018. 21(9): p. 1209-1217. PMID:30150661

648   51. Kawai, K., et al., *CD11b-mediated migratory property of peripheral blood B cells.* J Allergy Clin
649   Immunol, 2005. 116(1): p. 192-7. PMID:15990794

650   52. Kawai, K., et al., *CD11b-mediated migratory property of peripheral blood B cells.* Journal of Allergy and
651   Clinical Immunology, 2005. 116(1): p. 192-197.

652   53. Weitbrecht, L., et al., *CD4+ T cells promote delayed B cell responses in the ischemic brain after*
653   *experimental stroke.* Brain, Behavior, and Immunity, 2021. 91: p. 601-614.