CNS-resident B cells develop locally into a pro-inflammatory age-associated phenotype during aging and after stroke

Annabel M. McAtee[1]*, Mathew Kenwood[2]*, Thomas Ujas[1], Mary K. Colson[1], Julie Watkins[1], Edric D. Winford[3], Katherine M. Cotter[1], Daimen Britsch[1], Dene Betz[2], Jadwiga Turchan-Cholewo[4], Jenny Lutshumba[4], Connor Stuart[1], Gargi Shah[1], Loganann Runice[1], Mark Ebbert[1], Salvatore Cherra[1], Peter T. Nelson[5], Jamie L. Sturgill[6], Nancy L. Monson[7,8,9], Mark P. Goldberg[2], Ann M. Stowe[1,4]

[1]Department of Neuroscience, University of Kentucky, 741 S. Limestone, BBSRB 479, Lexington, Kentucky 40508, USA

[2]Department of Neurology, Long School of Medicine, University of Texas at San Antonio, 7703 Floyd Curl Drive, MSC 7883, San Antonio, Texas 78229, USA

[3]Department of Neurology, Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Gertrude H. Sergievsky Center, Vagelos College of Physicians and Surgeons, Columbia University, New York, New York 10032 USA

[4]Department of Neurology, University of Kentucky, 740 S. Limestone, Kentucky Clinic J-455 Lexington, Kentucky 40536, USA

[5]Sanders-Brown Center on Aging and Department of Pathology, University of Kentucky, 800 S. Limestone, Rm 111B, Lexington, Kentucky 40536, USA

[6]Department of Microbiology, Immunology, and Molecular Genetics, University of Kentucky, 760 Press Ave., HKRB 532, Lexington, Kentucky 40506, USA

[7]Department of Neurology, University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Suite NL9.114, Dallas, Texas 75390-8813, USA

[8]Department of Immunology, University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Suite NL9.114, Dallas, Texas 75390-8813, USA

[9]O'Donnell Brain Institute, University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Suite NL9.114, Dallas, Texas 75390-8813, USA

*equal contribution

Corresponding author:
Ann M. Stowe, PhD, FAHA
Department of Neurology, BBSRB 479
741 S. Limestone St., Lexington, KY USA 40508
Phone: 001-859-323-8420
e-mail: ann.stowe@uky.edu
ORCID iD: 0000-0001-8111-4429

Preprint Server: BioRxiv

Key words: B lymphocyte, ischemic stroke, autoreactivity, B cell receptor, progenitor, age-associated B cell

The authors have no competing interests to declare.

Abstract

Aging and age-related diseases like ischemic stroke induce chronic lymphocyte recruitment into the CNS. Conflicting effects on post-stroke functional recovery, however, are secondary to the differences in responding lymphocyte populations that shift immunophenotype with injury and age. To better define CNS-localized B cell subsets, we used flow cytometry, single-cell RNA sequencing, and B cell receptor sequencing on B cells isolated from uninjured and post-stroke brains of aged male and female mice. We identified a novel B1b cell progenitor pool distinct from canonical pleural/peritoneal niches. Trajectory analysis showed progenitor transition into age-associated B cell (ABC) subsets, and clonal expansion of IgM$^+$ ABCs (ABC/B1b) and plasma cells following ischemic stroke. We also confirmed analogous ABCs and developing B cell populations in post-mortem human parenchymal tissue isolated from aged brain donors. These studies reveal unique B cell populations that can proliferate within the aging CNS and exhibit a conserved inflammatory signature across species.

Introduction

Ischemic stroke risk increases with age,[2] and older patients have significantly reduced post-stroke recovery.[1,2] Rodent models of ischemic stroke replicate the age-related decline in functional recovery attributed, in part, to the aging immune system.[3,4] Acutely after stroke, there is an influx of innate immune cells in response to damage-associated molecular patterns (DAMPs) originating from CNS cell stress and death.[5,6] However, during chronic stroke there is a switch to adaptive immune cell infiltration (B and T cells) that persists for months to years after injury.[7] Throughout life, the majority of B cells arise from hematopoietic stem cells in the bone marrow,[8] though B1 progenitors also exist in peripheral compartments (e.g. peritoneal and pleural cavities).[9] B cells progress through developmental stages leading to diverse B cell receptors (BCRs)[9] before leaving the bone marrow and travelling to secondary lymphoid organs, including the spleen and lymph nodes, where they develop into mature B cells and await antigen exposure.[8] Aging induces the additional formation of a Tbet$^+$ B cell population[10] known as age-associated B cells (ABCs). ABCs are found in many peripheral comparments[14] and comprise 30-50% of splenic B cells in aged animals. ABCs are memory-like, antigen-experienced, or quiescent B cells that have inflammatory effects depending on disease and tissue context.[11] Despite thorough characterization of ABC phenotypes outside of the CNS,[14,16] few studies have investigated brain-resident ABCs in aging and after stroke.[12]

Previous studies illustrate a detrimental effect of post-stroke B cell populations, including worsened cognitive outcome and CNS-directed autoreactivity,[13-15] while others showed a protective role for anti-inflammatory regulatory B cells.[16-18] Remarkably, chronic depletion of CD20$^+$ B cell subsets with anti-CD20 antibody after stroke improved cognitive outcomes in young male mice.[14] Aged male mice reconstituted with young-male bone marrow also demonstrate improved post-stroke deficits.[3] ABCs specifically induce microglia phagocytosis in male mice, with unclear effects on stroke outcome,[12] while previous ABC characterization via single-cell RNA sequencing (scRNAseq) in the periphery only used aged female mice.[16] Of note, oligoclonal bands found in the cerebrospinal fluid (CSF) of stroke patients suggest local antibody synthesis by brain/CSF-resident B cells,[19] and patients with elevated CSF antibodies for myelin are at higher risk of post-stroke cognitive decline.[20] Taken together, larger preclinical studies are needed to characterize brain-resident B cell development into ABCs during aging, across sexes, and after ischemic stroke.

To better understand how brain-resident B cells contribute to post-stroke outcomes, we utilized flow cytometry, single cell (sc)RNAseq, and BCR sequencing (scBCRseq) of brain-localized B cells from healthy aged and post-stroke mice. We show that ABC populations expand after in both sexes. Additionally, we identify novel progenitor B1b cell subsets that directly develop into ABCs with pro-inflammatory phenotypes and expanded clonality after stroke, highlighting a conserved CNS-resident population that may not be exposed to peripheral immunotherapies. Finally, we identify ABC populations in aged post-mortem human brains. Our results reveal that with aging, progenitor brain-localized B cells develop into a proinflammatory ABC phenotype, shared between sexes, that may inhibit recovery after ischemic stroke, with conserved populations present in the aging human brain.

Results

ABCs are present in aged mouse brains and increase after stroke

Flow cytometry on aged post-stroke male and female mouse brains 3 weeks following a 30-min transient middle cerebral artery occlusion (tMCAo) identified B and T cell populations (Extended Data Fig.1), with specific CD11c[+] and CD11b[+]CD11c[+] ABCs increasing in all three brain regions (injured cortex, uninjured cortex, cerebellum) after stroke, but by a much larger magnitude in males than in females (Fig.1a,b: statistics in Extended Data Table 1). A CD23[+]CCR7[+] B cell memory/activated subset expressing the chemokine receptor CCR7 (facilitating tissue diapedesis),[21] however, exhibited the greatest sex-specific interactions following stroke (Fig.1c). It has been suspected that ABCs may differentiate into autoantibody-secreting cells[5,16] and transition directly into plasma cells after exposure to CNS antigens.[22] Interestingly, CD138[+] plasma cells were highest in the aged uninjured female left hemispheres, but also demonstrated significant effects of injury, brain region, and sex interactions (Fig.1d).

To identify associations of brain-localized B cell subsets with functional recovery in aging mice, we performed open field, to assess anxiety like-behavior, and rotarod, to assess motor coordination recovery. Open field showed that aged mice, in general, remain in the periphery, indicating anxiety, with no effect of stroke on exploration or anxiety-like behaviors (Extended Data Fig.1e,f; Extended Data Table 2). For rotarod, mice were pre-trained for 14 days (Extended Data Fig.1g) and assessed at baseline and once weekly for 3 weeks post-stroke. Post-stroke aged male mice showed sustained deficits in motor coordination, as reported previously in aged animals[3] (Fig.1e). In both sexes, 3-week motor recovery showed no association with ABC subsets but instead negatively correlated with the number of plasma cells in the ischemic hemisphere (Fig.1f; males R^2 =0.978, females R^2 = 0.535). B cells isolated from the post-stroke brain confirmed Tbet[+] expression[10] in a subset of B220[+] cells, including larger antibody-producing B cells (Fig.1g). These findings confirm that ABC subsets are present in uninjured aged brains and increase after stroke, with plasma cells, possibly differentiated from ABCs, contributing negatively to functional recovery.

Single-cell atlas of aged brain-resident B cells reveals developmental stages within the CNS

To gain an in-depth understanding of the heterogeneity of brain-resident B cells in aging and after stroke, we next created a multi-omic single-cell atlas using B cell-enriched single cell suspensions isolated from aged healthy or ischemic brains 3 weeks after tMCAo (Fig.2a). Aged post-stroke males were not included in this study due to the disproportionately high incidence of cell death that did not meet 10x Genomics requirement for minimum number of live cells per animal. After pre-processing and *in silico* filtering for inclusion of CD45[+] non-microglia only, 12,081 total cells were obtained from uninjured aged males (n=4, 6073 cells), uninjured aged females (n=4, 3919 cells), and 3-week post-stroke aged female mice (n=4, 2089 cells; Extended Data Fig.2a). CD45[+] cells contained hematopoietic stem cells (HSCs), monocytes, plasma cells, and a wide diversity of B cells (10,966 cells; CD19[+]/CD20[+] (*Ms4a1*); Fig.2b,c) spanning multiple developmental stages defined by canonical markers for each population (Extended Data Fig.2b). This includes *Mki67*-expressing mitotic progenitor B cells and early pro- and pre-B cells (Fig.2d), in line with previously coarsely reported early B cells in the brain.[22]

To expand on the phenotypic resolution of developing B cells (Fig. 2e) and validate our scRNAseq data finding that CNS-resident progenitor B cells are present in aged and post-stroke brains, we identified early developing B cell subsets (Fig.2f; Extended Data Table 3). To avoid the confounding nature of immune infiltration resulting from blood-brain barrier breakdown of ischemic regions, we performed flow cytometry on the cerebellum, a remote uninjured region of the brain. Pre-pro-, pro-, pre-, and immature B cell populations were present in all groups, with a significant injury effect in pre-pro- and pro-B cell populations (p<0.05; Fig.2f). Thus, the aging

CNS houses early developmental B cell subsets creating the potential for local expansion and
differentiation within the parenchyma.

Age-associated B cells develop locally within the CNS

We next reclustered the CNS-resident CD19+ population, resulting in 5,835 cells from aged
males, 3,646 cells from aged females, and 1,855 cells from post-stroke aged females (Extended
Data Fig.2d-j). B cells were individually labeled via an anchor label transfer with previously
published aged mouse B cell annotations (GSE174834)[23] ranging in developmental stage from
mitotic B cells to ABCs (Fig.3a; Extended Data Fig.3). We confirmed these populations with
canonical marker genes (Fig.3b).[24] Cell annotation identified similar population composition
shared amongst all three groups, excluding ABCs where post-stroke females had higher mean
proportion of ABCs compared to uninjured females (13.4% in post-stroke vs. 4.4% in uninjured;
Fig.3c) consistent with our flow cytometry data (Fig.1).

To assess changes across the continuum of B cell development and differentiation, we
performed pseudotime trajectory analysis[25] leveraging the known mitotic B cell cluster. This
analysis produced a *single trajectory* from mitotic B cells, traversing pro- and pre-B cells, through
immature and mature stages to finally culminate in ABCs (Fig.3d,e). These results demonstrate a
continuum of B cell development in the aging brain culminating in transcriptionally unique
ABC/B1b cells that are significantly enriched in the 3-week post-stroke period.

To better understand the ABC/B1b phenotype, we reclustered these cells and obtained two
resulting populations (Fig.3f). The ABC cluster, enriched for *Tbx21* (Fig.3g), comprises a greater
proportion in post-stroke females compared to uninjured females (Fig.3h). Cluster markers
outline genes that differentiate ABCs from B1b cells (Fig.3i; Extended Data Fig.4). To confirm
terminal differentiation into an ABC phenotype, we integrated a publicly available dataset with B
cells from aged mouse blood, bone marrow, and dura[23] with our own brain data (Fig.3j; Extended
Data Fig 5a-c). CNS-resident B cell composition mirrors the composition of bone marrow and
dura B cells, namely via a large population of pre-B cells (Extended Data Fig 5d-f) compared to
blood. Pseudotime trajectories of combined compartments show the same population of ABCs at
the end of one trajectory as found in the aged brain (Fig.3k). Furthermore, expression of *Fgl2*
(fibrinogen-like protein 2), a consistent cluster marker of ABCs, represents a novel marker of
mouse ABCs irrespective of compartment (Extended Data Fig.5g).

ABCs exhibit activated phenotypes

Differential gene expression (DEG) identified gene changes secondary to stroke (uninjured
vs. post-stroke females) or with aging in uninjured male vs. female mice (Extended Data
Supplement 1). Post-stroke females compared to uninjured females showed 519 genes (up: 471;
down: 48) to be differentially expressed after stroke (Fig.4a) while aged mice demonstrated 593
genes (up: 430; down: 163) differentially expressed between uninjured males versus females
(Fig.4b). DEGs after stroke within female mice and sex differences with aging resulted in an
overlap of 228 (33.9%) upregulated genes and 40 (23.4%) downregulated genes, with 83%
downregulated genes after stroke shared with aging males (Fig.4c). Biological process gene
ontology (GO) analysis of upregulated genes after stroke revealed terms enriched for
cytoplasmic translation, mature B cell differentiation, cytokine production involved in
inflammatory response, and positive regulation of B cell proliferation. Biological process gene
ontology of stroke upregulated genes was very similar to GO terms seen in between males and
females in healthy aging, with terms enriched for increased cytoplasmic translation, protein
metabolism, positive regulation of cytokine production, both suggesting increased metabolic
activity and activation in response to inflammation (Extended Data Fig.6a). Taken together, these
results suggest a link between brain-resident B cell activation responses of stroke-induced
inflammation and sex differences in aging, with high pro-inflammatory markers in males even in
the uninjured state.

We utilized high dimensional Weighted Gene Co-expression Analysis (hdWGCNA)[26] to
create correlated gene expression modules to investigate gene-specific changes across the B
cell developmental trajectory. After dynamic tree splitting, 7 modules were present that
represented both B cell population-localized and diffuse gene expression networks (Fig.4d;
Extended Data Fig. 6b,c). Interrogation of module-specific functionality revealed modules
significantly enriched for B cell activation and proliferation (M3), chromosome reorganization (M1,
M5), cellular respiration and mitochondrial activity (M4), Rho GTPase activity (M2, M6), and
ribosomal genes (M7; Fig.4e,f). Genes encompassed within M3 relating to B cell activity were
highly enriched in mature B cells (*Stap-1, Fcrl1, Gpr183*) and ABCs (*Cd81, Cd22, Themis2*;
Extended Data Fig.6d). We examined CNS-resident B cell module enrichment across stroke and
aging conditions by computing differential module expression scores, utilizing the top 1000
genes in each module. Module M3, containing genes associated with B cell activation and BCR
signaling and enriched in mature B cells and ABCs, was significantly upregulated in post-stroke
vs. uninjured females ($Log_2FC=0.583$; P_{adj}=5.19E-45) as well as in uninjured males vs. uninjured
females ($Log_2FC=0.588$; P_{adj}=2.68E-81). The same pattern of upregulation can be seen in M6
with stroke ($Log_2FC=0.564$; P_{adj}=2.79E-16) and in uninjured males ($Log_2FC=0.969$; P_{adj}=2.67E-
106; Fig.4g). M6 is also enriched in mature B cells and ABCs. M7, albeit not B cell lineage-
specific, demonstrates a conserved global increase in GTPase motility and gene expression
similarly upregulated with stroke and in uninjured males. Utilizing Rank-Rank Hygrometric
Overlap, we found a significantly strong correlation (r=0.64; $p=2.2*10^{-16}$) between transcriptomic
profiles of post-stroke females vs. uninjured females and uninjured male vs. uninjured female
(Fig.4g; Extended Data Fig.6e,f). These results suggest that this transcriptional signature, shared
between stroke and aging sex differences and seen in coarse DEG analysis, is related to B cell
activation in mature B cell and ABC/B1b CNS-localized populations.

<u>Transcription factor regulation network analysis implicates *Cebpb* regulon driving ABC activation</u>

To understand if the B cell activation transcriptional signature is driven by specific
transcription factors (TF), we further utilized hdWGCNA[26] to investigate the transcriptional
programs within the identified modules by utilizing machine learning models to infer regulatory
relationships between TFs and their predicted targets.[27,28] The increase in B cell activation in
uninjured aged males over uninjured aged females (module M3) is driven by a TF regulon
ensemble encompassing *Cebpb, Junb, Fos, Nr4a1*, and *Bcl6.* These TFs are not only
differentially expressed, excluding *Bcl6*, but their positive target genes are upregulated and
differentially expressed in uninjured male B cells compared to females (Fig.4h; Extended Data
Fig.7a-e). Comparing differential TF regulons between post-stroke females and uninjured
females reveals *Cebpb* and *Irf8* (Fig.4i; Extended Data Fig.7f). Their target genes are also
differentially expressed in post-stroke females relative to uninjured females, specifically – and
again – in B cell activation module M3. *Cebpb* was previously implicated in a B cell phagocytic
phenotype after injury.[29] These findings outline a potential TF program that drives sex-specific B
cell activation via a shared TF, *Cebpb*, that is also activated after stroke in females.

We calculated hdWGCNA module pseudotime trajectories to investigate how the modules
changed within our developmental trajectory. Modules M3, M6, and M7, previously identified to
be rich in B cell activation genes, have the highest module expression scores in the ABC/B1b
population (Extended Data Fig.8a), suggesting increased B cell activation in ABCs after stroke
and in uninjured males compared to uninjured females. We also investigated if any previously
identified TFs were significantly localized across pseudotime, with a specific focus on mature B
cells and ABC/B1b cells. Numerous TFs were enriched in distinct populations; however, within
the top 20, only *JunB* and *Klf2* were localized to mature B cells and ABC/B1b (Extended Data
Fig.8b). Investigation of TF network dynamics across differentiation demonstrates that early B
cell populations are Myb^+ & $Bach2^{high}$ and, during transition, are marked by a loss of *Myb*,
previously shown to induce Tbet (*Tbx21*) after its loss while decreasing *Bach2*.[30] Concurrently,
the induction of *Myc* and *Cebpb* occurs in mature and late-stage ABC/B1b populations (Fig.4j),

suggesting the development into a phagocytic phenotype.[22] We further confirmed the previous
observation of B cell development phenotype-specific TF recruitment across compartments *of*
Myb, Bach2, Myc and *Cebpb* (Extended Data Fig.8c). These findings implicate multiple TFs in
the concerted activation of mature B cells and ABCs in the brain.

Stroke induces clonal expansion of IgM$^+$ B cells in the aged brain

We performed scBCRseq to investigate the effects of aging and stroke on brain-resident
BCR repertoires. Density plots laid over the initial UMAP with all CD45$^+$ cell populations confirms
BCRs are present in our identified B cell clusters across all groups (Fig.5a). We defined BCR
clones by identical VDJ gene usage in both heavy and light chains, allowing for amino acid
changes and somatic hypermutation. Hyperexpanded clones (defined as comprising greater than
10% of the total BCR repertoire) were found primarily in the ABC/B1b and plasma cell clusters
(Fig.5b). Clonal network analysis identified BCR sequence similarity amongst cell populations
using either plasma cells or ABC/B1b cells (Fig 5c) as the starting node. Both the plasma cell
cluster and ABC/B1b cell cluster share clonal similarity with other B cell developmental
populations, again indicating the potential for CNS-specific proliferation. Post-stroke females
exhibited the largest proportion of hyperexpanded clones, while uninjured males had no
hyperexpansion (Fig.5d,e). Shannon index scores confirmed BCR clonal diversity,[31] in which a
low score indicates decreased diversity presumably due to clonal expansion. Uninjured females
had a Shannon index of 4.12 vs. post-stroke females (4.00; Fig.5f), with the differences between
groups largely driven by one mouse. Interestingly, in all groups, the predominant heavy chain
constant region was IgM (Fig.5g), indicating minimal class switching. Amongst the
hyperexpanded clonotypes, *IGHV7-3* was the most expressed variable gene in both post-stroke
and uninjured females (Fig.5h).

We generated ABodyBuilder2 Fv models for the six representative antibodies (top two
clonotypes per group) and predicted paratope residues with proABC-2 (Fig.5i). The top clone in
uninjured animals of both sexes were in the ABC/B1b cell cluster, while the top post-stroke clone
was in the plasma cell cluster (Fig.5i). Using a conservative cutoff (pt ≥ 0.45), antibodies
contained a median of 6.5 predicted contact residues (range 3–7), with most contacts on the
heavy chain (median 4 residues; range 3–5). At the sensitivity cutoff (pt ≥ 0.40), the predicted
interface expanded to a median of 18.5 residues (range 18–22) while remaining strongly
complimentary-determining region (CDR)-enriched (median 89.2% of residues in CDRs).
Predicted contacts consistently involved CDR-H3 (positions 95–102; present in all six antibodies),
which is the largest portion of the binding surface and is key to determining antibody specificity,
whereas no CDR-H1 residues exceeded pt ≥ 0.45. Residues above each threshold were
mapped onto the Fv structures to visualize candidate binding surfaces for subsequent antigen-
focused experiments.

Identification of ABC populations in human post-mortem brains

The presence of brain-localized ABCs was confirmed in humans who were participants in the
University of Kentucky Alzheimer's Disease Research Center (UK-ADRC) autopsy cohort.
Details of the recruitment, inclusion/exclusion criteria, and follow-up from the UK-ADRC cohort
have been published previously.[32,33] An emphasis of this cohort is on the rapid post-mortem
(RPM) biobank, with tissue collection near time of death (autopsies usually performed ~2-3 hours
postmortem). Samples of left and right motor cortex and cerebellum (lacking dura) were collected
from 14 RPM donors with an average age of death at 86 years (Fig.6a; Extended Data Table 4).
Among the included participants, 77% (n=10) had cognitive impairment documented before
death, with 69% having the diagnosis of dementia (n=9). Three study donors (23%) had a
documented prior history of stroke. Results of left and right brain samples were averaged after
confirming no cell population differences between hemispheres. Both CD11b$^+$ and CD11c$^+$ ABCs
(Fig. 6b-d) were broadly present within the CD19$^+$ B cell populations isolated from every brain

region, which also showed high CD23$^+$ expression (Fig.6e). The proportion of CD11b$^+$ ABCs in
both the cerebellum and motor cortex showed no significant sex differences, though males
tended to have a lower proportion than females in the cerebellum (p=0.093; Extended Data
Table 5). CD11c$^+$ ABCs comprised a wide range, with proportions ranging from 2-75% of B cells.
There were no significant differences detected in ABC proportion between sexes in either
location, though males exhibited a significant increase in cerebellar CD11c$^+$ ABCs with
increasing age, a correlation not found within the same donors for the CD11b$^+$ ABCs (Fig.6f,g).
The 3 donors with prior stroke exhibited some of the lowest representative values for CD11c$^+$
ABCs. We confirmed ABC presence in one of the RPM brain donors with prior stroke (82-year-
old male) using immunohistochemistry to confirm CD11c$^+$ ABCs in parenchyma and not in
unperfused vasculature or leptomeninges (Fig.6h). Investigation of a scRNAseq dataset of
human brain and leptomeningeal B cells (GSE288856)[34] in Alzheimer's Disease (AD) verified
that CD11c$^+$ (*Itgax*) T-Bet$^+$ (*Tbx21*) ABCs are proportionally enriched in AD patients (40%)
compared to case controls (18%), along with phagocytic and immature B cell populations (Fig.6i).

Discussion

We identified that stroke, in the context of aging, results in an expansion of pro-inflammatory ABCs, and that the aged brain houses early parenchymal-based B cell progenitors. Trajectory analysis demonstrated a continuum of B cell development from these early progenitors through immature and mature subsets to terminally differentiate into the stroke-expanded *Bach2*[low] *Cebpb*[+] ABC/B1b phenotype. We corroborated the finding of ABCs in human cortical and cerebellar tissue, identifying populations that are specifically affected by sex and age. If there is an early developmentally derived CNS B cell pool maintained by local progenitors as our study supports, treatments can be developed to specifically target this niche, particularly in aged individuals with impaired peripheral B cell development.

The *de novo* development of B1 cells is considered to occur from the yolk sac (first wave) and fetal liver (second wave and predominant source) prior to bone marrow (third wave), though populations are largely replenished with age through self-renewal driven by Bach2.[35] Of critical note, IgM[+] B1 cells canonically reside in pleural and peritoneal cavities, and the presence of Bach2 is a survival factor for B1 cells in peripheral niches.[9,36] To our knowledge, B1 cells have not been studied in the brain. Interestingly, B1a and B1b B cells are increased in the cervical lymph nodes of 3xTgAD Alzheimer phenotype disease mice,[37] and innate-like marginal zone B cells play a role in post-stroke infection prevention.[38] However, our data are the first to localize the development of these functionally-relevant innate-like B cells to the CNS.

ABCs are memory-like or antigen-experienced B cells which have been most extensively studied in models of autoimmune diseases like multiple sclerosis and lupus.[11] Little is known about their role(s) in the healthy aged brain and after stroke,[12] though a previous study showed that CD11b[high] ABCs induce changes in microglia that result in higher rates of microglial phagocytosis 7 days post-stroke.[12] We, however, are the first to show that these B cells may terminally differentiate locally in the brain as opposed to the periphery. This is significant as it implicates the brain as a functioning ectopic lymphoid tissue both in normal aging and after stroke, supporting sustained B cell development. Further studies should determine if enhanced post-stroke differentiation requires the help of T cells, as prior studies have shown,[14] or if this is a T cell-independent process more akin to B1 maturation. Interestingly, our flow cytometry data show that these populations increase in both hemispheres after stroke, not only the ischemic hemisphere, a finding in line with our prior studies that identified increased B cell populations remote from the infarct that support neurorecovery.[16,39]

Meningeal B cells originate predominantly from the calvaria bone marrow and can enter the brain directly, bypassing systemic circulation.[40] We are limited to conclude that early immature B cells divide and transition through maturity *exclusively* within the brain and do not migrate from the periphery or dura.[22] However, BCR analysis shows expansion of IgM[+] brain-resident mature B cells, ABCs, and plasma cells, suggesting a within-brain transition for all B cell subsets. Additionally, we found a strong negative correlation between total plasma cells in the CNS and post-stroke motor performance. Antibody-producing B cells contribute to post-stroke cognitive impairment in preclinical models,[14] and IgM antibodies remain in the brains of stroke patients.[14] Interestingly, our work did not identify class-switched hyperexpanded IgG or IgA BCRs, which confirms similarity to the polyreactive, nonspecific IgM[+] autoantibodies traditionally formed by B1b cells. Clinical data show a detrimental association of self-reactive antibodies and cognitive decline.[41] It remains unknown what the antigen targets are for the expanded clones identified in this study; future studies will be required to elucidate possible B cell-derived CNS autoantibodies and how they may hinder stroke recovery. But of note, only B cells from females were hyperexpanded and may contribute to brain-localized plasma antibody-producing populations though the caveat remains we did not scBCRseq post-stroke males.

Male post-stroke mice exhibit the greatest increases in ABC populations in flow cytometry
data. However, as mentioned we could not isolate sufficient live male post-stroke B cells at 3
weeks post-stroke to perform scRNAseq in this group, highlighting a limitation of this study.
Advances in single cell sequencing technologies that allow for sequencing of fixed whole cells
may overcome this technical hurdle.[42] Large sequencing efforts in other brain-resident immune
cells, such as microglia, demonstrate similar age-associated sex differences in activation of male
immune cells.[43] This transcriptional signature of activation was also upregulated after stroke in
females in ABCs and, after TF analysis, was inferred to be controlled by *Cebpb. Cebpb* was
previously identified to be increased in peripheral blood mononuclear cells of elderly patients[44]
and has been implicated in inducing post-injury B cells into a myeloid like phagocytic
phenotype,[45] consistent with previously described ABC function.[46] Future studies that directly
perturb B cell *Cebpb* in aged rodent models are required to elucidate how *Cebpb* and its target
genes work to increase B cell activation, specifically in ABCs, and their functional relevance to
health and recovery in the aging brain.

Our RPM data confirmed the presence of both CD11b$^+$ and CD11c$^+$ ABC populations in the
human brain, with a range of expression up to 80% of total CD19$^+$ B cells. Most B cells also
expressed CD23, which is associated with mature and activated B cells.[47] We validated these
populations in a publicly available human scRNAseq data set to show a significant ABC
enrichment in Alzheimer's disease. We found gender-specific and brain location-specific
differences in ABC populations, with caveats related to sample sizes that limited statistical power.
We also noted that CD11b$^+$ ABCs trended to be higher in females when isolated from the
cerebellum. CD11b$^+$ ABCs in healthy human donor blood exhibit migratory properties,[48] and
CD11b facilitates B cell migration by aiding in adhesion to endo- and sub-endothelium in the
vasculature via the intercellular adhesion molecule 1(ICAM1).[49] These peripheral B cells express
CD11b, regardless of chemokine presence and attraction, and also expressed CD27, a memory
B cell marker.[49] Future studies should include both CD27, CCR7, and CXCR5, the latter
receptors instrumental in homing for CNS-specific migration from parenchymal ectopic lymphoid
structures,[50] to confirm migratory versus CNS-localized progenitor B cell pools in aged humans.

In conclusion, this study is the first to identify a B cell resident lineage in the aged brain that
develops from early progenitors to ABCs. We also show that aged male mice have a
substantially increased baseline B cell inflammatory state in the brain over female counterparts.
Stroke induced clonal expansion of IgM$^+$ ABC/B1b and plasma cells that exhibit a phagocytic
phenotype. We also confirmed developing, mature and ABC B cell subsets in human
parenchyma. These findings have important implications for future work examining sex
differences, brain injuries, and neurodegenerative diseases in the aging brain. Stroke, traumatic
brain injury, and Alzheimer's disease have all had clinical trials with peripheral immunotherapies
that have shown a broad range of results in terms of efficacy and safety. It is possible that
detrimental – or perhaps even some beneficial – immune effects originate from CNS-localized,
self-renewing B cell pools sequestered from peripheral immunotherapies and thus may require
unique CNS targeting for therapeutic efficacy in diseases of aging.

Acknowledgments

Thank you to the brain donors and their supportive families for their invaluable contribution to this work. These studies are supported by grants from the National Institutes of Health to AMS: R01NS088555, RF1NS088555; AMM: F30NS141486; MKC: T32AG057461; TU: T32NS077889, 3RF1NS088555-07A1S1; KMC: T32NS077889; SJC: R01NS129668; UK-ADRC: P30 AG072946. Services and support for scRNAseq experiments were provided by the College of Arts & Sciences Imaging Center at the University of Kentucky. This work was supported by the UK Light Microscopy core facility (RRID: SCR_026405), Flow Cytometry & Immune Monitoring core (RRID: SCR_026358) and Biospecimen Procurement & Translational Pathology Shared Resource Facility (P30CA177558).

Methods

All experiments were performed in accordance with the University of Kentucky IACUC under accreditation of AAALAC. Aged (>18 months) male and female C57BL/6JN mice were obtained from the National Institute on Aging. Mice were socially housed and kept in a 14:10 light-dark cycle room with free access to food and water. Methods for tMCAo, behavioral testing, brain processing, B cell isolation, flow cytometry, immunohistochemistry, scRNAseq, scBCRseq, scRNAseq processing & data analysis, high dimensional weighted gene co-expression analysis, transcription factor regulatory network analysis in hdWGCNA, pseduotime trajectory analysis, external data processing, label transfer, and multicompartment aged B cell analysis, clonal analysis, Alpha Fold 3 structural prediction, and human rapid postmortem brain processing are included in the Extended Methods.

Data Availability:

Publicly available B cell sequencing data was obtained from GSE174834. Raw and Processed data from this study will be available upon publication or request.

Figure 1: B cell subsets demonstrate sex- and brain area-dependent expression patterns following ischemic stroke that impairs long-term motor recovery.

Flow cytometry in immune cells isolated from the left and right hemispheres and cerebellum (designated x axis) in uninjured mice and mice at 3-weeks following a transient middle cerebral artery occlusion (tMCAo; red symbols) show sex-specific interactions for aged females (circles; green bars) vs. males (squares; blue bars). This includes **a,** CD11c+ and **b,** CD11b+CD11c+ ABCs, with additional interactions for **c,** CD23+CCR7+ memory B cell subsets and **d,** plasma cells. **e,** tMCAo induced motor deficits in both aged males and females without an effect of sex. Of all B cell subsets, only **f,** plasma cells showed a negative correlation with functional recovery in both females (top graph) and males (bottom graph). **g,** To confirm expression of the canonical ABC transcription factor Tbet, immunostaining in B cells isolated from the post-stroke brain show high B220 expression (green; left panel) and more reduced Tbet expression (red, middle, right panels) with very large B cell subsets remaining with high Tbet expression.

Figure 2: scRNAseq of B cells in healthy aging and after ischemic stroke

a, Experiment design of B cell enriched cell suspension scRNAseq & scBCRseq isolated from
healthy aged (Female n = 4; Male n = 4) and 21-day post-stroke mouse brains (Female n = 4). **b,**
UMAP of 12,081 total CD45+ cells annotated via canonical marker genes. Hemopoietic stem
cells (HSCs; n = 656), monocytes (n = 180), plasma cells (n = 279), all B cells (n = 10,966). **c,**
Gene feature plots of Ms4a1 (Cd20) & Cd19 on all Cd45$^+$ cells confirm B cell lineage. **d,** Dotplot
of immature B cell markers across CD45$^+$ cells show all developmental stages to age-associated
B cell (ABC) populations all unique from the monocyte lineage. **e,** Schematic of B cell
development phenotypes. **f,** Number of immature B cell subsets in uninjured aged mouse
cerebellum and at 3wks after transient middle cerebral artery occlusion (tMCAo). Blue circle =
naïve for both males (left) and females (right); purple squares = tMCAo.

4
451

Figure 3: B cells develop into ABCs in the brain and other compartments

a, UMAP of 10,966 brain-resident B cells annotated via canonical marker genes and via an anchor-based label transfer from a published multi-compartment B cell scRNAseq dataset. Mitotic B cells (n = 509), Pro-B cells (n = 203), Pre-B cells (n = 2251), Immature B cells (n = 6141), Mature naïve B cells (n = 659), ABCs (n = 710), B1b cells (n = 493). **b,** Top 5 B cell population gene markers per identified B cell subset ranked by Log2FC. **c,** B cell UMAP split by treatment condition (left), with B cell population proportions split by treatment condition and

shown in distribution graphs (right). **d,** UMAP of 10,966 B cells colored Monocole3 pseudotime
with trajectories branch tree plotted. Root node labeled by white circled 1 (in dark purple area),
determined by identifying earliest branch node inside of *KI67*+ mitotic B cells. **e,** Violin nested
boxplot of pseudotime values across B cell phenotypes ordered increasingly by mean
pseudotime**. f,** UMAP of 1018 ABC/B1b cells colored by phenotype (ABCs, purple; B1b cells,
pink). **g,** Gene feature plot of *Tbx21* (Tbet), the canonical ABC transcription factor. **h,** ABC/B1b
cells split by treatment condition in both UMAP (left) and distribution bar graphs (right). **i,** Dotplot
of top 10 ABCs & B1b cell marker genes (ABCs, bottom row). **j,** UMAP of a multicompartment
integrated 29,806 B cells split across 4 compartments (Blood, Bone Marrow, Dura, Brain) and
annotated via identified marker genes to identify B cell subsets. **k,** Monocole3 pseudotime
trajectories of integrated atlas with root node selected from earliest tree node designated as #1 in
the dark purple mitotic cell population.
471

Figure 4: Gene and transcription factor regulator networks reveal shared inflammatory
signaling between uninjured male and post-stroke female B cells

a-b, Volcano plots of differentially expressed genes between treatment conditions (Log2FC >
0.25 or < -0.25; Padj < 0.05, min.pct > 0.1; Top 10 genes by Log2FC labeled). **c,** Venn diagram
of significant up- and downregulated genes shared between stroke female vs uninjured female
and uninjured male vs uninjured female. **d,** hdWGCNA Module expression score feature plots of
first 1000 genes in each module on 10,966 brain resident B cells. **e,** Top 5 significant Gene
Ontology (GO) Biological Process 2025 terms for each module via Enrichr. Dot size is Enrichr
enrichment score and colored by -Log10 adjusted P value. **f,** Top 5 significant Reactome
Pathway 2024 terms for each module via Enrichr. Dot size is Enrichr enrichment score and
colored by -Log10 adjusted P value. **g,** hdWGCNA differential module score heatmap (Log2FC >
0.2 or < -0.2, Padj < 0.05, Wilcox rank sum, pseudocount = 0.01; ** Padj < 0.01, *** Padj, <
0.001). **h-i,** Transcription Factor (TF) differential regulon scatter plot across treatment conditions
where predicted positive relationship regulons (promoting) plotted on the X-axis and predicted
negative relationship (suppressing) regulons plotted on the Y-axis. TFs are colored by module
association (See hdWGCNA methods for more detail). **j,** Dotplot of *Myb*, *Bach2*, *Myc*, *Cebpb*
across B cell phenotypes.

Figure 5: Stroke induces hyperexpansion of IgM⁺ ABC/B1b B cells and plasma cells

a, UMAP of B cell clusters overlayed with BCR density contour, confirming sequenced BCRs were present in the expected populations across groups. **b,** Clone size across the identified B cell subtypes. **c,** BCR clonal network analysis showing the network with plasma cells as the primary node (left) and the ABC/B1 cluster as the primary node (right). **d,** Clonal expansion status across groups. Injured females showed the highest proportion of hyperexpanded cells, followed by uninjured females. **e,** UMAP of B cells showing expansion status across. **f,** Shannon diversity indices show minimal difference in diversity across groups. **g,** IgH constant chain usage is large IgM across groups. h, variable gene usage of hyperexpanded clones in uninjured and post-stroke females. Uninjured males did not have any hyperexpanded clones. I, top expanded clone in each group depicted on UMAP of B cells. post-stroke female top clone is in the plasma cell cluster while uninjured males and females are ABC/B1b cluster. Structural predictions of binding regions are depicted. Red region is 0.45pt confidence, orange region is 0.40pt confidence.

Figure 6: ABCs are present in human rapid post-mortem post-stroke brain

a, Experimental overview for collection of rapid post-mortem human brain samples. **b,** Gating strategy for flow cytometry of human brain samples. Samples were first gated for time, single cells, and live cells (not shown). **c-e,** CD11b⁺ **(c)** and CD11c⁺ **(d)** ABCs and CD23⁺ mature B

cells **(e)** in human motor cortex and cerebellum expressed as percent of live CD45$^+$CD19$^+$ cells.
Females = green circles. Males = blue squares. Red depicts donors with a prior reported history
of stroke. **f-g,** correlation of age with percentage of CD11b and CD11c ABCs in cerebellar tissue
show male-only increase with age for CD11c$^+$ ABCs. **h,** Immunohistochemistry staining of post-
mortem human brain for MAP2 (green, neurons), B220 (red), CD11c (purple) identifies B220$^+$
CD11c$^+$ MAP2$^-$ B cells (arrow). **i,** UMAP of human B cells from GSE288856 of brain and
leptomeninges with proportion of cell types grouped by case diagnosis of Alzheimer's disease
(AD) vs. cognitively normal control (CTRL). Dotplot of B cell and ABC markers across B cell
subsets for the B cell-related genes *Cd19*, *MS4A1*, *Tbx21, ITGAX*, and *ITGAM*.

References

Winkelmeier, L. *et al.* Age and Functional Outcomes in Patients With Large Ischemic Stroke Receiving Endovascular Thrombectomy. *JAMA Netw Open* **7**, e2426007, (2024). PMID: 39133490.

Knoflach, M. *et al.* Functional recovery after ischemic stroke--a matter of age: data from the Austrian Stroke Unit Registry. *Neurology* **78**, 279-285, (2012). PMID: 22238419.

Ritzel, R. M. *et al.* Aging alters the immunological response to ischemic stroke. *Acta Neuropathol* **136**, 89-110, (2018). PMID: 29752550.

Tuohy, M. C., Hillman, E. M. C., Marshall, R. & Agalliu, D. The age-dependent immune response to ischemic stroke. *Curr Opin Neurobiol* **78**, 102670, (2023). PMID: 36586305.

Iadecola, C., Buckwalter, M. S. & Anrather, J. Immune responses to stroke: mechanisms, modulation, and therapeutic potential. *The Journal of Clinical Investigation* **130**, 2777-2788, (2020). PMID:

Iadecola, C. & Anrather, J. The immunology of stroke and dementia. *Immunity* **58**, 18-39, (2025). PMID: 39813992.

Malone, M. K. *et al.* The immunopathology of B lymphocytes during stroke-induced injury and repair. *Seminars in Immunopathology* (2022). PMID:

LeBien, T. W. & Tedder, T. F. B lymphocytes: how they develop and function. *Blood* **112**, 1570-1580, (2008). PMID: 18725575.

Morris, G. *et al.* Emerging role of innate B1 cells in the pathophysiology of autoimmune and neuroimmune diseases: Association with inflammation, oxidative and nitrosative stress and autoimmune responses. *Pharmacological Research* **148**, 104408, (2019). PMID:

Hao, Y., O'Neill, P., Naradikian, M. S., Scholz, J. L. & Cancro, M. P. A B-cell subset uniquely responsive to innate stimuli accumulates in aged mice. *Blood* **118**, 1294-1304, (2011). PMID: 21562046.

Cancro, M. P. Age-Associated B Cells. *Annual Review of Immunology* **38**, 315-340, (2020). PMID:

Korf, J. M. *et al.* CD11b(high) B Cells Increase after Stroke and Regulate Microglia. *J Immunol* **209**, 288-300, (2022). PMID: 35732342.

Doyle, K. P. & Buckwalter, M. S. Does B lymphocyte-mediated autoimmunity contribute to post-stroke dementia? *Brain Behav Immun* (2016). PMID: 27531189.

Doyle, K. P. *et al.* B-lymphocyte-mediated delayed cognitive impairment following stroke. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **35**, 2133-2145, (2015). PMID: 25653369.

Ortega, S. B. *et al.* Stroke induces a rapid adaptive autoimmune response to novel neuronal antigens. *Discov Med* **19**, 381-392, (2015). PMID: 26105701.

Ortega, S. B. *et al.* B cells migrate into remote brain areas and support neurogenesis and functional recovery after focal stroke in mice. *Proceedings of the National Academy of Sciences* **117**, 4983-4993, (2020). PMID:

Offner, H. *et al.* Experimental stroke induces massive, rapid activation of the peripheral immune system. *J Cereb Blood Flow Metab* **26**, 654-665, (2006). PMID: 16121126.

Ren, X. *et al.* Regulatory B cells limit CNS inflammation and neurologic deficits in murine experimental stroke. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **31**, 8556-8563, (2011). PMID: 21653859.

19 Prüss, H. *et al.* Evidence of Intrathecal Immunoglobulin Synthesis in Stroke: A
Cohort Study. *Archives of Neurology* **69**, 714-717, (2012). PMID:
20 Becker, K. J., Tanzi, P., Zierath, D. & Buckwalter, M. S. Antibodies to myelin basic
protein are associated with cognitive decline after stroke. *Journal of*
Neuroimmunology **295-296**, 9-11, (2016). PMID:
21 Förster, R. *et al.* CCR7 Coordinates the Primary Immune Response by
Establishing Functional Microenvironments in Secondary Lymphoid Organs. *Cell*
99, 23-33, (1999). PMID:
22 Wang, R. *et al.* A novel phenotype of B cells associated with enhanced phagocytic
capability and chemotactic function after ischemic stroke. *Neural Regeneration*
Research **18** (2023). PMID:
23 Brioschi, S. *et al.* Heterogeneity of meningeal B cells reveals a lymphopoietic
niche at the CNS borders. *Science* **373**, 408-+, (2021). PMID:
WOS:000679232100025.
24 Morgan, D. & Tergaonkar, V. Unraveling B cell trajectories at single cell resolution.
Trends Immunol **43**, 210-229, (2022). PMID: 35090788.
25 Cao, J. *et al.* The single-cell transcriptional landscape of mammalian
organogenesis. *Nature* **566**, 496-502, (2019). PMID: 30787437.
26 Morabito, S., Reese, F., Rahimzadeh, N., Miyoshi, E. & Swarup, V. hdWGCNA
identifies co-expression networks in high-dimensional transcriptomics data. *Cell*
Rep Methods **3**, 100498, (2023). PMID: 37426759.
27 Childs, J. E. *et al.* Relapse to cocaine seeking is regulated by medial habenula
NR4A2/NURR1 in mice. *Cell Rep* **43**, 113956, (2024). PMID: 38489267.
28 Vorontsov, I. E. *et al.* HOCOMOCO in 2024: a rebuild of the curated collection of
binding models for human and mouse transcription factors. *Nucleic Acids Res* **52**,
D154-d163, (2024). PMID: 37971293.
29 Wang, R. *et al.* A novel phenotype of B cells associated with enhanced phagocytic
capability and chemotactic function after ischemic stroke. *Neural Regen Res* **18**,
2413-2423, (2023). PMID: 37282471.
30 Piovesan, D. *et al.* c-Myb Regulates the T-Bet-Dependent Differentiation Program
in B Cells to Coordinate Antibody Responses. *Cell Rep* **19**, 461-470, (2017).
PMID: 28423310.
31 Pelissier, A., Luo, S., Stratigopoulou, M., Guikema, J. E. J. & Rodríguez Martínez,
M. Exploring the impact of clonal definition on B-cell diversity: implications for the
analysis of immune repertoires. *Front Immunol* **14**, 1123968, (2023). PMID:
37138881.
32 Schmitt, F. A. *et al.* Celebrating 40 years of the University of Kentucky Alzheimer's
Disease Research Center. *Alzheimers Dement* **21**, e70181, (2025). PMID:
40365904.
33 Schmitt, F. A. *et al.* University of Kentucky Sanders-Brown healthy brain aging
volunteers: donor characteristics, procedures and neuropathology. *Curr Alzheimer*
Res **9**, 724-733, (2012). PMID: 22471862.
34 Hobson, R. *et al.* Clonal CD8(+) T□cells populate the leptomeninges and
coordinate with immune cells in human degenerative brain diseases. *Nat Immunol*
27, 323-335, (2026). PMID: 41593242.
35 Oh, S. G., Noh, J., Jang, E. & Youn, J. The transcription repressor Bach2 is
required for maintaining the B-1 cell population by regulating self-renewal. *Front*
Immunol **16**, 1553089, (2025). PMID: 40170867.

36   Oh, S.-G., Noh, J., Jang, E. & Youn, J. The transcription repressor Bach2 is
619   required for maintaining the B-1 cell population by regulating self-renewal.
620   *Frontiers in Immunology* **Volume 16 - 2025** (2025). PMID:

621   37   Kim, K. *et al.* Therapeutic B-cell depletion reverses progression of Alzheimer's
622   disease. *Nature Communications* **12**, 2185, (2021). PMID:

623   38   McCulloch, L., Smith, C. J. & McColl, B. W. Adrenergic-mediated loss of splenic
624   marginal zone B cells contributes to infection susceptibility after stroke. *Nature*
625   *Communications* **8**, 15051, (2017). PMID:

626   39   Torres, V. O. *et al.* B cells upregulate NMDARs, respond to extracellular
627   glutamate, and express mature BDNF to protect the brain from ischemic injury.
628   *Neurobiology of Disease* **207**, 106819, (2025). PMID:

629   40   Cugurra, A. *et al.* Skull and vertebral bone marrow are myeloid cell reservoirs for
630   the meninges and CNS parenchyma. *Science* **373** (2021). PMID: 34083447.

631   41   Becker, K. J., Tanzi, P., Zierath, D. & Buckwalter, M. S. Antibodies to myelin basic
632   protein are associated with cognitive decline after stroke. *J Neuroimmunol* **295-**
633   **296**, 9-11, (2016). PMID: 27235342.

634   42   Phan, H. V. *et al.* High-throughput RNA sequencing of paraformaldehyde-fixed
635   single cells. *Nat Commun* **12**, 5636, (2021). PMID: 34561439.

636   43   Li, X. *et al.* Transcriptional and epigenetic decoding of the microglial aging
637   process. *Nat Aging* **3**, 1288-1311, (2023). PMID: 37697166.

638   44   Wootla, B., Denic, A., Warrington, A. E. & Rodriguez, M. A monoclonal natural
639   human IgM protects axons in the absence of remyelination. *J Neuroinflammation*
640   **13**, 94, (2016). PMID: 27126523.

641   45   Budeus, B., Kibler, A. & Kuppers, R. Human IgM-expressing memory B cells.
642   *Front Immunol* **14**, 1308378, (2023). PMID: 38143767.

643   46   Xie, G. *et al.* Age-Associated B Cells in Autoimmune Diseases: Pathogenesis and
644   Clinical Implications. *Clinical Reviews in Allergy & Immunology* **68**, 18, (2025).
645   PMID:

646   47   Herisson, F. *et al.* Direct vascular channels connect skull bone marrow and the
647   brain surface enabling myeloid cell migration. *Nat Neurosci* **21**, 1209-1217, (2018).
648   PMID: 30150661.

649   48   Kawai, K. *et al.* CD11b-mediated migratory property of peripheral blood B cells. *J*
650   *Allergy Clin Immunol* **116**, 192-197, (2005). PMID: 15990794.

651   49   Kawai, K. *et al.* CD11b-mediated migratory property of peripheral blood B cells.
652   *Journal of Allergy and Clinical Immunology* **116**, 192-197, (2005). PMID:

653   50   Weitbrecht, L. *et al.* CD4+ T cells promote delayed B cell responses in the
654   ischemic brain after experimental stroke. *Brain, Behavior, and Immunity* **91**, 601-
655   614, (2021). PMID:

656