1 **Proteome landscape of B-cell malignancies identifies mantle cell lymphoma**

2 **protein signature**

3 Samantha A. Swenson[1,2,3,*], C. Bea Winship[4,5], Kasidy K. Dobish[2,3,4,5], Karli J. Wittorf[2,3,6],
4 Henry Chun Hin Law[3,7], Julie M. Vose[3,8], Timothy Greiner[3,8], Michael R. Green[9,10,11],
5 Nicholas T. Woods[3,7], and Shannon M. Buckley[4,5,*]

6
[1]Department of Biochemistry and Molecular Biology, University of Nebraska Medical
Center, Omaha, NE, USA; [2]Department of Genetics, Cell Biology and Anatomy,
University of Nebraska Medical Center, Omaha, NE, USA; [3]Fred and Pamela Buffett
Cancer Center, University of Nebraska Medical Center, Omaha, NE, [4]Department of
Internal Medicine, Division of Hematology & Hematopoietic Malignancies, University of
Utah, Salt Lake City, UT USA; [5]Huntsman Cancer Institute, University of Utah, Salt Lake
City, UT USA; [6]Current Address: Creighton University School of Medicine, Omaha, NE,
USA; [7]Eppley Institute, University of Nebraska Medical Center, Omaha, NE, USA;
[8]Department of Pathology and Microbiology, University of Nebraska Medical Center,
Omaha, NE, USA; [9]Department of Lymphoma and Myeloma, Division of Cancer Medicine,
The University of Texas MD Anderson Cancer Center, Houston, TX, USA; [10]Department
of Genomic Medicine, University of Texas MD Anderson Cancer Center, Houston, TX,
USA; [11]Center for Cancer Epigenetics, University of Texas MD Anderson Cancer Center,
Houston, TX, USA.
21

22 *****Address Correspondence To:**

23 **Shannon M. Buckley, Ph.D.**
24 Email: Shannon.Buckley@hci.utah.edu
25
26 **Samantha A. Swenson, Ph.D.**
27 Email: samantha.swenson@unmc.edu
28

29 **CONFLICTS OF INTEREST**

30 The authors have no conflicts of interest related to this work.

ABSTRACT

Mantle cell lymphoma (MCL) is one of the deadliest forms of Non-Hodgkin's B-cell lymphoma. Typically, patients present with both overexpression of CyclinD1 and secondary mutations identified by genomic sequencing. Although MCL patients may initially respond to treatment, they eventually relapse and succumb to disease, highlighting the essential need to identify new targets for treatment. Here we performed proteomic profiling of healthy B cells and three different forms of B-cell malignancies, including MCL, to define the proteomic signature of MCL. We compared the proteome of each to MCL and identified 10 proteins that are specifically upregulated in MCL. Of these 10 proteins, seven of them show no transcriptional changes and have been overlooked by conventional RNA expression analysis. Further analysis of the proteomic signature reveals potential avenues for dual targeting in CAR T-cell therapy and provides guidance for personalized therapeutics based on protein expression.

STATEMENT OF SIGNIFICANCE

We present a resource defining the protein landscape of MCL, CLL, and FL as compared to healthy b cells identified utilizing quantitative proteomics from primary patient samples. Applied to MCL, our results identify 10 proteins specifically upregulated in MCL that may prove to be therapeutic targets to treat the disease.

INTRODUCTION

Malignancies affecting the hematopoietic system make up approximately 8-10% of all human cancers.[1] The two most common adult hematopoietic malignancies in western countries are chronic lymphocytic leukemia (CLL) and follicular lymphoma (FL). Both are B-cell malignancies with incidences of 25-30% of all adult leukemias and 20-40% of all Non-Hodgkin's Lymphomas (NHL), respectively.[2] Current therapies have improved treatment options and overall survival (OS). CLL has an ~87% 5-year OS, in part due to the BTK inhibitor Ibrutinib being approved by the FDA in 2016 for frontline treatment.[3] Likewise, FL has an ~90% 5-year OS associated with the addition of the monoclonal antibody rituximab being used in conjunction with chemotherapeutics.[4]

Unfortunately, not all B-cell malignancies have similar survival rates. In the case of mantle cell lymphoma (MCL), an aggressive NHL that accounts for ~4-6% percent of all lymphoma cases in the United States, patients have only an ~40% 5-year survival.[2] Besides the standard high dose chemotherapy regimen, limited treatments are available, resulting in poor outcomes.[2] Recently, chimeric antigen receptor T-cell (CAR-T) therapy targeting the B-cell marker CD19 has been approved by the FDA for treatment in relapse/refractory MCL.[5] Although this provides a new therapeutic option for MCL patients, it is not considered a universal cure and many patients still relapse.[5] Additionally, the proteasome inhibitor Bortezomib was approved for MCL patients with relapse or refractory disease in 2013; however, discrepancies between survival of these patients show a significant need for new first-line therapeutics.[6]

This study aims to garner a better understanding of MCL pathogenesis and create a resource that can be used for the identification of potential novel therapeutic targets to

improve patient prognosis. Currently, most research has focused on gene expression

profiling (GEP) of patients to identify molecular predictors of disease; however, these

studies are not extensively validated at the protein level where targeted therapies take

effect. Additionally, current proteomic studies are heavily reliant on cell lines which cannot

accurately mimic patient tumors.[7, 8] When patient samples have been utilized, the

comparison has only been made to healthy donor (HD) and not between disease types.[9,

10] The one instance where disease types have been compared (MCL to CLL), only

transcriptomics was utilized.[11] To address this gap in knowledge, we utilized 35 primary

patient samples to identify an MCL-specific proteomic signature by comparing MCL to

CLL, FL, and HD B cells by quantitative mass spectrometry. The minimal overlap

observed between proteomic and GEP analyses emphasizes the importance of

developing such a resource. Of particular interest are proteins we have identified as

differentially expressed on the plasma membrane, as these may be ideal targets for CAR-

T therapy. We also identified patient-specific variations in the protein abundance of

proteasomal subunits suggesting that addition of Bortezomib to the treatment regimen

may benefit some patients, but not others. Finally, our results identify 10 proteins

specifically upregulated in MCL. Of these 10 proteins, seven are unchanged at the mRNA

level, and would therefore not be identified by GEP. Importantly, three of these proteins

have never been associated with any lymphoid malignancy, making them a novel finding.

107
RESULTS

Proteomics signatures of MCL, CLL, and FL

To identify the protein landscape of MCL, CLL, and FL, we performed quantitative

tandem mass tag (TMT) mass spectrometry on B cells from 5 HD, 7 CLL, 7 FL, and 16

MCL patients (**Supplemental Table S1**). We identified 8,027 proteins in total with 1,502

proteins present in all batches (**Figure 1A and B**, **Supplemental Table S2**). Principal

component analysis (PCA) shows that HD and individual malignancies cluster among

themselves and lymphomas capable of forming solid tumors (MCL and FL) cluster

similarly while CLL, which remains predominantly in the peripheral blood, clusters

separately (**Figure 1C**).

Heat map of all disease protein abundances shows that samples cluster by

disease sub-type and form five distinct protein clusters associated with various Gene

Ontology (GO) pathways (**Figure 1D, Supplemental Tables S3-S8**). The most distinct

clusters being one, four, and five. Cluster one proteins, upregulated in MCL and FL,

primarily contain proteins associated with the immune response, but also contain proteins

associated with cell adhesion and angiogenesis. Because these tumors can infiltrate

secondary lymphoid tissues to form solid tumors, targeting these proteins may be

significant for slowing disease progression. Cluster four proteins, upregulated in CLL, are

associated with mitochondrial health and function. It has been reported that CLL cells are

heavily reliant on mitochondrial metabolism.[12] Therefore, proteins within this cluster may

prove to be vulnerabilities within the disease ideal for targeting. Cluster five proteins are

downregulated in FL and MCL and are associated with RNA processing. We have

previously reported that B cells lacking the catalytic domain of the E3 ubiquitin ligase

component *n*-recognin 5 (UBR5) have splicing defects.[13] Given that UBR5 is mutated in

~18% of MCL patients, mutations to UBR5's catalytic domain are specific to MCL tumors,

and UBR5 is the second most mutated gene in our MCL patient cohort, present in 31%

of samples, proteins in this cluster may provide insight into how UBR5 mutations affect

MCL pathogenesis (**Supplemental Figure S1A**).[14, 15]

As compared to HD, there are 131 significantly dysregulated proteins (104

upregulated, 27 downregulated) in MCL, 101 in CLL (70 up, 31 down), and 152 in FL (131

up, 21 down) (**Figure 1E, Supplemental Table S9**). Of the dysregulated proteins, only 6

are consistently upregulated and 3 are consistently downregulated across all three

malignancies. GO analysis of dysregulated proteins in each malignancy identifies unique

and similar pathways between each lymphoma type (**Figure 1F**). As discussed above,

both MCL and FL have upregulation of proteins associated with cell-cell adhesion,

angiogenesis, and inflammatory response. Proteins of interest include intercellular

adhesion molecule 1 (ICAM1) and transforming growth factor β-induced protein (TGFBI).

(**Figure 1G, orange**). High ICAM1 surface levels in FL and MCL, but low levels in CLL is

linked to unfavorable patient prognosis.[16] Although it has not been studied in MCL,

transcriptional silencing of TGFBI is considered to be a potential mechanism of B-cell

transformation resulting in FL.[17] These data are in line with our proteomics data, validating

our findings.

Upregulated protein pathways unique to MCL are actin filament organization and

positive regulation of cell migration. They include SH3-domain binding protein 1

(SH3BP1) and myristoylated alanine-rich C-kinase substrate (MARCKS) (**Figure 1G,**

purple). While upregulation of SH3BP1 is a new finding in lymphoma, it's upregulation is

associated with poor prognosis in acute myeloid leukemia (AML) patients.[18] *MARCKS* is

upregulated in MCL and has an important role in cancer spread, invasion, metastasis,

and Bortezomib resistance in multiple myeloma (MM).[11]

FL has one unique upregulated pathway, signal transduction, that is associated

with 19 upregulated genes. Of these genes, four of them are specific to FL: Rho GTPase-

activating protein 1 (ARHGAP1), complement component c3 (C3), Ras-related protein

Rap-1b (RAP1B), and guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit Beta-2

(GNB2) (**Figure 1G, pink**). The most significant of these proteins, ARHGAP1, has not

previously been associated with lymphoma; however, one study found that it is

alternatively spliced in a TRAF6 dependent manner in myelodysplastic syndrome.[19]

Upregulated pathways in CLL are very different from MCL and FL. Some proteins

associated with these pathways include glutathione peroxidase 1 (GPX1) and thiol S-

methyltransferase (METTL7A) (**Figure 1G, light blue**). High GPX1 expression correlates

with poor prognosis in lymphoma patients.[20] METTL7A has not previously been

associated with lymphoma but overexpression is associated with enhanced human bone

marrow stem cell viability.[21]

Pathways downregulated in each malignancy are highly unique. Some potentially

relevant downregulated proteins from each malignancy include methyl-CpG-binding

protein 2 (MECP2) in MCL, tyrosine-protein phosphatase non-receptor type 6 protein

(PTPN6) in FL, and cluster of differentiation 22 (CD22) in CLL (**Figure 1G, light green**).

MECP2 interacts with the SIN3a/histone deacetylase 1/2 corepressor complex to silence

BIM, a proapoptotic member of the BLC-2 family.[22] PTPN6 is an important repressor of

STAT3 activation, a transcription factor constitutively activated in a variety of cancers,

including lymphoma, and is considered a tumor suppressor in AML and MM.[23]

Overall, this data identifies similarities and unique aspects of three different types

of lymphoid malignancies, which have been previously unidentified or underexplored.

These pathways and proteins may prove to be prognostic markers for their respective

diseases and highlight potential vulnerabilities for possible treatment strategies.

Identification of cell surfaceome

Recent success of CAR-T therapy in B-cell malignances targeting specific cell

surface receptors reinforces the importance of the cell surfaceome. Of the 1,502 proteins

identified in our MS analysis, 120 of them localize to the plasma membrane (**Figure 2A**).

One of the defining characteristics of CLL is that it is CD5$^+$, CD19$^+$, and CD23$^+$ with weak

amounts of CD22, CD79b, and surface immunoglobulin and is FMC7$^-$.[24] FL cells are

CD19$^+$, CD20$^+$, CD10$^+$, and BCL-6$^+$, but are CD5$^-$ and CD23$^-$.[25] MCL cells are

characterized by B-cell surface markers CD19$^+$, CD20$^+$, CD22$^+$, CD79a$^+$, CD5$^+$, FMC-7$^+$,

and CD10$^-$.[26] Cell surface protein pattern of CD5, CD19 and CD22 was consistent with

diagnostic markers (**Figure 2B**). Because CAR-T therapy is rapidly becoming the

treatment of choice for patients that stop responding to other therapies, and the high

incidence of relapse in lymphoid malignancies – MCL has a 100% relapse rate following

initial treatments – plasma membrane proteins may prove to be new targets for CAR-T

treatment. Of these 120 proteins, 48 of them are either significantly up- or downregulated

in each of the three malignancies individually as compared to HD B cells (**Figure 2C**). Of

these 48 proteins, 11 of them are specifically upregulated in MCL (**red text**), 8 in FL

(**green text**), and 7 in CLL (**dark blue text**). 10 of these proteins overlap between MCL

and FL (**orange text**), 1 overlaps between CLL and FL (**light blue text**), and two are

upregulated between all three malignancies (**purple text**). We measured abundance of

cell surface proteins identified in our analysis in MCL and CLL cell lines – Jeko1 and

Granta519 (MCL) and HG3 (CLL) – and compared them to peripheral blood in HD (**Figure**

2D). We found that in these cell lines, some proteins, such as LGALS9, CD81, and ICAM1

matched our proteomic findings, while others such as CD5, CD22, and B2M were more

variable. These differences are likely due to differences between the *in vitro* and *in vivo*

microenvironments, emphasizing the importance of utilizing patient samples when

identifying therapeutic targets of interest.

Current CAR-T therapy in lymphoma targets CD19[+] cells.[5] It has been suggested

that low CD19 may diminish efficacy of CAR-T treatment suggesting dual receptor

targeting could improve outcomes.[27] We found that CD81 was increased in MCL, but not

CLL. When we stratified MCL patients based on CD19 expression, we found that CD19

and CD81 had an inverse relationship to one another, but no significant difference in

CD22 expression (**Figure 2E and 2F**). This suggests that using CD81 as a CAR-T therapy

target in combination with CD19 may improve patient outcomes. Together, these findings

provide additional potential receptors for cellular therapy.

217
Genetic heterogeneity of MCL corresponds to proteomic heterogeneity

It is thought that genetic heterogeneity is one of the reasons for the poor survival

and limited treatments in MCL.[14] The 16 MCL patients included in our study had typical

age and male to female distribution. While there were mutations in 88 different genes,

with patients having up to 15 mutations, we saw the expected frequency in MCL for TP53

and ATM mutations (**Supplemental Figure S1A**). PCA analysis revealed no distinct

clustering in the MCL samples (**Figure 3A**). Overall mutational profile or number of

mutations again revealed no specific clustering in the MCL cohort (**Supplemental Figure**

S1B).

Besides standard high dose of cytarabine regime, limited treatments are available

for patients with MCL, resulting in poor survival rates and outcomes.[2] The proteasome

inhibitor Bortezomib has recently been approved for patients with relapse or refractory

disease, however, there has been mixed results.[6] It has been reported that expression of

the proteasome subunits correlates with proteasome activity and increased sensitivity to

proteosome inhibition.[28] Clustering MCL, CLL, and FL patients based on protein

expression of proteasome components shows diverse expression profiles among the

MCL cohort (**Figure 3B**). A number of patients clustered with HD samples with low

expression of proteasome components (MCL-006, MCL-003, and MCL-038) suggesting

that the patients would have decreased sensitivity to Bortezomib. These findings suggest

that for a heterogenous disease like MCL, precision targeted therapeutics could benefit

patients and highlight the importance of using proteomics to identify new therapies that

may improve patient outcomes.

Novel post transcriptionally regulated proteins in MCL

Because MCL has the lowest 5-year survival and a high need for new therapeutics,

we compared protein expression of MCL patients to both HD B cells and to the other B-

cell malignancies. While there were no proteins consistently downregulated in MCL as

compared to all groups, comparison of the upregulated proteins found that there are 10

proteins specifically upregulated in MCL (**Figure 4A and B, Supplemental Table S10**).

These proteins are high mobility group box 3 (HMGB3), EH domain containing 1 (EHD1),

MARCKS, galectin 9 (LGALS9), PDZ and LIM domain protein 1 (PDLIM1), drebrin 1

(DBN1), glutathione S-transferase Pi (GSTP1), dimethylarginine dimethylaminohydrolase

2 (DDAH2), stathmin 1 (STMN1), and four and a half LIM domains protein 1 (FHL1).

Importantly, all 10 of these proteins had 3 or more unique peptides identified.

(**Supplemental Table S2, Supplemental Figure S2)**. Of these, seven have never been

associated with MCL and three have never been associated with any form of lymphoid

malignancy (**Supplemental Figure S2**). Using previously published transcriptional data

that includes these same MCL patients, we compared mRNA expression for these 10

proteins.[29] We found that seven of them are unchanged transcriptionally (**Figure 4C**). The

RNA expression level of these 10 genes shows no significant difference in patient

survival, which is not surprising given that the transcripts are unchanged (**Supplemental**

Figure S3).

To validate the upregulation of these proteins in MCL, we performed western blots

for three these 10 proteins (STMN1, GSTP1, and HMGB3) in both primary patients and

patient-derived cell lines including three MCL cell lines: Mino, Jeko1, and Granta519; two

CLL cell lines: Mec1 and HG3; and two other lymphoma cell lines: REH, a B-cell acute

lymphoblastic leukemia (B-ALL) cell line; and Ramos, a Burkitt's lymphoma cell line

(**Figure 4D and E)**. Our results in both models show these proteins are increased in MCL

as compared to HD B cells and to samples from other lymphoma types.

These findings identify novel proteins that are differentially expressed at the protein

level that have previously not been identified in classic transcriptional profiling, suggesting

potential novel targets for drug discovery or new translational avenues for PROTAC-

based drug development tailored to highly expressed proteins in MCL.

271
DISCUSSION

Historically, therapeutic targets for cancer treatments have been identified using

GEP. While these studies have identified many relevant targets, many of them have not

been confirmed at the proteomic level. Additionally, most of these studies – including

those that examine the changes to the proteome – examine only one type of cancer and

compare that cancer only to HD samples. Furthermore, these analyses are heavily reliant

on cell lines and when patient samples are used, the number of patients included is very

small.

To gain a better understanding of the differences between hematological

malignancies, we have taken samples from patients suffering from three different B-cell

malignancies and utilized quantitative proteomics to identify protein signatures of each

malignancy as compared to both healthy B cells and the other disease types. With the

identification of over 8,000 proteins across all samples, this study provides an in-depth

analysis of the proteomes of each malignancy type and utilizes the findings in MCL to

validate the results. Overall, the study identifies proteins that are specifically dysregulated

in MCL that may prove to be therapeutic targets for MCL therapies in the future.

Given the ever-increasing prevalence of CAR-T therapy in the treatment of B-cell

malignancies, our study identified a significant number of surface proteins that may prove

to be novel CAR-T targets. We also found that patients may benefit from proteomic

analysis of proteasome subunits prior to Bortezomib-based treatments. One of the most

striking findings of this study is the differences between the proteomic signature and the

previously identified GEPs of these patients. Because most MCL patients exhibit a

heterogenous mixture of genomic mutations, it is difficult to treat the disease, and this

lends to the high mortality rate of these patients. We had expected to see that patients

who displayed alterations in genes frequently mutated in MCL such as ATM, TP53, and

UBR5 would cluster together and show similar changes in their proteome. However, we

found that the number and type of mutations present in these patients had little to no

effect on the proteins that are either up or downregulated. Furthermore, the finding that

only three of the 10 proteins specifically upregulated in MCL also exhibited increases in

mRNA reinforces the importance of examining the patient's proteome in addition to their

genome to identify therapeutic targets for treatment.

The presence of MARCKS and EHD1 in our study, two proteins whose

corresponding gene expression has previously been reported to be upregulated in MCL

as compared to CLL and in NHL as a whole, respectively, validates these findings.[11, 30] In

addition to MARCKS and EHD1, only one of the other 10 identified proteins had a

transcriptional change, DBN1. It has previously been identified that DBN1 is a direct target

of SOX11, a protein known to be aberrantly expressed in MCL.[31] The role that these three

proteins play in actin cytoskeleton formation, membrane formation, and the formation of

cellular projections suggest that they may be important for disease progression and

infiltration into secondary lymphoid tissues. Although it does not have a transcriptional

change present like the other three, identification of GSTP1 as overexpressed at the

proteomic level also validates the study. GSTP1 has been shown to be chronically

overexpressed in MCL and may provide resistance to drug therapies as its inhibition

increases chemotherapy sensitivity in MCL patients.[32] Because MCL patients chronically

relapse following current drug regimens, targeting GSTP1 in conjunction with these

therapies may prove to be an effective strategy to improve patient outcomes.

The other six proteins specifically upregulated in MCL are also transcriptionally

unchanged, indicating that they would have remained unidentified by typical patient

screening methods. These six proteins have never been associated with MCL previously.

STMN1 is known to be overexpressed in other types of leukemias and lymphomas.[33]

LGALS9 has been shown to be upregulated on the surface of obese B-ALL patients, and

targeting this protein was reported to promote cell death, disrupt cell cycle progression,

and prolong survival in B-ALL patients.[34] FHL1 stabilization leads to Ibrutinib resistance

in diffuse large B-cell lymphoma.[35] HMGB3, PDLIM1, and DDAH2 have never been

associated with any form of lymphoma previously, making them of particular interest to

this study.

In conclusion, this study highlights the importance of utilizing proteomics when

determining the treatment strategy for lymphoma patients. This data can be used as a

resource for identifying novel treatment strategies not only in MCL, but in other B-cell

malignancies. It identifies several new potential CAR-T therapy targets as well as novel

proteins of interest for drug discovery or PROTAC-based drug development, highlighting

the importance of the work.

334
METHODS

Patient Samples

The study was approved by the institutional review board of the University of Nebraska

Medical Center (UNMC) (203-15-EP) and performed in accordance with the Declaration

of Helsinki. Matched gene expression microarray data are available through the Gene

Expression Omnibus, accession GSE132929.[14, 29] Samples obtained from the UNMC

Lymphoma Tissue Bank were prioritized for inclusion in this study if they had previously

undergone pathology review, been interrogated by Affymetrix U133 Plus 2.0 gene

expression microarrays, and exome sequenced.[9,28] Patient samples are at diagnosis, and

annotated for gender, age at diagnosis, OS, and mutational profile (**Supplemental Table**

S1). Protein was isolated from 16 MCL, 7 FL, and 7 CLL patients. For the 5 HD patient

samples, either adult CD19$^+$ B cells were purchased from Stem Cell Technologies or

magnetic selected B220$^+$ B cells were isolated from tonsils. All samples were de-identified

and accompanied by the diagnosis from the medical records, plus overall survival and

patient mutations. Mutation and CNA data are publicly viewable through cBioPortal:

https://www.cbioportal.org/study/summary ?id=mbn_mdacc_2013.[9]

Mass Spectrometry

Solid tumors from MCL and FL were ground to powder in liquid nitrogen. Samples were

prepared for TMT labelling, as previously described.[13] Samples were prepared in five

separate batches each containing a combined sample control (4μg from each sample)

and a mixture of HD, CLL, FL, and MCL samples. Protein identification was performed

using proteome discoverer software version 2.4 (ThermoFisher, Waltham, MA) by

searching MS/MS data against the UniProt human protein database downloaded

February 2022. Data was normalized to the average abundance of the batch control for

each of the five batches. Analyses were performed on the ratio of the abundance of

normalized protein in batch control to individual patient sample.   Data are available via

ProteomeXchange with identifier PXD038937.

363
364 **Flow Cytometry**

365 Cells were stained for 1 h on ice with indicated antibodies in 3% FBS in PBS. Data was

366 then analyzed using FlowJo software (Treestar, Ashland, OR). Antibody information is

367 available in supplementary materials and methods.

Cell Culture

Jeko-1, Mino, REH, and Ramos cells were purchased from ATCC (Manassa, VA) and Granta519, HG-3, and MEC-1 cells were purchased from DSMZ (Leibniz Institute, Germany). Cell lines were cultured following company specifications. All cells were confirmed mycoplasma negative monthly while in culture.

Western Blot

Samples were prepared and analyzed as previously described.[13] Antibody information is available in supplemental materials and methods.

Statistical analysis

All experiments were performed in triplicate unless noted. Proteins with a fold change ≥ 1.5 were considered as upregulated, ≤ 0.5 as downregulated, and with a p-value ≤ 0.05 as significant. Statistical analyses were performed using unpaired two-tailed Student's t-test assuming experimental samples of unequal variance (with Welch's correction). Error bars depict the standard deviation ± the mean. PCA plots were generated using ClustVis online tool (https://biit.cs.ut.ee/clustvis/). Heat maps were completed using Morpheus online tool (https://software.broadinstitute.org/morpheus), using hierarchical clustering and $\log_2$ transformation of protein abundance. Volcano plots were made using GraphPad Prism v10.0.2 (GraphPad, La Jolla, CA). Gene ontology (GO) pathway analysis was performed using DAVID Bioinformatics Database v6.8 (https://davidbioinformatics.nih.gov/home.jsp). Outputs were first sorted by count, and then by p-value. All outputs with ≥ 5 associated genes and a p-value ≤ 0.05 are included for reference in supplementary tables. Graphs were made using the top 5 hits for both count and p-value. GO terms with similar names and gene were only listed once under

394 the first instance. For survival curve analyses, the p-value was calculated using a Log-

395 rank (Mantel-cox) test.

397 **AUTHOR CONTRIBUTIONS**

403
ACKNOWLEDGEMENTS

We would like to thank the University of Utah Flow Cytometry Shared Resource

Laboratory, and UNMC Mass Spectrometry and Proteomics Core Facility for expert

assistance. The Nebraska Lymphoma Tissue Bank and core facilities are

administrated through the Office of the Vice Chancellor for Research and supported by

state funds from the Nebraska Research Initiative (NRI) and The Fred and Pamela Buffett

Cancer Center's National Cancer Institute Cancer Support Grant (P30CA036727).

S.M.B. is supported by the National Institutes of Health P20GM121316, R37CA262635,

and R01AI53090. This publication was supported by the Huntsman Cancer Institute at

the University of Utah, supported by the National Cancer Institute of the National Institute

of Health under award number P30CA042014.

FIGURE LEGENDS

Figure 1: Quantitative proteomics identifies protein landscapes of lymphoid

neoplasms as compared to healthy B cells

A. Schematic of sample processing workflow of protein from either healthy donor (HD) B

cells or lymphoid tumors from three different B-cell malignancies: mantle cell lymphoma

(MCL), chronic lymphoblastic leukemia (CLL), and follicular lymphoma (FL). Combined

protein from all samples were processed the same as individual samples and used as a

control in each run. **B.** Filtering parameters of LC-MS/MS outputs from the five different

MS batches. 24,794 peptides corresponding to 8,027 proteins were filtered to only include

those with an abundance output (5,114). Analysis was performed on proteins identified in

all batches (1,502). **C.** PCA plot using ClustVis software of normalized abundances for all

samples. **D.** Heat map of the Log_2 value of normalized abundances for all samples.

Hierarchical clustering was performed using Euclidian distance for both rows and

columns. **E.** Venn Diagram of common upregulated ($\geq$1.5-fold) (above) and

downregulated ($\leq$0.5-fold) (below) proteins between MCL, FL, and CLL all as compared

to HD samples. **F.** GO terms for upregulated ($\geq$1.5-fold) and downregulated ($\leq$0.5-fold)

proteins for each cancer as compared to HD. **G.** Volcano plots of the Log_2 fold change

for each cancer as compared to HD showing significant (p$\leq$0.05) upregulated ($\geq$1.5-fold)

and downregulated ($\leq$0.5-fold) proteins. Proteins upregulated in both MCL and FL

depicted in orange, proteins upregulated in MCL in purple, proteins upregulated in FL in

pink, proteins upregulated in CLL in light blue, and downregulated proteins of interest in

light green.

438
Figure 2: Cell surfaceome identifies potential CAR-T cell therapy targets in MCL

A. Pie chart of localization of 1,502 identified proteins highlighting 120 plasma membrane

proteins. **B.** Relative protein abundance of cell surface markers associated with B-

lymphoid malignancies (CD5, CD19, CD22). **C.** Heat map of plasma membrane proteins

that are differentially expressed in at least one of the three different B lymphoid

malignancies as compared to HD. **D.** Cell surface expression by flow cytometry in normal

human peripheral blood (PB), and B-Lymphoid derived cell lines Jeko1 (MCL), Granta519

(MCL), and HG3 (CLL). **E.** Relative abundance of CD81 and **F.** CD22 in CD19 high and

low expression patients. ns = not significant, * = p-value ≤ 0.05, ** = p-value ≤ 0.01.

Figure 3: Analysis of proteasome components suggest precision targeted

therapeutics could improve patient outcomes

A. PCA plot of MCL patient samples with clustering determined by ClustVis software. **B.**

Heatmap of proteasome subunits in all patients clustered by Pearson's Correlation.

453
454
Figure 4: Identification of 10 post-transcriptionally regulated proteins specifically

upregulated in MCL

A. Venn Diagram of proteins significantly (p≤0.05) upregulated (≥1.5-fold) (left) and

downregulated (≤0.5-fold) (right) in MCL samples as compared to HD, FL, and CLL

samples. **B.** Violin plots of relative protein abundance for 10 proteins specifically

upregulated in MCL as compared to HD, CLL, and FL. **C.** Violin plots of relative gene

expression as determined by RNA microarray of the 10 proteins from A in MCL as

compared to HD. **D.** Western blot confirmation in select MCL, CLL, and FL primary patient

samples and **E**. MCL, CLL, B-cell Acute Lymphoblastic Leukemia (B-ALL), and Burkitt's

Lymphoma (BL) patient derived cell lines as compared to HD. * = p-value ≤ 0.05, ** = p-

value ≤ 0.01, *** = p-value ≤ 0.001, **** = p-value ≤ 0.0001.

466
REFERENCES

1.   Mak TW, Saunders ME, and Jett BD. Primer to The immune response. 2nd
edition / ed. Amsterdam ; Boston: Elsevier/AP Cell, AP Cell is an imprint of Elsevier;
2014. xxvii, 674 pages p.

2.   Swerdlow SH, Campo E, Pileri SA, Harris NL, Stein H, Siebert R, et al. The 2016
revision of the World Health Organization classification of lymphoid neoplasms. Blood.
2016;127(20):2375-90. Epub 20160315. doi: 10.1182/blood-2016-01-643569. PubMed
PMID: 26980727; PMCID: PMC4874220.

3.   Burger JA, Barr PM, Robak T, Owen C, Ghia P, Tedeschi A, et al. Long-term
efficacy and safety of first-line ibrutinib treatment for patients with CLL/SLL: 5 years of
follow-up from the phase 3 RESONATE-2 study. Leukemia. 2020;34(3):787-98. Epub
20191018. doi: 10.1038/s41375-019-0602-x. PubMed PMID: 31628428; PMCID:
PMC7214263.

4.   Ghielmini M, Schmitz SF, Cogliatti SB, Pichert G, Hummerjohann J, Waltzer U, et
al. Prolonged treatment with rituximab in patients with follicular lymphoma significantly
increases event-free survival and response duration compared with the standard weekly
x 4 schedule. Blood. 2004;103(12):4416-23. Epub 20040219. doi: 10.1182/blood-2003-
10-3411. PubMed PMID: 14976046.

5.   Wang M, Munoz J, Goy A, Locke FL, Jacobson CA, Hill BT, et al. KTE-X19 CAR
T-Cell Therapy in Relapsed or Refractory Mantle-Cell Lymphoma. N Engl J Med.
2020;382(14):1331-42. doi: 10.1056/NEJMoa1914347. PubMed PMID: 32242358;
PMCID: PMC7731441.

6.   Schieber M, Gordon LI, and Karmali R. Current overview and treatment of mantle
cell lymphoma. F1000Res. 2018;7. Epub 20180725. doi:
10.12688/f1000research.14122.1. PubMed PMID: 30109020; PMCID: PMC6069726.

7.   Psatha K, Kollipara L, Voutyraki C, Divanach P, Sickmann A, Rassidakis GZ, et
al. Deciphering lymphoma pathogenesis via state-of-the-art mass spectrometry-based
quantitative proteomics. J Chromatogr B Analyt Technol Biomed Life Sci. 2017;1047:2-
14. Epub 20161115. doi: 10.1016/j.jchromb.2016.11.005. PubMed PMID: 27979587.

8.   Boyd RS, Dyer MJ, and Cain K. Proteomic analysis of B-cell malignancies. J
Proteomics. 2010;73(10):1804-22. Epub 20100324. doi: 10.1016/j.jprot.2010.03.010.
PubMed PMID: 20346427.

9.   Ghobrial IM, McCormick DJ, Kaufmann SH, Leontovich AA, Loegering DA, Dai
NT, et al. Proteomic analysis of mantle-cell lymphoma by protein microarray. Blood.

2005;105(9):3722-30. Epub 20050113. doi: 10.1182/blood-2004-10-3999. PubMed PMID: 15650054; PMCID: PMC1895014.

10.   Johnston HE, Carter MJ, Larrayoz M, Clarke J, Garbis SD, Oscier D, et al. Proteomics Profiling of CLL Versus Healthy B-cells Identifies Putative Therapeutic Targets and a Subtype-independent Signature of Spliceosome Dysregulation. Mol Cell Proteomics. 2018;17(4):776-91. Epub 20180124. doi: 10.1074/mcp.RA117.000539. PubMed PMID: 29367434; PMCID: PMC5880099.

11.   Vargova J, Vargova K, Dusilkova N, Kulvait V, Pospisil V, Zavadil J, et al. Differential expression, localization and activity of MARCKS between mantle cell lymphoma and chronic lymphocytic leukemia. Blood Cancer J. 2016;6(9):e475. Epub 20160923. doi: 10.1038/bcj.2016.80. PubMed PMID: 27662204; PMCID: PMC5056972.

12.   Jitschin R, Hofmann AD, Bruns H, Giessl A, Bricks J, Berger J, et al. Mitochondrial metabolism contributes to oxidative stress and reveals therapeutic targets in chronic lymphocytic leukemia. Blood. 2014;123(17):2663-72. Epub 20140219. doi: 10.1182/blood-2013-10-532200. PubMed PMID: 24553174.

13.   Swenson SA, Gilbreath TJ, Vahle H, Hynes-Smith RW, Graham JH, Law HC, et al. UBR5 HECT domain mutations identified in mantle cell lymphoma control maturation of B cells. Blood. 2020;136(3):299-312. doi: 10.1182/blood.2019002102. PubMed PMID: 32325489; PMCID: PMC7365918.

14.   Ma MCJ, Tadros S, Bouska A, Heavican T, Yang H, Deng Q, et al. Subtype-specific and co-occurring genetic alterations in B-cell non-Hodgkin lymphoma. Haematologica. 2022;107(3):690-701. Epub 20220301. doi: 10.3324/haematol.2020.274258. PubMed PMID: 33792219; PMCID: PMC8883549.

15.   Meissner B, Kridel R, Lim RS, Rogic S, Tse K, Scott DW, et al. The E3 ubiquitin ligase UBR5 is recurrently mutated in mantle cell lymphoma. Blood. 2013;121(16):3161-4. Epub 20130213. doi: 10.1182/blood-2013-01-478834. PubMed PMID: 23407552.

16.   Wang W, Li Y, Ali H, Zhao L, Mei D, Hu W, et al. Aberrant expression of CD54 detected by flow cytometry is a characteristic of B-lymphoma cells in bone marrow specimens. BMC Cancer. 2021;21(1):1315. Epub 20211208. doi: 10.1186/s12885-021-09061-3. PubMed PMID: 34879826; PMCID: PMC8653582.

17.   Manara F, Jay A, Odongo GA, Mure F, Maroui MA, Diederichs A, et al. Epigenetic Alteration of the Cancer-Related Gene TGFBI in B Cells Infected with Epstein-Barr Virus and Exposed to Aflatoxin B1: Potential Role in Burkitt Lymphoma Development. Cancers (Basel). 2022;14(5). Epub 20220302. doi: 10.3390/cancers14051284. PubMed PMID: 35267594; PMCID: PMC8909323.

18.   Yang L, Xu Q, and Li J. Prognostic impact of ARHGAP43(SH3BP1) in acute myeloid leukemia. J Formos Med Assoc. 2024;123(9):992-1003. Epub 20240405. doi: 10.1016/j.jfma.2024.04.002. PubMed PMID: 38582737.

19.   Fang J, Bolanos LC, Choi K, Liu X, Christie S, Akunuru S, et al. Ubiquitination of
hnRNPA1 by TRAF6 links chronic innate immune signaling with myelodysplasia. Nat
Immunol. 2017;18(2):236-45. Epub 20161226. doi: 10.1038/ni.3654. PubMed PMID:
28024152; PMCID: PMC5423405.

20.   Wang S, Lu Y, Woods K, Di Trapani G, and Tonissen KF. Investigating the
Thioredoxin and Glutathione Systems' Response in Lymphoma Cells after Treatment
with [Au(d2pype)2]CL. Antioxidants (Basel). 2021;10(1). Epub 20210113. doi:
10.3390/antiox10010104. PubMed PMID: 33451071; PMCID: PMC7828567.

21.   Lee E, Kim JY, Kim TK, Park SY, and Im GI. Methyltransferase-like protein 7A
(METTL7A) promotes cell survival and osteogenic differentiation under metabolic
stress. Cell Death Discov. 2021;7(1):154. Epub 20210630. doi: 10.1038/s41420-021-
00555-4. PubMed PMID: 34226523; PMCID: PMC8257615.

22.   Piazza R, Magistroni V, Mogavero A, Andreoni F, Ambrogio C, Chiarle R,
Mologni L, et al. Epigenetic silencing of the proapoptotic gene BIM in anaplastic large
cell lymphoma through an MeCP2/SIN3a deacetylating complex. Neoplasia.
2013;15(5):511-22. doi: 10.1593/neo.121784. PubMed PMID: 23633923; PMCID:
PMC3638354.

23.   Di Crosta M, Arena A, Benedetti R, Gilardini Montani MS, and Cirone M. 5-AZA
Upregulates SOCS3 and PTPN6/SHP1, Inhibiting STAT3 and Potentiating the Effects of
AG490 against Primary Effusion Lymphoma Cells. Curr Issues Mol Biol.
2024;46(3):2468-79. Epub 20240314. doi: 10.3390/cimb46030156. PubMed PMID:
38534772; PMCID: PMC10968859.

24.   Huang PY, Best OG, Almazi JG, Belov L, Davis ZA, Majid A, et al. Cell surface
phenotype profiles distinguish stable and progressive chronic lymphocytic leukemia.
Leuk Lymphoma. 2014;55(9):2085-92. Epub 20140124. doi:
10.3109/10428194.2013.867486. PubMed PMID: 24289109.

25.   Freedman A, and Jacobsen E. Follicular lymphoma: 2020 update on diagnosis
and management. Am J Hematol. 2020;95(3):316-27. Epub 2019/12/10. doi:
10.1002/ajh.25696. PubMed PMID: 31814159.

26.   Pileri SA, and Falini B. Mantle cell lymphoma. Haematologica. 2009;94(11):1488-
92. doi: 10.3324/haematol.2009.013359. PubMed PMID: 19880776; PMCID:
PMC2770958.

27.   Majzner RG, Rietberg SP, Sotillo E, Dong R, Vachharajani VT, Labanieh L, et al.
Tuning the Antigen Density Requirement for CAR T-cell Activity. Cancer Discov.
2020;10(5):702-23. Epub 20200319. doi: 10.1158/2159-8290.CD-19-0945. PubMed
PMID: 32193224; PMCID: PMC7939454.

28.   Vilchez D, Boyer L, Morantte I, Lutz M, Merkwirth C, Joyce D, et al. Increased
proteasome activity in human embryonic stem cells is regulated by PSMD11. Nature.

2012;489(7415):304-8. Epub 2012/09/14. doi: 10.1038/nature11468. PubMed PMID:
22972301.

29.   Iqbal J, Shen Y, Liu Y, Fu K, Jaffe ES, Liu C, et al. Genome-wide miRNA profiling
of mantle cell lymphoma reveals a distinct subgroup with poor prognosis. Blood.
2012;119(21):4939-48. Epub 20120405. doi: 10.1182/blood-2011-07-370122. PubMed
PMID: 22490335; PMCID: PMC3367895.

30.   Izykowska K, Przybylski GK, Gand C, Braun FC, Grabarczyk P, Kuss AW, et al.
Genetic rearrangements result in altered gene expression and novel fusion transcripts
in Sezary syndrome. Oncotarget. 2017;8(24):39627-39. Epub 2017/05/11. doi:
10.18632/oncotarget.17383. PubMed PMID: 28489605; PMCID: PMC5503638.

31.   Wang X, Bjorklund S, Wasik AM, Grandien A, Andersson P, Kimby E, et al. Gene
expression profiling and chromatin immunoprecipitation identify DBN1, SETMAR and
HIG2 as direct targets of SOX11 in mantle cell lymphoma. PLoS One.
2010;5(11):e14085. Epub 20101122. doi: 10.1371/journal.pone.0014085. PubMed
PMID: 21124928; PMCID: PMC2989913.

32.   Rolland D, Raharijaona M, Barbarat A, Houlgatte R, and Thieblemont C.
Inhibition of GST-pi nuclear transfer increases mantle cell lymphoma sensitivity to
cisplatin, cytarabine, gemcitabine, bortezomib and doxorubicin. Anticancer Res.
2010;30(10):3951-7. PubMed PMID: 21036708.

33.   Alsagaby SA. Biological roles of THRAP3, STMN1 and GNA13 in human blood
cancer cells. 3 Biotech. 2024;14(10):248. Epub 20240925. doi: 10.1007/s13205-024-
04093-5. PubMed PMID: 39345963; PMCID: PMC11424602.

34.   Lee M, Hamilton JAG, Talekar GR, Ross AJ, Michael L, Rupji M, et al. Obesity-
induced galectin-9 is a therapeutic target in B-cell acute lymphoblastic leukemia. Nat
Commun. 2022;13(1):1157. Epub 2022/03/05. doi: 10.1038/s41467-022-28839-y.
PubMed PMID: 35241678; PMCID: PMC8894417.

35.   Tao Y, Yin H, Shen R, He S, and Miao X. USP13 mediates resistance to Ibrutinib
in diffuse large B-cell lymphoma via augmenting FHL1 stabilization. Am J Cancer Res.
2025;15(7):2932-48. Epub 20250715. doi: 10.62347/IMFU7145. PubMed PMID:
40814378; PMCID: PMC12344158.
607

Swenson, et al.- Figure 1

Swenson, et al. MCL Patients Figure 2

Swenson, et al. MCL Patients Figure 3

A.

B.

Swenson, et al. MCL Patients Figure 4