1   **A Multidimensional Framework for Behavioral Persistence: Dissociable Dimensions of**

2   **Effort, Endurance, and Sequence Stability in Mice**

3   Tianyu Cao, William R. Johnston, Stephen Christensen, Samuel Crouse and Qian-Quan Sun*

4   *Department of Zoology and Physiology, University of Wyoming, Laramie, WY 82071

5   *Corresponding Author, email: neuron@uwyo.edu

Abstract

Behavioral persistence, the maintenance of goal-directed action despite obstacles, is a
fundamental adaptive process, yet its scientific study remains fragmented across disciplines
with disparate operational definitions. Here, we introduce the Persistence Spectrum (PERCS)
framework, a five-dimensional model deconstructing persistence into Perseverance of Effort (P),
Strategic Endurance (E), Resistance to Extinction (R), Temporal Consistency (C), and
Repetitive Sequence Stability (S). Using programmable operant schedules via the FED3
system, we induced a continuum of persistent food-seeking in mice across four paradigms:
Fixed Ratio (FR), Alternating 2×2 and 5×5, and Random Progressive Ratio (RPR). To
objectively identify persistence periods, we developed a session-specific Gaussian mixture
model, providing a data-driven alternative to arbitrary frequency thresholds. We found that
persistence bouts were fundamentally driven by unrewarded effort, not reward delivery:
instantaneous frequency for incorrect pokes significantly exceeded that for correct pokes in FR,
2×2, and 5×5 (all $p < 0.05$). This pattern was most pronounced in the high-demand RPR
schedule, where high-rate poking continued unabated even after successes, directly challenging
reinforcement-centric models and providing strong empirical support for frustration theory.
Linear mixed-effects modeling revealed that frequency increased with consecutive unrewarded
pokes across all paradigms, with a quadratic (inverted-U) relationship specific to FR and 2×2 ($p
< 0.001$), suggesting effort invigoration followed by strategic disengagement in simpler tasks,
whereas high-demand schedules maintained linear increases. Notably, the collapse of
Sequence Stability (S) in RPR is partly constrained by the task's environmental contingency:
random reward rules mathematically restrict stable sequence learning. Aggregate analysis
showed total poke counts and unrewarded effort scaled with task difficulty (all $p < 2e-16$), yet
pellet retrieval rates remained stable, indicating goal achievement despite increased challenge.

Crucially, PERCS profiles were robust across independent and continuous training histories,

demonstrating they reflect stable phenotypes shaped by current contingencies rather than

training artifacts. Application of PERCS revealed distinct fingerprints: FR produced near-zero P,

E, R, and C but maximal S, characteristic of an efficient habit; 2×2 and 5×5 elevated P, E, and R

while reducing S; and RPR generated highest P and R, lowest S, and marked inter-individual

variability. These findings demonstrate that operant schedules dissociably shape distinct

persistence dimensions, with unrewarded effort acting as a key motivational trigger, positioning

frustrative nonreward as a primary engine of persistent behavior. The PERCS framework

provides a unified, quantitative language for characterizing persistent behavior across species

and paradigms, offering a powerful tool for linking dimensions to neural circuits and

understanding their breakdown in neuropsychiatric disorders.

Introduction

Behavioral persistence, the maintenance of goal-directed actions despite obstacles, delay, or diminishing returns, is a fundamental adaptive process across species, essential for survival, foraging, and long-term achievement. Despite its centrality, the scientific study of persistence is conceptually fragmented, scattered across disciplines with disparate operational definitions and terminologies that hinder a unified, mechanistic understanding (Nevin and Grace 2000); (Amsel and Roussel 1952). In comparative psychology, persistence is often studied through specific, narrow lenses. Behavioral momentum theory conceptualizes it as resistance to change when reinforcement is disrupted or withheld (Nevin and Grace 2005, Podlesnik and Shahan 2008). Progressive ratio (PR) schedules reduce it to a single motivational "breakpoint" (Hodos 1961); (Richardson and Roberts 1996, Bradshaw and Killeen 2012), while foraging models describe it as the scalable continuation of feeding under threat. In parallel, human psychology has developed constructs like grit, defined as perseverance for long-term goals (Duckworth, Peterson et al. 2007), and delay of gratification (Mischel, Shoda et al. 1989), which focus on trait-level or cognitive self-regulation but lack direct neurobiological translatability. Clinical research further complicates the picture by studying pathological extremes, such as compulsivity in obsessive-compulsive disorder (Graybiel and Rauch 2000) or amotivation in depression (Treadway and Zald 2011), and cognitive rigidity in autism spectrum disorder (Geurts, Corbett et al. 2009, Lord, Brugha et al. 2020),without a dimensional framework that connects these states to adaptive persistence. This fragmentation presents a fundamental challenge: we lack a common operational vocabulary and a controlled experimental paradigm capable of inducing and dissecting the full continuum of persistent states, from efficient goal-pursuit to maladaptive perseveration(Eisenberger 1992, Carver and Scheier 1998).

Recent technological advances in behavioral monitoring offer a path forward. The Feeding

Experimentation Device version 3 (FED3) has emerged as a transformative tool for high-

resolution, home-cage measurement of feeding microstructure and operant behavior,

minimizing stress and enabling rich, longitudinal phenotyping (Matikainen-Ankney, Earnest et al.

2021). It has been used descriptively to study threat-suppressed feeding (de Araujo Salgado, Li

et al. 2023), effort-based motivation in obesity models (Barrett, Raineri Tapies et al. 2020), and

neural correlates of consumption . However, its potential extends beyond description. Crucially,

the FED3 is programmable. We propose to leverage this feature not merely to measure

persistence, but to actively engineer it by systematically manipulating reinforcement

contingencies(Skinner 1988) (Thorndike 1933). By designing operant schedules that

parametrically vary effort cost, reward uncertainty, and sequential complexity, we can directly

induce a broad spectrum of persistent behavioral states within the same subjects. This creates

a controlled experimental "persistence continuum," a necessary substrate for rigorous,

multidimensional analysis.

Quantifying this continuum requires moving beyond single metrics like breakpoints or total

responses (Stevens 1946; Cronbach and Meehl 1955). A mouse pressing a lever hundreds of

times on a PR schedule, a bird vigilantly foraging under predation risk, and a human working

diligently on a long-term project may all be described as "persistent," but they engage

qualitatively different behavioral strategies (Tinbergen 1963, 1972; McFarland and Sibly 1975;

Schleidt 1974). Deconstructing persistence into its core, dissociable components is therefore

essential for linking specific behavioral patterns to their underlying neural circuits and for

understanding individual differences in adaptive and maladaptive behavior (Eisenberger 1992,

Sayar-Atasoy, Yavuz et al. 2024)(Sayar-Atasoy et al., 2024). To address this, we introduce the

Persistence Spectrum (PERCS) framework, a five-dimensional model designed to provide a

unified, quantitative characterization of persistent behavior. PERCS deconstructs persistence

into the core dimensions of Perseverance of Effort (P), Strategic Endurance (E), Resistance to

Extinction (R), Temporal Consistency (C), and Repetitive Sequence Stability (S). Temporal

Consistency (C) draws on extensive work in biological rhythms and circadian behavior (Boulos

and Terman 1980, Frank, Kupfer et al. 2005, Wang and Sun 2024)(Boulos and Terman 1980;

Wang and Sun 2023; Frank, Kupfer et al. 2005), while Repetitive Sequence Stability (S) builds

on ethological analyses of action syntax(Markowitz, Gillis et al. 2023) (Jin, Tecuapetla et al.

2014, Wiltschko, Johnson et al. 2015, Weinreb, Pearl et al. 2024).

Our primary objective was to test the hypothesis that distinct operant schedules would

differentially shape these PERCS dimensions, generating unique behavioral "fingerprints" or

profiles. We predicted that schedules imposing higher effort costs (e.g., progressive ratio) would

elevate P (Perseverance of Effort); schedules with unpredictable rewards or requiring complex

sequences would modulate E (Strategic Endurance) and S (Sequence Stability); and the R

(Resistance to Extinction) and C (Temporal Consistency) dimensions would capture

fundamental differences between habit-like, automated responding and flexible, goal-directed

persistence (Squire and Zola 1996, Graybiel 2008). This work establishes a new methodological

paradigm: using a programmable device (FED3) to induce a persistence continuum and

applying a multidimensional computational framework (PERCS) to deconstruct it. By bridging

controlled operant paradigms, naturalistic ethology, and clinical concepts, this approach

provides a common quantitative language for persistence research. Here, we demonstrate that

a series of FED3 operant schedules, Fixed Ratio (FR), Alternating (2×2, 5×5), and Random

Progressive Ratio (RPR), reliably produce a wide range of persistent feeding behaviors in mice.

Our multidimensional analysis reveals that these schedules produce distinct, quantifiable

PERCS profiles. For instance, the low-cost FR schedule produces high S but low P,

characteristic of an efficient habit (Squire and Zola 1996, Graybiel 2008) , while the high-cost

RPR schedule elevates P and R but collapses S, indicative of effortful but less structured

responding. These findings validate the PERCS framework as a powerful tool for mapping the

adaptive landscape of persistent behavior and set the stage for future research to identify the

specific neurobiological substrates governing each persistence dimension.

Results

Threshold Determination Using Gaussian Mixture Models

To determine the high and low instantaneous frequency thresholds used to define persistence

periods objectively, we fitted a two-component Gaussian mixture model (GMM) separately for

each mouse and for each training session, using the flexmix R package. This session-specific,

mouse-specific approach avoids pooling data across sessions or individuals and accounts for

both inter-individual and within-animal variability in poking behavior. This data-driven method

provides an objective alternative to arbitrary frequency thresholds, thereby enhancing the

reproducibility of "persistence" as a behavioral construct. Let y_i denote the instantaneous

frequency of the ith poke within a given mouse and session. We assumed that y_i arises from a

mixture of two normal distributions, $y_i \sim \pi_1 N(\mu_1, \sigma_1^2) + \pi_2 N(\mu_2, \sigma_2^2)$, where π_1 and π_2 = 1-

π_1 are the mixture proportions, and μ_k and σ_k denote the mean and variance of component k ,

for k = 1,2 . Although the true distribution of instantaneous frequencies is unknown, this two-

component Gaussian mixture model provided a reasonable and robust separation between low-

and high-frequency pokes. Model parameters (π_k, μ_k, σ_k^2) were estimated using the

expectation–maximization (EM) algorithm. For each poke i , the posterior probability of

belonging to component k was computed as $Pr(z_i = k \mid y_i) = (\pi_k N(y_i \mid \mu_k, \sigma_k^2)) / (\sum_j \pi_j N(y_i \mid \mu_j,$

$\sigma_j^2))$, where z_i is a latent component indicator. Each poke was assigned to the component with

the higher posterior probability. The high instantaneous frequency threshold was defined

as High bound = (max{y_i | z_i = low}1 + min{y_i | z_i = high}) / 2 , and the low instantaneous

frequency threshold was defined as $Low\ bound = \mu_{low}$. These session-specific thresholds were

then used to identify persistence periods. To visualize the mixture model and the resulting

thresholds, we added fitted Gaussian envelopes to the histograms of instantaneous

frequencies. An example from a single session of a randomly selected mouse from the 2×2

paradigm shows that the two-component Gaussian mixture model, while not perfectly capturing

all features of the data, provides sufficiently distinct components to determine the threshold for

persistence periods (Supplementary Figure 1).

Expression of Persistence Behavior Across Paradigms

To dissect the architecture of persistent pursuit, we aligned poking behavior to the onset of

persistence and non-persistence periods across schedules of escalating cost (Figure 1B). In the

low-effort FR schedule, poking was characterized by sparse, low-frequency events. Conversely,

the high-cost RPR schedule evoked the highest overall poking density, marked by sustained

high instantaneous frequencies, indicating maximal energetic investment. Relative to the

intermediate 2×2 and 5×5 schedules, RPR mice initiated persistence periods more frequently,

though each bout was shorter in duration. Across all paradigms, persistence periods were

characterized by a striking elevation in poke frequency compared to equivalent non-persistence

epochs. Critically, high-frequency poking clusters were composed almost exclusively of

incorrect pokes, revealing that these bouts represent phasic bursts of unrewarded, cost-tolerant

effort rather than reward-oriented correct responses.

Persistence Periods Are Driven by Unrewarded Effort

To determine whether persistence periods represent bursts of unrewarded effort, we compared

instantaneous poking frequencies for correct versus incorrect pokes within each schedule

(Figure 2A). In the FR, 2×2, and 5×5 schedules, instantaneous frequency was significantly lower

for correct pokes than for other pokes (FR: $t_{(4)}$ = -9.76, p = 0.0006, mean difference = -0.705;

2×2: $t_{(4)}$ = -21.98, p < 0.0001, mean difference = -0.108; 5×5: $t_{(4)}$ = -3.50, p = 0.025, mean

difference = -0.148), confirming that high-frequency clusters are composed primarily of

unrewarded responses. Notably, this pattern held even in the FR schedule, which lacks formal

persistence periods, suggesting that the drive for high-frequency poking is not contingent on

schedule complexity. In sharp contrast, the RPR schedule revealed no significant difference

between correct and other pokes ($t_{(4)}$ = 2.49, p = 0.067, mean difference = 0.034), with incorrect

pokes exhibiting a marginally lower mean frequency than correct pokes. This divergence

reflects the unique contingency of the RPR schedule, in which reward delivery depends on a

cumulative poke count across both ports, rather than on consecutive correct responses.

Consequently, rewards do not reliably terminate a persistence bout, and mice continue poking

at high rates even after successes, accounting for the frequent but brief persistence periods

observed in Figure 1. Together, these results are consistent with frustration theory(Amsel and

Roussel 1952), wherein the omission of an expected reward energizes ongoing behavior,

driving phasic bursts of unrewarded effort. This finding directly challenges reinforcement-centric

models of persistence, which would predict that rewarded actions are the primary drivers of

behavior. Instead, our data position frustrative nonreward as a primary engine of persistent

responding.

Influence of Consecutive Unrewarded Pokes on Effort

To further examine how unrewarded actions influence poking behavior, we introduced a new

variable, "Number of Consecutive Other Pokes," which counts how many consecutive pokes do

not yield a pellet. Specifically, this variable is reset to zero whenever a correct poke resulting in

a pellet occurs or when a pellet is retrieved. Counting begins only after the first unrewarded

("other") poke following a correct poke or pellet retrieval, and it increments with each

subsequent consecutive unrewarded poke until a rewarded action occurs. This measure allows

us to investigate how mice adjust their effort when repeated pokes fail to produce a reward

(Figure 2B). We fitted a linear mixed-effects model separately for each paradigm: $Y_{ij} = \beta_0 + \beta_1 X_{ij}$

$+ \beta_2 X_{ij}^2 + b_i + \epsilon_{ij}$, where Y_{ij} = average instantaneous frequency for mouse i at a given

consecutive-other-poke count, X_{ij} = number of consecutive other pokes, $\beta_0, \beta_1, \beta_2$ = fixed effects

(intercept, linear, quadratic slope), b_i = random intercept for mouse i , and ϵ_{ij} = random error. To

account for heteroscedasticity, residual variances were allowed to differ across levels of X_{ij}

using a *varIdent* variance structure. Specifically,

$$\varepsilon_{ij} \sim \mathcal{N}(0, \sigma_k^2),$$

where the residual variance σ_k^2 depends on the level k of X_{ij}. This heteroscedastic variance

structure was implemented for all paradigms.

In Table 1, we can see that for all four paradigms, the slope of the linear term (consec_Other)

was positive and statistically significant (p < 0.05), indicating effort increases as failures rise.

However, the quadratic (inverted-U) term was only significant in the FR and 2×2 paradigms (p <

0.001). We found a quadratic (inverted-U) relationship in simpler tasks (FR, 2×2), revealing an

initial effort invigoration followed by strategic disengagement (p < 0.001), whereas high-demand

tasks (5×5, RPR) maintained linear effort increases, likely driven by the energizing effects of

frustrative nonreward. Overall, these results imply that for most paradigms, average

instantaneous poking frequency increases approximately linearly with consecutive other poke

counts, while in some cases (e.g., FR, 2×2) the relationship exhibits a nonlinear, inverted-U

shape.

Effectiveness of Effort in Pellet Retrieval

To quantify how effectively mice converted their poking effort into rewards, we analyzed the

instantaneous frequency of pellet retrieval across the different paradigms (Supplementary

Figure 2). Four panels, arranged from top to bottom corresponding to the FR, 2×2, 5×5, and

RPR paradigms, show the instantaneous frequency of pellet retrieval for the randomly selected

mice in Figure 1. The instantaneous frequency for each pellet was calculated as the reciprocal

of the time interval between the current pellet and the previous pellet. Across all paradigms, the

instantaneous frequency of pellet retrieval was markedly lower and more stable than the

instantaneous frequency of poking behavior. Unlike poking, which exhibited substantial

variability and paradigm-dependent structure, pellet retrieval occurred at a relatively constrained

rate with limited fluctuation over time. This pattern was consistently observed in the FR, 2×2,

5×5, and RPR paradigms, indicating that despite large differences in response dynamics, the

temporal structure of reinforcement remained comparatively uniform across task designs. This

stability in reward rate despite varying effort demands echoes findings from studies of

behavioral regulation and homeostasis (McFarland and Sibly 1975, Boulos and Terman 1980).

Aggregate Comparison Across All Mice and Paradigms

To assess the cumulative impact of response cost on effort and reward acquisition, we

compared total pokes, other (incorrect) pokes, and pellet retrieval counts across all schedules

(Figure 3A; n = 5 mice per paradigm). Total poke counts increased monotonically with task

difficulty, with the FR schedule yielding the lowest counts and the RPR schedule the highest;

the 2×2 and 5×5 schedules showed intermediate and comparable levels. A parallel pattern was

observed for other pokes, indicating that unrewarded effort scales with schedule demands. In

contrast, pellet retrieval counts remained stable across all four paradigms, suggesting that

despite escalating costs, mice maintained consistent food intake.

Generalized linear mixed-effects models (GLMM) with a negative binomial distribution confirmed

these observations. Relative to the FR baseline, total poke counts were significantly elevated in

all schedules (2×2: estimate = 0.653, p < 2e-16; 5×5: estimate = 0.664, p < 2e-16; RPR:

estimate = 1.635, p < 2e-16). Similarly, other-poke counts were markedly higher in all schedules

compared to FR (2×2: estimate = 3.100, p < 2e-16; 5×5: estimate = 3.117, p < 2e-16; RPR:

estimate = 4.477, p < 2e-16). Crucially, pellet retrieval counts did not differ significantly from FR

in any schedule (2×2: p = 0.098; 5×5: p = 0.113; RPR: p = 0.460), confirming that reward

acquisition rate was preserved despite increased effort and failure tolerance. Together, these

results demonstrate that mice adapt to rising behavioral obstacles by investing greater

unrewarded effort, yet successfully defend baseline levels of goal attainment.

Training History Does Not Confound Behavioral Phenotypes

To rule out order effects, we compared behavioral output between mice trained on independent

versus continuous schedules. Mann–Whitney tests revealed no significant differences in total

pokes, other pokes, or pellet counts within any paradigm (Figures 3B–3D; all p > 0.05),

confirming that task history did not influence performance and that subsequent analyses could

be collapsed across training cohorts.

The PERCS Framework Reveals Multidimensional Effort Strategies

We next applied the PERCS framework to quantify behavior across five core dimensions—

Persistence (P), Endurance (E), Resistance to Extinction (R), Consistency (C), and Sequence

Stability (S)—with Resistance to Extinction grounded in behavioral momentum theory via

devaluation-based satiety probes (Nevin and Grace 2005, Podlesnik and Shahan 2008). In the

FR schedule, persistence bouts were virtually absent, yielding near-zero values for P, E, R, and

C, while S approached ceiling (near 100% correct pokes), reflecting a habitual, low-effort

response structure(Squire and Zola 1996, Graybiel 2008). The 2×2 and 5×5 schedules

displayed similar multidimensional profiles, characterized by elevated P, E, and R alongside

reduced S, indicating that alternating response demands require sustained effort and strategic

endurance at the expense of sequence stability. The RPR schedule exhibited the lowest S—

consistent with the mathematical unpredictability of a random environment that precludes stable

sequence learning—while showing the highest P and R, reflecting maximal effort and continued

responding despite uncertainty. Notably, RPR also revealed marked inter-individual variability in

E and C, suggesting that high-demand conditions unmask divergent strategies in temporal

consistency and endurance.

PERCS Profiles Are Robust Across Training Histories

To assess reproducibility, we examined PERCS profiles separately in the continuous training

cohort (Figure 4B) and in the combined dataset (Figure 4C). In both cases, the multidimensional

signatures remained qualitatively identical to those observed in the independent cohort: FR was

dominated by high S and minimal persistence-related dimensions, 2×2 and 5×5 showed

intermediate elevation of P, E, and R with reduced S, and RPR exhibited low S with maximal P

and R. The stability of these profiles across independent and continuous training histories

demonstrates that PERCS dimensions capture stable behavioral phenotypes shaped by current

task contingencies rather than by training artifacts.

Multivariate Analysis Reveals Paradigm-Specific Persistence Architectures

To determine whether distinct task contingencies recruit dissociable combinations of

persistence dimensions, we performed multivariate analyses of PERCS profiles. Analysis of

multivariate dispersion revealed significant differences in within-paradigm variability across

schedules (global permutation test, $p < 0.05$; Table 5). RPR exhibited significantly greater

dispersion than FR ($p < 0.05$), while comparisons between RPR and the alternating schedules

(2×2, $p = 0.119$; 5×5, $p = 0.151$) did not reach statistical significance, likely reflecting limited

sample size ($n = 5$ per paradigm) for high-dimensional comparisons. While this lack of

significance with the alternating schedules is acknowledged as a limitation of the current sample

size, the consistent pattern of median deviations (Figure 5A) suggests that RPR may drive

greater individual variability in persistence strategies. This trend is conceptually significant, as it

implies that environments with high uncertainty and effort demands may unmask or amplify pre-

existing individual differences in coping strategies, rather than producing a uniform behavioral

response. Nonetheless, median deviations (Figure 5A) indicated that RPR consistently showed

the greatest overall spread, suggesting that high-demand, stochastic schedules amplify

individual variability in persistence strategies.

Unsupervised clustering of raw PERCS vectors ($k = 4$) yielded three empirically distinct

groupings (Figure 5B). Cluster 1 comprised 16 mice, 94% of which (15 of 16) were from the FR

schedule, confirming FR as a qualitatively distinct, low-effort phenotype. Clusters 2 and 3

consisted exclusively of RPR mice, revealing that the RPR schedule supports at least two

separable behavioral subtypes. This divergence within the RPR schedule further supports the

idea that high-demand conditions reveal distinct behavioral phenotypes, a finding that aligns

with the trend towards increased multivariate dispersion observed in Figure 5A. Cluster 4

contained only 2×2 and 5×5 mice, indicating that these intermediate schedules occupy an

overlapping region of PERCS space and recruit similar multidimensional persistence profiles.

Principal component analysis (PCA) corroborated this structure (Figure 5C). FR formed a

compact, isolated cluster. The 2×2 and 5×5 schedules extensively overlapped, forming a

second cluster. RPR displayed the widest dispersion, occupying a region partially separated

from the other paradigms, consistent with its elevated multivariate variability. Together, these

multivariate patterns demonstrate that different task contingencies evoke distinct combinations

of persistence dimensions, supporting theoretical frameworks that posit multiple, dissociable

mechanisms underlying persistent behavior (Carver and Scheier 1982, Eisenberger 1992)..

Discussion

Our findings demonstrate that operant conditioning schedules delivered via the FED3 system

produce a systematic, quantifiable continuum of persistent food-seeking behaviors. This

continuum is directly shaped by the interplay of effort cost, sequential rule complexity, and

outcome uncertainty inherent in each schedule, validating the hypothesis that persistence is not

a binary state but a multidimensional behavioral phenotype that can be experimentally sculpted.

The results align with and extend prior work using operant paradigms to study persistence. The

finding that higher effort costs, as in the Random Progressive Ratio (RPR), elevate overall

response output (Perseverance of Effort, P) and the frequency of persistence bouts is

consistent with progressive ratio (PR) breakpoint assays, a gold standard for measuring

motivational strength ((Hodos 1961, Richardson and Roberts 1996)). However, traditional PR

schedules measure only the terminal point of effort expenditure. Our multidimensional analysis

reveals that the RPR schedule not only increases P but also drastically reduces Repetitive

Sequence Stability (S), a dimension not captured by a breakpoint. This aligns with behavioral

momentum theory (Nevin and Grace 2000), which posits that behavior under rich, predictable

reinforcement histories becomes more resistant to change. Our RPR schedule, by introducing

high uncertainty, weakens this "behavioral mass," leading to less structured, more variable

responding, a profile captured by low S, a pattern that reflects the mathematical constraints of a

random environment where stable sequence learning is impossible, thereby capturing effortful

but unstructured responding. The structured alternation schedules (2×2, 5×5) produced a

distinct profile, with elevated Strategic Endurance (E) and intermediate S. This mirrors findings

from complex sequential learning and set-shifting tasks, where animals must maintain a rule in

working memory and execute a specific action chain (Floresco, Zhang et al. 2009). We found a

quadratic (inverted-U) relationship between consecutive unrewarded pokes and instantaneous

frequency in simpler tasks (FR, 2×2), revealing an initial effort invigoration followed by strategic

disengagement at very high failure counts. In contrast, high-demand tasks (5×5, RPR)

maintained linear effort increases. The significant quadratic relationship between consecutive

unrewarded pokes and instantaneous frequency in the 2×2 paradigm suggests an "inverted-U"

of effort, an initial increase in vigor following failure, consistent with Amsel's frustration theory

(Amsel and Roussel 1952)), followed by a strategic disengagement at very high failure counts.

This nuanced pattern would be invisible in a simple response-count analysis.

A key novel finding is that persistence bouts are triggered by unrewarded effort, not reward

delivery. This was most evident in the RPR schedule, where high-rate poking continued even

after occasional rewarded trials, and in the paired comparisons showing that "other" (incorrect)

pokes had higher instantaneous frequencies than correct pokes in most paradigms. This directly

challenges a purely reinforcement-driven account of persistence, which would predict that

rewarded actions are the primary drivers of behavioral vigor. Instead, it strongly supports a

motivational view where the absence of expected reward (frustrative nonreward) energizes

ongoing behavior (Amsel and Roussel 1952). This mechanistic insight into the trigger of

persistence is a significant advancement beyond descriptive schedules like PR, positioning

frustrative nonreward as a primary, rather than secondary, engine of persistent responding. This

energizing effect is most directly captured by the Perseverance of Effort (P) dimension, which

quantifies total work output, and the inverted-U pattern in simpler tasks reveals the dynamic

interplay between P and Strategic Endurance (E), where initial invigoration gives way to

adaptive disengagement. This directly challenges a purely reinforcement-driven account of

persistence and supports a motivational view where the absence of expected reward (frustrative

nonreward) energizes ongoing behavior (Amsel and Roussel 1952). This mechanistic insight

into the trigger of persistence is a significant advancement beyond descriptive schedules like

PR. Furthermore, our use of Gaussian mixture modeling (GMM) to define persistence periods

session-by-session provides objective, data-driven alternative to arbitrary frequency thresholds,

enhancing the reproducibility of "persistence" as a behavioral construct. This method accounts

for individual and session variability, making the identification of "persistent states" more robust

and comparable across subjects and time, a methodological improvement over fixed criteria.

Finally, the Fixed Ratio (FR) schedule served as a critical control, generating a PERCS profile of

near-zero P, E, R, and C, but maximal S. This profile represents a pure, efficient habit (Graybiel

2008), highly automated, temporally irregular, and requiring minimal cognitive engagement. It

starkly illustrates that high sequence stability can exist independently of other persistence

dimensions, a dissociation made explicit by our framework. The neural substrates of such

habitual behavior likely involve corticostriatal loops and dopamine signaling (Jin, Tecuapetla et

al. 2014, Goedhoop, van den Boom et al. 2022, Goedhoop, Arbab et al. 2023). In summary, our

FED3 results confirm that schedule parameters dissociably shape distinct facets of persistence.

They bridge classic operant concepts (breakpoint, resistance to extinction) with a richer,

multidimensional characterization, providing a more precise map of how challenge structure

dictates behavioral strategy.

Understanding behavioral persistence is of paramount importance. It is a transdiagnostic feature

spanning adaptive success, from foraging and migration to academic and professional

achievement, and debilitating psychopathology, including the compulsions of OCD, the apathy

of depression (Marin 1991, Treadway and Zald 2011), the perseveration of frontal lobe

syndromes (Sandson and Albert 1984) , the inattention of ADHD (Barkley 1997), and the

cognitive rigidity of ASD ((Graybiel and Rauch 2000); (Treadway and Zald 2011, Lord, Brugha

et al. 2020)). Despite this centrality, the field has been historically fragmented, as detailed in our

introduction. This fragmentation arises from divergent disciplinary roots: behaviorism's focus on

reinforcement history (Nevin and Grace 2000), human psychology's emphasis on traits like grit

(Duckworth, Peterson et al. 2007) and willpower (Baumeister, Bratslavsky et al. 1998), and

clinical neuroscience's categorical approach to disorders. Consequently, the field lacks a

common operational language. The "persistence" measured by a PR breakpoint in a rat, the

"tenacity" of a student studying for exams, and the "perseveration" of a patient with frontal lobe

damage are often discussed in isolation, impeding the integration of knowledge across levels of

analysis and the translation of mechanistic insights from model systems to human conditions.

Our work directly addresses this gap. By employing the FED3, a tool capable of inducing a full

continuum of persistent states under controlled conditions, we move persistence from a

descriptive phenomenon observed in response to external manipulations (e.g., threat, drugs) to

an experimentally tractable dependent variable. This allows for the systematic investigation of

its causes and components, a necessary step for a mature science.

The PERCS framework represents more than a novel description of our data; it offers a

transformative roadmap for future research in behavioral neuroscience, ethology, and

psychiatry. First, it provides a dimensional, quantitative phenotyping tool. Notably, our

framework identifies Persistence (P), Endurance (E), Resistance to Extinction (R), Consistency

(C), and Sequence Stability (S) as core dimensions, with Resistance to Extinction (R) measured

through devaluation-based satiety probes, grounding this dimension in behavioral momentum

(Nevin and Grace 2005, Podlesnik and Shahan 2008). PERCS offers a common metric for

cross-species, cross-paradigm, and cross-disorder comparison. Future studies can now quantify

whether a genetic mutation, pharmacological treatment, or environmental insult specifically

alters Perseverance of Effort (P) versus Repetitive Sequence Stability (S). For example, does a

dopamine agonist increase effort tolerance (P) in a PR task and strategic vigilance (E) in a

sustained attention task? Such questions can now be addressed using mouse models of

neuropsychiatric disorders(Peca, Feliciano et al. 2011, Carcea, Caraballo et al. 2021). PERCS

allows these ostensibly different outcomes to be recognized as potentially related manipulations

of the persistence spectrum. This dimensional approach aligns with the NIH's Research Domain

Criteria (RDoC) initiative (Onnela and Rauch 2016, Insel 2018),moving beyond diagnostic labels

toward dysfunction in core behavioral systems. Second, PERCS enables disentangling adaptive

from maladaptive persistence. A core promise of PERCS is its ability to distinguish functional

persistence from dysfunctional perseveration. As shown in our results and theoretical model,

pathology often involves an imbalance or extreme expression of one dimension (e.g.,

pathologically high R and S in OCD) or a deficit in another (e.g., pathologically low P and E in

depression). By profiling patients along these five dimensions, clinicians could move toward a

more precise, mechanism-based diagnosis. Treatment could then be targeted: cognitive-

behavioral therapy with exposure and response prevention might specifically aim to reduce

maladaptive R and S in OCD(Azrin and Nunn 1973, Davis, Ressler et al. 2006) (Azrin and Nunn

1973; Davis, Ressler et al. 2006), while behavioral activation therapy aims to augment deficient

P and E in depression(Lejuez, Hopko et al. 2011).Cognitive remediation approaches (Wykes,

Huddy et al. 2011) and executive function training could target specific dimensional

deficits. Third, PERCS facilitates bridging levels of analysis from behavior to circuit. The

dissociable nature of PERCS dimensions suggests they may have partially distinct neural

substrates. P (effort) is heavily linked to dopamine signaling in the ventral striatum and anterior

cingulate cortex (Salamone et al., 2007). E (endurance/attention) relies on norepinephrine and

prefrontal networks (Aston-Jones and Cohen 2005). S (sequence stability) involves

corticostriatal loops responsible for habit formation (Jin, Tecuapetla et al. 2014). C (temporal

consistency) engages circadian and ultradian rhythm generators (Boulos and Terman 1980,

Frank, Kupfer et al. 2005, Wang and Sun 2024). Future research can test these mappings

directly by correlating neural activity (e.g., via calcium imaging or electrophysiology in mice

performing our FED3 schedules) with simultaneously derived PERCS profiles. This would allow

us to move from asking "what brain region is active during a hard task?" to "what circuit dynamic

encodes a specific dimension of persistent challenge?" Fourth, PERCS enables computational

psychiatry and personalized prediction. The tiered normalization strategy of PERCS allows raw

behavioral data from diverse sources, home-cage monitoring, clinical task batteries, even digital

phenotyping from smartphones (Barnett and Onnela 2020), to be transformed into comparable

dimension scores. This opens the door for large-scale computational phenotyping. Machine

learning models could be trained on these multidimensional profiles to predict individual risk for

disorders, trajectory of illness, or response to specific treatments, fostering a new era of

personalized medicine. While the current study establishes the normative adaptive landscape

across these schedules, a critical next step involves identifying the "pathological tipping point"

where adaptive persistence transitions into maladaptive perseveration. Although diseased

models were initially considered, future research will utilize the baseline established here to

explicitly profile such models along the PERCS dimensions to differentiate, for example, high S

in OCD from low P in depression. While the current study establishes the normative adaptive

landscape across these schedules, a critical next step involves identifying the "pathological

tipping point" where adaptive persistence transitions into maladaptive perseveration. The

diseased model mice mentioned in the Methods were part of a preliminary exploration, but the

data are not included here as the focus of this foundational paper is to establish the normative

PERCS framework and validate its sensitivity to schedule parameters. Future research will

explicitly profile such disease models along the PERCS dimensions using the baseline

established here to differentiate, for example, high S in OCD from low P in depression.

A limitation of the current study is the relatively small cohort used for independent training (N=5

per paradigm). While the primary effects on total pokes and dimension rankings were highly

significant, the smaller sample size likely reduced the power of our multivariate dispersion tests.

This is particularly relevant for the comparisons between RPR and the alternating schedules

(2×2 and 5×5) in Figure 5A, where non-significant p-values may reflect a Type II error rather

than a true absence of effect. The consistent pattern of median deviations and the emergence

of two distinct RPR clusters in Figure 5B, however, suggest that these trends are biologically

meaningful and warrant investigation in larger cohorts. Environments with high uncertainty and

effort demands may unmask pre-existing individual differences in coping strategies, leading to

divergent behavioral phenotypes. Crucially, however, the multidimensional behavioral

signatures we observed were robust across both independent and continuous training histories

(Figures 3B-D, 4A-C), demonstrating that the PERCS profiles are stable phenotypes shaped by

current task contingencies rather than mere training artifacts. This robustness strengthens

confidence that our findings reflect fundamental properties of the schedules themselves. Future

studies with larger populations will be required to further refine the statistical boundaries of

these behavioral clusters. In conclusion, this study establishes a synergistic methodology: using

programmable operant schedules to induce a persistence continuum and applying the

multidimensional PERCS framework to characterize it. We have demonstrated that schedule

uncertainty and structure dissociably shape core dimensions of persistent behavior, with

unrewarded effort acting as a key trigger, challenging reinforcement-centric models and

positioning frustrative nonreward as a primary engine of persistence. By providing a unified,

quantitative, and neurobiologically-grounded model, the PERCS framework transcends

historical fragmentation. It offers a powerful new lens through which to view goal-directed

behavior, from the rhythmic lick of a mouse to the sustained striving of a human, and a precise

toolkit for deconstructing its breakdown in disease. This work lays the foundation for a unified,

mechanistic science of behavioral persistence.

Methods and Materials

Subject details

All experimental procedures were approved by the Institutional Animal Care and Use Committee

(IACUC) and the Biosafety Committee of the University of Wyoming. 10 male and 10 female

immunocompetent mice of postnatal day 101.5±23.3 days on a B6.Cg-Snap25tm3.1Hze/J

background were included in the datasets. All mice were bred on a C57/BL6J background. Mice

older than 30 days were housed with same-sex littermates or alone in a vivarium at 21-23 °C

with 25%-30% humidity and a 12-h light/dark cycle.

Food Restriction Procedure

To elicit goal-oriented behavior, we implemented a food restriction protocol not exceeding 30

days, with body weight maintained no lower than 85% of baseline, consistent with IACUC

guidelines and established standards for behavioral neuroscience research (National Research

Council 2003). Mice typically consume 10-15% of body weight daily (3-5 grams of food). Mice

were acclimated over a 7-day period with gradual reduction to the target level of 2 grams per

day (Chevee, Kim et al. 2023). To ensure adequate caloric reserves and mitigate nutritional

deficits, infant formula soymilk (SMA Wysoy) was provided as a supplement during handling,

post-behavioral training, and post-experimentation. During training and experimental sessions,

food pellets served as positive reinforcement and contributed to the daily total allowance of 2

grams (Goltstein, Reinert et al. 2018). Mice were monitored daily throughout the restriction

period, with body weight recorded pre- and post-consumption. Health assessments included

systematic evaluation of skin coat condition, posture, urine and feces output, and activity level.

Any mouse exhibiting signs of distress or body weight drop below 85% of baseline was

immediately removed from the study and allowed to recover prior to any potential re-entry

(National Research Council 2003).

Behavioral apparatus

We used a FED3-based behavioral apparatus(Matikainen-Ankney, Earnest et al. 2021) to

implement multiple feeding paradigms. The FED3 chamber consisted of three ports: two lateral

nose-poke ports (left and right) and a central pellet port for food delivery and retrieval. Task

rules governing pellet delivery were fully programmable, allowing precise control over the

response-outcome contingencies.

Operant feeding paradigms

In this study, four distinct paradigms were employed. The fixed-ratio (FR) control condition was

a paradigm with minimal response cost, in which a pellet was delivered to the central port

following a single nose poke to either lateral port. The alternating 2×2 paradigm required

alternating activation of the left and right nose-poke ports; when one lateral port was active, a

single nose poke to that port immediately triggered pellet delivery, followed by pellet retrieval,

and this poke-pellet-retrieval sequence constituted one response cycle. Two such response

cycles were required before the active port switched to the opposite side, after which the same

two-cycle sequence was repeated. The alternating 5×5 paradigm followed the same alternating

structure as 2×2 but required five consecutive response cycles before switching to the opposite

port. The random progressive ratio (RPR) condition imposed the highest response cost,

requiring animals to perform a variable total number of nose pokes across both ports before

pellet delivery; the required number of pokes was randomly drawn from a uniform distribution

between 1 and 10 and was re-sampled after each pellet retrieval. These schedules were

designed to systematically vary effort cost, sequential complexity, and reward uncertainty,

building on classic operant conditioning principles(Skinner 1988).

Response classification

523   To characterize operant responses, pokes were categorized based on their effectiveness in

524   obtaining pellets. A poke that matched the current programmed rule and successfully triggered

525   pellet delivery was classified as a correct poke (or rewarded poke). All other pokes were

526   classified as other pokes (or incorrect pokes), including those directed to an inactive port and

527   unnecessary additional pokes made after a correct poke but before pellet retrieval.

Experimental design

529   We conducted two main experimental designs. In the independent paradigm training, each of

530   the four paradigms was trained independently with five healthy, naive mice per paradigm,

531   resulting in a total of 20 mice. In the consecutive paradigm training, three healthy naive mice

532   and three diseased naive mice were each trained consecutively through all four paradigms,

533   allowing within-subject comparison across paradigms; this cohort comprised a total of six mice.

Definition and detection of persistence periods

For mice undergoing reversal training, a persistence period was defined as follows: it begins

when a mouse's poke instantaneous frequency exceeds a high threshold, continues while the

poke repeats at least four times without falling below a lower bound, and ends when the

instantaneous frequency drops below that lower bound. If the poke repeats fewer than four

consecutive times at a high instantaneous frequency, it is not considered a persistence period.

This explains why the FR paradigm, characterized by low-frequency, single pokes, does not

contain formal persistence periods. This definition captures the essence of behavioral bouts as

units of persistent action(Markowitz, Gillis et al. 2023) (Jin, Tecuapetla et al. 2014, Wiltschko,

Johnson et al. 2015, Weinreb, Pearl et al. 2024).

Threshold determination using Gaussian mixture models

To determine the high and low instantaneous frequency thresholds used to define persistence

periods objectively, we fitted a two-component Gaussian mixture model (GMM) separately for

each mouse and for each training session, using the flexmix R package. This session-specific,

mouse-specific approach avoids pooling data across sessions or individuals and accounts for

both inter-individual and within-animal variability in poking behavior. Let y_i denote the

instantaneous frequency of the i th poke within a given mouse and session. We assumed

that y_i arises from a mixture of two normal distributions, $y_i \sim \pi_1 N(\mu_1, \sigma_1^2) + \pi_2 N(\mu_2, \sigma_2^2)$,

where π_1 and $\pi_2 = 1-\pi_1$ are the mixture proportions, and μ_k and σ_k denote the mean and

variance of component k , for k = 1,2 . Model parameters (π_k, μ_k, σ_k^2) were estimated using the

expectation–maximization (EM) algorithm. For each poke i , the posterior probability of

belonging to component k was computed as $Pr(z_i = k \mid y_i) = (\pi_k N(y_i \mid \mu_k, \sigma_k^2)) / (\sum_j \pi_j N(y_i \mid \mu_j,$

$\sigma_j^2))$, where z_i is a latent component indicator. Each poke was assigned to the component with

the higher posterior probability. The high instantaneous frequency threshold was defined

as High bound = (max{$y_i \mid z_i$ = low} + min{$y_i \mid z_i$ = high}) / 2 , and the low instantaneous

frequency threshold was defined as *Low bound* = μ_{low}. These session-specific thresholds were

then used to identify persistence periods as described above.

The PERCS model: a formal, cross-species definition of persistence

562 We formally define behavioral persistence as the functional maintenance of goal-directed

563 behavior over time despite challenges, moving beyond topographic description to a functional

564 analysis of behavioral regulation(Nevin and Grace 2000). The PERCS model quantifies this as a

565 vector in a five-dimensional adaptive strategy space, where the overall behavioral profile *P* is: *P*

566 *= [P, E, R, C, S]*. Each dimension is defined as a theoretically dissociable component of

567 persistence, with quantification methods drawn from established, often cross-species,

568 experimental paradigms. *Perseverance of Effort (P)* is the total work output or energy

569 expenditure directed toward a goal against increasing effort costs or diminishing returns,

570 reflecting the intensity of engagement with a behavioral final common path (Salamone, Correa

571 et al. 2007). It is quantified as *P = Σ(Goal-Directed Responses)* or *P = Breakpoint* on a

572 progressive ratio (PR) schedule (Hodos 1961, Richardson and Roberts 1996); the PR

573 breakpoint operationalizes the point at which the cost of effort outweighs the common currency

574 value of the reward, providing a ratio-scale measure (Killeen, Posadas-Sanchez et al.

575 2009). *Strategic Endurance (E)* is the sustained, focused application of effort toward a goal

576 under conditions of acute challenge, high cognitive load, or frustrative nonreward (e.g., delay

577 discounting, distraction). This measures the stability of the goal representation in working

578 memory against competing valuations and is quantified as *E = f(Time-on-Task, Accuracy | High*

579 *Difficulty or Distraction)*, often indexed by latency to succumb to distraction or quit a demanding

580 task (Mischel, Shoda et al. 1989). *Resistance to Extinction (R)* is the tendency to continue

581 emitting a previously reinforced response after the reinforcement contingency is withdrawn. This

582 captures a fundamental learning dynamic where adaptive persistence must be distinguished

583 from maladaptive failure to update behavior(Nevin and Grace 2000) and is quantified as *R =*

584 *Number of Responses during Extinction* or as resistance to change via disruptor probes (e.g.,

585 pre-feeding, novel stimuli) within behavioral momentum theory. *Temporal Consistency (C)* is the

586 rhythmic stability, regularity, and predictability of a goal-directed behavioral stream over

587 extended time scales (days, weeks, seasons); high C indicates habit-like, automated patterning

(Graybiel 2008) behavior (Boulos and Terman 1980, Frank, Kupfer et al. 2005, Wang and Sun

2024). It is quantified as $C ≈ 1$ / Variance(Inter-Response Intervals or Session-to-Session

Frequency) , where low temporal variance indicates high rhythmicity and schedule

control. *Repetitive Sequence Stability (S)* is the structured, ordered chaining of behavioral

modules ("syllables" or actions) into a stable, repeatable sequence. This dimension quantifies

the syntax of action and its rigidity vs. flexibility (Markowitz, Gillis et al. 2023); (Wiltschko,

Johnson et al. 2015) and is quantified as *S = Markov Chain Stability* (e.g., high transition

probability from A→B→C) or low sequence entropy (Markowitz, Gillis et al. 2023); (Wiltschko,

Johnson et al. 2015).

PERCS framework quantification for the present study

For the current experiment, each PERCS dimension was operationalized using the collected

behavioral data as follows. *Perseverance of Effort (P)* was quantified as the total number of

persistence bouts, reflecting total effort investment across the session. *Strategic Endurance*

(E) was defined as the proportion of time spent in persistence periods relative to the total

duration of each training paradigm, multiplied by the proportion of correct pokes among all

pokes; this measure captures the ability to sustain effective effort toward goal

attainment. *Resistance to Extinction (R)* was operationalized as the number of persistence

bouts occurring after satiety, defined here as the point at which the mouse had obtained 120

pellets, reflecting continued effort even after the primary goal was achieved. *Temporal*

Consistency (C) was measured as the variance in the duration of individual persistence bouts,

indexing the stability of persistence over time. *Repetitive Sequence Stability (S)* was quantified

as the correct poke rate; under the task rules, higher correct poke rates necessarily reflect

stronger and more stable sequential behavioral structure.

Normalization: a tiered framework for cross-context and cross-species comparison

Raw PERCS metrics exist on disparate scales (counts, distances, entropy values). To enable

unified quantitative comparison and create a common currency for persistence (Tinbergen,

1963), we introduce a tiered normalization framework. This transforms raw dimension scores

into comparable, unitless indices ranging from 0 (minimal observed or plausible persistence) to

1 (maximal), relative to a defined reference frame. The core theoretical principle is that an

organism's persistence is represented as a normalized vector in five-dimensional space: P_{norm} =

[P_{norm}, E_{norm}, R_{norm}, C_{norm}, S_{norm}]. Each score's interpretation depends on the reference criterion

used for normalization. Three normalization strategies are applied based on the scope of

inference and research question. *Within-Study (Norm-Referenced) Normalization* is used for

comparing experimental groups within a controlled cohort (e.g., mice on different reinforcement

schedules). Raw scores are Z-scored or min-max scaled relative to the observed minimum and

maximum across the study sample; this answers which treatment group shows relatively more

or less of a persistence dimension (Cronbach and Meehl 1955). *Biologically Anchored*

(Criterion-Referenced) Normalization is used for naturalistic data or when study samples may

not capture the full biological range. Scores are scaled using a priori defined, species-typical, or

theoretically plausible minima and maxima (e.g., physiological limits, observed species

maxima). This prevents compression artifacts and asks how an individual's persistence

compares to the adaptive range for its phenotype. *Conceptual Strategy (Ideal-Type)*

Normalization is used for cross-species or cross-context strategy comparison (e.g., mouse

operant persistence vs. deer migration vs. human academic diligence). Here, normalized scores

are mapped to a common conceptual scale (minimal–typical–maximal adaptive persistence)

based on functional equivalence (Tinbergen 1972)). This constructs idealized persistence

profiles, such as "efficient forager" or "compulsive responder", to which real behavioral vectors

can be compared. Implementation and validation require that the choice of method is a critical,

theory-driven decision. A decision guide and complete algorithms are provided in

Supplementary Methods S1. Validation follows psychometric principles, including sensitivity

analysis and tests of convergent/discriminant validity across normalization tiers ((Cronbach and

Meehl 1955). In the present study, for the independent training data (Figure 14), we applied

within-study normalization. For each PERCS dimension, values were normalized such that the

maximum value across all 20 mice was set to 1, and all other values were scaled relative to this

maximum. All dimensions are thus constrained to the range of 0 to 1, facilitating direct

comparison across dimensions and paradigms.

Statistical analysis

For descriptive and comparative statistics, paired t -tests were used to compare the

instantaneous frequency of correct vs. other pokes within each paradigm. To analyze the effect

of consecutive unrewarded pokes on instantaneous frequency, we fitted a linear mixed-effects

model separately for each paradigm: $Y_{ij} = \beta_0 + \beta_1 X_{ij} + \beta_2 X_{ij}^2 + b_i + \epsilon_{ij}$, where Y_{ij} = average

instantaneous frequency for mouse i at a given consecutive-other-poke count, X_{ij} = number of

consecutive other pokes, $\beta_0, \beta_1, \beta_2$ = fixed effects (intercept, linear, quadratic slope), b_i =

random intercept for mouse i , and ϵ_{ij} = random error. To compare total poke counts, other-poke

counts, and pellet counts across paradigms and training designs, we fitted separate generalized

linear mixed-effects models (GLMMs) with a negative binomial distribution to account for over-

dispersed count data. Models included Paradigm and, where applicable, Treatment and their

interaction as fixed effects, with mouse identity as a random intercept to account for repeated

measures. Mann-Whitney U tests were used to compare outcomes (poke counts, other poke

counts, pellet counts) between the independent and continuous training designs for each

paradigm. All analyses were performed in R (version 4.x.x). A significance threshold of $p<0.05$

was used.

Table 1: Fixed Effects from Linear Mixed-Effects Models Across Paradigms

Paradigm	Fixed Effect	Estimate	Std. Error	DF	t-value	p-value
FR	Intercept	0.02742	0.00224	16	12.239	<0.001
	consec_Other	0.72071	0.08097	16	8.901	<0.001
	(consec_Other)^2	-0.12415	0.01978	16	-6.278	<0.001
2×2	Intercept	0.11582	0.03602	63	3.216	0.0021
	consec_Other	0.04661	0.00894	63	5.212	<0.001
	(consec_Other)^2	-0.00212	0.00052	63	-4.061	0.0001
5×5	Intercept	0.14280	0.05626	80	2.538	0.0131
	consec_Other	0.03708	0.01066	80	3.479	0.0008
	(consec_Other)^2	-0.00086	0.00056	80	-1.527	0.131
RPR	Intercept	0.24484	0.07922	75	3.091	0.0028
	consec_Other	0.04413	0.01320	75	3.343	0.0013
	(consec_Other)^2	-0.00068	0.00054	75	-1.262	0.211

664　　Table 2: Generalized Linear Mixed-Effects Model (GLMM, negative binomial) of poke counts

665　　across paradigms.

Effect	Estimate	Std. Error	z value	p-value
(Intercept, FR)	6.370	0.056	113.81	<2e-16 ***
Paradigm 2x2	0.653	0.078	8.36	<2e-16 ***
Paradigm 5x5	0.664	0.078	8.50	<2e-16 ***
Paradigm RPR	1.635	0.077	21.12	<2e-16 ***

667　　*Table 3: Generalized Linear Mixed-Effects Model (GLMM, negative binomial) of other-poke*

668　　*counts across paradigms.*

Effect	Estimate	Std. Error	z value	p-value
(Intercept, FR)	3.332	0.114	29.25	<2e-16 ***
Paradigm 2x2	3.100	0.138	22.41	<2e-16 ***
Paradigm 5x5	3.117	0.138	22.53	<2e-16 ***
Paradigm RPR	4.477	0.138	32.57	<2e-16 ***

671   Table 4: Generalized Linear Mixed-Effects Model (GLMM, negative binomial) of pellet counts

672   across paradigms

Effect	Estimate	Std. Error	z value	p-value
(Intercept, FR)	6.319	0.045	140.18	<2e-16 ***
Paradigm 2x2	-0.106	0.064	-1.65	0.098 .
Paradigm 5x5	-0.102	0.064	-1.59	0.113
Paradigm RPR	-0.047	0.064	-0.74	0.460

675   Table 5: Global and pairwise permutation tests for multivariate dispersion of PERCS vectors

676   across paradigms

Comparison	F	p-value	Interpretation
Overall (all 4 paradigms)	3.56	0.001	Significant difference in dispersion among groups
FR vs RPR	4.29	0.001	RPR significantly more dispersed than FR
2x2 vs RPR	3.47	0.119	No significant difference
5x5 vs RPR	3.07	0.151	No significant difference

Acknowledgements

We thank C. Zhang for animal husbandry, animal genotyping, histology assistance and items

purchasing. This work is supported by grants from National Institute of Mental Health

(1R21MH131363), and from National Institute of General Medical Sciences (2P20GM121310).

Author contributions

Q.Q.S. designed the experiment and supervised the project, acquired the funding, and wrote the

manuscript. T.C. analyzed data, performed statistical analysis and wrote the manuscript. W.J.

S.C. and SC designed and performed FED3 experiments, analyzed the data and wrote the

manuscript.

Declaration of interests

The authors declare no competing interests.

Figures and Tables

Figure 1: **Instantaneous poke frequency and raster dynamics across feeding paradigms.**

The figure consists of four rows and two columns. Each row corresponds to one feeding

paradigm (from top to bottom: FR, 2×2, 5×5, and RPR). For each paradigm, **A (left column)**

and **B (right column)** present complementary views of poke behavior from the same

representative mouse. **(A) Instantaneous frequency of poking.** For each paradigm, the

instantaneous poke frequency was calculated as the reciprocal of the inter-poke interval (time

between the current poke and the preceding poke). Each dot represents a single poke plotted

over time. Blue dots indicate correct pokes, and red dots indicate other pokes. Gray shaded

regions denote persistence periods (persistence bouts). **(B) Raster-like plot aligned to**

behavioral state transitions. Raster plots show poke timing across persistence and non-

persistence periods. Each dot represents a single poke, with time on the x-axis. For persistence

periods, onset times were aligned to the vertical dashed line. For non-persistence periods, the

beginning of each non-persistence period was aligned to zero. This alignment allows direct

comparison of poke dynamics relative to the onset of each behavioral state. Blue and red dots

again indicate correct and other pokes, respectively, and gray shading denotes persistence

bouts.

Figure 2: Instantaneous frequency differences between poke types and modulation by consecutive unrewarded pokes across paradigms.

The figure contains two main panels: **(A, left)** and **(B, right)**, showing analyses across the four feeding paradigms (FR, 2×2, 5×5, and RPR). **(A) Paired comparison of instantaneous frequency for correct and other pokes.** For each mouse within each paradigm, the average instantaneous frequency was calculated separately for correct and other (unrewarded) pokes. Each dot represents one mouse, and gray lines connect paired values from the same mouse. Statistical significance was assessed using a two-sided paired t-test comparing correct versus other pokes. **(B) Instantaneous frequency as a function of consecutive other pokes.** For each mouse within each paradigm, the average instantaneous frequency is plotted against the number of consecutive other (unrewarded) pokes. Dots represent the mean instantaneous frequency at each consecutive other poke value, and lines connect data points from the same

724   mouse. The black solid curve shows the quadratic fit from the mixed-effects model, reflecting

725   the overall trend across mice.

Figure 3: **Poke and pellet counts across paradigms and training types.**

(A) Total poke, other poke, and pellet retrieval counts for independent training.

Data are shown for all four paradigms, with five mice per paradigm. For each paradigm, three

color-coded boxplots represent total poke count, other poke count, and pellet retrieval count.

Each box summarizes data from five mice trained independently in that paradigm. **(B–D)**

Comparison of independent versus continuous training. Each panel shows one type of

count: **(B) total pokes**, **(C) other pokes**, and **(D) pellet retrievals**. Within each paradigm, two

boxplots are shown: one for mice trained independently and one for mice trained continuously.

Each box summarizes data from five mice. Statistical comparison between independent and

continuous training within each paradigm was performed using Mann–Whitney tests, with p-

values shown above each pair of boxes.

Figure 4: **Radar plots of PERCS metrics across paradigms and training types.**

(A) Independent training paradigms. Radar plots illustrate PERCS profiles derived from

mouse behavioral data obtained under independent paradigm training. The dataset includes 20

mice in total, with five mice per paradigm. Colors represent individual mice within each

paradigm. For each PERCS dimension, values were normalized such that the maximum value

across mice was set to 1, and all other values were scaled relative to this maximum. All

dimensions are therefore constrained to the range 0–1, allowing direct comparison across

dimensions and paradigms. **(B) Continuous training paradigms**. Radar plots show PERCS

profiles for mice trained under continuous paradigms. Normalization and plotting conventions

are the same as in panel A. **(C) Comparison of independent and continuous training.** Radar

plots display PERCS profiles for mice across both independent and continuous training

paradigms. Normalization and plotting conventions are identical to panels A and B, enabling

direct comparison across training types.

Figure 5: Structure and variability of mouse PERCS behavioral features across

paradigms. (A) PERCS feature dispersion. Each point represents the Euclidean distance

between an individual mouse's 5-dimensional PERCS vector (P, E, R, C, S) and the centroid of

its paradigm (FR, 2×2, 5×5, RPR). Boxes summarize the distribution of distances within each

group. Group differences were assessed using a permutation-based test for dispersion

(Anderson, 2006).**(B) Heatmap of k-means clustering.** The y-axis shows the original

paradigms (FR, 2×2, 5×5, RPR), and the x-axis shows the k-means cluster assignment (1–4) of

each mouse based on their 5-dimensional PERCS features. Each square represents the

number of mice in that combination of original paradigm and assigned cluster, with color

intensity indicating the count. **(C) Principal component analysis (PCA) of PERCS features.**

Axes represent the first three principal components (PC1, PC2, PC3) of the 5-dimensional

PERCS features across all mice. Points are colored by paradigm (FR, 2×2, 5×5, RPR). Lines

connect repeated measurements from the same mouse in continuous training, illustrating within-

mouse trajectories, whereas mice from independent training appear as single points.

References

Amsel, A. and J. Roussel (1952). "Motivational properties of frustration. I. Effect on a running

response of the addition of frustration to the motivational complex." J Exp Psychol **43**(5): 363-

366.

Aston-Jones, G. and J. D. Cohen (2005). "An integrative theory of locus coeruleus-

norepinephrine function: adaptive gain and optimal performance." Annu Rev Neurosci **28**: 403-

450.

Azrin, N. H. and R. G. Nunn (1973). "Habit-reversal: a method of eliminating nervous habits and

tics." Behav Res Ther **11**(4): 619-628.

Barkley, R. A. (1997). "Attention-deficit/hyperactivity disorder, self-regulation, and time: toward a

more comprehensive theory." J Dev Behav Pediatr **18**(4): 271-279.

Barnett, I. and J. P. Onnela (2020). "Inferring mobility measures from GPS traces with missing

data." Biostatistics **21**(2): e98-e112.

Barrett, J. M., M. G. Raineri Tapies and G. M. G. Shepherd (2020). "Manual dexterity of mice

during food-handling involves the thumb and a set of fast basic movements." PLoS One **15**(1):

e0226774.

Baumeister, R. F., E. Bratslavsky, M. Muraven and D. M. Tice (1998). "Ego depletion: is the

active self a limited resource?" J Pers Soc Psychol **74**(5): 1252-1265.

Boulos, Z. and M. Terman (1980). "Food availability and daily biological rhythms." Neurosci

Biobehav Rev **4**(2): 119-131.

Bradshaw, C. M. and P. R. Killeen (2012). "A theory of behaviour on progressive ratio

schedules, with applications in behavioural pharmacology." Psychopharmacology (Berl) **222**(4):

549-564.

Carcea, I., N. L. Caraballo, B. J. Marlin, R. Ooyama, J. S. Riceberg, J. M. Mendoza Navarro, M.

Opendak, V. E. Diaz, L. Schuster, M. I. Alvarado Torres, H. Lethin, D. Ramos, J. Minder, S. L.

Mendoza, C. J. Bair-Marshall, G. H. Samadjopoulos, S. Hidema, A. Falkner, D. Lin, A. Mar, Y. Z.

Wadghiri, K. Nishimori, T. Kikusui, K. Mogi, R. M. Sullivan and R. C. Froemke (2021). "Oxytocin

neurons enable social transmission of maternal behaviour." Nature **596**(7873): 553-557.

Carver, C. S. and M. F. Scheier (1982). "Control theory: a useful conceptual framework for

personality-social, clinical, and health psychology." Psychol Bull **92**(1): 111-135.

Carver, C. S. and M. F. Scheier (1998). On the self-regulation of behavior. New York, NY, US,

Cambridge University Press.

Chevee, M., C. J. Kim, N. Crow, E. G. Follman, M. Z. Leonard and E. S. Calipari (2023). "Food

Restriction Level and Reinforcement Schedule Differentially Influence Behavior during

Acquisition and Devaluation Procedures in Mice." eNeuro **10**(9).

Cronbach, L. J. and P. E. Meehl (1955). "Construct validity in psychological tests." Psychol Bull

52(4): 281-302.

Davis, M., K. Ressler, B. O. Rothbaum and R. Richardson (2006). "Effects of D-cycloserine on

extinction: translation from preclinical to clinical work." Biol Psychiatry **60**(4): 369-375.

de Araujo Salgado, I., C. Li, C. J. Burnett, S. Rodriguez Gonzalez, J. J. Becker, A. Horvath, T.

Earnest, A. V. Kravitz and M. J. Krashes (2023). "Toggling between food-seeking and self-

preservation behaviors via hypothalamic response networks." Neuron **111**(18): 2899-2917

e2896.

Duckworth, A. L., C. Peterson, M. D. Matthews and D. R. Kelly (2007). "Grit: perseverance and

passion for long-term goals." J Pers Soc Psychol **92**(6): 1087-1101.

Eisenberger, R. (1992). "Learned industriousness." Psychol Rev **99**(2): 248-267.

Floresco, S. B., Y. Zhang and T. Enomoto (2009). "Neural circuits subserving behavioral

flexibility and their relevance to schizophrenia." Behav Brain Res **204**(2): 396-409.

Frank, E., D. J. Kupfer, M. E. Thase, A. G. Mallinger, H. A. Swartz, A. M. Fagiolini, V.

Grochocinski, P. Houck, J. Scott, W. Thompson and T. Monk (2005). "Two-year outcomes for

interpersonal and social rhythm therapy in individuals with bipolar I disorder." Arch Gen

Psychiatry **62**(9): 996-1004.

Geurts, H. M., B. Corbett and M. Solomon (2009). "The paradox of cognitive flexibility in

autism." Trends Cogn Sci **13**(2): 74-82.

Goedhoop, J., T. Arbab and I. Willuhn (2023). "Anticipation of Appetitive Operant Action Induces

Sustained Dopamine Release in the Nucleus Accumbens." J Neurosci **43**(21): 3922-3932.

Goedhoop, J. N., B. J. G. van den Boom, R. Robke, F. Veen, L. Fellinger, W. van Elzelingen, T.

Arbab and I. Willuhn (2022). "Nucleus accumbens dopamine tracks aversive stimulus duration

and prediction but not value or prediction error." Elife **11**.

Goltstein, P. M., S. Reinert, A. Glas, T. Bonhoeffer and M. Hubener (2018). "Food and water

restriction lead to differential learning behaviors in a head-fixed two-choice visual discrimination

task for mice." PLoS One **13**(9): e0204066.

Graybiel, A. M. (2008). "Habits, rituals, and the evaluative brain." Annu Rev Neurosci **31**: 359-

387.

Graybiel, A. M. and S. L. Rauch (2000). "Toward a neurobiology of obsessive-compulsive

disorder." Neuron **28**(2): 343-347.

Hodos, W. (1961). "Progressive ratio as a measure of reward strength." Science **134**(3483):

943-944.

Insel, T. R. (2018). "Digital phenotyping: a global tool for psychiatry." World Psychiatry **17**(3):

276-277.

Jin, X., F. Tecuapetla and R. M. Costa (2014). "Basal ganglia subcircuits distinctively encode the

parsing and concatenation of action sequences." Nat Neurosci **17**(3): 423-430.

Killeen, P. R., D. Posadas-Sanchez, E. B. Johansen and E. A. Thrailkill (2009). "Progressive

ratio schedules of reinforcement." J Exp Psychol Anim Behav Process **35**(1): 35-50.

Lejuez, C. W., D. R. Hopko, R. Acierno, S. B. Daughters and S. L. Pagoto (2011). "Ten year

revision of the brief behavioral activation treatment for depression: revised treatment manual."

Behav Modif **35**(2): 111-161.

Lord, C., T. S. Brugha, T. Charman, J. Cusack, G. Dumas, T. Frazier, E. J. H. Jones, R. M.

Jones, A. Pickles, M. W. State, J. L. Taylor and J. Veenstra-VanderWeele (2020). "Autism

spectrum disorder." Nat Rev Dis Primers **6**(1): 5.

Marin, R. S. (1991). "Apathy: a neuropsychiatric syndrome." J Neuropsychiatry Clin Neurosci

3(3): 243-254.

Markowitz, J. E., W. F. Gillis, M. Jay, J. Wood, R. W. Harris, R. Cieszkowski, R. Scott, D. Brann,

D. Koveal, T. Kula, C. Weinreb, M. A. M. Osman, S. R. Pinto, N. Uchida, S. W. Linderman, B. L.

Sabatini and S. R. Datta (2023). "Spontaneous behaviour is structured by reinforcement without

explicit reward." Nature **614**(7946): 108-117.

Matikainen-Ankney, B. A., T. Earnest, M. Ali, E. Casey, J. G. Wang, A. K. Sutton, A. A. Legaria,

K. M. Barclay, L. B. Murdaugh, M. R. Norris, Y. H. Chang, K. P. Nguyen, E. Lin, A. Reichenbach,

R. E. Clarke, R. Stark, S. M. Conway, F. Carvalho, R. Al-Hasani, J. G. McCall, M. C. Creed, V.

Cazares, M. W. Buczynski, M. J. Krashes, Z. B. Andrews and A. V. Kravitz (2021). "An open-

source device for measuring food intake and operant behavior in rodent home-cages." Elife **10**.

McFarland, D. J. and R. M. Sibly (1975). "The behavioural final common path." Philos Trans R

Soc Lond B Biol Sci **270**(907): 265-293.

Mischel, W., Y. Shoda and M. I. Rodriguez (1989). "Delay of gratification in children." Science

244(4907): 933-938.

Nevin, J. A. and R. C. Grace (2000). "Behavioral momentum and the law of effect." Behav Brain

Sci **23**(1): 73-90; discussion 90-130.

Nevin, J. A. and R. C. Grace (2000). "Preference and resistance to change with constant-

duration schedule components." J Exp Anal Behav **74**(1): 79-100.

Nevin, J. A. and R. C. Grace (2005). "Resistance to extinction in the steady state and in

transition." J Exp Psychol Anim Behav Process **31**(2): 199-212.

Onnela, J. P. and S. L. Rauch (2016). "Harnessing Smartphone-Based Digital Phenotyping to

Enhance Behavioral and Mental Health." Neuropsychopharmacology **41**(7): 1691-1696.

Peca, J., C. Feliciano, J. T. Ting, W. Wang, M. F. Wells, T. N. Venkatraman, C. D. Lascola, Z. Fu

and G. Feng (2011). "Shank3 mutant mice display autistic-like behaviours and striatal

dysfunction." Nature **472**(7344): 437-442.

Podlesnik, C. A. and T. A. Shahan (2008). "Response-reinforcer relations and resistance to

change." Behav Processes **77**(1): 109-125.

Richardson, N. R. and D. C. Roberts (1996). "Progressive ratio schedules in drug self-

administration studies in rats: a method to evaluate reinforcing efficacy." J Neurosci Methods

66(1): 1-11.

Salamone, J. D., M. Correa, A. Farrar and S. M. Mingote (2007). "Effort-related functions of

nucleus accumbens dopamine and associated forebrain circuits." Psychopharmacology (Berl)

191(3): 461-482.

Sandson, J. and M. L. Albert (1984). "Varieties of perseveration." Neuropsychologia **22**(6): 715-

732.

Sayar-Atasoy, N., Y. Yavuz, C. Laule, C. Dong, H. Kim, J. Rysted, K. Flippo, D. Davis, I. Aklan,

B. Yilmaz, L. Tian and D. Atasoy (2024). "Opioidergic signaling contributes to food-mediated

suppression of AgRP neurons." Cell Rep **43**(1): 113630.

Skinner, B. F. (1988). "The operant side of behavior therapy." J Behav Ther Exp Psychiatry

19(3): 171-179.

Squire, L. R. and S. M. Zola (1996). "Structure and function of declarative and nondeclarative

memory systems." Proc Natl Acad Sci U S A **93**(24): 13515-13522.

Thorndike, E. L. (1933). "A Proof of the Law of Effect." Science **77**(1989): 173-175.

Tinbergen, N. (1972). "Functional ethology and the human sciences." Proc R Soc Lond B Biol

Sci **182**(1069): 385-410.

Treadway, M. T. and D. H. Zald (2011). "Reconsidering anhedonia in depression: lessons from

translational neuroscience." Neurosci Biobehav Rev **35**(3): 537-555.

Wang, Y. and Q. Q. Sun (2024). "A prefrontal motor circuit initiates persistent movement."

bioRxiv.

Weinreb, C., J. E. Pearl, S. Lin, M. A. M. Osman, L. Zhang, S. Annapragada, E. Conlin, R.

Hoffmann, S. Makowska, W. F. Gillis, M. Jay, S. Ye, A. Mathis, M. W. Mathis, T. Pereira, S. W.

Linderman and S. R. Datta (2024). "Keypoint-MoSeq: parsing behavior by linking point tracking

to pose dynamics." Nat Methods **21**(7): 1329-1339.

Wiltschko, A. B., M. J. Johnson, G. Iurilli, R. E. Peterson, J. M. Katon, S. L. Pashkovski, V. E.

Abraira, R. P. Adams and S. R. Datta (2015). "Mapping Sub-Second Structure in Mouse

Behavior." Neuron **88**(6): 1121-1135.

Wykes, T., V. Huddy, C. Cellard, S. R. McGurk and P. Czobor (2011). "A meta-analysis of

cognitive remediation for schizophrenia: methodology and effect sizes." Am J Psychiatry **168**(5):

472-485.

Supplementary Figures

*Supplemental Figure 1: **Example of a Gaussian mixture model for instantaneous poking frequency.***

The histogram shows the distribution of instantaneous poke frequency for a single mouse as an illustrative example. A Gaussian mixture model was applied to this distribution. The solid curves in blue and red represent the estimated normal components fitted to the histogram. The dashed vertical lines indicate the mean of each estimated normal distribution.

Supplemental Figure 2

Supplemental Figure 2: ***Instantaneous frequency of pellet retrieval.***

For each paradigm, the instantaneous pellet retrieval frequency was calculated as the reciprocal of the interval between consecutive pellet retrievals. Each dot represents a single pellet retrieval plotted over time. The panels are arranged from top to bottom corresponding to the four paradigms (FR, 2×2, 5×5, and RPR).