A macrocyclic peptide-based fusion inhibitor targeting SARS-CoV-2 Spike S2 subunit

Hirofumi Ohashi[1#], Tatsuro Kawamura[2#], Masaki Ohuchi[2], Haruaki Kurasaki[2], Naoko Iwata-Yoshikawa[3], Yuichiro Hirata[3], Saya Moriyama[4], Kaho Shionoya[1], Kazutaka Nagatomo[2], Takayuki Nagasawa[2], Junpei Yamamoto[2], Kei Sudo[2], Naoko Nakamura[2], Katsuma Matsui[2], Haruhiko Ogawa[2], Kouji Yoshida[2], Yoshiaki Shimada[2], Tomoko Maruyama[2], Tomoaki Higuchi[2], Shoko Ito[2], Yoshimasa Takahashi[4], Naoki Kawamura[2], Patrick C. Reid[2], Masato Murakami[2,5], Tadaki Suzuki[3,6], Noriyo Nagata[3*], Hidetomo Kitamura[2*], Koichi Watashi[1*]

[1]Department of Drug Development, National Institute of Infectious Diseases, Japan Institute for Health Security, Japan

[2]PeptiDream Inc., Japan

[3]Department of Infectious Disease Pathology, National Institute of Infectious Diseases, Japan Institute for Health Security, Japan

[4]Research Center for Vaccine Development, National Institute of Infectious Diseases, Japan Institute for Health Security, Japan

[5]PeptiAid Inc., Japan

[6]Department of Infectious Disease Pathobiology, Graduate School of Medicine, Chiba University, Japan

[#]These authors contributed equally: Hirofumi Ohashi, Tatsuro Kawamura.

*Corresponding authors:
E-mail: nagata.no@jihs.go.jp (N.Nagata), hidetomo-kitamura@peptidream.com (H.Kitamura), watashi.k@jihs.go.jp (K.W.)

Abstract

Continuous emergence of SARS-CoV-2 variants carrying mutations in Spike presents a significant challenge for durable antiviral agents.   Here we screen for random 13-amino acid non-mimetic macrocyclic peptides that bind to Spike and identify PA-001 that inhibits SARS-CoV-2 infection with high potency at sub-nanomolar as 50% inhibitory concentration (IC_{50}).   PA-001 bound to Spike S2 subunit and inhibited the membrane fusion during virus entry.   Through drug-resistant selection, we revealed that PA-001 targeted the fusion peptide proximal region (FPPR) in S2, which has not been recognized as a drug target to date.   Consistent with its highly conserved amino acid sequences beyond strains, PA-001 exhibited broad antiviral activity against all tested SARS-CoV-2 variants, in contrast to clinically-approved S1-targeting antibodies that lost activity to Omicron variants.   PA-001 suppressed SARS-CoV-2 propagation and disease progression in mouse- and hamster-infection models, both by administration prophylactically and therapeutically.   Combination therapy with remdesivir further enhanced antiviral profiles.   In clinical phase-I trial, PA-001 was well-tolerated and showed high systemic exposure, with over 28,000-fold concentration of IC_{50} as maximum plasma concentration by single administration to healthy volunteers.   These evidence propose FPPR as an unexpected antiviral drug target accessible by macrocyclic peptides and identify PA-001 as a potent anti-SARS-CoV-2 fusion inhibitor.

Introduction

Coronavirus disease 2019 (COVID-19), caused by infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has had a profound impact on global public health, resulting in more than 770 million cumulative cases and over 7 million deaths worldwide (https://covid19.who.int/). Despite the development and widespread deployment of vaccines and antiviral drugs, COVID-19 continues to impose a substantial disease burden, especially among people including older adults, individuals with underlying medical conditions, and immunocompromised populations[1-3]. Throughout and beyond the pandemic period, SARS-CoV-2 has exhibited continuous viral evolution, given rise to numerous variants characterized by the accumulated mutations[4]. These genetic changes have altered viral pathogenicity and transmissibility and, critically, reduced susceptibility to approved antiviral drugs. There is a need to develop novel antiviral agents sustainably retaining their efficacy against ongoing emergence of SARS-CoV-2 variants.

The Spike glycoprotein of SARS-CoV-2 mediates viral entry to host cells through receptor binding and membrane fusion. Spike constitutes a trimeric precursor and comprises two functional subunits, S1 and S2. S1 consists of N-terminal domain (NTD), receptor binding domain (RBD), and C-terminal domains (CTDs), and is essential for recognition of host entry receptor, angiotensin converting enzyme 2 (ACE2), through its RBD[5,6]. S2 contains multiple functional domains including fusion peptide (FP) and its proximal region (FPPR), heptad repeat 1 (HR1), central helix (CH), connector domain (CD), heptad repeat 2 (HR2), and transmembrane domain (TMD) (see Fig. 2E)[4,6]. Upon RBD in the S1 subunit engagement to ACE2, S2 is processed at its N-terminal S2′ site by host proteases, either transmembrane serine protease 2 (TMPRSS2) on the cell surface or cathepsin L in the endosome, which then induces large-scale conformational rearrangements of S2 in the membrane fusion process (Supplementary Fig. 1)[6-12]: The S1-ACE2 complex dissociates from the S2 trimer (Supplementary Fig. 1a-c), and the original conformation (pre-fusion form) of S2 is converted to forms a long central coiled coil with blunted cone-shaped structure (involving FP and the adjacent FPPR) at its top, which allows insertion into the target cell membrane (Intermediation form; Supplementary Fig. 1d). It subsequently refolds into the stable structure by interaction of HR1 and HR2 to form a six-helix bundle that constitutes a rigid, rod-like shape (Post-fusion form; Supplementary Fig. 1e-f) to undergo fusion of viral envelope to cellular membranes.

This viral entry process is an attractive antiviral target [13]. The clinically-approved antibodies for COVID-19 treatment, such as casirivimab, imdevimab, sotrovimab, tixagevimab, cilgavimab, and bebtelovimab, recognize the S1 subunit and are designed to strongly block the S1 RBD-ACE2 binding[14,15]. However, their neutralization activities have been lost by accumulated mutations, especially in RBD, in the recent Omicron subvariants such as BQ.1 and XBB[16]. Other S1-targeting antibodies, including those targeting NTD, also face the similar challenges for emerging escape variants due to high sequence variability in this region. Given its highly conserved

sequence among circulating SARS-CoV-2 variants, S2 is expected as an alternative target for broadly active antivirals.   So far, S2-targeting antibodies have been identified, including those recognizing the region downstream of the S2′ cleavage site (used to be referred as FP, but is presently defined an intermediate loop) and the stem helix at the upstream of HR2[17-19], but their neutralizing activities were typically low compared to those targeting RBD in S1.   So far, peptides mimicking the HR2 sequence aiming to inhibit HR1-HR2 interaction during the conversion to the post-fusion form have been under clinical development.   These peptides have shown broad anti-SARS-CoV-2 activity, but their original activities of pre-optimized peptides were 500-3,000 nM ranges as 50% inhibitory concentrations (IC_{50})[18,20].   Later, these S2-mimicking linear peptides have been optimized for improving their activities through polyethylene glycol or cholesterol conjugations.   These antibody- and linear peptide-based approaches face a challenge in which their target site as well as antiviral activities are limited due to the S2's inherent characteristics including limited surface area accessibility by large molecules as well as low immunogenicity and high flexibility.

Cyclic peptide-based drugs are expected to provide an alternative approach[21] that can have drug-like properties and advantages including accessibility to narrow target sites, high specificity, tunable stability and efficacy, low immunogenicity, and relatively less cost compared to antibodies. In this study, we show that a non-mimetic macrocyclic peptide, PA-001, inhibits infection of broad SARS-CoV-2 variants via targeting FPPR within the S2 subunit, which is highly conserved in sequence but previously undescribed as a druggable site.   By using SARS-CoV-2-inoculated mouse and hamster models, PA-001 suppressed viral propagation and ameliorated disease progression under administrations either prophylactically or therapeutically.   A first-in-human Phase I dose-escalation trial demonstrated a favorable safety profile with high systemic exposure in pharmacokinetics (achieving maximum plasma concentration of over 28,000-fold concentration of in vitro IC_{50} against Omicron strains) in healthy volunteers.   Our findings present PA-001 as an anti-SARS-CoV-2 drug candidate and that FPPR is a potential druggable target for SARS-CoV-2 fusion inhibitor.

Results

Identification of an anti-SARS-CoV-2 cyclic peptide, PA-001

To explore novel COVID-19 therapy targeting SARS-CoV-2 Spike, we discovered and developed anti-SARS-CoV-2 agents following the scheme as summarized in Fig. 1A.   Through screening of a library consisting of ~10^{13} random peptides using Peptide Discovery Platform System (PDPS), which we have established so far[22-24] (Fig. 1B), and the following rational optimization process through medicinal-chemistry program, we developed PA-001 (Fig. 1C) as an investigational agent currently in clinical development as a therapy against COVID-19.   In the present study, we report the in vitro and in vivo antiviral profile of PA-001.

Peptides that were identified to bind Spike in vitro were optimized to improve antiviral potency

and pharmacokinetics (Fig. 1A).   We examined i) affinity with Spike in vitro (Supplementary Table

1), ii) inhibition activity to SARS-CoV-2 infection in a cell-based assay (Fig. 1D), and iii)

pharmacokinetics profiles in mice (Fig. 1E).   In surface plasmon resonance, PA-001 had high

affinity to recombinant Spike, with 1.3 nM as a K_D (Supplementary Table 1).   Activity of peptides

to inhibit SARS-CoV-2 infection was examined in a cell-based SARS-CoV-2 infection assay:

VeroE6/TMPRSS2 cells[25] were inoculated with a SARS-CoV-2 clinical isolate (Wk-5.2.1, Wuhan

type) at a multiplicity of infection (MOI) of 0.003 for 1 h and were cultured in the presence or

absence of peptides for 48 h to assess SARS-CoV-2-induced cytopathology (Fig. 1D).   The cell

line exhibited robust cytopathology by SARS-CoV-2 propagation at 48 h after infection, which was

however completely blocked by treatment with 1.0 μM PA-001, as 10 μM remdesivir (RDV), a

clinically-approved anti-SARS-CoV-2 polymerase inhibitor[26] used as a positive control (Fig. 1D-

b).   PA-001 reduced the production of SARS-CoV-2 proteins, Spike and N as representatives

detected by immunoblotting, at 24 h post-inoculation, the time before showing cytopathic effect

(Fig. 1D-c).   PA-001 reduced viral RNA at 24 h post-inoculation in a dose-dependent manner (Fig.

1D-d).   Pharmacokinetics of a single intraperitoneal administration with PA-001 at 10 mg/kg to

BALB/c mice showed 76,667 ng/mL (41,500 nM) as maximum plasma concentration (C_{max}) and a

gradual decline but sustained plasma concentration, with 1.0 h as half-life, and 282,236 ng·h/mL

(153,000 nM·h) as area under the curve (AUC_{0-24}) in plasma (Fig. 1E, Supplementary Table 2).

From these data showing the strong potential of PA-001 as an anti-SARS-CoV-2 agent, we

focused on PA-001 in the following analysis.

PA-001 targets S2 and inhibits Spike-dependent membrane fusion

We determined the target of PA-001 by in vitro binding assay that incubated PA-001 at varying

concentrations with recombinant either full-length (S1+S2), S1, or S2 protein as targets.   As

shown in Fig. 2A, PA-001 presented the binding signal with the full-length Spike and S2 in a dose-

dependent manner, but not with S1 at any treated concentrations up to 100 nM (Fig. 2A),

suggesting that PA-001 binds to S2.   In the SARS-CoV-2 life cycle, cell entry is initiated by viral

attachment to cells through binding of S1 RBD to ACE2 on cell surface (Fig. 2B, red); This is

followed by the fusion of viral envelope with cellular membranes either on cell surface or at

intracellular vesicles, triggered by cleavage of the S2 internal site (S2′ site) by cellular proteases

(Fig. 2B, brown).   To confirm the target step of PA-001 during the entry process, we separately

assessed the activity of viral attachment to cell surface (Fig. 2C-a) and membrane fusion (Fig.

2C-b).   Attachment assay was performed by incubating VeroE6/TMPRSS2 cells with SARS-CoV-

2 at 4°C to allow viral-cell attachment but not the following processes, followed by quantification

of viral RNA on the cell surface.   Cell-attached SARS-CoV-2 RNA was dose-dependently

reduced upon treatment with casirivimab + imdevimab (CAS + IMD) cocktail antibodies, clinically-

approved SARS-CoV-2 neutralizing antibodies[27] used as a positive control, whereas that was not

significantly affected by treatment with PA-001, as with mefloquine, a reported post-attachment

entry inhibitor[28] used as a negative control (Fig. 2C-a).   On the other hand, in the cell fusion

assay, which coincubated Spike-overexpressing HEK293T cells with endogenously ACE2-

expressing VeroE6/TMPRSS2 cells to induce cell membrane fusion, essentially as previously

described[29], PA-001 clearly reduced Spike-ACE2-induced fusion in a dose-dependent manner,

and its IC_{50} was calculated to be 3.6 nM (Fig. 2C-b).   These results are consistent with that PA-

001 targets S2, which mediates the membrane fusion process.

To identify the S2 target site responsible for the PA-001 sensitivity, we isolated PA-001-resistant

SARS-CoV-2 variants by continuous exposure of SARS-CoV-2-infected cells with PA-001 and

serial passages, essentially described previously[30] (Fig. 2D-a):   VeroE6/TMPRSS2 cells infected

with SARS-CoV-2 were continuously treated with PA-001 and their culture supernatant was

reinfected to naive VeroE6/TMPRSS2 cells.   After seven passages, we collected SARS-CoV-2

in the culture supernatant from eight independent cell batches (A8, B8, C8, D8, E8, F8, G8, and

H8).   We then examined the sensitivity of these virus clones to PA-001, as well as RDV as a

reference, by inoculating the cells with these viruses or the parental wild type SARS-CoV-2 and

treated with PA-001 or RDV at varying concentrations to quantify viral RNA produced into the

culture supernatant.   As shown in Fig. 2D-b, all the virus clones (A8-H8) showed no or limited

reduction in viral RNA levels by treatment with PA-001 up to 500 nM, in contrast to the drastic

reduction in RNA of the parental SARS-CoV-2 (Fig. 2D-b left, and Supplementary Fig. 2A-a).   In

contrast, all these variants remained sensitive to RDV, as the parental SARS-CoV-2 (Fig. 2D-b

right, Supplementary Fig. 2A-b).   The calculated 90% inhibitory concentrations (IC_{90}s) to PA-001

for all the clones were >500 nM (v.s. 8.4 nM for the parental virus), whereas those to RDV for all

clones remained 3,000-4,500 nM that were equivalent to the parental virus (Supplementary Fig.

2B), indicating that all the isolated variants acquired specific resistance to PA-001.

Next-generation sequencing of viral RNA derived from these eight clones identified 1-3 amino

acid substitutions for each clone, with enriched substitutions in the S2 region (Supplementary Fig.

2C), consistent with that PA-001 targets S2.   Interestingly, D8 and H8 had the identical and single

mutation that caused amino acid substitution, L841R in S2 protein (Fig. 2E-a, Supplementary Fig.

2C), suggesting that the L841R substitution is responsible for acquiring the PA-001 resistance.

Similarly, E8 and G8 possessed the identical but another single amino acid substitution, P862Q

in S2 (Fig. 2E-a, Supplementary Fig. 2C).   Remarkably, all the remaining clones also acquired a

substitution cluster at the neighboring amino acids, Q836 (in F8), Y837 (in A8 and F8), G838 (in

B8), and L858 (in C8) (Fig. 2E-a highlighted by pink, Supplementary Fig. 2C).   All these amino

acids reside at the upstream of the fusion peptide (FP: 866-910 aa) in S2, referred as the fusion

peptide proximal region (FPPR)[4,6-9] (Fig. 2E-a).   Based on the reported three-dimensional

structure of Spike protein at pre-fusion, intermediation, and post-fusion forms[10-12], FPPR

dynamically moves and changes its structure, together with FP, during the conversion from the

pre-fusion to the post-fusion form[4]: FPPR is originally exposed on the surface in a concave area

of the central part of the pre-fusion trimer, and undergoes a dynamic locational change to the top
of S2, forming the cone-shaped structure to assist FP insertion into target membrane in the post-
fusion form[11] (Fig. 2E-b).

As the above critical amino acids, Q836, Y837, G838, L841, L858, and P862 are exposed on
the surface of the Spike pre-fusion trimer, we assume these amino acids constitute the interface
to PA-001 binding, of which amino acid substitutions lead to escape from PA-001 binding.   We
then examined the binding activity of recombinant Spike having a substitution of either L841R (in
D8 and H8) or P862Q (in E8 and G8) to PA-001 by ELISA: Spike proteins were immobilized on
the well, and incubated with varying amount of HA-tagged PA-001, which was then visualized by
adding anti-HA-conjugated horseradish peroxidase (HRP).   Specific Spike-PA-001 binding was
evaluated by the difference in the signal from that under coincubation with and without excess
amount of non-HA-tagged PA-001 as a competitor (Fig. 2F-a).   This specific Spike-PA-001
binding was observed along with the increasing amount of PA-001-HA by using the wild type Spike,
however, this signal completely disappeared for Spike carrying the substitution at either L841R or
P862Q (Fig. 2F-b), suggesting that these amino acid substitutions abrogate PA-001 binding.
Consistently, PA-001's dose-dependent inhibition of the Spike-ACE2-mediated cell membrane
fusion was totally cancelled up to 1,000 nM by mutations of L841R or P862Q in Spike (Fig. 2G).
These data cumulatively suggest that PA-001 binds to FPPR to inhibit S2-dependent membrane
fusion.

PA-001 shows antiviral activity to broad range of SARS-CoV-2 variants
Multiple neutralizing antibodies have been approved for clinical use, however, all of them have
lost antiviral efficacy by amino acid substitutions in Spike observed in naturally-occurring SARS-
CoV-2 variants[14,31].   Most of these substitutions were concentrated in the S1 subunit.   To clarify
the breadth of the antiviral activity of PA-001, we infected VeroE6/TMPRSS2 cells with SARS-
CoV-2 Wuhan, Alpha, Beta, Gamma, Delta, and Omicron (BA.1.18 and XBB.1.16), and treated
with varying concentrations of PA-001 as well as CAS + IMD as a reference to quantify viral RNA
in the culture supernatant.   As previously reported, CAS + IMD showed antiviral effects on
Wuhan to Delta variants, but not Omicron variants (Fig. 3A, lower graphs), because of the amino
acid substitutions in RBD of S1[32,33].   In contrast, PA-001 reduced SARS-CoV-2 infection for all
the variants examined including Omicron variants, BA.1.18 and XBB.1.16, in a dose-dependent
manner (Fig. 3A, upper graphs).   Anti-SARS-CoV-2 activities of PA-001 against all these variants
ranged 0.34-3.2 nM as IC_{50}s and 1.7-9.6 nM as IC_{90}s, in contrast to the complete loss of activity
of CAS + IMD against BA.1.18 and XBB.1.16 (IC_{50} and IC_{90} >1,000 ng/mL) (Supplementary Table
3).   Furthermore, PA-001 showed the inhibitory activity to the recent Omicron subvariants, JN.1
and XFG, in contrast to loss of activity of CAS + IMD, examined by infection of VSV-based
pseudovirus carrying Spike derived from each SARS-CoV-2 variant, as previously described[34,35]
(Supplementary Fig. 3A).   Consistently, the amino acid sequences of all these variants from 834

to 863 aa are almost completely conserved, except for a single non-essential substitution at
position 856 only in BA.1.18 (Fig. 3B, Supplementary Fig. 3B).　Thus, PA-001 showed an
advantage that was effective on broad range of naturally-occurring SARS-CoV-2 variants, in clear
contrast to the clinically-approved anti-S1 antibodies.
242

PA-001 inhibits SARS-CoV-2 propagation and disease progression in vivo by either
pretreatment or posttreatment after virus infection
We assessed the antiviral effect of PA-001 in infection animal models.　We initially examined
the effect of prophylactic treatment with PA-001 before virus inoculation on propagation of
infectious SARS-CoV-2 in a Syrian hamster infection model[36,37] (Fig. 4A-a);　Syrian hamsters
were pretreated with PA-001 at 0.3, 1.0, or 3.0 mg/kg, or saline as a vehicle control
intraperitoneally twice [at 12 and 2 h before virus inoculation (-0.5 and -0.08 day)], followed by
intranasal inoculation with SARS-CoV-2 (Delta variant), and then posttreatment with PA-001 at
the same concentrations three times [at 12, 24, and 36 h (0.5, 1, and 1.5 day)].　Hamsters were
sacrificed at day 2 for collecting lungs to quantify the infectious titer of SARS-CoV-2.　SARS-
CoV-2 infectious titers detected in these tissues derived from vehicle-administered hamsters were
significantly reduced by PA-001 treatment in a dose-dependent manner, by up to 3.5-log (at 3.0
mg/kg of PA-001) (Fig. 4A-b).　These data indicate the potential anti-SARS-CoV-2 effect of PA-
001 in hamsters by prophylactic treatment.
To further evaluate the therapeutic efficacy by posttreatment after infection, we examined the
pathogenicity induced by SARS-CoV-2 propagation in mice upon posttreatment with PA-001 after
infection; BALB/cCrSlc mice intranasally infected with a mouse-adapted SARS-CoV-2 (QHmusX
strain)[38] were treated intraperitoneally with PA-001 at 10 or 30 mg/kg, saline as a vehicle control,
or CAS + IMD as a positive control at 1, 8, 24, 48, 72, and 96 h after virus inoculation (0.04, 0.3,
1, 2, 3, and 4 days), and were chased to measure body weight as well as survival rate daily up to
day 10 (Fig. 4B-b, c).　Seven mice out of eight treated with the vehicle were dead or reached the
humane endpoint (survival rate = 12.5%) at day 6, and body weights of vehicle-treated mice were
gradually decreased following SARS-CoV-2 infection to 73-80% of the original until day 5 (Fig. 4B-
b, c).　In contrast, all mice treated with PA-001 after virus inoculation at either 10 or 30 mg/kg
survived (survival rate = 100%), whereas one mouse was dead upon CAS + IMD treatment
(survival rate = 87.5%) (Fig. 4B-b).　Furthermore, PA-001-treated group showed only a transient
body weight reduction by up to around 11 % at day 3, followed by a recovery to the original body
weight level at day 10, as the case with CAS + IMD-treated mice (Fig. 4B-c).
We further examined histopathology in the lung as well as body weight and SARS-CoV-2 viral
load in the lung in Syrian hamsters posttreated with PA-001 after SARS-CoV-2 infection (Fig. 4C);
Hamsters after intranasal SARS-CoV-2 infection were posttreated with PA-001 (1 or 5 mg/kg) or
saline at 2, 8, 24, 48, and 72 h and were collected at day 5 to analyze histopathology and viral
RNA levels in the lung as well as chased for measuring body weight up to day 5 (Fig. 4C-a).

Body weight loss induced by SARS-CoV-2 propagation (to 88% in the vehicle-control group) was
significantly recovered by PA-001 posttreatment (to 94% in PA-001-treated group) (Fig. 4C-b).
SARS-CoV-2 RNA levels in the lung were reduced by PA-001 treatment in a dose-dependent
manner by up to 1.6-log reduction (by 5.0 mg/kg of PA-001) (Fig. 4C-c).   Histopathological
examination revealed severe pneumonia with consolidation from vehicle-treated hamsters,
whereas such severe illness in histopathology was improved by PA-001 treatment, especially at
5 mg/kg (Fig. 4C-d).   Thus, PA-001 inhibited SARS-CoV-2 propagation in the lung and
suppressed the resultant disease progression by post-treatment after infection in mice and
hamsters.
285

Combination treatment with PA-001 and remdesivir augments the antiviral effects
Combination treatment with multiple drugs with different mode of action generally enhances
the antiviral effect and decreases the risk for selecting drug-resistant variants[39].   We then
examined the combination treatment of PA-001 with RDV at various concentrations to evaluate
the antiviral effects in cell-based infection assay.   As shown in Fig. 5A, RDV reduced SARS-CoV-
2 production in a dose-dependent manner by single treatment, and this RDV's antiviral effects
were enhanced by addition of PA-001 (Fig. 5A): e.g., RDV at 1,500 nM and PA-001 at 0.70 nM
single treatment reduced viral production to 51% and 43%, respectively, but combination with
RDV at 1,500 nM plus PA-001 at 0.70 nM decreased viral production to 11% of the control (Fig.
5A), indicating that PA-001 potentiated the antiviral activity of RDV.
We next examined the effect of combination therapy with PA-001 and GS-441524 (RDV-m),
the active metabolite of RDV[40], in SARS-CoV-2 infection mouse model.   BALB/cCrSlc mice
intranasally inoculated with a mouse-adapted SARS-CoV-2 strain were intraperitoneally treated
with RDV-m (25, 12.5, 12.5 mg/kg), RDV-m (25, 12.5, 12.5 mg/kg) + PA-001 (26 mg/kg), or vehicle
control (Fig. 5B-a).   The vehicle-control group mice showed a continuous decrease in body
weight and all the mice died or reached the humane end point at day 5 after infection (Fig. 5B-b,
c, black).   In contrast, body weight loss of mice treated with RDV-m alone was only transient until
day 3 to 4, followed by a gradual recovery, with all the mice survived until day 10 (Fig. 5B-b, c,
blue).   Consistently, virus infectious titer in the lung was significantly decreased by 1.2-log,
although the lung weight, as a marker of tissue damage and inflammation was not significantly
improved by single treatment with RDV-m (Fig. 5B-d, e).   Combination treatment with RDV-m
and PA-001 more rapidly recovered the body weight loss at day 4 to 7 compared with RDV-m
single treatment, with no death during the whole experimental period (Fig. 5B-b, c, red), and
showed additional reduction in SARS-CoV-2 infectious titer in the lung (by 1.8-log) (Fig. 5B-d, red).
Furthermore, lung weight increase by SARS-CoV-2 was significantly improved by RDV-m + PA-
001 cotreatment (Fig. 5B-e, red).
We further examined the production profile of cytokines and chemokines, especially those
reportedly associated with the COVID-19 severe disease progression, including interleukin (IL)-6,

tumor necrosis factor-alpha (TNF-α), IL-1β, granulocyte colony stimulating factor (GCSF), interferon (IFN)-γ, IFNγ-induced protein 10 (IP10), macrophage inflammatory protein 1-alpha (MIP-1α), and monocyte chemoattractant protein-1 (MCP1)[41-44], in both the serum and the lung of mice at day 3 postinfection.   As shown in Fig. 5C, SARS-CoV-2-induced production of IL-6, TNF-α, IL-1β, IP-10, MIP-1α, and MCP-1 at least either in the serum or the lung was significantly inhibited by RDV-m monotreatment, and most of these expressions were more remarkably reduced by combination treatment with RDV-m and PA-001 (Fig. 5C).   Furthermore, induction of IP-10 in serum was significantly reduced in PA-001 and RDV-m combination treatment from RDV-m single treatment; Production of TNF-α and IFN-γ in the serum was significantly reduced under only combination treatment with PA-001 and RDV-m, but not single treatment with RDV-m (Fig. 5C).   Thus, combination treatment with PA-001 potentiated the anti-SARS-CoV-2 effect of RDV and the resultant disease progression.

Pharmacokinetics and safety of PA-001 in healthy volunteers

A double-blind, randomized, placebo-controlled phase I study for PA-001 was conducted in 2 parts (Part A and B) to examine the safety, tolerability, and pharmacokinetics of PA-001 in healthy volunteers.

Part A was a single-dose sequential-group study with a total of 32 randomized participants. Baseline characteristics were balanced across cohorts, in which mean age of participants was 47 years (range 22–75 years) and the majority of participants were female (59.4%), white (65.6%), and not Hispanic or Latino (62.5%), with a mean BMI of 26.17 kg/m^2 (Fig. 6A).   These volunteers were grouped into five cohorts who were administered intravenously with a single administration of either i) placebo (n = 8), ii) 18 mg PA-001 (n = 6), iii) 64 mg PA-001 (n = 6), or iv, v) 128 mg PA-001 (n = 6 to younger and n = 6 to elderly).   Administration with PA-001 at 128 mg had two cohorts, one with younger (median age of 37.2 years, iv) and one with elder volunteers (median age of 69.8 years, v).   These cohorts were chased to observe adverse effects as well as to collect the plasma in a time course up to 48 h post-administration for measuring the concentration of PA-001.   As a result, PA-001 was well-tolerated, showing no severe adverse effects related with treatment (treatment emergent adverse effects, TEAEs) in all individuals (Fig. 6B and Supplementary Table 4A).   Observed events were mild, including headache (1 subject in Placebo, 2 subjects in 64 mg PA-001) and vomiting (1 subject in 64 mg PA-001) (Fig. 6B and Supplementary Table 4A).   Pharmacokinetics profiles were presented with plasma concentrations against the time course up to 48 h after single dosing (Fig. 6C), and pharmacokinetic parameters of PA-001 at each dose calculated based on these profiles (Fig. 6D and Supplementary Table 5).   Dose proportional increases in the peak concentrations and exposures of PA-001 were observed across the evaluated dose range (Fig. 6C).   PA-001 was readily detectable in plasma at 0.5 h after administration with an average maximal concentrations (C_{max}) of up to 18,300 ng/mL (9,610 nM) at 128 mg treatment, and was gradually declined, with 3.61-4.79 h as half-life ($T_{1/2}$).   Given the

IC_{50} of 0.34 nM in the cell-based infection assay against the Omicron strains, BA.1.18 and

XBB.1.16 (Supplementary Table 3), 128 mg PA-001 group exhibited more than 28,000-fold

concentration of IC_{50} in plasma at peak (C_{max}) at 1 h after administration, and sustained more than

340-fold concentration of IC_{50} at 24 h post-administration (Fig. 6C). The overall exposure of PA-

001 ($AUC_{0\text{-tlast}}$) was up to 74,200 ng·h /mL at 128 mg administration (younger).

Part B was a multi-dose sequential-group study, in which total 15 participants were randomized

into three groups who received multiple dosing of either placebo (n = 4), 64 mg PA-001 (n = 5), or

128 mg PA-001 (n = 6) once a day until day 5 (Supplementary Table 6). The mean age of

participants was 37 years (range 24–60 years); Eight (53.3%) males and seven (46.7%) females,

with a mean BMI of 27.29 kg/m^2. The majority (66.7%) of subjects were White (Supplementary

Table 6). Pharmacokinetics parameters on day 1 and day 5 (Supplementary Table 7) were

similar to those of single dosing in part A (Fig. 6D and Supplementary Table 5). Following

multiple doses of 64 and 128 mg for 5 days, $AUC_{0\text{-}\tau}$ and C_{max} increased in a supra proportional

manner with respect to dose. The AR_{AUC} (Accumulation ratio based on $AUC_{0\text{-}\tau}$) was 1.01 for the

64 mg dose level and 1.03 for the 128 mg dose level (Supplementary Table 7B). Because there

was no accumulation, steady state concentrations of PA-001 were achieved from the onset of the

multiple dosing period. No serious adverse effects related with multi-dosing of PA-001 was

observed, although seven (46.7%) subjects had at least 1 mild TEAE, of whom 2 received 64 mg

PA-001 (Grade 1), 4 received 128 mg PA-001 (3 of Grade 1, 1 of Grade 2), and 1 received placebo

(Grade 1) (Supplementary Table 4B). Three (20.0%) subjects had Grade 1 TEAE possibly

related to study treatment (2 received 64 mg, 1 received 128 mg), but all these TEAEs had

resolved by the end of the study (Supplementary Table 4B). There were no deaths, serious

adverse effects, or discontinuations because of adverse effects.

Thus, PA-001 provided the sustained plasma concentration in human individuals above those

required for antiviral activity, with preferable safety profiles.

Discussion

In this study, we identified a 13-amino acid macrocyclic peptide, PA-001, specifically binding to

FPPR on the S2 subunit of the SARS-CoV-2 Spike protein, which has not been targeted by

antibodies or antiviral agents so far. Despite extensive efforts of cryo-EM analysis, we were

unable to obtain a high-resolution structure of the PA-001/Spike complex, probably due to the

characteristic of FPPR, having high conformational flexibility, as mentioned below[7,45,46].

Alternatively, to demonstrate the binding site, we created SARS-CoV-2 variants more than 100-

fold resistance to PA-001 and found an amino acid substitution cluster within residues 836-862,

which resides at FPPR, upstream of FP. Mutations in this region completely abolished Spike's

capacity to bind to PA-001, confirming this segment as the PA-001-binding site. In the pre-fusion

form, FPPR is positioned on the surface of Spike trimer between the convex CTD1 and CTD2,

forming a shallow depression at the surface.   FPPR is suggested to control the RBD movement

between its up and down conformations[4]: FPPR is structured and clamps RBD in the down

conformation of the pre-fusion form through interaction of Asp614 (in RBD) with Lys854 (in FPPR),

whereas it moves out and unstructured in the RBD down conformation[4,47-49].   Upon ACE2-RBD

engagement and the following S1 release, FPPR, together with the adjacent FP, undergoes an

overall translocation during the conformational rearrangement to be positioned at the top of the

S2 trimer and form a cone-shaped structure, which functions to be inserted into the target cellular

membrane.   The binding of PA-001 to FPPR at the pre-fusion form (Fig. 2A and F) would

sterically interfere with this mobility of FPPR toward the formation of the cone-shaped structure

(intermediation form; Supplementary Fig. 1) that is required for the fusion process.   These

findings highlight FPPR as a previously unexplored but potential target for antiviral drug

development.

Previously developed Spike-targeting therapeutics largely act through antibodies directed

against S1 such as casirivimab and imdevimab, amubarvimab and romlusevimab, bamlanivimab,

bamlanivimab and etesevimab, sotrovimab, tixagevimab and cilgavimab, and bebtelovimab[14,15].

These antibodies primarily exert anti-SARS-CoV-2 activity by neutralization of infection through

blocking Spike-ACE2 interaction and/or by engaging Fc-mediated effector functions that promote

viral clearance.   However, their neutralization activity has been severely impaired or completely

lost by accumulated mutations in RBD of the S1 subunit among the recent Omicron sublineages.

The NTD in S1, another target of antibodies, also has high sequence variability.   In contrast, the

S2 region is more conserved in sequence and represents a target for developing broadly active

antivirals.   S2-targeting antibodies reported so far mainly recognize the region downstream of

the S2′ site (residues 811-825, previously referred to as FP, but recently considered an

intermediate loop) or the stem helices upstream of HR2, however, they had relatively modest

antiviral potency, likely due to the limited surface accessibility by antibodies as well as low

immunogenicity[17,50,51].   To overcome such limitations, non-antibody modalities have been

applied so far, such as peptides and aptamers, which have smaller molecular size that can bind

to regions inaccessible to conventional antibodies.   HR2-mimetic peptides, such as EK1 and

HY3000, designed to interfere with the HR1-HR2 interaction during the conformational change to

the post-fusion form, have shown broad antiviral activity across SARS-CoV-2 variants, with

reported original IC_{50} at 500-3,000 nM, and their activities were further optimized through the lipid

conjugation[17,18].   S2-targeting DNA aptamers have also been reported, though most of their

antiviral activities in infection models remains to be characterized[52].   Among all these S2-directed

agents, PA-001 exhibited one of the most potent anti-SARS-CoV-2 activities reported to date (IC_{50}

= 0.34 − 3.2 nM of PA-001 v.s. 2468 nM of EK1[18] and 580 nM of HY3000[20].   The second key

characteristics of PA-001 lies in its target region, FPPR, as a highly conserved region with minimal

variations.   The third notable feature of PA-001 is its 13-residue macrocyclic structure, markedly

smaller than the previously reported mimicking peptides mentioned above (typically more than 30

residues), making it accessible to FPPR positioned in shallow depression at the surface of Spike

pre-fusion form.   In addition, its cyclic architecture generally confers improved proteolytic stability,

reduced immunogenicity, and favorable pharmacokinetics.

So far, macrocyclic peptides have been applied to target SARS-CoV-2 Spike to develop anti-

SARS-CoV-2 agents, majority of which target RBD in S1[53-57].   Among them, in vivo anti-SARS-

CoV-2 effect was shown for three series of peptides, 6L3-1F3P11hR (against RBD), multivalent

bicyclic peptides (against RBD), and S-20-1 (targeting both RBD and HR1)[54,56,57].   However,

antiviral activity of S-20-1 was relatively mild, with 540-10,230 nM as IC_{50} in cell-based infection

assay, and that of 6L3-1F3P11hR varied at 1-332 nM depending on the target SARS-CoV-2

variants, consistent with the property of RBD-targeting agents.   None of them have been

reported to be administered in human.   In addition to its distinct target (FPPR), PA-001 shows an

advantage among these macrocyclic peptides with its high activity in vivo and administration

profile in human.

In infection mouse and hamster models, PA-001 effectively inhibited SARS-CoV-2-induced

disease progression when administered before or after infection, underscoring its potential for

both prophylactic and therapeutic use.   Additionally, combination treatment with PA-001 with

RDV elevated therapeutic profiles, highlighting its well compatibility with already-approved

antivirals.   This is consistent with that PA-001 has a distinct mode of action from those of any

clinically approved anti-SARS-CoV-2 agents so far.   Given the demand for effective anti-COVID-

19 therapies, especially for immunocompromised patients who respond poorly to current

treatments and evolve drug-resistance[1-3], peptide-based antivirals with different mode of action

would provide an additional choice for COVID-19 therapy.   Encouragingly, phase I clinical trial of

PA-001 in 32 healthy volunteers demonstrated well tolerability (up to 128 mg administration of PA-

001) without serious adverse events, regardless of age.   In the single-dose cohort, the C_{max}

reached 18,300 ng/mL [9,610 nM, more than 28,000-fold higher than IC_{50} (0.34 nM) against

Omicron variants] with half-life of 4.46 h, and drug exposure was sufficient to maintain

concentrations above the cellular IC_{50} for up to 24 h post-dose in Groups of 128 mg administration.

Collectively, these findings suggest that PA-001 achieves robust and sustained systemic exposure,

supporting a favorable pharmacokinetic profile.   Therefore, our data raise PA-001 as a promising

anti-SARS-CoV-2 drug that targets S2 and inhibits fusion process, and forthcoming clinical trials

will assess its efficacy in SARS-CoV-2-infected patients.

In summary, our study shows the usefulness of macrocyclic peptides as a potential therapeutic

modality against COVID-19, that demonstrates FPPR in the S2 region as a druggable target to

inhibit virus fusion process.   The strong antiviral potency warrants PA-001 as a candidate seed

for the development of an anti-SARS-CoV-2 fusion inhibitor.

Methods

Synthesis of PA-001

PA-001 was synthesized using standard Fmoc solid-phase peptide synthesis (SPPS), as described in the Supplementary Notes.   After coupling of all amino acids, the N-terminus was deprotected and subsequently chloroacetylated on resin, followed by global deprotection using a standard deprotection cocktail containing trifluoroacetic acid (TFA).   The linear peptide was precipitated by the adding the deprotection solution to more than a 10-fold excess of diethyl ether and hexane.   The resulting crude peptide pellet was resuspended in diethyl ether and re-pelleted three times.   After the final wash, the pellet was dried and dissolved in DMSO/H_2O, followed by the addition of triethylamine to induce intramolecular cyclization via thioether bond formation between the cysteine thiol and the N-terminal chloroacetyl group.   Upon completion of cyclization, the cyclic peptide was purified by standard reverse-phase HPLC.   The molecular mass was confirmed by single-quadrupole LC/MS (LCMS-2020 system, Shimadzu).

Cell culture

VeroE6/TMPRSS2 cells, a VeroE6 cell line overexpressing transmembrane protease, serine 2 (TMPRSS2) and highly susceptible to SARS-CoV-2 infection[25], were cultured in Dulbecco's modified Eagle's medium (DMEM; Life Technologies) supplemented with 10% fetal bovine serum (FBS; Cell Culture Bioscience), 10 units/mL penicillin, 10 mg/mL streptomycin, 10 mM HEPES (pH 7.4), and 1 mg/mL G418 (Nacalai) at 37°C in 5% CO_2.   During the infection assay, 10% FBS was replaced with 2% FBS, with removing G418.   HEK293T cells were cultured in DMEM supplemented with 10% FBS (heat-inactivated) and 50 μg/mL gentamicin at 37°C in 5% CO_2.

Reagents

Anti-SARS-CoV-2 Spike S1 antibodies, casirivimab (CAS) and imdevimab (IMD), were cloned into human IgG1 and kappa expression vectors and co-transfected into Expi293F cells (Thermo Fisher Scientific) as described previously[58].   Mutant spike proteins were generated by introducing substitutions into expression vectors coding codon-optimized spike trimeric protein[59] by PCR, and expressed with Expi293F cells.   Antibodies and mutant spike proteins were purified from the culture supernatant using Protein G columns (Thermo Fisher Scientific) and TALON columns (Clontech), respectively, and dialyzed against PBS.   Nucleotide analogs, RDV and its active metabolite, GS-441524 (RDV-m), were purchased from Chemscene Inc (CS-0028115) and Cayman Chemical (30469), respectively.   PA-001 and HA tagged PA-001 were produced by PeptiDream Inc.   DMSO was purchased from Sigma-Aldrich.   Mefloquine was purchased from Tokyo Chemical Industry.

Infection assay

SARS-CoV-2 was handled in a biosafety level 3 (BSL3).   We used the SARS-CoV-2 Wk-521

(Wuhan type), QK002 (Alpha), TY8-612 (Beta), TY7-503 (Gamma), TY11-927 (Delta), TY38-873 (Omicron; BA.1.18), and TY41-984 (Omicron; XBB.1.16) strains, clinical isolates from a COVID-19 patient[25].   Virus infectious titers were measured by inoculating cells with a 10-fold serial dilution of virus and cytopathology measured to calculate $TCID_{50}$/mL[25].   For the infection assay, VeroE6/TMPRSS2 cells were inoculated with virus at an MOI of 0.003 for 1 h and washed to remove unbound virus.   Cells were cultured for an additional 24, 36, and 48 h prior to detection of extracellular viral RNA or intracellular viral N protein (24 or 36 h), and cytopathic effects (CPE) (48 h), respectively.   Compounds were added during virus inoculation (1 h) and incubation after washing (24, 36, or 48 h), or incubation after washing only.

Isolation and quantification of viral RNA

Viral RNA was extracted with RNeasy mini kit (QIAGEN) from cells and MagMax Viral/Pathogen II Nucleic Acid Isolation kit (Thermo Fisher Scientific) or QIAamp Viral RNA mini kit (QIAGEN) from culture supernatant.   Viral RNA was quantified by real time RT-PCR analysis with a one-step qRT-PCR kit (THUNDERBIRD Probe One-step qRT-PCR kit, TOYOBO) using 5'-ACAGGTACGTTAATAGTTAATAGCGT-3', 5'- ATATTGCAGCAGTACGCACACA-3' as a primer set, and 5'-FAM-ACACTAGCCATCCTTACTGCGCTTCG-TAMRA-3' (E-set) as a probe[60].

Detection of viral proteins

Viral proteins were detected using rabbit anti-SARS-CoV N antibody[61] and mouse anti-SARS-CoV-2 spike antibody (abcam, ab273433) as primary antibodies, with anti-rabbit IgG-HRP (Cell Signaling, 7074) and anti-mouse IgG-HRP (Cell Signaling, 7076) as secondary antibodies by immunoblot analyses as previously reported[62].

Virus-cell binding assay

SARS-CoV-2 preincubated with the indicated compounds for 30 min was exposed to VeroE6/TMPRSS2 cells at an MOI of 0.01 at 4°C for 15 min.   After extensive wash, total RNA was extracted from cells and viral RNA was quantified by real time RT-PCR to measure cell-bound virus, as described [63].

PA-001-Spike binding assay

Binding of HA-tagged PA-001 to SARS-CoV-2 Spike proteins was quantified by ELISA as follows: Recombinant wild type (from Sino biological, 40589-V08B1, 40591-V08B1, and 40590-V08B) or mutated Spike proteins were immobilized on the plates, and then incubated with PA-001-HA at various concentrations together with or without excess amount of non-tagged PA-001, used as a binding competitor.   After washing with PBST, horseradish peroxidase-labeled mouse anti-HA tag antibody (MBL Life Science, M180-7, 1:5000 or 1:2500) was added, and visualized with reacting with a substrate solution (SUMITOMO BAKELITE) to detect absorbance at 450 nm

with a microplate reader (TECAN).

Cell membrane fusion assay

The S-ACE2 interaction-mediated cell-cell fusion assay was performed essentially based on
the previous report[29], with some modifications. Spike proteins expressed in HEK293T cells
mediate the fusion with VeroE6/TMPRSS2 cells, and the cell-cell fusion was quantified based on
the measurement of luminescence intensity derived from reconstituted NanoLuc reporter
(composed of HiBiT stably expressed in VeroE6/TMPRSS2 cells and LgBiT transiently expressed
in HEK293T cells). HEK293T cells were co-transfected with the plasmid expressing Spike in
combination with the HaloTag-inserted pBiT1.1-N[TK/LgBiT] plasmid (Promega) using
Lipofectamine 3000 Transfection Reagent according to the manufacturer's instruction (Thermo
Fisher Scientific).   At 48 h post-transfection, the cells were co-incubated with
VeroE6/TMPRSS2/HiBiT cells in the presence or absence of varying concentrations of PA-001 for
1 h.   The media were replaced with Opti-MEM containing 0.5% FBS and the substrate of
NanoLuc from Nano-Glo Live Cell Assay System (Promega) to measure luminescence.

Generation of PA-001 resistant viruses

To obtain PA-001-resistant SARS-CoV-2 variants, SARS-CoV-2 (WK-521) was inoculated to
VeroE6/TMPRSS2 cells at an MOI of 0.003, followed by incubation of the cells in the presence of
1,000 nM PA-001 for 48 h and recovery of culture supernatant.   The 1:1 diluted culture
supernatant was then inoculated to VeroE6/TMPRSS2 cells to repeat the same procedure until
the second passage.   The 1:10 diluted culture supernatant was inoculated to the cells in the
presence of 10 μM PA-001, followed by recovery of culture supernatant at 24 h later, which was
repeated for the three additional passages.   Virus was further amplified with two additional
passages in the absence of PA-001.

SARS-CoV-2 pseudovirus infection assay

SARS-CoV-2 pseudoviruses were produced using vesicular stomatitis virus (VSV)-based
pseudovirus system, using the expression plasmid for the G-deficient VSV carrying the luciferase
gene and that encoding SARS-CoV-2 Spike derived from the Wuhan type Wk-521 or the Omicron
sublineages, JN.1 or XFG, essentially as previously described[64,65].   VeroE6/TMPRSS2 cells
were inoculated with the SARS-CoV-2 pseudovirus in the presence or absence of various
concentrations of the indicated compounds, peptide, or antibodies followed by incubation with or
without the compounds, peptide, or antibodies for 24 h to measure intracellular luciferase activity
with Luciferase Assay System (Promega), as described[62].

Sequence analysis of PA-001 resistant viruses

cDNA libraries were generated from RNA extracted from SARS-CoV-2 isolates according to

the ARTIC nCoV-2019 sequencing protocol v3 (LoCost) (https://www.protocols.io/view/ncov-
2019-sequencing-protocol-v3-locost-bp2l6n26rgqe/v3).   Whole-genome sequencing was
performed using the MinION platform (Oxford Nanopore Technologies), followed by consensus
sequence generation using the Medaka pipeline (https://github.com/nanoporetech/medaka).
Lineage and clade assignments, as well as mutation identification, were conducted using the
Pangolin   COVID-19   Lineage   Assigner   (https://pangolin.cog-uk.io/)   and   Nextclade
(https://clades.nextstrain.org/).

587 Pharmacokinetic analysis of PA-001 in mice

The study was conducted under accreditation by the Association for Assessment and
Accreditation of Laboratory Animal Care (AAALAC) with approval of Institutional Animal Care and
Use Committee (IACUC, protocol No. PK-T-07182021).   Fasted female BALB/c mice (8 weeks
old, Si Bei Fu Laboratory Animal Technology) were intraperitoneally administered with PA-001 at
the dose of 10 mg/kg, and were sacrificed to collect blood at 1, 7, and 24 h post-administration.
The blood was centrifuged to prepare plasma, mixed with acetonitrile containing formic acid, and
centrifuged.   The supernatant was injected onto LC/MS/MS system (LCMS-8060, Shimadzu) to
determine plasma concentrations of PA-001.

597 SARS-CoV-2 infection in vivo

For infection to the mouse model in Fig. 4B and Fig. 5B and C, all the experiments were
approved by the Animal Care and Use Committee of the National Institute of Infectious Diseases
in Japan, and handled in BSL3 animal facilities (ABSL3) that were certified by the Japan Health
Sciences Foundation according to the guidelines of this committee.   BALB/c mice (13 weeks old,
female, Japan SLC) were maintained for 3 weeks in the specific pathogen-free facility and then
transferred to the ABSL3.   After a week of acclimation to the facility, the mice (17 weeks old)
were anesthetized by intraperitoneal injection with a mixture of 0.1 ml/10 g body weight of 1.0 mg
ketamine and 0.02 mg xylazine.   Animals were then inoculated intranasally with 3×10^3 $TCID_{50}$
(30 µl, 5 LD_{50}/mouse, 100% lethal dose) QHmusX, a mouse-adopted SARS-CoV-2 clone[38].
Body weight was measured daily for 10 days (n = 10–11 per group), and animals were sacrificed
at 3 and 10 days post-inoculation to analyze viral replication, cytokine expression, and disease
pathology (n = 4 or 6–7 per group).   Clinical signs were observed up until 10 days post-
inoculation.   The humane endpoint was defined as the appearance of clinically diagnostic signs
of respiratory stress, including respiratory distress and more than 25% weight loss.   Animals
were euthanized under anesthesia with an overdose of isoflurane if severe disease symptoms or
weight loss were observed.   In Fig. 4B, PA-001 was administered intraperitoneally at a dose of
10 or 30 mg/kg, at 0.04, 0.3, 1, 2, 3, and 5 days post-inoculation.   The CAS + IMD-treatment
group received the cocktail CAS + IMD antibodies (10 mg/kg) at 1 h post-inoculation.   The
vehicle control group was administered with PBS.   In, Fig. 5B and C, RDV-m was administered

intraperitoneally at a dose of 25 mg/kg (on day 1 post-infection) and 12.5 mg/kg (on days 2 and 3 post-infection), with a volume of 6.25 mL/kg per dose, once daily.   Concurrently, PA-001 was administered intraperitoneally at a dose of 26 mg/kg, with a volume of 10 mL/kg per dose, once daily.   The control group was administered with PBS.

Infection studies to the hamster model shown in Fig. 4A and C were conducted in the animal BSL-3 laboratory, accredited by AAALAC, under IACUC-approved protocol (22-032, 16103.01). In Fig. 4A, seven-week-old male Syrian hamsters (Envigo) were intranasally inoculated with 3.48 x 10^3 $TCID_{50}$ of SARS-CoV-2 (Delta variant, BEI Resources, NR-56116) under anesthesia with ketamine/xylazine at day 0.   PA-001 was intraperitoneally administered to hamsters twice daily at 0.3, 1, or 3 mg/kg, or PBS as a vehicle control, at 12 h and 2 h prior to infection and 12 h, 24 h, and 36 h following infection (-0.5, -0.08, 0.5, 1, and 1.5 day post infection).   Lung in the hamsters were collected at 2 days post-infection to examine the infectious titer of SARS-CoV-2.   In Fig. 4C, Syrian Golden hamsters (female, 8 week-old, Charles Reiver Laboratories) were intranasally infected with 2.11 x 10^5 $TCID_{50}$ SARS-COV2 (USA_WA1/2020, CDC, kindly provided by UTMB Galveston), and then administered intraperitonially with either PBS as a vehicle control or PA-001 1 or 5 mg/kg (n=6/group) (7.5 mL/kg) at 2, 8, 24, 48 and 72 h post-infection.   Body weight and clinical observation were evaluated every day, and the lungs were collected at five day postinfection.   The left lung lobes were used for quantifying SARS-CoV-2 RNA by real time RT-PCR, whereas the right lung lobes were fixed with 10% neutral buffered formalin for histopathological evaluation.

Viral titration from mouse and hamster tissues

Lung tissue samples from BALB/c mice were collected at the time of postmortem examination and stored at -80°C.   Tissue homogenates (10% w/v) were prepared in Dulbecco's Modified Eagle medium (DMEM) containing 2% FBS, 50 IU/ml penicillin G, and 50 µg/ml streptomycin, and samples were inoculated onto VeroE6/TMPRSS2 cell cultures, which were then examined for observing cytopathic effects (CPEs) for 5 days to determine the viral infectivity titers ($TCID_{50}$) as described above.

Lung tissue samples from Syrian hamsters were homogenized and centrifuged to collect the supernatant.   VeroE6/TMPRSS2 cells were treated with the supernatant to evaluate CPEs for 4 days to determine the viral infectivity titers with the Read-Muench formula[66].

Detection of inflammatory cytokines and chemokines

Homogenized mouse lung tissue samples (10% w/v) were diluted 1:1 in cell extraction buffer [10 mM Tris (pH 7.4), 100 mM NaCl, 1 mM EDTA, 1 mM EGTA, 1 mM NaF, 20 mM $Na_4P_2O_7$, 2 mM Na_3VO_4, 1% Triton X-100, 10% glycerol, 0.1% SDS, and 0.5% deoxycholate (BioSource International, Camarillo, CA)], incubated for 10 min on ice with vortexing, irradiated for 10 min with UV-C light to inactivate infectious virus.   Cytokine and chemokine levels were measured with

mouse cytokine–chemokine magnetic bead panel 96-well plate assay kit (Milliplex MAP kit, Merck Millipore), detected on a Luminex 200 instrument with xPONENT software (Merck Millipore), according to the manufacturer's instruction.

Phase I study

1. Study design and oversight

This phase I, double-blind, randomized, placebo-controlled, single- and multiple-ascending dose study of the safety, tolerability, and pharmacokinetics in healthy and elderly subjects was conducted at a single clinical site, Fortrea Clinical Research Unit Inc.   The protocol (PA-001-101) was approved by Salus institutional review board (IRB).   Study began in Sep 2024 and was completed in Jun 2025 (from FSFV to LPLV).   All participants provided written informed consent before enrollment.

2. Study Design

This study was conducted in 2 parts of Part A and B, as shown below.

Part A comprised a single-dose, sequential-group study of PA-001.   A total of up to 40 subjects were planned to be studied in up to 4 groups (Groups A1-4, PA-001 doses were 18, 64, 128 and 128 mg, respectively, with each group comprising 8 subjects. In each group, 6 subjects were to receive a single dose of PA-001 and 2 subjects were to receive a single dose of placebo. Age range was 18 to 65 years for Group A1-3, and >65 years for Group A4.   Doses of PA-001 or placebo were administered in accordance with a randomization schedule on the morning of Day 1, after a fast of at least 10 h.   Sentinel dosing: Groups A1 to A4 were divided into 2 subgroups, with each subgroup being dosed at least 24 h apart.   The first subgroup comprised 2 subjects, with 1 subject receiving PA-001 and 1 subject receiving placebo.   The second subgroup comprised 6 subjects, with 5 subjects receiving PA-001 and 1 subject receiving placebo. Continuation to dose the remaining 6 subjects was at the investigator's discretion.

Part B comprised a multiple-dose, sequential group study.   Part B was permitted to start after completion of Group A3 (in Part A), at a dose level less than or equal to that administered in Groups A1 to A3.   A total of 16 subjects were planned to be studied in 2 groups, with each group comprising 8 subjects.   In each group, 6 subjects were to receive PA-001, and 2 subjects were to receive placebo once per day until day 5.   PA-001 or placebo were administered in accordance with a randomization schedule after a fast of at least 10 h.

Inclusion criteria and exclusion criteria were described in Supplementary Notes.

3. Safety assessments

The condition of each subject was monitored from the time of signing the ICF to final discharge from the study.   Subjects were observed for any signs or symptoms and asked about their condition by open questioning, such as "How have you been feeling since you were last asked?" at least once each day while resident at the study site and at each study visit.   Subjects were also encouraged to spontaneously report adverse effects occurring at any other time during the

study. All nonserious adverse effects, whether reported by the subject voluntarily or on questioning, or noted on physical examination, were recorded from initiation of study treatment until study completion. Serious adverse effects were recorded from the time the subject signs the ICF until study completion. The nature, time of onset, duration, and severity were documented, together with an investigator's (or designee's) opinion of the relationship to study treatment. Adverse events recorded during the study will be followed up, where possible, until resolution or until the unresolved adverse effects were judged by the investigator (or designee) to have stabilized. This was completed at the investigator's (or designee's) discretion.

4. Administration

In Part A, subjects received a single 50mL intravenous infusion over 1 h (± 2 min) containing 18 mg (Group A1), 64 mg (Group A2), or 128 mg (Groups A3 and A4) of PA-001 or placebo on day 1. In part B, subjects received multiple 50mL intravenous infusions over 1 h (± 2 min) containing 64 mg (Group B1) or 128 mg (Group B2) of PA-001 or placeboonce per day on day 1 through 5, inclusive.

5. Analysis of PA-001 concentrations

Concentration of PA-001 in plasma and urine samples was measured with liquid chromatography tandem mass spectrometry (LC/MS/MS, QTRAP® 5500 System, SCIEX).

Statistics and reproducibility

Bar graphs are presented as mean ± standard deviation with individual data sets, and experiments were repeated in at least triplicates or more unless indicated otherwise. In this study, GraphPad Prism (v.10.5.0) was used for the ordinary one-way ANOVA followed by Dunnett's multiple comparisons with control (Fig. 2C-a, Fig. 4A-b, and Fig. 4C-c) or followed by tukey's multiple comparisons (Fig. 5B-d, e, and C-a, b), and the ordinary two-way ANOVA followed by Dunnett's multiple comparisons with control (Fig. 4B-c, Fig. 4C-b) or followed by tukey's multiple comparisons (Fig. 2F-b and Fig. 5B-c).

Acknowledgments

We appreciate Drs. Setsuo Hasegawa (Pharma Spur), Akihito Shimoi (SNBL), Takuma Yonemura, Takeshi Chiyoda, Asuka Yamamoto (Sumida Hospital) for valuable scientific advice based on clinical research experience, which contributed to the rationale for the starting dose in the Phase I study. We also thank Dr. Keiichi Masuya for PA-001 generation. We are grateful to Drs. Dai Izawa, Midori Ozaki, Takiko Yoshida, for their technical assistance. We thank the members of the Management Department of Biosafety and Laboratory Animals for support with the BSL3 facility. We also appreciate leadership of Dr. Gene Voskuhl (Fortrea Clinical Research) to operate Phase I study. This study was supported by Japan Agency for Medical Research and Development (AMED) under grand numbers JP25fk0310525 (K.W.), JP23fk0108590 (K.W.), JP25ak0101271 (K.W.), 25wm0325079 (K.W.), JP253fa727002 (N.Nagata, T.S.),

731 JP25wm0125008 (T.S.), JP25fk0108732 (T.S.), JP223fa627003 (T.S.), JP22fk0108521 (M.M.);

732 JSPS/MEXT under KAKENHI grant numbers JP24K18457 (H.Ohashi), JP24K02290 (K.W.),

733 JP25K21782 (K.W.), JP25K21767 (K.W.), JP25K02691 (K.W.); JST MIRAI program grant number

734 JPMJMI22G1 (K.W.); Takeda Science Foundation (K.W.).

Author Contributions

Experimental systems were developed by M.O., T.S., N.Nagata and K.W. Compound

screening was supervised by M.O. and P.C.R. M.O., K.N.,T.N., K.S.,N.Nakamura and J.Y.

performed screening for lead compound of PA-001.   M.O., K.M. and H.Kurasaki optimized

peptides and identified PA-001.   Molecular biological and cell-based experiments were

performed by H.Ohashi, Y.H., K.Shionoya, T.K., H.Ogawa, and K.W.   Animal experiments using

mice were designed and performed by N.IY., N.Nagata, and H.Kitamura, and those using

hamsters were designed and managed by T.K., K.N. and H.Kitamura.   Peptide synthesis was

performed by M.O., H.Kurasaki, K.N., T.N., J.Y., K.Sudo, N.Nakamura, and K.M.   Non-clinical

ADME analyses was conducted by K.Y., Y.S. and T.H.   Preparing materials was managed by

H.Kurasaki, S.M., T.M., T.H., and S.I.   Experimental design was developed by N.Nagata, M.O.,

P.C.R., T.S., and K.W.   Phase I clinical trial was designed and managed by S.I., N.K., and M.M.

Original draft was written by H.Ohashi, N.Nagata, S.M., H.Kitamura, and K.W. with critical

discussion by T.K, M.O., H.Kurasaki, N.IY., Y.H., S.M., K.Shionoya, K.N., T.N., J.Y., K.Sudo.,

N.Nakamura, K.M., H.Ogawa, K.Yoshida, Y.S., T.M., T.H., S.I., Y.T., N.K., P.C.R., M.M., and T.S.

Fundings were acquired by H.Ohashi, M.M., T.S., N.Nagata, and K.W.   The project was

supervised by H.Kitamura and K.W.

Competing Interests

757 T.K., M.O., H.Kurasaki, K.N., T.N., J.Y., K.Sudo, N.Nakamura, K.M., H.Ogawa, K.Y., Y.S., T.M.,

758 T.H., S.I., N.K., H.Kitamura and M.M. are employees of PeptiDream Inc.   P.C.R. is CEO of

759 PeptiDream Inc.   M.M. is CMO of PeptiDream and CEO of PeptiAid Inc.

Data Availability

763 Source data are provided with this paper.

References

References

1   Iriyama, C. *et al.* Clinical and molecular landscape of prolonged SARS-CoV-2 infection with
resistance to remdesivir in immunocompromised patients. *PNAS Nexus* **4**, pgaf085 (2025).

2   Nooruzzaman, M. *et al.* Emergence of transmissible SARS-CoV-2 variants with decreased
sensitivity to antivirals in immunocompromised patients with persistent infections. *Nat*
Commun **15**, 7999 (2024).

3   Gandhi, S. *et al.* De novo emergence of a remdesivir resistance mutation during treatment of
persistent SARS-CoV-2 infection in an immunocompromised patient: a case report. *Nat*
Commun **13**, 1547 (2022).

4   Chen, B., Farzan, M. & Choe, H. SARS-CoV-2 spike protein: structure, viral entry and variants.
Nat Rev Microbiol **23**, 455–468 (2025).

5   Hoffmann, M. *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked
by a Clinically Proven Protease Inhibitor. *Cell* **181**, 271–280 e278 (2020).

6   Shi, W. *et al.* Cryo-EM structure of SARS-CoV-2 postfusion spike in membrane. *Nature* **619**,
403–409 (2023).

7   Calvaresi, V. *et al.* Structural dynamics in the evolution of SARS-CoV-2 spike glycoprotein. *Nat*
Commun **14**, 1421 (2023).

8   Lei, R. *et al.* Functional and antigenic characterization of SARS-CoV-2 spike fusion peptide by
deep mutational scanning. *Nat Commun* **15**, 4056 (2024).

9   Shi, W. *et al.* Effect of the S2' site cleavage on SARS-CoV-2 spike. *Nat Commun* **16**, 11675
(2025).

10   Marcink, T. C. *et al.* Intermediates in SARS-CoV-2 spike-mediated cell entry. *Sci Adv* **8**,
eabo3153 (2022).

11   Xing, L. *et al.* Early fusion intermediate of ACE2-using coronavirus spike acting as an antiviral
target. *Cell* **188**, 1297–1314 e1224 (2025).

12   Grunst, M. W. *et al.* Structure and inhibition of SARS-CoV-2 spike refolding in membranes.
Science **385**, 757–765 (2024).

13   Dimitrov, D. S. Virus entry: molecular mechanisms and biomedical applications. *Nat Rev*
Microbiol **2**, 109–122 (2004).

14   Dong, M., Galvan Achi, J. M., Du, R., Rong, L. & Cui, Q. Development of SARS-CoV-2 entry
antivirals. *Cell Insight* **3**, 100144 (2024).

15   Chen, Y. *et al.* Broadly neutralizing antibodies to SARS-CoV-2 and other human coronaviruses.
Nat Rev Immunol **23**, 189–199 (2023).

16   Imai, M. *et al.* Efficacy of Antiviral Agents against Omicron Subvariants BQ.1.1 and XBB. *N*
Engl J Med **388**, 89–91 (2023).

17   Guo, L. *et al.* Targetable elements in SARS-CoV-2 S2 subunit for the design of pan-coronavirus
fusion inhibitors and vaccines. *Signal Transduct Target Ther* **8**, 197 (2023).

18   Xia, S. *et al.* Inhibition of SARS-CoV-2 (previously 2019-nCoV) infection by a highly potent

pan-coronavirus fusion inhibitor targeting its spike protein that harbors a high capacity to
mediate membrane fusion. *Cell Res* **30**, 343–355 (2020).

19   Yan, W., Zheng, Y., Zeng, X., He, B. & Cheng, W. Structural biology of SARS-CoV-2: open the
door for novel therapies. *Signal Transduct Target Ther* **7**, 26 (2022).

20   Sun, H. *et al.* Structural basis of HCoV-19 fusion core and an effective inhibition peptide against
virus entry. *Emerg Microbes Infect* **9**, 1238–1241 (2020).

21   Vinogradov, A. A., Yin, Y. & Suga, H. Macrocyclic Peptides as Drug Candidates: Recent
Progress and Remaining Challenges. *J Am Chem Soc* **141**, 4167–4181 (2019).

22   Ravel, G. *et al.* Additive effects of ALXN2420, a GH receptor antagonist, and octreotide on
IGF1 suppression in vivo. *Eur J Endocrinol* **192**, 299–307 (2025).

23   Sun, D. *et al.* The discovery and structural basis of two distinct state-dependent inhibitors of
BamA. *Nat Commun* **15**, 8718 (2024).

24   Brousseau, M. E. *et al.* Identification of a PCSK9-LDLR disruptor peptide with in vivo function.
Cell Chem Biol **29**, 249–258 e245 (2022).

25   Matsuyama, S. *et al.* Enhanced isolation of SARS-CoV-2 by TMPRSS2-expressing cells. *Proc*
Natl Acad Sci U S A **117**, 7001–7003 (2020).

26   Wang, M. *et al.* Remdesivir and chloroquine effectively inhibit the recently emerged novel
coronavirus (2019-nCoV) in vitro. *Cell Res* **30**, 269–271 (2020).

27   Hansen, J. *et al.* Studies in humanized mice and convalescent humans yield a SARS-CoV-2
antibody cocktail. *Science* **369**, 1010–1014 (2020).

28   Jan, J. T. *et al.* Identification of existing pharmaceuticals and herbal medicines as inhibitors of
SARS-CoV-2 infection. *Proc Natl Acad Sci U S A* **118** (2021).

29   Xia, S. *et al.* A pan-coronavirus fusion inhibitor targeting the HR1 domain of human coronavirus
spike. *Sci Adv* **5**, eaav4580 (2019).

30   Matsuyama, S. *et al.* The Inhaled Steroid Ciclesonide Blocks SARS-CoV-2 RNA Replication
by Targeting the Viral Replication-Transcription Complex in Cultured Cells. *J Virol* **95** (2020).

31   Li, G., Hilgenfeld, R., Whitley, R. & De Clercq, E. Therapeutic strategies for COVID-19:
progress and lessons learned. *Nat Rev Drug Discov* **22**, 449–475 (2023).

32   Hoffmann, M. *et al.* The Omicron variant is highly resistant against antibody-mediated
neutralization: Implications for control of the COVID-19 pandemic. *Cell* **185**, 447–456 e411
(2022).

33   Planas, D. *et al.* Considerable escape of SARS-CoV-2 Omicron to antibody neutralization.
Nature **602**, 671–675 (2022).

34   Lu, Y., Ao, D., He, X. & Wei, X. The rising SARS-CoV-2 JN.1 variant: evolution, infectivity,
immune escape, and response strategies. *MedComm (2020)* **5**, e675 (2024).

35   He, Q. *et al.* Neutralization of EG.5, EG.5.1, BA.2.86, and JN.1 by antisera from dimeric
receptor-binding domain subunit vaccines and 41 human monoclonal antibodies. *Med* **5**, 401–
413 e404 (2024).

36   Sia, S. F. *et al.* Pathogenesis and transmission of SARS-CoV-2 in golden hamsters. *Nature*
583, 834–838 (2020).

37   Imai, M. *et al.* Syrian hamsters as a small animal model for SARS-CoV-2 infection and
countermeasure development. *Proc Natl Acad Sci U S A* **117**, 16587–16595 (2020).

38   Iwata-Yoshikawa, N. *et al.* A lethal mouse model for evaluating vaccine-associated enhanced
respiratory disease during SARS-CoV-2 infection. *Sci Adv* **8**, eabh3827 (2022).

39   Abdelnabi, R., Maes, P., de Jonghe, S., Weynand, B. & Neyts, J. Combination of the parent
analogue of remdesivir (GS-441524) and molnupiravir results in a markedly potent antiviral
effect in SARS-CoV-2 infected Syrian hamsters. *Front Pharmacol* **13**, 1072202 (2022).

40   Sheahan, T. P. *et al.* Broad-spectrum antiviral GS-5734 inhibits both epidemic and zoonotic
coronaviruses. *Sci Transl Med* **9** (2017).

41   Zhang, Z. *et al.* Associations of immunological features with COVID-19 severity: a systematic
review and meta-analysis. *BMC Infect Dis* **21**, 738 (2021).

42   Chen, L. D. *et al.* Association between cytokine profiles and lung injury in COVID-19
pneumonia. *Respir Res* **21**, 201 (2020).

43   Jontvedt Jorgensen, M. *et al.* Increased interleukin-6 and macrophage chemoattractant
protein-1 are associated with respiratory failure in COVID-19. *Sci Rep* **10**, 21697 (2020).

44   Liao, M. *et al.* Single-cell landscape of bronchoalveolar immune cells in patients with COVID-
19. *Nat Med* **26**, 842–844 (2020).

45   Akil, C., Xu, J., Shen, J. & Zhang, P. Unveiling the structural spectrum of SARS-CoV-2 fusion
by in situ cryo-ET. *Nat Commun* **16**, 5150 (2025).

46   Miner, J. C. *et al.* Integrative structural studies of the SARS-CoV-2 spike protein during the
fusion process (2022). *Curr Res Struct Biol* **4**, 220–230 (2022).

47   Cai, Y. *et al.* Distinct conformational states of SARS-CoV-2 spike protein. *Science* **369**, 1586–
1592 (2020).

48   Xiong, X. *et al.* A thermostable, closed SARS-CoV-2 spike protein trimer. *Nat Struct Mol Biol*
27, 934–941 (2020).

49   Zhang, J. *et al.* Structural impact on SARS-CoV-2 spike protein by D614G substitution.
Science **372**, 525–530 (2021).

50   Muthuraman, K. *et al.* Human antibody targeting of coronavirus spike S2 subunit is associated
with protection mediated by Fc effector functions. *J Virol* **99**, e0152325 (2025).

51   Wu, W. L. *et al.* Monoclonal antibody targeting the conserved region of the SARS-CoV-2 spike
protein to overcome viral variants. *JCI Insight* **7** (2022).

52   Silwal, A. P. *et al.* DNA aptamers inhibit SARS-CoV-2 spike-protein binding to hACE2 by an
RBD- independent or dependent approach. *Theranostics* **12**, 5522–5536 (2022).

53   Wang, S. *et al.* Identification of Covalent Cyclic Peptide Inhibitors Targeting Protein-Protein
Interactions Using Phage Display. *J Am Chem Soc* **147**, 7461–7475 (2025).

54   Xue, S. *et al.* A novel cyclic gamma-AApeptide-based long-acting pan-coronavirus fusion

inhibitor with potential oral bioavailability by targeting two sites in spike protein. *Cell Discov* **8**,
88 (2022).

55   Thijssen, V. *et al.* A broad-spectrum macrocyclic peptide inhibitor of the SARS-CoV-2 spike
protein. *Proc Natl Acad Sci U S A* **120**, e2303292120 (2023).

56   Wang, M. *et al.* Intranasal administration of broad-spectrum macrocyclic peptide inhibitor
protects against SARS-CoV-2 Omicron variants. *Nat Commun* **17**, 1753 (2026).

57   Gaynor, K. U. *et al.* Multivalent bicyclic peptides are an effective antiviral modality that can
potently inhibit SARS-CoV-2. *Nat Commun* **14**, 3583 (2023).

58   Moriyama, S. *et al.* Temporal maturation of neutralizing antibodies in COVID-19 convalescent
individuals improves potency and breadth to circulating SARS-CoV-2 variants. *Immunity* **54**,
1841–1852 e1844 (2021).

59   Onodera, T. *et al.* A SARS-CoV-2 antibody broadly neutralizes SARS-related coronaviruses
and variants by coordinated recognition of a virus-vulnerable site. *Immunity* **54**, 2385–2398
e2310 (2021).

60   Corman, V. M. *et al.* Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR.
Euro Surveill **25** (2020).

61   Mizutani, T., Fukushi, S., Saijo, M., Kurane, I. & Morikawa, S. Phosphorylation of p38 MAPK
and its downstream targets in SARS coronavirus-infected cells. *Biochem Biophys Res*
Commun **319**, 1228–1234 (2004).

62   Saso, W. *et al.* Significant role of host sialylated glycans in the infection and spread of severe
acute respiratory syndrome coronavirus 2. *PLoS Pathog* **18**, e1010590 (2022).

63   Ohashi, H. *et al.* Potential anti-COVID-19 agents, cepharanthine and nelfinavir, and their usage
for combination treatment. *iScience* **24**, 102367 (2021).

64   Fukushi, S. *et al.* Vesicular stomatitis virus pseudotyped with severe acute respiratory
syndrome coronavirus spike protein. *J Gen Virol* **86**, 2269–2274 (2005).

65   Tani, H. *et al.* Involvement of ceramide in the propagation of Japanese encephalitis virus. *J*
Virol **84**, 2798–2807 (2010).

66   Frilli, E., Amodeo, D., Cevenini, G., Nante, N. & Messina, G. Effectiveness of near-UVA in
SARS-CoV-2 inactivation. *Epidemiol Infect* **151**, e76 (2023).

Figure legends

Fig. 1. Identification of PA-001, a cyclic peptide showing potent anti-SARS-CoV-2 activity.

(A) Schematic diagram of the flow to identify Spike-binding macrocyclic peptides and evaluate their antiviral activity.   The bottom arrows show the figures corresponding to the analyses shown above.   (B) Schematic representation of the mRNA display system to screen for Spike-binding macrocyclic peptides.   (C) Structure of PA-001.   (D) PA-001 inhibited the propagation of SARS-CoV-2.   (a) Schematic representation of SARS-CoV-2 infection assay.   VeroE6/TMPRSS2 cells were inoculated with or without SARS-CoV-2 (Wk-521; wuhan strain) at an MOI of 0.003 for 1 h, followed by washing out free virus and culturing cells with the medium supplemented with indicated compounds for additional 24 h to detect viral proteins in cells by immunoblot (c) and to quantify viral RNA in the culture supernatant by real time RT-PCR (d), as well as culturing cells for 48 h to observe virus-induced cytopathic effect (CPE) by microscopic observation (b).   In (c), viral Spike (upper) and N (middle) proteins as well as cellular actin as an internal control (lower) were detected.   In (d), quantified viral RNAs were plotted against the concentration of treated PA-001 (nM).   (E) Pharmacokinetics of PA-001 in mice. (a) Scheme for treatment with PA-001 BALB/c mice and time points (1, 7, and 24 h post IP injection) of plasma collection.   (b) BALB/c mice were intraperitoneally administered with PA-001 at 10 mg/kg and were chased to obtain plasma at 1, 7, and 24 h post-administration to quantify PA-001 concentration (mean$\pm$SD).

Fig. 2. PA-001 bound to FPPR in the S2 subunit and inhibited the membrane fusion.

(A) Binding of PA-001 to recombinant SARS-CoV-2 Spike protein of the full length (S1+S2), S1, or S2 regions.   Each recombinant Spike protein was immobilized and incubated with HA-tagged PA-001 at various concentrations up to 100 nM, followed by washing out and detection with horseradish peroxidase (HRP)-conjugated anti-HA antibody.   (B) Schematic model for the SARS-CoV-2 entry process, in which viral attachment to cells via Spike-ACE2 binding is followed by the fusion of viral envelope with cellular membrane either on the cell surface or in the endosome/lysosome.   (C) (a) Virus-cell attachment assay.   VeroE6/TMPRSS2 cells were exposed to SARS-CoV-2 at an MOI of 0.01 at 4°C for 15 min in the presence or absence of casirivimab (CAS) + imdevimab (IMD) (50, 500, or 5,000 ng/mL), mefloquine (MFQ) (0.4, 2, or 10 µM), or PA-001 (40, 200, or 1,000 nM), and were then washed out to quantify cell-bound viral RNA by real-time RT-PCR.   Ordinary one-way ANOVA followed by Dunnett's multiple comparisons with the DMSO control was performed (*: $p<0.05$, **: $p<0.01$, ***: $p<0.001$, ****: $p<0.0001$).   (b) Cell-cell fusion assay.   Spike-overexpressing HEK293T cells carrying the LgBiT gene were co-cultured with ACE2-expressing VeroE6/TMPRSS2 cells carrying the HiBiT gene upon treatment with or without varying concentrations (up to 200 nM) of PA-001 for 1 h to detect luciferase activity to evaluate fusion activity mediated by Spike and ACE2. IC_{50} of PA-001 is also indicated.   (D) Induction of PA-001-resistant SARS-CoV-2 variants.   (a) Schematic representation of the infection assay for inducing PA-001-resistant SARS-CoV-2 clones.   SARS-CoV-2-infected

VeroE6/TMPRSS2 cells were continuously cultured with PA-001 at 1,000 nM for 24-48 h (Passage 0). The resultant culture supernatants were collected and diluted to 10-fold, which were then inoculated to naive VeroE6/TMPRSS2 cells in the presence of 1,000 nM PA-001 and cultured for 24 h (Passage 1). The culture supernatants were repeatedly collected and then inoculated to the cells upon 1,000 nM PA-001 for the next passage. At passage 7, the culture supernatants were extracted from eight independent cultures (A8 – H8). (b) PA-001 sensitivity of the collected supernatant clones. VeroE6/TMPRSS2 cells were inoculated with these collected supernatants or the parental virus (WT) upon treatment with varying concentrations of PA-001 (left) or RDV as a reference (right) to determine virus RNA levels in the culture supernatants as shown in Fig. 1D-d. (E) Amino acid sequence of the PA-001-resistant clones. (a) Upper, schematic structure for the SARS-CoV-2 Spike protein. Lower, amino acid sequence of the clones, A8–H8, as well as the parental virus (WT). The substituted amino acids are highlighted by red. The amino acid numbers are indicated. (b) Structure of Spike trimer in the pre-fusion (left, PDB: 6XR8), intermediation (center, PDB: 8Z7P), and post-fusion (right, PDB: 8FDW) forms. Red color indicates the mutated amino acids shown in (a). Blue color indicates FP. (F) Binding capacity of recombinant SARS-CoV-2 Spike carrying the L841R and P862Q substitutions to PA-001. (a) Each recombinant Spike (WT, L841R, and P862Q) was immobilized and incubated with HA-tagged PA-001 (PA-001-HA) at varying concentrations in the presence (gray, competition) or absence (red, S-PA001 binding) of non-tagged PA-001 at 10 µM. After washing out, bound HA-PA-001 was visualized with HRP-conjugated anti-HA antibody. Samples without Spike-immobilization were also prepared to define the background signal (green). (b) Binding signals with Spike WT (left), Spike L841R (center), and Spike P862Q (right) are plotted against the concentrations of treated PA-001-HA. Ordinary two-way ANOVA followed by tukey's multiple comparisons was performed (*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$). (G) Cell fusion assay performed as described in Fig. 2C-b using the cells overexpressing either Spike WT, L841R, or P862Q co-cultured with ACE2-expressing cells upon PA-001 treatment at various concentrations.

Fig. 3. PA-001 showed a broad anti-SARS-CoV-2 activity against multiple variants.

(A) Antiviral activity of PA-001 and casirivimab + imdevimab against the SARS-CoV-2 variants (Wuhan, Alpha, Beta, Gamma, Delta, and Omicrons BA.1.18 and XBB.1.16). Drug response was evaluated as described in Fig. 1D-d, using each SARS-CoV-2 variant as inoculum. (B) Sequence homology from aa 834 to 863 in Spike among the SARS-CoV2 variants. Asterisks indicate the fully homologous sites among all the variants.

Fig. 4. PA-001 inhibited viral propagation and disease progression in mice and hamsters.

(A) (a) Scheme for prophylactic treatment with PA-001 to SARS-CoV-2-infected hamsters. Golden Syrian hamsters (n = 7 for infected groups and n = 5 for uninfected group) were

intraperitoneally treated with PA-001 (0.3, 1.0, or 3.0 mg/kg) or the vehicle at day -0.5, -0.08,
0.5,1.0, and 1.5 (-12, -2, 12, 24, and 36 h) post-inoculation, and were inoculated with SARS-CoV-
2 (3.48×10^3 $TCID_{50}$/hamster) intranasally at day 0.   Lungs were collected at day 2 post-
inoculation.   (b) Infectious viral titer in the lungs at day 2 was quantified.   Black, vehicle; light
red, 0.3 mg/kg PA-001; red, 1.0 mg/kg PA-001; dark red, 3.0 mg/kg PA-001.   Ordinary one-way
ANOVA followed by Dunnett's multiple comparisons with the vehicle control was performed (*:
$p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$).   (B) (a) Scheme for therapeutic treatment with PA-001 to
SARS-CoV-2-infected mice.   BALB/cCrSlc mice (n = 8) were intranasally inoculated with mouse-
adopted SARS-CoV-2 (QHmusX) (3×10^3 $TCID_{50}$/mouse) (day 0) and then administered
intraperitoneally with PA-001 (10 or 30 mg/kg), CAS + IMD (10 mg/kg), or the vehicle control at
day 0.04, 0.3, 1, 2, 3, and 4 (1, 8, 24, 48, 72, and 96 h).   The mice were chased to examine the
body weight as well as survival ratio.   (b, c) Survival rate (b) and body weight (body weight at
day 0 as 100%) (c) along with the time course after virus inoculation (day) was plotted.   Black,
vehicle; gray, 10 mg/kg CAS + IMD; red, 10 mg/kg PA-001; dark red, 30 mg/kg PA-001.   Ordinary
two-way ANOVA followed by Dunnett's multiple comparisons with the PBS control was performed
(*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$).   (C) (a) Schematic diagram for PA-001
treatment after SARS-CoV-2 infection to hamsters.   Golden Syrian hamsters (n = 6) intranasally
inoculated with SARS-CoV-2 (2.11×10^5 $TCID_{50}$/hamster) (day 0) were intraperitoneally treated
with PA-001 (1.0 or 5.0 mg/kg) or the vehicle at day 0.08, 0.3, 1.0, 2.0, and 3.0 (2, 8, 24, 48, and
72 h).   Lungs were collected at day 5.   (b) Body weight was observed daily, defining those at
day 0 as 100%.   Black, vehicle; red, PA-001 1.0 mg/kg; dark red, PA-001 5.0 mg/kg.   Ordinary
two-way ANOVA followed by Dunnett's multiple comparisons with the vehicle control was
performed (*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$).   (c) SARS-CoV-2 RNA levels in
the lung were quantified at day 5.   Ordinary one-way ANOVA followed by Dunnett's
multiple comparisons with the vehicle control was performed (*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$).
(d) Histopathology of the lungs at day 5.   Lungs from hamsters of each treatment group were
stained by HE.

1014

Fig. 5. Augmented anti-SARS-CoV-2 profiles by combination treatment with PA-001 and
RDV.
(A) Combination treatment with PA-001 and RDV in cell-based infection assay.   Antiviral activities
were evaluated as described in Fig. 1D-d, using SARS-CoV-2-infected VeroE6/TMPRSS2 cells
that were treated with combinations of various concentrations of PA-001 (0, 0.04, 0.18, 0.70, 2.8,
and 11.2 nM) and RDV (0, 750, 1,500, 3,000, 6,000, and 12,000 nM).   (B) (a) Scheme for post-
administration of combination with PA-001 and the RDV metabolite, GS-441524 (RDV-m).
BALB/cCrSIc mice were intranasally inoculated with SARS-CoV-2 (2.3×10^3 $TCID_{50}$/mouse) and
then intraperitonially administered with either GS-441524 (25 mg/kg (day 1) and 12.5 mg/kg (days
2, 3)) alone, GS-441524 (25 mg/kg (day 1) and 12.5 mg/kg (days 2, 3)) with PA-001 (26 mg/kg),

or the vehicle once daily from day 1, to day 3. † Mice were euthanized at day 3 without dosing on the day for collecting lungs and sera (n = 4) and day 10 after monitoring survival rate and body weight (n = 5-7). (b, c) Survival rate (b) and body weight (c) until day 10 for each group are shown. Black, vehicle; blue, GS-441524; red, GS-441524 + PA-001. Ordinary two-way ANOVA followed by tukey's multiple comparisons was performed (*: $p<0.05$, **: $p<0.01$, ***: $p<0.001$, ****: $p<0.0001$). (d, e) Infectious viral titer in the lung (d) and the lung weight (e, lung weight-to-body weight ratio) at day 3 are shown. Ordinary one-way ANOVA followed by tukey's multiple comparisons was performed (*: $p<0.05$, **: $p<0.01$, ***: $p<0.001$). (C) Production of the cytokines related with the disease severity, IL-6, TNF-α, IL-1β, GCSF, IFN-γ, IP-10, MIP-1α, and MCP-1, from the serum and the lung at day 3 were analyzed. Ordinary one-way ANOVA followed by tukey's multiple comparisons was performed (*: $p<0.05$, **: $p<0.01$, ***: $p<0.001$).

Fig. 6. Safety and pharmacokinetics profiles of PA-001 single administration to healthy volunteers.

(A) Baseline characteristics of 32 participants in the phase I trial for PA-001. The indicated profiles are presented as mean ± standard deviation for each cohort group (Placebo, 18 mg PA-001, 64 mg PA-001, 128 mg PA-001, and 128 mg PA-001 to elderly adults). Numbers of participants for each group are also shown (n = 6 or 8 for each group). (B) Number of participants as well as percentage showing the treatment-related treatment emergent adverse effects (TEAEs) as indicated for each cohort group. (C) Plasma concentrations of PA-001 for each cohort (closed circle, 18 mg; closed diamond, 64 mg; closed square, 128 mg; open square,128 mg to elderly adults) in the single-dose sequential-group study (Part A) are plotted against the time post intravenous infusions in healthy subjects. (D) Pharmacokinetic parameters calculated based on the profile shown in (C). $AUC_{0-tlast}$, area under the concentration-time curve from time 0 to the endpoint; AUC_{0-24}, area under the concentration-time curve from 0 to 24 h; C_{max}, maximum concentration; T_{max}, time for maximum concentration; $T_{1/2}$, half-life. Data are represented as mean ± standard deviation.

Fig. 1

A

10¹³ candidates — PDPS — Peptide screening — Infection inhibition — Binding — Antiviral activity — Optimization — PK — PA-001 — MOA Fusion — S2-FPPR — Target motif — Pathogenesis — Phase I study

Table. 1 — Fig. 1 — Fig. 2, 3 — Fig. 4, 5 — Fig. 6

B In vitro peptide screening

Random DNA library (~10¹³) — Translation — Spike — Binding — Washout — Elution — Amplification

C PA-001

D

a) Cell-based SARS-CoV-2 infection assay

SARS-CoV-2 — PA-001 — viral RNA (d) — viral protein (c) — cytopathic effect (b)

VeroE6/TMPRSS2 cells — 1 h — 24 h — 48 h

b) SARS-CoV-2-induced cytopathic effect

DMSO — RDV — PA-001

(-) — (+) — (+) — (+) : SARS-CoV-2

c) SARS-CoV-2 protein production

SARS-CoV-2 : — DMSO — PA-001 — RDV

Spike

N

Actin

d) PA-001 – dose response

viral RNA (%) vs PA-001 (nM)

E

a) Pharmacokinetics in mice

Mice (BALB/c) (n = 3)

PA-001 10 mg/kg, IP

collection : 1 7 24 (h)

b) PA-001 in plasma

PA-001 (ng/mL) vs Time (h)

Fig. 2

A PA-001 binding to spike

B SARS-CoV-2 entry

- Attachment Fig. 2C a)
- Fusion Fig. 2C b), 2G
- ACE2
- pre-fusion form in Fig. 2E(b)
- post-fusion form in Fig. 2E(b)

C a) Viral attachment

b) Membrane fusion — IC50 = 3.6 nM

D a) Induction of PA-001-resistant variants

Passages (x 7)

A8, B8, C8, D8, E8, F8, G8, H8

VeroE6/TMPRSS2 cells

PA-001 sensitivity : Fig. 2D-b
Sequence : Fig. 2E
Spike binding : Fig. 2F
Fusion : Fig. 2G

b) SARS-CoV-2 production

E a) SARS-CoV-2 Spike

S1 — NTD, RBD, CTD, FPPR; S2 — FP, HR1, CH, CD, HR2, TMD

b) pre-fusion form, intermediation form, post-fusion form

F a) Spike-PA-001 binding assay

b) Spike (WT), Spike (L841R), Spike (P862Q)

G Membrane fusion — Spike (L841R), Spike (P862Q)

Fig. 3

Fig. 4

A Prophylactic PA-001 treatment in hamster

a)

Syrian hamsters (n = 7)

SARS-CoV-2 inoculation → 0

Tissue collection

PA-001 : -0.5 -0.08 0.5 1 1.5 2 (day)

b)

SARS-CoV-2 infectious titer in lung

PA-001 0.3 mg/kg
PA-001 1.0 mg/kg
PA-001 3.0 mg/kg

B Therapeutic PA-001 treatment in mice

a)

Mice (BALB/cCrSlc) (n = 8)

SARS-CoV-2 (QHmusX) inoculation

PA-001 : 0.04 0.3 1 2 3 4 10 (day)

b) Survival ratio

c) Body weight change

CAS + IMD 10 mg/kg, PA-001 10 mg/kg, PA-001 30 mg/kg

C Therapeutic PA-001 treatment in hamster

a)

Syrian hamsters (n = 6)

SARS-CoV-2 inoculation

Tissue collection

PA-001 : 0.08 0.3 1 2 3 5 (day)

b) Body weight change

c) SARS-CoV-2 copies in lung

PA-001 1.0 mg/kg
PA-001 5.0 mg/kg

d) Histopathology in lung

vehicle   PA-001 1.0 mg/kg   PA-001 5.0 mg/kg

Fig. 5

Fig. 6

A

Subject demographics and baseline characteristics

Demographic	Placebo (N = 8)	18 mg PA-001 (N = 6)	64 mg PA-001 (N = 6)	128 mg PA-001 (N = 6)	128 mg PA-001 (Elderly) (N = 6)	Overall (N = 32)
Age [years; range]	46.6 ± 14.20 [32-66]	38.7 ± 9.05 [30-55]	41.7 ± 14.18 [22-57]	37.2 ± 15.72 [23-59]	69.8 ± 3.71 [66-75]	46.8 ± 16.49 [22-75]
Sex						
Female	4 (50.0%)	2 (33.3%)	3 (50.0%)	5 (83.3%)	5 (83.3%)	19 (59.4%)
Male	4 (50.0%)	4 (66.7%)	3 (50.0%)	1 (16.7%)	1 (16.7%)	13 (40.6%)
Race						
White	7 (87.5%)	3 (50.0%)	2 (33.3%)	3 (50.0%)	6 (100%)	21 (65.6%)
Black or African American	1 (12.5%)	2 (33.3%)	4 (66.7%)	3 (50.0%)	---	10 (31.3%)
Multiple	---	1 (16.7%)	---	---	---	1 (3.1%)
Ethnicity						
Not Hispanic or Latino	5 (62.5%)	3 (50.0%)	5 (83.3%)	3 (50.0%)	4 (66.7%)	20 (62.5%)
Hispanic or Latino	3 (37.5%)	3 (50.0%)	1 (16.7%)	3 (50.0%)	2 (33.3%)	12 (37.5%)
Height (cm)	166.86 ± 8.522	172.03 ± 7.989	174.17 ± 10.809	169.28 ± 12.378	163.53 ± 6.087	169.03 ± 9.495
Body weight (kg)	74.640 ± 12.6180	74.700 ± 9.4240	84.650 ± 8.3050	76.770 ± 21.1180	65.130 ± 11.0530	75.140 ± 13.8040
Body Mass Index (kg/m^2)	26.75 ± 3.492	25.15 ± 1.231	27.97 ± 2.373	26.42 ± 4.118	24.38 ± 4.090	26.17 ± 3.299

B

Adverse effect of PA-001

Treatment-related TEAEs	Placebo (N = 8)	18 mg PA-001 (N = 6)	64 mg PA-001 (N = 6)	128 mg PA-001 (N = 6)	128 mg PA-001 (Erderly) (N = 6)
Overall	1 (12.5%)	-	2 (33.3%)	-	-
mild	1 (12.5%)	-	2 (33.3%)	-	-
Serious	-	-	-	-	-
Discontinuation	-	-	-	-	-
Death	-	-	-	-	-

C

PA-001 in plasma

D

Pharmacokinetic parameters of PA-001

Parameter	18 mg PA-001	64 mg PA-001	128 mg PA-001	128 mg PA-001 (Erderly)
$AUC_{0-tlast}$ (h*ng/mL)	7,050.0 ± 910.0	28,200.0 ± 3,190.0	74,200.0 ± 21,900.0	85,100.0 ± 12,500.0
AUC_{0-24} (h*ng/mL)	7,050.0 ± 910.0	27,900.0 ± 3,260.0	72,800.0 ± 21,100.0	83,200.0 ± 12,100.0
C_{max} (ng/mL)	2,130 ± 279	7,860 ± 814	18,300 ± 4,240	19,000 ± 2,220
T_{max} (h)	1.04 ± 0.04	1.02 ± 0.01	1.06 ± 0.02	1.05 ± 0.07
$T_{1/2}$ (h)	3.61 ± 0.63	4.41 ± 0.51	4.46 ± 0.54	4.79 ± 0.35