A Machine Learning Framework for Serogroup Classification of pathogenic species of *Leptospira* Based on *rfb* Locus Profiles

Authors

Edson de Carvalho Ferreira Filho[a], Pedro Melo Arruda[a], Leonardo Cabral Afonso Ferreira[a], Maria Raquel Venturim Cosate[b], Tetsu Sakamoto[a*]

Affiliation

[a]Bioinformatics Multidisciplinary Environment (BioME), Instituto Metrópole Digital (IMD), Universidade Federal do Rio Grande do Norte (UFRN), Brazil
[b]Independent Researcher, Boston, Massachusetts, USA

*Corresponding author: tetsu@imd.ufrn.br +55 84 98158-8225

Abstract

Leptospira is a highly diverse genus traditionally classified by serological assays into more than 30 serogroups and over 300 serovars. However, this classification system is often complex and inconsistent, as cross-reactions between antigens can lead to ambiguous results. Moreover, serological tests such as MAT and CAAT are labor-intensive, require live cultures, and are difficult to standardize across laboratories. To overcome these limitations, we compiled genomic data from 721 pathogenic *Leptospira* samples obtained from NCBI RefSeq and BIGSdb (Institut Pasteur) to develop a machine learning framework capable of predicting serological classification directly from genomic information. Our approach focuses on the *rfb* locus, a genomic region associated with lipopolysaccharide biosynthesis and antigenic diversity, and was designed to operate in two stages: the first stage assigns samples to one of four major serological classes, while the second stage classifies them into their respective serogroups. Models from both classification stages achieved high predictive performance, with perfect score in the first and a mean F1-score of 0.931 in the second stage. Feature importance analysis revealed non-random clustering of highly informative genes within the *rfb* locus and demonstrated that serogroup discrimination is driven by combinatorial patterns of gene presence and absence. Based on the strong genetic coherence observed at the *rfb* locus and shared antigenic features, we propose the term "seroclass" to designate these higher-order groupings. This approach provides a scalable and reproducible alternative to traditional serological testing and offers valuable applications for epidemiological surveillance, outbreak investigation, and vaccine development within the *Leptospira* genus.

Keywords

Leptospirosis; Serotyping; O-antigen; Lipopolysaccharide; Epidemiology; Random Forest

Highlights

- Genomic data enable accurate inference of *Leptospira* serological classification.
- The approach predicts serogroups directly from *rfb* locus gene composition.
- The model provides a scalable alternative to labor-intensive serological assays.
- We introduce the term "seroclass", a higher level of serological organization.
- The proposed framework supports epidemiological surveillance and vaccine design.

1. Introduction

The *Leptospira* genus comprises spirochete bacteria (Phylum Spirochaetes) characterized by their helical morphology and high motility (Ko et al., 2009; Levett, 2001). Pathogenic species of *Leptospira* are responsible for leptospirosis, one of the most widespread zoonotic diseases worldwide (Costa et al., 2015). Infection occurs through contact with water or soil contaminated with the urine of infected animals, leading to a wide spectrum of clinical outcomes ranging from mild febrile illness to severe Weil's disease and pulmonary hemorrhage (Haake and Levett, 2015).

There are two main approaches for classifying *Leptospira* isolates. The first is taxonomic classification, which is based on the evolutionary relationships among isolates. The advent of molecular and genomic methods prompted a taxonomic reorganization of the genus *Leptospira*. Initially divided into "pathogenic" and "saprophytic" species (Levett, 2001), the genus is now recognized to comprise at least 77 species (Dos Santos Ribeiro et al., 2023; Fernandes et al., 2022; Hamond et al., 2025; Korba et al., 2021), distributed across four phylogenetic clades with distinct ecological and pathogenic characteristics: P1 (pathogenic *sensu stricto*), P2 (intermediate), S1 (saprophytic *sensu stricto*), and S2 (environmental) (Vincent et al., 2019). Pathogenic species such as *L. interrogans*, *L. kirschneri*, and *L. borgpetersenii* are primarily associated with human and animal disease, whereas intermediate and saprophytic species are mainly found in environmental niches.

The second approach is serological classification, which relies on antigenic differences among isolates. *Leptospira* classification has traditionally been based on serological tests, primarily the microscopic agglutination test (MAT) and the cross-agglutination absorption test (CAAT), which led to the description of more than 300 serovars grouped into 30 serogroups (Caimi and Ruybal, 2020; Faine et al., 1999; Picardeau, 2017). It is important to note that serological classification does not necessarily reflect genetic relatedness, as distinct species may contain strains belonging to the same serogroup, and conversely, a single species may comprise multiple serogroups (Faine et al., 1999; Kalambaheti et al., 1999; Mitchison et al., 1997). Accurate identification of *Leptospira* strains and their serogroups or serovars is crucial for understanding epidemiological dynamics, tracking outbreaks, and supporting vaccine development (Levett, 2001; Picardeau, 2017). Although MAT and CAAT are considered the gold standard (Terpstra et al., 2003), they are laborious, require maintaining large reference collections, and are prone to cross-reactivity and subjective interpretation.

The development of alternative methods for serological classification is therefore of great interest in the field. In this context, molecular approaches have emerged as promising tools to complement traditional serological assays (Ciurariu et al., 2025; Giraud-Gatineau et al., 2025). At the molecular level, the *rfb* locus, which encodes enzymes involved in the biosynthesis of the O-antigen component of lipopolysaccharides (LPS), is largely responsible for *Leptospira*'s antigenic diversity and therefore underlies serogroup and serovar classification (de la Peña-Moctezuma et al., 2001, 1999). Variation in

gene content and organization within the *rfb* cluster determines structural differences in LPS, leading to distinct serological profiles among strains. Consequently, the *rfb* region represents a genetic link between traditional serological classification and genomic data.

Despite significant advances in sequencing and comparative genomics (Chinchilla et al., 2023; Ferreira et al., 2024; Medeiros et al., 2022; Nieves et al., 2022), connecting the extensive genomic diversity of *Leptospira* to its antigenic properties remains challenging. Machine learning (ML) methods have emerged as powerful tools for uncovering complex, nonlinear patterns in high-dimensional genomic data, enabling phenotype prediction such as antimicrobial resistance, host specificity, or serotype (Drouin et al., 2019; Lees et al., 2020). These approaches can detect subtle genomic signatures that are not easily captured by phylogenetic trees or pairwise similarity analyses. However, ML-based serogroup prediction in *Leptospira* remains largely unexplored.

Here, we propose a machine learning framework to classify *Leptospira* isolates into serogroups using genome-derived features. By integrating supervised learning algorithms with genomic representations, we aim to evaluate whether genomic variation—particularly in loci such as *rfb*—can reliably predict serogroup identity. Our results demonstrate that genomic data can reproduce and extend traditional serological classifications, providing a scalable and objective approach for *Leptospira* characterization and surveillance.

2. Materials and Methods

2.1 Genome and serological dataset collection

Leptospira genome sequences were obtained from the NCBI RefSeq Genomes database (Goldfarb et al., 2025) and the BIGSdb–Institut Pasteur repository (Jolley and Maiden, 2010) in October, 2024. Serological annotations were also retrieved from the corresponding databases. Only non-mutant isolates of the P1 group, with genome assemblies comprising fewer than 300 contigs and with available serogroup annotations, were included in the analysis. Potential redundancies between the two databases were identified and removed. Consistency of serogroup annotation between NCBI and BIGSdb entries was verified, and any discrepancies were resolved through manual curation and cross-checking with the literature. Additionally, we removed the sample *L. interrogans* str. UT053 (srg. Sejroe, srv. Medanensis) from the dataset, as it has been previously reported to deviate from the canonical *rfb* locus organization (Ferreira et al., 2024). Following these procedures, 721 *Leptospira* genomes were retained for downstream analyses (Supplementary Table S1). Additionally, we retrieved 30 genomic samples from NCBI RefSeq Genomes with serological information that were published after the development of the classification models to serve as validation dataset (Supplementary Table S2).

2.2 Feature matrix construction based on the *rfb* locus

The feature matrix used as input for the machine learning model comprised the identity values shared by each genome sample with the genes of the *rfb* locus as features (Supplementary Table S3). To construct the matrix, we first selected one representative strain from each *Leptospira* serogroup (Table 1), prioritizing genomes of high sequencing quality and that contained the full *rfb* locus assembled within a single contig. From each selected genome, amino acid sequences corresponding to the *rfb* locus genes were extracted. The resulting set of all protein sequences extracted was then clustered

using CD-HIT (Fu et al., 2012) with a similarity threshold of 80% to reduce redundancy. The clustering analysis resulted in a total of 592 clusters. Representative sequences from each cluster with at least 100 amino acids long, totaling 549 proteins, were retrieved and used as queries in TBLASTN (Altschul et al., 1990) searches against the genome assemblies of the isolates in the dataset. For each query–subject comparison, the percentage of amino acid identity of the best hit was corrected by the query length to account for partial alignments. The corrected identity values were used to fill the numerical feature matrix, which served as input for training the machine learning classification models.

Table 1: Representative leptospiral samples of each serogroup

Seroclass	Serogroup	Genome accession	Sample
	Ballum	GCF_009884235.1	*Leptospira borgpetersenii* str. R14L
	Canicola	GCF_008831465.1	*Leptospira interrogans* str. 611
	Celledoni	GCF_022344045.1	*Leptospira weilii* str. FMAS_PD2
	Icterohaemorrhagiae	GCF_000231175.1	*Leptospira interrogans* serovar Lai str. IPAV
1	Javanica	GCF_024722275.1	*Leptospira borgpetersenii* str. I2
	Manhao	GCF_000243815.2	*Leptospira alexanderi* serovar Manhao 3 str. L 60
	Pyrogenes	GCF_022559725.1	*Leptospira interrogans* str. FMAS_KG2
	Ranarum	GCF_000332415.1	*Leptospira weilii* serovar Ranarum str. ICFT
	Sarmin	GCF_030023765.1	*Leptospira santarosai* str. MMD3
	Australis	GCF_022819715.1	*Leptospira noguchii* str. IP1611024
	Autumnalis	GCF_022819425.1	*Leptospira noguchii* str. IP1709037
	Cynopteri	GCF_000243695.2	*Leptospira kirschneri* serovar Cynopteri str. 3522 CT
2	Djasiman	GCF_000216195.1	*Leptospira interrogans* serovar Djasiman str. LT1649
	Grippotyphosa	GCF_024526015.1	*Leptospira kirschneri* str. FMAS_PN5
	Panama	GCF_000306255.2	*Leptospira noguchii* serovar Panama str. CZ214
	Pomona	GCF_001857845.1	*Leptospira interrogans* serovar Pomona str. AKRFB
	Hebdomadis	GCF_000244115.1	*Leptospira interrogans* serovar Hebdomadis str. R499
3	Mini	GCF_000306675.2	*Leptospira mayottensis* str. 200901116
	Sejroe	GCF_003254845.1	*Leptospira borgpetersenii* serovar Hardjo-bovis str. 203
	Bataviae	GCF_014858865.1	*Leptospira interrogans* serovar Bataviae str. 1489
4	Shermani	GCF_000332395.2	*Leptospira santarosai* serovar Shermani str. 1342KT
	Tarassovi	GCF_024704545.1	*Leptospira borgpetersenii* str. MN900

To further minimize redundancy among samples, we performed a clustering analysis based on Pearson correlation coefficients computed from the feature matrix (Supplementary Table S4). Samples exhibiting extremely high correlation ($\rho > 0.999$) were grouped into clusters, and a single representative genome from each cluster was retained for subsequent analyses (Supplementary Table S1). This stringent threshold was adopted because isolates belonging to the same serogroup, even when originating from different geographic locations or years of isolation, frequently exhibited correlation values approaching 0.99. Nearly all clusters generated through this procedure contained isolates from the same species and serogroup. The only exception was a cluster comprising three isolates from serogroup Icterohaemorrhagiae and two from serogroup Javanica; these samples were excluded from further analyses to avoid potential misclassification bias. After this final filtering step, the dataset consisted of 384 *Leptospira* genomes (rows) and 549 *rfb*-encoded protein features (columns).

2.3 Exploratory Data Analysis and Visualization

To explore the structure and variability of the genomic feature matrix, a principal component analysis (PCA) was performed using the *scikit-learn* package (Pedregosa et al., 2011). PCA was applied to reduce the dimensionality of the dataset and to visualize the distribution of *Leptospira* isolates in a

two-dimensional space. Data visualization was carried out in Python using the Matplotlib and Seaborn libraries (Hunter, 2007; Waskom, 2021).

2.4 Machine Learning Models and Evaluation

The procedure for predicting the serogroup of a *Leptospira* isolate was designed as a two-step hierarchical classification pipeline. In the first step, each isolate was assigned to one of four major serological classes, hereafter referred to as seroclasses, as defined in previous work (Ferreira et al., 2024). In the second step, the isolate was classified into one of the serogroups belonging to the seroclass to which it had been assigned in the first stage.

All classifiers were trained as binary Balanced Random Forest (BRF) models to determine whether a given isolate belonged to a specific seroclass or serogroup. In other words, each model was trained to discriminate between "member" and "non-member" instances for a given target group. The hyperparameters were set as follows: n_estimators=200, random_state=2, max_leaf_nodes=5, class_weight="balanced".

For the first-stage classification, models were trained using the filtered dataset, encompassing 384 selected genomes. For the second stage, separate BRF models were trained using only isolates corresponding to the serogroups within a single seroclass. Likewise, only the similarity data derived from proteins from representative samples of serogroups of that seroclass were included as input features for model training in this stage.

Model evaluation was performed using distinct cross-validation schemes appropriate to the sample size of each classification level. The first-stage models were evaluated using 5-fold cross-validation, while the second-stage models were assessed using a leave-one-out (LOO) validation approach, chosen to account for the limited number of positive samples available for certain serogroups. The predictive performance of all models was evaluated based on accuracy, precision, recall, and F1-score metrics.

2.5 Reproducibility and data availability

All scripts, trained models, and feature matrices are available in a public GitHub repository (https://github.com/Spidey2004/Modelo_Leptospira). Genome accession numbers and associated serogroup metadata are listed in the Supplementary Tables S1 and S2.

3. Results

3.1 Dataset composition and exploratory analysis of the *rfb*-derived feature space

The feature matrix used to develop a machine learning model capable of serologically classifying pathogenic *Leptospira* isolates based on the gene composition of the *rfb* locus was comprised of 384 leptospiral samples (rows) and 549 proteins encoded in the *rfb* locus (columns). The selected samples were distributed across four major seroclasses and 22 distinct serogroups (Table 2). Among these, seroclass 1 was the most represented, encompassing 138 genomes assigned to nine serogroups. Within this class, the serogroups *Icterohaemorrhagiae* (43 genomes) and *Pyrogenes* (28 genomes) were the most prevalent, followed by *Canicola* (23 genomes) and *Ballum* (19 genomes). Seroclass 2 comprised 129 genomes distributed across six serogroups, with *Pomona* (43 genomes) and *Grippotyphosa* (39

genomes) being the most represented. Seroclass 3 contained 86 genomes assigned to three serogroups, primarily from the *Sejroe* (39 genomes). Finally, Seroclass 4 included 31 genomes, divided among three serogroups, with *Tarassovi* (18 genomes) as the most frequent representatives. Across all samples, *Leptospira interrogans* was the most represented species (199 genomes), followed by *L. borgpetersenii* (83 genomes), *L. kirschneri* (40 genomes), and *L. santarosai* (38 genomes). Among the 22 serogroups analyzed, 15 included isolates from at least two distinct species, indicating substantial interspecies distribution of serogroup classification. Only serogroups Ballum, Celledoni, Ranarum, Cynopteri, Djasiman, Panama, and Shermani were each represented by a single species. Similarly, among the nine *Leptospira* species included in the dataset, only *L. alstoni* and *L. mayottensis* were restricted to a single serogroup.

Table 2: Distribution of *Leptospira* isolates among classes, serogroups and species included in the machine learning analyses.

Seroclass*	Serogroup	*Lale*	*Lals*	*Lbor*	*Lint*	*Lkir*	*Lmay*	*Lnog*	*Lsan*	*Lwei*	Total
1 (138)	Ballum			19							19
	Canicola				22	1					23
	Celledoni									1	1
	Icterohaemorrhagiae				42	1					43
	Javanica	1		13					1		15
	Manhao	2								1	3
	Pyrogenes				22			1	5		28
	Ranarum		2								2
	Sarmin								3	1	4
2 (129)	Australis				14			3			17
	Autumnalis				19	3		1			23
	Cynopteri					2					2
	Djasiman				2						2
	Grippotyphosa				13	21			5		39
	Panama							3			3
	Pomona			3	30	10					43
3 (86)	Hebdomadis	1			19				1		21
	Mini			14	1	2	3		5	1	26
	Sejroe			27	9				3		39
4 (31)	Bataviae				6			1			7
	Shermani								6		6
	Tarassovi	1		7					9	1	18
	Total	5	2	83	199	40	3	9	38	5	384

*total number of samples in each seroclass indicated in parentheses.

Lale: *L. alexanderi*; *Lals*: *L. alstonii*; *Lbor*: *L. borgpetersenii*; *Lint*: *L. interrogans*; *Lkir*: *L. kirschneri*; *Lmay*: *L. mayottensis*; *Lnog*: *L. noguchii*; *Lsan*: *L. santarosai*; *Lwei*: *L. weilii*.

A principal component analysis (PCA) was performed to investigate the overall distribution of *Leptospira* samples based on the gene composition of the *rfb* locus (Figure 1). The first two principal components explained the largest proportion of variance in the dataset and revealed clustering patterns largely consistent with the seroclass structure defined by Ferreira et al. (2024) (Figure 1A). PCA was further applied separately to each seroclass to examine the distribution and clustering of their respective serogroups (Figure 1B–E). In most cases, serogroups formed well-defined and cohesive clusters. However, within seroclass 2, serogroups such as Autumnalis, Pomona, and Grippotyphosa displayed greater dispersion and partial overlap, indicating reduced separation compared to other serogroups.

3.2. *Leptospira* serogroup machine learning classifier

The procedure for predicting the serogroup of a *Leptospira* isolate was designed as a two-step hierarchical classification pipeline. In the first step, each isolate is assigned to one of four major seroclasses previously defined by Ferreira et al. (2024). Then, In the second step, the isolate is classified into one of the serogroups belonging to the seroclass to which it had been assigned in the first stage. Below, we present the analysis of the data and models generated in each stage.

3.2.1. First stage classification (seroclass level)

In this stage, classifiers were generated to assign *Leptospira* isolates to one of the four seroclasses defined by Ferreira et al. (2024). For each seroclass, an independent binary classification model was trained using the complete dataset (384 samples), with targets indicating whether a sample belonged or not to that specific seroclass. This binary approach was chosen to account for potential cases in which a new isolate might not belong to any of the previously characterized seroclasses. A conventional multiclass model would force such samples into one of the existing seroclasses, potentially leading to misclassification.

After model training using the BRF algorithm, performance was evaluated through 5-fold cross-validation. The models correctly assigned all samples to their respective seroclasses, achieving perfect classification performance across all evaluated metrics. This result is consistent with the clear separation of seroclasses observed in the PCA analysis (Figure 1a), which indicated minimal inter-class overlap in the *rfb*-derived feature space.

3.2.2. Second stage classification (serogroup level)

In the second stage of the classification pipeline, BRF models were trained to assign *Leptospira* isolates to specific serogroups within each of the four seroclasses. Each model was evaluated individually using the leave-one-out cross-validation strategy, and its performance was assessed through accuracy, precision, recall, and F1-score metrics (Table 3). Models for a few serogroups from seroclass 1 (Celledoni, Manhao, Ranarum, and Sarmin) and seroclass 2 (Cynopteri, Djasiman, and Panama) were not generated due to insufficient data for proper training.

Overall, the models achieved high predictive performance, with an average accuracy of 0.967, precision of 0.961, recall of 0.909, and F1-score of 0.931. Within Class 1 (n = 138), which contained the largest number of serogroups, the majority of models exhibited excellent performance, with F1-scores above 0.95 for serogroups Ballum, Icterohaemorrhagiae and Pyrogenes. Only the models for the Javanica and Canicola serogroup had a low F1-score compared to other models (above 0.85), but it was still considered high.

For Class 2 (n = 129), F1-scores ranged between 0.895 and 0.975, with the best results observed for the Grippotyphosa (F1 = 0.975), Autumnalis (F1 = 0.957) and Pomona (F1 = 0.966) serogroups. Slightly lower performance was recorded for the Australis serogroup, which yielded F1-scores of 0.895. Class 3 (n = 86) included serogroups Sejroe, Mini, and Hebdomadis, all of which achieved robust performance with F1-scores above 0.95, demonstrating consistent discriminative signal across these closely related serogroups. Finally, Class 4 (n = 31), which contained fewer samples, showed more variable results, with F1-scores ranging from 0.8 for Shermani to 0.914 for Tarassovi.

Table 3: Performance of the models from the second-stage (serogroup classification)

Seroclass*	Serogroup	# samples	Accuracy	Precision	Recall	F1-score
	Ballum	19	1.000	1.000	1.000	1.000
	Canicola	23	0.949	0.870	0.833	0.851
	Celledoni	1	not evaluated**			
	Icterohaemorrhagiae	43	0.971	0.953	0.953	0.953
1 (138)	Javanica	15	0.971	0.867	0.867	0.867
	Manhao	3	not evaluated**			
	Pyrogenes	28	1.000	1.000	1.000	1.000
	Ranarum	2	not evaluated**			
	Sarmin	4	not evaluated**			
	Australis	17	0.969	1.000	0.810	0.895
	Autumnalis	23	0.984	0.957	0.957	0.957
	Cynopteri	2	not evaluated**			
2 (129)	Djasiman	2	not evaluated**			
	Grippotyphosa	39	0.984	1.000	0.951	0.975
	Panama	3	not evaluated**			
	Pomona	43	0.977	0.977	0.955	0.966
	Hebdomadis	21	0.977	0.905	1.000	0.950
3 (86)	Mini	26	0.977	1.000	0.929	0.963
	Sejroe	39	1.000	1.000	1.000	1.000
	Bataviae	7	0.935	1.000	0.778	0.875
4 (31)	Shermani	6	0.903	1.000	0.667	0.800
	Tarassovi	18	0.903	0.889	0.941	0.914
	Mean performance		0.967	0.961	0.909	0.931

*total number of samples in each seroclass indicated in parentheses.
**not evaluated because of the low number of samples in this serogroup

A more detailed evaluation was performed by examining the prediction outcome for each sample during leave-one-out cross-validation (Supplementary Table S5) across all models. A summary of these results is presented in Table 4, where rows represent the predictive models trained for individual serogroups, whereas columns correspond to the true serogroup assignment of the samples. The values indicate the number of samples predicted as positive in each category. Overall, the models predominantly assigned positive predictions to samples belonging to their respective serogroups, as evidenced by the higher counts observed along the main diagonal within each seroclass block. Off-diagonal entries represent cross-classifications, reflecting instances in which samples from one serogroup were predicted as positive by a model trained on another serogroup. These misclassifications were generally infrequent, supporting the overall discriminatory capacity of the models. However, notable exceptions were observed among serogroups represented by a limited number of samples. Specifically, all samples from the Celledoni, Cynopteri, and Panama serogroups were systematically misclassified as Javanica, Pomona, and Australis, respectively. This complete reassignment suggests a strong overlap in *rfb* locus gene composition between these pairs of serogroups, indicating reduced discriminatory signal within the selected genomic features. At the same time, the small sample size of these underrepresented serogroups likely contributed to this behavior by limiting the models' ability to learn robust and distinctive patterns. Increasing the number of representative genomes for these serogroups would likely improve model calibration and enhance classification reliability.

Table 4: Distribution of samples classified as positive by each model across serogroups.

	Models	Ball (n=19)	Cani (n=23)	Cell (n=1)	Icte (n=43)	Java (n=15)	Manh (n=3)	Pyro (n=28)	Rana (n=2)	Sarm (n=4)
							Number of positive samples			
1	Ballum	19	0	0	0	0	0	0	0	0
	Canicola	0	20	0	4	0	0	0	0	0
	Icterohaemorrhagiae	0	2	0	41	0	0	0	0	0
	Javanica	0	0	1	0	13	1	0	0	0
	Pyrogenes	0	0	0	0	0	0	28	0	0

	Models	Aust (n=17)	Autu (n=23)	Cyno (n=2)	Djas (n=2)	Grip (n=39)	Pana (n=3)	Pomo (n=43)
2	Australis	17	1	0	0	0	3	0
	Autumnalis	0	22	0	1	0	0	0
	Grippotyphosa	0	0	0	1	39	0	1
	Pomona	0	0	2	0	0	0	42

	Models	Hebd (n=21)	Mini (n=26)	Sejr (n=39)
3	Hebdomadis	19	0	0
	Mini	2	26	0
	Sejroe	0	0	39

	Models	Bata (n=7)	Sher (n=6)	Tara (n=18)
4	Bataviae	7	0	2
	Shermani	0	6	3
	Tarassovi	0	1	16

Ball: Ballum; Cani: Canicola; Cell: Celledoni; Icte: Icterohaemorrhagiae; Java: Javanica; Manh: Manhao; Pyro: Pyrogenes; Rana: Ranarum; Sarm: Sarmin; Aust: Australis; Autu: Autumnalis; Cyno: Cynopteri; Djas: Djasiman; Grip: Grippotyphosa; Pana: Panama; Pomo: Pomona; Hebd: Hebdomadis; Sejr: Sejroe; Bata: Bataviae; Sher: Shermani; Tara: Tarassovi.

3.3. Validation of the serogroup classification model

Genome samples that were not included in the training dataset and were published after the development of the classification model were used as an independent validation set. The prediction results obtained for these genomes are presented in Table 5. Overall, the models demonstrated high predictive performance on this validation dataset. For most genomes, the predicted serogroup matched the true serogroup, with consistently high prediction probabilities, frequently above 0.90. Correct classifications were observed across multiple species, including *L. interrogans*, *L. kirschneri* and *L. borgpetersenii*, demonstrating that the predictive signal derived from the *rfb* locus is consistent across different pathogenic lineages. This pattern indicates strong generalization ability and suggests that the model successfully captures robust genomic signatures associated with serogroup classification. A single discrepancy was observed for strain L4 (serogroup Djasiman), which was predicted as Grippotyphosa. This misclassification is likely attributable to the limited representation of the Djasiman serogroup in the training dataset (n = 2), which prevented the development of a dedicated classification model for this serogroup and restricted the learning of distinctive genomic patterns. Consequently, Djasiman isolates were classified based on their similarity to models trained for other serogroups. Notably, the relatively low prediction probability assigned to this sample (0.501) indicates classification uncertainty and may serve as a useful indicator for flagging potential misclassifications in practical applications.

3.4. Relevant genes for serogroup classification

Because we employed a tree-based algorithm (Random Forest), it was possible to identify which genes (features) were most frequently used in the decision-making process across the ensemble of trees composing each model. By projecting the genomic positions of the most informative genes onto the *rfb* locus of representative strains from each serogroup (Figure 2), we observed that the majority of high-importance features clustered within the first half of the locus. This non-random distribution suggests that the proximal region of the *rfb* locus concentrates key genetic determinants underlying serogroup specificity.

Table 5: Classification of leptospiral samples from validation dataset

Accession	Species	Strain	True serogroup	Prediction	Prediction probability
GCF_041647705.1	*L. kirschneri*	H9	Australis	Australis	0.780
GCF_041647755.1	*L. interrogans*	H44	Pomona	Pomona	0.998
GCF_045159415.1	*L. interrogans*	Ballico	Icterohaemorrhagiae	Icterohaemorrhagiae	0.999
GCF_046529585.1	*L. interrogans*	Jez Bratislava	Australis	Australis	0.980
GCF_046848885.1	*L. interrogans*	Djasiman	Australis	Australis	1.000
GCF_046848895.1	*L. interrogans*	Van Tienen	Bataviae	Bataviae	0.940
GCF_046848905.1	*L. interrogans*	L4	Djasiman	Grippotyphosa	0.501
GCF_048973995.1	*L. borgpetersenii*	BU-20	Sejroe	Sejroe	0.990
GCF_048974015.1	*L. borgpetersenii*	BU-18	Sejroe	Sejroe	0.990
GCF_048974035.1	*L. borgpetersenii*	BU-19	Sejroe	Sejroe	0.990
GCF_048974055.1	*L. borgpetersenii*	BS-39	Sejroe	Sejroe	0.998
GCF_049953905.1	*L. interrogans*	L2	Mini	Mini	0.930
GCF_049954495.1	*L. interrogans*	L3	Hebdomadis	Hebdomadis	1.000
GCF_049954505.1	*L. borgpetersenii*	USDA CAL-010	Sejroe	Sejroe	0.967
GCF_049955055.1	*L. interrogans*	USDA BTL-020	Australis	Australis	1.000
GCF_049955065.1	*L. borgpetersenii*	USDA ICL-020	Tarassovi	Tarassovi	1.000
GCF_049955695.1	*L. interrogans*	L7	Pyrogenes	Pyrogenes	1.000
GCF_049955705.1	*L. interrogans*	RJ19115	Pomona	Pomona	0.999
GCF_049956375.1	*L. interrogans*	USDA ARL-010	Icterohaemorrhagiae	Icterohaemorrhagiae	0.999
GCF_049956385.1	*L. interrogans*	USDA GRL-020	Sejroe	Sejroe	0.988
GCF_049956975.1	*L. interrogans*	USDA MIL-020	Grippotyphosa	Grippotyphosa	0.929
GCF_049957005.1	*L. interrogans*	USDA PYL-010	Canicola	Canicola	1.000
GCF_049957935.1	*L. interrogans*	USDA ATL-010	Bataviae	Bataviae	0.940
GCF_049957945.1	*L. borgpetersenii*	USDA ARL-050	Ballum	Ballum	1.000
GCF_049958755.1	*L. interrogans*	USDA SJL-060	Autumnalis	Autumnalis	0.938
GCF_049958765.1	*L. interrogans*	USDA POL-010	Australis	Australis	0.980
GCF_052161685.1	*L. interrogans*	USDA HBL-010	Icterohaemorrhagiae	Icterohaemorrhagiae	0.945
GCF_052161695.1	*L. interrogans*	USDA BML-011	Canicola	Canicola	1.000
GCF_052161705.1	*L. interrogans*	USDA TAL-010	Canicola	Canicola	1.000
GCF_052161715.1	*L. interrogans*	USDA SJL-010	Canicola	Canicola	1.000

Proteins exhibiting feature importance values above 0.03, the threshold adopted to define the informative features, totaled 91 (Supplementary Table S6). Most of these proteins are associated with carbohydrate biosynthesis and modification pathways, including glycosyltransferases, class I SAM-dependent methyltransferases, aminotransferases of the DegT/DnrJ/EryC1/StrS family, and multiple oxidoreductases such as SDR family proteins and NAD-dependent epimerase/dehydratases. For some serogroups, including Sejroe, Pomona, Bataviae, and Shermani, discrimination was driven by highly specific markers that were consistently present in isolates belonging to these serogroups but largely absent in others. In contrast, for serogroups such as Australis and Grippotyphosa, classification performance was primarily supported by the consistent absence of particular genes in the ingroup coupled with their high prevalence among other serogroups. Notably, some protein IDs emerged as

important features across multiple serogroup-specific models, while certain serogroups lacked strictly exclusive markers. These observations indicate that discrimination is not solely dependent on uniquely serogroup-specific genes. Rather, they support a model in which serogroup differentiation is driven by combinatorial patterns of gene presence and absence within the *rfb* locus, constituting the underlying genomic basis of serological diversity.

4. Discussion

In this study, we proposed and validated a genomic-based machine learning framework for serogroup classification in *Leptospira*. To the best of our knowledge, this is the first study to apply machine learning methods to develop predictive classifiers specifically targeting serogroup-level identification in this genus based solely on genomic data. By leveraging the antigenic diversity encoded in the *rfb* locus, which underlies the biosynthesis of the O-antigen component of lipopolysaccharides (LPS) (Mitchison et al., 1997), our models effectively captured the genetic determinants that distinguish major serological groups. The two-step hierarchical strategy, first classifying samples into four broader antigenic classes and subsequently refining the prediction at the serogroup level, demonstrated robust performance, with perfect score for the first stage and average F1-scores of 0.931 for second stage. These values indicate that the genomic features derived from the *rfb* locus encode sufficient discriminatory power to accurately predict serological identity at serogroup level. In this sense, our results provide genomic validation of the long-assumed link between *rfb* organization and serological phenotype (Adler and de la Peña Moctezuma, 2010; Bulach et al., 2000; de la Peña-Moctezuma et al., 1999; Lehmann et al., 2014), bridging classical and molecular serological classification.

An important consideration when interpreting the misclassification for some models concerns the accuracy of serological annotations used for model training. Because traditional methods such as MAT and CAAT are susceptible to cross-reactivity and subjective interpretation (Adler and de la Peña Moctezuma, 2010; Levett, 2001), inconsistencies in serogroup labeling may be present in public databases. Such misclassifications introduce noise into the training dataset, which can artificially limit the model's performance or produce apparent misclassifications. Therefore, the performance metrics reported here likely represent a lower bound of the true predictive potential of genomic data, emphasizing the importance of improving and standardizing serological metadata for future studies.

Despite its promising results, this study has some limitations. First, the representation of certain serogroups remains limited in public genomic databases, which may restrict model generalizability, particularly for rare or newly emerging variants. Second, although the binary classification strategy reduces the risk of forcing isolates into predefined categories, it underscores the need for complementary mechanisms capable of detecting potentially novel serogroups that fall outside established classifications. Future research may address these limitations through the continued expansion of whole-genome sequencing efforts for *Leptospira* isolates, conducted in parallel with standardized serological characterization, and the incorporation of unsupervised clustering approaches, which could facilitate the detection of previously unrecognized antigenic patterns and contribute to the refinement of the proposed hierarchical classification structure.

Since the proposed models rely on genomic sequence data as input, such information may not be readily available in routine clinical settings, potentially limiting their immediate applicability in diagnostic workflows. However, the analysis of feature importance provides an important translational advantage. In particular, genes consistently exhibiting high importance and strong serogroup

specificity could serve as candidates for the development of PCR-based diagnostic tools, enabling rapid and cost-effective serological classification without the need for whole-genome sequencing. Such an approach could complement and refine previously proposed molecular typing strategies (Cai et al., 2010; Wenderlein et al., 2024) by providing a data-driven framework for marker selection grounded in machine learning–based feature prioritization. Therefore, beyond predictive performance, the feature importance analysis offers a pathway toward improving and modernizing molecular diagnostic methods for *Leptospira* serogroup classification.

Finally, we propose the term "seroclass" to designate a higher-order grouping of *Leptospira* serogroups. Serogroups within the same seroclass exhibit greater gene composition similarity at the *rfb* locus and frequent antigenic cross-reactions than those in different groups, indicating an intermediate hierarchical level not captured by the current serological framework. This terminology refines previous use of broader terms such as "cluster" (Chinchilla et al., 2023) or "class" (Ferreira et al., 2024) and provides a clearer structure for interpreting serological diversity.

5. Conclusion

In conclusion, our results demonstrate that machine learning models trained on *rfb*-derived genomic features can accurately reproduce the traditional serological classification of *Leptospira*. This approach offers a scalable, standardized, and reproducible alternative to serological assays, and contributes in paving the way toward a genomics-informed system for classification and surveillance of this complex and diverse bacterial genus.

Acknowledgements

We would like to thank the teams from Bioinformatics Multidisciplinary Environment (BioME/IMD) at UFRN, Centro de Processamento de Alto Desempenho (CEPAD/ICB) at UFMG, and High-Performance Computing Center (NPAD) at UFRN for the support on the computational resource. We are also grateful to NCBI and the Institut Pasteur teams for the curation and maintenance of the data used in this work.

Data statement

All genomic data used in this work are hosted in NCBI and BIGSdb, which are public databases.

Declaration of competing interest

The authors declare that they have no competing interests.

References

Adler, B., de la Peña Moctezuma, A., 2010. Leptospira and leptospirosis. Vet. Microbiol., Zoonoses: Advances and Perspectives 140, 287–296. https://doi.org/10.1016/j.vetmic.2009.03.012

Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. J. Mol. Biol. 215, 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Bulach, D.M., Kalambaheti, T., de la Peña-Moctezuma, A., Adler, B., 2000. Lipopolysaccharide biosynthesis in Leptospira. J. Mol. Microbiol. Biotechnol. 2, 375–380.

Cai, C.-S., Zhu, Y.-Z., Zhong, Y., Xin, X.-F., Jiang, X.-G., Lou, X.-L., He, P., Qin, J.-H., Zhao, G.-P., Wang, S.-Y., Guo, X.-K., 2010. Development of O-antigen gene cluster-specific PCRs for rapid typing six epidemic serogroups of Leptospira in China. BMC Microbiol. 10, 67. https://doi.org/10.1186/1471-2180-10-67

Caimi, K., Ruybal, P., 2020. Leptospira spp., a genus in the stage of diversity and genomic data expansion. Infect. Genet. Evol. 81, 104241. https://doi.org/10.1016/j.meegid.2020.104241

Chinchilla, D., Nieves, C., Gutiérrez, R., Sordoillet, V., Veyrier, F.J., Picardeau, M., 2023. Phylogenomics of Leptospira santarosai, a prevalent pathogenic species in the Americas. PLoS Negl. Trop. Dis. 17, e0011733. https://doi.org/10.1371/journal.pntd.0011733

Ciurariu, E., Prodan-Barbulescu, C., Mateescu, D.-M., Tutac, P., Sorop, V.-B., Susan, M., Varga, N.-I., 2025. Diagnostic Advances in Leptospirosis: A Comparative Analysis of Paraclinical Tests with a Focus on PCR. Microorganisms 13. https://doi.org/10.3390/microorganisms13030667

Costa, F., Hagan, J.E., Calcagno, J., Kane, M., Torgerson, P., Martinez-Silveira, M.S., Stein, C., Abela-Ridder, B., Ko, A.I., 2015. Global Morbidity and Mortality of Leptospirosis: A Systematic Review. PLoS Negl. Trop. Dis. 9, e0003898. https://doi.org/10.1371/journal.pntd.0003898

de la Peña-Moctezuma, A., Bulach, D.M., Adler, B., 2001. Genetic differences among the LPS biosynthetic loci of serovars of Leptospira interrogans and Leptospira borgpetersenii. FEMS Immunol. Med. Microbiol. 31, 73–81. https://doi.org/10.1111/j.1574-695X.2001.tb01589.x

de la Peña-Moctezuma, A., Bulach, D.M., Kalambaheti, T., Adler, B., 1999. Comparative analysis of the LPS biosynthetic loci of the genetic subtypes of serovar Hardjo: Leptospira interrogans subtype Hardjoprajitno and Leptospira borgpetersenii subtype Hardjobovis. FEMS Microbiol. Lett. 177, 319–326.

Dos Santos Ribeiro, P., Carvalho, N.B., Aburjaile, F., Sousa, T., Veríssimo, G., Gomes, T., Neves, F., Blanco, L., Lima, J.A., de Oliveira, D., Jaiswal, A.K., Brenig, B., Soares, S., Ramos, R., Matiuzzi, M., Góes-Neto, A., Figueira, C.P., Costa, F., Ristow, P., Azevedo, V., 2023. Environmental Biofilms from an Urban Community in Salvador, Brazil, Shelter Previously Uncharacterized Saprophytic Leptospira. Microb. Ecol. 86, 2488–2501. https://doi.org/10.1007/s00248-023-02253-3

Drouin, A., Letarte, G., Raymond, F., Marchand, M., Corbeil, J., Laviolette, F., 2019. Interpretable genotype-to-phenotype classifiers with performance guarantees. Sci. Rep. 9, 4071. https://doi.org/10.1038/s41598-019-40561-2

Faine, S., Adler, B., Bolin, C., Perolat, P., 1999. "Leptospira" and leptospirosis. MediSci, Melbourne Vic Australia.

Fernandes, L.G.V., Stone, N.E., Roe, C.C., Goris, M.G.A., van der Linden, H., Sahl, J.W., Wagner, D.M., Nally, J.E., 2022. Leptospira sanjuanensis sp. nov., a pathogenic species of the genus Leptospira isolated from soil in Puerto Rico. Int. J. Syst. Evol. Microbiol. 72. https://doi.org/10.1099/ijsem.0.005560

Ferreira, L.C., Ferreira Filho, L. de F., Cosate, M.R.V., Sakamoto, T., 2024. Genetic structure and diversity of the rfb locus of pathogenic species of the genus Leptospira. Life Sci. Alliance 7, e202302478. https://doi.org/10.26508/lsa.202302478

Fu, L., Niu, B., Zhu, Z., Wu, S., Li, W., 2012. CD-HIT: accelerated for clustering the next-generation sequencing data. Bioinforma. Oxf. Engl. 28, 3150–3152. https://doi.org/10.1093/bioinformatics/bts565

Giraud-Gatineau, A., Dagbo, K., Pĕtrošová, H., Werts, C., Veyrier, F.J., Picardeau, M., 2025. Shaping the future of Leptospira serotyping. J. Med. Microbiol. 74, 002059. https://doi.org/10.1099/jmm.0.002059

Goldfarb, T., Kodali, V.K., Pujar, S., Brover, V., Robbertse, B., Farrell, C.M., Oh, D.-H., Astashyn, A., Ermolaeva, O., Haddad, D., Hlavina, W., Hoffman, J., Jackson, J.D., Joardar, V.S., Kristensen, D., Masterson, P., McGarvey, K.M., McVeigh, R., Mozes, E., Murphy, M.R., Schafer, S.S., Souvorov, A., Spurrier, B., Strope, P.K., Sun, H., Vatsan, A.R., Wallin, C., Webb, D., Brister, J.R., Hatcher, E., Kimchi, A., Klimke, W., Marchler-Bauer, A., Pruitt, K.D., Thibaud-Nissen, F., Murphy, T.D., 2025. NCBI RefSeq: reference sequence standards through 25 years of curation and annotation. Nucleic Acids Res. 53, D243–D257. https://doi.org/10.1093/nar/gkae1038

Haake, D.A., Levett, P.N., 2015. Leptospirosis in humans. Curr. Top. Microbiol. Immunol. 387, 65–97. https://doi.org/10.1007/978-3-662-45059-8_5

Hamond, C., Tibbs-Cortes, B., Fernandes, L.G.V., LeCount, K., Putz, E.J., Anderson, T., Camp, P., Stuber, T., Hicks, J., van der Linden, H., dos Santos Ribeiro, P., Bayles, D.O., Schlater, L.K., Nally, J.E., 2025. Leptospira gorisiae sp. nov, L. cinconiae sp. nov, L. mgodei sp. nov, L. milleri sp. nov and L. iowaensis sp. nov: five new species isolated from water sources in the Midwestern United States. Int. J. Syst. Evol. Microbiol. 75, 006595. https://doi.org/10.1099/ijsem.0.006595

Hunter, J.D., 2007. Matplotlib: A 2D Graphics Environment. Comput. Sci. Eng. 9, 90–95. https://doi.org/10.1109/MCSE.2007.55

Jolley, K.A., Maiden, M.C., 2010. BIGSdb: Scalable analysis of bacterial genome variation at the population level. BMC Bioinformatics 11, 595. https://doi.org/10.1186/1471-2105-11-595

Kalambaheti, T., Bulach, D.M., Rajakumar, K., Adler, B., 1999. Genetic organization of the lipopolysaccharide O-antigen biosynthetic locus of Leptospira borgpetersenii serovar Hardjobovis. Microb. Pathog. 27, 105–117. https://doi.org/10.1006/mpat.1999.0285

Ko, A.I., Goarant, C., Picardeau, M., 2009. Leptospira: the dawn of the molecular genetics era for an emerging zoonotic pathogen. Nat. Rev. Microbiol. 7, 736–747. https://doi.org/10.1038/nrmicro2208

Korba, A.A., Lounici, H., Kainiu, M., Vincent, A.T., Mariet, J.-F., Veyrier, F.J., Goarant, C., Picardeau, M., 2021. Leptospira ainlahdjerensis sp. nov., Leptospira ainazelensis sp. nov., Leptospira abararensis sp. nov. and Leptospira chreensis sp. nov., four new species isolated from water sources in Algeria. Int. J. Syst. Evol. Microbiol. 71. https://doi.org/10.1099/ijsem.0.005148

Lees, J.A., Mai, T.T., Galardini, M., Wheeler, N.E., Horsfield, S.T., Parkhill, J., Corander, J., 2020. Improved Prediction of Bacterial Genotype-Phenotype Associations Using Interpretable Pangenome-Spanning Regressions. mBio 11, e01344-20. https://doi.org/10.1128/mBio.01344-20

Lehmann, J.S., Matthias, M.A., Vinetz, J.M., Fouts, D.E., 2014. Leptospiral pathogenomics. Pathog. Basel Switz. 3, 280–308. https://doi.org/10.3390/pathogens3020280

Levett, P.N., 2001. Leptospirosis. Clin. Microbiol. Rev. 14, 296–326. https://doi.org/10.1128/CMR.14.2.296-326.2001

Medeiros, E.J. de S., Ferreira, L.C.A., Ortega, J.M., Cosate, M.R.V., Sakamoto, T., 2022. Genetic basis underlying the serological affinity of leptospiral serovars from serogroups Sejroe, Mini and Hebdomadis. Infect. Genet. Evol. 103, 105345. https://doi.org/10.1016/j.meegid.2022.105345

Mitchison, M., Bulach, D.M., Vinh, T., Rajakumar, K., Faine, S., Adler, B., 1997. Identification and characterization of the dTDP-rhamnose biosynthesis and transfer genes of the lipopolysaccharide-related rfb locus in Leptospira interrogans serovar Copenhageni. J. Bacteriol. 179, 1262–1267.

Nieves, C., Vincent, A.T., Zarantonelli, L., Picardeau, M., Veyrier, F.J., Buschiazzo, A., 2022. Horizontal transfer of the rfb cluster in Leptospira is a genetic determinant of serovar identity. Life Sci. Alliance 6, e202201480. https://doi.org/10.26508/lsa.202201480

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M., Perrot, M., Duchesnay, É., 2011. Scikit-learn: Machine Learning in Python. J. Mach. Learn. Res. 12, 2825−2830.

Picardeau, M., 2017. Virulence of the zoonotic agent of leptospirosis: still terra incognita? Nat. Rev. Microbiol. 15, 297–307. https://doi.org/10.1038/nrmicro.2017.5

Terpstra, W.J., World Health Organization, International Leptospirosis Society (Eds.), 2003. Human leptospirosis: guidance for diagnosis, surveillance and control. World Health Organization, Geneva.

Vincent, A.T., Schiettekatte, O., Goarant, C., Neela, V.K., Bernet, E., Thibeaux, R., Ismail, N., Mohd Khalid, M.K.N., Amran, F., Masuzawa, T., Nakao, R., Amara Korba, A., Bourhy, P., Veyrier, F.J., Picardeau, M., 2019. Revisiting the taxonomy and evolution of pathogenicity of the genus Leptospira through the prism of genomics. PLoS Negl. Trop. Dis. 13, e0007270. https://doi.org/10.1371/journal.pntd.0007270

Waskom, M., 2021. seaborn: statistical data visualization. J. Open Source Softw. 6, 3021. https://doi.org/10.21105/joss.03021

Wenderlein, J., Zitzl, T., Dufay-Simon, N., Cachet, N., Pantchev, N., Le Guyader, M., Fontana, C., Bomchil, N., Tronel, J.-P., Cupillard, L., Straubinger, R.K., 2024. Detection and Identification of Pathogenic Leptospira spp. Serogroups in Europe between 2017 and 2020 Applying a Novel Gene-Based Molecular Approach. Transbound. Emerg. Dis. 2024, 1101841. https://doi.org/10.1155/2024/1101841

Figure captions

Figure 1: Principal Component Analysis (PCA) of *Leptospira rfb*-locus genomic features. (A) PCA including 384 *Leptospira* genomes representing four distinct seroclasses. (B–E) PCA performed independently for genomes assigned to seroclasses 1 (B), 2 (C), 3 (D), and 4 (E). Each point represents a single genome. In panel A, colors and marker shapes denote seroclass membership, whereas in panels B–E they indicate serogroup classification within each seroclass.

Figure 2. Feature importance profiles for each serogroup-specific model. Each panel represents one serogroup model. The y-axis indicates the feature importance values assigned by the model, whereas the x-axis corresponds to the genomic position of genes along the *rfb* locus. Higher importance values indicate genes that contributed more strongly to the classification decision. The red horizontal line in all panels represents the importance threshold of 0.03 adopted for highlighting relevant features. Only features associated with gene presence are displayed; genes for which discriminatory power was driven by their characteristic absence in a given serogroup are described in Supplementary Table S6.

Supplementary Tables

Supplementary Table S1: Metadata for the 721 *Leptospira* strains included in this study.

Supplementary Table S2: Metadata for the *Leptospira* strains included in the validation dataset.

Supplementary Table S3: Gene identity feature matrix derived from the *rfb* locus.

Supplementary Table S4: Pairwise correlation matrix among *Leptospira* strains based on *rfb* locus gene profiles.

Supplementary Table S5: Serogroup classification probabilities for each sample during leave-one-out cross-validation.

Supplementary Table 6: High-importance features for each serogroup-specific model