Using Variable Window Sizes for Phylogenomic Analyses of Whole Genome

Alignments

Jeremias Ivan[1], Robert Lanfear[1]

[1]Research School of Biology, Australian National University, ACT, Australia

Corresponding author: Jeremias Ivan

Research School of Biology, Australian National University, ACT, Australia

Ivan.Jeremias@anu.edu.au

ABSTRACT

Many phylogenomic studies used non-overlapping windows to address gene tree discordance

across a set of aligned genomes. Recently, Ivan et al. (2025) proposed an information

theoretic approach to choose an optimal window size given the alignment. However, this

approach selects only a single fixed window size per chromosome, which is a useful first step

but fails to account for variation in the size of non-recombining regions along each

chromosome. Such variation is expected to occur due to the stochastic nature of

recombination as well as the variation in recombination rates along chromosomes. In this

study, we extend the approach of Ivan et al. (2025) to allow window sizes to vary across the

chromosome, using a splitting-and-merging strategy that allows for each window to be of an

arbitrary length. We showed that the new method outperformed the fixed-window approach

in recovering gene tree topologies on a wide range of simulated datasets. Applying the new

method on the genomes of seven *Heliconius* butterflies, we found that the average window

sizes for the group ranged between 538-808bp, but with a very similar distribution of gene

tree topologies compared to previous studies that used fixed window sizes. For the genomes

of great apes, the average window sizes ranged from 4.2kb to 6.2kb, with the proportion of

the major topology (i.e., grouping human and chimpanzee together) reaching approximately

80%. In conclusion, our study highlights the limitations of using a fixed window size when

recombination rates vary across the chromosomes, and proposes a splitting-and-merging

approach that allows for variable window sizes across whole genome alignments.

Keywords: gene tree discordance, recombination rate, simulation, splitting, merging, AIC

INTRODUCTION

Many phylogenomic studies of closely-related taxa begin by aligning the genomes of species

or individuals, and then use those alignments to infer evolutionary events such as speciation,

introgression, and demographic changes (e.g., Pease et al. 2016; Morales-Briones et al. 2021;

Lescroart et al. 2023; Herrig et al. 2024). As different sites in the alignments may reflect

different evolutionary histories (Maddison 1997; Degnan and Rosenberg 2009; Mallet et al.

2016; Hibbins et al. 2023; Steenwyk et al. 2023), methods that account for this variation

become important. For example, Thawornwattana et al. (2018, 2022) extracted 100-locus

blocks (where a locus was defined as a coding or non-coding region with a minimum length

of 100bp and spaced at least 2kb apart) and inferred the species tree for each block using BPP

(Flouri et al. 2018). This approach is useful for identifying introgressed regions, but it

excludes most parts of the genomes. To infer the complete histories of whole genome

alignments – which is important for analyses such as PhyloGWAS (Pease et al. 2016) –

methods that use Hidden Markov Models (HMMs) and Ancestral Recombination Graphs

(ARGs) are powerful for inferring site-specific histories by considering the similarities

between neighbouring sites, potentially under a population genetic model of evolutionary

change. However, these approaches have their own limitations, including accuracy and

scalability (Siepel and Haussler 2005; Brandt et al. 2024). A relatively simpler alternative is

to partition the genome alignment into non-overlapping blocks (or windows) and infer a

single gene tree for each window. This approach is commonly-used in phylogenomic studies

to assess various evolutionary processes, such as incomplete lineage sorting (ILS) and

introgression (e.g., Fontaine et al. 2015; Pease et al. 2016; Edelman et al. 2019; Meleshko et

al. 2021; Yang et al. 2021; Feng et al. 2023; Lescroart et al. 2023; Herrig et al. 2024).

However, one of the main challenges for this approach is to select a window size that best

reflects the recombination breakpoints of given alignment, with a trade-off between

concatenation preferring the primary tree topology (when the window size is too long, thus

recombination is ignored) and gene tree estimation error (when the window size is too short,

thus limiting the information available to estimate a tree for each window). Most previous

studies have selected a single window size somewhat arbitrarily, in an attempt to balance

these tradeoffs. However, the selection of window size can have significant effects on

downstream inferences. For example, Ivan et al. (2025) showed that in the genomes of great

apes (i.e., human, chimpanzee, gorilla, orangutan), only 50% of gene trees grouped human

and chimpanzee together when using 250bp non-overlapping windows across all

chromosomes. However, this proportion increased to almost 100% when the window size

was 64kb. This highlights the importance of carefully selecting a window size in such

analyses.

Recently, Ivan et al. (2025) proposed that the AIC (Akaike Information Criteraion; Akaike

1974) – which reflects how the substitution models and the trees fit the observed alignment

data – could be used to select a window size in phylogenomic non-overlapping window

analyses. They showed that the AIC was a good predictor of the accuracy of different

window sizes in recovering the true histories of individual sites along simulated

chromosomes. Applying this approach on empirical datasets, Ivan et al. (2025) further

showed that the best window sizes for chromosome alignments of *Heliconius* butterflies

ranged between ≤125-250bp, while the window sizes for great apes ranged between 500bp-

1kb (and 4kb for the mitochondrial genome). These windows are significantly shorter than

the ones used in previous studies (e.g., Edelman et al. (2019) used 10kb and 50kb windows to

analyse the genomes of *Heliconius* butterflies), but still longer than the estimated lengths of

non-recombining blocks – sometimes called 'coalescent genes' or 'c-genes' – of each group

(e.g., Springer and Gatesy (2016) estimated the average length of c-genes between human,

chimpanzee, and gorilla to be ~109bp, but see also Edwards et al. (2016)). While the AIC

provides a less arbitrary way to choose a window size for non-overlapping window analysis,

the method proposed by Ivan et al. (2025) has two key limitations: it was evaluated using

simulated chromosomes with constant recombination rates, and it selects a single fixed

window size for each chromosome. A single fixed window size is problematic as some non-

recombining blocks will be much longer than others even under constant recombination rate,

due to the stochastic nature of recombination. And since recombination rates vary along the

chromosomes for a variety of reasons (Jensen-Seaman et al. 2004; McVean et al. 2004; Chan

et al. 2012; Edelman et al. 2019), the best window size may also vary systematically along

chromosomes.

In this study, we extend the method proposed in Ivan et al. (2025) to allowing the window

size to vary along chromosomes via a splitting-and-merging approach (Fig. 1). We do this by

first partitioning the alignment into non-overlapping windows of a fixed size. Then, we split

each window into two sub-windows and build gene trees for both the initial window and the

two sub-windows. If the total AIC of the two sub-windows is better than the AIC of the initial

window, we keep the split; otherwise, we retain the initial window. Once no further AIC

improvement could be achieved through splitting, we then iteratively merge neighbouring

windows. Similar to the splitting step, for every pair of neighbouring windows, we compare

the total AIC of the two initial windows and the AIC of the newly-merged window, keeping

the merge only if the AIC improves. Through this splitting-and-merging procedure, we

generate a set of non-overlapping windows that can vary in length along the chromosome and

should better represent the underlying recombination breakpoints of the alignment compared

to using a single window size. This approach is similar to GARD (Genetic Algorithm for

Recombination Detection) (Kosakovsky Pond et al. 2006) that aims to detect recombination

breakpoints from multiple sequence alignments. However, our method differs in using: (i)

Maximum Likelihood (ML) instead of Neighbour-Joining (NJ) to build the gene trees; (ii) the

AIC instead of the corrected AIC (AIC_c) to score the split because both criteria yielded very

similar results in prior work (Ivan et al. 2025, results not shown); and (iii) a greedy algorithm

instead of a genetic algorithm to locate the putative genomic breakpoints, which allows our

method to scale to whole genome alignments in reasonable amount of time.

Figure 1. *Overview of the splitting-and-merging approach proposed in this study. The splitting step is iteratively applied to every window in the alignment – selecting split configuration with the best AIC score for each window – until no further improvement can be achieved from any new split. The merging step is then iteratively applied to every possible pair of neighbouring windows in the alignment, prioritising merges with the biggest AIC improvement for each iteration until no further improvement can be achieved from any new merge.*

In order to assess if using variable windows sizes would improve the accuracy of non-

overlapping window method to recover the true topology of every site in the alignment (as

compared to when using only one fixed window size), we benchmarked our splitting-and-

merging approach with two simulation schemes: one with homogeneous recombination rates

from Ivan et al. (2025), and one with heterogeneous recombination rates generated in this

study. For each approach, we measured the accuracy of the final set of window sizes using

the two metrics used in Ivan et al. (2025): *site accuracy* (i.e., the proportion of sites from

non-overlapping windows that recover the true simulated topology) and *RMSE* (i.e., the root

mean squared error of the observed tree topology distribution from non-overlapping windows

compared to the true simulated distribution). Finally, we applied our method to the empirical

datasets from *erato-sara* clade of *Heliconius* butterflies and great apes to infer the variation

of window sizes along each chromosome alignment, and assess how this variation affects the

distribution of gene tree topologies of each group.

MATERIALS & METHODS

Overview

Our method works by first partitioning the genome alignment into non-overlapping windows

of a fixed size. We then iteratively refine this partitioning by either splitting each window

into two smaller sub-windows or merging two neighbouring windows together, keeping the

split and/or merge only if it improves the AIC of the gene trees. This approach generates a

partitioning scheme in which the window size can be longer in some regions due to merging,

and shorter in other regions due to splitting. In order to evaluate if using variable window

sizes improves the accuracy of non-overlapping windows to recover the true histories from

the alignment (as compared to using only one fixed window size), we ran our splitting-and-

merging approach on simulated alignments with varying recombination rates and measured

the accuracy of the final sets of window sizes using the two metrics used in Ivan et al. (2025):

site accuracy and RMSE. We then applied our method on the genomes of *erato-sara* clade of

Heliconius butterflies (Edelman et al., 2019) and great apes (Waterson et al. 2005; Locke et

al. 2011; Scally et al. 2012; Schneider et al. 2017). All the code necessary to reproduce the

methods in this paper is available at https://github.com/jeremiasivan/SimNOW.

Simulating Chromosomes with Varying Recombination Rates

We designed seven scenarios to simulate 10Mb alignments: four with homogeneous (i.e.,

constant) recombination rates and three with heterogeneous recombination rates (Fig. 2). We

used ms (Hudson 2002) to simulate the gene trees and AliSim (Ly-Trong et al. 2022) to

simulate the alignments in every scenario. The four scenarios with homogeneous

recombination rates (Fig. 2) were taken directly from Ivan et al. (2025). The three scenarios

with heterogenous recombination rates comprise: (i) *Scenario 1* (Fig. 2), a simple artificial

scenario in which the first half of the 10Mb chromosome was simulated with no

recombination, and the second half was simulated with an intermediate recombination rate of

$\rho=100$; (ii) *Scenario 2* (Fig. 2) crudely mimics a pattern of recombination often seen in

empirical datasets where the recombination rates are higher towards the ends of the

chromosomes and lower towards the center, in which 2.5Mb at each end of the chromosome

was simulated with $\rho=50$, and the central 5Mb region was simulated with $\rho=0$; (iii) *Scenario*

3 (Fig. 2) mimics the same scenario, but with a 1Mb region at each end of the chromosome

was simulated with $\rho=200$, the next 2.5Mb region at each end simulated with $\rho=50$, and the

central 3Mb of the chromosome simulated with $\rho=0$.

Figure 2. Scenarios for simulating chromosome alignments. Simulated chromosomes with

homogenous recombination rates were derived from Ivan et al. (2025), while ones with

heterogeneous recombination rates were generated in this study. Gray: zero recombination

rate; green: low recombination rate; blue: intermediate recombination rate; red: high

recombination rate.

For each simulation, we simulated gene trees following the ms (Hudson 2002) parameters

used in Ivan et al. (2025) to mimic the evolutionary histories between seven *Heliconius*

butterflies with medium level of ILS. Then, we used these sets of gene trees as input for

AliSim (Ly-Trong et al. 2022) to simulate alignments under the Jukes-Cantor (JC) model

(Jukes and Cantor 1969) for computational efficiency. Note that we multiplied the coalescent

branch lengths used in ms by a factor of 0.005 to produce branch lengths in substitutions per

site used in AliSim. This scaling factor was shown previously to produce alignments that

closely mimic empirical alignments, with roughly 5% informative sites (Ivan et al. 2025). We

did ten independent replicates for each scenario, resulting in 70 simulated chromosomes (i.e.,

10 replicates for each of the 7 simulated chromosomes depicted in Figure 2).

Running Non-overlapping Window Analyses with Fixed Window Sizes

We ran non-overlapping window analysis on each simulated chromosome with 16 different

fixed window sizes, ranging from 100bp to 10Mb. Then, we built the gene tree for each

window using IQ-TREE v2.4.0 (Minh et al. 2020) with the following command: `iqtree2 -`

`S alndir -m JC -blmin 1/window_size`, where `-S alndir` refers to the directory

storing all window alignments, `-m JC` sets the substitution model to be JC (Jukes and

Cantor 1969), and `-blmin 1/window_size` sets the minimum branch length for all trees

to represent at least one substitution per branch, penalising gene trees that were inferred from

windows with limited phylogenetic signal (Ivan et al. 2025).

For each simulated chromosome, we then calculated the accuracy of each window size using

site accuracy (i.e., the proportion of sites from non-overlapping window analysis that

correctly recover the true simulated topology) and the RMSE of the gene tree distribution

from the non-overlapping window analysis compared to the simulated truth (Ivan et al. 2025).

As Ivan et al. (2025) proposed that the AIC is a good predictor of both accuracy metrics, we

also retrieved the AIC of each window size from each simulation and calculated the loss of

accuracy if we always chose the window size with the lowest AIC score (compared to the

window sizes with the highest site accuracy and/or lowest RMSE). To avoid unnecessary

computation, we retrieved the results for the 40 simulated chromosomes with homogenous

recombination rates from Ivan et al. (2025) and did not redo these analyses.

Running the Splitting-and-Merging Approach on Simulated Datasets

The splitting-and-merging approach requires a starting window size to be chosen by the user.

We evaluated two starting window sizes for this study: (i) the best fixed window size selected

by the AIC (Ivan et al. 2025) for each simulated chromosome, and (ii) a single window

comprising the entire chromosome (10Mb in this case). These relatively extreme values

represent two ends of a spectrum – the former tends to be very small as it should account for

the short non-recombining loci in the chromosome, while the latter is the largest possible

starting value. Applying both allows us to evaluate the sensitivity of the splitting-and-

merging approach to the choice of starting window size.

Once a starting window size has been selected, the splitting-and-merging approach proceeds

as follows for each simulated chromosome. Every initial window is divided into two sub-

windows using three size ratios: 0.25:0.75, 0.50:0.50, and 0.75:0.25 (e.g., if the initial

window size is 1kb and the ratio is 0.25:0.75, then the first sub-window would be 250bp

while the second sub-window would be 750bp). If the initial window was not divisible by the

given ratio, the length of the first sub-window was rounded up (e.g., if the initial window size

is 250bp and the ratio is 0.25:0.75, then the first sub-window would be 63bp while the second

sub-window would be 187bp). Next, we inferred the gene trees for the original window and

each of the three split ratios using IQ-TREE2 with the following command: `iqtree2 -s`

`file_fasta -blmin 1/window_size`, where `-s` specifies the alignment file and `-`

`blmin` sets the minimum branch length based on the window size. We also set the minimum

number of parsimony informative sites per window to be equal to the total number of

branches on the (unrooted) tree, ensuring, in principle, that each branch is supported by at

least one informative site. If either of the sub-windows from a split could not be analysed

(e.g., due to missing data or lack of parsimony informative sites), we skipped that splitting

configuration. We then compared the AIC scores of the resulting four partitioning schemes

(i.e., the original unsplit window and up to three split configurations) for each window and

kept the scheme with the lowest AIC for the next iteration. After applying this approach to

every window in the initial alignment, we iteratively repeated the splitting procedure until no

further splits led to an improvement in the AIC.

Following this, we used the AIC to guide a procedure in which neighbouring windows are

iteratively merged. For each pair of neighbouring windows in the chromosome, we first

calculated the change in AIC that would result from merging them into a single window. To

do this, we concatenated their alignments using SeqKit v2.4.0 (Shen et al. 2016) and inferred

the gene tree of the newly-merged window using IQ-TREE2 with the same parameters as

above. We then extracted the AIC of the merged window and compared it with the total AIC

of the two original (unmerged) windows. This procedure provided a list of neighbouring

windows whose merging would improve the AIC score. However, not all these merging

events were necessarily mutually compatible, because the AIC could be improved by

merging the same window with the neighbouring window to the left *and* to the right of it. To

resolve these conflicts, we used a greedy algorithm that prioritises merges with the biggest

AIC improvements. The algorithm proceeded by merging the pair of windows with the

largest AIC improvement, then removing the pair and all conflicting merges from the list of

merges that could improve the AIC score. This procedure was iterated until all possible

merges that improved the AIC score were exhausted. We then iteratively repeated the entire

merging step until we could find no pair of neighbouring windows whose merging would

improve the AIC score.

We measured the accuracy of the final partitioning scheme for each simulated chromosome

using site accuracy and RMSE (Ivan et al. 2025), and assessed if using variable window sizes

improved the two accuracy measures of non-overlapping window analysis compared to using

a single fixed window size. This method is available in the GitHub repository under

https://github.com/jeremiasivan/SimNOW/tree/main/codes/3_variable_window_size/.

Empirical Analyses

Erato-sara clade of Heliconius butterflies

We downloaded the genome alignments from six species of *erato-sara* clade of *Heliconius*

butterflies (with *H. melpomene* outgroup) from Edelman et al. (2019) following the steps

described in Ivan et al. (2025). While the best window sizes for this group ranged between

≤125-250bp according to the stepwise non-overlapping window approach (Ivan et al. 2025),

we found that using very short window sizes as the starting window sizes could lead to low

accuracy (see Results). Thus, we ran the splitting-and-merging approach on each of the 21

chromosome alignments with the whole chromosome (i.e., full concatenation) as the starting

window size.

After no further AIC improvement could be achieved from either splitting or merging, we

extracted the final set of window sizes for each chromosome and built the gene tree of each

window using IQ-TREE2 with the following command: `iqtree2 -s file_fasta -`

`blmin 1/window_size -bb 1000`, where `-bb` reflects the number of UFBoot replicates

(Hoang et al. 2018). We rebuilt the gene trees to extract the bootstrap support for each gene

tree. We then retrieved the topology of every gene tree, and compared the proportion of each

topology across the chromosomes to those from Edelman et al. (2019) and Ivan et al. (2025).

Great apes

We downloaded genome alignments of four great apes (human, chimpanzee, gorilla, and

orangutan) from UCSC Genome Browser (Kent et al. 2002) following the steps described in

Ivan et al. (2025). Similar to the analyses on the *Heliconius* butterflies, we ran the splitting-

and-merging approach on each of the 25 chromosomes (22 autosomes, chromosome X,

chromosome Y, and mitochondrial DNA) but with a minimum of 5 parsimony informative

sites per window, using the entire chromosome as the starting point. We then built the gene

trees based on the final partitioning scheme with 1,000 UFBoot replicates using IQ-TREE2

and calculated the topology distribution of the group across the chromosomes.

RESULTS

Variable window sizes outperform fixed window sizes on simulated datasets

For simulated chromosomes with zero recombination rate, non-overlapping window analyses

using either fixed window sizes (from Ivan et al. 2025) or variable window sizes (from the

splitting-and-merging approach proposed in this study) always prefer full concatenation, as

expected (Fig. 3-4). However, as variable window sizes consistently outperform fixed

window sizes in any other scenario (see below), we recommend to always use variable

window sizes when running non-overlapping window approach. For simulations with non-

zero homogeneous recombination rates, the variable window size approach increases the site

accuracy of non-overlapping window method by on average 3.0-4.7% across scenarios

compared to using fixed window sizes from the AIC (Fig. 3), and decreases the RMSE by on

average 0.05-0.35% across scenarios (except for $\rho = 2000$ with on average 0.14% RMSE

increase; Fig. 4). For simulated chromosomes with heterogeneous recombination rates, the

variable window size approach also outperforms the fixed window size method. Using

variable window sizes increases the site accuracy of non-overlapping window method by on

average 4.7-8.3% across scenarios compared to using fixed window sizes from the AIC (Fig.

3), and decreases the RMSE by on average 0.57-0.81% (Fig. 4). In one replicate from

Scenario 3, running the variable window size approach results in an 22.3% site accuracy

increase (Fig. 3) and 4.1% RMSE decrease (Fig. 4) compared to using a single fixed window

size. This trend is also reflected by the AIC scores, where partitioning schemes from the

variable window size approach have better AIC than the fixed window sizes (Fig. S1-S2).

Figure 3. *Site accuracy from running the non-overlapping window analyses with fixed and*

variable window sizes across 7 simulation scenarios (Fig. 2). Colour denotes the set of

window sizes used when running the non-overlapping window method, either using a fixed

window size (Ivan et al. 2025) or the splitting-and-merging approach with different starting

window sizes. Black lines connect the same replicates.

Figure 4. *RMSE from running the non-overlapping window analyses with fixed and variable*
window sizes across 7 simulation scenarios (Fig. 2). Colour denotes the set of window sizes
used when running the non-overlapping window method, either using a fixed window size
(Ivan et al. 2025) or the splitting-and-merging approach with different starting window sizes.
Black lines connect the same replicates.

Full concatenation is preferable as starting window size for splitting-and-merging

approach

In general, our simulations show that there is very little difference in accuracy when using

different starting window sizes for the splitting-and-merging algorithm. On average, the two

methods differ in site accuracy by 0.1-1.4% (Fig. 3) and in RMSE by 0.01-0.23% (Fig. 4).

However, there is one simulated chromosome from Scenario 3 where using full concatenation

as the starting window size has 12.48% higher site accuracy and 2.34% lower RMSE

compared to using the fixed window sizes from AIC. This suggests that in certain cases,

starting the splitting-and-merging algorithm with a small window size can lead it to become

stuck in a local optimum. We also found that starting the splitting-and-merging approach with

short windows often fails to recover the long chromosomal regions with zero recombination

rate as a single window (Fig. S3-S5 (left), Table S1). This is improved when starting the

algorithm with the whole chromosome as a single window (Fig. S3-S5 (right)). Additionally,

starting with a single concatenated window has the advantage of being far more

computationally efficient, as no prior computation needs to be done to choose the starting

window size. Because of this, we used this approach on the empirical datasets whose analyses

we describe in the following sections.

Fixed window sizes do not represent the size variation of non-recombining loci across

empirical chromosomes

As the stepwise non-overlapping window method from Ivan et al. (2025) excluded windows

that cannot be analysed due to missing data, we could not compare the AIC of the

partitioning schemes between the fixed window sizes and the variable window sizes. For

erato-sara clade of *Heliconius* butterflies, the variable window size approach produces

substantially different window size results compared to the fixed window size method of Ivan

et al. (2025). Using the variable window size method, the shortest window on each

chromosome ranges from 18bp to 40bp, while the longest window ranges from 31kb to

106kb (Table S1; Fig. S6). Based on the distributions of window sizes across chromosomes,

the best fixed window sizes selected by the stepwise non-overlapping window method (Ivan

et al. 2025) are 2-5x shorter than the average window sizes from the variable window size

approach, and much closer the first quartile of the distribution than the median (Fig. S7;

Table S1). Despite these differences, the ten most common topologies recovered from the

variable window size approach are consistent with those from previous studies (Edelman et

al. 2019; Ivan et al. 2025), accounting for 53.0% of the sites (or 40.8% of the windows) when

including all gene trees (Fig. 5, S8). The most common topology is T2 (which is regarded as

the species tree topology (Edelman et al. 2019)), followed by T1 and T3. When we filtered

out gene trees with <95 average UFBoot support, the top ten topologies account for 97.9% of

the sites (or 92.9% of the windows) from the highly-supported windows (Fig. 5, S8), with T1

as the most common topology. Our results highlight that the distribution of tree topologies

can differ substantially depending on whether it is quantified by individual windows or

individual sites. For example, the species tree topology (T2) that dominates the sex

chromosome (chr21) is only recovered by 15.37% of windows on that chromosome (followed

by 12.68% of T3 and 12.19% of T7; Table S2). However, these windows account for 26.85%

of the sites (followed by 18.44% of T3 and 13.16% of T7; Table S3), reflecting that windows

that are associated with these topologies tend to be longer than average.

Figure 5. Comparison of topology frequency of the most common tree topologies from the genomes of erato-sara clade of Heliconius between 50kb sliding windows (SW) and 10kb non-overlapping windows (NOW) from Edelman et al. (2019), stepwise NOW from Ivan et al. (2025), and the splitting-and-merging approach from this study. For splitting-and-merging approach, the percentages reflect the proportions of sites that support each topology, while the numbers in bracket reflect the proportions of windows that support each topology. Topologies

are rooted with H. melpomene (not shown) as the outgroup. The number under each method

reflects the total number of windows. Black dotted lines refer to other topologies not listed.

For the genomes of great apes, the shortest window ranges from 17bp to 141bp across

chromosomes, while the longest window ranges from 138kb to 4.5Mb, except for chrY

(chromosome Y) and mtDNA (mitochondrial DNA; Table S6; Fig. S9). For chrY, a major

proportion of the alignment was not split due to missing data (Ivan et al. 2025), driving the

average window size to reach almost 200kb. On the other hand, the 16kb-long mtDNA was

partitioned into 8 windows, with an average window size of 2.1kb (Table S6). Similar to the

results from the *Heliconius* butterflies, the best fixed window sizes from the stepwise non-

overlapping window method (Ivan et al. 2025) are 4-11x shorter than the average window

sizes from the splitting-and-merging approach (Fig. S10). When we consider all gene trees

across all chromosomes, the major topology (grouping human and chimpanzee together) is

supported by 78.9% of the sites (or 75.3% of the windows), while the minor topologies

account for 10.8% and 10.3% of the sites (or 12.5% and 12.2% of the windows), respectively

(Fig. 6, S11; Table S7-S8). This contrasts with the topology distribution of 60.7%, 19.8%,

and 19.2% when using the fixed window size approach from Ivan et al. (2025) (Fig. 6).

Similar to the results from Ivan et al. (2025), filtering out gene trees with <95 average

UFBoot support drives the proportion of the major topology to approximately 92%, while the

minor topologies account for 4.7% and 3.7% of the sites, respectively (Fig. 6; Table S7-S8).

Interestingly, for mtDNA, six of the eight windows recover the major topology (comprising

80.08% of the sites), while the other two windows result in each of the minor topologies

(Table S7-S8; Fig. S11). This variation of gene trees in mtDNA is likely caused systematic

error rather than biological processes (Richards et al. 2018), and serves as a useful reminder

of some of the limitations of current phylogenetic inference methods.

Figure 6. *Comparison of unrooted tree topology frequency from the genomes of great apes*

using different approaches: single gene tree inference and MAST (Wong et al. 2024), the

stepwise non-overlapping windows (NOW) that includes all gene trees and only highly-

supported trees (Ivan et al. 2025), and the splitting-and-merging approach from this study.

For splitting-and-merging approach, the percentages reflect the proportions of sites that

support each topology, while the numbers in bracket reflect the proportions of windows that

support each topology. The number under each method reflects the total number of windows.

Topologies are visualised using FigTree v1.4.4 (http://tree.bio.ed.ac.uk/software/figtree)

without branch lengths.

Variable window sizes can identify regions of biological importance

Similar to Edelman et al. (2019), we found that the proportion of T1 on the chromosomes of

Heliconius butterflies generally goes down as the chromosome becomes longer (R^2=0.743,

$\rho<0.01$ for all gene trees; and R^2=0.520, $\rho<0.01$ for highly-supported trees), while T2 is

positively correlated with the chromosome length (R^2=0.712, $\rho<0.01$ for all gene trees; and

R^2=0.392, ρ<0.01 for highly-supported trees) (Fig. S12). When each chromosome was

divided into 10 equally-sized bins along its length, we also observed that T1 has lower

proportion on the ends of the chromosomes, in contrast to T2 that shows the opposite pattern

(Fig. S12). Moreover, we identified two genomic regions that are dominated by the minor

topologies (i.e., neither T1 nor T2; Fig. 7). The first region spans nearly 2Mb on chromosome

2 and is dominated by T3 (Fig. 7a), while the second region comprises 376 windows and is

dominated by T4, covering approximately 400kb on chromosome 15 (Fig. 7b). The latter is

of particular interest as it spans the loci that control the wing colour pattern in *Heliconius*

butterflies (Edelman et al. 2019). To assess whether the region is best fit by a single tree

topology (as our splitting-and-merging approach recovered multiple different topologies

along this region despite being dominated by the T4 topology), we analysed the whole

~400kb region as one partition, and compared the AIC to the partitioning scheme recovered

from our algorithm. This analysis shows that analysing the region as a single partition has a

much worse AIC score than the partitioning scheme recovered by the variable window size

method (AIC score of 1928355 compared to 1913217, a difference of 15,138 units).

Figure 7. *Inversion topologies on (a) chromosome 2 and (b) chromosome 15 of erato-sara*

clade of Heliconius butterflies (Edelman et al. 2019). Colouring is based on Fig. S8.

DISCUSSION

The non-overlapping window method is a common approach to infer gene trees across a set

of aligned chromosomes. While the selection of window size is often arbitrary, a recent study

by Ivan et al. (2025) showed that the total AIC score of the gene trees is a good indicator of

the ability of different window sizes to recover the 'true' topology of each site in the aligned

chromosomes. In this study, we extended this approach by allowing the window size to vary

across the chromosomes. We achieved this by iteratively splitting each window and merging

neighbouring windows until the total AIC of the whole alignment stops improving from any

new split and/or merge (Fig. 1). Our results showed that using variable window sizes from

the splitting-and-merging algorithm had better accuracy in recovering the 'true' gene tree

topology from each site of simulated datasets compared to using fixed window sizes. The

fixed window sizes selected by the AIC had at least 70% site accuracy across simulated

chromosomes with homogeneous and heterogenous recombination rates (Fig. 3-4). Using

variable window sizes consistently increased this accuracy further by on average 3-8% across

scenarios (Fig. 3), reflecting that variable window sizes better fit the size variation of non-

recombining loci across the chromosomes (Fig. S3-S5). While the different starting window

sizes tested in this study did not significantly affect the accuracy of the splitting-and-merging

approach (Fig. 3-4), we proposed future phylogenomic analyses to start with full

concatenation compared to the best window size from the AIC (Ivan et al. 2025) because: (i)

we found one simulation with low accuracy when starting with short window size (Fig. 3),

which might be caused by initial over-splitting of the chromosome alignment that makes it

harder to improve the AIC via merging (i.e., if a non-recombining locus is split into multiple

windows, merging two neighbouring windows might not improve the AIC due to lack of

phylogenetic signal, as opposed to merging *all* windows together); (ii) it better captured the

long non-recombining loci from the simulated chromosomes as splitting these regions would

make the AIC worse (Fig. S3-S5); and (iii) it is computationally more efficient because we do

not have to test and compare the AIC of various fixed window sizes for the given alignment.

When we ran the splitting-and-merging algorithm on the six species from *erato-sara* clade of

Heliconius butterflies (with *H. melpomene* outgroup), we found that the window sizes varied

substantially across the chromosomes (Fig. S6-S7). For instance, the window sizes on

chromosome 1 ranged from 29bp to 70kb with an average of 581 ± 11bp (95% CI), while the

best fixed window size from the stepwise non-overlapping window method (Ivan et al. 2025)

was 250bp (Table S1). This best fixed window size was ~10x longer than the smallest

window – which prompted the concatenation of multiple windows during gene tree inference

– but 280x smaller than the biggest window – which might increase gene tree estimation error

during gene tree inference. As both concatenation and gene tree estimation error might bias

tree inferences (Ivan et al. 2025), this result further highlights the importance of using

variable window sizes when running non-overlapping window method on chromosome

alignments. Unfortunately, we cannot compare the AIC between the two methods, because

the stepwise non-overlapping window method (Ivan et al. 2025) excluded windows that

cannot be analysed while the splitting-and-merging approach keep them in the analysis.

Interestingly, the topology distribution of the group is generally consistent across studies

when considering the top ten most common topologies (Fig. 5), with either T1 or T2 as the

most common topology depending on the methods being used. This might infer that the gene

tree inference for this group is quite robust to the selection of window sizes. However, we

note that the *Heliconius* chromosomes from Edelman et al. (2019) were not phased. Huang et

al. (2022) demonstrated that random phasing can bias species tree inference, whereas allelic

phasing may improve this inference (Andermann et al. 2019; but see also Kates et al. 2018).

We hypothesise that running the splitting-and-merging approach on phased chromosomes

would generally result in shorter windows (particularly in highly heterozygous regions)

because our approach would also consider the allelic variation derived from phasing. This

observation highlights the importance of using variable window sizes on phased genome

alignments whenever possible to accurately identify recombination breakpoints across the

genomes.

Similar to the results from *Heliconius* butterflies, running the splitting-and-merging approach

on the genomes of great apes showed that the window sizes vary substantially across the

chromosomes (Fig. S9-S10). Interestingly, the mtDNA was partitioned into eight windows

(Table S6), where six of the windows recovered the major topology (grouping human and

chimpanzee together) while the other two windows recovered the two minor topologies (Fig.

S11). Gene tree discordance in mitochondrial genomes has been observed in different groups,

such as tetrapods (Toups et al. 2025), branchiopods (Sun and Cheng 2023), and new world

monkeys (Finnegan et al. 2025). In our case, the first window with the minor topology is

3.1kb-long and spans the 12S and 16S ribosomal RNA genes. A recent study by DeSalle and

Tessler (2025) observed that the two rRNA genes had low degree of phylogenetic congruence

with the other genes in the mtDNA genomes across 89 animal taxonomic groups, which

likely resulted in different tree topology. On the other hand, the second window is part of the

mtDNA control region. While previous studies have shown the utility of this region as a

phylogenetic marker (e.g., Randi et al. 2001; Bronstein et al. 2018), this window only

contains six parsimony informative sites in our four-taxon dataset, which is likely not enough

to infer a robust tree.

Across the great apes chromosomes, the estimated proportion of the major topology that

groups human and chimpanzee together (~79% of the sites) was generally higher than

previous estimates that ranged between 60-77% (Fig. 6; Ebersberger et al. 2007; Vanderpool

et al. 2020; Wong et al. 2024; Ivan et al. 2025). There are three potential explanations for the

differences in the estimates. First, both window-based and gene-based inferences are likely to

be affected by trade-offs between concatenation and gene tree estimation error (Ivan et al.

2025). While our method should more accurately reflect chromosomal recombination

breakpoints – and thus provides a more reliable estimate of topology distribution than gene-

based or fixed-window approaches – it is not entirely free from these competing factors. For

instance, we did not split a window if it cannot be analysed due to missing data or an

insufficient number of parsimony informative sites (as clearly shown for chromosome Y;

Table S6), which might concatenate multiple non-recombining loci together and favour the

topology that is supported by the majority of sites. In contrast, methods like MAST assign the

tree for individual sites, allowing weaker incongruent signals to be accounted in the final

estimates. Second, our analyses were based on whole genome alignments that include both

coding and non-coding regions. In great apes, coding regions were shown to undergo stronger

selection than non-coding regions (Cagan et al. 2016). Under the Hill-Robertson effect (Hill

and Robertson 1966), regions with stronger selection tend to have reduced effective

population size (N_e) (Comeron et al. 2008), which correlates with shortened coalescent time

and increased gene tree discordance. However, we cannot fully evaluate this explanation

because we did not explicitly differentiate the estimates between the coding and non-coding

regions. Third, Reddy et al. (2017) proposed that the current substitution models might be

inadequate to explain the evolution of coding regions, and potentially better-fit the non-

coding regions. As model violations have been shown to affect phylogenetic analysis (Naser-

Khdour et al. 2019), it is possible that analyses of coding regions – on which most previous

estimates were based – might underestimate the proportion of the major topology due to

higher level of discordance caused by model misspecification. In order to assess the relative

contributions of these explanations to our results, further studies should incorporate these

factors into the analyses (e.g., running MAST on whole genome alignments and estimating

the topological proportions separately for coding and non-coding regions).

Using variable window sizes, we further confirmed the findings from Edelman et al. (2019),

where the proportion of T1 was higher on shorter chromosomes, while T2 was more common

on longer chromosomes (Fig. S12 (left)). Martin et al. (2019) showed that chromosome

length is negatively correlated with recombination rate in *Heliconius* butterflies (i.e., shorter

chromosomes have higher recombination rates). On the other hand, recombination rate is also

correlated with the estimated admixture proportion between the three focal *Heliconius*

species in that study (Martin et al. 2019). This pattern arises because genomic regions with

low recombination rate tend to exhibit increased linkage among loci, which can act as a

barrier to gene flow (Barton and Bengtsson 1986; Aeschbacher et al. 2017). In our dataset, T2

is the regarded as the species tree topology, while T1 reflects hybridisation between *H.*

hecalesia and *H. telesiphe* (Edelman et al. 2019). As a result, T1 is enriched on shorter

chromosomes with higher recombination rates and weaker barrier to hybridisation, while T2

is enriched on longer chromosomes with lower recombination rates and stronger barrier to

gene flow. A similar pattern was observed along individual chromosomes (Fig. S12 (right)):

T1 was more frequent towards chromosomal ends (where recombination rates are higher),

while T2 had higher proportion towards the chromosomal centers (where recombination rates

are lower; Martin et al. 2019). We also identified two genomic regions that are dominated by

the minor topologies (Fig. 7). The first region spanned approximately 2Mb on chromosome 2

and was dominated by T3, likely caused by ILS and ancestral polymorphism (Edelman et al.

2019). The second region was 400Mb-long and spanned the *cortex* locus on chromosome 15,

which controls the wing colour patterns in *Heliconius* butterflies (Joron et al. 2006; Nadeau et

al. 2016; Jay et al. 2018). This region reflects an inversion in *H. demeter*, *H. sara*, *H.*

telesiphe, and *H. hecalesia*, but not in *H. erato*, *H. himera*, and the outgroup *H. melpomene*

(Edelman et al. 2019). As a result, *H. telesiphe* and *H. hecalesia* formed a sister pair, and

together become the sister clade of *H. demeter* and *H. sara* (T4; Edelman et al. 2019).

Overall, our AIC-based splitting-and-merging approach provides a method that uses variable

window sizes to infer gene trees from whole genome alignments. The windows (i.e., loci)

selected by our method should better reflect recombination breakpoints and thus serve as

better approximations of c-genes than simply choosing a fixed window size, or selecting loci

based on known genomic boundaries such as exons and introns. These loci can then be used

in a number of downstream applications. For example, if the aim is to reconstruct different

species trees across a genome alignment, loci from our method could be used as the starting

point for coalescent-based approaches such as those used by Thawornwattana et al. (2018,

2022). If the aim is to infer the evolutionary history of every site in the genome (as may be

required for applications such as PhyloGWAS (Pease et al. 2016)), then gene trees could be

estimated for each locus. If stochastic error is a concern, gene trees inferred from these loci

could be further refined using more complex substitution models that better-explain empirical

datasets (e.g., GHOST (Crotty et al. 2020) that models heterotachy, MixtureFinder (Ren et al.

2025) that models variation of transition rates among sites), and/or more sophisticated

methods that estimate the trees for individual sites (e.g., MAST (Wong et al. 2024)) or refine

the gene trees in relation to each other (e.g., Espalier (Rasmussen and Guo 2023), HMMs

(Siepel and Haussler 2005)). In all these cases, more accurate approximations of c-genes can

benefit downstream analyses, and have the potential to improve our understanding of

complex genomic histories.

SUPPLEMENTARY MATERIALS

Supplementary material will be available online upon the acceptance of the paper.

ACKNOWLEDGEMENTS

We thank Minh Bui, Justin Borevitz, the members of Lanfear and Bui Lab from Australian

National University, as well as Matt Hahn from Indiana University, for comments on the

study design and the manuscript.

FUNDING

This research is supported by Australian Government Research Training Program (AGRTP)

scholarship via The Australian National University Higher Degree by Research (HDR)

program.

REFERENCES

Aeschbacher S., Selby J.P., Willis J.H., Coop G. 2017. Population-genomic inference of the

strength and timing of selection against gene flow. Proceedings of the National

Academy of Sciences. 114:7061–7066.

Akaike H. 1974. A new look at the statistical model identification. IEEE Transactions on

Automatic Control. 19:716–723.

Andermann T., Fernandes A.M., Olsson U., Töpel M., Pfeil B., Oxelman B., Aleixo A.,

Faircloth B.C., Antonelli A. 2019. Allele Phasing Greatly Improves the Phylogenetic

Utility of Ultraconserved Elements. Systematic Biology. 68:32–46.

Barton N., Bengtsson B.O. 1986. The barrier to genetic exchange between hybridising

populations. Heredity. 57:357–376.

Brandt D.Y.C., Huber C.D., Chiang C.W.K., Ortega-Del Vecchyo D. 2024. The Promise of

Inferring the Past Using the Ancestral Recombination Graph. Genome Biology and

Evolution. 16:evae005.

Bronstein O., Kroh A., Haring E. 2018. Mind the gap! The mitochondrial control region and

its power as a phylogenetic marker in echinoids. BMC Evol Biol. 18:80.

Cagan A., Theunert C., Laayouni H., Santpere G., Pybus M., Casals F., Prüfer K., Navarro

A., Marques-Bonet T., Bertranpetit J., Andrés A.M. 2016. Natural Selection in the

Great Apes. Mol Biol Evol. 33:3268–3283.

Chan A.H., Jenkins P.A., Song Y.S. 2012. Genome-Wide Fine-Scale Recombination Rate

Variation in Drosophila melanogaster. PLOS Genetics. 8:e1003090.

Comeron J.M., Williford A., Kliman R.M. 2008. The Hill–Robertson effect: evolutionary

consequences of weak selection and linkage in finite populations. Heredity. 100:19–

31.

Crotty S.M., Minh B.Q., Bean N.G., Holland B.R., Tuke J., Jermiin L.S., Haeseler A.V.

2020. GHOST: Recovering Historical Signal from Heterotachously Evolved

Sequence Alignments. Syst Biol. 69:249–264.

Degnan J.H., Rosenberg N.A. 2009. Gene tree discordance, phylogenetic inference and the

multispecies coalescent. Trends Ecol Evol. 24:332–340.

DeSalle R., Tessler M. 2025. Mitochondrial Gene Phylogenetic Incongruencies Are Linked

to Chromosomal Position and Function. Genome Biol Evol. 17:evaf209.

Ebersberger I., Galgoczy P., Taudien S., Taenzer S., Platzer M., von Haeseler A. 2007.

Mapping Human Genetic Ancestry. Molecular Biology and Evolution. 24:2266–2276.

Edelman N.B., Frandsen P.B., Miyagi M., Clavijo B., Davey J., Dikow R.B., García-

Accinelli G., Van Belleghem S.M., Patterson N., Neafsey D.E., Challis R., Kumar S.,

Moreira G.R.P., Salazar C., Chouteau M., Counterman B.A., Papa R., Blaxter M.,

Reed R.D., Dasmahapatra K.K., Kronforst M., Joron M., Jiggins C.D., McMillan

W.O., Di Palma F., Blumberg A.J., Wakeley J., Jaffe D., Mallet J. 2019. Genomic

architecture and introgression shape a butterfly radiation. Science. 366:594–599.

Edwards S.V., Xi Z., Janke A., Faircloth B.C., McCormack J.E., Glenn T.C., Zhong B., Wu

S., Lemmon E.M., Lemmon A.R., Leaché A.D., Liu L., Davis C.C. 2016.

Implementing and testing the multispecies coalescent model: A valuable paradigm for

phylogenomics. Molecular Phylogenetics and Evolution. 94:447–462.

Feng C., Wang J., Liston A., Kang M. 2023. Recombination Variation Shapes Phylogeny and

Introgression in Wild Diploid Strawberries. Molecular Biology and Evolution.

40:msad049.

Finnegan N., Lima M.G.M., Lynch J.W. 2025. Mitochondrial DNA for Phylogeny Building:

Assessing Individual and Grouped mtGenes as Proxies for the mtGenome in

Platyrrhines. American Journal of Primatology. 87:e70017.

Flouri T., Jiao X., Rannala B., Yang Z. 2018. Species Tree Inference with BPP Using

Genomic Sequences and the Multispecies Coalescent. Molecular Biology and

Evolution. 35:2585–2593.

Fontaine M.C., Pease J.B., Steele A., Waterhouse R.M., Neafsey D.E., Sharakhov I.V., Jiang

X., Hall A.B., Catteruccia F., Kakani E., Mitchell S.N., Wu Y.-C., Smith H.A., Love

R.R., Lawniczak M.K., Slotman M.A., Emrich S.J., Hahn M.W., Besansky N.J. 2015.

Extensive introgression in a malaria vector species complex revealed by

phylogenomics. Science. 347:1258524.

Herrig D.K., Ridenbaugh R.D., Vertacnik K.L., Everson K.M., Sim S.B., Geib S.M.,

Weisrock D.W., Linnen C.R. 2024. Whole Genomes Reveal Evolutionary

Relationships and Mechanisms Underlying Gene-Tree Discordance in Neodiprion

Sawflies. Systematic Biology.:syae036.

Hibbins M.S., Breithaupt L.C., Hahn M.W. 2023. Phylogenomic comparative methods:

Accurate evolutionary inferences in the presence of gene tree discordance.

Proceedings of the National Academy of Sciences. 120:e2220389120.

Hill W.G., Robertson A. 1966. The effect of linkage on limits to artificial selection. Genetics

Research. 8:269–294.

Hoang D.T., Chernomor O., von Haeseler A., Minh B.Q., Vinh L.S. 2018. UFBoot2:

Improving the Ultrafast Bootstrap Approximation. Molecular Biology and Evolution.

35:518–522.

Huang J., Bennett J., Flouri T., Leaché A.D., Yang Z. 2022. Phase Resolution of

Heterozygous Sites in Diploid Genomes is Important to Phylogenomic Analysis under

the Multispecies Coalescent Model. Systematic Biology. 71:334–352.

Hudson R.R. 2002. Generating samples under a Wright–Fisher neutral model of genetic

variation. Bioinformatics. 18:337–338.

Ivan J., Frandsen P., Lanfear R. 2025. Selecting a Window Size for Phylogenomic Analyses

of Whole Genome Alignments using AIC. Syst Biol.:syaf053.

Jay P., Whibley A., Frézal L., Cara M.Á.R. de, Nowell R.W., Mallet J., Dasmahapatra K.K.,

Joron M. 2018. Supergene Evolution Triggered by the Introgression of a

Chromosomal Inversion. Current Biology. 28:1839-1845.e3.

Jensen-Seaman M.I., Furey T.S., Payseur B.A., Lu Y., Roskin K.M., Chen C.-F., Thomas

M.A., Haussler D., Jacob H.J. 2004. Comparative Recombination Rates in the Rat,

Mouse, and Human Genomes. Genome Res. 14:528–538.

Joron M., Papa R., Beltrán M., Chamberlain N., Mavárez J., Baxter S., Abanto M.,

Bermingham E., Humphray S.J., Rogers J., Beasley H., Barlow K., ffrench-Constant

R.H., Mallet J., McMillan W.O., Jiggins C.D. 2006. A Conserved Supergene Locus

Controls Colour Pattern Diversity in Heliconius Butterflies. PLOS Biology. 4:e303.

Jukes T.H., Cantor C.R. 1969. CHAPTER 24 - Evolution of Protein Molecules. In: Munro

H.N., editor. Mammalian Protein Metabolism. Academic Press. p. 21–132.

Kates H.R., Johnson M.G., Gardner E.M., Zerega N.J.C., Wickett N.J. 2018. Allele phasing

has minimal impact on phylogenetic reconstruction from targeted nuclear gene

sequences in a case study of Artocarpus. American Journal of Botany. 105:404–416.

Kent W.J., Sugnet C.W., Furey T.S., Roskin K.M., Pringle T.H., Zahler A.M., Haussler and

D. 2002. The Human Genome Browser at UCSC. Genome Res. 12:996–1006.

Kosakovsky Pond S.L., Posada D., Gravenor M.B., Woelk C.H., Frost S.D.W. 2006.

Automated Phylogenetic Detection of Recombination Using a Genetic Algorithm.

Molecular Biology and Evolution. 23:1891–1901.

Lescroart J., Bonilla-Sánchez A., Napolitano C., Buitrago-Torres D.L., Ramírez-Chaves

H.E., Pulido-Santacruz P., Murphy W.J., Svardal H., Eizirik E. 2023. Extensive

Phylogenomic Discordance and the Complex Evolutionary History of the Neotropical

Cat Genus Leopardus. Molecular Biology and Evolution. 40:msad255.

Locke D.P., Hillier L.W., Warren W.C., Worley K.C., Nazareth L.V., Muzny D.M., Yang S.-

P., Wang Z., Chinwalla A.T., Minx P., Mitreva M., Cook L., Delehaunty K.D.,

Fronick C., Schmidt H., Fulton L.A., Fulton R.S., Nelson J.O., Magrini V., Pohl C.,

Graves T.A., Markovic C., Cree A., Dinh H.H., Hume J., Kovar C.L., Fowler G.R.,

Lunter G., Meader S., Heger A., Ponting C.P., Marques-Bonet T., Alkan C., Chen L.,

Cheng Z., Kidd J.M., Eichler E.E., White S., Searle S., Vilella A.J., Chen Y., Flicek

P., Ma J., Raney B., Suh B., Burhans R., Herrero J., Haussler D., Faria R., Fernando

O., Darré F., Farré D., Gazave E., Oliva M., Navarro A., Roberto R., Capozzi O.,

Archidiacono N., Valle G.D., Purgato S., Rocchi M., Konkel M.K., Walker J.A.,

Ullmer B., Batzer M.A., Smit A.F.A., Hubley R., Casola C., Schrider D.R., Hahn

M.W., Quesada V., Puente X.S., Ordoñez G.R., López-Otín C., Vinar T., Brejova B.,

Ratan A., Harris R.S., Miller W., Kosiol C., Lawson H.A., Taliwal V., Martins A.L.,

Siepel A., RoyChoudhury A., Ma X., Degenhardt J., Bustamante C.D., Gutenkunst

R.N., Mailund T., Dutheil J.Y., Hobolth A., Schierup M.H., Ryder O.A., Yoshinaga

Y., de Jong P.J., Weinstock G.M., Rogers J., Mardis E.R., Gibbs R.A., Wilson R.K.

2011. Comparative and demographic analysis of orang-utan genomes. Nature.

469:529–533.

Ly-Trong N., Naser-Khdour S., Lanfear R., Minh B.Q. 2022. AliSim: A Fast and Versatile

Phylogenetic Sequence Simulator for the Genomic Era. Molecular Biology and

Evolution. 39:msac092.

Maddison W.P. 1997. Gene Trees in Species Trees. Systematic Biology. 46:523–536.

Mallet J., Besansky N., Hahn M.W. 2016. How reticulated are species? BioEssays. 38:140–

149.

Martin S.H., Davey J.W., Salazar C., Jiggins C.D. 2019. Recombination rate variation shapes

barriers to introgression across butterfly genomes. PLOS Biology. 17:e2006288.

McVean G.A.T., Myers S.R., Hunt S., Deloukas P., Bentley D.R., Donnelly P. 2004. The

Fine-Scale Structure of Recombination Rate Variation in the Human Genome.

Science. 304:581–584.

Meleshko O., Martin M.D., Korneliussen T.S., Schröck C., Lamkowski P., Schmutz J.,

Healey A., Piatkowski B.T., Shaw A.J., Weston D.J., Flatberg K.I., Szövényi P.,

Hassel K., Stenøien H.K. 2021. Extensive Genome-Wide Phylogenetic Discordance Is

Due to Incomplete Lineage Sorting and Not Ongoing Introgression in a Rapidly

Radiated Bryophyte Genus. Molecular Biology and Evolution. 38:2750–2766.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A.,

Lanfear R. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. Molecular Biology and Evolution. 37:1530–1534.

Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A., Brockington

S.F., Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2021. Disentangling Sources

of Gene Tree Discordance in Phylogenomic Data Sets: Testing Ancient

Hybridizations in Amaranthaceae s.l. Systematic Biology. 70:219–235.

Nadeau N.J., Pardo-Diaz C., Whibley A., Supple M.A., Saenko S.V., Wallbank R.W.R., Wu

G.C., Maroja L., Ferguson L., Hanly J.J., Hines H., Salazar C., Merrill R.M., Dowling

A.J., ffrench-Constant R.H., Llaurens V., Joron M., McMillan W.O., Jiggins C.D.

2016. The gene cortex controls mimicry and crypsis in butterflies and moths. Nature.

534:106–110.

Naser-Khdour S., Minh B.Q., Zhang W., Stone E.A., Lanfear R. 2019. The Prevalence and

Impact of Model Violations in Phylogenetic Analysis. Genome Biol Evol. 11:3341–

3352.

Pease J.B., Haak D.C., Hahn M.W., Moyle L.C. 2016. Phylogenomics Reveals Three Sources

of Adaptive Variation during a Rapid Radiation. PLOS Biology. 14:e1002379.

Randi E., Lucchini V., Hennache A., Kimball R.T., Braun E.L., Ligon J.D. 2001. Evolution

of the Mitochondrial DNA Control Region and Cytochrome *b* Genes and the

Inference of Phylogenetic Relationships in the Avian Genus *Lophura* (Galliformes).

Molecular Phylogenetics and Evolution. 19:187–201.

Rasmussen D.A., Guo F. 2023. Espalier: Efficient Tree Reconciliation and Ancestral

Recombination Graphs Reconstruction Using Maximum Agreement Forests.

Systematic Biology. 72:1154–1170.

Reddy S., Kimball R.T., Pandey A., Hosner P.A., Braun M.J., Hackett S.J., Han K.-L.,

Harshman J., Huddleston C.J., Kingston S., Marks B.D., Miglia K.J., Moore W.S.,

Sheldon F.H., Witt C.C., Yuri T., Braun E.L. 2017. Why Do Phylogenomic Data Sets

Yield Conflicting Trees? Data Type Influences the Avian Tree of Life more than

Taxon Sampling. Systematic Biology. 66:857–879.

Ren H., Wong T.K.F., Minh B.Q., Lanfear R. 2025. MixtureFinder: Estimating DNA Mixture

Models for Phylogenetic Analyses. Mol Biol Evol. 42:msae264.

Richards E.J., Brown J.M., Barley A.J., Chong R.A., Thomson R.C. 2018. Variation Across

Mitochondrial Gene Trees Provides Evidence for Systematic Error: How Much Gene

Tree Variation Is Biological? Systematic Biology. 67:847–860.

Scally A., Dutheil J.Y., Hillier L.W., Jordan G.E., Goodhead I., Herrero J., Hobolth A.,

Lappalainen T., Mailund T., Marques-Bonet T., McCarthy S., Montgomery S.H.,

Schwalie P.C., Tang Y.A., Ward M.C., Xue Y., Yngvadottir B., Alkan C., Andersen

L.N., Ayub Q., Ball E.V., Beal K., Bradley B.J., Chen Y., Clee C.M., Fitzgerald S.,

Graves T.A., Gu Y., Heath P., Heger A., Karakoc E., Kolb-Kokocinski A., Laird

G.K., Lunter G., Meader S., Mort M., Mullikin J.C., Munch K., O'Connor T.D.,

Phillips A.D., Prado-Martinez J., Rogers A.S., Sajjadian S., Schmidt D., Shaw K.,

Simpson J.T., Stenson P.D., Turner D.J., Vigilant L., Vilella A.J., Whitener W., Zhu

B., Cooper D.N., de Jong P., Dermitzakis E.T., Eichler E.E., Flicek P., Goldman N.,

Mundy N.I., Ning Z., Odom D.T., Ponting C.P., Quail M.A., Ryder O.A., Searle

S.M., Warren W.C., Wilson R.K., Schierup M.H., Rogers J., Tyler-Smith C., Durbin

R. 2012. Insights into hominid evolution from the gorilla genome sequence. Nature.

483:169–175.

Schneider V.A., Graves-Lindsay T., Howe K., Bouk N., Chen H.-C., Kitts P.A., Murphy

T.D., Pruitt K.D., Thibaud-Nissen F., Albracht D., Fulton R.S., Kremitzki M.,

Magrini V., Markovic C., McGrath S., Steinberg K.M., Auger K., Chow W., Collins

J., Harden G., Hubbard T., Pelan S., Simpson J.T., Threadgold G., Torrance J., Wood

J.M., Clarke L., Koren S., Boitano M., Peluso P., Li H., Chin C.-S., Phillippy A.M.,

Durbin R., Wilson R.K., Flicek P., Eichler E.E., Church D.M. 2017. Evaluation of

GRCh38 and de novo haploid genome assemblies demonstrates the enduring quality

of the reference assembly. Genome Res. 27:849–864.

Shen W., Le S., Li Y., Hu F. 2016. SeqKit: A Cross-Platform and Ultrafast Toolkit for

FASTA/Q File Manipulation. PLoS ONE. 11:e0163962.

Siepel A., Haussler D. 2005. Phylogenetic Hidden Markov Models. In: Nielsen R., editor.

Statistical Methods in Molecular Evolution. New York, NY: Springer. p. 325–351.

Springer M.S., Gatesy J. 2016. The gene tree delusion. Molecular Phylogenetics and

Evolution. 94:1–33.

Steenwyk J.L., Li Y., Zhou X., Shen X.-X., Rokas A. 2023. Incongruence in the

phylogenomics era. Nat Rev Genet. 24:834–850.

Sun X., Cheng J. 2023. Conflicts in Mitochondrial Phylogenomics of Branchiopoda, with the

First Complete Mitogenome of Laevicaudata (Crustacea: Branchiopoda). Current

Issues in Molecular Biology. 45:820–837.

Thawornwattana Y., Dalquen D., Yang Z. 2018. Coalescent Analysis of Phylogenomic Data

Confidently Resolves the Species Relationships in the Anopheles gambiae Species

Complex. Molecular Biology and Evolution. 35:2512–2527.

Thawornwattana Y., Seixas F.A., Yang Z., Mallet J. 2022. Full-Likelihood Genomic Analysis

Clarifies a Complex History of Species Divergence and Introgression: The Example

of the erato-sara Group of Heliconius Butterflies. Systematic Biology. 71:1159–1177.

Toups B.S., Thomson R.C., Brown J.M. 2025. Complex Models of Sequence Evolution

Improve Fit, But Not Gene Tree Discordance, for Tetrapod Mitogenomes. Syst Biol.

74:86–100.

Vanderpool D., Minh B.Q., Lanfear R., Hughes D., Murali S., Harris R.A., Raveendran M.,

Muzny D.M., Hibbins M.S., Williamson R.J., Gibbs R.A., Worley K.C., Rogers J.,

823   Hahn M.W. 2020. Primate phylogenomics uncovers multiple rapid radiations and

824    ancient interspecific introgression. PLOS Biology. 18:e3000954.

825 Waterson R.H., Lander E.S., Wilson R.K., The Chimpanzee Sequencing and Analysis

826    Consortium. 2005. Initial sequence of the chimpanzee genome and comparison with

827    the human genome. Nature. 437:69–87.

828 Wong T.K.F., Cherryh C., Rodrigo A.G., Hahn M.W., Minh B.Q., Lanfear R. 2024. MAST:

829    Phylogenetic Inference with Mixtures Across Sites and Trees. Systematic

830    Biology.:syae008.

831 Yang W., Feiner N., Pinho C., While G.M., Kaliontzopoulou A., Harris D.J., Salvi D., Uller

832    T. 2021. Extensive introgression and mosaic genomes of Mediterranean endemic

833    lizards. Nat Commun. 12:2762.

835