Addressing antibody validation failures: a multi-stakeholder Delphi consensus study on actionable solutions

Short title: Delphi consensus on antibody validation reform

Authors: Katherine Blades[1], Michael Biddle[1], Robert Froud[2,3], Eva M. Krockow[1]*, Harvinder Virk[1]*

*joint corresponding authors

Harvinder Virk – hsv6@leicester.ac.uk, Eva Krockow – emk12@leicester.ac.uk

Affiliations:

1. University of Leicester, College of Life Sciences, Leicester, England, UK

2. School of Health Sciences, Kristiania University of Applied Sciences, Oslo, Norway

3. Warwick Medical School, University of Warwick, United Kingdom

Abstract

The experimental use of antibodies that have not been validated for context-specific use frequently misdirects biomedical research. Experimental results that derive from the use of inadequately validated antibodies are estimated to waste over $1 billion annually in the United States alone and to consume millions of animal and human biological samples in experiments whose conclusions may be unreliable. Community validation frameworks, reporting standards, and independent characterisation initiatives have made important progress, and multi-stakeholder coordination efforts are emerging. However, the research community lacks a formally developed, consensus-based action plan that specifies what each stakeholder group should do, by when, and with what priority. We conducted a modified Delphi study with international experts representing academic researchers, scientific publishers, research funders, antibody manufacturers, and institutional research leaders to develop actionable recommendations for improving antibody validation,

selection, and reporting practices. Thirty-two participants rated 33 proposed actions on effectiveness and feasibility using 9-point scales, with consensus assessed using the RAND/UCLA Appropriateness Method. Over two rounds, the panel achieved consensus on 15 items as both effective and feasible for implementation by 2030. These spanned institutional actions (training in antibody validation, integration into research integrity frameworks, support for local expertise networks), funder actions (dedicated validation budgets, grant application requirements, endorsement of community reporting standards), publisher actions (complete antibody reporting packages, clear validation standards), manufacturer actions (assignment of unique identifiers at source), and cross-stakeholder coordination (a shared roadmap for improvement). An additional 15 items were rated as effective but with uncertain feasibility, reflecting a consistent pattern in which the panel agreed on the value of proposed interventions but expressed reservations about realistic implementation timelines. One item was rejected by the panel with concerns around effectiveness and feasibility. Participants described four interconnected barriers to progress: diffuse ownership of the problem across stakeholders; market dynamics that inadequately reward antibody quality; difficulty justifying investment when returns are distributed across the research system; and coordination challenges among actors with different incentive structures. These barriers are addressable through coordinated action, and the findings complement existing technical and data-sharing initiatives by providing the structured, stakeholder-endorsed policy framework needed to translate awareness of the problem into concrete practice and policy changes.

Introduction

Antibodies are among the most widely used tools in biomedical research, enabling researchers to detect, quantify, and isolate specific proteins within complex biological samples. However, research antibodies do not always bind exclusively to their intended targets. Poor antibody specificity has repeatedly misdirected biomedical research across diverse fields, leading to incorrect conclusions about protein function, unreliable biomarker discovery, and misidentification of drug targets [1-10]. These failures represent not isolated occurrences but a persistent and systemic issue affecting the reliability of the published literature.

The scale of this problem is substantial. An estimated $28.2 billion per year is spent in the United States on preclinical research that is not reproducible, with issues relating to biological reagents likely being the biggest contributor [11]. Antibody-related failures alone account for an estimated $1 billion annually in the US in direct economic waste [12, 13], a figure that excludes the substantial opportunity cost of misdirected research effort. Beyond the economic dimension, there is a large and largely unquantified ethical cost: the waste of animals both in antibody production and in research studies that employ unsuitable antibodies, and the waste of donated human biological materials in experiments whose conclusions may be unreliable [14]. Recent estimates suggest that at least 4 million animal samples and 6 million human tissue samples have been consumed globally in

experiments using antibodies that failed independent testing, without context-specific validation

since December, 2000 [14].

How well an antibody performs in binding to its target varies between applications, protocols, and

the cell or tissue types used for each experiment. Consequently, it is important that performance is

assessed for each context of use, a process referred to as validation, and it is the responsibility of the

researcher to carry out these studies for their specific protocol [15]. The complexity and scale of

antibody use have contributed to a marketplace containing a high proportion of poorly performing

antibodies [1]. Recent independent benchmarking by the YCharOS (Antibody Characterisation by

Open Science) consortium quantified the likely scale of the problem: greater than 50% of 614

commercial antibodies against 65 neuroscience-related proteins failed characterisation experiments

in three commonly used applications, and 88.4% of papers using poorly performing antibodies in

immunofluorescence presented no relevant validation data [1]. Moreover, each protein target was

linked to an average of approximately 12 published papers that presented data using poorly

performing antibodies, perpetuating the use of these reagents and the associated issues of

reproducibility.

Efforts to address this problem have operated on multiple fronts. Technical advances include the

IWGAV five-pillar validation framework, which outlines complementary strategies (genetic,

orthogonal, independent antibody, tagged protein, and immunocapture mass spectrometry), for

establishing antibody selectivity [15], and the large-scale independent characterisation work of the

YCharOS consortium, which has provided open, head-to-head comparisons of commercial

antibodies using knockout cell lines as isogenic controls [1]. Reporting standards such as the

Materials Design Analysis Reporting (MDAR) framework have sought to improve transparency at

the point of publication [16], and tools including the Research Resource Identifier (RRID) initiative

have enabled more precise tracking of antibody use across the literature [17]. Individual

manufacturers have also invested in more rigorous characterisation, in some cases removing or

relabelling products found to perform poorly [1, 14].

In parallel, multi-stakeholder coordination has begun to emerge. Kahn et al. [12] assembled a

history of efforts to address the antibody characterisation crisis and proposed sector-specific actions

for researchers, universities, journals, vendors, scientific societies, disease foundations, and funding

agencies. The Only Good Antibodies (OGA) community, , of which members of the present study

team are founding contributors [18], has brought together stakeholders across the research

ecosystem to work toward making best practice in antibody use more feasible, easy, and rewarding,

drawing on the Centre for Open Science's framework for culture change in research. These

initiatives represent important progress. However, the field still lacks a formally developed,

consensus-based action plan in which proposed interventions have been systematically rated for

both effectiveness and feasibility by a diverse expert panel. Without such structured prioritisation, it

is difficult for stakeholders to know which actions to implement first, where the strongest

agreement exists, and where feasibility barriers require targeted attention.

The underlying reasons for the persistence of this problem are complex and multifaceted,

encompassing the inherent challenges of producing and characterising high-quality antibodies, but

also researcher awareness, education, and incentive structures [14]. Researchers identify time, cost,

and lack of supervisor support as primary barriers to better validation practices, while strongly

supporting structural solutions including open data sharing, dedicated validation funding, and

publisher requirements [14]. Behavioural analysis drawing on the COM-B (Capabilities +

Opportunities + Motivation Behaviour) model suggests that individual capability and motivation

interact with the opportunity provided by the research environment, such that improvements to

practice require changes not only in individual behaviour but in the systems and structures that

shape it [14, 18]. This suggests that the problem requires coordinated action across the research

ecosystem.

Having established the scale and drivers of the problem [14], and situated it within the broader

landscape of ongoing technical and community efforts [1, 12, 18], this study sought to move from

identifying what needs to change, to building structured agreement on how to change it. We

convened a multi-stakeholder Delphi consensus process to systematically develop and prioritise

recommendations for improving antibody validation, selection, and reporting practices. Drawing on

structured assessment by 32 international experts representing researchers, publishers, funders,

manufacturers, and institutional leaders, this study presents a consensus process for each expert

stakeholder group alongside an analysis of implementation barriers and opportunities for

coordinated action.

Methods

Study Design and Ethical Approval

We conducted a modified Delphi study to develop stakeholder-informed recommendations for

improving antibody validation, selection, and reporting practices in biomedical research. The

Delphi method is a structured consensus-building approach that systematically collects and

synthesises expert opinion through iterative questionnaires with controlled feedback [19]. This

study was approved by the University of Leicester College of Life Sciences Research Ethics

Committee (reference number [2025-1122-5743]) and conducted in accordance with the General

Data Protection Regulation (GDPR, 2018). All participants provided informed consent before

participation. The study protocol anticipated up to four rounds of consultation between August 2025

and February 2026, with each round allowing 2-3 weeks for completion. Data collection and

management were conducted using the Clinvivo platform (https://www.clinvivo.com; Clinvivo Ltd,

London, UK), a secure GDPR-compliant system designed specifically for Delphi studies.

Participant Recruitment

We identified expert stakeholders from five key sectors involved in antibody use and research

governance: academic biomedical research, scientific publishing, research funding, antibody

manufacturing, and institutional research leadership. Experts were defined as individuals who are

representative of their profession, have the power to implement findings, or who would not be

challenged as experts in their field [21]. Thirty-eight experts were identified through networks

based on their expertise in antibody validation, editorial roles at major biomedical journals,

leadership positions at research funding organisations, executive roles in commercial antibody

production, or responsibility for research integrity policies at academic institutions. Tailored

recruitment emails were sent to each stakeholder group. Participants were informed that the study

aimed to develop a consensus-based action plan to improve antibody validation and reporting across

the research ecosystem, and that their anonymised responses would be aggregated to inform policy

recommendations for journals, funders, institutions, and manufacturers.

The Delphi Process

The Delphi process consisted of iterative rounds of anonymous questionnaire completion with

structured feedback. Initial draft statements for Round 1 were developed from two sources:

recommendations emerging from a stakeholder meeting convened by the National Centre for the

Replacement, Refinement and Reduction of Animals in Research (NC3Rs) in February 2024 [20],

and themes identified through focus group discussions (and validated with a survey) with antibody-

using researchers conducted in the summer of 2023 [14]. These sources yielded 23 proposed actions

spanning four domains: journal and publisher policies (n=9 items covering reporting requirements,

validation transparency, and editorial capacity building), research funder actions (n=9 items

covering grant requirements, validation funding, and manufacturer engagement), institutional and

educational interventions (n= 3 items covering training, ethics frameworks, and local champions),

and cross-stakeholder coordination mechanisms (n= 2 items covering roadmaps, shared

infrastructure, and benchmarking initiatives). Each item described a specific action that one or more

stakeholder groups could adopt to improve antibody practices (S6 Text).

Round 1 was conducted from August 12 to September 12, 2025, with two deadline extensions to maximise response rates. Participants received weekly email reminders from Clinvivo, and the research team sent a final reminder on the originally scheduled closing date. Round 2 was conducted from November 6 to November 23, 2025, following the same reminder protocol. The decision to proceed to additional rounds was based on pre-specified criteria: lack of consensus, persistence of significant disagreement requiring resolution, or evidence that additional rounds would meaningfully shift ratings (e.g., based on qualitative feedback).

Rating System and Assessment of Consensus

Participants rated each proposed action on two 9-point rating scales. The effectiveness scale permitted panellists to rate how meaningfully they thought the item would improve the selection, validation, or reporting of antibodies in research (1=extremely ineffective, 9=extremely effective). The feasibility scale permitted panellists to rate how realistically they thought items could be adopted across relevant settings by 2030 (1=extremely infeasible, 9=extremely feasible). Participants could select "don't know" if an item fell outside their area of expertise. For each item, participants could provide free-text comments to share their rationale, identify potential risks or unintended consequences, or suggest alternative wordings.

To assess disagreement, we employed the RAND/UCLA Appropriateness Method [19]. For each item, we calculated the median rating, the inter-percentile range (IPR; difference between the 30th and 70th percentiles), and the inter-percentile range adjusted for symmetry (IPRAS). The IPRAS accounts for asymmetric rating distributions using the equation: $IPRAS = IPRr + (AI \times CFA)$, where IPRr is the inter-percentile range required for disagreement when perfect symmetry exists (2.35), AI is the asymmetry index (calculated as the distance between the central point of the IPR and the scale midpoint of 5), and CFA is the correction factor for asymmetry (1.5). These values of 2.35 and 1.5 were empirically derived by Fitch et al. [19] to best reproduce classic RAND definitions of disagreement for 9-point scales. Percentiles were calculated using Stata version 18

(StataCorp LLC, College Station, TX; RRID:SCR_012763) with the R-2 formula (Stata's default

method).

An item was classified as having disagreement when IPR > IPRAS for that item. Items were

classified according to median ratings: effective/feasible (median 7-9), ineffective/infeasible

(median 1-3), or uncertain (median 4-6). For medians falling on category boundaries (3.5 or 6.5)

due to even participant numbers, we applied a "round down" rule, classifying 3.5 as

ineffective/infeasible and 6.5 as uncertain. This meant a conservative approach was taken to

describing ratings as effective or feasible. Qualitative free-text comments were reviewed by the

research team to inform item refinement, identify new proposed actions, and develop narrative

summaries for feedback to participants in subsequent rounds.

Progression Decisions Between Rounds

Items were progressed, removed, or modified between rounds based on their ratings, the presence or

absence of disagreement, or qualitative feedback. Items rated as both effective (median $\geq$7) and

feasible (median $\geq$7) without disagreement were accepted as recommendations and did not require

re-rating in subsequent rounds unless there were suggestions for rewording in the qualitative

comments. Items rated as both ineffective (median $\leq$3) and infeasible (median $\leq$3) without

disagreement were rejected and removed from further consideration. Items classified as uncertain

on either dimension, or those with disagreement, were considered for re-rating in the next round,

particularly if qualitative comments suggested that rewording, clarification, or presentation of

divergent perspectives might shift opinions.

Between Round 1 and Round 2, panel members' free-text comments were reviewed to identify: (1)

suggestions for new action items to add to the consensus process, (2) proposed modifications to

existing item wording, and (3) substantive concerns or alternative perspectives that should be shared

with the panel. Items were reworded when comments indicated ambiguity or suggested

semantically meaningful improvements. New items suggested by multiple participants or

representing distinct actionable proposals were added to Round 2. For items with disagreement or uncertainty, we prepared feedback summaries that synthesised the range of panel perspectives to support informed re-rating. Items with substantive wording changes based on panel feedback were re-rated regardless of initial ratings, to ensure consensus reflected the revised formulation.

Results

Participant Characteristics and Response Rates

Thirty-eight individuals were invited to participate in Round 1. Thirty-two completed the round (response rate: 84.2%). All Round 1 participants were invited to Round 2, and 30 completed this round (response rate: 93.8%; overall retention: 78.9% of original invitees).

The 32 Round 1 participants included 13 females (40.6%), 17 males (53.1%), and 2 who preferred not to disclose sex (6.3%). Among the 30 participants who reported their age, the mean was 51.5 years (SD=11.3). Participants were primarily based in the United Kingdom (n=16, 50.0%) and United States (n=13, 40.6%), with additional representation from Canada (n=2, 6.3%) and Germany (n=1, 3.1%). The panel was highly educated, with 27 participants (84.4%) holding PhD-level qualifications, 2 (6.3%) holding master's degrees, and 3 (9.4%) holding bachelor's degrees. Participants represented diverse fields within the biomedical research ecosystem: academic biomedical research (n=7, 21.9%), scientific publishing (n=6, 18.8%), antibody manufacturing (n=6, 18.8%), research funding (n=4, 12.5%), industrial biomedical research (n=2, 6.3%), and other fields (n=7, 21.9%) including antibody informatics, infrastructure provision, and the charity sector. Within these sectors, participants held senior decision-making roles: 37.5% (n=12) reported full autonomy to make decisions on behalf of their organisations, 34.4% (n=11) made decisions with input from senior leaders, and 28.1% (n=9) made recommendations with final decisions made by senior leaders. Detailed breakdowns of career stages within each employment sector are provided in S1 Table.

Overview of Items and Consensus Development

A total of 33 proposed action items were rated across two rounds (Figure 1). Round 1 presented 23 items spanning four domains: journal and publisher policies (9 items, R1–R9), research funder actions (9 items, R10–R18), institutional and educational interventions (3 items, R19–R21), and cross-stakeholder coordination (2 items, R22–R23), see S6 Text for full item wording. All 32 participants rated effectiveness and feasibility for each of the 23 proposed actions. Panel ratings demonstrated strong consensus on most items, with disagreement identified on only 2 of 46 ratings (effectiveness and feasibility combined).

Following Round 1 analysis, 11 items (R1, R2, R10–R15, R19, R20, R22) achieved consensus as both effective (median ≥7) and feasible (median ≥7) without disagreement and were immediately accepted as recommendations without requiring re-rating. The remaining 12 items from Round 1 were re-presented in Round 2, including six with revised wording based on panel comments (R3, R8, R9, R17, R18, R21). Based on participant free-text suggestions, 10 additional action items (A1–A10) were added to Round 2. Round 2 thus presented 22 items for rating: 12 re-rated items from Round 1 and 10 newly proposed items.

In Round 2 disagreement only occurred on 1 of 44 ratings. Among the 22 items presented in Round 2, four achieved agreement as both effective and feasible (R3, R21, A1, A7), joining the 11 items accepted in Round 1 for a total of 15 consensus recommendations at the close of Round 2. One item (R5) shifted from uncertain in Round 1 to consensually infeasible in Round 2 (median feasibility declined from 4.0 to 3.0) and was the only item to be rated either infeasible or ineffective. Two items (A4, A5) were rated as uncertain on both effectiveness and feasibility.

Final Consensus Outcomes

Across the two rounds, the panel evaluated 33 proposed actions, achieving consensus on 15 items as both effective (median ≥7) and feasible (median ≥7), without disagreement (Table 1). These 15 recommendations span all four stakeholder domains and represent actions that expert panellists

agreed would meaningfully improve antibody practices and could realistically be implemented by 2030.

The panel rated 17 items as uncertain on at least one dimension (Table 2). In 15 of these 17 cases, items were rated as effective (median ≥7) but with uncertain feasibility (median 4–6), indicating that while panellists believed these actions would improve antibody practices, they had reservations about realistic implementation by 2030. This pattern, high confidence in effectiveness, uncertainty about feasibility, was the dominant finding for items not achieving consensus, and is a recurring theme in the stakeholder-specific recommendations presented below. One item was rejected by the panel, with concerns around both effectiveness and feasibility (Table 3). A summary of the ratings across all items is presented in Figure 2.

Following analysis of Round 2 results, the research team decided not to proceed to a third round. This decision was based on three convergent factors. First, a high proportion of items (15 of 33, 45.5%) had achieved clear consensus as both effective and feasible across the two rounds. Second, disagreement remained minimal, with disagreement being present on only one item in Round 2. Third, rating stability between rounds was high: among the 12 items re-rated in Round 2, median effectiveness ratings remained identical for 9 items (75.0%) and median feasibility ratings remained identical for 10 items (83.3%), with no item shifting by more than 1 point on either scale. The minimal rating changes between rounds indicated that further iteration would be unlikely to resolve the persistent pattern of items rated as effective but with uncertain feasibility. Full quantitative results for all items across both rounds are presented in S2 and S3 Tables, with complete qualitative commentary provided in S7 and S8 Texts.

Figure 1. Overview of the Delphi process including formulation of recommendations and their

assessment across rounds.

290   **TABLE 1: Consensus Recommendations - Items Rated Both Effective and Feasible with agreement**

Item	Full Recommendation	Median Effectiveness	Median Feasibility	Stakeholder Group targeted by recommendation
Journal and Publisher Policies				
R1	Authors should include the Research Resource Identifier (RRID) for each antibody used, where available.	7.0	8.0	Journals, Authors
R2	Authors should provide sufficient metadata to allow unambiguous identification of each antibody. This includes: Clone ID (for monoclonal antibodies), Catalogue number and lot number (for polyclonal antibodies), Vendor or source.	8.0	8.0	Journals, Authors
R3	Authors should report the dilution ratio at which each antibody is used. In addition, authors should state the antibody protein concentration where possible. Examples include dilution ratios such as 1:1000 or 1:500, or concentrations expressed in µg/mL.	8.0	8.0	Journals, Authors

A1	Journals should have a clearly stated standard for antibody validation and reporting.	7.5	7.0	Journals
Research Funder Actions				
R10	Funders should include a section in the application form where applicants must detail the steps they will take to validate the antibodies used.	7.5	8.0	Funders
R11	Applicants to biomedical funding schemes should include a specific line item within their budget for resources to conduct robust antibody validation (if relevant to their proposal).	8.0	7.0	Funders
R12	Funders should create or expand targeted schemes to support the development and validation of critical antibody-based research tools. These could include, for example: Characterisation of under-studied antibodies, Generation of KO/KD cell lines, Cross-platform benchmarking of antibody performance, Replacement of animal-derived antibodies.	8.0	7.0	Funders

R13	Funders should share information within the applicant guidance that antibody performance is an important limitation in many experimental methods. Applicants should be encouraged to address this in methodological or reproducibility sections, as this will be part of how these sections are evaluated by reviewers and panels.	7.0	7.0	Funders
R14	Funders should encourage grantees to deposit antibody validation data in open-access repositories, ideally linked to RRIDs or registry entries. This expectation could be embedded in end-of-grant reporting or incentivised through visibility or reproducibility initiatives.	7.0	7.0	Funders
R15	Funders should formally endorse community-developed reporting standards that promote antibody transparency and validation (e.g., IWGAV, MDAR). Such endorsements can help reinforce aligned expectations across funders and journals.	7.0	7.0	Funders
Institutional and Educational Actions				
R19	Training in antibody validation should be offered in relevant bioscience courses and programmes. This could include: Undergraduate education (e.g., in	8.0	7.0	Institutions, Learned Societies

	immunology, neuroscience, or molecular biology programmes), Postgraduate or doctoral training (e.g., through doctoral training centres), Postdoctoral or continuing professional development. Training should be field-appropriate, not universally required across all domains.			
R20	Universities and research institutions should incorporate antibody validation expectations into research integrity and ethics review frameworks. This may include: Asking researchers to consider antibody validation as part of research ethics approval processes, Providing guidance or templates for documenting validation plans.	7.0	7.0	Institutions
R21	Universities, research institutions and learned societies should recognise and support the Antibody Champions Scheme. This UK-based initiative, co-funded by the Medical Research Council (MRC) and the National Centre for the replacement, refinement and reduction of animals in research (NC3Rs), will be launched in 2026. It will be a voluntary scheme of local champions or experts in antibody validation. These individuals will serve as: Points of contact for advice on antibody selection, validation design, and data interpretation; Trainers or facilitators in	7.0	7.0	Institutions, Learned Societies

	antibody validation best practice workshops or research methods courses; Advocates for antibody good practice within departments or core facilities. Support could include protected time, acknowledgement in performance reviews, or funding for training and outreach.			
Cross-Stakeholder Coordination				
R22	Stakeholders should work together to develop a shared roadmap for improving antibody validation practices by 2030. This would outline: Stakeholder-specific actions and responsibilities, Target timelines and milestones, Mechanisms for accountability and feedback.	7.0	7.0	All Stakeholders
Manufacturer Actions				
A7	Antibody manufacturers should be encouraged to assign RRIDs to their products at source.	8.0	7.0	Manufacturers

 TABLE 2: Items with Agreed Uncertain Effectiveness and/or Feasibility

Item	Full Recommendation	Median Eff	Median Feas	Pattern
Effective but Feasibility Uncertain				
R4	Journals should require authors to deposit antibody validation data in an open-access repository, linked to a persistent identifier (e.g., RRID, DOI). Repositories might include: Zenodo, institutional repositories, manufacturer portals, or The Antibody Registry (linked to Biomed Resource Watch).	8.0	5.0	Effective, Uncertain Feasibility
R6	Publishers should provide training resources for reviewers and editors to help them evaluate antibody validation information in submitted manuscripts.	7.0	5.0	Effective, Uncertain Feasibility
R7	Publishers and other stakeholders should support the development and use of automated tools to flag potential antibody validation issues in submitted manuscripts – e.g., for antibodies that have already been withdrawn from the market because of performance issues.	8.0	5.0	Effective, Uncertain Feasibility

R8	As part of the submission process, authors should describe how antibody specificity was assessed, making reference to the International Working Group on Antibody Validation (IWGAV) framework. A brief paragraph, checklist, or table should indicate which of the IWGAV pillars were used, if any (e.g., genetic, orthogonal, independent antibody, tagged protein, IP-MS).	7.0	6.0	Effective, Uncertain Feasibility
R9	Where antibodies are critical to key experimental conclusions, where technically feasible, authors should either: present validation data based on a genetic strategy (e.g., knockout or knockdown), or cite previously published genetic validation data.	8.0	5.0	Effective, Uncertain Feasibility
R16	Funders should directly engage with manufacturers to encourage them to: Publish antibody validation datasets, Adopt consistent metadata standards (e.g., RRID, clone ID, lot number).	7.0	5.0	Effective, Uncertain Feasibility
R17	Funders should collaborate with independent benchmarking initiatives such as YCharOS or equivalent open-science consortia. They should directly co-fund the validation of antibodies against key targets to create suites of validated tools that can be used by researchers to improve data quality and thereby increase return on investment for the funder.	8.0	6.0	Effective, Uncertain Feasibility

R18	Funders should promote industry participation in independent benchmarking initiatives by asking applicants to consider working with initiatives such as YCharOS or equivalent open-science consortia as part of their work. This could be achieved either by including the recommendation in the guidance notes or flagging to applicants at the panel review stage.	7.0	6.0	Effective, Uncertain Feasibility
R23	A coordinated, shared infrastructure should be developed (or expanded) to: Aggregate antibody validation data, Provide standardised formats and tools for sharing, Enable journals, funders, and institutions to access validation status or red flags. This may involve expanding existing platforms such as YCharOS, or building interoperability across initiatives like RRID, Benchsci, CiteAb, and Antibody Registry.	8.0	6.0	Effective, Uncertain Feasibility
A2	Detailed Protocols and Validation data should be included in the manuscript or supplemental data.	8.0	6.0	Effective, Uncertain Feasibility
A3	Journals should appoint a specialist editor that focuses on reproducibility.	7.0	5.5	Effective, Uncertain Feasibility

A6	Funders should mandate that if funds are used to develop an antibody as part of a research project, the reagent must be a recombinant antibody and must be made available to the wider community.	7.0	6.0	Effective, Uncertain Feasibility
A8	Manufacturers should perform a standard set of validation experiments on their reagent antibodies and make the data available.	8.0	6.5	Effective, Uncertain Feasibility
A9	Manufacturers should shift production to all recombinant antibodies.	7.0	5.0	Effective, Uncertain Feasibility
A10	A learned society for antibody validation should be established.	7.0	5.0	Effective, Uncertain Feasibility
Uncertain on Both Dimensions				
A4	Funders should maintain and provide researchers with a list of manufacturers who validate their antibodies.	6.0	5.0	Uncertain Both

| A5 | Funders should mandate the use of recombinant antibodies (or at least monoclonals) when available. Justification should be required if a polyclonal is used. | 6.0 | 5.0* | Uncertain Both |

294 *Panel disagreement on feasibility (IPR=4 > IPRAS=2.35)

296 **TABLE 3: Rejected Recommendation - Item Rated Uncertain effectiveness, and Infeasible, with agreement**

Item	Full Recommendation	Median Effectiveness	Median Feasibility
R5	Journals should introduce a tiered scoring system or transparency badge to recognise papers that meet higher standards of antibody reporting and validation.	4.0	3.0

Figure 2. Effectiveness and feasibility ratings for all 33 proposed actions. Each point represents one proposed action, plotted by median effectiveness (x-axis) and median feasibility (y-axis) ratings on 9-point scales. Points are colour-coded by stakeholder group. Dashed lines indicate the boundary between ratings classified as effective/feasible (median ≥7) and uncertain (median 4–6), following our RAND/UCLA round-down rule for boundary medians of 6.5. The upper-right quadrant contains items rated as both effective and feasible with agreement (15 items). The lower-right quadrant contains items rated as effective but with uncertain feasibility (15 items). Item with panel disagreement (A5, IPR > IPRAS) is shown as open circle with dashed borders; all other items achieved agreement. One item (R5) was rejected by the panel (uncertain effectiveness, infeasible). Points at identical coordinates have been slightly offset for readability.

Stakeholder-Specific Recommendations

The following sections present the consensus recommendations, predicted to be both effective and feasible, and items predicted to be effective but with uncertain feasibility for implementation by 2030, organised by stakeholder group. For each recommendation, we provide the panel's quantitative assessment alongside key themes from the qualitative feedback that contextualise the ratings and inform implementation considerations. Full qualitative analysis for the 15 consensus recommendations, is presented in S1 Text, and detailed stakeholder consultation drafts including implementation options are provided in S2–5 Texts.

Key Recommendations for Institutions

> **Box 1. Summary of Institutional Recommendations**
>
> *Consensus recommendations (effective and feasible by 2030):*
>
> • R19: Training in antibody validation in relevant bioscience courses and programmes (Eff 8.0 / Feas 7.0)
>
> • R20: Incorporate antibody validation into research integrity and ethics review frameworks (Eff 7.0 / Feas 7.0)
>
> • R21: Recognise and support local champions or experts in antibody validation (Eff 7.0 / Feas 7.0)
>
> *All three institutional items achieved full consensus (ie rated effective and feasible without disagreement). No items with uncertain feasibility. This is unique among stakeholder groups.*
>
> Median ratings: Eff = Effectiveness; Feas = Feasibility; rated on 9-point scales.

The panel viewed institutional actions as foundational to the broader reform effort, with one panellist noting that "without laying such a groundwork of training and advocacy, antibody validation will never become the norm."

Training in antibody validation received strong endorsement (median effectiveness 8.0, feasibility 7.0;

R19). The panel recommended that training should be offered in relevant bioscience courses and

programmes, including undergraduate education in fields such as immunology, neuroscience, and

molecular biology, postgraduate and doctoral training, and continuing professional development.

Importantly, the panel emphasised that training should be field-appropriate rather than universally

mandated across all domains. Panel members highlighted the need to address laboratory culture

alongside formal training, noting that "*in many cases the emphasis on validation and proper controls*

comes down to lab culture. If a student is trained to validate antibodies but the PI/senior lab members

do not agree this is a priority, it will not be done." The need to train established researchers as well as

trainees was a recurring theme: "*Need to train the senior staff as well!*"

The panel endorsed incorporating antibody validation expectations into institutional research integrity

and ethics review frameworks (median effectiveness 7.0, feasibility 7.0; R20). This could include

asking researchers to consider antibody validation as part of research ethics approval processes and

providing guidance or templates for documenting validation plans. However, panel members debated

the appropriate framing, with some questioning whether antibody validation constitutes an ethical issue

in the traditional sense: "*I don't really see this as an ethical issue. Lumping it into ethics with big issues*

like image manipulation, data fabrication, animal use, etc doesn't seem appropriate." Others noted that

the waste of animal and human samples resulting from poor antibody validation carries clear ethical

implications. Panel members also noted that "*a lot of research does not go through ethics committees,*"

emphasising the importance of complementary mechanisms beyond ethics review alone.

The third institutional recommendation called for universities, research institutions, and learned

societies to recognise and support local champions or experts in antibody validation (median

effectiveness 7.0, feasibility 7.0; R21). These individuals could serve as points of contact for advice on

antibody selection and validation design, trainers or facilitators in reproducibility workshops, advocates

for good practice within departments or core facilities, and contributors to cross-institutional

reproducibility networks. Support could include protected time, acknowledgement in performance

reviews, or funding for training and outreach. The panel viewed this as foundational, for example

noting that "*all of the questions in this section are not going to make antibody research more*

reproducible per se but will lay the foundation for increased awareness and adoption of best practice."

However, concerns about institutional buy-in and sustainability were raised, with panellists questioning

how to maintain expertise networks given high turnover of research staff and limited institutional

awareness of the problem.

Key Recommendations for Funders

Box 2. Summary of Funder Recommendations

Consensus recommendations (effective and feasible by 2030):

- R10: Require antibody validation plans in grant applications (Eff 7.5 / Feas 8.0)

- R11: Dedicated budget line for antibody validation in funded projects (Eff 8.0 / Feas 7.0)

- R12: Create or expand targeted tool development and validation schemes (Eff 8.0 / Feas 7.0)

- R13: Signal importance of antibody validation in applicant guidance (Eff 7.0 / Feas 7.0)

- R14: Encourage deposition of validation data in open-access repositories (Eff 7.0 / Feas 7.0)

- R15: Formally endorse community-developed reporting standards (e.g., IWGAV, MDAR) (Eff 7.0 / Feas 7.0)

Effective but feasibility uncertain (4 items):

- R17: Co-fund independent benchmarking initiatives (Eff 8.0 / Feas 6.0)

- R18: Promote researcher participation in benchmarking (Eff 7.0 / Feas 6.0)

> • R16: Directly engage manufacturers on transparency (Eff 7.0 / Feas 5.0)
>
> • A6: Mandate funded antibodies be recombinant and publicly available (Eff 7.0 / Feas 6.0)
>
> Median ratings: Eff = Effectiveness; Feas = Feasibility; rated on 9-point scales.

The panel viewed funders as occupying a critical position in the research lifecycle: before experiments

are conducted, when validation planning and budgeting can be embedded from the outset (full

qualitative comments in S7 and S8 Texts).

The panel reached consensus that funders should provide dedicated financial support for antibody

validation. This encompasses two complementary actions: applicants to biomedical funding schemes

should include a specific budget line item for resources to conduct robust antibody validation where

relevant to their proposal (median effectiveness 8.0, feasibility 7.0; R11), and funders should create or

expand targeted schemes to support the development and validation of critical antibody-based research

tools (median effectiveness 8.0, feasibility 7.0; R12). Both items received the highest effectiveness

ratings among funder recommendations. Panel members noted, however, that budget constraints could

pose challenges, with one participant observing that *"research proposals typically employ multiple*

antibodies (5–10+). Would the applicant be required to validate all antibodies prior to use? This

would result in a pretty significant budget increase."

The panel also endorsed embedding antibody validation expectations into grant application processes

through escalating approaches. At a minimum, funders should include information in applicant

guidance that antibody performance is an important limitation in many experimental methods and

encourage applicants to address this in methodological sections (median effectiveness 7.0, feasibility

7.0; R13). More substantively, funders should include a section in application forms where applicants

must detail the steps they will take to validate the antibodies used (median effectiveness 7.5, feasibility

8.0; R10). Panel members nevertheless cautioned against box-ticking: *"Another box ticking exercise for*

the applicant, which is filled with little care and using generic text each time." Panel members

cautioned that effectiveness would depend on whether validation plans are genuinely evaluated during

review.

Two further consensus recommendations addressed community standards and data sharing. Funders

should formally endorse community-developed reporting standards that promote antibody transparency

and validation, such as those from the IWGAV and MDAR frameworks (median effectiveness 7.0,

feasibility 7.0; R15). Funders should also encourage grantees to deposit antibody validation data in

open-access repositories, ideally linked to RRIDs or antibody registry entries (median effectiveness

7.0, feasibility 7.0; R14). Panel members noted that endorsement alone, while a *"feasible and low lift"*

action, may be insufficient without accompanying enforcement mechanisms, and expressed concern

about the quality of deposited data without robust review processes.

Beyond the consensus recommendations, co-funding of independent benchmarking initiatives such as

YCharOS were deemed effective (median effectiveness 8.0), but feasibility remained uncertain

(median 6.0; R17). Panel members agreed that creating suites of validated tools through independent

benchmarking would substantially improve data quality and increase return on investment, but

expressed concern about alignment with funder mandates and funding mechanism constraints. Several

additional funder recommendations were rated as effective but with feasibility concerns, including

promoting researcher participation in benchmarking (median effectiveness 7.0, feasibility 6.0; R18),

mandating that funded antibodies be recombinant and publicly available (median effectiveness 7.0,

feasibility 6.0; A6), and engaging manufacturers directly on transparency (median effectiveness 7.0,

feasibility 5.0; R16).

Key Recommendations for Manufacturers

<div style="border:1px solid blue; padding:1em;">

Box 3. Summary of Manufacturer Recommendations

Consensus recommendation (effective and feasible by 2030):

  • A7: Assign RRIDs to products at source (Eff 8.0 / Feas 7.0)

Effective but feasibility uncertain (2 items):

  • A8: Perform standard validation experiments and make data available (Eff 8.0 / Feas 6.5)

  • A9: Shift production toward recombinant antibodies (Eff 7.0 / Feas 5.0)

One consensus item; feasibility barriers for remaining items reflect commercial viability and market dynamics. S5 Text for consultation draft.

Median ratings: Eff = Effectiveness; Feas = Feasibility; rated on 9-point scales.

</div>

The manufacturer landscape presented a distinctive pattern: one recommendation achieved full

consensus, while the remaining items were rated as effective but with significant feasibility barriers

reflecting the commercial context in which manufacturers operate. Panel members, including

manufacturer representatives, repeatedly acknowledged that leading manufacturers already exemplify

good practice, while others lag behind.

The panel reached consensus that antibody manufacturers should assign Research Resource Identifiers

(RRIDs) to their products at source (median effectiveness 8.0, feasibility 7.0; A7). This was the only

manufacturer recommendation to achieve full consensus, and the response was notably enthusiastic,

with some panellists describing it as *"really, an easy one"* and noting it *"would be fantastic."* RRIDs

assigned at source would enable researchers to uniquely identify and cite specific antibodies,

facilitating systematic tracking of antibody performance across the literature and supporting automated

screening of submissions by journals. Panel members acknowledged, however, that some suppliers

may not cooperate: *"There are some suppliers that will ignore this, in the same way they have ignored*

the calls to improve validation."

Manufacturers performing a standard set of validation experiments and making the data available was

rated as effective (median effectiveness 8.0) but with uncertain feasibility (median 6.5; A8). The

tension between rigorous context-specific validation and commercial viability was central to panel

deliberations. Some panellists were emphatic that this was unrealistic, e.g. with one noting "*Not going*

to happen. Too expensive", while others noted that "*reputable manufacturers are already doing this*"

and that "*validation is a continual/on-going process for every antibody*." A shift toward recombinant

antibody production was similarly rated as effective (median 7.0) but with uncertain feasibility (median

5.0; A9), with the panel noting that manufacturer production decisions are ultimately driven by

customer demand.

 Key Recommendations for Publishers and Journals

Box 4. Summary of Publisher and Journal Recommendations

Consensus recommendations (effective and feasible by 2030):

   • R1: Authors include RRIDs for each antibody used (Eff 7.0 / Feas 8.0)

   • R2: Authors provide complete antibody metadata for unambiguous identification (Eff 8.0 / Feas 8.0)

   • R3: Authors report dilution ratio and protein concentration (Eff 8.0 / Feas 8.0)

   • A1: Journals establish clear antibody validation and reporting standards (Eff 7.5 / Feas 7.0)

Effective but feasibility uncertain (4 items, all Eff ≥7.0):

   • R4: Require validation data in open-access repositories (Eff 8.0 / Feas 5.0)

   • A2: Include validation protocols in manuscripts or supplements (Eff 8.0 / Feas 6.0)

   • R7: Automated tools to flag antibody validation issues (Eff 8.0 / Feas 5.0)

   • R9: Genetic validation for critical antibodies (Eff 8.0 / Feas 5.0)

   • R6: Train reviewers and editors on validation assessment (Eff 7.0 / Feas 5.0)

   • A3: Appoint specialist reproducibility editors (Eff 7.0 / Feas 5.5)

Four consensus items centred on reporting requirements. Other effective items face barriers around editorial resources, cost, and infrastructure. See S1 Text for full analysis of consensus items; S2 Text for consultation draft.

Median ratings: Eff = Effectiveness; Feas = Feasibility; rated on 9-point scales.

Publisher recommendations reflected a common pattern: the panel agreed that several interventions

would substantially improve research quality but identified practical barriers around editorial resources,

cost, and infrastructure. Four items achieved full consensus, while four additional items deemed

effective had uncertain feasibility.

The panel reached consensus on a package of reporting requirements that together enable unambiguous

antibody identification and experimental replication. Authors should include Research Resource

Identifiers for each antibody used where available (median effectiveness 7.0, feasibility 8.0; R1),

provide sufficient metadata to allow unambiguous identification of each antibody including clone ID,

catalogue number, lot number, and vendor (median effectiveness 8.0, feasibility 8.0; R2), and report the

dilution ratio and protein concentration where possible for each antibody used (median effectiveness

8.0, feasibility 8.0; R3). These three items work synergistically: complete metadata enables antibody

identification, while RRIDs provide persistent identifiers that link to characterisation databases and the

broader literature. The panel also reached consensus that journals should establish clear standards for

antibody validation and reporting that authors must meet (median effectiveness 7.5, feasibility 7.0; A1),

noting that this would only be maximally effective *"if there were community standards already in*

place or consensus across most journals."

The panel rated both the deposition of validation data in open-access repositories (median effectiveness

8.0, feasibility 5.0; R4) and the inclusion of detailed validation protocols in manuscripts or

supplementary data (median effectiveness 8.0, feasibility 6.0; A2) as effective, but with uncertain

feasibility. Similarly, the development and use of automated tools to flag potential antibody validation

issues in submitted manuscripts were predicted to be effective, but again uncertain feasibility (median

effectiveness 8.0, feasibility 5.0; R7). Panel members viewed automated tools as potentially

transformative but questioned who would fund their development and maintenance, and noted that *"not*

all publishers/journals have the financial means" to implement them. The requirement that authors

present genetic validation data for critical antibodies (median effectiveness 8.0, feasibility 5.0; R9) was

rated as effective but faced feasibility barriers around technical limitations (essential genes cannot be

knocked out), cost, and the difficulty of defining which antibodies are "critical" to a study's

conclusions.

Building editorial capacity for validation assessment was rated as effective but with feasibility

concerns. Both training reviewers and editors to evaluate antibody validation information (median

effectiveness 7.0, feasibility 5.0; R6) and appointing specialist reproducibility editors (median

effectiveness 7.0, feasibility 5.5; A3) faced scepticism about reviewer engagement and publisher

resources. Panellists doubted that reviewers would invest time in additional training given already

onerous review obligations, and many journals were seen as lacking resources for specialist editor

positions.

Cross-Stakeholder Consensus Items

Box 5. Summary of Cross-Stakeholder Items

Consensus recommendation:

  • R22: Develop a shared roadmap for improving antibody validation practices by 2030
  (Eff 7.0 / Feas 7.0)

Effective but feasibility uncertain:

  • R23: Coordinated shared infrastructure for aggregating antibody validation data (Eff 8.0
  / Feas 6.0)

  • A10: Establish a learned society for antibody validation (Eff 7.0 / Feas 5.0)

Rejected:

  • R5: Tiered scoring system or transparency badge (Eff 4.0 / Feas 3.0) — the only item
  rejected by the panel (uncertain effectiveness, infeasible)

Median ratings: Eff = Effectiveness; Feas = Feasibility; rated on 9-point scales.

Two cross-stakeholder items merit specific attention. The panel reached consensus that stakeholders

should work together to develop a shared roadmap for improving antibody validation practices by 2030

(median effectiveness 7.0, feasibility 7.0; R22). This roadmap would outline stakeholder-specific

actions and responsibilities, target timelines and milestones, and mechanisms for accountability and

feedback.

The development of coordinated, shared infrastructure for aggregating antibody validation data was

deemed potentially effective (median effectiveness 8.0) but uncertain feasibility (median 6.0; R23).

This infrastructure would aggregate validation data, provide standardised formats and tools for sharing,

and enable journals, funders, and institutions to access validation status or identify known problem

antibodies. Panel members viewed this as potentially transformative, supporting several other

recommendations, from automated manuscript screening to informed grant review, but recognised the

substantial coordination and resource challenges involved. Several recommendations across

stakeholder groups would benefit significantly from such shared infrastructure, suggesting that its

development could unlock a cascade of downstream improvements.

The establishment of a learned society for antibody validation was rated as effective (median 7.0) but

with uncertain feasibility (median 5.0; A10). While the panel recognised the potential value of a

professional body to coordinate standards, training, and advocacy, concerns were raised about

sustainability and the risk of duplicating existing networks. Conversely, the only item rejected by the

panel across both rounds was the introduction of a tiered scoring system or transparency badge for

antibody reporting (R5), which declined from uncertain to infeasible between rounds, reflecting

scepticism about badge-based incentive schemes in this context.

Discussion

This Delphi study engaged 32 international experts from across the research ecosystem to develop a

formally prioritised, consensus-based set of recommendations for addressing antibody validation

failures in biomedical research. The panel achieved consensus on 15 items as both effective and

feasible for implementation by 2030, spanning institutional, funder, manufacturer, and publisher

actions as well as cross-stakeholder coordination. An additional 15 items were rated as effective but

with uncertain feasibility, reflecting a consistent pattern in which experts were confident that proposed

interventions would improve antibody practices but less certain about realistic implementation within

the target timeframe. Together with existing technical solutions [1], data-sharing initiatives [17, 21],

and emerging stakeholder coordination efforts [12, 18], these findings provide a structured framework

for multi-stakeholder action.

Barriers to Implementation

Four interconnected barriers emerge from the panel's assessment that help explain both the persistence

of the antibody validation problem and the challenges of implementing solutions.

First, diffuse ownership enables collective inaction. Antibody validation failures impose costs across

the research system — wasted funding, unreliable literature, consumed biological materials — but no

single stakeholder bears the full cost or has authority to mandate change unilaterally. Researchers lack

incentives to validate when publication and funding systems do not reward the practice; publishers

hesitate to impose requirements that could divert submissions to less demanding competitors; funders

may view validation as outside their mandate; and manufacturers competing on price and catalogue

breadth face limited incentives to invest in validation when researchers continue to purchase on the

basis of previous use and supplier reputation [14]. Panel members described a pattern of mutual

deflection, with responsibility for validation perceived as belonging to other stakeholder groups rather

than one's own (S7 Text, R22). Several characterised the broader landscape as too fragmented for

meaningful alignment to feel achievable. However, the panel's endorsement of a shared roadmap (R22)

and the interdependencies across recommendations point toward a countervailing dynamic: participants

noted that journal requirements would create *de facto* mandates for manufacturers, that researcher

demand would shift manufacturer behaviour, and that funder investment in infrastructure would enable

practices that other stakeholders could then require.

Second, market dynamics create structural tensions. The research antibody market operates

predominantly under a Research Use Only model that prioritises catalogue breadth and low unit cost

over rigorous validation. Manufacturer representatives acknowledged significant economic barriers to

comprehensive validation, describing the costs as prohibitive and warning that extensive validation

programmes would reduce revenue without a corresponding market incentive (S7 Text, A8). At the

same time, one manufacturer described their company's ongoing commitment to validation, illustrating

that the barrier is structural rather than universal. Yet the apparent low cost of inadequately validated

antibodies is illusory when downstream costs are considered: failed experiments, repeated purchasing,

wasted samples, and unreliable findings impose far greater costs on the research system than better

validation would. Creating market conditions in which quality is rewarded — through purchasing

decisions informed by independent characterisation data, journal requirements that make validation

data visible, and funder signals that prioritise quality — is necessary to shift the equilibrium.

Third, resource justification challenges persist despite clear return on investment. Participants noted

competing demands on limited resources, with some questioning whether funders would prioritise

antibody validation given the breadth of competing demands on their budgets, even while

acknowledging the value of the proposed actions (S8 Text, A10). The economic case is compelling at a

system level — the estimated \$1 billion annually in direct waste in the US alone [12, 13], plus far

greater opportunity costs — but the return is diffuse and difficult for individual stakeholders to capture.

A funder investing in validation infrastructure benefits the entire community, not just its own grantees;

an institution investing in training produces graduates who carry improved practices elsewhere. The

benefits are real but distributed across time and across the ecosystem, and compounded by the

difficulty of demonstrating negative outcomes that were prevented.

Fourth, coordination challenges reflect the scale of the problem. With more than 22,000 human protein

targets, millions of antibody products, and a global research community, the coordination challenge is

immense. No single authority can mandate standards, and voluntary adoption requires alignment across

actors with different incentives, timescales, and resources. The panel's findings suggest, however, that

complete alignment is not a prerequisite for progress. The 15 consensus recommendations represent

actions individual stakeholders can implement without waiting for others, and participants emphasised

sequencing over simultaneity: several argued that funders must first act as enablers — establishing

resources, infrastructure, and training — before mandates can be effective (S8 Text, A5), while others

suggested that even imperfect initial standards would allow journals to demonstrate a credible

commitment to improving antibody reporting practices (S8 Text, A1).

These barriers are consistent with recent multi-stakeholder commentaries. Kahn et al. [12] noted that

the origins of the antibody crisis are 'many and varied' and that 'it is unrealistic to expect a complete

solution in the near future', while emphasising that significant progress is achievable through concerted

action. In earlier work by members of this study team [18], we argued that behavioural and cultural

change could be supported by engaging stakeholders at every level using established frameworks for

research culture change. The Delphi findings complement these analyses by providing quantitative

evidence of where expert agreement is strongest and where feasibility barriers remain.

Next Steps and Implementation

The findings lend themselves to a programme of targeted stakeholder consultation. To support this, we

have prepared separate documents for each stakeholder group — publishers, funders, institutions, and

manufacturers — presenting consensus recommendations alongside implementation options derived from the panel's qualitative feedback (S2–5 Texts). These present options rather than prescriptions, recognising that the optimal approach will vary across organisations, countries, and contexts. Items rated as effective with uncertain feasibility are not dismissed but treated as implementation challenges to be addressed through consultation. Interdependencies across stakeholder groups are made explicit throughout, supporting coordinated rather than siloed implementation.

Several elements of the recommended infrastructure already exist. The NC3Rs Antibody Champions programme provides a model for the local champions recommendation (R21) and delivers masterclasses for doctoral training aligned with R19. The OGA Academy offers structured online training [18], the OGA Antibody Database provides searchable characterisation data, and YCharOS continues to expand its open characterisation pipeline [1]. The eLife stakeholder roadmap [12] proposed actions that align closely with many consensus items, and the RRID initiative [17] provides identification infrastructure on which several publisher-focused recommendations depend. These resources demonstrate that the recommendations can be implemented by building on proven approaches. Notably, several recommendations are interdependent: shared infrastructure for aggregating validation data (R23) would support automated manuscript screening (R7), informed grant review (R10), and evidence-based purchasing decisions, suggesting that investment in infrastructure could unlock a cascade of downstream improvements across stakeholder groups.

Limitations

Several limitations should be considered. Participants were identified through existing professional networks, and the panel may therefore share perspectives that do not fully represent the diversity of views across the global research community. The panel was weighted toward the United Kingdom and United States, and the biomedical research ecosystem in other countries may face different challenges. Some sectors (particularly research funders, n=4) had few representatives, meaning feasibility

assessments for funder-directed recommendations may not capture the full diversity of perspectives.

The high retention rate (93.8%) and low disagreement suggest a relatively cohesive panel, which may

underestimate the range of views from a broader consultation.

The Delphi method captures expert opinion at a specific point in time and does not constitute empirical

evidence of effectiveness. The consensus recommendations represent the panel's assessment of what is

likely to be effective and feasible, but actual effectiveness will depend on implementation context. The

panel reached consensus after two rounds, which limited iterative refinement where uncertainty

persisted. However, rating stability between rounds suggests that additional rounds would have been

unlikely to shift results substantially. The qualitative data were synthesised narratively rather than

through formal thematic analysis, consistent with RAND/UCLA practice where free-text comments

inform item interpretation rather than serving as independent qualitative data. Full qualitative

commentary is provided in S7 and S8 Texts.

Conclusions

The rating of 'effective but uncertain feasibility' that characterises many of the proposed actions

should not be interpreted as reason for inaction. It reflects an honest assessment that the pace

and sequencing of change matter as much as the direction. The 15 consensus recommendations

provide an evidence-based starting point for immediate action; the remaining items rated as

effective represent a second wave whose feasibility is likely to increase as early implementation

builds infrastructure, shifts incentives, and establishes new norms. This staged approach —

enablement to expectation to mandate — was consistently endorsed by the panel and is reflected

in the consultation documents (S2–5 Texts). Together with existing technical and community

initiatives, these findings provide a structured basis for coordinated multi-stakeholder action to

address a persistent and costly source of unreliability in biomedical research.

Supporting Information

S1 Table. Detailed participant characteristics by employment sector.

S2 Table. Full quantitative results for Round 1 (n=32 participants, 23 items), including median

effectiveness and feasibility ratings, inter-percentile range (IPR), IPR adjusted for symmetry (IPRAS),

consensus classification, and disagreement assessment.

S3 Table. Full quantitative results for Round 2 (n=30 participants, 22 items: 12 re-rated from Round 1,

10 new), including median effectiveness and feasibility ratings, IPR, IPRAS, consensus classification,

and disagreement assessment.

S1 Text. Full item-by-item qualitative analysis with detailed recommendations and implementation
considerations for consensus items.

S2 Text. Publisher and journal consultation draft.

S3 Text. Research funder consultation draft.

S4 Text. Institutional and educational body consultation draft.

S5 Text. Antibody manufacturer consultation draft.

S6 Text. Full item sets for Round 1 and Round 2.

S7 Text. Round 1 qualitative feedback (participant comments).

S8 Text. Round 2 qualitative feedback (participant comments).

Data Availability

All data underlying the findings reported in this study are available within the article and its

Supporting Information files. Full quantitative results including median ratings, inter-percentile

ranges, and consensus classifications for all items across both Delphi rounds are provided in S2

and S3 Tables. Complete qualitative feedback from participants is provided in S7 and S8 Texts.

624 Individual-level raw ratings are not provided to protect participant anonymity, consistent with

625 the ethical approval for this study.

Competing Interests

HV and MB have received funding for a research studentship from Abcam Ltd, and contributions in

kind for manufacturers that contribute to the YCharOS Inc. consortium. RF is a director and

shareholder of Clinvivo Limited, a company specialising in Delphi studies. Clinvivo Limited

administered this Delphi study. The remaining authors declare no competing interests. These

relationships did not influence the study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Funding

634 This work was supported by a grant from the National Centre for the Replacement, Refinement and

635 Reduction of Animals in Research (NC3Rs) and Medical Research Council (MRC) (NC3Rs Ref:

636 NC/NAM0019/1, MRC UKRI076) alongside support from the Leicester Impact Accelerator Account

637 (BBSRC and MRC).

Acknowledgements

We thank the following individuals for their expert participation as Delphi panellists. Their

inclusion in this list acknowledges their contribution to the consensus process and does not

necessarily imply endorsement of the final manuscript or its recommendations:

Alex Ball (Senior Scientist, GeneTex); Anita Bandrowski (CEO and Co-founder, SciCrunch);

Andrew Bradbury (Founder and CSO, Specifica, an IQVIA business); Andrew Chalmers (CEO,

CiteAb); Alejandra Clark (Managing Editor, The Company of Biologists); Brian Dickie (Chief

Scientist, Motor Neurone Disease Association); Carly Dix (Associate Director, Biologics

References

1. Ayoubi R, Ryan J, Biddle MS, Alshafie W, Fotouhi M, Bolivar SG, et al. Scaling of an

antibody validation procedure enables quantification of antibody performance in major research

applications. Elife. 2023;12:RP91645. https://doi.org/10.7554/eLife.91645

2.   Elliott S, Busse L, Bass MB, Lu H, Sarosi I, Sinclair AM, et al. Anti-Epo receptor antibodies do

not predict Epo receptor expression. Blood. 2006;107(5):1892-5. https://doi.org/10.1182/blood-

2005-10-4066

3.   Berglund L, Björling E, Oksvold P, Fagerberg L, Asplund A, Al-Khalili Szigyarto C, et al. A

genecentric Human Protein Atlas for expression profiles based on antibodies. Mol Cell

Proteomics. 2008;7(10):2019-27. https://doi.org/10.1074/mcp.R800013-MCP200

4.   Schrohl AS, Pedersen HC, Jensen SS, Nielsen SL, Brünner N. Human epidermal growth factor

receptor 2 (HER2) immunoreactivity: specificity of three pharmacodiagnostic antibodies.

Histopathology. 2011;59(5):975-83. https://doi.org/10.1111/j.1365-2559.2011.04034.x

5.   Lukinavičius G, Lavogina D, Gönczy P, Johnsson K. Commercial Cdk1 antibodies recognize

the centrosomal protein Cep152. Biotechniques. 2013;55(3):111-4.

https://doi.org/10.2144/000114074

6.   Baker M. Reproducibility crisis: Blame it on the antibodies. Nature. 2015;521(7552):274-6.

https://doi.org/10.1038/521274a

7.   Andersson S, Sundberg M, Pristovsek N, Ibrahim A, Jonsson P, Katona B, et al. Insufficient

antibody validation challenges oestrogen receptor beta research. Nat Commun. 2017;8:15840.

https://doi.org/10.1038/ncomms15840

8.   Laflamme C, McKeever PM, Kumar R, Schwartz J, Kolahdouzan M, Chen CX, et

al. Implementation of an antibody characterization procedure and application to the major

ALS/FTD disease gene C9ORF72. Elife. 2019;8:e48363. https://doi.org/10.7554/eLife.48363

9.   Virk HS, Rekas MZ, Biddle MS, Wright AKA, Sousa J, Weston CA, et al. Validation of

antibodies for the specific detection of human TRPA1. Sci Rep. 2019;9(1):18500.

https://doi.org/10.1038/s41598-019-55133-7

10.   Kwon D. The antibodies don't work! The race to rid labs of molecules that ruin experiments.

Nature. 2024;635(8037):26-8. https://doi.org/10.1038/d41586-024-03590-0

11.   Freedman LP, Cockburn IM, Simcoe TS. The economics of reproducibility in preclinical

research. PLoS Biol. 2015;13(6):e1002165. https://doi.org/10.1371/journal.pbio.1002165

12.   Kahn RA, Virk H, Laflamme C, Houston DW, Polinski NK, Meijers R, et al. Antibody

characterization is critical to enhance reproducibility in biomedical research. Elife.

2024;13:e100211. https://doi.org/10.7554/eLife.100211

13.   Kahn RA, Virk HS, McPherson PS. Heed a decade of calls for antibody validation. Nature.

2023;620(7974):492. https://doi.org/10.1038/d41586-023-02566-w

14.   Biddle M, Cooper J, Blades K, Ruddy D, Krockow EM, Virk H. Drivers and ethical impacts of

insufficient validation of antibodies in research. bioRxiv [Preprint]. 2026 Feb 19 [cited 2026

Feb 20]. Available from: https://doi.org/10.64898/2026.02.19.706766

15.   Uhlen M, Bandrowski A, Carr S, Edwards A, Ellenberg J, Lundberg E, et al. A proposal for

validation of antibodies. Nat Methods. 2016;13(10):823-7. https://doi.org/10.1038/nmeth.3995

16.   Macleod M, Collings AM, Graf C, Kiermer V, Mellor D, Swaminathan S, et al. The MDAR

(Materials Design Analysis Reporting) Framework for transparent reporting in the life sciences.

Proc Natl Acad Sci U S A. 2021;118(17):e2103238118.

https://doi.org/10.1073/pnas.2103238118

17. Bandrowski AE, Martone ME. RRIDs: a simple step toward improving reproducibility through

rigor and transparency of experimental methods. Neuron. 2016;90(3):434-6.

https://doi.org/10.1016/j.neuron.2016.04.030

18. Biddle M, Stylianou P, Rekas M, Wright A, Sousa J, Ruddy D, et al. Improving the integrity

and reproducibility of research that uses antibodies: a technical, data sharing, behavioral and

policy challenge. mAbs. 2024;16(1):2323706. https://doi.org/10.1080/19420862.2024.2323706

19. Fitch K, Bernstein SJ, Aguilar MD, Burnand B, LaCalle JR, Lazaro P, et al. The RAND/UCLA

Appropriateness Method User's Manual. Santa Monica, CA: RAND Corporation; 2001.

20. Virk H, Eyre R, Holmes A, Biddle M, Krockow E. Defining the role of antibodies in improving

research reproducibility: NC3Rs and Only Good Antibodies community meeting report.

London: NC3Rs; 2024 [cited 2026 Feb 8]. Available from:

https://nc3rs.org.uk/sites/default/files/2024-07/NC3Rs-

OGA%20Meeting%20report%20%E2%80%93%20Defining%20the%20role%20of%20antibod

ies%20in%20improving%20research%20reproducibility.pdf

21. Fink A, Kosecoff J, Chassin M, Brook RH. Consensus methods: characteristics and guidelines

for use. Am J Public Health. 1984;74(9):979-83. https://doi.org/10.2105/ajph.74.9.979