Mast Cell Specific *Cyp11a1* Deficiency Promotes T Cell Mediated Immunity and Suppresses Tumour Metastasis in a Mouse Model of Melanoma

M. Abdul Alim [1, 2]*, Clara Veiga-Villauriz[1], Katie Butcher[1], Hosni A.M. Hussein[1, 3],

[1]Department of Pathology, University of Cambridge, Cambridge, CB2 1QP, UK

[2]Uppsala University, Department of Medical Biochemistry and Microbiology, Uppsala, Sweden

[3]Department of Microbiology, Faculty of Science, Al-Azhar University, Assiut 71524, Egypt

* Corresponding author: M. Abdul Alim, abdul.alim@imbim.uu.se

Keywords

Mast cells; Steroidogenesis; Cyp11a1; Immune suppression; Lung metastasis; Tumour microenvironment; Immune evasion; CD8☐ T cells; Melanoma

ABSTRACT
Mast cells are emerging players in malignant conditions, but the underlying molecular
mechanisms remain poorly defined. Based on previous studies showing that steroids can
impact on tumour progression in various settings, we here investigated whether mast cell-
derived steroid synthesis can have an impact on tumour metastasis in a melanoma model. To
this end, we used mice with mast cell-specific ablation of Cyp11a1, a key enzyme in steroid
synthesis. We show that lung colonization of melanoma nodules was markedly diminished in
mice with mast cell-specific ablation of *Cyp11a1*, accompanied by reduced infiltration of
mast cells into the lungs. *Cyp11a1* gene expression was significantly decreased in lungs of
mice with mast cell-specific ablation of Cyp11a1, indicating that mast cells account for a
substantial fraction of the total Cyp11a1 expression. Our results also revealed that the mast
cell-specific deletion of *Cyp11a1* led to an overall increase in CD107a/LAMP1 staining
intensity of the lung tissue, suggesting that mast cell-derived steroids can suppress immune
cell activation/degranulation. A further dissection of this finding by flow cytometry analysis
of individual immune cell populations revealed that CD8$^+$ T cells, NK cells and basophils
were activated to a higher extent in lungs from mice with mast cell-specific *Cyp11a1*
ablation. We also demonstrate that both CD8$^+$ and CD4$^+$ T cells in lungs of mice with mast
cell-specific deletion of *Cyp11a1* expressed elevated levels of IFN-γ in comparison with
controls. Altogether, these findings introduce a hitherto unrecognized role of a mast cell-
derived steroid axis in regulating tumour metastasis.

INTRODUCTION

Although being traditionally recognized for their role in allergic responses and pathogen defence, mast cells are increasingly recognized as active contributors in the tumour microenvironment, where they impact disease progression through a range of immunological and structural mechanisms [1-3]. Mast cells are key players in neuroimmune communication, mediating interactions with the peripheral nervous system and the immune system. A role for mast cells has also been suggested in range of additional settings, including tissue healing responses and inflammation, and metabolic disorders such as diabetes, hyperlipidemia and thyroiditis [4-8]. Given their known ability to interact with other immune cells and their capacity to influence extracellular matrix parameters, it is conceivable that mast cells can have an impact on tumour development and metastasis. However, this issue has not been studied to a large extent, and there is thus a need for a deeper understanding of mast cell functions in cancer immunology. A number of studies have approached these issues, and have revealed that mast cells can both suppress and promote tumourigenesis depending on the context [9, 10]. By releasing a wide range of preformed and newly synthesized mediators including cytokines (e.g., TNF□α, IL□6, IL□10, IL-33), proteases (e.g., tryptase, chymase), lipid mediators (e.g., prostaglandins, leukotrienes), and growth factors such as VEGF, mast cells may have the capacity to shape angiogenesis, immune recruitment, and extracellular matrix dynamics [11-13]. However, the precise molecular pathways through which mast cells influence tumour progression and metastatic niche formation remain poorly understood [10, 14, 15].

De novo steroidogenesis has emerged as a non-classical immunomodulatory mechanism. In this process, the mitochondrial rate limiting enzyme CYP11A1 catalyses the conversion of cholesterol into pregnenolone - a reaction that initiates steroid hormone biosynthesis [16, 17]. Steroid biosynthesis, though classically attributed to endocrine tissues, has increasingly been documented in immune populations, where locally synthesized steroids can act as immunoregulatory signals [18-23]. Further, it has recently been shown that tumour-derived pregnenolone promotes bone metastasis by enhancing osteoclastogenesis through prolyl 4□hydroxylase subunit beta (P4HB), in breast cancer and melanoma models [24]. It has also been demonstrated that tumour-associated immune cells can utilize the steroidogenesis pathway to promote immune suppression [25, 26]. However, the role of mast cell steroidogenesis and its relevance in shaping the immune landscape within tumour contexts has not been explored.

In this study, we investigated the role of mast cell–derived steroidogenesis in shaping the immune microenvironment during pulmonary metastasis of melanoma. Using a genetically engineered mouse model with mast cell–specific ablation of Cyp11a1, we demonstrate that mast cell–derived steroidogenesis mediates immune suppression within lung tumours, which is associated with enhanced lung tumour colonization. Hence, this study introduces a novel mast cell-steroid axis in regulating tumour progression.

MATERIALS AND METHODS

Mice and Ethical Approval
In this study, we used C57BL/6J mice, originally sourced from The Jackson Laboratory. Mast cell–specific Cyp11a1 knockout mice (cKO) were generated by crossing Cpa3-Cre transgenic mice (JAX stock #026828, Jackson laboratory) with Cyp11a1$^{fl/fl}$ animals (generated by Sanger Institute) [19, 27]. Littermates possessing the Cyp11a1$^{fl/fl}$ genotype but lacking the Cpa3-Cre transgene served as wild-type (WT) controls. All mice were housed in a specific pathogen–free (SPF) facility under controlled conditions, including a 12-hour light/12-hour dark cycle. Both male and female mice were included, with random allocation to experimental groups to minimize bias. All procedures involving animals adhered to the UK Animals (Scientific Procedures) Act 1986 and were conducted under Home Office Project License (PPL PP2448972). Ethical approval was obtained from the institutional Animal Welfare and Ethical Review Body, described elsewhere [19].

Melanoma Lung Metastasis Model
For metastatic challenge, 0.5×10^6 B16-F10 melanoma cells were injected intravenously via the tail vein into 12-18-weeks-old control (n=8) and cKO (n=8) mice. Animals were euthanized at day 13 post-injection for endpoint analyses unless otherwise indicated (**Fig. 1**). To assess general health and systemic effects of tumour burden, body weight was recorded at baseline (day 0), day 7, and day 13 following tumour cell injection. Weight measurements were performed at the same time each day using a digital scale. The weight range of the mice was 19.9-27.7 g for WT controls, and 23.8-34.6 g for cKO mice, and the age of the mice was between 10-18 weeks at the start of the experiment. At the conclusion of the study, all mice were euthanized by cervical dislocation in accordance with institutional guidelines, ensuring a rapid and ethical procedure.

Tumour Burden Quantification and Tissue Processing
At endpoint, lungs were perfused by ice-cold PBS and harvested in PBS supplemented with 2.5% FBS and 2 mM EDTA and stored on ice until further processing. Lungs were rinsed in ice-cold PBS to enhance nodule visibility. Visible surface tumour nodules were counted manually with two independent observers (blinded) as well as under a microscope after basic histological staining, and representative images were captured and quantified in a semi-quantitative fashion.

For histological analysis, a small portion of lung tissue was immediately frozen (-80°C). The remaining tissue was placed in PBS within a 2 ml microcentrifuge tube and manually chopped into small pieces using sterile scissors. A digestion cocktail was freshly prepared with DMEM/F12 supplemented with125 µl (1 mg/ml) of collagenase A (Roche), 125 µl (1 mg/ml) of collagenase D (Roche), and 35 µl of DNase I (0.2–0.4 mg/ml). Samples were incubated at 37°C with gentle agitation (Eppendorf ThermoMixer C) for 30 minutes. To terminate enzymatic activity, cells were washed in PBS containing 2% FBS and 2 mM EDTA (FACS buffer). Digested tissue was filtered through 70 µm nylon mesh strainers (Falcon) into 50 ml conical tubes and washed with 30 ml cold PBS. Cell suspensions were centrifuged at $400\times$ g for 5 min at 4°C. Pellets were briefly vortexed and incubated with 3 ml of $1\times$ red blood cell lysis buffer at room temperature for 3 min. The lysis reaction was halted by dilution with PBS, followed by centrifugation under the same conditions. Cells

128 were resuspended in PBS, filtered through 30 μm strainers, and kept on ice for immediate
129 downstream processing.

Histology and Toluidine Blue Staining

Frozen lung tissues were sectioned at 7 μm thickness under a cryostat (-20°C) and the slices
were fixed with ice-cold acetone (–20°C) for 20 min. Sections were stained with
hematoxylin and eosin (H&E) to evaluate tumour architecture. For mast cell visualization,
adjacent sections were stained with Toluidine blue (0.1% in 0.5N HCl) for 3 min at room
temperature. Mast cells were quantified in peritumoural regions from at least three-five high-
power fields (HPF) per section. Staining was imaged using a Zeiss AxioObserver.Z1
Widefield microscope.

Immunofluorescence and Confocal Microscopy

Lung tissue slices from control (Cyp11a1-sufficient) and mast cell–specific Cyp11a1
knockout (cKO i.e., Cyp11a1- deficient) mice were fixed in cold acetone (–20 °C) for 20
min. Staining with antibodies was performed following protocols described previously [4, 28]
Briefly, tissue sections were blocked with normal horse and/or goat serum, followed by three
washes in PBS. Slides were then incubated overnight at 4°C in the dark with the following
primary antibodies: Monoclonal Mouse Mast Cell Tryptase (ab2378, Abcam) diluted 1:100,
CYP11A1 (D8F4F) Rabbit mAb (Cell Signaling Technology, #14217) diluted 1:200, Mouse
CD45 (FITC) diluted 1:200, and CD107a PE diluted 1:200.

After incubation with primary antibodies, tissue sections were washed three times with PBS
and subsequently incubated for 60 min at room temperature with secondary antibodies. These
included anti-rabbit IgG F(ab')□ Alexa Fluor 488 (Cell Signaling Technology, #4412S) or
goat anti-rabbit multi-rAb CoraLite Plus 555 (Proteintech, #RGAR003), both diluted 1:200,
and F(ab')□ goat anti-mouse IgG Alexa Fluor 647 (Proteintech, #SA00014-10), also diluted
1:200. Nuclei were counterstained with DAPI (4′,6-diamidino-2-phenylindole; Invitrogen).
Confocal microscopy was performed using a Zeiss LSM700 confocal microscope. Images
were captured at original magnifications of ×200 or ×400. Image processing and analysis
were conducted using ImageJ software (Fiji, ImageJ2).

Flow Cytometry

The resulting single-cell suspensions were stained for flowcytometry using FACS buffer
(PBS containing 2% FBS and 2 mM EDTA). Cells were incubated with anti-CD16/32 (Fc
block; eBioscience or BioLegend) for 10 min at 4°C, followed by surface staining in FACS
buffer with fluorochrome-conjugated antibodies against the following markers: CD45
(1:1,000), CD11c or I-A/I-E (1:1,000), CD4 (1:1,000), CD8 (1:400), CD11b, FcεRI (1:400),
NK1.1 (CD161) (1:500), CD117 (c-Kit), CD170 (Siglec-F), CD19 (1:800), and CD107a
(1:200). Dead cells were excluded using Ghost Dye UV 450 (CYTEK, 1:1,000).

For intracellular cytokine staining, cells were stimulated overnight at 37°C with PMA
(50 ng/mL) and ionomycin (500 ng/mL) in the presence of brefeldin A (5 μg/mL; BD
Biosciences). After surface staining and fixation/permeabilization (BD Cytofix/Cytoperm
kit), cells were stained with antibodies against CD3 (1:800), IFN-γ (1:400), and TNF-α
(1:400). All antibodies were titrated for optimal performance. Data acquisition was
performed on a Cytek Aurora spectral flow cytometer (Cytek Biosciences), and spectral
unmixing and data analysis were conducted using FlowJo v10. Immune cell subsets were
quantified as a percentage of live CD45□ cells. Mast cells were identified as FcεRI□
CD117□.

RNA Extraction, cDNA Synthesis and Quantitative PCR

Frozen lung tissue samples were homogenized, and total RNA was extracted using the
RNeasy Mini Kit (Qiagen) following the manufacturer's guidelines. RNA quantity was
determined using a NanoDrop spectrophotometer (Thermo Fisher Scientific), and RNA
integrity was assessed with the Agilent 2100 Bioanalyzer (Agilent Technologies).

For cDNA synthesis, 250 ng of total RNA was reverse transcribed using the iScript cDNA
Synthesis Kit (Bio-Rad), according to the provided protocol. Quantitative PCR was carried
out using SYBR Green Supermix (Bio-Rad) on a CFX384 Real-Time PCR System (Bio-
Rad). Primer sequences were either designed using PrimerBank or obtained as pre-validated
primer assays from Bio-Rad (Supplementary Table 1). All primer pairs were tested for
amplification efficiency prior to use to ensure accurate quantification.

Relative gene expression levels were calculated using the comparative Ct ($\Delta\Delta$Ct) method.
Actb and Gapdh were used as endogenous reference genes for normalization. All reactions
were run in technical duplicates. Statistical analysis was performed using unpaired two-tailed
t-tests to assess differences in gene expression between groups.

Statistical analyses

Data are presented as mean $\pm$ standard error of the mean (SEM). Comparisons between
two groups were made using unpaired, two-tailed Student's t-test. P values < 0.05 were
considered statistically significant. All analyses were performed using GraphPad Prism
(v10.0).

RESULTS

Mast cell–derived Cyp11a1 promotes lung metastatic colonization of melanoma cells and mast cell infiltration

To investigate the role of mast cell–expressed Cyp11a1 in metastatic tumour growth, we compared lung tumour burden in control (Cyp11a1-sufficient) and mast cell–specific *Cyp11a1* knockout (cKO) mice following intravenous injection of melanoma cells (**Fig. 1**). Control mice exhibited a significantly greater number of lung tumour nodules relative to cKO counterparts, indicating that mast cell-derived Cyp11a1 facilitates metastatic colonization and tumour outgrowth in the lung microenvironment (**Fig. 2a-b**). Importantly, body weight monitoring over the course of tumour progression revealed no significant differences between groups, excluding systemic illness or cachexia as confounding factors for tumour burden disparities (**Fig. 2c**). Histological analysis of lung sections using hematoxylin/eosin and toluidine blue staining demonstrated enhanced tumour nodule formation and increased mast cell infiltration in control lungs compared to those from cKO mice (**Fig. 2d-e**). In agreement with this, quantitative real time PCR (qPCR) measurements demonstrated that the expression of mast cell-specific genes (Mcpt1, Mcpt6) was profoundly diminished in lungs of cKO vs. control mice (**Fig. 2f-g**). These findings indicate that mast cell-expressed Cyp11a1 promotes melanoma colonization of the lung, and that mast cell-expressed Cyp11a1 has an autocrine function to support mast cell accumulation in the context of melanoma metastasis to the lung.

Cyp11a1 is expressed in the mast cell niche in lung tumours

Confocal microscopy analysis of lung tumour sections revealed a large extent of colocalization of tryptase (a mast cell marker) and the steroidogenic enzyme Cyp11a1 in control (Cyp11a1-sufficient) mice, indicating a high abundance of steroidogenic mast cells within the metastatic lung microenvironment (**Fig. 3a-b**). In contrast, tumours from mast cell–specific Cyp11a1 knockout (cKO) mice exhibited markedly reduced Cyp11a1: tryptase colocalization, consistent with diminished Cyp11a1 expression in mast cells (**Fig. 3a-b**).

Targeted deletion of *Cyp11a1* in Mast cells is associated with reduced CD45☐ immune cell activation in lung tumours

Next, we asked whether mast cell-specific deletion of Cyp11a1 has an effect on the overall population of Cyp11a1-positive cells, and on the overall accumulation of leukocytes in the lung tumours. To this end, tumour sections were stained for the pan-leukocyte marker CD45 (red) and Cyp11a1(green) (**Fig. 4a**). These analyses revealed that the overall intensity of Cyp11a1 staining was markedly diminished in tumours from cKO vs. control mice, indicating that mast cells account for a large fraction of the total population of Cyp11a1-positive cells. The reduction of overall Cyp11a1 expression in lungs of cKO mice was also supported by qPCR analysis, which revealed a profound reduction in *Cyp11a1* mRNA levels in lungs of cKO vs. control mice (**Fig. 3c**).

In contrast, similar intensities of CD45 staining were seen in tumours of both control- and cKO mice (**Fig. 4b**; lower panel), suggesting that the absence of Cyp11a1 expression in mast cells does not affect the overall immune cell infiltration to the lung. To assess for the extent of overall immune cell degranulation, we stained for CD107a (also known as LAMP1), a protein that is transported to the cell surface upon degranulation of immune cells (including mast cells, cytotoxic T lymphocytes, basophils, eosinophils). Interestingly, the overall positivity for CD107a was markedly higher in cKO vs. control mice (**Fig. 4c-d**). indicating

249 that mast cell-dependent steroidogenesis regulates immune cell activation within the lung
250 tumours.
251
Effect of mast cell–derived Cyp11a1 on lung immune cell populations
To provide further insight into how mast cell-dependent steroidogenesis regulates the
immune landscape of the melanoma tumours of the lung, we performed flow cytometry
analysis of various immune cells subsets. These analyses demonstrated comparable
frequencies of T cells ($CD4^+$ and $CD8^+$), B cells, NK cells, eosinophils and basophils
between control- and mast cell-specific *Cyp11a1* knockout (cKO) mice during baseline
conditions (**Fig. 5**).
259
Loss of mast cell–derived Cyp11a1 enhances immune effector cell activation
Based on the demonstration that mast cell-specific expression of Cyp11a1 suppresses the
overall population of CD107a-positive (degranulated) immune cells in the lung melanoma
tumours, we next assessed corresponding effects on individual immune cell types. As seen in
Fig. 6, the proportions of activated (degranulated) $CD8^+$ T-cells, NK cells and basophils were
significantly higher in the tumours from cKO vs. control mice at baseline (unstimulated; US),
hence indicating that mast cells, via Cyp11a1, suppressed the activation of these immune cell
populations. After PMA stimulation (S), the expression of CD107a remained significantly
higher in $CD8^+$ T cells and basophils in the lungs of cKO vs. control mice, whereas CD107a
expression in NK cells was not significantly different between the genotypes after PMA
stimulation (**Fig. 6**). Notably, although the depletion of mast cell-expressed *Cyp11a1* did not
affect the cell surface expression of CD107a in mast cells at baseline conditions, we noted
that PMA-activated cKO mast cells exposed lower levels of cell surface CD107a than control
cells, possibly reflecting an effect of intrinsic steroidogenesis on their capacity to undergo
activation.
275
Mast cell-derived steroidogenesis restrains T cell effector responses in the tumour
microenvironment
The results above indicate that mast cell steroidogenesis impacts on the ability of $CD8^+$ T
cells to undergo degranulation. Given the central role of T cells in cytotoxicity against
tumour cells, we next performed experiments to further assess the effects of mast cell-
expressed Cyp11a1 on T cell functionality. To this end, we performed *ex vivo* stimulation of
lung immune cells isolated from control and mast cell–specific *Cyp11a1* knockout (cKO)
mice. Following PMA activation, both CD4☐ helper and CD8☐ cytotoxic T cells from cKO
mice produced significantly higher levels of interferon-gamma (IFN-γ) compared to those
from control animals (**Fig. 7a**). These findings suggest that Cyp11a1-mediated steroid
synthesis in mast cells exerts an immunosuppressive influence on T cell activation. In
contrast, no significant effect of the depletion of mast cell-expressed Cyp11a1 on IFN-γ
expression was seen in NK cells. We also assessed whether mast cell Cyp11a1 has effects on
the expression of TNF-α. These analyses revealed that mast cells deficient in *Cyp11a1* had a
profoundly reduced ability to express TNF-α protein, whereas no corresponding effects on
TNF-α levels were seen in either $CD4^+$ T-cells, $CD8^+$ T cells, NK cells or B cells (**Fig. 7a**).
In agreement with these findings, qPCR analyses confirmed increased expression of IFN-γ
mRNA in the lungs of cKO vs. control mice, whereas no significant effects of the mast cell-
expressed Cyp11a1 on TNF-α mRNA levels were seen (**Fig. 7b**).
295

DISCUSSION

Numerous previous studies have implicated mast cells in melanoma progression. In clinical studies, the presence and abundance of mast cells has frequently been associated with poor prognosis, but there are also clinical studies in which the presence of mast cells has been linked to favourable outcomes [29-34]. The impact of mast cells on melanoma progression has also been studied in animal models. In melanoma models similar to the model used herein, two independent studies have shown that total deficiency in mast cells is associated with a more pronounced lung colonization of melanoma cells, i.e. pointing to an altogether harmful role of mast cells in such settings [35, 36]. In contrast, the growth of the primary tumour was not affected by the total absence of mast cells [35, 36]. Notably, these studies hence reflect the total impact of mast cells in these settings, taking into account the combined effect of all of the compounds that mast cells express under a given condition. As another approach, effects of individual mast cell-expressed compounds can be studied. Intriguingly, such studies have revealed that the absence of mast cell-specific proteases is associated with enhanced lung metastasis and growth of primary tumours in melanoma models [35, 37], pointing to anti-tumour effects mediated by these mast cell-specific enzymes. Hence, this indicates that mast cells can express both tumor-promoting and suppressive effects, with a conceivable scenario being that the net effect of mast cells on tumour progression reflects the balanced output of such pro- and anti-tumour factors under given tumour conditions.

As noted above, several previous reports have suggested that mast cells may promote melanoma progression. This could potentially be linked to the ability of mast cells to express various tumour-promoting growth factors, including VEGF, FGF and TGF-β [38-40]. However, in this study we have evaluated an alternative scenario by which mast cells may halter melanoma progression. In previous studies it has been shown T cells can be induced to produce steroids in tumour settings, and that such T cell-derived steroids can cause immunosuppression leading to enhanced tumour growth in a melanoma model [19]. Given the reported role of mast cells in regulating melanoma metastasis to the lung [35, 36], we here asked whether the impact of mast cells on melanoma progression can be attributable to mast cell-derived steroid synthesis. This was approached by using mice in which *Cyp11a1* expression was conditionally ablated in mast cells.

In support for a role of mast cell-derived steroids in regulating melanoma metastasis, our data reveal a substantial reduction of metastatic colonisation of the lung in mice in which mast cells lack the expression of Cyp11a1, a key rate-limiting enzyme in steroid synthesis [16, 17]. Intriguingly, mast cell specific deletion of *Cyp11a1* was associated with an overall decrease in the mast cell numbers in the lungs of tumour-bearing mice, as shown both by staining for mast cells and by analysing the levels of mast cell-specific transcripts. One potential explanation for this finding could be that mast cell-derived steroids act in an autocrine, stimulatory effects on mast cell accumulation in the lung. However, the molecular mechanism behind this effect remains to be explored.

Notably, Cyp11a1 was not ubiquitously expressed among the lung mast cell populations, as evidenced by the existence of both *Cyp11a1*[+] and *Cyp11a1*[-] mast cells. It was also notable that numerous cells of the lung expressed Cyp11a1 but were negative for mast cell markers. However, the expression of the *Cyp11a1* gene was considerably reduced in lungs of mice with mast cell-specific deletion of Cyp11a1, indicating that mast cells in fact account for a substantial fraction of the total Cyp11a1 expression. The latter was also supported by

immunohistochemical staining, which revealed a clear reduction in the overall Cyp11a1
expression in lungs of mice with mast cell-specific deletion of *Cyp11a1*.

An intriguing finding in this study was that the overall positivity for CD107a/LAMP1 was
substantially higher in lungs of tumour bearing mice with mast cell-specific deletion of
Cyp11a1 vs. control mice. CD107a/LAMP1 is a protein present in the membrane of secretory
granules/lysosomes of various immune cell types. In their resting state, CD107a/LAMP1 is
hence intracellular, but when the cells are activated, this can lead degranulation, accompanied
by fusion of the granule/lysosome membranes with the cell membrane, accompanied by cell
surface exposure of CD107a/LAMP1. Cell surface exposure of CD107a/LAMP1 can thus
serve as a marker for immune cell activation/degranulation [41-43]. Hence, the increase in
CD107a positivity as a consequence of mast cell-specific deletion of *Cyp11a1* indicates that
mast cell-derived steroids may suppress immune cell activation/degranulation. To further
substantiate this finding, we employed flow cytometry analysis to evaluate the effects of mast
cell Cyp11a1 expression on individual immune cell populations. These analyses did not
reveal any significant effects of mast-cell Cyp11a1 on the numbers of either T cells (CD4$^+$ or
CD8$^+$), B cells, NK cells eosinophils or basophils within the lungs of tumour-bearing mice.
However, when assessing the activation status of the respective immune cell populations by
assessing CD107a/LAMP1 positivity, we noted that CD8$^+$ T cells, NK cells and basophils
were activated/degranulated to a higher extent in lungs of mice with mast cell-specific
ablation of *Cyp11a1* expression in comparison with controls. Hence, these findings suggest
that mast cell-derived steroid synthesis has the capacity to suppress the activation of these
respective immune cell populations. Notably, CD8$^+$ T cells and NK cells are known to have
major roles in the regulation of tumour cell growth [44], and the relieved suppression of these
cell populations in the absence of mast cell-derived Cyp11a1 may consequently account, at
least partly, for the reduced tumour development seen in these mice.

To provide deeper insight into this issue, we also assessed whether mast cell-specific deletion
of *Cyp11a1* had an impact on the expression of classical markers for immune cell activation,
by focusing on IFN-γ and TNF-α. Indeed, we found that CD8$^+$ T cells in lungs from mice
with mast cell-specific deletion of *Cyp11a1* exhibited elevated levels of IFN-γ, as assessed by
flow cytometry analysis. Elevated IFN-γ levels were also seen in CD4$^+$ T cells in lungs from
mice with mast cell-specific depletion of *Cyp11a1*.

Altogether, this study introduces a hitherto unrecognised impact of mast cell-derived steroid
synthesis on melanoma metastasis to the lung. These findings may thus be related to the
emerging awareness that immunosuppression by steroids can be linked to increased
resistance to cancer therapy [45, 46]. Based on the present data, we may envision that
targeting of mast cells by regimes that suppress their capacity to release steroids may become
explored as potential therapeutic options to suppress melanoma metastasis in clinical settings.
However, future work is required to evaluate the impact of our present findings in human
clinical settings. It is also important to note that, while we implicate mast cell–derived
steroids as modulators of immune suppression, the precise steroid species involved and their
downstream signalling pathways remain to be elucidated. Further investigations are also
warranted to assess the impact of a mast cell-steroid axis in tumour settings other than
melanoma.

ACKNOWLEDGEMENTS

We would like to thank the Flow Cytometry Facility (Joana Cerveira and Sameen Khan), Confocal Microscopy Facility (Ben Sutcliffe and Jonathan Howe), the Histology Facility at the Department of Pathology, and the UBS Animal Facility (Polly Seaton) at the Gurdon Institute, for technical assistance and animal support. We would like to thank Prof. Heike Laman for support and constructive feedback on the study. We would like to thank Fabio Melo, Uppsala University, for ordering validated primers from the Bio-Rad.

FUNDING

This work was supported by the Swedish Society for Medical Research (SSMF) Postdoctoral Grant (PG-22-0477-H-01, project no: 465602001AA/SSMF) and the Swedish Cancer Society (GP; 22 1988 Pj).

AUTHOR CONTRIBUTIONS

AA: Planned, designed, and performed experiments; analysed, assembled, visualized, and quantified data; wrote the manuscript.

CV and KB: Assisted with intravenous injections and animal studies.

SKS: Contributed to tissue processing.

HH: Assisted with tissue processing, FACS analysis and RNA extraction.

FINANCIAL INTERESTS

The authors have no relevant financial or non-financial interests to disclose.

REFERENCES

1. Galli, S.J. and M. Tsai, *IgE and mast cells in allergic disease.* Nature Medicine, 2012. **18**(5): p. 693-704.

2. Shi, S., et al., *Focus on mast cells in the tumor microenvironment: Current knowledge and future directions.* Biochim Biophys Acta Rev Cancer, 2023. **1878**(1): p. 188845.

3. Panagi, M., et al., *Stabilizing Tumor-Resident Mast Cells Restores T-Cell Infiltration and Sensitizes Sarcomas to PD-L1 Inhibition.* Clinical Cancer Research, 2024. **30**(11): p. 2582-2597.

4. Alim, M.A., et al., *Glutamate triggers the expression of functional ionotropic and metabotropic glutamate receptors in mast cells.* Cell Mol Immunol, 2021. **18**(10): p. 2383-2392.

5. Alim, M.A., M. Peterson, and G. Pejler, *Do Mast Cells Have a Role in Tendon Healing and Inflammation?* Cells, 2020. **9**(5).

6. Grujic, M., et al., *Mast Cells are Dependent on Glucose Transporter 1 (GLUT1) and GLUT3 for IgE-mediated Activation.* Inflammation, 2024. **47**(5): p. 1820-1836.

7. Toda, S., et al., *Growth factor-expressing mast cells accumulate at the thyroid tissue-regenerative site of subacute thyroiditis.* Thyroid, 2000. **10**(5): p. 381-6.

8. Cheng, Y., et al., *Large Accumulation of Collagen and Increased Activation of Mast Cells in Hearts of Mice with Hyperlipidemia.* Arq Bras Cardiol, 2017. **109**(5): p. 404-409.

9. Marichal, T., M. Tsai, and S.J. Galli, *Mast Cells: Potential Positive and Negative Roles in Tumor Biology.* Cancer Immunology Research, 2013. **1**(5): p. 269-279.

10. Longo, V., et al., *Controversial role of mast cells in NSCLC tumor progression and angiogenesis.* Thorac Cancer, 2022. **13**(21): p. 2929-2934.

11. Jackson, C.W., et al., *Mast Cells in and Their Cellular Interactions Within the Tumor Environment*, in *Handbook of Cancer and Immunology*, N. Rezaei, Editor. 2025, Springer Nature Switzerland: Cham. p. 1-28.

12. Liu, X., et al., *Mast cells in colorectal cancer tumour progression, angiogenesis, and lymphangiogenesis.* Front Immunol, 2023. **14**: p. 1209056.

13. Komi, D.E.A. and F.A. Redegeld, *Role of Mast Cells in Shaping the Tumor Microenvironment.* Clin Rev Allergy Immunol, 2020. **58**(3): p. 313-325.

14. Guo, X., et al., *Role of mast cells activation in the tumor immune microenvironment and immunotherapy of cancers.* Eur J Pharmacol, 2023. **960**: p. 176103.

15. Wensman, H., et al., *Tumor-mast cell interactions: induction of pro-tumorigenic genes and anti-tumorigenic 4-1BB in MCs in response to Lewis Lung Carcinoma.* Mol Immunol, 2012. **50**(4): p. 210-9.

16. Libert, C. and L. Dejager, *How Steroids Steer T Cells.* Cell Reports, 2014. **7**(4): p. 938-939.

17. Hanukoglu, I., *Steroidogenic Enzymes - Structure, Function, and Role in Regulation of Steroid-Hormone Biosynthesis.* Journal of Steroid Biochemistry and Molecular Biology, 1992. **43**(8): p. 779-804.

18. Taves, M.D., C.E. Gomez-Sanchez, and K.K. Soma, *Extra-adrenal glucocorticoids and mineralocorticoids: evidence for local synthesis, regulation, and function.* Am J Physiol Endocrinol Metab, 2011. **301**(1): p. E11-24.

19. Mahata, B., et al., *Tumors induce de novo steroid biosynthesis in T cells to evade immunity.* Nature Communications, 2020. **11**(1): p. 3588.

20. Chakraborty, S., J. Pramanik, and B. Mahata, *Revisiting steroidogenesis and its role in immune regulation with the advanced tools and technologies.* Genes & Immunity, 2021. **22**(3): p. 125-140.

21. Stanczyk, F.Z., et al., *Biosynthesis, Transport, and Metabolism of Steroid Hormones*,
in *Hormones, Genes, And Cancer*. 2003, Oxford University Press. p. 0.
22. Schmidt, K.L., et al., *Neurosteroids, immunosteroids, and the Balkanization of*
endocrinology. Gen Comp Endocrinol, 2008. **157**(3): p. 266-74.
23. Slominski, R.M., et al., *Extra-adrenal glucocorticoid biosynthesis: implications for*
autoimmune and inflammatory disorders. Genes Immun, 2020. **21**(3): p. 150-168.
24. Sandor, L.F., et al., *De novo steroidogenesis in tumor cells drives bone metastasis and*
osteoclastogenesis. Cell Rep, 2024. **43**(3): p. 113936.
25. Arner, E.N. and J.C. Rathmell, *Metabolic programming and immune suppression in*
the tumor microenvironment. Cancer Cell, 2023. **41**(3): p. 421-433.
26. Kim, I.S., et al., *Immuno-subtyping of breast cancer reveals distinct myeloid cell*
profiles and immunotherapy resistance mechanisms. Nature Cell Biology, 2019.
21(9): p. 1113-1126.
27. Lilla, J.N., et al., *Reduced mast cell and basophil numbers and function in Cpa3-Cre;*
Mcl-1fl/fl mice. Blood, 2011. **118**(26): p. 6930-8.
28. Alim, M.A., et al., *Pleckstrin Levels Are Increased in Patients with Chronic*
Periodontitis and Regulated via the MAP Kinase-p38α Signaling Pathway in Gingival
Fibroblasts. Front Immunol, 2021. **12**: p. 801096.
29. Grimbaldeston, M.A., et al., *Association between melanoma and dermal mast cell*
prevalence in sun-unexposed skin. Br J Dermatol, 2004. **150**(5): p. 895-903.
30. Ribatti, D., et al., *Tumor vascularity and tryptase-positive mast cells correlate with a*
poor prognosis in melanoma. Eur J Clin Invest, 2003. **33**(5): p. 420-5.
31. Ribatti, D., et al., *Neovascularisation, expression of fibroblast growth factor-2, and*
mast cells with tryptase activity increase simultaneously with pathological
progression in human malignant melanoma. Eur J Cancer, 2003. **39**(5): p. 666-74.
32. Tóth-Jakatics, R., et al., *Cutaneous malignant melanoma: correlation between*
neovascularization and peritumor accumulation of mast cells overexpressing vascular
endothelial growth factor. Hum Pathol, 2000. **31**(8): p. 955-60.
33. Dyduch, G., K. Okoń, and E. Pescarini, *Mast cells in melanocytic skin lesions. An*
immunohistochemical and quantitative study. Pol J Pathol, 2011. **62**(3): p. 139-44.
34. Annese, T., et al., *Autocrine/Paracrine Loop Between SCF(+)/c-Kit(+) Mast Cells*
Promotes Cutaneous Melanoma Progression. Front Immunol, 2022. **13**: p. 794974.
35. Öhrvik, H., et al., *Mast cells promote melanoma colonization of lungs.* Oncotarget,
2016. **7**(42): p. 68990-69001.
36. Benito-Martin, A., et al., *Mast cells impair melanoma cell homing and metastasis by*
inhibiting HMGA1 secretion. Immunology, 2023. **168**(2): p. 362-373.
37. Grujic, M., et al., *The combined action of mast cell chymase, tryptase and*
carboxypeptidase A3 protects against melanoma colonization of the lung. Oncotarget,
2017. **8**(15): p. 25066-25079.
38. Marichal, T., M. Tsai, and S.J. Galli, *Mast cells: potential positive and negative roles*
in tumor biology. Cancer Immunol Res, 2013. **1**(5): p. 269-79.
39. Varricchi, G., et al., *Are Mast Cells MASTers in Cancer?* Front Immunol, 2017. **8**: p.
424.
40. Oldford, S.A. and J.S. Marshall, *Mast cells as targets for immunotherapy of solid*
tumors. Mol Immunol, 2015. **63**(1): p. 113-24.
41. Betts, M.R., et al., *Sensitive and viable identification of antigen-specific CD8+ T cells*
by a flow cytometric assay for degranulation. J Immunol Methods, 2003. **281**(1-2): p.
65-78.
42. Lorenzo-Herrero, S., et al., *CD107a Degranulation Assay to Evaluate Immune Cell*
Antitumor Activity. Methods Mol Biol, 2019. **1884**: p. 119-130.

43.   Betts, M.R. and R.A. Koup, *Detection of T-Cell Degranulation: CD107a and b*, in
515   *Methods in Cell Biology*. 2004, Academic Press. p. 497-512.
516   44.   Mahata, B., et al., *Tumors induce de novo steroid biosynthesis in T cells to evade*
517   *immunity.* Nat Commun, 2020. **11**(1): p. 3588.
518   45.   Prekovic, S., et al., *Glucocorticoid receptor triggers a reversible drug-tolerant*
519   *dormancy state with acquired therapeutic vulnerabilities in lung cancer.* Nat
520   Commun, 2021. **12**(1): p. 4360.
521   46.   Poinot, H., et al., *Activation of endogenous glucocorticoids by HSD11B1 inhibits the*
522   *antitumor immune response in renal cancer.* OncoImmunology, 2024. **13**(1): p.
523   2286820.

Figure Legends

Figure. 1 Schematic illustration of the experimental design for the melanoma lung metastasis model. Control and conditional knockout (cKO) mice were intravenously injected with 0.5 × 10☐ B16-F10 melanoma cells, and assessed for pulmonary tumour nodule formation at day 13. Body weight was monitored on days 0, 7, and 13. Mice were euthanized on day 13, and total tumour nodules were counted. Next, lung tissue samples were processed for characterization of immune cells, following ex vivo stimulation with PMA.

Figure 2. Mast cell–derived Cyp11a1 promotes lung tumour colonization and is associated with increased mast cell infiltration. (**a**) Representative images of lung tumour nodules from control (Cyp11a1-sufficient) and mast cell–specific *Cyp11a1* knockout (cKO) mice following metastatic challenge. Note the greater tumour burden (black arrows) in control vs. cKO animals. (**b**) Quantification of lung metastatic nodules. (**c**) Body weight monitoring of mice during metastatic progression. (**d**) Histological (H&E) and toluidine blue staining showing tumour tissue morphology with enhanced nodule formation (red arrows) and greater mast cell infiltration (black arrows) in lung tissues from control (Cyp11a1-sufficient) mice compared to mast cell–specific *Cyp11a1* knockout mice. (**e**) Quantitative assessment of mast cells in lungs of control- and cKO mice (at least three-five high-power fields per section were counted). **(f-g)** Quantitative RT-PCR (qPCR) analysis of the expression of mast cell marker genes (*Mcpt1* and *Mcpt6*) in lungs of tumour-bearing control (Ctrl) and cKO mice. Gene expression was normalized to *GAPDH* and *Actb* using the ΔΔCt method and presented as fold change. Statistical analysis was performed using unpaired *t*-tests (For tumour nodule enumeration, n = 8; histology, n = 3; qPCR, n = 4). *P* < 0.05 was considered significant.

Figure 3. Expression of Cyp11a1 in mast cells of tumours from control mice but not in mice with mast cell-specific depletion of *Cyp11a1*. (**a**) Confocal microscopy images showing localization of the mast cell marker tryptase (red) and Cyp11a1 (green) in lung tumours from control (Ctrl; Cyp11a1-sufficient) and mast cell–specific Cyp11a1 knockout (cKO) mice. (**b**) Quantitative analysis of Cyp11a1/tryptase co-localization using ImageJ software confirming a significantly higher proportion of Cyp11a1☐ mast cells in control tumours compared to cKO tumours (**c**) Quantitative RT-PCR (qPCR) analysis of *Cyp11a1* gene expression in lungs of tumour-bearing control (Ctrl) and cKO mice. *Cyp11a1* gene expression was normalized to *Gapdh* and *Actb* expression using the ΔΔCt method and presented as fold change. Statistical analysis was performed using unpaired t-tests (For IF staining, n = 3; for qPCR, n= 4). P < 0.05 was considered significant.

Figure 4. Expression of Cyp11a1, CD45 and CD107a in lung cells from tumour-bearing control mice and mice with Cyp11a1-deficient mast cells. (**a, c**) Confocal microscopy images showing expression of the general leukocyte marker CD45 (red), the activation marker CD107a (magenta) and Cyp11a1 in lung cells isolated from control (Cyp11a1-sufficient) and mast cell–specific *Cyp11a1* knockout (cKO) tumour-bearing mice. (**b, d**) Quantitative analysis of the intensity of Cyp11a1, CD45 and CD107a staining intensity in lung tissue cells from control mice (Ctrl) and mice with mast cell-specific depletion of *Cyp11a1* (cKO), using the ImageJ software. VoF, view of field. Data represent mean values ±☐SEM; *P*☐<☐0.05 by unpaired two-tailed *t*-test (n = 3).

Figure 5. Immune profiling of lungs from tumour-bearing control mice and mice with Cyp11a1-deficient mast cells. Immunophenotyping of lung tumours by flow cytometry. Note that no significant effect of mast cell-specific depletion of *Cyp11a1* on the proportion of T cells, B cells, NK cells, eosinophils or basophils among the CD45$^+$ population were seen (gating strategy shown in **Suppl. Fig. 1**). Data represent mean values ±SEM; $P < 0.05$ by unpaired two-tailed *t*-test (Ctrl, n = 8; cKO, n = 7 - 8).

Figure 6. Enhanced immune cell activation in lungs from mice with Cyp11a1-deficient mast cells. CD8$^+$ T cells, CD4$^+$ T cells, NK cells, basophils and mast cells were assessed for expression of the general activation/degranulation marker CD107a by flow cytometry. Assessments were performed at baseline conditions (unstimulated; US) or after activation with PMA (stimulated; S). The gating strategy is shown. Results are expressed as % CD107a-positve cells out the total cells in the respective population. Data represent mean values ±SEM; $P < 0.05$ by unpaired two-tailed *t*-test (at least n = 6/group). US, unstimulated; S, stimulated with PMA.

Figure 7. Increased interferon-γ production by T cells in lungs from mice with Cyp11a1-deficient mast cells. Total dispersed lung cell populations were prepared from tumour-bearing control mice (Ctrl) and mice with *Cyp11a1*-deficient mast cells (cKO), and were activated by incubation with PMA. (**a**) After activation of cells, the expression of IFN-γ and TNFα in CD8$^+$ T cells, CD4$^+$ T cells, NK cells, B cells and mast cells was assessed by flow cytometry as indicated. (**b**) qPCR analysis of IFN-γ and TNFα gene expression in lungs from tumour-bearing control mice (Ctrl) and mice with *Cyp11a1*-deficient mast cells (cKO). Data represent mean ±SEM; $P < 0.05$ by unpaired two-tailed *t*-test (at least n = 6/group).

Suppl Fig. 1. Gating strategy used for Figs 5-7.

Suppl. Table 1: List of Primers with sequences

1. In vitro culture

B16-F10 melanoma

0.5*10^6 cells in 0.1 ml PBS

2. In vivo injection

1. Control (Cyp11a1$^{fl/fl}$)

2. cKO (Cyp11a1$^{fl/fl}$;Cpa3-Cre)

Day 0　　　　　Day 7　　　Day 13

Tumor nodules formation period

- Monitor mice health/body weight

Sacrifice

Count total nodules

Right　　Left

3. Ex vivo Stimulation

PMA

Flow cytometry IHC/IF/qPCR

- Immune cells phenotypes
- Quantification & visualization
- mRNA analysis

a

Unstimulated (US) | Stimulated (S) with PMA

INF-γ+ CD8 cells
0.0377

TNF-α+ CD8 cells
0.0761

IFN-γ+ CD4 T cells
0.0177

TNF-α+ CD4 cells
0.4350

IFN-γ+ NK cells (S)
0.2823

TNF-α+ NK cells (S)
0.2135

TNFα+ B cells (S)
0.6492

TNFα+ mast cells (S)
0.0416

b

Ifn-γ
*

Tnf-α
ns