1

Investigating the use of human COVID-19 rapid assays to detect antibody and antigen in domesticated dogs (*Canis lupus familiaris*) and cats (*Felis catus*)

Lauren C. Cybulska[1*], Sarah A. Hamer[2], Chloe Teasdale[3], Glen Johnson[1], Gabriel L. Hamer[4], Jean

Grassman[1]

[1] Department of Environmental, Occupational, and Geospatial Health Sciences, City University of New

York Graduate School of Public Health and Health Policy, 55 West 125th Street, New York, New York,

10027, United States of America

[2] Department of Veterinary Integrative Biosciences, College of Veterinary Medicine & Biomedical

Sciences, Texas A&M University, 4458 TAMU, College Station, Texas, 77843, United States of America

[3] Department of Epidemiology and Biostatistics, City University of New York Graduate School of Public

Health and Health Policy, 55 West 125th Street, New York, New York, 10027, United States of America

[4] Department of Entomology, Texas A&M University, 370 Olsen Boulevard, 2475 TAMU, College

Station, Texas, 77843, United States of America

*Corresponding author

E-mail: Lauren.Cybulska@gmail.com (LC)

Abstract

Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), the virus that causes

coronavirus disease-2019 (COVID-19) in humans, is also known to infect animals including

dogs (*Canis lupus familiaris*) and cats (*Felis catus*). This study evaluated the efficacy of human

COVID-19 rapid antigen and antibody tests in dogs and cats. Nasal/oral swabs from 60 animals

(32 dogs, 28 cats) and serum from 40 animals (20 dogs, 20 cats) were tested. Rapid antigen tests

used on respiratory swabs showed low-to-moderate sensitivity (75% dogs, 57% cats) and

moderate-to-high specificity (79% dogs, 95% cats) compared to RT-PCR. Rapid antibody tests

used on serum samples demonstrated low-to-moderate sensitivity (70% dogs, 50% cats) and

moderate-high specificity (60% dogs, 100% cats) compared to PRNT. While imperfect, these

test kits may have some utility for field surveillance studies, particularly when species-specific

rapid SARS-CoV-2 assays for dogs and cats are unavailable. These test characteristics in dogs

and cats are similar to the findings from studies of the same types of tests in humans which have

found an average sensitivity and specificity of common commercially available kits in the US

range from 50.0-84.3% and 64.5-74.3%, respectively, when used with human samples (1,2).

Introduction

Severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) is known to cause

coronavirus disease-2019 (COVID-19) in humans as well as other species including domestic

dogs (*Canis lupus familiaris*) and cats (*Felis catus*), among other animals (3). In some cases,

these animals can develop symptoms similar to those seen in human COVID-19 patients (4–8),

but most SARS-CoV-2-infected animals are thought to be asymptomatic (9–13). Evidence

suggests that pets living in homes with COVID-positive humans are more likely to test positive,

pointing to human-to-animal transmission as the likely cause (14–18). SARS-CoV-2 infections

in pets are not only a risk to animal health; coronaviruses mutate easily, which represents a

potential for the generation of novel viruses within non-human hosts that can then be transmitted

to humans (i.e., "spillover"). Some authors have stressed the importance of continued

surveillance of SARS-CoV-2 in domesticated dogs, cats, and other susceptible species, to help

mitigate the threat of zoonotic (i.e., animal-to-human) and reverse-zoonotic (human-to-animal)

transmission through early detection and response (19–26).

Similar to their use in humans (27–29), real-time polymerase chain reaction (RT-PCR),

plaque reduction neutralization testing (PRNT) (i.e., viral neutralization), and laboratory-based

enzyme-linked immunoassays (ELISAs) methods are commonly used to detect SARS-CoV-2

viral ribonucleic acid (RNA) and antibodies in dogs and cats (30–32). RT-PCR and PRNT are

considered gold-standard testing methods. However, these methods can be costly and require

specialized laboratory equipment and trained personnel making these tests inaccessible and

impractical for widespread use, particularly in low-resource settings. This highlights the

importance of identifying alternative methods of testing that are more economical and allow for

rapid diagnoses and accessible surveillance techniques.

While rapid antibody and rapid antigen tests (RATs) have been developed for SARS-

CoV-2 detection in humans and are affordable, no such similar tests have been developed for

non-human species. In the United States (US), each assay costs as little as $9.00 and they are

widely available across the country online or over the counter (33). If such tests were found to be

effective when used in non-human subjects, this would be a cost-effective and portable solution

to aid in widespread surveillance testing or the individual testing of animals at local veterinary

clinics, animal shelters, or homes. The average sensitivity and specificity of common

commercially available RATs in the US range from 50.0-84.3% and 64.5-74.3%, respectively,

when used with human swab samples (1,2).

With zoonotic diseases becoming an ever-increasing threat to global health – such as

avian influenza and Ebola (34,35) – it is important to consider alternative testing methods to aid

in widespread surveillance of these diseases across all susceptible animal populations. It is not

uncommon for researchers, veterinarians, and wildlife disease specialists to use human

diagnostic tests with animal samples. For example, human rapid influenza diagnostic tests

(RIDTs) have been used to detect swine influenza (H1N1 and H3N2 strains) in pigs in Southeast

Asia (36). To date, there is limited information about the use of human rapid tests in animals for

detecting SARS-CoV-2. Since March 2020, some authors have reported the use of human

SARS-CoV-2 ELISA and PCR assays with animal samples in laboratory settings (37–39), and

others have successfully conducted experiments using human rapid antigen and antibody tests

with dog or cat saliva, nasal swabs, and serum samples (40–43). In this study, we investigated

whether commercial rapid human assays, both antibody and antigen tests, could be used to detect

SARS-CoV-2 in dogs and cats in an off-label assessment.

Materials and methods

Study methods

We used rapid COVID-19 tests developed for humans to test dog and cat biological

samples to identify whether these assays could be used by veterinarians or public health officials

in surveillance studies. Dog and cat nasal and oral swab samples collected in viral transport

media (VTM), a solution used to preserve biological samples, were tested with commercially

available human COVID-19 RATs (Abbott BinaxNOW® Covid-19 Antigen Card Tests, Abbott

Diagnostics, Scarborough, ME, USA) (44). Dog and cat serum samples were tested with human

COVID-19 rapid antibody assays (EcoTest® COVID-19 IgM/IgG Rapid Test Device,

Hangzhou, China) (45). Sensitivity, specificity, positive predictive value (PPV), and negative

predictive value (NPV) were estimated for the rapid assays when used with animal samples. The

performance characteristics of rapid antibody and rapid antigen assays were assessed by species.

The current laboratory study was conducted in a Biosafety Level 2 (BSL-2) facility from

November 2021 to March 2022. Respiratory swabs and blood samples were collected from pet

dogs and cats living in COVID-positive households from 2020-2021 by a field team from Texas

A&M University in College Station, Texas, USA (46). Nasal swab material was stored in VTM

(CDC SOP#: DSR-052-02) (47). Oral swab material was collected from cats when nasal swab

samples could not be obtained (e.g., if the sterile swab was too large for a cat's nares).

The whole blood was spun in a centrifuge to obtain sera. Gold-standard antibody and

molecular testing were performed at Hamer Laboratory prior to the current study. Antibody titers

were measured in serum by PRNT, and the presence of viral RNA was measured by the

molecular method of RT-PCR using VTM. This involves extracting RNA and converting it to

deoxyribonucleic acid (DNA) followed by amplification by PCR. The animal sera and VTM

samples were stored at -80°C prior to the initiation of the current study; freezing and thawing

were kept at a minimum to preserve the quality of the samples (48). The investigator was blinded
to the positivity status of the VTM and serum samples.

Real-time polymerase chain reaction protocols

The viral RNA samples were extracted at the Texas laboratory as previously described
using a MagMAX CORE Nucleic Acid Purification Kit on a 96-well Kingfisher Flex System
(ThermoFisher Scientific, Waltham, MA, USA) (31). A subset of samples was tested for RNA
concentration on an Epoch Microplate Spectrophotometer (BioTek, Winooski, VT, USA).
Negative controls were included in the nucleic acid purification plates (VTM) and qPCR plates
(PCR-grade water) with no indication of contamination (31).

Plaque-reduction neutralization testing protocols

117   At the Texas laboratory, serum was tested for SARS-CoV-2 by $PRNT_{90}$ as previously
118 described using Isolate USAIL1/2020, NR 52381 (49,50). The use of an early lineage of SARS-
119 CoV-2 for the PRNT was appropriate for minimizing the occurrence of false seronegative
120 results, given findings in humans that neutralizing antibodies mounted for recently emerged
121 variants are capable of neutralizing ancestor lineages (50).

Antigen testing using human COVID-19 RAT test kits with VTM solution

The anterior nasal and oral swab samples stored in VTM solution (400 µL aliquots) were
used to detect the presence of SARS-CoV-2 nucleocapsid antigen with a commercially available
RAT (Abbott BinaxNOW® Covid-19 Antigen Card Test) (44). A curated set of pet swabs

(n=60) stored in VTM, previously subjected to RT-qPCR methods, was selected for inclusion,

including 25% (8/32) dog samples and 25% (7/28) cat samples that were RT-qPCR positive.

These samples were stored in -80 degree freezer for up to 6 months prior to preparing aliquots

for the current study. We transferred 100 µL of VTM to the RAT device in the location where

the buffer is normally administered. The buffer solution that was included in the test kit was not

used. The sterile swab included in each test kit was inserted in the test card after the VTM

solution was administered to the device; the sterile swab was rotated in the test device, and the

test card was closed. The results were read at least 15, and no more than 30, minutes after

initiating the test, per the instruction manual (44).

Antibody testing using human COVID-19 rapid antibody test kits with sera

A curated set of pet serum samples (n=40) previously subjected to PRNT methods was

selected for inclusion, were tested using EcoTest® COVID-19 IgM/IgG Rapid Test Device to

detect IgG and IgM antibodies. The set included 40-50 µL aliquots of serum that had been in

freezer storage for up to 6 months, with 50% (10/20) dog and 50% (10/20) cat samples being true

positives. The tests were used in the same manner described in the manufacturer's instructions

with the use of animal serum samples instead of human serum (45).

Sample size and calculations

The sample size used for the study was based on the available SARS-CoV-2 positive

animal samples. A sample was considered a true positive if it was positive on RT-qPCR or

PRNT. The sensitivity (proportion of all true positives that showed positive test results),

specificity (proportion of all true negatives that showed negative results), positive predictive

value (PPV; probability that a positive test result is truly positive), and negative predictive value

(NPV; probability that a negative test result is truly negative) were calculated for all test results.

Ethical considerations

152   The animal samples were collected by the Hamer Laboratory team at Texas A&M

153 University with prior approval from the Texas A&M University Institutional Animal Care and

154 Use Committee (IACUC), located in College Station, Texas, USA, Animal Use Protocol

155 #IACUC 2022-0001 CA.

Results

For dog serum samples, the rapid antibody assay was found to have 70% sensitivity and

60% specificity, the PPV was 64% and the NPV was 67% in comparison to PRNT methods

(Table 1). For cat serum samples, the rapid antibody assay was found to have 50% sensitivity

and 100% specificity, the PPV was 100% and the NPV was 40%. All the positive tests had

confirmed IgM indicator lines and there were no IgG-positive samples. Some of the IgM test

bands were very faint, while other test bands were as dark as, or darker than, the control line.

Table 1. Sensitivity, specificity, positive predictive value (PPV), and negative predictive value

(NPV) estimates using human COVID-19 rapid antibody and antigen assays for dog and cat

samples, in comparison to PRNT and RT-qPCR methods, respectively.

	Species	**Sensitivity (%)**	**Specificity (%)**	**PPV (%)**	**NPV (%)**

Antibody test	Dogs	70	60	64	67
	Cats	50	100	100	40
Antigen test	Dogs	75	79	55	90
	Cats	57	95	80	87

For dog VTM samples, the RAT was found to have 75% sensitivity and 80% specificity, the PPV was 55% and the NPV was 91%, using the qRT-PCR as the gold standard method. For cat VTM samples, the RAT was found to have 57% sensitivity and 95% specificity, the PPV was 80% and the NPV was 87%.

Discussion

This study employed human rapid antibody and antigen test kits with dog and cat serum and nasal swab samples, to estimate sensitivity and specificity compared to PRNT and RT-qPCR testing. The test characteristics we observed in dogs and cats are similar to the findings from studies of the same types of tests in humans which have found an average sensitivity and specificity of common commercially available kits in the US range from 50.0-84.3% and 64.5-74.3%, respectively (1,2). Overall, sensitivity of human antibody and antigen tests was moderate in dogs and low in cats, and the specificity was low-to-moderate in dogs and high in cats. Considering the absence of readily available species-specific rapid SARS-CoV-2 assays for dogs and cats, the test kits in the current study may still be beneficial for field surveillance studies, particularly in low-resource settings.

In our study, all positive animals had positive IgM test bands on the antibody assays

suggesting that these animals were recently infected with SARS-CoV-2; this is consistent with

the timeframe of animal sample collection occurring during the owners' acute COVID-19

infections. Previous studies found the opposite; they solely reported the detection of IgG

antibodies in animal samples which could mean their samples were collected later in the animals'

infections (51–55). The different results across studies may be due to the variations in collection

times from the point of initial exposure of an animal to an infected owner. In the current study,

the animal samples were collected while the owners were quarantined at home during an acute

COVID-19 infection, supporting the IgM-positive findings. Conversely, other studies collected

samples after an owner's COVID-19 infection had already resolved or the owner's COVID-19

history was unknown, suggesting that these animals may not have had acute or recent infections.

Consequently, the detection of IgM or IgG antibodies in the current or previous studies would

align with the animals' infection stage. –

Several factors could have led to human antigen and antibody SARS-CoV-2 tests

exhibiting low sensitivity in dogs and cats, and low specificity in dogs in this study. For

example, the test kits were not used in the manner described in the instruction manuals which

could have potentially decreased the efficacy of the tests (44,45). Firstly, the COVID-19

antibody test and RAT devices were intended for use in human subjects but instead were used

with animal biological samples during this study. Secondly, the standard COVID-19 RAT

protocols normally involve collecting a fresh respiratory swab sample (i.e., nasal or throat swab)

directly from a human subject, submerging the swab in the buffer solution provided with the test

kit, and administering the solution to the test device immediately after collection. Additionally,

the nasal and oral swab materials were not collected from animals immediately before

administering the tests, which was a major limitation in this study. In the current study, we did

not have access to fresh biological samples; instead, we used respiratory swab sample material

from animals that were previously collected in VTM, frozen, and thawed prior to administration

in the test devices. It is possible that sensitivity could have been increased if the samples were

tested immediately after collection from animal subjects.

Congruency between positive RAT and RT-PCR results in respiratory swab samples has

been described in previous studies (41,42). This suggests that further investigation employing

rapid COVID-19 assays *in vivo* in conjunction with gold standard methods may help determine

whether they would produce increased sensitivity and specificity when used with live animal

subjects. It is also possible that our storage conditions (i.e., undergoing multiple freeze-thaw

cycles) could have resulted in degradation of the SARS-CoV-2 virus and antibodies in the VTM

and serum, respectively (56–58). Cats were found to be more susceptible to SARS-CoV-2

infection in multiple studies (59–61). However, it remains unclear whether differences in viral

susceptibility between species may have played a role in the higher assay specificity in cats

compared to dogs in this study, or if other biological differences between these species

influenced viral or antibody detection using these devices. Lastly, there is subjectivity in reading

tests and it can be challenging to visually determine whether a test result is positive as the test

bands can be very faint, potentially due to low viral load and possibly due to other unknown

factors (42).

There is a need for affordable and accessible testing methods for SARS-CoV-2 in

domestic dogs and cats to integrate a One Health approach in disease surveillance. The "One

Health" concept posits that human, animal, and environmental health are interconnected, with an

emphasis on preventing diseases directly or indirectly arising between animals and humans at the

human-animal interface (62,63). Considering the proximity of domestic animals and humans, the

surveillance methods in the current study could be implemented through a One Health

framework to detect disease trends resulting from zoonotic or reverse-zoonotic transmission.

High seroprevalence (21-57%) of SARS-CoV-2 has been found among dogs and cats, with some

of the highest rates in COVID-19-positive households, providing evidence of suspected human-

to-animal transmission (14,16,18,31,37,42,64–69). It is therefore critical to monitor SARS-CoV-

2 and other coronaviruses in susceptible species (e.g., dogs and cats) to mitigate the threat of

viral evolution in non-human hosts that could potentially lead to more pathogenic strains that

could result in the next pandemic ("Disease X") (19–26).

In this study of 60 dogs and cats, we found that commonly available human rapid

COVID-19 assays could be used to test animal samples and have similar but somewhat lower

performance as compared with human samples (1,2). This suggests that commercial RAT and

antibody assays could potentially have real-world applications in veterinary settings, surveillance

investigations, or households with dogs and cats. It is important to find accessible and affordable

testing methods that would allow veterinary and public health professionals to conduct

surveillance of SARS-CoV-2 in domesticated dogs and cats because of the potential for cross-

species transmission that could negatively affect human or animal health.

Author contributions

LC conceptualized and designed the study, conducted the experiment, curated the data,

performed statistical analysis, wrote the original draft, revised the manuscript, secured funding,

and managed project logistics and coordination. SH, GH, and the Hamer Laboratory team at

Texas A&M University collected biological samples from animal subjects in the field, tested

them using gold-standard RT-PCR and PRNT methods, and provided sera and VTM samples to
the partnering investigator in New York. SH also contributed to study conceptualization, design,
and manuscript review and editing. CT and GJ assisted with data analysis and manuscript review
and editing. JG supervised the research project, contributed to study conceptualization and
design, conducted the experiment, secured funding, and participated in manuscript review and
editing.

Acknowledgments

We thank Lisa Auckland and Wendy Tang at Texas A&M University for their significant
contributions to this work, including the provision of biological samples and the conduct of RT-
PCR and PRNT analyses. We also acknowledge Dr. Catherine Machalaba, Dr. William B.
Karesh, and the staff at the City University of New York Graduate School of Public Health and
Health Policy for their mentorship and guidance during the development of this manuscript. We
are especially grateful to Dr. Xinyin Jiang for permitting us to use her laboratory facilities at
Brooklyn College.

Funding

266 This research received grant funding from the Department of Environmental, Occupational, and
267 Geospatial Health Sciences at City University of New York (CUNY) Graduate School of Public
268 Health and Health Policy (EOGHS Spring Grants 2022 and 2023), and the Research Foundation
269 (RF) of CUNY (PSC CUNY TRADB-53-259 award #65492-00 53, and PSC CUNY TRADB-
270 54-121 award #66413-00 54).

Competing interests

The author declares no competing interests.

References

1. Chu VT, Schwartz NG, Donnelly MAP, Chuey MR, Soto R, Yousaf AR, et al. Comparison of Home Antigen Testing With RT-PCR and Viral Culture During the Course of SARS-CoV-2 Infection. JAMA Intern Med. 2022 July 1;182(7):701.

2. Sugiharto VA, Gatrell SK, Chen HW, Blazek GR, Cherry AM, Schilling M. Performance Evaluation of Five Rapid At-Home COVID-19 Antigen Tests against the Omicron Variant. Jhaveri TA, editor. Microbiol Spectr. 2023 Feb 14;11(1):e02286-22.

3. Centers for Disease Control and Prevention (CDC). Animals and COVID-19 [Internet]. 2022. Available from: https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/animals.html

4. Silva Filho AP, Kmetiuk LB, Bicalho GC, Castillo AP, Soares LF, Vasconcelos P, et al. First reported cases of SARS-CoV-2 infection in owned dogs in Belo Horizonte, Brazil. Arq Bras Med Vet Zootec. 2023 June;75(3):531–4.

5. Hamdy ME, El-Deeb AH, Hagag NM, Shahein MA, Liyanage NPM, Shalaan M, et al. SARS-CoV-2 infection of companion animals in Egypt and its risk of spillover. Veterinary Medicine and Science. 2023 Jan;9(1):13–24.

6. Carpenter A, Ghai RR, Gary J, Ritter JM, Carvallo FR, Diel DG, et al. Determining the role of natural SARS-CoV-2 infection in the death of domestic pets: 10 cases (2020–2021). Journal of the American Veterinary Medical Association. 2021 Nov 1;259(9):1032–9.

7. Segalés J, Puig M, Rodon J, Avila-Nieto C, Carrillo J, Cantero G, et al. Detection of SARS-CoV-2 in a cat owned by a COVID-19−affected patient in Spain. Proc Natl Acad Sci USA. 2020;117(40):24790–3.

8. Hosie MJ, Hofmann-Lehmann R, Hartmann K, Egberink H, Truyen U, Addie DD, et al. Anthropogenic Infection of Cats during the 2020 COVID-19 Pandemic. Viruses. 2021 Jan 26;13(2):185.

9. Chiba S, Halfmann PJ, Hatta M, Maemura T, Fan S, Armbrust T, et al. Protective Immunity and Persistent Lung Sequelae in Domestic Cats after SARS-CoV-2 Infection. Emerg Infect Dis. 2021 Feb;27(2):660–3.

10. Barroso-Arévalo S, Sánchez-Morales L, Barasona JA, Rivera B, Sánchez R, Risalde MA, et
al. Evaluation of the clinical evolution and transmission of SARS-CoV-2 infection in cats by
simulating natural routes of infection. Vet Res Commun. 2022;46(3):837–52.

11. Gaudreault NN, Trujillo JD, Carossino M, Meekins DA, Morozov I, Madden DW, et al.
SARS-CoV-2 infection, disease and transmission in domestic cats. Emerging Microbes &
Infections. 2020 Jan;9(1):2322–32.

12. Jairak W, Chamsai E, Udom K, Charoenkul K, Chaiyawong S, Techakriengkrai N, et al.
SARS-CoV-2 delta variant infection in domestic dogs and cats, Thailand. Sci Rep. 2022 May
19;12(1):8403.

13. Jairak W, Charoenkul K, Chamsai E, Udom K, Chaiyawong S, Bunpapong N, et al. First
cases of SARS-CoV-2 infection in dogs and cats in Thailand. Transbounding Emerging Dis.
2022 July;69(4).

14. Michelitsch A, Allendorf V, Conraths FJ, Gethmann J, Schulz J, Wernike K, et al. SARS-
CoV-2 Infection and Clinical Signs in Cats and Dogs from Confirmed Positive Households
in Germany. Viruses. 2023 Mar 24;15(4):837.

15. Miró G, Regidor-Cerrillo J, Checa R, Diezma-Díaz C, Montoya A, García-Cantalejo J, et al.
SARS-CoV-2 Infection in One Cat and Three Dogs Living in COVID-19-Positive
Households in Madrid, Spain. Front Vet Sci. 2021 Nov 10;8:779341.

16. Bae J, Ro C, Kang Y, Ga E, Na W, Song D. Human-to-Animal Transmission of SARS-CoV-
2, South Korea, 2021. Emerg Infect Dis [Internet]. 2023 May;29(5). Available from:
https://wwwnc.cdc.gov/eid/article/29/5/22-1359_article

17. Akhtardanesh B, Jajarmi M, Shojaee M, Salajegheh Tazerji S, Khalili Mahani M, Hajipour
P, et al. Molecular screening of SARS-CoV-2 in dogs and cats from households with
infected owners diagnosed with COVID-19 during Delta and Omicron variant waves in Iran.
Veterinary Medicine and Science. 2023 Jan;9(1):82–90.

18. Fritz M, Rosolen B, Krafft E, Becquart P, Elguero E, Vratskikh O, et al. High prevalence of
SARS-CoV-2 antibodies in pets from COVID-19+ households. One Health. 2020
Dec;11:100192.

19. Barroso-Arévalo S, Díaz-Frutos M, Domínguez L, Sánchez-Vizcaíno JM. Importance of
genomic surveillance of SARS-CoV-2 in cats during reverse zoonosis events: potential viral
evolution may occur. Jones CJ, editor. Microbiol Spectr. 2023;11(5):e00680-23.

20. Molini U, Coetzee LM, Engelbrecht T, De Villiers L, De Villiers M, Mangone I, et al.
SARS-CoV-2 in Namibian Dogs. Vaccines. 2022;10(12):2134.

21. Guo R, Wolff C, Prada JM, Mughini-Gras L. When COVID-19 sits on people's laps: A
systematic review of SARS-CoV-2 infection prevalence in household dogs and cats. One
Health. 2023 June;16:100497.

22. Galhardo JA, Barbosa DS, Kmetiuk LB, De Carvalho OV, Teixeira AIP, Fonseca PLC, et al.
Molecular detection and characterization of SARS-CoV-2 in cats and dogs of positive
owners during the first COVID-19 wave in Brazil. Sci Rep. 2023 Sept 2;13(1):14418.

23. Sánchez-Montes S, Ballados-González GG, Gamboa-Prieto J, Cruz-Romero A,
Romero-Salas D, Pérez-Brígido CD, et al. No molecular evidence of SARS-CoV-2 infection
in companion animals from Veracruz, Mexico. Transbounding Emerging Dis. 2022
July;69(4):2398–403.

24. Barroso-Arévalo S, Barneto A, Ramos ÁM, Rivera B, Sánchez R, Sánchez-Morales L, et al.
Large-scale study on virological and serological prevalence of SARS-CoV-2 in cats and dogs
in Spain. Transbounding Emerging Dis [Internet]. 2022;69(4). Available from:
https://onlinelibrary.wiley.com/doi/10.1111/tbed.14366

25. Fernández-Bastit L, Marfil S, Pradenas E, Valle R, Roca N, Rodon J, et al. Severe acute
respiratory syndrome coronavirus 2 (SARS-CoV-2) infection and humoral responses against
different variants of concern in domestic pet animals and stray cats from North-Eastern
Spain. Transbounding Emerging Dis. 2022;69(6):3518–29.

26. Amman BR, Cossaboom CM, Wendling NM, Harvey RR, Rettler H, Taylor D, et al. GPS
Tracking of Free-Roaming Cats (Felis catus) on SARS-CoV-2-Infected Mink Farms in Utah.
Viruses. 2022 Sept 27;14(10):2131.

27. Horbach IS, De Souza Azevedo A, Schwarcz WD, Alves NDS, De Moura Dias B, Setatino
BP, et al. Plaque Reduction Neutralization Test (PRNT) Accuracy in Evaluating Humoral
Immune Response to SARS-CoV-2. Diseases. 2024 Jan 18;12(1):29.

28. Fereidouni A, Safari H, Rezapoor H, Mahmoudzadeh S, Fereidouni M. Evaluation of SARS-
CoV-2 Specific Antibodies in Recovered Patients by Different ELISA Kits. Iranian Journal
of Immunology. 2023 July;

29. Rahbari R, Moradi N, Abdi M. rRT-PCR for SARS-CoV-2: Analytical considerations.
Clinica Chimica Acta. 2021 May;516:1–7.

30. Barroso R, Vieira-Pires A, Antunes A, Fidalgo-Carvalho I. Susceptibility of Pets to SARS-
CoV-2 Infection: Lessons from a Seroepidemiologic Survey of Cats and Dogs in Portugal.
Microorganisms. 2022;10(2):345.

31. Hamer SA, Pauvolid-Corrêa A, Zecca IB, Davila E, Auckland LD, Roundy CM, et al.
SARS-CoV-2 Infections and Viral Isolations among Serially Tested Cats and Dogs in
Households with Infected Owners in Texas, USA. Viruses. 2021;13(5):938.

32. Calvet GA, Pereira SA, Ogrzewalska M, Pauvolid-Corrêa A, Resende PC, Tassinari WDS, et
al. Investigation of SARS-CoV-2 infection in dogs and cats of humans diagnosed with
COVID-19 in Rio de Janeiro, Brazil. Hemida MG, editor. PLoS ONE. 2021 Apr
28;16(4):e0250853.

33. CVS Pharmacy. At-Home COVID-19 Tests. 2024; Available from:
https://www.cvs.com/shop/home-health-care/home-tests/at-home-covid-19-tests

34. Jahid MJ, Nolting JM. Dynamics of a Panzootic: Genomic Insights, Host Range, and
Epidemiology of the Highly Pathogenic Avian Influenza A(H5N1) Clade 2.3.4.4b in the
United States. Viruses. 2025 Feb 25;17(3):312.

35. Naa Momo Lokko C, Obodai Lokko F. Epidemiology of human Ebola virus disease outbreak
in Sub-Saharan Africa. International Journal of General Medicine and Pharmacy (IJGMP).
2025 Jan 1;14(1):37–58.

36. Saipinta D, Chuammitri P, Ohkura S, Luangwaranan B, Tadee P. Human rapid influenza
diagnostic test: efficacy and significance for using in animal. Veterinary Integrative
Sciences. 2023 Feb 20;21(2):265–71.

37. Sirakov I, Rusenova N, Rusenov A, Gergova R, Strateva T. Human ELISA Detects anti-
SARS-CoV-2 Antibodies in Cats: Seroprevalence and Risk Factors for Virus Spread in
Domestic and Stray Cats in Bulgaria. Veterinary Sciences. 2023 Jan 6;10(1):42.

38. Yamayoshi S, Ito M, Iwatsuki-Horimoto K, Yasuhara A, Okuda M, Hamabata T, et al.
Seroprevalence of SARS-CoV-2 antibodies in dogs and cats during the early and mid-
pandemic periods in Japan. One Health. 2023 Dec;17:100588.

39. Zambrano-Mila MS, Freire-Paspuel B, Orlando SA, Garcia-Bereguiain MA. SARS-CoV-2
infection in free roaming dogs from the Amazonian jungle. One Health. 2022;14:100387.

40. Santos HO, Santos EMS, Da Silva De Oliveira H, Dos Santos WS, Tupy AA, Souza EG, et
al. Screening for canine coronavirus, canine influenza virus, and severe acute respiratory
syndrome coronavirus 2 in dogs during the coronavirus disease-2019 pandemic. Vet World.
2023 Sept;1772–80.

41. Gosling M, Bacon J. Acute-onset respiratory signs in a Labrador Retriever with a positive
SARS-CoV-2 rapid antigen test and infection confirmed by RT-PCR analysis: a case report.
J Vet Sci. 2022;23(6):e80.

42. Hagag IT, Weber S, Sadeghi B, Groschup MH, Keller M. Impact of animal saliva on the
performance of rapid antigen tests for detection of SARS-CoV-2 (wildtype and variants
B.1.1.7 and B.1.351). Veterinary Microbiology. 2021;262:109243.

43. Shorunke FO, Okolocha EC, Kia GS, Usman A, Akano O, Awosanya EJ. Prevalence and
risk factors associated with SARS-CoV-2 infections among veterinary practitioners and dogs
patients, June-August 2020, Lagos, Nigeria. One Health Outlook. 2022 Dec;4(1):4.

44. BTTN USA. Abbott BinaxNOW Covid-19 Antigen Card Test (POC) [Internet]. 2022.
Available from: https://www.globalpointofcare.abbott/us/en/lp/binaxnow-self-test/covid-19-
antigen.html

45. Code 1 Supply. 15-Minute Rapid IgG IgM COVID Antibody Test Kits [Internet]. 2021.
Available from: https://www.medline.com/product/Ecotest-COVID19-IgG/IgM-Rapid-Test-
Control/Z05-PF269810

46. Pauvolid-Corrêa A, Davila E, Auckland LD, Zecca IB, Busselman RE, Tang W, et al. Active
surveillance of cats and dogs from households with human COVID-19 cases reveals over
one quarter of pets infected with SARS-CoV-2 in 2020-2021 in Texas, United States
[Internet]. 2025. Available from: http://biorxiv.org/lookup/doi/10.1101/2025.02.03.636361

47. Centers for Disease Control and Prevention (CDC). Preparation of Viral Transport Medium.
Standard Operating Procedure SOP #: DSR-052-02 (Archived). 2020; Available from:
https://archive.cdc.gov/#/details?q=preparation%20of%20viral%20transport%20medium&st
art=0&rows=10&url=https://www.cdc.gov/coronavirus/2019-ncov/downloads/Viral-
Transport-Medium.pdf

48. World Health Organization (WHO). Surveillance Guidelines for Measles, Rubella and
Congenital Rubella Syndrome in the WHO European Region. Annex 3 Collection, storage
and shipment of specimens for laboratory diagnosis and interpretation of results [Internet].
2012. Available from: https://www.ncbi.nlm.nih.gov/books/NBK143256/

49. Beaty B, Calisher C, Shope R. Arboviruses. Viral, Rickettsial, and Chlamydial infections.
Washington, DC: American Public Health Association; 1995. 189–212 p.

50. Hamer SA, Nunez C, Roundy CM, Tang W, Thomas L, Richison J, et al. Persistence of
SARS-CoV-2 neutralizing antibodies longer than 13 months in naturally infected, captive
white-tailed deer (*Odocoileus virginianus*), Texas. Emerging Microbes & Infections.
2022;11(1):2112–5.

51. Dileepan M, Di D, Huang Q, Ahmed S, Heinrich D, Ly H, et al. Seroprevalence of SARS-
CoV-2 (COVID-19) exposure in pet cats and dogs in Minnesota, USA. Virulence. 2021 Dec
31;12(1):1597–609.

52. Jaramillo Hernández DA, Chacón MC, Velásquez MA, Vásquez-Trujillo A, Sánchez AP,
Salazar Garces LF, et al. Seroprevalence of exposure to SARS-CoV-2 in domestic dogs and
cats and its relationship with COVID-19 cases in the city of Villavicencio, Colombia.
F1000Res. 2023;11:1184.

53. Udom K, Jairak W, Chamsai E, Charoenkul K, Boonyapisitsopa S, Bunpapong N, et al.
Serological survey of antibodies against SARS-CoV-2 in dogs and cats, Thailand.
Transbounding Emerging Dis. 2022 July;69(4):2140–7.

54. Kleinerman G, Gross S, Topol S, Ariel E, Volokh G, Melloul S, et al. Low serological rate of
SARS-CoV-2 in cats from military bases in Israel. Comparative Immunology, Microbiology
and Infectious Diseases. 2022 Nov;90–91:101905.

55. Imanishi I, Asahina R, Hayashi S, Uchiyama J, Hisasue M, Yamasaki M, et al. Guest edited
collection serological study of SARS-CoV-2 antibodies in japanese cats using protein-A/G-
based ELISA. BMC Vet Res. 2022 Dec 21;18(1):443.

56. Takahashi H, Ichinose N, Okada Y. False-negative rate of SARS-CoV-2 RT-PCR tests and
its relationship to test timing and illness severity: A case series. IDCases. 2022;28:e01496.

57. Yang Y, Yang M, Yuan J, Wang F, Wang Z, Li J, et al. Laboratory Diagnosis and
Monitoring the Viral Shedding of SARS-CoV-2 Infection. The Innovation.
2020;1(3):100061.

58. Cardona S, Eck A, Cassellas M, Gallart M, Alastrue C, Dore J, et al. Storage conditions of
intestinal microbiota matter in metagenomic analysis. BMC Microbiol. 2012;12(1):158.

59. Islam A, Ferdous J, Islam S, Sayeed MdA, Rahman MdK, Saha O, et al. Transmission
dynamics and susceptibility patterns of SARS-CoV-2 in domestic, farmed and wild animals:
Sustainable One Health surveillance for conservation and public health to prevent future
epidemics and pandemics. Transbounding Emerging Dis. 2022;69(5):2523–43.

60. Stanojevic S, Radojicic S, Misic D, Srejić D, Vasiljevic DV, Prokic K, et al. Frequency of
SARS-CoV-2 infection in dogs and cats: Results of a retrospective serological survey in
Šumadija District, Serbia. Preventive Veterinary Medicine. 2022;208:105755.

61. Cardoso NP, Rivero C, Castillo M, Mansilla FC, Pastorino F, Piccirilli G, et al. Serological
screening of SARS-CoV-2 infection in companion animals of Buenos Aires suburbs. Front
Vet Sci. 2023;10:1161820.

62. World Health Organization. Initiatives: One Health [Internet]. 2023. Available from:
https://www.who.int/europe/initiatives/one-
health#:~:text=One%20Health%20is%20an%20approach,animal%2Dhuman%2Denvironme
nt%20interface

63. PREDICT Consortium, Saylors K, Wolking DJ, Hagan E, Martinez S, Francisco L, et al.
Socializing One Health: an innovative strategy to investigate social and behavioral risks of
emerging viral threats. One Health Outlook. 2021;3(1):11.

64. Pandit R, Matthews QL. A SARS-CoV-2: Companion Animal Transmission and Variants
Classification. Pathogens. 2023 May 29;12(6):775.

65. Zhang Q, Zhang H, Gao J, Huang K, Yang Y, Hui X, et al. A serological survey of SARS-
CoV-2 in cat in Wuhan. Emerging Microbes & Infections. 2020 Jan;9(1):2013–9.

66. Stevanovic V, Tabain I, Vilibic-Cavlek T, Mauric Maljkovic M, Benvin I, Hruskar Z, et al.
The Emergence of SARS-CoV-2 within the Dog Population in Croatia: Host Factors and
Clinical Outcome. Viruses. 2021 July 22;13(8):1430.

67. Zhang Y, You L, Zou Y, He X, Wu S, Yang F, et al. Epidemiology and Molecular
Characterizations of Coronavirus from Companion Animals Living in Chengdu, Southwest
China. Hernandez Lopez J, editor. Transboundary and Emerging Diseases. 2023 Mar
30;2023:1–9.

68. Jairak W, Charoenkul K, Chamsai E, Udom K, Chaiyawong S, Hangsawek A, et al. Survey
of SARS-CoV-2 in dogs and cats in high-risk areas during the second wave of COVID-19
outbreak, Thailand. Zoonoses and Public Health. 2022 Sept;69(6):737–45.

69. Hamer SA, Ghai RR, Zecca IB, Auckland LD, Roundy CM, Davila E, et al. SARS-CoV-2
B.1.1.7 variant of concern detected in a pet dog and cat after exposure to a person with
COVID-19, USA. Transbounding Emerging Dis. 2022;69(3):1656–8.

Supporting information

S1 Table. Comparative diagnostic performance of antibody and antigen testing in canines and felines. This table summarizes the sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) for both antibody-based and antigen-based diagnostic assays across dog and cat populations.

S2 Dataset. Diagnostic accuracy of serum testing in veterinary samples. Summary of diagnostic performance metrics. This file contains the contingency table values (TP, FP, FN, TN) and calculated performance metrics including prevalence, sensitivity, specificity, and predictive values for all samples combined, as well as canine and feline subgroups.

S3 Dataset. Diagnostic accuracy of VTM testing in veterinary samples. This table presents the diagnostic performance metrics for VTM-based testing. It includes raw counts for True Positives (TP), False Positives (FP), False Negatives (FN), and True Negatives (TN), alongside calculated values for prevalence, sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) for all samples and species-specific subgroups.