Multi-omic responses to acute exercise in abdominal subcutaneous adipose tissue of sedentary adults: findings from MoTrPAC

Cheehoon Ahn[1]*, Christopher A. Jin[2]*, Katie L. Whytock[1], Gina M. Many[3], Tyler J. Sagendorf[3], James A. Sanford[3], Zhenxin Hou[4], Mark R. Viggars[5], Jia Nie[6], Sara E. Espinoza[6], Nicolas Musi[6], Yifei Sun[7], Maria F. Pino[1], Patrick Hart[8], Dan H. Katz[9], Hasmik Keshishian[8], Gregory R. Smith[7], Scott Trappe[10], Natalie M. Clark[8], Sue C. Bodine[11], Laurie J. Goodyear[12], Karyn A. Esser[5], Christopher B. Newgard[13], Bryan C. Bergman[14], Joshua N. Adkins[3]**, Martin J. Walsh[7]**, Lauren M. Sparks[1]**, The MoTrPAC Group

*Co-first **Co-corresponding

[1]Translational Research Institute, AdventHealth, Orlando, FL, USA
[2]Department of Genetics, School of Medicine, Stanford University, Stanford, California, USA
[3]Biological Sciences Division, Pacific Northwest National Laboratory, Richland, WA, USA
[4]Department of Biochemistry, Emory University School of Medicine, Atlanta, GA, USA
[5]Department of Physiology and Aging, University of Florida, Gainesville, FL, USA
[6]Department of Medicine, Cedars-Sinai Medical Center, Los Angeles, CA, USA
[7]Icahn School of Medicine at Mount Sinai, New York, NY, USA
[8]Proteomics Platform, Broad Institute of MIT and Harvard, Cambridge, MA, USA
[9]Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA
[10]Human Performance Laboratory, Ball State University, Muncie, IN, USA
[11]Aging and Metabolism Research Program, Oklahoma Medical Research Foundation, Oklahoma City, OK, USA
[12]Section on Integrative Physiology and Metabolism, Joslin Diabetes Center, Harvard Medical School, Boston, MA, USA
[13]Duke Molecular Physiology Institute and Sarah W. Stedman Nutrition and Metabolism Center, Duke University Medical Center, Durham, NC, USA
[14]Division of Endocrinology, Diabetes, and Metabolism, University of Colorado Anschutz Medical Campus, Aurora, CO, USA

SUMMARY

Exercise induces widespread health benefits across multiple tissues, yet the acute molecular responses in human adipose tissue remain poorly defined. The Molecular Transducers of Physical Activity Consortium (MoTrPAC) profiled temporal molecular changes in abdominal subcutaneous adipose tissue (ASAT) following a single bout of exercise. Healthy sedentary adults were randomized to endurance (EE), resistance (RE), or control (CON) groups. ASAT biopsies were collected pre-exercise and at 45min, 4hr, and 24hr post-exercise, followed by transcriptomic, proteomic, phosphoproteomic, and metabolomic analyses. EE and RE elicited distinct, time-resolved molecular programs involving angiogenesis, extracellular matrix remodeling, mitochondrial metabolism, substrate utilization, and circadian regulation. Phosphoproteomics revealed acute changes in cytoskeletal and branched-chain amino acid metabolism proteins associated with glycemic control. Temporal metabolomic shifts were cell-type-specific. Finally, we identified candidate adipose-derived exerkines with predicted

42 endocrine actions. This multi-omic map of acute ASAT responses offers insight into adipose-
43 specific mechanisms by which exercise promotes metabolic health.

INTRODUCTION

The cardiometabolic health benefits of regular exercise are well-established[1,2], yet the molecular
mechanisms that drive these effects are incompletely understood. Many of these benefits—such
as enhanced whole-body and tissue-specific insulin sensitivity and reduced blood pressure[3-5]—
are initiated by single bouts of exercise, even in individuals who exercise regularly[6-8]. It is widely
accepted that the cumulative effects of these repeated, episodic molecular responses to acute
exercise bouts underlie the long-term health benefits of training[9]. As such, mapping the
molecular landscape of acute exercise responses offers a promising strategy to uncover 'how'
exercise promotes cardiometabolic health.

While much of the current and historical focus has been on skeletal muscle and the
cardiovascular system, adipose tissue is emerging as a key target of exercise adaptations.
Abdominal subcutaneous adipose tissue (ASAT) responds dynamically to both acute exercise[10-
12] and chronic training[10,13-17], contributing to improved cardiometabolic outcomes. Moreover,
similar to other tissues, ASAT may also mediate the systemic effects of exercise[18] by serving as
a rich source of numerous secreted factors—often referred to as exerkines—that act across
tissues[19]. However, the physiological adaptations and molecular-level responses in human
ASAT, particularly in the context of temporal and modality-specific dynamics of acute exercise,
remains a significant gap.

The Molecular Transducers of Physical Activity Consortium (MoTrPAC) was launched to
address this knowledge gap by systematically mapping the molecular effects of physical activity.
In this study, which leveraged pre-COVID MoTrPAC human cohort data, we profiled temporal,
multi-omic responses in ASAT following a single bout of either endurance exercise (EE) or
resistance exercise (RE)—both of which represent clinically relevant exercise prescriptions at
an unprecedented scale. Here, we reveal temporal-, modality-, and cell-type-specific molecular
responses in the ASAT transcriptome, proteome, phosphoproteome, and metabolome following
acute exercise. We examine the clinical relevance of these molecular signatures and, through *in
silico* modeling, explore the potential endocrine role of ASAT-derived exerkines. Together, our
findings provide new insights into the mechanisms underlying early ASAT-specific molecular
responses to exercise and identify candidate mediators of long-term metabolic benefits.

RESULTS

Participant characteristics and multi-omic sex differences

173 healthy, sedentary individuals were enrolled and randomized into one of three groups: an
endurance exercise group (EE, N=63), a resistance exercise group (RE, N=73), or a non-
exercise control group (CON, N=37) (Figure 1A). The cohort was predominantly female (72%),
with 46 women in EE, 49 in RE, and 31 in CON, with an average age of 41±15 years and a BMI
of 27.1±4.0 kg/m² (Table 1). Participants in EE completed a 40-minute bout of endurance
exercise at 65% VO2peak, while those in RE performed an 8-exercise resistance circuit at
volitional fatigue (~10RM), and the CON group rested in the supine position for 40 minutes
(Figure 1A). Abdominal subcutaneous adipose tissue (ASAT) biopsy samples were collected
before and at post 45min, 4hr, and 24hr (Figure 1A). To mitigate the confounding effects that
may stem from a prior biopsy on the same side (e.g., biopsy-induced inflammatory response),
participants were randomized into either "Early" (n=43; Pre-Exercise and post 45min), "Middle"
(n=80; Pre-Exercise and post 4hr), or "Late" (n=50; Pre-Exercise and post 24hr) groups that
underwent a total of two ASAT biopsies (one on either side of the abdomen) that included a pre-
exercise (all groups) and one post-exercise timepoint (Figure 1A). Collected ASAT samples
were subjected to four omic analyses: transcriptomics, global proteomics, phosphoproteomics,
and metabolomics (Table S1).

No pre-exercise differences were found in participant characteristics across the three groups
except HOMA-IR (Table 1). We observed relationships between some cardiometabolic health
traits and circulating factors. As expected, adiposity measurements such as BMI and waist
circumference (WC) were positively associated with higher blood pressure, whole-body and
adipose-specific insulin resistance indices (HOMA-IR and Adipo-IR, respectively), and HbA1c
(p<0.01; Figure S1A). WC was also positively associated with circulating branched-chain amino
acids (BCAAs; isoleucine, leucine, and valine), which are known to be elevated in humans and
rodents with insulin resistance and obesity[20] (Figure S1A). More detailed clinical characteristics
are available here (*REF: MoTrPAC Molecular Landscape, in review*).

To explore the relationship between baseline clinical characteristics and ASAT cell types, we
performed a deconvolution analysis—estimating ASAT cell type proportions using a previously
published ASAT single-nuclei (sn) RNAseq dataset and computational pipeline[21,22] (Figure 1B,
see 'Deconvolution' in Methods). As expected, the estimated proportion of macrophages was
strongly correlated with higher adiposity, insulin resistance indices, and HbA1c, supporting the
established link between ASAT macrophages and obesity and insulin resistance[23,24] (p<0.001;
Figure 1C). In contrast, the estimated proportions of adipocytes and vascular cells were
negatively correlated with adiposity and insulin resistance indices, highlighting the complex role
of ASAT cellular heterogeneity in cardiometabolic health traits (p<0.01; Figure 1C).

A significant degree of sexual dimorphism was observed across all ASAT omic layers measured
at pre-exercise: 6150 transcripts (3039 up and 3111 down in males relative to females), 755
proteins (423 up and 332 down), 2133 phosphosites (1041 up and 1092 down), and 143
metabolites (36 up and 107 down) were significantly different between sexes (adjusted p<0.05;
Figure 1D). To gain insight into sex-specific biological processes, we performed gene set

enrichment analysis using CAMERA-PR[25]. At the transcriptomic level, females showed
enrichment of lipid metabolism, branched-chain amino acid metabolism, and oxidative
phosphorylation pathways, whereas males exhibited upregulation of protein translation and
PDGFRβ signaling (adjusted p < 0.05; Figure 1E). Proteomic analysis revealed higher oxidative
phosphorylation in females, while males displayed upregulation of pathways involved in actin
cytoskeleton remodeling (e.g., actin filament, stress fiber, and myosin) and SUMOylation (Figure
1E). Kinase–substrate enrichment analysis (KSEA) of phosphoproteomic data identified sex-
specific kinase activity, with all differentially enriched kinases upregulated in females. These
included MAPK pathway kinases (e.g., PRKCH, RAF1, MAPK9, MAPK3, MAP3K8) and AKT
signaling components (e.g., PDPK, GSK3A, GSK3B), both of which are known to strongly
influence adipose cell proliferation, differentiation, and insulin signaling[26,27] (Figure 1F).
Metabolomic analysis showed higher abundances of phosphatidylethanolamines (PE) in
females and increased saturated fatty acids (FA) in males (Figure 1G). Deconvolution analysis
of the transcriptomics dataset suggested that the estimated proportion of Adipocyte-1, a subtype
of mature adipocyte, previously characterized as 'anti-oxidative' with upregulation of genes
related to anti-oxidative activity (*GPX1 & GPX4*)[21], was higher in females compared to males
(p=0.0007, Figure S1B). Conversely, the estimated proportion of vascular cells was lower in
females (p=0.0003, Figure S1B), indicating intrinsic cellular heterogeneity by sex.

Differentially expressed features following acute exercise

Differential analysis (DA) was performed across omic layers to investigate acute exercise
effects. Importantly, we accounted for temporal changes observed in the non-exercising control
(CON) group—where ASAT showed the most changes over time compared with muscle and
blood even without exercise (*REF: MoTrPAC Molecular Landscape, in review*)—to better isolate
the effects of acute endurance (EE) and resistance exercise (RE) (see 'Differential Analysis' in
Methods). For example, in the CON group, 2801 transcripts (1306 upregulated, 1495
downregulated; Figure S2A) were differentially expressed at post 4hr, including core circadian
clock genes *CLOCK*, *NR1D1/2,* and *PER1/2* (Figure S2B), despite the absence of any exercise
stimulus. These genes have previously been reported as responsive to acute aerobic exercise
in ASAT[10,12] (Figure S2C). However, those earlier studies did not include non-exercising control
groups, raising the possibility that the observed transcriptomic changes may have been at least
partially driven by non-exercise factors, such as time of day, fasting, or biopsy-induced
inflammation. Notably, temporal changes in these and other key clock genes like *CLOCK* were
not differentially regulated across CON, EE, or RE (Figure S2B). Controlling for CON yielded
markedly different outcomes across omic layers (Figure 2A), compared to differential analysis in
one exercise group disregarding controls (Figure S2A). The number of differentially regulated
transcripts peaked at post 45min (EE: n=57, RE: n=34), declined by post 4hr (EE: n=3, RE: 31),
with these changes almost entirely returning to baseline by post 24hr (EE: n=3, RE: n=0),
suggesting that transcriptional responses to acute exercise are largely transient (Figure 2A).
Acute exercise triggered changes in proteins (EE: n=305, RE: n=2) and phosphosites (EE:
n=280, RE: n=43) at post 4hr (adjusted p<0.05; Figure 2A). Differential metabolomic responses
in ASAT also showed distinct temporal- and modality-specific patterns. Notably, the majority of
differentially regulated metabolites were observed at post 4hr following EE (EE: n=93, RE: n=0)
(adjusted p<0.05; Figure 2A). Principal component analysis (PCA) of $\log_2$fold changes further

supported these distinct temporal and modality-dependent signatures in the transcriptome and
metabolome (Figure 2B).

Controlling for CON enabled more accurate pathway enrichment interpretation. Several
pathways were upregulated at 45min post-EE in both adjusted and unadjusted models (e.g.,
'mRNA processing', 'VEGF signaling', 'BCAA catabolism', and certain lipid classes like FA and
dicarboxylic acids; Figure 2C, purple). However, CAMERA-PR applied to CON-adjusted DA
revealed effects uniquely attributable to acute exercise when accounting for controls (Figure 2C,
pink). For example, a reduction in neutrophil migration at 45 min post-EE was only evident after
adjustment to the control group, which independently exhibited an increase in neutrophil
migration thus masking the effect in the EE group without adjustment (Figure 2C). This
distinction was even more pronounced at 4hr, where the directionality and magnitude of many
enriched pathways were altered after CON adjustment. Notably, OXPHOS appeared
downregulated at post 4hr in RE prior to CON adjustment, but this was primarily driven by
stronger downregulation in the CON group; after adjustment, RE showed a relative upregulation
of OXPHOS (Figure 2C). CON-adjusted enrichment of various lipid classes (e.g., TG, SM, CER,
O-PC, O-PE) became evident post 4hr in both EE and RE (Figure 2C).

Focusing on CON-adjusted DA, upregulation of pathways related to angiogenesis, ECM
remodeling, cell growth/differentiation, substrate utilization, and mitochondrial respiration at
45min post-EE, followed by protein translation at 4hr (adjusted p<0.05; Figure S2D). In RE,
pathway enrichment predominantly emerged at 4hr and included mRNA processing, translation,
ribosome biogenesis, substrate utilization, and mitochondrial respiration. By 24hr, most
transcriptomic pathways were no longer significant, although some remained suppressed
(translation in EE; mitochondrial biogenesis in RE). Proteomic analysis revealed suppression of
substrate utilization and mitochondrial respiration pathways at 4hr in both EE and RE. In the
metabolome, upregulation of FAs, N-acylethanolamines (NAEs), and dicarboxylic acids at
45min post-EE (adjusted p<0.05; Figure S2F). At 4hr, EE induced upregulation of sphingolipids
(CER, SM) and glycerophospholipids (PC, LPC, O-PC, PE, O-PE), while suppressing acyl-CoAs
and TGs. By 24hr, EE continued to alter lipid classes (LPC, PC, TG) and pyrimidine rNDPs
(Figure S2F). RE upregulated glycerophospholipids at 45min, with some (e.g., O-PC, LPC)
remaining elevated at 4hr and 24hr. Other lipid species (CER, LPC, DG, acyl-CoAs) were
altered at 24hr post-RE, suggesting a prolonged metabolomic response.

PTM Signature Enrichment Analysis (PTMSEA) identified kinase activity differences at 4hr post-
exercise (Figure 2D). Canonical MAPK pathway kinases ERK1/2 and P38, known to regulate
adipogenesis and adipocyte structure[26], were positively enriched in both EE and RE (adjusted
p<0.1). Additional kinases enriched in EE or RE included those involved in cytoskeletal
remodeling (e.g., PAK1/3, ROCK1, PKCζ, DYRK2, MARK2, CAMK2A/D, AURKC, LYN[28-31]).
These kinase signatures occurred with enriched cytoskeletal pathways from overrepresentation
analysis (ORA) of collapsed phosphoproteins (Figure S2G) and differentially expressed
phosphosites mapping to processes such as 'Cell junction assembly', 'Cell-matrix adhesion',
'Cell motility', and 'Cytoskeleton organization' (Figure 2E). Several phosphosites in Tensin-1
(TNS1) were differentially phosphorylated or showed trends toward significance (Figure S2H).
TNS1 is a scaffold protein anchoring actin filaments to integrins and mediates signal

transduction between the ECM and cytoskeleton[32]. Notably, TNS1 has been reported as a
downstream target of ROCK1[33], and their protein abundances were significantly correlated in
our data (p<0.001; Figure S2I), supporting the link between ROCK1 and TNS1 activation.

Modality-specific multi-omic responses following acute exercise

Given the limited overlap in differentially expressed features between DA completed on EE and
RE (Figure S3), we directly compared their effects on ASAT omic profiles to identify modality-
specific responses (Figure 3A). Despite differences in exercise type, duration, and energy
expenditure, the abundance of differentially expressed transcripts was minimal. Only one
transcript, NR4A1 (also known as Nur77) — transcription activator that is known to regulate
systemic glucose metabolism [34] and adipocyte differentiation [35]— showed significantly greater
upregulation in EE vs. RE at post 45min, with no significant differences at later time points
(adjusted p<0.05; Figure 3B, C). The sphingomyelin species SM 42:2;O2 exhibited divergent
responses at 45 minutes post-exercise, decreasing in EE but remaining unchanged in RE
(Figure 3B, C). The reduced phosphorylation of NDRG1-S336 was more pronounced by RE at
post 4hr. NDRG1 is required for the activation of a key regulator of adipogenesis, CEBPα, and
its phosphorylation promotes adipocyte formation[36]. However, the functional role of the specific
S336 phosphosite remains largely unexplored in the current literature. Meanwhile, only EE led
to a robust increase in SLC44A1 (a choline transporter) protein abundance, which may explain
the elevated PC species seen at post 4hr (Figure 2A).

Compared with RE, EE triggered greater early transcriptional enrichment of substrate
metabolism, BCAA catabolism, ribosome biogenesis, oxidative phosphorylation, and MAPK
signaling at post 45min, followed by immune cell activation and inflammatory responses at post
4 hr (adjusted p<0.05; Figure 3D). In contrast, RE induced greater enrichment in ECM signaling
and remodeling pathways in post 45min, followed by Slits and Robos signaling (which has been
partially linked to sympathetic activity and energetics in adipose[37]). Additionally, oxidative
phosphorylation was induced at post 4hr with RE, and immunoglobulin complex at post 24 hr
compared with EE (adjusted p<0.05; Figure 3D). The early transcriptional pathways enriched in
EE vs. RE—particularly related to mitochondrial, ribosomal, and cytoskeletal structures (Figure
3D), were reflected in proteome-level enrichment at post 4hr (adjusted p<0.05, Figure 3E). RE,
on the other hand, showed proteomic enrichment of the exosome complex, ribosome
biogenesis, and spliceosome-related pathway at post 4 hr (adjusted p<0.05, Figure 3E). In
metabolomics, EE induced greater alterations in amino acids, DG, ceramides, and fatty acids at
post 45min, which then shifted toward changes in PE, SM, O-PE, and ceramide at post 4hr
(adjusted p<0.05; Figure 3F). By post 24hr, amino acids, DG, and PC remained more
significantly altered by EE vs. RE (adjusted p<0.05, Figure 3F). Conversely, RE induced greater
changes in PC, O-PC, PE, LPC, SM, and O-PE at post 45 min compared with EE, followed by
altered saturated FA and phenolic acids at post 4hr and pyrimidine rNDP at post 24 hr (adjusted
p<0.05, Figure 3F). A notable EE-specific response was the increase in canonical ceramide
species (C14:0–C24:0) at post 4hr (Figure 3G), aligning with upregulation of inflammatory
pathways (Figure 3D). Acute inflammation is known to cause tissue ceramide accumulation
through TNFa and TLR4 activation of ceramide synthesis[38]. However, ASAT ceramides did not
remain elevated 24 hours after exercise, likely due to increased clearance, which would parallel

results from acute exercise in skeletal muscle[39]. Enhanced clearance of ceramide post-exercise
may perhaps result in lower ASAT ceramide content, which is associated with favorable
cardiometabolic health[38]. Overall, these findings highlight the unique transcriptomic, proteomic,
phosphoproteomic, and metabolomic responses induced by EE compared with RE, which may
underlie modality-specific adaptations in ASAT by exercise training.

Acute Exercise Induces Cell Type-Specific Transcriptional and Post-Translational Responses in
ASAT

Given the heterogeneous relationship between ASAT cell types and cardiometabolic health
traits observed in our deconvolution analysis (Figure 1C), we hypothesized that acute exercise
would elicit distinct transcriptional responses across ASAT cell types. To test this, we curated a
list of ASAT cell-type marker genes by integrating two of our previously published ASAT
snRNAseq datasets[21,40], capturing nine major cell types (see 'Cell Type Analysis' in Methods,
see cell markers in Table S2). Cell type enrichment analysis of transcriptomic data using
CAMERA-PR revealed modality- and time-dependent transcriptional activation across ASAT cell
types (Figure 4A). Notably, at post 45 min, marker genes for Adipocyte-2—previously described
as an insulin-responsive adipocyte subtype due to its upregulation of genes involved in lipolysis
suppression (PDE3B), lipid metabolism (ABCA5), and insulin signaling[21]—were significantly
upregulated in both training modalities (Figure 4A). In contrast, enrichment of Lipid-Associated
Macrophages (LAM) was downregulated at post 45min in both modalities, suggesting early
inhibition of pro-inflammatory macrophage activation following acute exercise. However, this
suppression was reversed at post 4hr and post 24hr in response to EE, suggesting a delayed
activation of macrophage-associated pathways (Figure 4A).

Additionally, transcriptional upregulated responses were observed in pre-adipocytes, stem cell,
and vascular cell markers post 45min-RE, whereas enrichment of Adipocyte-1 was suppressed
in response to RE, suggesting subtype-specific adipocyte responses occurring shortly after
resistance exercise (Figure 4A). Furthermore, RE uniquely activated marker genes for both
populations of mature adipocytes and progenitor cells (pre-adipocytes and stem cells) at post
24hr, indicating a long-lasting effect of RE on adipose progenitor and mature adipocyte
transcriptional programs (Figure 4A). Importantly, despite these transcriptional changes, cell-
type proportions assessed by deconvolution remained unchanged, suggesting that these
transcriptional shifts may reflect activation states rather than changes in cell abundance (Figure
S4A).

We next mapped differentially expressed features across omic layers to specific cell types. At
45 min, many transcripts were linked to mast cells and vascular cells (Figure 4B), even though
these cell types were not enriched overall. In both proteomics and phosphoproteomics,
differentially expressed features were primarily markers of adipocytes and vascular cells,
suggesting these cells are major sites of functional change. Many differentially expressed
phosphoproteins associated with cytoskeleton remodeling (e.g., SYNPO, FLNA, TNS1, TLN1)
were attributed to Adipocyte-1 and vascular cells (e.g., SYNE2, NES, AFDN, PKP4), while
metabolic regulators (e.g., PECR, BCL6) were linked to Adipocyte-2 (Figure 4B). Differentially
phosphorylated sites between EE and RE were predominantly mapped to adipocytes at post 4

hr (Figure S4B). NR4A1 is known to activate mast cells, which release histamine[41], a molecular
transducer of exercise responses[42] with wide-ranging beneficial effects in human skeletal
muscle[43]. In ASAT, our data suggest that NR4A1 is a mast cell marker (Figure 4B, S4B). The
upregulated gene expression of NR4A1 at post 45min-EE was followed by increased
abundance of histamine in ASAT at post 4hr-EE (Figure S4C), suggesting the same exercise-
induced histamine release — via mast cells — may therefore also be involved in responses of
ASAT. To investigate whether exercise-induced changes in the ASAT metabolome mirror shifts
in cell type activity, we performed fuzzy c-means clustering on a combined transcriptomics and
metabolomics dataset. We identified 13 clusters per modality, each characterized by a distinct
trajectory over time and unique patterns of cell type and metabolite class enrichment. For
instance, in the EE group, Cluster 3 peaked at 4 hr post-exercise following a modest dip at 45
min, and was enriched for immune cell types including LAMs, mast cells, and NK/T cells, which
aligns well with findings from the cell-type enrichment (Figure 4A). This immune-associated
trajectory was accompanied by increased representation of metabolite classes such as O-PC,
O-PE, and sphingomyelin, supporting the link between phospholipid metabolism and
macrophages[44] (Figure 4C). In RE, Clusters 2, 3, and 6 showed overlapping enrichment for
LAMs and Adipocyte-1, with corresponding decreases post 45min—matching their
transcriptional downregulation (Figure 4A). These clusters were also enriched for amino acids
and diacylglycerols. Unsaturated fatty acids followed distinct temporal trajectories by modality:
EE (Clusters 7, 9, 11) peaked at 45 min, while RE (Clusters 4, 5, 7) peaked at 4 hr. These
clusters were associated with Adipocyte-2 in EE and Adipocyte-1 in RE (Figure 4C, D),
suggesting modality-specific engagement of adipocyte subtypes in lipid metabolism. Similarly,
ceramide-enriched clusters at post 4 hr exercise - cluster 1 in EE and cluster 7 in RE - were
linked to Adipocyte-2 and Adipocyte-1, respectively, further indicating that EE and RE may
differentially regulate lipid metabolism through distinct adipocyte subtypes.

Multi-Omic Network Analysis Links ASAT Features to Cardiometabolic Health and Exercise
Responses

To explore the clinical relevance of multi-omic features in ASAT and their regulation by exercise,
we applied weighted gene co-expression network analysis (WGCNA; Figure 5A)[45] to identify co-
regulated feature clusters (modules; Table S3) within each omic layer using baseline data: 13
transcriptomic modules (T0–T12), 15 proteomic modules (Pr0–Pr14), 19 phosphoproteomic
modules (Ph0–Ph18), and 8 metabolomic modules (M0–M7) (Figure 5B; see WGCNA in
Methods). ORA was performed to identify enriched pathways in each module. Next, we
correlated module eigengenes—the first principal component summarizing overall module
feature expression[46]—with cardiometabolic health traits and examined the module membership
of features differentially regulated by exercise to assess the biological relevance of these
modules. Several modules were significantly associated with adiposity (BMI, WC), insulin
resistance (IR) indices (HOMA-IR, Adipo-IR), and HbA1c across omic layers, including T2, T5,
Pr7, Pr12, Ph10, and M1 (Figure 5B), which showed high degrees of cross-omic intercorrelation
(Figures S5A). Many of these modules were positively correlated with estimated macrophage
proportion but inversely correlated with estimated adipocyte and vascular cell proportions
(Figure S5B), and were enriched in pathways previously implicated in adipose-related metabolic
complications, including inflammation (T2, Pr12)[47], branched-chain amino acid (BCAA)

metabolism (Pr7)[20], post-transcriptional/translational modification (T5)[14], cytoskeleton
organization (Ph10)[48], and ceramides and phosphatidylcholine (M1)[38].

We next examined whether acute exercise may impact features tightly associated with
cardiometabolic health. Ph10 remained significantly associated with HOMA-IR and HbA1c after
adjusting for key covariates (age, sex, WC, p<0.05, Figure S5C), suggesting a direct link
between Ph10 and glycemic control. Among phosphosites differentially regulated by exercise in
Ph10, those with positive intra-module connectivity (kME)—such as AHNAK2-S294 and
SPTBN1-S2165;S2172—were positively correlated with IR indices and HbA1c, whereas
phosphosites with negative kME—including SLC9A3R1-S280, TPH52L1-S148, and GLUL-
S343—were inversely correlated with IR and HbA1c (p<0.05, Figure 5E, F). Strikingly,
phosphosites with positive kME were downregulated, whereas those with negative kME were
mostly upregulated, by acute exercise, particularly following EE at post 4 hr (adjusted p<0.05,
Figure 5C). Exercise may therefore modulate phosphorylation of proteins involved in ASAT
cytoskeletal organization acutely in a manner that contributes to favorable metabolic health.

BCAA catabolism is suppressed in ASAT in obesity, contributing to elevated circulating BCAA
and IR[20,49]. In our data, module Pr5 was inversely correlated with circulating BCAAs (isoleucine,
leucine, and valine) after adjusting for covariates (p < 0.05; Figure S5C). Within Pr5, proteins
with negative kME values were all upregulated by EE at post 4 hr, whereas those with positive
kME values were mostly downregulated (Figure 5D). Functional annotation of Pr5 highlighted
enrichment for RNA processing, protein biosynthesis, cytoskeletal remodeling, and
mitochondrial ribosomal components (Figure 5B), suggesting that acute exercise promotes
proteomic remodeling in ASAT linked to these pathways.

In parallel, acute EE and RE significantly increased circulating levels of branched-chain keto
acids (BCKAs)—specifically ketoleucine and 2-keto-3-methylvalerate—which are derived from
BCAAs through transamination by branched-chain amino transaminase (Figure 5F). Notably,
prior work by Walejko et al. showed that acute exposure of perfused, isolated working hearts to
elevated BCKA levels (mirroring those seen in obesity) was sufficient to induce phosphoprotein
changes, cytoskeletal remodeling, and activation of protein synthesis[50] (Figure 5E). Strikingly,
our results echo these findings: translational pathways were upregulated at post 4 hr in both EE
and RE (Figure 5G), and nearly half of the phosphoproteins differentially phosphorylated in the
BCKA-exposed heart overlapped with those regulated by exercise in ASAT. These overlapping
phosphoproteins were enriched for cytoskeleton remodeling functions (Figure 5H), consistent
with the remodeling signatures reported by Walejko et al. Together, this supports the hypothesis
that transient, exercise-induced increases in BCKAs may trigger shared phosphoproteomic and
cell structural remodeling programs in adipose tissue. While chronic elevation of BCKAs—as
seen in obesity—may contribute to maladaptive cardiac hypertrophy through sustained
activation of protein synthesis, acute exercise-induced BCKA elevations may be health-
promoting by virtue of their intermittent rather than chronic effects.

Acute exercise can stimulate the release of exerkines—secreted factors that mediate metabolic
and functional adaptations in distant tissues—with ASAT serving as a key contributor[19]. To
characterize the exercise-responsive secretome of ASAT, we filtered differentially expressed
transcripts and proteins using curated secretome resources, including MoTrPAC's blood
proteomics panel (*REF: MoTrPAC Blood Companion, in review*) and The Human Protein Atlas[51],
which includes 3,277 known secreted proteins. This yielded 60 unique candidate proteins
(Figure 6B, S6A), the majority of which were identified through proteomics and induced at 4□hr
post-exercise (Figure S6A, Table S4). For metabolites, 92 differentially expressed ASAT
features overlapped with plasma metabolomics (Figure 6C). At baseline, ASAT abundances of
TIMP4 and CCN1 were significantly correlated with their circulating protein levels (p□<□0.05;
Figure S6B), suggesting that ASAT may contribute to their systemic concentrations at rest.
Likewise, ASAT-derived glycerophospholipids (e.g., PC, PE) and sphingolipids (e.g., SM,
ceramide) showed strong correlations with their plasma counterparts (p□<□0.05; Figure S6C).

To explore potential endocrine roles of ASAT-derived exerkines, we applied inter-tissue gene-
gene correlation analysis using GTEx—the most comprehensive multi-tissue human
transcriptomics resource[52] (see Gene-to-Gene Correlations in Methods). We also assigned
each transcript an "extracellular score (0-5)" using COMPARTMENTS (see 'Extracellular score
in Methods')[53] to estimate its secretory likelihood. Secreted candidate genes expressed in SAT
displayed distinct correlations with transcriptional programs in distant tissues—most notably
visceral adipose tissue, skeletal muscle, heart, and thyroid (adjusted p<0.05, Figure 6D). To
infer potential functional pathways in recipient tissues, we applied Gene-Derived Correlations
Across Tissues (GD-CAT)[54], which assumes that if a gene encoding a secreted protein in one
tissue correlates with specific transcriptional programs in another, the protein may mediate inter-
tissue crosstalk[55]. We focused on two exerkine candidates significantly altered in plasma during
or after exercise: *CEACAM8* (CD66b) and *BST2* (adjusted p<0.05, Figure 6E). *CEACAM8*, a
marker gene for LAM (Figure 4C) and known for its immunoregulatory role[56], had a high
extracellular score of 5 and was acutely increased in the plasma during EE, with increased
protein abundance in ASAT at post 4hr-EE (Figure 6E). In GTEx, SAT-*CEACAM8* was
associated with upregulated immune responses (e.g., MHC assembly, antimicrobial humoral
response) and downregulated ribosome biogenesis in visceral adipose tissue (VAT) and thyroid
(adjusted p<0.05, Figure 6F, Table S5). In contrast, SAT-CEACAM8 correlated with enhanced
protein translation and dampened B cell receptor signaling in skeletal muscle, suggesting
possible tissue-specific endocrine functions (adjusted p<0.05, Figure 6F). BST2, a
transmembrane glycoprotein linked to obesity-associated inflammation[57] with extracellular score
of 4.5, decreased systemically at post 4hr-RE (adjusted p<0.05). Interestingly, this coincided
with the prevention of an increase in ASAT BST2 post-exercise (EE, adjusted p<0.05; RE,
adjusted p=0.14), contrasting with its rise in CON (Figure 6E). In GTEx, SAT-BST2 was linked
to downregulated mitochondrial energetics (proton motive force-driven ATP synthesis) in VAT,
heart, and thyroid, but not skeletal muscle (adjusted p<0.05, Figure 6F, Table S5). Conversely,
SAT-BST2 was associated with upregulated complement activation in skeletal muscle (adjusted
p<0.05, Figure 6F). These data suggest that exercise-induced reductions in BST2 may confer
systemic metabolic benefits by limiting inflammation and preserving mitochondrial function

across tissues. Together, these findings reveal novel ASAT-derived exerkine candidates
modulated by acute exercise and demonstrate how integrated transcriptomic correlations can
uncover potential endocrine effects of adipose tissue on distant organs.

Exercise-Induced ASAT Secretomes: Adaptations in the Endocrine Network with Exercise
Training

To investigate whether ASAT-derived exerkine candidates contribute to systemic adaptations
during endurance training, we leveraged the MoTrPAC rat training dataset[58], which includes
transcriptomic profiles of subcutaneous white adipose tissue (WATSC) and 17 other tissues at
multiple time points (1, 2, 4, and 8 weeks; TR1W–TR8W), along with age-matched non-trained
controls (CON). This dataset enabled us to assess how WATSC exerkine candidates correlate
with target tissues and how these relationships evolve with training. We applied Quantitative
Endocrine Network Interaction Estimation[59], a systems genetics framework that computes a
"secretome score" (Ssec) to estimate the endocrine influence of secretory genes across origin–
target tissue pairs. Briefly, for each gene in the origin tissue, a -log10 sum of the correlation p-
values with all genes in the target tissue is computed to generate Ssec, where higher scores
indicate broader or stronger endocrine connectivity (see QENIE in Methods). Using this
framework, we previously identified WATSC as one of the most responsive endocrine tissues to
endurance training[60]. We designated WATSC as the origin tissue and calculated Ssec for 2,248
secretory protein-coding genes found in humans (Figure 6A) across multiple target tissues
(Figure 7A; see Methods; full dataset in Table S6). Given the strong sexual dimorphism of
WATSC in this cohort[61], all correlations were adjusted for sex. Ssec distributions revealed
tissue- and time-dependent endocrine connectivity changes (Figure 7B). Out of 60 ASAT protein
exerkine candidates we identified (Figure 6), 53 of them were mappable in rat WATSC.
Focusing on these 53 candidates, we found significant alteration in their Ssec rank distributions
between control and 8-week trained rats for key endocrine connections: WATSC-to-vastus
lateralis, liver, and kidney (adjusted p<0.05, Figure 7C). These rank shifts were paralleled by
increases in average Ssec values in 8-week trained rats compared to controls in all three tissue
connections (adjusted p<0.05, Figure S7A), indicating strengthened endocrine influence of
WATSC with training.

Time-course analysis of individual Ssec trajectories showed progressive increases for several
exerkines. For example, *CX3CL1* (fractalkine)—a known exerkine also released from skeletal
muscle[62]—displayed increased Ssec over time, peaking at 8 weeks in WATSC-to-colon,
hypothalamus, and lung connections (Figure 7D). Feature-level comparisons revealed
heterogeneous adaptations across candidates and tissues (Figure S7B). While overall Ssec
ranks for WATSC-to-vena cava were unchanged (Figure 7C), several candidates (e.g., *TIMP4,*
LHPP, PFKFB2, CX3CL1) showed dramatically elevated Ssec values at 8 weeks of training,
suggesting a role in vascular crosstalk during habitual exercise adaptation (Figure S7B). We
next highlighted top candidates showing the greatest increases in Ssec rank between control
and 8-week trained rats across the vastus lateralis, liver, and kidney (Figure 7E). *IGFBP7*
(Insulin-like Growth Factor Binding Protein 7), which was induced by EE at 4 hr (Figure 7F),
emerged as a notable candidate in WATSC-to-vastus lateralis and WATSC-to-liver connections
(Figure 7E). In the WATSC-to-vastus lateralis axis, *IGFBP7* rose from rank 1477 in CON to 95

in trained rats. Similarly, in the WATSC-to-liver axis, *IGFBP7* jumped from 1942 to 121 (Figure
7E). GD-CAT analysis revealed that *IGFBP7* was associated with enhanced mitochondrial
metabolism pathways in the vastus lateralis of 8-week trained rats, while control rats showed
enrichment of inflammatory signaling and mitochondrial suppression (adjusted p<0.05; Figure
7G). These findings align with *in vitro* evidence showing improved mitochondrial respiration in
myotubes treated with IGFBP7-conditioned media[63], supporting its potential endocrine role in
exercise adaptations. In the liver, WATSC-*IGFBP7* correlated with upregulated mitochondrial
complex expression, translation, and lipid oxidation, along with downregulated immune
responses in 8-week trained rats (adjusted p<0.05; Figure 7H)—associations absent in control
rats. Notably, IGFBP7 is a known tumor suppressor; its deletion or downregulation in the liver
induces hepatocellular carcinoma[64,65], raising the possibility of additional protective roles in
exercise-adapted crosstalk mediated by WATSC.

Together, these systems genetics analyses suggest that acutely-induced ASAT exerkines can
exert tissue-specific endocrine effects, and that repeated endurance training progressively
reshapes the WATSC-mediated secretome landscape. This endocrine network remodeling may
underlie key systemic adaptations induced by habitual exercise.

DISCUSSION

In this study, we captured modality- and time-specific multi-omic molecular responses in human
abdominal subcutaneous adipose tissue (ASAT) following acute exercise using both endurance
(EE) and resistance (RE) modalities across multiple post-exercise time points in sedentary
adults. Acute exercise triggered diverse transcriptional and proteomic responses involving
substrate metabolism, angiogenesis, adipogenesis, ECM remodeling, protein synthesis, cell
signaling, cytoskeletal remodeling, and inflammation and metabolite-specific changes. Using
integrative bioinformatics, we identified cell-type-dependent responses, novel exercise-
responsive features linked to cardiometabolic health, and a set of putative adipose-derived
exerkines. In addition to these post-acute-exercise changes, baseline sex differences and
variations in cell-type proportions were examined, offering insights into how acute exercise
responses may initiate early adaptations towards long-term benefits.

Adipose tissue exhibits circadian rhythmicity in gene expression patterns and physiological
functions. This includes mediating key metabolic enzymes and transport systems involved in
lipid metabolism, lipolysis/lipogenesis and the secretion of adipokines, thus influencing systemic
metabolism, energy homeostasis and inflammatory processes[66,67]. By inclusion of a non-
exercising control group in our differential analysis, we were able to dissociate exercise-specific
effects from confounding time-of-day and fasting influences—particularly pronounced in adipose
tissue and more substantial than anticipated (*REF: MoTrPAC Molecular Landscape, in review*).
Although previous studies have reported post-acute-exercise changes in core clock genes[10,68],
we observed similar patterns in non-exercising controls, suggesting that some of these
transcriptomic shifts may be driven by time-of-day effects rather than exercise *per se*. This
comparison provides an additional layer of confidence in interpreting true exercise-responsive
signals. Our design also enabled us to detect exerkines such as BST2, whose significance
emerged specifically in relation to the control group trajectory. To our knowledge, this study

represents one of the first efforts to directly account for non-exercise-mediated changes when
assessing acute exercise-induced molecular responses in adipose tissue and may serve as a
reference point for future studies aiming to disentangle these overlapping influences.

The actin cytoskeleton plays a central role in insulin action and adipocyte size dynamics[48,69].
Although the precise functions of many phosphosites remain unknown, several within the Ph10
module were strongly associated with insulin resistance. Notably, acute exercise modulated
these phosphosites in a direction consistent with improved metabolic health—sites positively
associated with insulin resistance tended to be downregulated, while those inversely associated
with insulin resistance were upregulated. The overall pattern suggests that acute exercise may
activate phosphosignaling within cytoskeletal remodeling pathways that contribute to favorable
metabolic responses. Cell-type analysis further revealed that Adipocyte-1—characterized by
anti-oxidative and translational activity—and vascular cells were the predominant sources of
cytoskeleton-related phosphosignaling. This raises the possibility that these cell types may be
key mediators of early, metabolically beneficial responses to acute exercise.

A major advance of this study was the identification of adipose-derived exerkine candidates,
derived by filtering differentially regulated features through plasma proteomics, metabolomics,
and curated secretome databases. Using GD-CAT[54] and GTEx pan-tissue transcriptomics[52], we
inferred potential endocrine actions of several candidates—such as CEACAM8 and BST2—
whose predicted effects varied across tissues. Further, by applying QENIE[59] to the MoTrPAC rat
endurance training dataset[58], we found that adipose endocrine networks are remodeled with
training. Several exerkine candidates, particularly IGFBP7, appeared central to this adaptive
rewiring, especially in WATSC-to-vastus lateralis and WATSC-to-liver axes. Interestingly, 8
weeks of endurance training led to modest but significant downregulation of *IGFBP7* transcript
levels in WATSC (male rats: –0.11 $\log_2$FC; female rats: –0.22 $\log_2$FC; FDR = 0.008)[58], without
corresponding changes in protein abundance—suggesting that, like many exerkines[70,71], the
systemic effects of IGFBP7 may be primarily driven by acute induction rather than chronic
upregulation. While our systems genetics approach is correlation-based and requires
experimental validation, this framework has been successfully used to uncover novel crosstalk
regulators that have been validated in pre-clinical models[55,59,72]. Additionally, other candidates—
TIMP4 (Tissue Inhibitor of Metalloproteinase 4), LRP1 (Low-density lipoprotein receptor-related
protein 1), and RETN (Resistin)— also emerged among top-ranking factors across several
adipose-to-target tissue connections in trained rats, warranting further mechanistic investigation.

Despite the strengths of our design and depth of analysis, several limitations should be
acknowledged: Participants served as their own control within each exercise group, but
comparisons across groups were not conducted using a cross over design. By design the two
exercise modalities were not matched for exercise duration and intensity or energy
expenditure[73]. While this limits certain interpretations, the goal was to model real-world clinical
approaches rather than creating tightly matched interventions. Moreover, proteomic and
phosphoproteomic profiling was not performed at post 45 min or post 24 hr, possibly resulting in
missed windows of dynamic regulation. Given the robust changes at post 4hr, additional time
points could provide a more complete temporal map. Finally, our secretome and endocrine
modeling leveraged rodent training data, which may not fully translate to humans. Integration of

anticipated MoTrPAC 12-week training data in sedentary adults will allow deeper analysis of
secretome dynamics and tissue crosstalk in the context of long-term adaptation.

ACKNOWLEDGEMENTS

We would like to acknowledge Dr. Marcus Seldin at the University of California, Irvine, for
assisting GTEx and QENIE analysis. The graphical abstract and schematic figures were created
using BioRender (www.biorender.com), and confirmation of publication and licensing rights was
obtained. The MoTrPAC Study is supported by NIH grants U24OD026629 (Bioinformatics
Center), U24DK112349, U24DK112342, U24DK112340, U24DK112341, U24DK112326,
U24DK112331, U24DK112348 (Chemical Analysis Sites), U01AR071133, U01AR071130,
U01AR071124, U01AR071128, U01AR071150, U01AR071160, U01AR071158 (Clinical
Centers), U24AR071113 (Consortium Coordinating Center), U01AG055133, U01AG055137,
U01AG055135, U01AG070959, U01AG070960, and U01AG070928 (Pre-Clinical Animal Sites).

Additional grant funding: D.H.K: K23HL164980, 23CDA1040581; N.M and S.E: P30AG094848.

AUTHOR CONTRIBUTIONS

JNA, MJW, and LMS conceptualized and designed the study. CA, CAJ, and TJS performed
data analyses and developed figures. CA, CAJ, KLW, GMM, JAS, ZH, MRV, JN, SEE, NM, YS,
MFP, SCB, LJG, KAE, CBN, BCB, JNA, MJW, and LMS contributed to data interpretation. CA
and LMS drafted the work with input from other authors. All authors have participated in revising
the work. All authors have read and approved the final version of the manuscript.
551

MoTrPAC Study Group:

Bioinformatics Center: David Amar, Trevor Hastie, David Jimenez-Morales, Daniel H. Katz,
Malene E. Lindholm, Samuel Montalvo, Robert Tibshirani, Jay Yu, Jimmy Zhen, Euan A. Ashley,
Matthew T. Wheeler

Biospecimens Repository: Sandra T. May, Jessica L. Rooney, Russell Tracy

Data Management, Analysis, and Quality Control Center: Catherine Gervais, Fang-Chi Hsu,
Byron C. Jaeger, David Popoli, Joseph Rigdon, Courtney G. Simmons, Cynthia L. Stowe,
Michael E. Miller

Exercise Intervention Core: W. Jack Rejeski

NIH: Ashley Y. Xia

Preclinical Animal Study Sites: Sue C. Bodine, R. Scott Rector

Chemical Analysis Sites: Hiba Abou Assi, Mary Anne S. Amper, Brian J. Andonian, Isaac K.
Attah, Jacob L. Barber, Kevin Bonanno, Clarisa Chavez Martinez, Natalie M. Clark, Johanna Y.
Fleischman, David A. Gaul, Yongchao Ge, Marina A. Gritsenko, Joshua R. Hansen, Patrick Hart,
Zhenxin Hou, Chelsea M. Hutchinson-Bunch, Olga Ilkayeva, Gayatri Iyer, Pierre M. Jean-
Beltran, Christopher A. Jin, Maureen T. Kachman, Hasmik Keshishian, Damon T. Leach,
Minghui Lu, D. R. Mani, Gina M. Many, Nada Marjanovic, Nikhil Milind, Matthew E. Monroe,

Ronald J. Moore, Venugopalan D. Nair, German Nudelman, Nora-Lovette Okwara, Vladislav A.
Petyuk, Paul D. Piehowski, Hanna Pincas, Wei-Jun Qian, Prashant Rao, Abraham Raskind,
Alexander Raskind, Stas Rirak, Jeremy M. Robbins, Margaret Robinson, Tyler J. Sagendorf,
James A. Sanford, Gregory R. Smith, Kevin S. Smith, Yifei Sun, Mital Vasoya, Nikolai G. Vetr,
Alexandria Vornholt, Yilin Xie, Xuechen Yu, Elena Zaslavsky, Zidong Zhang, Bingqing Zhao,
Joshua N. Adkins, Charles F. Burant, Steven A. Carr, Clary B. Clish, Facundo M. Fernandez,
Robert E. Gerszten, Stephen B. Montgomery, Christopher B. Newgard, Eric A. Ortlund, Stuart C.
Sealfon, Michael P. Snyder, Martin J. Walsh

Clinical Sites: Cheehoon Ahn, Alicia Belangee, Bryan C. Bergman, Daniel H. Bessesen,
Gerard A. Boyd, Anna R. Brandt, Nicholas T. Broskey, Toby L. Chambers, Clarisa Chavez
Martinez, Maria Chikina, Alex Claiborne, Zachary S. Clayton, Paul M. Coen, Katherine A.
Collins-Bennett, Tiffany M. Cortes, Gary R. Cutter, Matthew Douglass, Daniel E. Forman, Will A.
Fountain, Aaron H. Gouw, Kevin J. Gries, Fadia Haddad, Joseph A. Houmard, Kim M. Huffman,
Ryan P. Hughes, John M. Jakicic, Catherine M. Jankowski, Neil M. Johannsen, Johanna L.
Johnson, Erin E. Kershaw, Dillon J. Kuszmaul, Bridget Lester, Colleen E. Lynch, Edward L.
Melanson, Cristhian Montenegro, Kerrie L. Moreau, Masatoshi Naruse, Jia Nie, Bradley C Nindl,
Maria F. Pino, Tuomo Rankinen, Ulrika Raue, Ethan Robbins, Kaitlyn R. Rogers, Renee J.
Rogers, Irene E. Schauer, Robert S. Schwartz, Chad M. Skiles, Lauren M. Sparks, Maja
Stefanovic-Racic, Andrew M. Stroh, Kristen J. Sutton, Anna Thalacker-Mercer, Todd A. Trappe,
Caroline S. Vincenty, Elena Volpi, Katie L. Whytock, Gilhyeon Yoon, Thomas W. Buford, Dan M.
Cooper, Sara E. Espinoza, Bret H. Goodpaster, Wendy M. Kohrt, William E. Kraus, Nicolas
Musi, Shlomit Radom-Aizik, Blake B. Rasmussen, Eric Ravussin, Scott Trappe

MoTrPAC Study Group Acknowledgements

Nicole Adams, Abdalla Ahmed, Andrea Anderson, Carter Asef, Arianne Aslamy, Marcas M.
Bamman, Jerry Barnes, Susan Barr, Kelsey Belski, Will Bennett, Amanda Boyce, Brandon
Bukas, Emily Carifi, Chih-Yu Chen, Haiying Chen, Shyh-Huei Chen, Samuel Cohen, Audrey
Collins, Gavin Connolly, Elaine Cornell, Julia Dauberger, Carola Ekelund, Shannon S. Emilson,
Jerome Fleg, Nicole Gagne, Mary-Catherine George, Ellie Gibbons, Jillian Gillespie, Aditi Goyal,
Bruce Graham, Xueyun Gulbin, Jere Hamilton, Leora Henkin, Andrew Hepler, Lidija Ivic, Ronald
Jackson, Andrew Jones, Lyndon Joseph, Leslie Kelly, Gary Lee, Adrian Loubriel, Ching-ju Lu,
Kristal M. Maner-Smith, Ryan Martin, Padma Maruvada, Alyssa Mathews, Curtis McGinity,
Lucas Medsker, Kiril Minchev, Samuel G. Moore, Michael Muehlbauer, Anne Newman, John
Nichols, Concepcion R. Nierras, George Papanicolaou, Lorrie Penry, June Pierce, Megan
Reaves, Eric W. Reynolds, Teresa N. Richardson, Jeremy Rogers, Scott Rushing, Santiago
Saldana, Rohan Shah, Samiya M. Shimly, Cris Slentz, Deanna Spaw, Debbie Steinberg,
Suchitra Sudarshan, Alyssa Sudnick, Jennifer W. Talton, Christy Tebsherani, Nevyana
Todorova, Jennifer Walker, Michael P. Walkup, Anthony Weakland, Gary Weaver, Christopher
Webb, Sawyer Welden, John P. Williams, Marilyn Williams, Leslie Willis, Yi Zhang, Frank Booth,
Andrea Hevener, Ian Lanza, Jun Li, K Sreekumaran Nair

DECLARATION OF INTERESTS

The authors declare no competing interests.

STAR Methods
Experimental model and study participant details
IRB
The Molecular Transducers of Physical Activity Consortium (MoTrPAC) (NCT03960827) is a
multicenter study designed to isolate the effects of structured exercise training on the molecular
mechanisms underlying the health benefits of exercise and physical activity. Methods are
described here in sufficient detail to interpret results, with references to supplemental material
and prior publications for detailed descriptions. The present work includes data from prior to the
suspension of the study in March 2020 due to the Covid-19 pandemic. The study was
conducted in accordance with the Declaration of Helsinki and approved by the Institutional
Review Board of Johns Hopkins University School of Medicine (IRB protocol # JHMIRB5;
approval date: 05/06/2019). All MoTrPAC participants provided written informed consent for the
MoTrPAC study indicating whether they agreed to open data sharing and at what level of
sharing they wanted. Participants could choose to openly share all de-identified data, with the
knowledge that they could be reidentified, and they could also choose to openly share limited
de-identified individual level data, which is lower risk of re-identification. All analyses and
resulting data and results are shared in compliance with the NIH Genomic Data Sharing (GDS)
policy and DSMB requirements for the randomized study.
Participant Characteristics
Volunteers were screened to: (a) ensure they met all eligibility criteria; (b) acquire phenotypic
assessments of the study population; and (c) verify they were healthy and medically cleared to
be formally randomized into the study (cite Clinical Landscape).34 Participants were then
randomized into intervention groups stratified by clinical site and completed a pre-intervention
baseline acute test. A total of 176 participants completed the baseline acute test (EE=66,
RE=73, and CON=37), and among those, 175 (99%) had at least one biospecimen sample
collected. Participants then began 12 weeks of exercise training or control conditions. Upon
completion of the respective interventions, participants repeated phenotypic testing and the
acute test including biospecimen sampling. Due to the COVID-19 suspension, only 45
participants (26%) completed the post-intervention follow-up acute exercise test (some with less
than 12 weeks of training), with 44 completing the post-intervention biospecimen collections.
See Clinical Landscape for further detail.
Participant Assessments
As described in Clinical Landscape (REF, MoTrPAC Clinical landscape, *in review*) and protocol
paper[73], prior to randomization, participant screening was completed via questionnaires, and
measurements of anthropometrics, resting heart rate and blood pressure, fasted blood panel,
cardiorespiratory fitness, muscular strength, and free-living activity and sedentary behavior.
Cardiorespiratory fitness was assessed using a cardiopulmonary exercise test (CPET) on a
cycle ergometer (Lode Excalibur Sport Lode BV, Groningen, The Netherlands) with indirect
calorimetry. Quadriceps strength was determined by isometric knee extension of the right leg at
a 60° knee angle using a suitable strength device[73]. Grip strength of the dominant hand was
obtained using the Jamar Handheld Hydraulic Dynamometer (JLW Instruments, Chicago, IL).
See Clinical Landscape (REF, MoTrPAC Clinical Landscape, *in review*) for participant
assessment results.

Acute Exercise Intervention

The pre-intervention baseline EE acute bouts were composed of three parts: (1) a 5min warm up at 50% of the estimated power output to elicit 65% VO2peak, (2) 40min cycling at ~65% VO2peak, and (3) a 1min recovery at ~25 W. The pre-intervention baseline RE acute bouts were composed of a 5min warm-up at 50-60% heart rate reserve (HRR) followed by completion of three sets each of five upper (chest press, overhead press, seated row, triceps extension, biceps curl) and three lower body (leg press, leg curl, leg extension) exercises to volitional fatigue (~10RM) with 90sec rest between each set. Participants randomized to CON did not complete exercise during their acute tests. Participants rested supine for 40min to mirror the EE and RE acute test schedule. See Clinical Landscape (REF, MoTrPAC Clinical Landscape, *in review*) for acute bout results.

Adipose Tissue Collection and Omics Profiling

Abdominal subcutaneous adipose tissue (ASAT) biopsies were collected from all participants as part of the acute exercise test. Prior to testing, participants adhered to pre-sampling restrictions, including abstention from caffeine, alcohol, NSAIDs, exercise, and specific dietary components[73]. Upon arrival, participants rested supine for at least 30 minutes prior to the pre-exercise biopsy. Post-exercise adipose collections were conducted according to randomized temporal profiles designed to span the acute recovery period while minimizing participant burden. These profiles included early (Pre and 45 min Post), middle (Pre and 4 hr Post), late (Pre and 24 hr Post), and full (all four time points, although ASAT was only collected at Pre and 4 hr in this group). Final sample sizes were 43 for early, 80 for middle, and 49 for late profiles. The post-exercise biopsy was timed immediately following completion of the final exercise component—40 minutes of cycling for EE, the third set of leg extensions for RE, or the 40-minute rest period for CON. ASAT biopsies were successfully collected in 172 participants (EE = 62, RE = 73, CON = 37), and only these participants were included in this study.

Adipose tissue was subjected to multi-omics profiling. Transcriptomics and metabolomics analyses were performed at all four time points (Pre, 45 min, 4 hr, and 24 hr post-exercise). Proteomics and phosphoproteomics were conducted at Pre and 4 hr Post only. All biospecimen processing and quality control procedures followed standardized protocols as described in the Clinical Landscape (*REF: MoTrPAC Clinical Landscape, in review*) and protocol design manuscript[73].

Molecular Assays

Transcriptomics

RNA Sequencing (RNA-Seq) was performed at Stanford University and the Icahn School of Medicine at Mount Sinai.

Extraction of total RNA

Approximately 50 mg of frozen ASAT were disrupted in Agencourt RNAdvance tissue lysis
buffer (Beckman Coulter, Brea, CA) using a tissue ruptor (Omni International, Kennesaw, GA,
#19-040E). Total RNA was extracted in a BiomekFX automation workstation according to the
manufacturer's instructions for tissue-specific extraction. The RNA was quantified by NanoDrop
(ThermoFisher Scientific, # ND-ONE-W) and Qubit assay (ThermoFisher Scientific), and the
quality was determined by either Bioanalyzer or Fragment Analyzer analysis.

mRNA Sequencing Library Preparation

Universal Plus mRNA-Seq kit from NuGEN/Tecan (# 9133) were used for generation of

RNA-Seq libraries derived from poly(A)-selected RNA according to the manufacturer's

instructions. Universal Plus mRNA-Seq libraries contain dual (i7 and i5) 8 bp barcodes and an 8
bp unique molecular identifier (UMI), which enable deep multiplexing of NGS sequencing
samples and accurate quantification of PCR duplication levels. Approximately 300ng of ASAT
was used. The Universal Plus mRNA-Seq workflow consists of poly(A) RNA selection, RNA
fragmentation and double-stranded cDNA generation using a mixture of random and oligo(dT)
priming, end repair to generate blunt ends, adaptor ligation, strand selection, AnyDeplete
workflow to remove unwanted ribosomal and globin transcripts, and PCR amplification to enrich
final library species. All library preparations were performed using a Biomek i7 laboratory
automation system (Beckman Coulter). Tissue-specific reference standards provided by the
consortium were included with all RNA isolations to QC the RNA.

RNA Sequencing, quantification, and normalization

RNA sequencing, quantification, and normalization Pooled libraries were sequenced on an
Illumina NovaSeq 6000 platform (Illumina, San Diego, CA, USA) to a target depth of 40 million
read pairs (80 million paired-end reads) per sample using a paired-end 100 base pair run
configuration. In order to capture the 8-base UMIs, libraries were sequenced using 16 cycles for
the i7 index read and 8 cycles for the i5 index read. Reads were demultiplexed with bcl2fastq2
(v2.20.0) using options --use-bases-mask Y*,I8Y*,I*,Y* --mask-short-adapter-reads 0 --
minimum-trimmed-read-length 0 (Illumina, San Diego, CA, USA), and UMIs in the index FASTQ
files were attached to the read FASTQ files. Adapters were trimmed with cutadapt (v1.18)[74], and
trimmed reads shorter than 20 base pairs were removed. FastQC (v0.11.8) was used to
generate pre-alignment QC metrics. STAR (v2.7.0d)[75] was used to index and align reads to
release 38 of the Ensembl Homo sapiens (hg38) genome and Gencode (Version 29). Default
parameters were used for STAR's genomeGenerate run mode; in STAR's alignReads run mode,
SAM attributes were specified as NH HI AS NM MD nM, and reads were removed if they did not
contain high-confidence collapsed splice junctions (--outFilterType BySJout). RSEM (v1.3.1)[76]
was used to quantify transcriptome-coordinate-sorted alignments using a forward probability of
0.5 to indicate a non-strand-specific protocol. Bowtie 2 (v2.3.4.3)[77] was used to index and align
reads to globin, rRNA, and phix sequences in order to quantify the percent of reads that
mapped to these contaminants and spike-ins. UCSC's gtfToGenePred was used to convert the
hg38 gene annotation (GTF) to a refFlat file in order to run Picard CollectRnaSeqMetrics

(v2.18.16) with options MINIMUM_LENGTH=50 and RRNA_FRAGMENT_PERCENTAGE=0.3.
UMIs were used to accurately quantify PCR duplicates with NuGEN's "nodup.py" script
(https://github.com/tecangenomics/nudup). QC metrics from every stage of the quantification
pipeline were compiled, in part with multiQC (v1.6). The openWDL-based implementation of the
RNA-Seq pipeline on Google Cloud Platform is available on Github
(https://github.com/MoTrPAC/motrpac-rna-seq-pipeline). Filtering of lowly expressed genes and
normalization were performed separately in each tissue. RSEM gene counts were used to
remove lowly expressed genes, defined as having 0.5 or fewer counts per million in at least 10%
of samples. These filtered raw counts were used as input for differential analysis with the
variancePartition::dream[78], as described in the statistical analysis methods section. To generate
normalized sample-level data for downstream visualization, filtered gene counts were TMM-
normalized using edgeR::calcNormFactors, followed by conversion to log counts per million with
edgeR::cpm.

Principal Component Analysis and calculation of the variance explained by variables of interest
were used to identify and quantify potential batch effects. Based on this analysis, processing
batch, clinical site, percentage of UMI duplication, and RNA Integrity Number (RIN) technical
effects were regressed out of the TMM-normalized counts via linear regression using
limma::RemoveBatchEffect[79] function in R. A design matrix including age, sex, and a
combination of group and timepoint was used during batch effect removal to avoid removing
variance attributable to biological effects.

ASAT RNA extraction was performed in separate batches according to exercise modality, so the
extraction batch and exercise modality were perfectly collinear. This collinearity was identified at
the RNA extraction step and samples were randomized prior to construction of cDNA libraries.
Ultimately, under this processing implementation, differences in gene expression attributable to
exercise group can be impossible to disentangle from RNA extraction batch effects, so the
batch variable was not regressed out in the technical effect stage or included as a covariate in
the differential analysis, and left as an experimental limitation.

RNA Quality Inclusion Criteria

In the blood transcriptomic data, 79/1032 processed adult-sedentary samples had RIN values
under 5. Based on established guidelines and internal quality control assessments, any samples
with a RIN score below 5 were excluded from further analysis due to concerns about potential
degradation artifacts. Additional visualizations and summary figures supporting this decision are
available in the quality control report at: https://github.com/MoTrPAC/precovid-
analyses/tree/main/QC/

Proteomics and Phosphoproteomics

Study design

LC-MS/MS analysis was performed at the Pacific Northwest National Laboratories (PNNL). 46
ASAT samples representing baseline and 4-hours post-intervention from all three groups were
analyzed.

Generation of common reference

Samples were digested at each site following sample processing protocol described below, and
centrally aliquoted. Common reference for adipose tissue was generated using bulk tissue from
6 individuals representing a 4:2 ratio of female:male. 250 µg aliquots of common reference
samples were made to be included in each multiplex (described below) and additional aliquots
are stored for inclusion in future MoTrPAC phases to facilitate data integration.

Sample processing

Proteomics analyses were performed using clinical proteomics protocols described
previously[80,81]. Samples were lysed in ice-cold, freshly-prepared lysis buffer (8 M urea (Sigma-
Aldrich, St. Louis, Missouri), 50 mM Tris pH 8.0, 75 mM sodium chloride, 1 mM EDTA, 2 µg/ml
Aprotinin (Sigma-Aldrich, St. Louis, Missouri), 10 µg/ml Leupeptin (Roche CustomBiotech,
Indianapolis, Indiana), 1 mM PMSF in EtOH, 10 mM sodium fluoride, 1% phosphatase inhibitor
cocktail 2 and 3 (Sigma-Aldrich, St. Louis, Missouri), 10 mM Sodium Butyrate, 2 µM SAHA, and
10 mM nicotinamide and protein concentration was determined by BCA assay. Protein lysate
concentrations were normalized within samples of the same tissue type, and protein was
reduced with 5 mM dithiothreitol (DTT, Sigma-Aldrich) for 1 hour at 37°C with shaking at 1000
rpm on a thermomixer, alkylated with iodoacetamide (IAA, Sigma-Aldrich) in the dark for 45
minutes at 25°C with shaking at 1000 rpm, followed by dilution of 1:4 with Tris-HCl, pH 8.0 prior
to adding digestion enzymes. Proteins were first digested with LysC endopeptidase (Wako
Chemicals) at a 1:50 enzyme:substrate ratio (2 hours, 25 °C, 850 rpm), followed by digestion
with trypsin (Promega) at a 1:50 enzyme:substrate ratio (or 1:10 ratio for adipose tissue; 14
hours, 25 °C, 850 rpm). The next day formic acid was added to a final concentration of 1% to
quench the reaction. Digested peptides were desalted using Sep-Pac C18 columns (Waters),
concentrated in a vacuum centrifuge, and a BCA assay was used to determine final peptide
concentrations. 250µg aliquots of each sample were prepared, dried down by vacuum
centrifugation and stored at -80°C.

Tandem mass tag (TMT) 16-plex isobaric labeling reagent (ThermoFisher Scientific) was used
for this study. Samples were randomized across the first 15 channels of TMT 16-plexes, and the
last channel (134N) of each multiplex was used for a common reference that was prepared prior
to starting the study (see above). Randomization of samples across the plexes within each site
was done using https://github.com/MoTrPAC/clinical-sample-batching, with the goal to have all
timepoints per participant in the same plex, and uniform distribution of groups (endurance,
resistance, control), sex and sample collection clinical site across the plexes. Peptide aliquots
(250 µg per sample) were resuspended to a final concentration of 5 µg/µL in 200 mM HEPES,
pH 8.5 for isobaric labeling. TMT reagent was added to each sample at a 1:2 peptide: TMT ratio,
and labeling proceeded for 1 hour at 25°C with shaking at 400 rpm. The labeling reaction was
diluted to a peptide concentration of 2 µg/µL using 62.5 µL of 200 mM HEPES and 20% ACN. 3

µL was removed from each sample to quantify labeling efficiency and mixing ratio. After labeling
QC analysis, reactions were quenched with 5% hydroxylamine and samples within each
multiplex were combined and desalted with Sep-Pac C18 columns (Waters).

Combined TMT multiplexed samples were then fractionated using high pH reversed phase
chromatography on a 4.6mm ID x 250mm length Zorbax 300 Extend-C18 column (Agilent) with
5% ammonium formate/2% Acetonitrile as solvent A and 5% ammonium formate/90%
acetonitrile as solvent B. Samples were fractionated with 96min separation gradient at flow rate
of 1mL/min and fractions were collected at each minute onto a 96-well plate. Fractions are then
concatenated into 24 fractions with the following scheme: fraction 1 = A1+C1+E1+G1, fraction 2
= A2+C2+E2+G2, fraction 3 = A3+C3+E3+G3, all the way to fraction 24 = B12+D12+F12+H12
following the same scheme. 5% of each fraction was removed for global proteome analysis, and
the remaining 95% was further concatenated to 12 fractions for phosphopeptide enrichment
using immobilized metal affinity chromatography (IMAC).

Phosphopeptide enrichment was performed through immobilized metal affinity chromatography
(IMAC) using Fe 3+ -NTA-agarose beads, freshly prepared from Ni-NTA-agarose beads
(Qiagen, Hilden, Germany) by sequential incubation in 100 mM EDTA to strip nickel, washing
with HPLC water, and incubation in 10 mM iron (III) chloride). Peptide fractions were
resuspended to 0.5 µg/uL in 80% ACN + 0.1% TFA and incubated with beads for 30 minutes in
a thermomixer set to 1000 rpm at room temperature. After 30 minutes, beads were spun down
(1 minute, 1000 rcf) and supernatant was removed and saved as flow-through for subsequent
enrichments. Phosphopeptides were eluted off IMAC beads in 3x 75 µL of agarose bead elution
buffer (500 mM K2HPO4 , pH 7.0), desalted using C18 stage tips, eluted with 50% ACN, and
lyophilized. Samples were then reconstituted in 3% ACN / 0.1% FA for LC-MS/MS analysis (9
µL reconstitution / 4 µL injection at the BI; 12 µL reconstitution / 5 µL injection at PNNL).

Data acquisition

For mass spectrometry analysis of the global proteome, online separation was performed using
a Dionex Ultimate 3000 UHPLC system (ThermoFisher) and a 25 cm x 75 µm i.d. picofrit
column packed in-house with C18 silica (1.7 µm UPLC BEH particles, Waters Acquity) with
solvent A of 0.1% formic acid/3% acetonitrile and solvent B of 0.1% formic acid/90% acetonitrile,
flow rate of 200nL/min and the following gradient: 1-8% B in 10min, 8-25% B in 90min, 25-35%
B in 10min, 35-75% B in 5min, and 75-5% in 3min. Proteome fractions were analyzed on a Q
Exactive HF-X Plus mass spectrometer (ThermoFisher Scientific) with MS1 scan across the
300-1800 m/z range at 60,000 resolution, AGC target of 3x106 and maximum injection time of
20ms. MS2 scans of most abundant 12 ions were performed at 30,000 resolution with AGC
target of 1x105 and maximum injection time of 100ms, isolation window of 0.7m/z and
normalized collision energy of 30.

For mass spectrometry analysis of the phosphoproteome, online separation was performed
using a Dionex Ultimate 3000 UHPLC system (ThermoFisher) and a 25 cm x 75 µm i.d. picofrit
column packed in-house with C18 silica (1.7 µm UPLC BEH particles, Waters Acquity) with
solvent A of 0.1% formic acid/3% acetonitrile and solvent B of 0.1% formic acid/90% acetonitrile,

flow rate of 200nL/min and the following gradient: 1-8% B in 10min, 8-25% B in 90min, 25-35%
B in 10min, 35-75% B in 5min, and 75-5% in 3min. Phosphoproteome fractions were analyzed
on a Q Exactive HF-X Plus mass spectrometer (ThermoFisher Scientific) with MS1 scan across
the 300-1800 m/z range at 60,000 resolution, AGC target of $3x10^6$ and maximum injection time
of 20ms. MS2 scans of most abundant 12 ions were performed at 45,000 resolution with AGC
target of $1x10^5$ and maximum injection time of 100ms, isolation window of 0.7m/z and
normalized collision energy of 30.

Data searching

Proteome and phosphoproteome data were searched against a composite protein database at
the Bioinformatics Center (BIC) using the MSGF+ cloud-based pipeline previously described[82].
This database comprised UniProt canonical sequences (downloaded 2022-09-13; 20383
sequences), UniProt human protein isoforms (downloaded 2022-09-13; 21982 sequences), and
common contaminants (261 sequences), resulting in 42,626 sequences.

QC/Filtering/Normalization

The log2 Reporter ion intensity (RII) ratios to the common reference were used as quantitative
values for all proteomics features (proteins and phosphosites). Datasets were filtered to remove
features identified from contaminant proteins and decoy sequences. Datasets were visually
evaluated for sample outliers by looking at top principal components, examining median feature
abundance and distributions of RII ratio values across samples, and by quantifying the number
of feature identifications within each sample. No outliers were detected in the muscle tissue
dataset. In the adipose datasets (proteome and phosphoproteome), four samples were flagged
as outliers based on inspection of the top principal components and evaluation of the raw
quantitative results (median feature abundance <-1.0). The Log2 RII ratio values were
normalized within each sample by median centering to zero. Principal Component Analysis and
calculation of the variance explained by variables of interest were used to identify and quantify
potential batch effects. Based on this analysis, TMT plex, and Clinical Site batch effects were
removed using Linear Models Implemented in the limma::RemoveBatchEffect() function in R[79].
A design matrix including age, sex, and group_timepoint was used during batch effect removal
in order to preserve the effect of all variables included in later statistical analysis. Correlations
between technical replicates analyzed within and across CAS (where applicable) were
calculated to evaluate intra- and inter-site reproducibility; the data from technical replicates were
then averaged for downstream analysis. Finally, features with quantification in less of 30% of all
samples were removed. For specific details of the process, see available code.

Metabolomics and Lipidomics

The metabolomic analysis was performed by investigators across multiple Chemical Analysis
Sites that employed different technical platforms for data acquisition. At the highest level, these
platforms were divided into two classes: Targeted and Untargeted. The data generated by the
untargeted platforms were further divided into Named (confidently identified chemical entities)
and Unnamed (confidently detected, but no chemical name annotation) compounds.

Targeted metabolomics data were generated at 3 analysis Sites: Duke University, the Mayo
Clinic, and Emory University. Duke quantified metabolites belonging to the metabolite classes
Acetyl-CoA, Keto Acids, and Nucleic Acids ("acoa", "ka", "nuc"), Mayo quantified amines and
TCA intermediates ("amines" and "tca"), while Emory quantified Oxylipins ("oxylipneg").

The untargeted metabolomics data were generated at 3 analysis Sites: the Broad Institute, the
University of Michigan, and the Georgia Institute of Technology (Georgia Tech). The Broad
Institute applied Hydrophilic interaction liquid chromatography (HILIC) in the positive ion mode
(hilicpos), Michigan applied reverse phase liquid chromatography in both positive and negative
ion modes ("rppos" and "rpneg") and ion-pairing chromatography in the negative mode
("ionpneg"), and Georgia Tech performed lipidomics assays using reverse phase
chromatography in both positive and negative ion modes ("lrppos" and "lrpneg").

LC-MS/MS analysis of branched-chain keto acids

Duke University conducted targeted profiling of branched-chain keto acid metabolites. Plasma
samples containing isotopically labeled ketoleucine (KIC)-d3, ketoisovalerate (KIV)-13C5 (from
Cambridge Isotope Laboratories), and 3-methyl-2-oxovalerate (KMV)-d8 (from Toronto
Research Chemicals, Canada) internal standards, were subjected to deproteinization with 3M
perchloric acid. Tissue homogenates were prepared at 100 mg/ml in 3M perchloric acid and 200
µL was centrifuged.

Next, 200 µL of 25 M o-phenylenediamine (OPD) in 3M HCl were added to both the plasma and
tissue supernatants. The samples were incubated at 80°C for 20 minutes. Keto acids were then
extracted using ethyl acetate, following a previously described protocol[83,84]. The extracts were
dried under nitrogen, reconstituted in 200 mM ammonium acetate, and subjected to analysis on
a Xevo TQ-S triple quadrupole mass spectrometer coupled to an Acquity UPLC (Waters),
controlled by the MassLynx 4.1 operating system.

The analytical column used was a Waters Acquity UPLC BEH C18 Column (1.7 µm, 2.1 × 50
mm), maintained at 30°C. 10 µL of the sample were injected onto the column and eluted at a
flow rate of 0.4 ml/min. The gradient consisted of 45% mobile phase A (5 mM ammonium
acetate in water) and 55% mobile phase B (methanol) for 2 minutes. This was succeeded by a
linear gradient to 95% B from 2 to 2.5 minutes, holding at 95% B for 0.7 minutes, returning to 45%
A, and finally re-equilibrating the column at initial conditions for 1 minute. The total run time was
4.7 minutes.

In positive ion mode, mass transitions of m/z 203 → 161 (KIC), 206 → 161 (KIC-d3), 189 → 174
(KIV), 194 → 178 (KIV-5C13), 203 → 174 (KMV), and 211 → 177 (KMV-d8) were monitored.
Endogenous keto acids were quantified using calibrators prepared by spiking dialyzed fetal
bovine serum with authentic keto acids (Sigma-Aldrich).

Flow injection MS/MS analysis of acyl CoAs

Duke University conducted targeted profiling of acyl CoAs. Here, 500 µL of tissue homogenate,
prepared at a concentration of 50 mg/ml in isopropanol/0.1 M KH2PO4 (1:1), underwent

extraction with an equal volume of acetonitrile. The resulting mixture was then centrifuged at
14,000 x g for 10 minutes, following a previously established procedure[85,86].

The supernatants were acidified with 0.25 ml of glacial acetic acid, and the acyl CoAs were
subjected to additional purification through solid-phase extraction (SPE) using 2-(2-pyridyl) ethyl
functionalized silica gel (Sigma-Aldrich), as outlined in Minkler et al. (2008)[87]. Prior to use, the
SPE columns were conditioned with 1 ml of acetonitrile/isopropanol/water/glacial acetic acid
(9/3/4/4 : v/v/v/v). After application and flow-through of the supernatant, the SPE columns
underwent a washing step with 2 ml of acetonitrile/isopropanol/water/glacial acetic acid (9/3/4/4 :
v/v/v/v). Acyl CoAs elution was achieved with 2 ml of methanol/250 mM ammonium formate
(4/1 : v/v), followed by analysis using flow injection MS/MS in positive ion mode on a Xevo TQ-S
triple quadrupole mass spectrometer (Waters). The mobile phase used was methanol/water
(80/20, v/v) with 30 mM ammonium hydroxide. Spectra were acquired in the multichannel
acquisition mode, monitoring the neutral loss of 507 amu (phosphoadenosine diphosphate) and
scanning from m/z 750 to 1100. As an internal standard, heptadecanoyl CoA was employed.
Quantification of endogenous acyl CoAs was carried out using calibrators created by spiking
tissue homogenates with authentic Acyl CoAs (Sigma-Aldrich) that covered saturated acyl chain
lengths from C0 to C18. Empirical corrections for heavy isotope effects, particularly 13C, to the
adjacent m+2 spectral peaks in a specific chain length cluster were made by referring to the
observed spectra for the analytical standards.

LC-MS/MS analysis of nucleotides

Duke University conducted targeted profiling of nucleotide metabolites. 300 µL of tissue
homogenates, prepared at a concentration of 50 mg/ml in 70% methanol, underwent spiking
with nine internal standards: 13C10,15N5-adenosine monophosphate, 13C10,15N5-guanosine
monophosphate, 13C10,15N2-uridine monophosphate, 13C9,15N3-cytidine monophosphate,
13C10-guanosine triphosphate, 13C10-uridine triphosphate, 13C9-cytidine triphosphate,
13C10-adenosine triphosphate, and nicotinamide-1, N6-ethenoadenine dinucleotide (eNAD) (all
from Sigma-Aldrich). Nucleotides were extracted using an equal volume of hexane, following the
procedures previously outlined[88,89]. After vortexing and centrifugation at 14,000 x g for 5 minutes,
the bottom layer was subjected to another centrifugation. Chromatographic separation and MS
analysis of the supernatants were performed using an Acquity UPLC system (Waters) coupled
to a Xevo TQ-XS quadrupole mass spectrometer (Waters). The analytical column employed
was a Chromolith FastGradient RP-18e 50-2mm column (EMD Millipore, Billerica, MA, USA),
that was maintained at 40°C. The injection volume was 2 µL. Nucleotides were separated using
a mobile phase A consisting of 95% water, 5% methanol, and 5 mM dimethylhexylamine
adjusted to pH 7.5 with acetic acid. Mobile phase B comprised 20% water, 80% methanol, and
10 mM dimethylhexylamine. The flow rate was set to 0.3 ml/min. The 22-minute gradient
(t=0, %B=0; t=1.2, %B=0; t=22, %B=40) was followed by a 3-minute wash and 7-minute
equilibration. Nucleotides were detected in the negative ion multiple reaction monitoring (MRM)
mode based on characteristic fragmentation reactions. Endogenous nucleotides were quantified
using calibrators prepared by spiked-in authentic nucleotides obtained from Sigma-Aldrich.

LC-MS/MS analysis of amino metabolites

The Mayo Clinic conducted LC-MS-based targeted profiling of amino acids and amino
metabolites, as previously outlined[90,91]. In summary, either 20 ml of plasma samples or 5 mg of
tissue homogenates were supplemented with an internal standard solution comprising
isotopically labeled amino acids (U-13C4 L-aspartic acid, U-13C3 alanine, U-13C4 L-threonine,
U-13C L-proline, U-13C6 tyrosine, U-13C5 valine, U-13C6 leucine, U-13C6 phenylalanine, U-
13C3 serine, U-13C5 glutamine, U-13C2 glycine, U-13C5 glutamate, U-13C6,15N2 lysine, U-
13C5,15N methionine, 1,1U-13C2 homocysteine, U-13C6 arginine, U-13C5 ornithine, 13C4
asparagine, 13C2 ethanolamine, D3 sarcosine, D6 4-aminobutyric acid). The supernatant was
promptly derivatized using 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate with a MassTrak
kit (Waters). A 10-point calibration standard curve underwent a similar derivatization procedure
after the addition of internal standards. Both the derivatized standards and samples were
subjected to analysis using a Quantum Ultra triple quadrupole mass spectrometer
(ThermoFischer) coupled with an Acquity liquid chromatography system (Waters). Data
acquisition utilized selected ion monitoring (SRM) in positive ion mode. The concentrations of 42
analytes in each unknown sample were calculated against their respective calibration curves.

GC-MS analysis of TCA metabolites

The Mayo Clinic conducted GC-MS-targeted profiling of tricarboxylic acid (TCA) metabolites, as
previously detailed[92,93], with some modifications. In summary, 5 mg of tissue were homogenized
in 1X PBS using an Omni bead ruptor (Omni International, Kennesaw, GA), followed by the
addition of 20 µL of an internal solution containing U-13C labeled analytes (13C3 sodium lactate,
13C4 succinic acid, 13C4 fumaric acid, 13C4 alpha-ketoglutaric acid, 13C4 malic acid, 13C4
aspartic acid, 13C5 2-hydroxyglutaric acid, 13C5 glutamic acid, 13C6 citric acid, 13C2,15N
glycine, 13C2 sodium pyruvate). For plasma, 50 µL were used. The proteins were precipitated
out using 300 µL of a mixture of chilled methanol and acetonitrile solution. After drying the
supernatant in the speedvac, the sample was derivatized first with ethoxime and then with
MtBSTFA + 1% tBDMCS (N-Methyl-N-(t-Butyldimethylsilyl)-Trifluoroacetamide + 1% t-
Butyldimethylchlorosilane) and then analyzed on an Agilent 5977B GC/MS (Santa Clara,
California) under single ion monitoring conditions using electron ionization. Concentrations of
lactic acid (m/z 261.2), fumaric acid (m/z 287.1), succinic acid (m/z 289.1), ketoglutaric acid
(m/z 360.2), malic acid (m/z 419.3), aspartic acid (m/z 418.2), 2-hydroxyglutaratic acid (m/z
433.2), cis-aconitic acid (m/z 459.3), citric acid (m/z 591.4), isocitric acid (m/z 591.4), and
glutamic acid (m/z 432.4) were measured against 7-point calibration curves that underwent the
same derivatization procedure.

Targeted lipidomics of low-level lipids

Lipid targeted profiling was conducted at Emory University following established methodologies
as previously described[94,95]. In summary, 20 mg of powdered ASAT samples were
homogenized in 100 µL PBS using Bead Ruptor (Omni International, Kennesaw, GA).
Homogenized tissue samples (or plasma samples) were diluted with 300 µL 20% methanol and
spiked with a 1% BHT solution to a final BHT concentration of 0.1% and pH of 3.0 by acetic acid
addition. After centrifugation (10 minutes, 14000 rpm), the supernatants were transferred to 96-
well plates for further extraction. The supernatants were loaded onto Isolute C18 solid-phase

extraction (SPE) columns that had been conditioned with 1000 µL ethyl acetate and 1000 µL 5%
methanol. The SPE columns were washed with 800 µL water and 800 µL hexane, and the
oxylipins were then eluted with 400 µL methyl formate. The SPE process was automated using
a Biotage Extrahera (Uppsala, Sweden). The eluate was dried with nitrogen and reconstituted
with 200 µL acetonitrile before LC-MS analysis. Sample blanks, pooled extract samples used as
quality controls (QC), and consortium reference samples were prepared for analysis using the
same methods. All external standards were purchased from Cayman Chemical (Ann Arbor,
Michigan) at a final concentration in the range 0.01-20 µg/ml and consisted of: prostaglandin E2
ethanolamide (Catalog No. 100007212), oleoyl ethanolamide (Catalog No. 90265), palmitoyl
ethanolamide (Catalog No. 10965), arachidonoyl ethanolamide (Catalog No. 1007270),
docosahexaenoyl ethanolamide (Catalog No. 10007534), linoleoyl ethanolamide (Catalog No.
90155), stearoyl ethanolamide (Catalog No. 90245), oxy-arachidonoyl ethanolamide (Catalog
No. 10008642), 2-arachidonyl glycerol (Catalog No. 62160), docosatetraenoyl ethanolamide
(Catalog No. 90215), α-linolenoyl ethanolamide (Catalog No. 902150), oleamide (Catalog No.
90375), dihomo-γ-linolenoyl ethanolamide (Catalog No. 09235), decosanoyl ethanolamide
(Catalog No. 10005823), 9,10 DiHOME (Catalog No. 53400), prostaglandin E2-1-glyceryl ester
(Catalog No. 14010), 20-HETE (Catalog No. 10007269), 9-HETE (Catalog No. 34400), 14,15
DiHET (Catalog No. 10007267), 5(S)-HETE (Catalog No. 34210), 12(R)-HETE (Catalog No.
10007247), 11(12)-DiHET (Catalog No. 10007266), 5,6-DiHET (Catalog No. 10007264),
thromboxane B2 (Catalog No. 10007237), 12(13)-EpOME (Catalog No. 52450), 13 HODE
(Catalog No. 38600), prostaglandin F2α (Catalog No. 10007221), 14(15)-EET (Catalog No.
10007263), 8(9)-EET (Catalog No. 10007261), 11(12)-EET (Catalog No. 10007262), leukotriene
B4 (Catalog No. 20110), 8(9)-DiHET (Catalog No. 10007265), 13-OxoODE (Catalog No. 38620),
13(S)-HpODE (Catalog No. 48610), 9(S)-HpODE (Catalog No. 48410), 9(S)-HODE (Catalog No.
38410), resolvin D3 (Catalog No. 13834), resolvin E1 (Catalog No. 10007848), resolvin D1
(Catalog No. 10012554), resolvin D2 (Catalog No. 10007279), 9(S)HOTrE (Catalog No. 39420),
13(S)HOTrE (Catalog No. 39620), 8-iso Progstaglandin F2α (Catalog No. 25903), maresin 1
(Catalog No. 10878), maresin 2 (Catalog No. 16369).

LC-MS data were acquired using an Agilent 1290 Infinity II chromatograph from Agilent (Santa
Clara, CA), equipped with a ThermoFisher Scientific AccucoreTM C18 column (100 mm × 4.6,
2.6 µm particle size), and coupled to Agilent 6495 mass spectrometer for polarity switch multiple
reaction monitoring (MRM) scan. The mobile phases consisted of water with 10 mM ammonium
acetate (mobile phase A) and acetonitrile with 10 mM ammonium acetate (mobile phase B). The
chromatographic gradient program was: 0.5 minutes with 95% A; 1 minute to 2 minutes with 65%
A; 2.1 minutes to 5.0 minutes with 45% A; 7 minutes to 20 minutes with 25% A; and 21.1
minutes until 25 minutes with 95% A. The flow rate was set at 0.40 ml/min. The column
temperature was maintained at 35°C, and the injection volume was 6 µL. For MS analysis, the
MRM analysis is operated at gas temperature of 290 °C, gas flow of 14 L/min, Nebulizer of 20
psi, sheath gas temperature of 300 °C, sheath gas flow of 11 L /min, capillary of 3000 V for both
positive and negative ion mode, nozzle voltage of 1500 V for both positive and negative ion
mode, high pressure RF of iFunnel parameters of 150V for both positive and negative ion mode,
and low pressure of RF of iFunnel parameters of 60 V for both positive and negative ion mode.
Skyline (version 25.1.0.142)154 was utilized to process raw LC-MS data. Standard curves were

constructed for each oxylipin/ethanolamide and scrutinized to ensure that all concentration
points fell within the linear portion of the curve with an R-squared value not less than 0.9.
Additionally, features exhibiting a high coefficient of variation (CV) among the quality control
(QC) samples were eliminated from the dataset. Pearson correlation among the QCs for each
tissue type was computed using the Hmisc R library, and the figures documented in the QC
report were generated and visualized with the corrplot R library[96,97].

Hydrophilic interaction LC-MS metabolomics

The untargeted analysis of polar metabolites in the positive ionization mode was conducted at
the Broad Institute of MIT and Harvard. The LC-MS system consisted of a Shimadzu Nexera X2
UHPLC (Shimadzu Corp., Kyoto, Japan) coupled to a Q-Exactive hybrid quadrupole Orbitrap
mass spectrometer (Thermo Fisher Scientific). Plasma samples (10 µL) were extracted using 90
µL of 74.9/24.9/0.2 v/v/v acetonitrile/methanol/formic acid containing valine-d8 and
phenylalanine-d8 internal standards. Following centrifugation, the supernatants were directly
injected onto a 150 x 2 mm, 3 µm Atlantis HILIC column (Waters). Tissue (10 mg)
homogenization was performed at 4°C using a TissueLyser II (QAIGEN) bead mill set to two 2
min intervals at 20 Hz in 300 µL of 10/67.4/22.4/0.018 v/v/v/v water/acetonitrile/methanol/formic
acid valine-d8 and phenylalanine-d8 internal standards. The column was eluted isocratically at a
flow rate of 250 µL/min with 5% mobile phase A (10 mM ammonium formate and 0.1% formic
acid in water) for 0.5 minute, followed by a linear gradient to 40% mobile phase B (acetonitrile
with 0.1% formic acid) over 10 minutes, then held at 40% B for 4.5 minutes. MS analyses
utilized electrospray ionization in the positive ion mode, employing full scan analysis over 70-
800 m/z at 70,000 resolution and 3 Hz data acquisition rate. Various MS settings, including
sheath gas, auxiliary gas, spray voltage, and others, were specified for optimal performance.
Data quality assurance was performed by confirming LC-MS system performance with a mixture
of >140 well-characterized synthetic reference compounds, daily evaluation of internal standard
signals, and the analysis of four pairs of pooled extract samples per sample type inserted in the
analysis queue at regular intervals. One sample from each pair was used to correct for
instrument drift using "nearest neighbor" scaling while the second reference sample served as a
passive QC for determination of the analytical coefficient of variation of every identified
metabolite and unknown. Raw data processing involved the use of TraceFinder software (v3.3,
Thermo Fisher Scientific) for targeted peak integration and manual review, as well as
Progenesis QI (v3.0, Nonlinear Dynamics, Waters) for peak detection and integration of both
identified and unknown metabolites. Metabolite identities were confirmed using authentic
reference standards.

Reversed Phase-High Performance & Ion-pairing LC metabolomics

Reversed-phase and ion pairing LC-MS profiling of polar metabolites was conducted at the
University of Michigan. LC-MS grade solvents and mobile phase additives were procured from
Sigma-Aldrich, while chemical standards were obtained from either Sigma-Aldrich or Cambridge
Isotope Labs. Plasma (50 µL aliquot) was extracted by addition of 200 µL of 1:1:1 v:v
methanol:acetonitrile:acetone containing the following internal standards diluted from stock
solutions to yield the specified concentrations: D4-succinic acid, 12.5 µM; :D3-malic acid,12.5

µM; D5 -glutamic acid,12.5 µM; D10 -leucine,12.5 µM; D5-tryptophan,12.5 µM;D5-
phenylalanine,12.5 µM; D3-caffeine,12.5 µM; D8-lysine,12.5 µM; D4-chenodeoxycholic
acid,12.5 µM; phosphatidylcholine(17:0/17:0),2.5 µM; phosphatidylcholine(19:0/19:0),2.5 µM;
D31-palmitic acid,12.5 µM; D35-stearic acid,12.5 µM; gibberellic acid, 2.5 µM; epibrassinolide,
2.5 µM.   The extraction solvent also contained a 1:400 dilution of Cambridge Isotope
carnitine/acylcarnitine mix NSK-B, resulting in the following concentrations of
carnitine/acylcarnitine internal standards: D9-L-carnitine, 380 nM; D3-L-acetylcarnitine, 95 nM;
D3-L-propionylcarnitine, 19 nM; D3-L-butyrylcarnitine, 19 nM; D9-L-isovalerylcarnitine, 19 nM;
D3-L-octanoylcarnitine, 19 nM; D9-L-myristoylcarnitine, 19 nM; D3-L-palmitoylcarnitine, 38 nM.
After addition of solvent, samples were vortexed, incubated on ice for 10 minutes, and then
centrifuged at 15,000 x g for 5 minutes. 150 µL supernatant were retrieved from the pellet,
transferred to a glass autosampler vial with a 250 uL footed insert, and dried under a constant
stream of nitrogen gas at ambient temperature. A QC sample was created by pooling residual
supernatant from multiple samples. This QC sample underwent the same drying and
reconstitution process as described for individual samples. Dried supernatants were stored at -
80 C until ready for instrumental analysis. On the day of analysis, dried samples were
reconstituted in 37.5 µL of 8:2 v:v water:methanol, vortexed thoroughly, and submitted for LC-
MS.

Frozen ASAT was rapidly weighed into pre-tared, pre-chilled Eppendorf tubes and extracted in
1:1:1:1 v:v methanol:acetonitrile:acetone:water containing the following internal standards
diluted from stock solutions to yield the specified concentrations: 13C3-lactic acid, 12.5 µM;
13C5-oxoglutaric acid, 125 nM;13C5-citric acid,1.25 µM; 13C4-succinic acid,125 nM; 13C4-
malic acid, 125 nM; U-13C amino acid mix (Cambridge Isotope CLM-1548-1), 2.5 µg/mL; 13C5-
glutamine, 6.25 µM;15N2-asparagine, 1.25 µM;15N2-tryptophan, 1.25 µM; 13C6-glucose, 62.5
µM; D4-thymine, 1 µM; 15N-anthranillic acid, 1 µM; gibberellic acid,1 µM; epibrassinolide, 1 µM.
The extraction solvent also contained a 1:400 dilution of Cambridge Isotope
carnitine/acylcarnitine mix NSK-B, resulting in the following concentrations of
carnitine/acylcarnitine internal standards: D9-L-carnitine, 380 nM; D3-L-acetylcarnitine, 95 nM;
D3-L-propionylcarnitine, 19 nM; D3-L-butyrylcarnitine, 19 nM; D9-L-isovalerylcarnitine, 19 nM;
D3-L-octanoylcarnitine, 19 nM; D9-L-myristoylcarnitine, 19 nM; D3-L-palmitoylcarnitine, 38 nM.
Sample extraction was performed by adding chilled extraction solvent to tissue sample at a ratio
of 1 ml solvent to 50 mg wet tissue mass. Immediately after solvent addition, the sample was
homogenized using a Branson 450 probe sonicator. Subsequently, the tubes were mixed
several times by inversion and then incubated on ice for 10 minutes. Following incubation, the
samples were centrifuged and 300 µL of the supernatant was carefully transferred to two
autosampler vials with flat-bottom inserts and dried under a constant stream of nitrogen gas. A
QC sample was created by pooling residual supernatants from multiple samples. This QC
sample underwent the same drying and reconstitution process as described for individual
samples. Dried supernatants were stored at -80 C until ready for instrumental analysis. On the
day of analysis, samples were reconstituted in 60 µL of 8:2 v:v water:methanol and then
submitted to LC-MS.

Reversed phase LC-MS samples were analyzed on an Agilent 1290 Infinity II / 6545 qTOF MS
system with a JetStream electrospray ionization (ESI) source (Agilent Technologies, Santa

Clara, California) using a Waters Acquity HSS T3 column, 1.8 µm 2.1 x 100 mm equipped with a
matched Vanguard precolumn (Waters Corporation). Mobile phase A was 100% water with 0.1%
formic acid and mobile phase B was 100% methanol with 0.025% formic acid. The gradient was
as follows: Linear ramp from 0% to 100% B from 0-10 minutes, hold 100% B until 17 minutes,
linear return to 0% B from 17 to 17.1 minutes, hold 0% B until 20 minutes. The flow rate was
0.45 ml/min, the column temperature was 55°C, and the injection volume was 5 µL. Each
sample was analyzed twice, once in positive and once in negative ion mode MS, scan rate 2
spectra/sec, mass range 50-1200 m/z. Source parameters were: drying gas temperature 350°C,
drying gas flow rate 10 L/min, nebulizer pressure 30 psig, sheath gas temperature 350°C and
flow 11 L/minute, capillary voltage 3500 V, internal reference mass correction enabled. A QC
sample run was performed at minimum every tenth injection.

Ion-pair LC-MS samples were analyzed on an identically-configured LC-MS system using an
Agilent Zorbax Extend C18 1.8 µm RRHD column, 2.1 x 150 mm ID, equipped with a matched
guard column. Mobile phase A was 97% water, 3% methanol. Mobile phase B was 100%
methanol. Both mobile phases contained 15 mM tributylamine and 10 mM acetic acid. Mobile
phase C was 100% acetonitrile. Elution was carried out using a linear gradient followed by a
multi-step column wash including automated (valve-controlled) backflushing, detailed as follows:
0-2min, 0%B; 2-11 min, linear ramp from 0-99%B; 12-16 min, 99%B, 16-17.5min, 99-0%B. At
17.55 minutes, the 10-port valve was switched to reverse flow (back-flush) through the column.
From 17.55-20.45 min the solvent was ramped from 99%B to 99%C. From 20.45-20.95 min the
flow rate was ramped up to 0.8 mL/min, which was held until 22.45 min, then ramped down to
0.6mL/min by 22.65 min. From 22.65-23.45 min the solvent was ramped from 99% to 0% C
while flow was simultaneously ramped down from 0.6-0.4mL/min. From 23.45 to 29.35 min the
flow was ramped from 0.4 to 0.25mL/min; the 10-port valve was returned to restore forward flow
through the column at 25 min. . Column temperature was 35°C and the injection volume was 5
µL. MS acquisition was performed in negative ion mode, scan rate 2 spectra/sec, mass range
50-1200 m/z. Source parameters were: drying gas temperature 250°C, drying gas flow rate 13
L/min, nebulizer pressure 35 psig, sheath gas temp 325°C and flow 12 L/min, capillary voltage
3500V, internal reference mass correction enabled. A QC sample run was performed at
minimum every tenth injection.

Iterative MS/MS data was acquired for both reverse phase and ion pairing methods using the
pooled sample material to enable compound identification. Eight repeated LC-MS/MS runs of
the QC sample were performed at three different collision energies (10, 20, and 40) with
iterative MS/MS acquisition enabled. The software excluded precursor ions from MS/MS
acquisition within 0.5 minute of their MS/MS acquisition time in prior runs, resulting in deeper
MS/MS coverage of lower-abundance precursor ions.

Feature detection and alignment was performed utilizing a hybrid targeted/untargeted approach.
Targeted compound detection and relative quantitation was performed by automatic integration
followed by manual inspection and correction using Profinder v8.0 (Agilent Technologies, Santa
Clara, CA.) Non-targeted feature detection was performed using custom scripts that automate
operation of the "find by molecular feature" workflow of the Agilent Masshunter Qualitative
Analysis (v7) software package. Feature alignment and recursive feature detection were

performed using Agilent Mass Profiler Pro (v8.0) and Masshunter Qualitative Analysis ("find by
formula" workflow), yielding an aligned table including m/z, RT, and peak areas for all features.

Data Cleaning and Degeneracy Removal: The untargeted features and named metabolites were
merged to generate a combined feature table. Features missing from over 50% of all samples in
a batch or over 30% of QC samples were then removed prior to downstream normalization
procedure. Next, the software package Binner was utilized to remove redundancy and
degeneracy in the data[98]. Briefly, Binner first performs RT-based binning, followed by clustering
of features by Pearson's correlation coefficient, and then assigns annotations for isotopes,
adducts or in-source fragments by searching for known mass differences between highly
correlated features.

Normalization and Quality Control: Data were then normalized using a Systematic Error
Removal Using Random Forest (SERRF) approach[99], which helps correct for drift in peak
intensity over the batch using data from the QC sample runs. When necessary to correct for
residual drift, peak area normalization to closest-matching internal standard was also applied to
selected compounds. Both SERRF correction and internal standard normalization were
implemented in R. Parameters were set to minimize batch effects and other observable drifts,
as visualized using principal component analysis score plots of the full dataset. Normalization
performance was also validated by examining relative standard deviation values for additional
QC samples not included in the drift correction calculations.

Compound identification: Metabolites from the targeted analysis workflow were identified with
high confidence (MSI level 1)[100] by matching retention time (+/- 0.1 minute), mass (+/- 10 ppm)
and isotope profile (peak height and spacing) to authentic standards. MS/MS data
corresponding to unidentified features of interest from the untargeted analysis were searched
against a spectral library (NIST 2020 MS/MS spectral database or other public spectral
databases) to generate putative identifications (MSI level 2) or compound-class level
annotations (MSI level 3) as described previously[101].

LC-MS/MS untargeted lipidomics

Sample preparation: Non-targeted lipid analysis was conducted at the Georgia Institute of
Technology. Powdered tissue samples (10 mg) were extracted in 400 µL isopropanol containing
stable isotope-labeled internal standards (IS) with bead homogenization with 2mm zirconium
oxide beads (Next Advance) using a TissueLyser II (10min, 30 Hz). Samples were then
centrifuged (5 min, 21,100xg), and supernatants were transferred to autosampler vials. Plasma
samples (25 µL) were extracted by mixing with 75 µL isopropanol containing the IS mix followed
by centrifugation. Sample blanks, pooled extract samples used as quality controls (QC), and
consortium reference samples, were prepared for analysis using the same methods. The IS mix
consisted of PC (15:0-18:1(d7)), Catalog No. 791637; PE (15:0-18:1(d7)), Catalog No. 791638;
PS (15:0-18:1(d7)), Catalog No. 791639; PG(15:0-18:1(d7)), Catalog No. 791640; PI(15:0-
18:1(d7)), Catalog No. 791641; LPC(18:1(d7)), Catalog No. 791643; LPE(18:1(d7)); Catalog No.
791644; Chol Ester (18:1(d7)), Catalog No. 700185; DG(15:0-18:1(d7)), Catalog No. 791647;
TG(15:0-18:1(d7)-15:0), Catalog No. 791648; SM(18:1(d9)), Catalog No. 791649; Cholesterol

(d7), Catalog No. 700041. All internal standards were purchased from Avanti Polar Lipids
(Alabaster, Alabama) and added to the extraction solvent at a final concentration in the 0.1-8
µg/ml range.

Data collection: Lipid LC-MS data were acquired using a Vanquish (ThermoFisher Scientific)
chromatograph fitted with a ThermoFisher Scientific AccucoreTM C30 column (2.1 × 150 mm,
2.6 µm particle size), coupled to a high-resolution accurate mass Q-Exactive HF Orbitrap mass
spectrometer (ThermoFisher Scientific) for both positive and negative ionization modes. For
positive mode analysis, the mobile phases were 40:60 water:acetonitrile with 10 mM ammonium
formate and 0.1% formic acid (mobile phase A), and 10:90 acetonitrile:isopropyl alcohol, with 10
mM ammonium formate and 0.1% formic acid (mobile phase B). For negative mode analysis,
the mobile phases were 40:60 water:acetonitrile with 10 mM ammonium acetate (mobile phase
A), and 10:90 acetonitrile:isopropyl alcohol, with 10 mM ammonium acetate (mobile phase B).
The chromatographic method used for both ionization modes was the following gradient
program: 0 minutes 80% A; 1 minute 40% A; 5 minutes 30% A; 5.5 minutes 15% A; 8 minutes
10% A; held 8.2 minutes to 10.5 minutes 0% A; 10.7 minutes 80% A; and held until 12 minutes.
The flow rate was set at 0.40 ml/min. The column temperature was set to 50°C, and the
injection volume was 2 µL.

For analysis of the organic phase the electrospray ionization source was operated at a vaporizer
temperature of 425°C, a spray voltage of 3.0 kV for positive ionization mode and 2.8 kV for
negative ionization mode, sheath, auxiliary, and sweep gas flows of 60, 18, and 4 (arbitrary
units), respectively, and capillary temperature of 275°C. The instrument acquired full MS data
with 240,000 resolution over the 150-2000 m/z range. LC-MS/MS experiments were acquired
using a DDA strategy. MS2 spectra were collected with a resolution of 120,000 and the dd-MS2
were collected at a resolution of 30,000 and an isolation window of 0.4 m/z with a loop count of
top 7. Stepped normalized collision energies of 10%, 30%, and 50% fragmented selected
precursors in the collision cell. Dynamic exclusion was set at 7 seconds and ions with charges
greater than 2 were omitted.

Data processing: Data processing steps included peak detection, spectral alignment, grouping
of isotopic peaks and adduct ions, drift correction, and gap filling. Compound Discoverer V3.3
(ThermoFisher Scientific) was used to process the raw LC-MS data. Drift correction was
performed on each individual feature, where a Systematic Error Removal using Random Forest
(SERRF) method builds a model using the pooled QC sample peak areas across the batch and
was then used to correct the peak area for that specific feature in the samples. Detected
features were filtered with background and QC filters. Features with abundance lower than 5x
the background signal in the sample blanks and that were not present in at least 50% of the QC
pooled injections with a coefficient of variance (CV) lower than 80% (not drift corrected) and 50%
(drift corrected) were removed from the dataset. Lipid annotations were accomplished based on
accurate mass and relative isotopic abundances (to assign elemental formula), retention time
(to assign lipid class), and MS2 fragmentation pattern matching to local spectral databases built
from curated experimental data. Lipid nomenclature followed that described previously[102].

Quality control procedures: System suitability was assessed prior to the analysis of each batch.
A performance baseline for a clean instrument was established before any experiments were
conducted. The mass spectrometers were mass calibrated, mass accuracy and mass resolution
were checked to be within manufacturer specifications, and signal-to-noise ratios for the suite of
IS checked to be at least 75% of the clean baseline values. For LC-MS assays, an IS mix
consisting of 12 standards was injected to establish baseline separation parameters for each
new column. The performance of the LC gradient was assessed by inspection of the column
back pressure trace, which had to be stable within an acceptable range (less than 30% change).
Each IS mix component was visually evaluated for chromatographic peak shape, retention time
(lower than 0.2 minute drift from baseline values) and FWHM lower than 125% of the baseline
measurements. The CV of the average signal intensity and CV of the IS (<=15%) in pooled
samples were also checked. These pooled QC samples were used to correct for instrument
sensitivity drift over the various batches using a procedure similar to that described by the
Human Serum Metabolome (HUSERMET) Consortium. To evaluate the quality of the data for
the samples themselves, the IS signals across the batch were monitored, PCA modeling for all
samples and features before and after drift correction was conducted, and Pearson correlations
calculated between each sample and the median of the QC samples.

Metabolomics/Lipidomics Data filtering and normalization

The untargeted metabolomics datasets were categorized as either "named", for chemical
compounds confidently identified, or "unnamed", for compounds with specific chemical
properties but without a standard chemical name. While the preprocessing steps were
performed on the named and unnamed portions together, only the named portions were utilized
for differential analysis. For each dataset i.e. each assay for each tissue type, the following
steps are performed:

Average rows that have the same metabolite ID.

Merge the "named" and "unnamed" subparts of the untargeted datasets.

Convert negative and zero values to NAs.

Remove features with > 20% missing values.

Features with < 20% missing values are imputed either using K-Nearest Neighbor imputation
(for datasets with > 12 features) or half-minimum imputation (for datasets with < 12 features).

All data are log2-transformed, and the untargeted data are median-MAD normalized if neither
sample medians nor upper quartiles were significantly associated with sex or sex-stratified
training group (Kruskal-Wallis p-value < 0.01). Note that for all log2 calculations, 1 is added to
each value before log-normalization. This allows metabolite values that fall between 0 and 1 to
have a positive log2 value.

Outlier detection was performed by examining the boxplot of each Principal Component and
extending its whiskers to the predefined multiplier above and below the interquartile range (5x

the IQR). Samples outside this range are flagged. All outliers were reviewed by Metabolomics
CAS, and only confirmed technical outliers were removed.

Redundant Metabolite/Lipid Management

To address metabolites measured on multiple platforms (e.g. a metabolite measured on the
HILIC positive and RP positive platforms), and metabolites with the same corresponding RefMet
ID (e.g. alpha-Aminoadipic-acid, Aminoadipic acid both correspond to RefMet name
'Aminoadipic acid'), we utilize the set of internal standards described above to make a decision
on which platform's measurement of a given feature to include in further analysis. For each
tissue, based on whichever platform has the lowest coefficient of variation across all included
reference standards for a given refmet id, that metabolite was chosen. The other platforms or
metabolites, for this tissue, corresponding to this refmet id were removed from further
downstream analysis. Data for all metabolites removed, including information about the
coefficient of variation in the standards, normalized data, or differential analysis results, is
available in the R Package MotrpacHumanPreSuspensionData, but is not loaded by default.

Statistical analysis

Differential analysis

To model the effects of both exercise modality and time, relative to non exercising control, each
measured molecular feature was treated as an outcome in a linear mixed effects model
accounting for fixed effects of exercise group (RE, EE, or CON), timepoint, as well as
demographic and technical covariates (see covariate selection below for more info). Participant
identification was treated as a random effect. For each molecular feature, a cell-means model is
fit to estimate the mean of each exercise group-timepoint combination, and all hypothesis tests
are done comparing the means of the fixed effect group-timepoint combinations. In order to
model the effects of exercise against non-exercising control, a difference-in-changes model was
used which compared the change from pre-exercise to a during or post-exercise timepoint in
one of the two exercise groups to the same change in control. This effect is sometimes referred
to as a "delta-delta" or "difference-in-differences" model.

Model selection via simulation

In order to determine the optimal statistical package/model for these data – given MoTrPAC's
unique sampling design (see 'acute exercise intervention' above) – a simulation study was
conducted to determine the impact of various approaches to account for correlation in repeated
measurements. The simulation evaluated type 1 error, power, and bias for multiple plausible
modeling approaches, to identify those that would obtain higher power while maintaining
nominal type 1 error rates. For a subset of molecular features in the datasets, mean, covariance,
skew, and kurtosis were summarized over the relevant timepoints in the acute exercise bout
within subgroups defined by sex and exercise type. Then, these summary values were used in
combination with the "covsim" R package to generate non-normally distributed simulated data[103].
Additionally, sample size and missingness patterns were aligned in the simulated data to match

those of the observed data. Using a total of 5 million simulated instances where data were
generated under the null hypothesis (i.e., no change in the mean values of the outcome over the
exercise timepoints) or alternative hypothesis (i.e., the mean value of the outcome differs
between at least two timepoints), type 1 error rate and power, respectively, were evaluated for
seven analytic strategies: (1) pairwise t-tests between timepoints of interest, (2) ordinary least
squares regression, (3) differential expression for repeated measures (i.e., dream) with random
intercepts,129 (4) linear mixed models with random intercepts, (5) mixed models for repeated
measures using an unstructured covariance matrix, and (6) generalized linear models using
generalized estimating equations (GEEs) and a first degree autoregressive correlation structure.
These analyses were replicated for adipose, blood, and muscle samples. The dream function
obtained type 1 error rates 5.2, 5.4, and 5.2% for adipose, blood, and muscle analyses,
respectively, with higher power than all other methods apart from generalized linear models with
GEEs. Although generalized linear models with GEEs had higher power than dream, they also
had higher type 1 error with 7.7%, 6.0%, and 6.5% type 1 error rates for adipose, blood, and
muscle analyses, respectively. Therefore, due to stability of type 1 error and relatively high
power compared to other approaches, the dream function from the variancePartition R package
was selected for the primary analyses for every omic platform except Methyl-Cap (see
methylation capture sequencing methods above for more details).129

Covariate selection

Covariates were selected through a combination of a priori knowledge about factors influencing
molecular levels as well through empiric screening. Factors were considered for model inclusion
by correlating the principal components of each tissue-ome feature set to demographic and
technical factors using variancePartition::canCorPairs[78]. Visualizations and computational
analysis that describe this process for the decisions for covariate selection can be found at:
https://github.com/MoTrPAC/precovid-analyses/tree/main/QC. Ultimately, the following fixed
effect covariates were included in the models for every omic platform, except Methyl Cap (see
methods above): group and timepoint in a cell means model, clinical site, age, sex, and BMI.
'Participant id' was included as a random effect in every model. Each ome then had ome-
specific covariates selected as described in prior sections.

A targeted analysis was conducted to assess whether and how race, ethnicity, and genetic
ancestry could be incorporated, given prior evidence that these factors can influence omic
data[104,105]. We evaluated multiple models that included patient-reported race and/or ethnicity, as
well as principal components derived from whole-genome sequencing (WGS), to determine their
impact on model fit using the BIC. However, no combinations of individual or joint inclusion of
race, ethnicity, or WGS-derived principal components led to an improvement in average model
BIC across all features for a platform.

The final set of covariates included in each model for each tissue and ome can be found in the
R Package MotrpacHumanPreSuspensionData.

Missingness

Data missingness varied for multiple reasons including experimental design (i.e. temporal
randomization which randomized some participants to have samples obtained at a subset of
timepoints. For some omic sets, imputation to alleviate missingness could be performed
(metabolomics) but for others was not, including proteomics. Analysis demonstrated proteomics
missingness was approximately at random (data not shown).

Given the above issues affecting the ultimate sample size for effect estimation, for a feature to
be included in the analysis, a minimum number of 3 participants were required to have a paired
pre-exercise sample for all groups and all during/post-exercise timepoints. Thus, all group
comparisons (e.g. RE vs CON at 3.5 hours post-exercise) required 3 participants with matched
pre- and post-exercise samples in each group. This requirement was satisfied in all omes and
tissues except a subset of MS-acquired proteomics and phosphoproteomics features. Features
not meeting this criterion in any group-timepoint set were excluded from differential analysis
entirely, though raw and QC-normalized values are available.

Specific omic-level statistical considerations

Models for all omes other than methylation were fit using 'variancePartition::dream'[78]. For the
transcriptomics and ATAC-seq, which are measured as numbers of counts, the mean-variance
relationship was measured using 'variancePartition::voomWithDreamWeights' as previously
described[78]. For all other omes the normalized values were used directly as input to the
statistical model.

The full implementation of the statistical models for each of non-methylation datasets can be
found in the R Package MotrpacHumanPreSuspensionData. Methylation data was processed
separately, and the methods can be found in the methylation methods section.

Contrast types

The specific contrasts made in the statistical analysis include three categories of comparison:
difference-in-changes relative to control, group-specific, and resistance vs endurance. All
comparisons were structured using 'variancePartition::makeContrastsDream'[78].

The difference-in-changes model as described at the opening of this section represents the
primary DA analysis, and any individual feature mentioned as statistically significantly changing
due to exercise will be referring to the difference-in-changes results unless otherwise
specifically stated.

The next category of contrasts is a group-specific comparison, which compares a given post-
intervention timepoint to pre-exercise within a given group, without comparing to the control
group. This contrast was implemented to more easily quantify effect sizes within each group
independently, but are not used as a general endpoint.

Finally, the RE vs EE is very similar to the difference-in-changes model, but sets the RE group
as a matched control to the EE group. In this comparison, the change in response to exercise at
a given timepoint in one exercise group is directly compared to the corresponding change in the

other exercise group. This contrast can directly test the hypothesis that the EE effect is different
from the RE effect. There are no samples obtained during RE bouts, so those timepoints in
blood were excluded from this comparison.

Significance thresholds

For each of the above contrasts, p-values were adjusted for multiple comparisons for each
unique contrast-group-tissue-ome-timepoint combination separately using the Benjamini-
Hochberg method to control False Discovery Rate (FDR)[106]. Features were considered
significant at a FDR of 0.05 unless otherwise stated.

Human feature to gene mapping

The feature-to-gene map links each feature tested in differential analysis to a gene, using
Ensembl version 105 (mapped to GENCODE 39)[107] as the gene identifier source. Proteomics
feature IDs (UniProt IDs) were mapped to gene symbols and Entrez IDs using UniProt's
mapping files[108]. Epigenomics features were mapped to the nearest gene using the
ChIPseeker::annotatePeak()[109,110] function with Homo sapiens Ensembl release 105 gene
annotations. Gene symbols, Entrez IDs, and Ensembl IDs were assigned to features using
biomaRt version 2.58.2 (Bioconductor 3.18)[111-113].

Metabolite features were mapped to KEGG IDs using KEGGREST, RefMet REST API, or web
scraping from the Metabolomics Workbench[112,114].

Enrichment analysis

Preparation of differential analysis results

The differential analysis results tables for each combination of tissue and ome were converted
to matrices of z-scores with either gene symbols, RefMet metabolite/lipid IDs, or
phosphorylation flanking sequences as rows and contrasts as columns. These matrices serve
as input for the enrichment analyses. For the transcriptomics, TMT proteomics, and Olink
proteomics results, transcripts and proteins were first mapped to gene symbols. To resolve
cases where multiple features mapped to a single gene, only the most extreme z-score for each
combination of tissue, contrast, and gene was retained. Metabolite and lipid identifiers were
standardized using the Metabolomic Workbench Reference List of Metabolite Names (RefMet)
database[115]. For the CAMERA-PR enrichment of phosphoproteomics results, since peptides
could be phosphorylated at multiple positions, phosphorylation sites were separated into single
sites with identical row information (e.g., protein_S1;S2 becomes protein_S1 and protein_S2,
both with the same data). For any combinations of contrast and phosphosite that were not
uniquely defined, only the most extreme z-score (maximum absolute value) was selected for
inclusion in the matrix. Singly phosphorylated sites were required to use the curated kinase–
substrate relationship information available from PhosphoSitePlus (PSP)[108], described below.

Gene set selection

Gene sets were obtained from the MitoCarta3.0 database, CellMarker 2.0 database[116], and the
C2-CP (excluding KEGG_LEGACY) and C5 collections of the human Molecular Signatures
Database (MSigDB; v2023.2.Hs)[116-118] Metabolites and lipids were grouped according to
chemical subclasses from the RefMet database[115]. This includes subclasses such as "Acyl
carnitines" and "Saturated fatty acids".Flanking sequences for protein phosphorylation sites
were grouped according to their known human protein kinases provided in the
Kinase_Substrate_Dataset   file   from   PhosphoSitePlus   (PSP;   v6.7.1.1;
https://www.phosphosite.org/staticDownloads.action; last modified 2023-11-17)[108]. In addition to
these kinase sets, sets from the directional Post-Translational Modification Signatures Database
(PTMsigDB) were included for analysis[119].

For each combination of tissue and ome, molecular signatures were filtered to only those genes,
metabolites/lipids, or flanking sequences that appeared in the differential analysis results. After
filtering, all molecular signatures were required to contain at least 5 features, with no restriction
on the maximum size of sets. Additionally, gene sets were only kept if they retained at least 70%
of their original genes to increase the likelihood that the genes that remain in a given set are
accurately described by the set label.

Analysis of Molecular Signatures

Analysis of molecular signatures was carried out with the pre-ranked Correlation Adjusted MEan
RAnk (CAMERA-PR) gene set test using the z-score matrices described in the "Preparation of
differential analysis results" section to summarize the differential analysis results for each
contrast at the level of molecular signatures[25,119]. Z-scores were selected as the input statistics
primarily to satisfy the normality assumption of CAMERA-PR, and the analysis was carried out
with the cameraPR.matrix function from the TMSig R/Bioconductor package[120,121].

Over-representation analysis (ORA)

Over-representation analysis (ORA) was conducted using the run_ORA function from the
MotrpacHumanPreSuspension R package (v0.0.1.53) which uses the hypergeometric test for
statistical significance. The input consisted of differentially expressed genes (adjusted p-value <
0.05) identified from one or more omic layers in a particular tissue. The background gene set
included all genes detected in each of the ome(s) under consideration.

Statistical significance thresholds

For both CAMERA-PR and ORA results, p-values were adjusted within each combination of
tissue, ome, contrast, and broad molecular signature collection (MitoCarta3.0, CellMarker 2.0,
C2, C5, RefMet, PSP, and PTMsigDB) using the Bejamini-Hochberg method. Gene sets and
RefMet subclasses were declared significant if their adjusted p-values were less than 0.05. This
threshold was raised to 0.1 for kinase sets.

Visualization methods

Bubble heatmaps were generated from the enrichment analysis results using the enrichmap
function from the TMSig R/Bioconductor package[120].

Clustering techniques

Fuzzy C-Means Clustering

The z-score matrices described in the Enrichment Analysis Methods were used as input for
fuzzy c-means (FCM) clustering[122,123]. The matrices for the different omes, excluding global
proteomics and phosphoproteomics (as these were only measured at two timepoints) were
stacked to form a single matrix with tissue-specific timepoints as columns and features being a
row. The rows of this matrix were then divided by their sample standard deviations, which were
calculated after two columns of zeros were temporarily included to represent the pre-exercise
timepoints; these zero columns were discarded before clustering. The Mfuzz R package was
used to perform FCM clustering with the parameter m set to the value of Mfuzz::mestimate[124,125].
The minimum distance between centroids was used as the cluster validity index to determine
the optimal number of clusters for each tissue. A scree/elbow plot of these distances for a range
of cluster numbers (3-14) was generated for each tissue, and the final cluster numbers were
determined through manual inspection (n=13). For visualization purposes, hard clustering was
performed by assigning each feature to the cluster with the highest membership probability;
features with probabilities below 0.3 are not clustered or displayed. Cluster numbers are
assigned arbitrarily.

Sex difference.

To compare the differential abundance of features between male and female participants, we fit
a linear model with a single covariate (participant sex). Several alternative models were fit for
models adjusting the effect of age, clinical site, technical covariates, etc. on molecular
expression. Each of these models had, on average across all features for a given platform, a
weaker model fit as determined by Bayesian Information Criteria (BIC), partially due to the
limited number of samples available to estimate these additional parameters. The results and
model fits used to guide this decision are available in: https://github.com/MoTrPAC/precovid-
analyses/blob/main/figures/adipose/baseline_sex_differences.Rmd

Cell type analysis

Cell type enrichment. ASAT cell marker genes were curated from an integrated object of two
previously published ASAT full-length single-nuclei (sn) RNAseq[21,40]. FindALLMarkers() from
Seurat R package was used to identify marker genes (min.pct=0.1, logfc.threshold=0.25), and
adjusted p-value<0.05 and average log2FC>1 were further applied to filter out non-significant
and non-specific marker genes. A total of nine cell types were included: two mature adipocyte
subtypes (i.e., Adipocyte-1 and -2), pre-adipocyte, stem cell, vascular cell, two macrophage
subtypes (i.e., M1-like lipid-associated macrophage, LAM and M2-like resident macrophage),
mast cell, and NK/T cell. Each cell type consisted of a list of 100 marker genes (Table S2).
CAMERA-PR was applied on ASAT transcriptomics data using curated cell-type marker genes
as a reference geneset.

Deconvolution of ASAT cell types. Deconvolution—i.e., in silico estimation of cell type
proportions—was performed on ASAT transcriptomic data as previously described[22]. Briefly, an
ASAT-specific cell type signature matrix was derived from full-length single-nucleus RNA-seq
data generated from ASAT of 20 adults[21]. Using the optimized pipeline for human ASAT
deconvolution[22], we estimated the relative proportions of seven distinct ASAT cell populations:
two mature adipocyte subtypes (Adipocyte-1, Adipocyte-2), vascular cells, stem cells, pre-
adipocytes, macrophages, and mast cells. Estimated proportions were computed across all bulk
RNA-seq samples using this reference signature.

Weighted Gene Co-expression Network Analysis (WGCNA). WGCNA (version 1.73) was
employed to identify distinct, non-overlapping clusters (modules) in each ASAT omic layer at
baseline samples (transcriptomics, n=172; proteomics and phosphoproteomics, n=42;
metabolomics, n=172). Connectivity for each omic feature was calculated by summing its
correlation strengths with all other features. A scale-free signed network was constructed using
an optimal soft-threshold power (β>0.8). A dynamic tree-cutting algorithm and a merging
threshold of 0.3 was used for all omic layers.

In each omic layer, module 0 (i.e., T0, Pr0, Ph0, and M0) often referred to as the 'grey' module,
comprises transcripts that could not be assigned to any other module due to weak correlation
patterns. Therefore, module 0 was excluded for downstream analyses. Top hub genes within
each module were determined based on their kME values (module eigengene-based
connectivity). Module eigengenes (MEs), representing the first principal eigenvector of each
module, were extracted and correlated with clinical and subclinical variables using biweight
midcorrelation. ORA was conducted on module features. For phosphoproteomics, phosphosites
were collapsed and the original protein was used. For metabolomics, REFMET was used as a
reference database.

Secretome analysis

A list of putative secreted proteins was curated by combining two sources: proteins detected in
plasma via Olink proteomics (n = 1,408) (*REF: MoTrPAC Blood Companion, in review*) and
the known secretome annotated in the Human Protein Atlas (n = 2,520)[51], resulting in 3,277
unique proteins. Any differentially regulated transcripts or proteins in ASAT that overlapped with
this curated list were considered candidate *exerkines*—putative adipose-derived secretory
factors responsive to acute exercise.

Extracellular score ("COMPARTMENTS"). Subcellular localization of proteins was annotated
using the COMPARTMENTS database[53], which provides scores ranging from 0 to 5, with higher
scores indicating greater confidence in localization. The "All channels integrated" file for human
proteins was downloaded
(https://download.jensenlab.org/human_compartment_integrated_full.tsv). To calculate the
extracellular score, we selected the highest score among the following compartments
associated with extracellular localization: *Extracellular region*, *Extracellular space*, *Extracellular*
exosome, and *Extracellular vesicle*.

Gene Derived Correlation Across Tissues (GD-CAT). Transcriptomic data from the GTEx
(Genotype-Tissue Expression) project (n = 310 individuals, 19 tissues) were accessed via the

GD-CAT pipeline[52,54]. GD-CAT computes gene-to-gene correlations between user-defined origin
and target tissues to infer potential inter-tissue regulatory relationships. Of the 60 ASAT-
exerkine candidate genes identified in our study, 48 were detected in the subcutaneous adipose
tissue (SAT) transcriptome of the GTEx cohort. These 48 SAT genes were correlated with the
expression of all genes in 18 distal target tissues. Correlation was performed using biweight
midcorrelation (bicor), and significance was determined at adjusted $p < 0.05$ following
Benjamini-Hochberg correction for multiple testing. To assess the biological pathways
associated with SAT-derived exerkine signaling, bicor values were z-score transformed, and
CAMERA-PR was applied to the z-scaled bicor vectors of correlated genes in each distal tissue.

Integration of MoTrPAC rat endurance training study

Quantitative Endocrine Network Interaction Estimation (QENIE) quantifies the potential
endocrine influence of secretory genes by calculating a secretome score (Ssec) for each gene
in a given origin tissue, based on gene–gene correlation with target tissues[59]. The pipeline
begins by computing a p-value matrix from biweight midcorrelations between gene expression
in the origin and target tissues. For each secretory gene in the origin tissue, the sum of -ln(p-
value) across all genes in the target tissue is computed as the raw Ssec. This score is then
normalized by the number of genes in the target tissue, allowing for comparisons across origin–
target tissue pairs. Higher Ssec values indicate a stronger overall correlation—and potential
endocrine influence—between a secretory gene in the origin tissue and transcriptional programs
in a given target tissue. QENIE was applied to multi-tissue transcriptomic datasets from the
MoTrPAC rat endurance training study as previously described[58]. Briefly, subcutaneous white
adipose tissue (WATSC) was designated as the origin tissue. Sex-adjusted correlation p-value
matrices were obtained from correlating WATSC transcript expression matrix against 16 other
tissues within each training group (i.e., TR1W, 1-week training; TR2W, 2-week training; TR4W,
4-week training; TR8W, 8-week training, and a non-exercising control group (CON) duration-
matched with TR8W; n=10 (5F, 5M) for each group). Ovary, testis (due to differing sample
sizes), and brown adipose tissue (due to its limited presence in adult humans) were excluded.
Subsequently, only secretory protein-coding genes were retained using the same curated list
described above, yielding a total of 2,248 secretory genes identified in WATSC for Ssec
calculation. Density plots of Ssec distributions were generated across all WATSC-to–target
tissue pairs. Wilcoxon signed-rank tests and paired t-tests were used to compare Ssec ranks
between CON and TR8W for the 53 mappable ASAT-exerkine candidate genes. As previously
described, fast gene set enrichment analysis (FGSEA) was performed on bicor vectors of genes
in vastus lateralis and liver that correlated with WATSC-IGFBP7 at each time point to assess
functional associations.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical approaches for differential and enrichment analyses are described in their respective
Methods sections. Biweight midcorrelation was performed using WGCNA::bicor() for Figures 1C,
S1A, 5B–D, S5A–C, 6D, S6B–C, as well as for the secretory score analyses (i.e., QENIE and
GD-CAT) in Figures 7 and S7. Spearman correlation was applied for Figure S2I. Student's t-test
(independent) was used for Figure S1B, and the paired t-test was used for Figure S7A. A two-

way linear mixed-effects model was conducted for Figure S4A using lme4::lmer(), with exercise
group and time point as fixed effects. Adjusted *p*-values were calculated using the Benjamini–
Hochberg method. All statistical analyses were performed in R (R Foundation for Statistical
Computing, Vienna, Austria)

Characteristic	Overall N = 173[1]	CON N = 37[1]	EE N = 63[1]	RE N = 73[1]
Age (yrs)	41 (15)	42 (15)	41 (14)	40 (15)
Height (cm)	168 (9)	167 (8)	167 (9)	168 (10)
Weight (kg)	76 (13)	73 (13)	76 (13)	77 (14)
BMI (kg/m²)	26.9 (4.0)	26.2 (3.9)	26.9 (3.9)	27.2 (4.0)
Waist Circumference (cm)	92 (11)	89 (13)	92 (10)	93 (12)
Systolic BP (mmHg)	116 (12)	114 (11)	117 (11)	116 (12)
Diastolic BP (mmHg)	72 (8)	72 (8)	73 (9)	72 (8)
Hemoglobin A1c (%)	5.30 (0.30)	5.29 (0.21)	5.28 (0.32)	5.31 (0.32)
eGFR (mL/min/1.73 m²)	101 (20)	99 (19)	103 (18)	101 (22)
Total Cholesterol (mg/dL)	192 (40)	197 (44)	192 (40)	189 (39)
HDL Cholesterol (mg/dL)	58 (15)	62 (17)	57 (14)	56 (15)
LDL Cholesterol (mg/dL)	117 (34)	119 (38)	116 (34)	116 (32)
HOMA-IR	2.21 (1.52)	1.99 (2.07)	2.07 (1.18)	2.44 (1.46)
NEFA (mmol/L)	0.54 (0.19)	0.57 (0.19)	0.56 (0.21)	0.51 (0.17)
Cardiopulmonary Exercise Test (CPET)				
VO2max (L)	1.86 (0.65)	1.73 (0.56)	1.89 (0.68)	1.89 (0.65)
VO2max (ml/min/kg)	24 (7)	24 (6)	25 (8)	25 (7)
Workload (Watts)	151 (50)	142 (40)	155 (53)	153 (52)
O2 Pulse (ml/beat)	10.6 (3.3)	10.0 (2.8)	10.8 (3.5)	10.8 (3.4)
RER	1.16 (0.08)	1.16 (0.08)	1.17 (0.07)	1.15 (0.08)
Strength Tests				

Isometric Knee Extension Torque (Nm)	147 (57)	137 (53)	143 (54)	156 (60)
Hand Grip Strength (kg)	30 (11)	29 (8)	30 (11)	31 (12)

[1] Mean (SD)

1604

1605 Table 1. Pre-exercise participant characteristics

1606 BMI, Body Mass Index; HR, Heart Rate; SBP, Systolic Blood Pressure; DBP, Diastolic Blood
1607 Pressure; eGFR, estimated glomerular filtration rate; HDL, High-Density Cholesterol; LDL, Low-
1608 Density Cholesterol; HOMA-IR, Homeostatic Model Assessment for Insulin Resistance; NEFA,
1609 Non-esterified fatty acids.

1610

FIGURE LEGENDS

Figure 1. Study design and baseline characteristics.

A) Overview of the study design. B) Workflow of pre-exercise cell type deconvolution of ASAT.
C) Heatmap showing correlation between estimated cell type proportions and clinical/subclinical
variables. "p<0.05, **p<0.01, ***p<0.001. D) Volcano plots showing sex-based differences (male
vs. female) in transcriptomics, proteomics, phosphoproteomics, and metabolomics. E)
CAMERA-PR results highlighting pathways enriched in males vs. females. Positive z-scores
indicate male-biased enrichment. For Kinase substrate enrichment analysis (KSEA), no kinases
were significantly upregulated in the males.

Figure 2. Multi-omic responses to acute exercise across modalities and time points.

A) Volcano plots illustrating differentially regulated features across omic layers—transcriptomics,
proteomics, phosphoproteomics, and metabolomics—stratified by exercise modality and time
point. Differential analysis (DA) model accounts for changes observed in the control (CON)
group. B) Principal component analysis of summarized log fold-change (logFC) values across
modalities and time points, highlighting global shifts in transcriptomic and metabolomic
responses to exercise. C) Scatter plots comparing significantly enriched biological terms from
CAMERA-PR (adjusted $p < 0.05$) between CON-adjusted (y-axis) and CON-unadjusted (x-
axis) DA results for transcriptomics, proteomics, and metabolomics. Hot pink indicates terms
enriched only in CON-adjusted analyses; gray, terms enriched only in unadjusted analyses;
purple, terms shared by both. Positive z.std values indicate upregulation; negative values
indicate downregulation. D) Kinome phylogenetic tree annotated with enriched kinases
(adjusted $p < 0.1$) from PTM-SEA at post 4 hr following EE or RE. Red branches denote
upregulated kinase activity; blue branches denote downregulation. See Methods for kinase
group classification. E) Scatter plot of comparing logFC of differentially expressed phosphosites
(adjusted $p < 0.05$) at post 4 hr (EE or RE), annotated with cytoskeleton remodeling-related
Gene Ontology terms. Shapes indicate whether a phosphosite was differentially regulated by
EE only, RE only, or both.

Figure 3. Acute exercise induces modality-specific responses in ASAT

A) Volcano plots illustrating differentially regulated features (adjusted p<0.05) between EE vs. RE across omic layers at each post-exercise time point. Positive logFC indicates greater enrichment in EE; negative logFC indicates greater enrichment in RE. B) Scatter plot comparing the z.std of differentially regulated features between EE and RE. Positive z.std values indicate upregulation; negative values indicate downregulation. Shape indicates time point. C) Individual line plots of features representing each quadrant of Figure 3B are shown. CAMERA-PR analysis comparing EE vs. RE in D) transcriptomic, E) proteomics, and F) metabolomics. Positive z.std values represent pathways more enriched in EE; negative values indicate enrichment in RE. G) Heatmap illustrating exercise responses in canonical ceramide species. *adjusted p<0.05.

Figure 4. Acute exercise induces cell-type-specific responses in ASAT

A) Cell-type enrichment analysis using a reference cell signature matrix generated from ASAT single-nucleus RNA-seq datasets. CAMERA-PR was applied to ASAT transcriptomic data using the signature matrix as the gene set reference. B) Volcano plots of differentially regulated transcripts, proteins, and phosphosites mapped to known cell-type markers in EE and RE. 'Non-marker' indicates features not specific to any known ASAT cell type. C) EE-specific and D) RE-specific fuzzy c-means clustering analysis showing 13 trajectory clusters of combined object of transcriptomes and metabolomes. ORA was conducted on each cluster using the curated cell type markers and REFMET.

Figure 5. Acute exercise may induce metabolically favorable molecular responses in ASAT

A) Schematic overview of weighted gene co-expression network analysis (WGCNA) and downstream analyses. B) WGCNA was independently performed for each omic layer—transcriptomics, proteomics, phosphoproteomics, and metabolomics—using baseline data. Each panel includes: over-represented pathways per module (*FDR$\square$<$\square$0.05), correlations between module eigengenes and z-scored clinical/subclinical traits (*p$\square$<$\square$0.05; **p$\square$<$\square$0.01; ***p$\square$<$\square$0.001), module size (feature count), and the number of differentially regulated features (adjusted p<0.05) per module following acute exercise. C) Left: Scatterplot of module membership (kME) for phosphosites in Ph10; phosphosites differentially regulated by exercise are highlighted in red. Middle: Heatmaps showing baseline Spearman correlations between these phosphosites and clinical markers of insulin resistance (HOMA-IR, ADIPO-IR, HbA1c) (*p$\square$<$\square$0.05; **p$\square$<$\square$0.01). Right: Corresponding phosphosite fold changes after acute exercise (*adjusted p$\square$<$\square$0.05). D) Left: Scatterplot of kME values for proteins in Pr5; red points indicate proteins differentially regulated by exercise. Middle: Heatmaps showing baseline correlations with circulating BCAAs (*p$\square$<$\square$0.05; **p$\square$<$\square$0.01). Right: Protein fold changes following exercise (*adjusted p$\square$<$\square$0.05). E) Schematic summary of the BCKA perfusion experiment by Walejko et al., in which isolated rat hearts perfused with BCKAs via the Langendorff apparatus showed increased phosphorylation of 4E-BP1 (a translational repressor), enhanced protein synthesis, and activation of cytoskeleton-related phosphosignaling. F) Changes in plasma BCKA levels during and following EE and RE. G) CAMERA-PR enrichment results for ASAT

transcriptomics filtered for protein translation-related pathways. H) ORA of differentially regulated phosphoproteins in ASAT at post 4□hr, overlapping with phosphoproteins identified in the BCKA heart perfusion experiment. An adjusted p-value threshold of 0.1 was used for ASAT phosphosite selection. BCAA, Branched-chain amino acids; BCKA, branched-chain ketoacids.

Figure 6. Identification of ASAT-secreted exerkine candidates and their potential endocrine role

A) Schematic overview of the secretome screening pipeline. Transcripts and proteins differentially regulated at any post-exercise timepoint (post 45 min, 4r, and 24hr) were filtered using MoTrPAC blood Olink proteomics data and secretome annotations from the Human Protein Atlas. Metabolites were filtered against MoTrPAC blood metabolomics. B) Bar plots showing the number of differentially regulated transcripts and proteins per time point. Purple bars indicate features annotated as secreted. C) Bar plots showing the number of differentially regulated metabolites per time point. Purple bars indicate metabolites also detected in plasma. D) Integration with GTEx: RNA-seq from n = 310 GTEx subjects across 18 solid tissues was used to correlate 48 GTEx-mappable ASAT-exerkine candidate genes with genes in distal tissues. The heatmap displays the number of significantly correlated genes (adjusted p<0.05) per exerkine candidate per target tissue. Green bars indicate the extracellular region score (range 0–5) from COMPARTMENTS, reflecting the strongest extracellular annotation across multiple categories (region, space, exosome, vesicle). E) Exercise responses of CEACAM8 and BST2 proteins in both ASAT and circulation. F) GD-CAT analysis showing pathways associated with SAT-CEACAM8 and SAT-BST2 in selected target tissues. CAMERA-PR was used for enrichment testing.

Figure 7. Contribution of ASAT-exerkine candidates to training-induced remodeling of adipose tissue endocrine network

A) Schematic workflow for applying QENIE to the MoTrPAC rat endurance training study. Human exerkine candidates were mapped to rat orthologs, and the same secretome filters as in Figure 6 were applied, yielding 2,248 WATSC secretory genes. B) Density plots showing distributions of secretome scores (Ssec) across WATSC-to-target tissue connections over 1, 2, 4, and 8 weeks of endurance training (TR1W, TR2W, TR4W, TR8W), compared with non-trained control rats (CON). C) Wilcoxon signed-rank test comparing Ssec rank distributions of 53 mappable ASAT-exerkine candidates between CON and TR8W for each WATSC-to-target tissue connection. Adjusted **p□<□0.01, ***p□<□0.001, **p□<□0.0001. D) Time-course trajectories of Ssec values for the 53 exerkine candidates across training weeks. Top 3 genes per tissue connection at TR8W (with extracellular score ≥□4) are labeled. E) String plots showing the top 15 exerkine candidates with the greatest Ssec rank shift between CON and TR8W for WATSC-to-vastus lateralis, liver, and kidney connections. Pink triangle, CON; dark green circle, TR8W. Vertical green bars denote extracellular scores. F) Exercise-induced response of IGFBP7 protein at post 4□hr. G) FGSEA results showing pathways associated with WATSC-Igfbp7 in vastus lateralis. H) FGSEA results showing pathways associated with WATSC-Igfbp7 in the liver. ADREN, adrenal gland; HIPPO, hippocampus; HYPOTH, hypothalamus; SKM-GN/VL, skeletal muscle gastrocnemius/vastus lateralis; SMLINT, small intestine; VENACV, vena cava; NES, normalized enrichment score.

REFERENCES

1. Ashcroft, S.P., Stocks, B., Egan, B., and Zierath, J.R. (2024). Exercise induces tissue-specific adaptations to enhance cardiometabolic health. Cell Metab *36*, 278-300. 10.1016/j.cmet.2023.12.008.

2. Knowler, W.C., Barrett-Connor, E., Fowler, S.E., Hamman, R.F., Lachin, J.M., Walker, E.A., Nathan, D.M., Watson, P.G., Mendoza, J.T., and Smith, K.A. (2002). Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin.

3. Goodyear, L.J., and Kahn, B.B. (1998). Exercise, glucose transport, and insulin sensitivity. Annual review of medicine *49*, 235-261.

4. Houmard, J.A., Tanner, C.J., Slentz, C.A., Duscha, B.D., McCartney, J.S., and Kraus, W.E. (2004). Effect of the volume and intensity of exercise training on insulin sensitivity. Journal of applied physiology *96*, 101-106.

5. Cardoso, C.G., Jr., Gomides, R.S., Queiroz, A.C., Pinto, L.G., da Silveira Lobo, F., Tinucci, T., Mion, D., Jr., and de Moraes Forjaz, C.L. (2010). Acute and chronic effects of aerobic and resistance exercise on ambulatory blood pressure. Clinics (Sao Paulo) *65*, 317-325. 10.1590/S1807-59322010000300013.

6. Ryan, B.J., Schleh, M.W., Ahn, C., Ludzki, A.C., Gillen, J.B., Varshney, P., Van Pelt, D.W., Pitchford, L.M., Chenevert, T.L., and Gioscia-Ryan, R.A. (2020). Moderate-intensity exercise and high-intensity interval training affect insulin sensitivity similarly in obese adults. The Journal of Clinical Endocrinology & Metabolism *105*, e2941-e2959.

7. Morrison, D.J., Kowalski, G.M., Grespan, E., Mari, A., Bruce, C.R., and Wadley, G.D. (2018). Measurement of postprandial glucose fluxes in response to acute and chronic endurance exercise in healthy humans. Am J Physiol Endocrinol Metab *314*, E503-E511. 10.1152/ajpendo.00316.2017.

8. Ross, R., Dagnone, D., Jones, P.J., Smith, H., Paddags, A., Hudson, R., and Janssen, I. (2000). Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men: a randomized, controlled trial. Annals of internal medicine *133*, 92-103.

9. Egan, B., and Zierath, J.R. (2013). Exercise metabolism and the molecular regulation of skeletal muscle adaptation. Cell Metab *17*, 162-184. 10.1016/j.cmet.2012.12.012.

10. Fabre, O., Ingerslev, L.R., Garde, C., Donkin, I., Simar, D., and Barres, R. (2018). Exercise training alters the genomic response to acute exercise in human adipose tissue. Epigenomics *10*, 1033-1050.

11. Ludzki, A.C., Pataky, M.W., Cartee, G.D., and Horowitz, J.F. (2018). Acute endurance exercise increases Vegfa mRNA expression in adipose tissue of rats during the early stages of weight gain. Applied Physiology, Nutrition, and Metabolism *43*, 751-754.

12. Ahn, C., Zhang, T., Rode, T., Yang, G., Chugh, O.K., Ellis, S., Ghayur, S., Mehta, S., Salzman, R., Jiang, H., et al. (2025). Molecular responses in abdominal subcutaneous adipose tissue after a session of endurance exercise: effects of exercise intensity. The Journal of Physiology. 10.1113/jp289339.

13. Ahn, C., Ryan, B.J., Schleh, M.W., Varshney, P., Ludzki, A.C., Gillen, J.B., Van Pelt, D.W., Pitchford, L.M., Howton, S.M., and Rode, T. (2022). Exercise training remodels

subcutaneous adipose tissue in adults with obesity even without weight loss. The
Journal of Physiology *600*, 2127-2146.

14.   Ahn, C., Zhang, T., Yang, G., Rode, T., Varshney, P., Ghayur, S.J., Chugh, O.K., Jiang,
H., and Horowitz, J.F. (2024). Years of endurance exercise training remodel abdominal
subcutaneous adipose tissue in adults with overweight or obesity. Nature Metabolism.
10.1038/s42255-024-01103-x.

15.   Pino, M.F., Dijkstra, P., Whytock, K.L., Ahn, C., Yu, G., Sanford, J.A., Hansen, J.,
Hutchinson, C., Gritsenko, M., Piehowski, P., et al. (2025). Exercise alters molecular
profiles of inflammation and substrate metabolism in human white adipose tissue. Am J
Physiol Endocrinol Metab *328*, E478-E492. 10.1152/ajpendo.00339.2024.

16.   Stinkens, R., Brouwers, B., Jocken, J.W., Blaak, E.E., Teunissen-Beekman, K.F.,
Hesselink, M.K., van Baak, M.A., Schrauwen, P., and Goossens, G.H. (2018). Exercise
training-induced effects on the abdominal subcutaneous adipose tissue phenotype in
humans with obesity. Journal of Applied Physiology *125*, 1585-1593.

17.   Riis, S., Christensen, B., Nellemann, B., Møller, A.B., Husted, A.S., Pedersen, S.B.,
Schwartz, T.W., Jørgensen, J.O.L., and Jessen, N. (2019). Molecular adaptations in
human subcutaneous adipose tissue after ten weeks of endurance exercise training in
healthy males. Journal of Applied Physiology *126*, 569-577.

18.   Stanford, K.I., Middelbeek, R.J., Townsend, K.L., Lee, M.-Y., Takahashi, H., So, K.,
Hitchcox, K.M., Markan, K.R., Hellbach, K., and Hirshman, M.F. (2015). A novel role for
subcutaneous adipose tissue in exercise-induced improvements in glucose homeostasis.
Diabetes *64*, 2002-2014.

19.   Chow, L.S., Gerszten, R.E., Taylor, J.M., Pedersen, B.K., van Praag, H., Trappe, S.,
Febbraio, M.A., Galis, Z.S., Gao, Y., Haus, J.M., et al. (2022). Exerkines in health,
resilience and disease. Nat Rev Endocrinol. 10.1038/s41574-022-00641-2.

20.   Newgard, C.B., An, J., Bain, J.R., Muehlbauer, M.J., Stevens, R.D., Lien, L.F., Haqq,
A.M., Shah, S.H., Arlotto, M., Slentz, C.A., et al. (2009). A branched-chain amino acid-
related metabolic signature that differentiates obese and lean humans and contributes to
insulin resistance. Cell Metab *9*, 311-326. 10.1016/j.cmet.2009.02.002.

21.   Whytock, K.L., Divoux, A., Sun, Y., Pino, M.F., Yu, G., Jin, C.A., Robino, J.J., Plekhanov,
A., Varlamov, O., Smith, S.R., et al. (2024). Aging human abdominal subcutaneous
white adipose tissue at single cell resolution. Aging Cell, e14287. 10.1111/acel.14287.

22.   Ahn, C., Divoux, A., Zhou, M., Seldin, M.M., Sparks, L.M., and Whytock, K.L. (2025).
Optimized RNA sequencing deconvolution illustrates the impact of obesity and weight
loss on cell composition of human adipose tissue. Obesity (Silver Spring).
10.1002/oby.24264.

23.   Weisberg, S.P., McCann, D., Desai, M., Rosenbaum, M., Leibel, R.L., and Ferrante,
A.W., Jr. (2003). Obesity is associated with macrophage accumulation in adipose tissue.
J Clin Invest *112*, 1796-1808. 10.1172/JCI19246.

24.   Kunz, H.E., Hart, C.R., Gries, K.J., Parvizi, M., Laurenti, M., Dalla Man, C., Moore, N.,
Zhang, X., Ryan, Z., and Polley, E.C. (2021). Adipose tissue macrophage populations
and inflammation are associated with systemic inflammation and insulin resistance in
obesity. American Journal of Physiology-Endocrinology and Metabolism *321*, E105-E121.

25. Wu, D., and Smyth, G.K. (2012). Camera: a competitive gene set test accounting for
inter-gene correlation. Nucleic Acids Res *40*, e133. 10.1093/nar/gks461.
26. Bost, F., Aouadi, M., Caron, L., and Binétruy, B. (2005). The role of MAPKs in adipocyte
differentiation and obesity. Biochimie *87*, 51-56.
27. Cong, L.-N., Chen, H., Li, Y., Zhou, L., McGibbon, M.A., Taylor, S.I., and Quon, M.J.
(1997). Physiological role of Akt in insulin-stimulated translocation of GLUT4 in
transfected rat adipose cells. Molecular endocrinology *11*, 1881-1890.
28. Szczepanowska, J. (2009). Involvement of Rac/Cdc42/PAK pathway in cytoskeletal
rearrangements. Acta Biochimica Polonica *56*.
29. Pasapera, A.M., Heissler, S.M., Eto, M., Nishimura, Y., Fischer, R.S., Thiam, H.R., and
Waterman, C.M. (2022). MARK2 regulates directed cell migration through modulation of
myosin II contractility and focal adhesion organization. Curr Biol *32*, 2704-2718 e2706.
10.1016/j.cub.2022.04.088.
30. Maddika, S., and Chen, J. (2009). Protein kinase DYRK2 is a scaffold that facilitates
assembly of an E3 ligase. Nat Cell Biol *11*, 409-419. 10.1038/ncb1848.
31. Even-Faitelson, L., and Ravid, S. (2006). PAK1 and aPKCzeta regulate myosin II-B
phosphorylation: a novel signaling pathway regulating filament assembly. Mol Biol Cell
17, 2869-2881. 10.1091/mbc.e05-11-1001.
32. Lo, S.H., Janmey, P.A., Hartwig, J.H., and Chen, L.B. (1994). Interactions of tensin with
actin and identification of its three distinct actin-binding domains. The Journal of cell
biology *125*, 1067-1075.
33. Bernau, K., Torr, E.E., Evans, M.D., Aoki, J.K., Ngam, C.R., and Sandbo, N. (2017).
Tensin 1 Is Essential for Myofibroblast Differentiation and Extracellular Matrix Formation.
Am J Respir Cell Mol Biol *56*, 465-476. 10.1165/rcmb.2016-0104OC.
34. Chao, L.C., Wroblewski, K., Zhang, Z., Pei, L., Vergnes, L., Ilkayeva, O.R., Ding, S.Y.,
Reue, K., Watt, M.J., Newgard, C.B., et al. (2009). Insulin resistance and altered
systemic glucose metabolism in mice lacking Nur77. Diabetes *58*, 2788-2796.
10.2337/db09-0763.
35. Qin, D.D., Yang, Y.F., Pu, Z.Q., Liu, D., Yu, C., Gao, P., Chen, J.C., Zong, C., Zhang,
Y.C., Li, X., et al. (2018). NR4A1 retards adipocyte differentiation or maturation via
enhancing GATA2 and p53 expression. J Cell Mol Med *22*, 4709-4720.
10.1111/jcmm.13715.
36. Cai, K., El-Merahbi, R., Loeffler, M., Mayer, A.E., and Sumara, G. (2017). Ndrg1
promotes adipocyte differentiation and sustains their function. Sci Rep *7*, 7191.
10.1038/s41598-017-07497-x.
37. Wang, Y.N., Tang, Y., He, Z., Ma, H., Wang, L., Liu, Y., Yang, Q., Pan, D., Zhu, C., Qian,
S., and Tang, Q.Q. (2021). Slit3 secreted from M2-like macrophages increases
sympathetic activity and thermogenesis in adipose tissue. Nat Metab *3*, 1536-1551.
10.1038/s42255-021-00482-9.
38. Chaurasia, B., Talbot, C.L., and Summers, S.A. (2020). Adipocyte Ceramides-The
Nexus of Inflammation and Metabolic Disease. Front Immunol *11*, 576347.
10.3389/fimmu.2020.576347.
39. Bergman, B.C., Brozinick, J.T., Strauss, A., Bacon, S., Kerege, A., Bui, H.H., Sanders,
P., Siddall, P., Wei, T., Thomas, M.K., et al. (2016). Muscle sphingolipids during rest and

exercise: a C18:0 signature for insulin resistance in humans. Diabetologia *59*, 785-798.
10.1007/s00125-015-3850-y.

40.   Whytock, K.L., Sun, Y., Divoux, A., Yu, G., Smith, S.R., Walsh, M.J., and Sparks, L.M.
(2022). Single cell full-length transcriptome of human subcutaneous adipose tissue
reveals unique and heterogeneous cell populations. iScience *25*, 104772.
10.1016/j.isci.2022.104772.

41.   Borriello, F., Iannone, R., and Marone, G. (2017). Histamine release from mast cells and
basophils. Histamine and histamine receptors in health and disease, 121-139.

42.   Van der Stede, T., Van de Loock, A., Turiel, G., Hansen, C., Tamariz-Ellemann, A.,
Ullrich, M., Lievens, E., Spaas, J., Yigit, N., Anckaert, J., et al. (2025). Cellular
deconstruction of the human skeletal muscle microenvironment identifies an exercise-
induced histaminergic crosstalk. Cell Metab *37*, 842-856 e847.
10.1016/j.cmet.2024.12.011.

43.   Luttrell, M.J., and Halliwill, J.R. (2017). The Intriguing Role of Histamine in Exercise
Responses. Exerc Sport Sci Rev *45*, 16-23. 10.1249/JES.0000000000000093.

44.   Snider, S.A., Margison, K.D., Ghorbani, P., LeBlond, N.D., O'Dwyer, C., Nunes, J.R.C.,
Nguyen, T., Xu, H., Bennett, S.A.L., and Fullerton, M.D. (2018). Choline transport links
macrophage phospholipid metabolism and inflammation. J Biol Chem *293*, 11600-11611.
10.1074/jbc.RA118.003180.

45.   Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation
network analysis. BMC Bioinformatics *9*, 559. 10.1186/1471-2105-9-559.

46.   Langfelder, P., and Horvath, S. (2012). Fast R functions for robust correlations and
hierarchical clustering. Journal of statistical software *46*.

47.   Kawai, T., Autieri, M.V., and Scalia, R. (2021). Adipose tissue inflammation and
metabolic dysfunction in obesity. American Journal of Physiology-Cell Physiology *320*,
C375-C391.

48.   Hansson, B., Moren, B., Fryklund, C., Vliex, L., Wasserstrom, S., Albinsson, S., Berger,
K., and Stenkula, K.G. (2019). Adipose cell size changes are associated with a drastic
actin remodeling. Sci Rep *9*, 12941. 10.1038/s41598-019-49418-0.

49.   Zhou, M., Shao, J., Wu, C.Y., Shu, L., Dong, W., Liu, Y., Chen, M., Wynn, R.M., Wang,
J., Wang, J., et al. (2019). Targeting BCAA Catabolism to Treat Obesity-Associated
Insulin Resistance. Diabetes *68*, 1730-1746. 10.2337/db18-0927.

50.   Walejko, J.M., Christopher, B.A., Crown, S.B., Zhang, G.F., Pickar-Oliver, A., Yoneshiro,
T., Foster, M.W., Page, S., van Vliet, S., Ilkayeva, O., et al. (2021). Branched-chain
alpha-ketoacids are preferentially reaminated and activate protein synthesis in the heart.
Nat Commun *12*, 1680. 10.1038/s41467-021-21962-2.

51.   Uhlén, M., Karlsson, M.J., Hober, A., Svensson, A.-S., Scheffel, J., Kotol, D., Zhong, W.,
Tebani, A., Strandberg, L., and Edfors, F. (2019). The human secretome. Science
signaling *12*, eaaz0274.

52.   Consortium, G.T., Laboratory, D.A., Coordinating Center -Analysis Working, G.,
Statistical Methods groups-Analysis Working, G., Enhancing, G.g., Fund, N.I.H.C.,
Nih/Nci, Nih/Nhgri, Nih/Nimh, Nih/Nida, et al. (2017). Genetic effects on gene expression
across human tissues. Nature *550*, 204-213. 10.1038/nature24277.

53. Binder, J.X., Pletscher-Frankild, S., Tsafou, K., Stolte, C., O'Donoghue, S.I., Schneider,
R., and Jensen, L.J. (2014). COMPARTMENTS: unification and visualization of protein
subcellular localization evidence. Database (Oxford) *2014*, bau012.
10.1093/database/bau012.
54. Zhou, M., Tamburini, I., Van, C., Molendijk, J., Nguyen, C.M., Chang, I.Y., Johnson, C.,
Velez, L.M., Cheon, Y., Yeo, R., et al. (2024). Leveraging inter-individual transcriptional
correlation structure to infer discrete signaling mechanisms across metabolic tissues.
Elife *12*. 10.7554/eLife.88863.
55. Cao, Y., Wang, Y., Zhou, Z., Pan, C., Jiang, L., Zhou, Z., Meng, Y., Charugundla, S., Li,
T., and Allayee, H. (2022). Liver-heart cross-talk mediated by coagulation factor XI
protects against heart failure. Science *377*, 1399-1406.
56. Singer, B.B., Opp, L., Heinrich, A., Schreiber, F., Binding-Liermann, R., Berrocal-
Almanza, L.C., Heyl, K.A., Muller, M.M., Weimann, A., Zweigner, J., and Slevogt, H.
(2014). Soluble CEACAM8 interacts with CEACAM1 inhibiting TLR2-triggered immune
responses. PLoS One *9*, e94106. 10.1371/journal.pone.0094106.
57. Mahauad-Fernandez, W.D., and Okeoma, C.M. (2016). The role of BST-2/Tetherin in
host protection and disease manifestation. Immun Inflamm Dis *4*, 4-23. 10.1002/iid3.92.
58. MoTr, P.A.C.S.G., Lead, A., and MoTr, P.A.C.S.G. (2024). Temporal dynamics of the
multi-omic response to endurance exercise training. Nature *629*, 174-183.
10.1038/s41586-023-06877-w.
59. Seldin, M.M., Koplev, S., Rajbhandari, P., Vergnes, L., Rosenberg, G.M., Meng, Y., Pan,
C., Phuong, T.M.N., Gharakhanian, R., Che, N., et al. (2018). A Strategy for Discovery of
Endocrine Interactions with Application to Whole-Body Metabolism. Cell Metab *27*, 1138-
1155 e1136. 10.1016/j.cmet.2018.03.015.
60. Ahn, C., Hevener, A.L., Goodyear, L.J., Bodine, S.C., Esser, K.A., Seldin, M.M., and
Sparks, L.M. (2025). Exercise training remodels inter-organ endocrine networks. Mol
Metab *99*, 102219. 10.1016/j.molmet.2025.102219.
61. Many, G.M., Sanford, J.A., Sagendorf, T.J., Hou, Z., Nigro, P., Whytock, K.L., Amar, D.,
Caputo, T., Gay, N.R., Gaul, D.A., et al. (2024). Sexual dimorphism and the multi-omic
response to exercise training in rat subcutaneous white adipose tissue. Nat Metab *6*,
963-979. 10.1038/s42255-023-00959-9.
62. Stromberg, A., Olsson, K., Dijksterhuis, J.P., Rullman, E., Schulte, G., and Gustafsson,
T. (2016). CX3CL1--a macrophage chemoattractant induced by a single bout of exercise
in human skeletal muscle. Am J Physiol Regul Integr Comp Physiol *310*, R297-304.
10.1152/ajpregu.00236.2015.
63. Gonzalez-Franquesa, A., Peijs, L., Cervone, D.T., Kocana, C., Zierath, J.R., and
Deshmukh, A.S. (2021). Insulin and 5-Aminoimidazole-4-Carboxamide Ribonucleotide
(AICAR) Differentially Regulate the Skeletal Muscle Cell Secretome. Proteomes *9*.
10.3390/proteomes9030037.
64. Akiel, M., Guo, C., Li, X., Rajasekaran, D., Mendoza, R.G., Robertson, C.L., Jariwala, N.,
Yuan, F., Subler, M.A., Windle, J., et al. (2017). IGFBP7 Deletion Promotes
Hepatocellular Carcinoma. Cancer Res *77*, 4014-4025. 10.1158/0008-5472.CAN-16-
2885.

65. Tomimaru, Y., Eguchi, H., Wada, H., Kobayashi, S., Marubashi, S., Tanemura, M.,
Umeshita, K., Kim, T., Wakasa, K., and Doki, Y. (2012). IGFBP7 downregulation is
associated with tumor progression and clinical outcome in hepatocellular carcinoma.
International journal of cancer *130*, 319-327.
66. Hepler, C., and Bass, J. (2023). Circadian mechanisms in adipose tissue bioenergetics
and plasticity. Genes Dev *37*, 454-473. 10.1101/gad.350759.123.
67. Dollet, L., and Zierath, J.R. (2019). Interplay between diet, exercise and the molecular
circadian clock in orchestrating metabolic adaptations of adipose tissue. The Journal of
Physiology *597*, 1439-1450.
68. Ahn, C., Zhang, T., Rode, T., Yang, G., Chugh, O.K., Ellis, S., Ghayur, S.J., Mehta, S.,
Salzman, R., Jiang, H., et al. (2025). Acute session of three endurance exercise
intensities alters subcutaneous adipose tissue transcriptome in regular exercisers.
bioRxiv, 2025.2005.2002.651890. 10.1101/2025.05.02.651890.
69. Tsakiridis, T., Tong, P., Matthews, B., Tsiani, E., Bilan, P.J., Klip, A., and Downey, G.P.
(1999). Role of the actin cytoskeleton in insulin action. Microscopy Research and
Technique *47*, 79-92. 10.1002/(sici)1097-0029(19991015)47:2<79::Aid-jemt1>3.0.Co;2-s.
70. Fischer, C.P., Plomgaard, P., Hansen, A.K., Pilegaard, H., Saltin, B., and Pedersen, B.K.
(2004). Endurance training reduces the contraction-induced interleukin-6 mRNA
expression in human skeletal muscle. Am J Physiol Endocrinol Metab *287*, E1189-1194.
10.1152/ajpendo.00206.2004.
71. Goj, T., Hoene, M., Fritsche, L., Schneeweiss, P., Machann, J., Petrera, A., Hauck, S.M.,
Fritsche, A., Birkenfeld, A.L., Peter, A., et al. (2023). The Acute Cytokine Response to
30-Minute Exercise Bouts Before and After 8-Week Endurance Training in Individuals
With Obesity. J Clin Endocrinol Metab *108*, 865-875. 10.1210/clinem/dgac623.
72. Strocchi, S., Liu, L., Wang, R., Haseli, S.P., Capone, F., Bode, D., Nambiar, N., Eroglu,
T., Santiago Padilla, L., Farrelly, C., et al. (2024). Systems Biology Approach Uncovers
Candidates for Liver-Heart Interorgan Crosstalk in HFpEF. Circ Res *135*, 873-876.
10.1161/CIRCRESAHA.124.324829.
73. Group, M.S., Jakicic, J.M., Kohrt, W.M., Houmard, J.A., Miller, M.E., Radom-Aizik, S.,
Rasmussen, B.B., Ravussin, E., Serra, M., and Stowe, C.L. (2024). Molecular
Transducers of Physical Activity Consortium (MoTrPAC): human studies design and
protocol. Journal of Applied Physiology *137*, 473-493.
74. Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput
sequencing reads. EMBnet J. *17*, 10. 10.14806/ej.17.1.200.
75. Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., and Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics *29*, 15-21. 10.1093/bioinformatics/bts635.
76. Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq
data with or without a reference genome. BMC Bioinformatics *12*, 323. 10.1186/1471-
2105-12-323.
77. Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2.
Nat. Methods *9*, 357-359. 10.1038/nmeth.1923.
78. Hoffman, G.E. (2017). variancePartition (Bioconductor).

79. Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015).
1981   limma powers differential expression analyses for RNA-sequencing and microarray
1982   studies. Nucleic Acids Res. *43*, e47. 10.1093/nar/gkv007.
1983 80. Mertins, P., Tang, L.C., Krug, K., Clark, D.J., Gritsenko, M.A., Chen, L., Clauser, K.R.,
1984   Clauss, T.R., Shah, P., Gillette, M.A., et al. (2018). Reproducible workflow for
1985   multiplexed deep-scale proteome and phosphoproteome analysis of tumor tissues by
1986   liquid chromatography-mass spectrometry. Nat. Protoc. *13*, 1632-1661. 10.1038/s41596-
1987   018-0006-9.
1988 81. Zecha, J., Satpathy, S., Kanashova, T., Avanessian, S.C., Kane, M.H., Clauser, K.R.,
1989   Mertins, P., Carr, S.A., and Kuster, B. (2019). TMT labeling for the masses: A robust and
1990   cost-efficient, in-solution labeling approach. Mol. Cell. Proteomics *18*, 1468-1478.
1991   10.1074/mcp.TIR119.001385.
1992 82. Jimenez-Morales, D., Samdarshi, M., and Hershman, S. (2023). MoTrPAC/motrpac-
1993   proteomics-pipeline: MoTrPAC proteomics pipeline v1.0.0 (Zenodo).
1994 83. Olson, K.C., Chen, G., and Lynch, C.J. (2013). Quantification of branched-chain keto
1995   acids in tissue by ultra fast liquid chromatography-mass spectrometry. Anal. Biochem.
1996   *439*, 116-122. 10.1016/j.ab.2013.05.002.
1997 84. Kwee, L.C., Ilkayeva, O., Muehlbauer, M.J., Bihlmeyer, N., Wolfe, B., Purnell, J.Q.,
1998   Xavier Pi-Sunyer, F., Chen, H., Bahnson, J., Newgard, C.B., et al. (2021). Metabolites
1999   and diabetes remission after weight loss. Nutr. Diabetes *11*, 10. 10.1038/s41387-021-
2000   00151-6.
2001 85. Deutsch, J., Grange, E., Rapoport, S.I., and Purdon, A.D. (1994). Isolation and
2002   quantitation of long-chain acyl-coenzyme A esters in brain tissue by solid-phase
2003   extraction. Anal. Biochem. *220*, 321-323. 10.1006/abio.1994.1344.
2004 86. Magnes, C., Sinner, F.M., Regittnig, W., and Pieber, T.R. (2005). LC/MS/MS method for
2005   quantitative determination of long-chain fatty acyl-CoAs. Anal. Chem. *77*, 2889-2894.
2006   10.1021/ac048314i.
2007 87. Minkler, P.E., Kerner, J., Ingalls, S.T., and Hoppel, C.L. (2008). Novel isolation
2008   procedure for short-, medium-, and long-chain acyl-coenzyme A esters from tissue.
2009   Analytical biochemistry *376*, 275-276.
2010 88. Cordell, R.L., Hill, S.J., Ortori, C.A., and Barrett, D.A. (2008). Quantitative profiling of
2011   nucleotides and related phosphate-containing metabolites in cultured mammalian cells
2012   by liquid chromatography tandem electrospray mass spectrometry. J. Chromatogr. B
2013   Analyt. Technol. Biomed. Life Sci. *871*, 115-124. 10.1016/j.jchromb.2008.07.005.
2014 89. Gooding, J.R., Jensen, M.V., Dai, X., Wenner, B.R., Lu, D., Arumugam, R., Ferdaoussi,
2015   M., MacDonald, P.E., and Newgard, C.B. (2015). Adenylosuccinate is an insulin
2016   secretagogue derived from glucose-induced purine metabolism. Cell Rep. *13*, 157-167.
2017   10.1016/j.celrep.2015.08.072.
2018 90. Gonsalves, W.I., Broniowska, K., Jessen, E., Petterson, X.-M., Bush, A.G., Gransee, J.,
2019   Lacy, M.Q., Hitosugi, T., and Kumar, S.K. (2020). Metabolomic and lipidomic profiling of
2020   bone marrow plasma differentiates patients with monoclonal gammopathy of
2021   undetermined significance from multiple myeloma. Scientific reports *10*, 10250.

91. Lanza, I.R., Zhang, S., Ward, L.E., Karakelides, H., Raftery, D., and Nair, K.S. (2010).
Quantitative metabolomics by 1H-NMR and LC-MS/MS confirms altered metabolic
pathways in diabetes. PloS one *5*, e10538.
92. Dutta, T., Kudva, Y.C., Persson, X.-M.T., Schenck, L.A., Ford, G.C., Singh, R.J., Carter,
R., and Nair, K.S. (2016). Impact of long-term poor and good glycemic control on
metabolomics alterations in type 1 diabetic people. The Journal of Clinical Endocrinology
& Metabolism *101*, 1023-1033.
93. Wilkins, J., Sakrikar, D., Petterson, X.-M., Lanza, I.R., and Trushina, E. (2019). A
comprehensive protocol for multiplatform metabolomics analysis in patient-derived skin
fibroblasts. Metabolomics *15*, 83.
94. Collins, J.M., Jones, D.P., Sharma, A., Khadka, M., Liu, K.H., Kempker, R.R., Prideaux,
B., Maner-Smith, K., Tukvadze, N., Shah, N.S., et al. (2021). TCA cycle remodeling
drives proinflammatory signaling in humans with pulmonary tuberculosis. PLoS Pathog.
17, e1009941. 10.1371/journal.ppat.1009941.
95. Maner-Smith, K.M., Goll, J.B., Khadka, M., Jensen, T.L., Colucci, J.K., Gelber, C.E.,
Albert, C.J., Bosinger, S.E., Franke, J.D., Natrajan, M., et al. (2020). Alterations in the
human plasma lipidome in response to tularemia vaccination. Vaccines (Basel) *8*, 414.
10.3390/vaccines8030414.
96. Harrell, F.E., Jr. (2003). Hmisc: Harrell Miscellaneous. The R Foundation.
97. Wei, T., and Simko, V. (2010). Corrplot: Visualization of a correlation matrix. The R
Foundation.
98. Kachman, M., Habra, H., Duren, W., Wigginton, J., Sajjakulnukit, P., Michailidis, G.,
Burant, C., and Karnovsky, A. (2020). Deep annotation of untargeted LC-MS
metabolomics data with Binner. Bioinformatics *36*, 1801-1806.
10.1093/bioinformatics/btz798.
99. Fan, S., Kind, T., Cajka, T., Hazen, S.L., Tang, W.H.W., Kaddurah-Daouk, R., Irvin, M.R.,
Arnett, D.K., Barupal, D.K., and Fiehn, O. (2019). Systematic error removal using
random forest for normalizing large-scale untargeted lipidomics data. Anal. Chem. *91*,
3590-3596. 10.1021/acs.analchem.8b05592.
100. Sumner, L.W., Amberg, A., Barrett, D., Beale, M.H., Beger, R., Daykin, C.A., Fan,
T.W.M., Fiehn, O., Goodacre, R., Griffin, J.L., et al. (2007). Proposed minimum reporting
standards for chemical analysis Chemical Analysis Working Group (CAWG)
Metabolomics Standards Initiative (MSI). Metabolomics *3*, 211-221. 10.1007/s11306-
007-0082-2.
101. Anderson, B.G., Raskind, A., Habra, H., Kennedy, R.T., and Evans, C.R. (2021).
Modifying chromatography conditions for improved unknown feature identification in
untargeted metabolomics. Anal. Chem. *93*, 15840-15849.
10.1021/acs.analchem.1c02149.
102. Fahy, E., Subramaniam, S., Brown, H.A., Glass, C.K., Merrill, A.H., Jr., Murphy, R.C.,
Raetz, C.R.H., Russell, D.W., Seyama, Y., Shaw, W., et al. (2005). A comprehensive
classification system for lipids. Eur. J. Lipid Sci. Technol. *107*, 337-364.
10.1002/ejlt.200405001.

103. Foldnes, N., and Olsson, U.H. (2016). A simple simulation technique for nonnormal data
with prespecified skewness, kurtosis, and covariance matrix. Multivariate Behav. Res. *51*,
207-219. 10.1080/00273171.2015.1133274.
104. Tahir, U.A., Katz, D.H., Avila-Pachecho, J., Bick, A.G., Pampana, A., Robbins, J.M., Yu,
Z., Chen, Z.-Z., Benson, M.D., Cruz, D.E., et al. (2022). Whole Genome Association
Study of the Plasma Metabolome Identifies Metabolites Linked to Cardiometabolic
Disease in Black Individuals. Nat. Commun. *13*, 4923. 10.1038/s41467-022-32275-3.
105. Katz, D.H., Tahir, U.A., Bick, A.G., Pampana, A., Ngo, D., Benson, M.D., Yu, Z., Robbins,
J.M., Chen, Z.-Z., Cruz, D.E., et al. (2022). Whole Genome Sequence Analysis of the
Plasma Proteome in Black Adults Provides Novel Insights Into Cardiovascular Disease.
Circulation *145*, 357-370. 10.1161/CIRCULATIONAHA.121.055117.
106. Benjamini, Y., and Hochberg, Y. (1995). Controlling the false discovery rate: a practical
and powerful approach to multiple testing. Journal of the royal statistical society series b-
methodological *57*, 289-300. 10.1111/J.2517-6161.1995.TB02031.X.
107. Dyer, S.C., Austine-Orimoloye, O., Azov, A.G., Barba, M., Barnes, I., Barrera-Enriquez,
V.P., Becker, A., Bennett, R., Beracochea, M., Berry, A., et al. (2025). Ensembl 2025.
Nucleic Acids Res. *53*, D948-D957. 10.1093/nar/gkae1071.
108. Hornbeck, P.V., Zhang, B., Murray, B., Kornhauser, J.M., Latham, V., and Skrzypek, E.
(2015). PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. Nucleic Acids Res
43, D512-520. 10.1093/nar/gku1267.
109. Wang, Q., Li, M., Wu, T., Zhan, L., Li, L., Chen, M., Xie, W., Xie, Z., Hu, E., Xu, S., and
Yu, G. (2022). Exploring epigenomic datasets by ChIPseeker. Curr. Protoc. *2*, e585.
10.1002/cpz1.585.
110. Yu, G., Wang, L.-G., and He, Q.-Y. (2015). ChIPseeker: an R/Bioconductor package for
ChIP peak annotation, comparison and visualization. Bioinformatics *31*, 2382-2383.
10.1093/bioinformatics/btv145.
111. Steffen Durinck, W.H. (2017). biomaRt (Bioconductor).
112. Workbench, M. (2016). PR000001. 10.21228/M8159B.
113. Durinck, S., Spellman, P.T., Birney, E., and Huber, W. (2009). Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt. Nat. Protoc.
4, 1184-1191. 10.1038/nprot.2009.97.
114. Tenenbaum, D., and Maintainer, B.P. (2025). KEGGREST: Client-side REST access to
the Kyoto Encyclopedia of Genes and Genomes (KEGG).
115. Fahy, E., and Subramaniam, S. (2020). RefMet: a reference nomenclature for
metabolomics. Nat. Methods *17*, 1173-1174. 10.1038/s41592-020-01009-y.
116. Hu, C., Li, T., Xu, Y., Zhang, X., Li, F., Bai, J., Chen, J., Jiang, W., Yang, K., Ou, Q., et al.
(2023). CellMarker 2.0: an updated database of manually curated cell markers in
human/mouse and web tools based on scRNA-seq data. Nucleic Acids Res. *51*, D870-
D876. 10.1093/nar/gkac947.
117. Rath, S., Sharma, R., Gupta, R., Ast, T., Chan, C., Durham, T.J., Goodman, R.P.,
Grabarek, Z., Haas, M.E., Hung, W.H.W., et al. (2021). MitoCarta3.0: an updated
mitochondrial proteome now with sub-organelle localization and pathway annotations.
Nucleic Acids Res *49*, D1541-D1547. 10.1093/nar/gkaa1011.

118. Liberzon, A., Birger, C., Thorvaldsdóttir, H., Ghandi, M., Mesirov, J.P., and Tamayo, P.
(2015). The Molecular Signatures Database (MSigDB) hallmark gene set collection. Cell
Syst. *1*, 417-425. 10.1016/j.cels.2015.12.004.
119. Krug, K., Mertins, P., Zhang, B., Hornbeck, P., Raju, R., Ahmad, R., Szucs, M., Mundt,
F., Forestier, D., Jane-Valbuena, J., et al. (2019). A curated resource for phosphosite-
specific signature analysis. Mol. Cell. Proteomics *18*, 576-593.
10.1074/mcp.TIR118.000943.
120. Sagendorf, T. (2025). TMSig: Tools for Molecular Signatures (Bioconductor).
121. Team, R.C. (2025). R: A Language and Environment for Statistical Computing.
122. Dunn, J.C. (1973). A fuzzy relative of the ISODATA process and its use in detecting
compact well-separated clusters. J. Cybern. *3*, 32-57. 10.1080/01969727308546046.
123. Bezdek, J.C. (1981). Pattern recognition with fuzzy objective function algorithms (Kluwer
Academic/Plenum).
124. Futschik, M.E., and Carlisle, B. (2005). Noise-robust soft clustering of gene expression
time-course data. J. Bioinform. Comput. Biol. *3*, 965-988. 10.1142/s0219720005001375.
125. Kumar, L., and E Futschik, M. (2007). Mfuzz: a software package for soft clustering of
microarray data. Bioinformation *2*, 5-7. 10.6026/97320630002005.
2124

Fig 3A

Fig 3B

Fig 3C

Fig 3D

Fig 3E

Fig 3F

Fig 3G

Fig 4A

Whytock et al., 2022, 2024

Modality
EE
RE

Timepoint
Post 45 min
Post 4 hr
Post 24 hr

z.std
7
0
-4.8

BH Adjusted P-Value
< 0.05
≥ 0.05

Fig 4B

Fig 4C

EE (Transcriptomics + Metabolomics)

Fig 4D

RE (Transcriptomics + Metabolomics)

Fig 7A

Fig 7B

Fig 7C

Fig 7D

Fig 7E

Fig 7F

Fig 7G

Fig 7H

Fig S1A

Fig S1B

Figure S1. Intercorrelation of clinical/sub-clinical variables and sexual dimorphism in ASAT cell type proportions at baseline (Supplement to Figure 1)

A) Intercorrelation heatmap across clinical and subclinical variables. "p<0.05, **p<0.01, ***p<0.001. B) Sex-based comparisons of deconvoluted ASAT cell type proportions. BMI, Body mass index; WC, Waist circumference; SBP, Systolic blood pressure; DBP, DIastolic blood pressure; HR, Heart rate; NEFA, Non-esterifed fatty acid; Total_C, Total cholesterol; HDL, High-density lipoprotein; LDL, Low-density lipoprotein.

Figure S2. Differential analysis and enrichment analysis (Supplement to Figure 2)
A) Volcano plots showing differentially regulated features across omic layers and timepoints in CON, and in CON-unadjusted EE and RE groups. B) Temporal expression patterns of core circadian transcripts (CLOCK, NR1D1/2, and PER1/2, and DBP) in response to exercise. C) Acute exercise effects on core clock genes from previous reports (Fabre et al., 2018 and Ahn et al., 2025). In Fabre et al., 15 sedentary males without obesity performed 15min of moderate intensity endurance exercise (80% VO_2 max) on a cycle ergometer. ASAT sampling occurred 4 hr post-exercise. In Ahn et al., 15 regular exercisers (7 females and 8 males) without obesity performed 45min of moderate intensity endurance exercise (65% VO_2 max) on a cycle ergometer. ASAT sampling occurred 1.5 hr post-exercise. In both studies, ASAT transcriptome was profiles via bulk RNA sequencing. Bar plots show changes in core clock genes post-exercise in two studies. *adjusted p<0.05, **adjusted p<0.01, ***adjusted p<0.001. D) CAMERA-PR analysis of transcriptomic, E) proteomic, and F) metabolomic changes post-exercise. G) ORA results on collapsed differentially regulated phosphoproteins, stratified by directionality. H) Heatmap illustrating exercise responses on TNS1 phosphorylation at post 4 hr. *adjusted p<0.05, †adjusted p<0.1. I) Spearman correlation between protein abundance of ROCK1 and TNS1.

Fig S3

Figure S3. Comparison of EE vs. RE (Supplementary to Figure 3)
Scatter plots comparing logFC between EE and RE for features differentially regulated (adjusted p<0.05) in EE or RE.d p<0.05.

Fig S4A

Fig S4B

Fig S4C

Figure S4. Cell type deconvolution and enrichment (Supplement to Figure 4)
A) Estimated ASAT cell type proportions (%) from deconvolution analysis at each timepoint, separated by exercise modality. No significant modality × timepoint interaction or main effect was detected. B) Volcano plots of differentially regulated phosphosites in EE vs. RE, focusing on features annotated as cell-type markers. logFC > 0 indicates greater enrichment in EE; logFC < 0 indicates greater enrichment in RE. C) Temporal expression patterns of *NR4A1* and histamine in ASAT.

Figure S5. Integrative network downstream correlation analyses (Supplement to Figure 5)
A) Heatmap illustrating cross-omic correlation of constructed modules. *p<0.05. B) Heatmap showing unadjusted biweight midcorrelations between module eigengenes and estimated ASAT cell type proportions from baseline deconvolution. *p<0.05; **p<0.01; ***p<0.001. C) Heatmap showing biweight midcorrelations between module eigengenes and clinical traits, adjusted for sex, age group, and waist circumference (WC). Covariate adjustment was performed by regressing both traits and eigengenes, and residuals were used for correlation. *p<0.05; **p<0.01; ***p<0.001.

Figure S6. Relationship between ASAT feature abundance and corresponding plasma feature abundance at baseline (Supplementary to Figure 6)

A) Heatmaps illustrating exercise responses on filtered exerkine candidates. *adjusted p<0.05. B) Biweight midcorrelation between ASAT-exerkine transcript expression and corresponding plasma protein abundance at baseline. *p<0.05. C) Biweight midcorrelation between ASAT-exerkine metabolites and corresponding plasma metabolites at baseline. *p<0.05, **p<0.01, ***p<0.001

Fig S7A

Fig S7B

Figure S7. ASAT-exerkine candidates in endurance training (Supplementary to Figure 7)
A) Paired T-test results comparing Ssec for each WATSC-to-target tissue connection between CON and TR8W. Pink boxplot refers to CON. Dark green boxplot refers to TR8W. B) Heatmap showing Ssec difference of 60 ASAT-secreted candidates between TR8W vs. CON. Positive z-scaled Ssec diff indicates higher Ssec observed in the 8-week trained rats compared with 8-week untrained rats for the given WATSC-to-target tissue connection.