Assembling a fully-dated complete tree of life

Jonathan D. Duke[1,2]*, Jialiang Guo[2,3], Félix Forest[4], Rikki Gumbs[5], Emily Jane McTavish[6¶],

James Rosindell[2¶]

[1] Science and Solutions for a Changing Planet DTP, Grantham Institute, Imperial College

London, London, United Kingdom

[2] Department of Life Sciences, Imperial College London, Silwood Park Campus, Ascot,

Berkshire, United Kingdom

[3] PTN Program, Academy for Interdisciplinary Studies, Peking University, Haidian District,

Beijing, China

[4] Royal Botanic Gardens, Kew, Richmond, United Kingdom

[5] Institute of Zoology, Zoological Society of London, Regent's Park, London, United

Kingdom

[6] Department of Life and Environmental Sciences, School of Natural Sciences, University

of California, Merced, California, United States of America

* Corresponding author

E-mail: j.duke24@imperial.ac.uk (JDD)

¶ These authors contributed equally to this work.

Abstract

Time-scaled phylogenetic trees summarising evolutionary relationships are fundamental to many analyses in biology, from diversification rate estimation to conservation prioritisation. The most comprehensive available summary of these relationships, the Open Tree of Life, synthesises information from over two thousand studies into a supertree covering the full range of global biodiversity, but its use in downstream analyses is limited by the lack of divergence times. Previous work has mapped dates from Open Tree's database of trees to certain nodes in the supertree, but for the majority of nodes no date is available. While algorithms exist to interpolate missing dates in a tree, we found that their time and memory requirements scaled quadratically with the number of nodes, which made it computationally infeasible to run them on the entire tree. In this work, we describe novel date interpolation algorithms that scale linearly with the number of nodes. These enabled us to produce a distribution of fully-dated trees containing 2.3 million extant described species, greatly expanding the scope of feasible phylogenetic analyses. We illustrate the utility of these trees by computing the most robust estimate yet of the phylogenetic diversity of the complete tree of life, incorporating both topological and temporal uncertainty.

Introduction

The tree of life has been an object of study for centuries [1, 2], telling the story of how the diverse array of species on Earth came into being. The advent of genomic sequencing has provided vast datasets [3, 4], which have been analyzed to map out ever-larger clades of the tree [5-7]. Many fundamental questions in biology require knowledge of the evolutionary history presented in the tree. For example, speciation, extinction, and the evolution of traits are all studied using a dated tree [8-10]. Ecological analyses must account for the shared ancestry of the species they examine [11-15]. Missing data in trait databases can be imputed using phylogenetic correlations [16-18]. And in conservation, the tree-based phylogenetic diversity (PD) measure has found widespread use [19, 20].

A complete understanding of evolution requires a complete tree. Direct estimation of the entire tree from genomic data has not yet been achieved, owing both to insufficient data and to the computational infeasibility of fitting phylogenetic statistical models on such a scale. Rather than infer the entire tree using a single model, the Open Tree of Life project instead synthesises a tree by collating and summarising individual studies [21, 22], producing the most complete tree of life available today. The Open Tree synthesis algorithm stitches together over 2,000 trees from a curated database to construct a backbone tree of 150,000 species [23, 24]. This phylogenetic synthesis is combined with their taxonomic database [25, 26], producing a "supertree" that covers 2.3 million described species.

The Open Tree team make regular updates to the tree incorporating new information, improving phylogenetic coverage over time. At present, however, substantial parts of the tree come primarily from taxonomic data, resulting in extensive polytomies [27] that can produce biases in downstream analyses [28, 29]. While the true fully-resolved topology is unknown and the Open Tree synthesis algorithm explicitly does not attempt to estimate it [24], for many analyses the varying degrees of topological uncertainty across the tree could be more transparently reflected using trees sampled from a set of plausible topologies to produce distributions of possible solutions [28, 30, 31].

Crucially, the Open Tree also lacks divergence times, i.e. the estimated historical dates
of the tree's nodes, which represent common ancestors or speciation events. The lengths
of the branches between nodes in units of time are used to compute phylogenetic
correlations, phylogenetic diversity, and to associate evolutionary events with geological
or climatic changes over the history of Earth. Previous work has mapped dates from Open
Tree's database of underlying phylogenetic trees to nodes in the supertree [6, 32, 33], but
the database includes dates for only 77,000 internal nodes. Work is ongoing to improve
coverage by incorporating more published dated trees - where they are available
electronically - into the Open Tree database, but for the majority of nodes in the complete
tree, no dates have yet been estimated.

Dating the tree therefore requires interpolation between dated nodes on a sparsely-dated
tree with no branch lengths. Such interpolation methods exist [34, 35], but in their
present implementations, the computational time and memory required to run them
scale with the square of the number of nodes in the tree [34]. This makes interpolation
across the millions of nodes in the entire tree of life infeasible. Methods that scale linearly
with the number of nodes have been developed for applications such as smoothing
molecular substitution rates across a tree with branch lengths in units of molecular
substitutions [36, 37], but to our knowledge the dynamic programming techniques used
by these algorithms have not been applied to interpolating dates across topology-only
trees.

In this work, we describe stochastic algorithms that resolve the polytomies in the Open
Tree by placing taxa with only taxonomic information into plausible locations in the
phylogenetic backbone, automating the methods of commonly-used tree-completion
tools [e.g., 38, 39, 40]. We use our algorithms to produce a distribution of bifurcating trees
covering all extant described species. We then compare five linear-time methods for
interpolating dates across any tree, and apply them to our distribution of bifurcating
trees, resulting in a set of fully-dated complete trees of life.

93 Results

94 Fully-dated complete trees

Our date interpolation algorithms ran in times that scaled linearly with the number of tree
nodes (**Fig 1**), which made it computationally feasible for us to produce bifurcating fully-
dated trees at the scale of the entire tree of life. For example, interpolating dates across
a tree of 2.3 million species using an "equal splits" interpolation method took 26 seconds
(**Fig 1**). In comparison, the same method as implemented in the BLADJ algorithm in the
software Phylocom scaled in both run time and memory requirement with the square of
the number of nodes (**Fig 1**). We were unable to run it successfully on a tree of 2.3 million
species owing to a memory requirement of 50 terabytes.

Fig 1. Run times for interpolation algorithms that space dates evenly along undated paths between input dates. The time taken in minutes to interpolate dates across a tree is plotted against the number of nodes in that tree, for (blue circles) the BLADJ implementation in Phylocom; (orange triangles) our implementation of the same algorithm, which we call EQS-L; (green crosses) our implementation including our improved data cleaning algorithm to make the input dates time-consistent. Each triplet of points at a given number of nodes represents the same subtree extracted from a partially-dated complete tree. The blue dotted line illustrates a growth trend in run times increasing with the square of the number of nodes, while the orange dashed line and green dash-dot line increase linearly with the number of nodes. The three blue open circles show hypothetical run times imputed from the trend, because Phylocom's BLADJ implementation did not run successfully on trees of such size - on the complete tree, our projected run time is 14 days, but in practice the run failed owing to a memory requirement of 50 terabytes.

Achieving temporal consistency of input dates

Our interpolation used calibration dates summarised from 334 source trees. The source trees differed from each other and from the supertree in their calibration dates, dating methodologies and topologies, so when these dates were mapped to the supertree they were not time-consistent. That is, some nodes were assigned dates younger than the

date assigned to a descendant node or older than an ancestor. The BLADJ
implementation includes a data cleaning step that ensures time consistency by
traversing the tree from root to leaves and removing the date from any node that has a
younger ancestor. In our tree, this step removed 9,121 of the 76,953 dates. Our alternative
algorithm (see Materials and methods) removes only the minimum number of dates
necessary to achieve time consistency across the tree. It removed only 5,000 dates,
retaining more of the original information while still operating with linear time complexity
(**Fig 1**).

Comparison of interpolation algorithms

We compared five approaches for interpolating missing dates on sparsely-dated trees
with no branch lengths. We tested the *equal splits* algorithm, which spaces nodes evenly
along undated paths between input dates; we gave this the abbreviation EQS, after Kuhn
et al. [28]. Sometimes, the algorithm must interpolate along several paths from the same
start node, all of which end in a leaf node. In this case, the choice of which path is
interpolated first will affect the result (see Materials and methods for further explanation
and details of all approaches). We therefore tested two variants of the EQS approach: one
which interpolates first along the longest available path (we call this EQS-L; this is the
approach used by the BLADJ algorithm in the Phylocom software) and one which
interpolates first along the shortest path (EQS-S). Our third approach used a linear
combination of the dates obtained from these methods, with the parameter choice of
0.25*EQS-L + 0.75*EQS-S determined by testing the algorithms on small fully-dated
trees. We called this linear combination EQS-LS. We also considered two approaches
that assign dates using the expected branch lengths from a birth model with
exponentially-distributed waiting times: the *log-N* (LnN) approach, which works down
each lineage in a clade independently and is deterministic, and an approach we called
birth model (BM), which randomly chooses the order of speciation events across an
undated clade.

To compare the algorithms, we first used four fully-dated trees that had been produced
using molecular data for every included species (see Materials and methods for details).

In each tree, we removed some of the input dates at random and applied the interpolation algorithms in turn to estimate dates for the undated nodes. We then computed the date reproduction error. We repeated this analysis for seven clades of our fully-resolved complete tree, each of which had a crown age in our input data and between 19% and 73% of internal nodes dated. For all trees, we also computed mean and median pendant edge lengths, to examine how closely the algorithms maintained the distribution of branch lengths in the tree.

We found that, in trees where fewer than 70% of nodes needed to be interpolated, the five interpolation methods all performed similarly, with an average absolute reproduction error of 1.5-3% of the crown age (**Fig 2** row 1). For trees where more than 70% of nodes required an interpolated date, the EQS-LS approach produced the lowest reproduction error of 3-4%, while the error for the other approaches increased more rapidly with the degree of interpolation, especially when interpolating more than 95% of dates.

The EQS-LS algorithm also kept the distribution of pendant edge lengths more stable than the other four approaches as the degree of interpolation rose (**Fig 2** rows 2-3). For analyses in the remainder of this section, we will use dates computed by the EQS-LS approach, but we have made distributions of trees available for both the EQS-LS algorithm and the birth model algorithm (see Data availability). A full breakdown of the results for each individual tree and algorithm is given in the supplementary material (S1 Fig, S2 Fig).

Fig 2: Performance of date interpolation algorithms. Row 1: the average error in date reproduction tests across (column A) four fully-dated phylogenetic trees and (column B) seven subtrees of one sample tree in our distribution of fully-resolved complete trees of life. We removed a proportion of calibration dates at random from the tree, then interpolated dates across all undated internal nodes, and compared the interpolated dates with the original dates we had removed. The full datasets are supplied in the supplementary materials. Row 2: mean pendant edge length in the tree, relative to (for column A) the mean pendant edge length in the fully-dated tree, or (for column B) the mean pendant edge length in the subtree when using all available calibration dates and missing dates interpolated using the EQS-LS algorithm. Row 3: as row 2, but for median pendant edge lengths rather than mean.

Phylogenetic diversity in the complete tree

Our linear-time algorithms allowed us to estimate dates for all 2.3 million internal nodes of a complete bifurcating tree of extant described species, and therefore to estimate phylogenetic diversity for the entire tree and for any clade (**Fig 3**). We made use of phylogenetic diversity as a summary statistic to explore two dimensions of uncertainty around our trees.

Fig 3. A pruned fully-dated complete phylogenetic tree of life from our distribution. The branch colours represent phylogenetic diversity (PD), illustrating how the PD of all life (upper right) is divided among clades in the tree. The underlying tree is ultrametric, but it is too large to display here. Instead we show a tree pruned to major clades: either infrakingdoms (where that rank exists in the Open Tree Taxonomy) or phyla, or the first taxon with rank lower than phylum if there is no phylum rootwards of it in the tree. The square nodes indicate the crown node and crown age of each clade. Circled nodes have calibration dates associated with them. PD estimates for selected clades of interest are given to the right of the colour bar; these estimates do not include the stem back to the origin of life.

First, we considered topological variation. We sampled 501 plausible topologies for the

complete tree using our stochastic polytomy resolution algorithms (see Methods and

materials). Where we had multiple sources for the date on a node, we used the median

date. This produced a median phylogenetic diversity estimate for the complete tree of

39.6 trillion years (95% CI: 38.4-41.3 tyr; **Fig 4**). Highly-studied vertebrate and plant

clades displayed very little topological variation (**Fig 5**), for example Chordata (98.5-

101.4% of the median for the clade) and Tracheophyta (98.4%-101.6%). Most clades

showed considerably wider variation (**Fig 5**), for example phylum Mollusca (82.5-116.1%)

and phylum Rhodophyta (84.7-127.3%).

Clade	A: Variation from topology		B: Variation from date sources		C: Variation from both	
	Median PD (Byr)	95% CI	Median PD (Byr)	95% CI	Median PD (Byr)	95% CI
All Life	39556	38375 - 41321	39696	38907 - 40591	39783	38152 - 41874
Eukaryota	28754	28371 - 29177	28976	28187 - 29870	28914	27945 - 29927
Metazoa	19384	19075 - 19683	19354	18964 - 20224	19385	18776 - 20303
Chordata	1081	1072 - 1092	1094	1066 - 1151	1098	1065 - 1152
Actinopteri	407	403 - 411	410	402 - 419	410	402 - 419
Squamata	174	172 - 177	175	158 - 221	177	159 - 223
Aves	91	90 - 91	96	78 - 106	96	79 - 106
Amphibia	128	125 - 130	130	128 - 131	127	124 - 130
Mammalia	55	55 - 56	57	48 - 66	57	49 - 66
Chondrichthyes	52	50 - 53	51	51 - 52	52	51 - 54
Testudines	6	6 - 6	6	6 - 6	6	6 - 6
Arthropoda	12770	12548 - 12986	12542	12205 - 13416	12676	12183 - 13604
Pancrustacea	9994	9822 - 10187	9677	9365 - 10556	9877	9446 - 10807
Insecta	8531	8359 - 8706	8187	7910 - 9078	8389	7968 - 9328
Coleoptera	2867	2726 - 3029	2408	2296 - 3298	2583	2304 - 3537
Lepidoptera	785	741 - 817	778	738 - 880	793	726 - 921
Diptera	1286	1251 - 1319	1261	1229 - 1316	1314	1242 - 1388
Hymenoptera	1084	1059 - 1109	1081	1070 - 1094	1085	1059 - 1115
Hemiptera	1241	1187 - 1310	1299	1270 - 1320	1258	1192 - 1338
Odonata	87	84 - 90	85	85 - 86	88	84 - 91
Malacostraca	580	547 - 615	554	540 - 594	584	530 - 637
Copepoda	464	427 - 501	473	471 - 506	465	417 - 512
Branchiopoda	49	45 - 54	50	50 - 51	49	45 - 54
Chelicerata	2186	2070 - 2272	2278	2255 - 2307	2198	2077 - 2293
Myriapoda	592	551 - 621	585	548 - 586	577	530 - 633
Mollusca	1754	1536 - 1933	1895	1895 - 1898	1754	1539 - 1931
Platyhelminthes	677	616 - 731	652	651 - 657	677	617 - 730
Annelida	672	638 - 700	678	621 - 685	670	606 - 700
Nematoda	576	558 - 597	568	568 - 568	576	558 - 598
Cnidaria	572	550 - 591	570	552 - 605	571	540 - 617
Porifera	366	353 - 380	371	365 - 387	364	346 - 384
Echinodermata	247	237 - 259	247	229 - 247	243	223 - 256
Bryozoa	151	145 - 156	151	151 - 151	151	145 - 156
Tardigrada	45	42 - 48	57	47 - 57	53	43 - 58
Chloroplastida	2689	2642 - 2738	2619	2456 - 2997	2803	2452 - 3020
Tracheophyta	1880	1859 - 1898	1813	1656 - 2190	2010	1650 - 2191
Magnoliopsida	1661	1642 - 1679	1589	1430 - 1966	1787	1429 - 1971
Polypodiopsida	139	136 - 142	140	138 - 142	140	137 - 145
Bryophyta	152	147 - 158	148	147 - 151	152	146 - 160
Marchantiophyta	147	138 - 158	143	141 - 158	152	137 - 170
Chlorophyta	373	336 - 413	369	369 - 369	373	336 - 413
Rhodophyta	307	277 - 361	320	320 - 320	307	277 - 361
Fungi	3667	3494 - 3865	3636	3468 - 3783	3678	3396 - 3955
Ascomycota	2244	2073 - 2424	2312	2150 - 2312	2236	2020 - 2455
Basidiomycota	1000	949 - 1054	1009	893 - 1016	996	904 - 1118
Microsporidia	134	123 - 147	140	134 - 162	142	125 - 179
SAR	2026	1893 - 2114	2080	2080 - 2082	2026	1893 - 2115
Stramenopiles	379	360 - 401	364	364 - 364	379	360 - 401
Alveolata	862	824 - 898	850	850 - 852	862	825 - 898
Rhizaria	785	660 - 855	865	865 - 865	785	657 - 855
Archaea	454	421 - 488	459	459 - 459	454	421 - 488
Bacteria	10313	9297 - 11973	10259	10259 - 10259	10313	9295 - 11979

Fig 4: Table of phylogenetic diversity for selected clades of the tree. Panel A: PD owing to topological variation, from 501 different tree topologies using the median of the available dates for each node. Panel B: PD using a single tree topology, sampling 501 times from the available date sources. We used the topology corresponding to the median PD for the complete tree in the topological distribution. Panel C: PD from 1503 trees: 501 different topologies, with 3 sets of date source samples for each. In the hierarchy listed on the left, the clades chosen are a non-exhaustive subset of their parent clade.

Fig 5. Phylogenetic diversity distributions for selected large clades of the tree. Panel A:
distributions of PD owing to topological variation, from 501 different tree topologies using the median
of the available dates for each node. Panel B: distributions of PD using a single tree topology, sampling
501 times from the available date sources. We used the topology corresponding to the median PD for
the complete tree in the topological distribution. These PD distributions are shown relative to a
"baseline" PD computed using median dates. Panel C: PD distributions from 1503 trees: 501 different
topologies, with 3 sets of date source samples for each. In the hierarchy listed on the left, the clades
chosen are a non-exhaustive subset of their parent clade.

Second, we considered temporal variation. Instead of using the median of the available

date sources at each node, we randomly selected a single source (see Methods and

materials). We repeated this 501 times, producing 501 different PD estimates for one tree

with fixed topology. There was wide variation in PD in clades with a large number of

sources such as birds (class Aves, 84.7-121.2% of the median for the clade) and

mammals (class Mammalia, 84.5-123.0%) (**Fig 5**), while in nine clades there was no

variation at all because no node in the clade had more than one date source (**Fig 5**).

Finally, we explored both dimensions of uncertainty simultaneously, sampling both

topologies and date sources at random. We sampled 501 trees with three different sets

of dates for each tree, giving a distribution of 1503 trees. This produced a median PD

estimate of 39.8 trillion years (95% CI: 38.2-41.9 tyr; **Fig 4**).

Discussion

Fully-dated complete trees

The linear time complexity of our algorithms allowed us to resolve and date trees containing millions of species (**Fig 1**), which to the best of our knowledge has not previously been computationally feasible. Our trees incorporate all 2.3 million extant species in the Open Tree Taxonomy [25], which aims to include all formally-described species across all domains of life [41]. A fully-dated complete tree will be made available in the OneZoom tree viewer, which will allow users to explore the entire tree interactively [42].

To our knowledge, the largest fully-dated tree previously published is the TimeTree of Kumar et al. [43], an ongoing project that has the objective of including every species for which molecular data is available, and whose most recent published version in 2022 covered 137,306 species [44], about 6% of formally-described species. As of 2025, NCBI's GenBank database contained at least some genetic information for 581,000 described species [45], a number that has increased by 15% over the past year [45]. The ongoing growth in data availability is encouraging, but it is clear that a significant proportion of species will remain unsequenced for the foreseeable future [46]. Our objective in this work was to give researchers access to a distribution of time-calibrated trees for the entire tree of life or any desired subset. We argue that efficient imputation and interpolation algorithms capable of filling in gaps in molecular coverage are, and will remain, an important tool in this endeavour, though we recognise that the resulting trees will not be suitable for every purpose [47-49].

Interpolation quality

For our test trees or subtrees where at least 30% of nodes had input dates assigned to them, results for all five interpolation algorithms were similar (**Fig 2**). In this case, there is sufficient phylogenetic and temporal structure in the synthetic tree that the scope of

interpolation error is relatively limited. In practice, only the highly-studied vertebrate

clades have such strong coverage in the Open Tree database, with ferns coming close

(29%) and other vascular plants falling short (8%). Elsewhere, few clades have date

coverage of more 2%. Our analyses suggest that our EQS-LS algorithm is able to produce

a reasonable distribution of branch lengths in trees with 5-10% of nodes dated (**Fig 2**),

which represents a less daunting target than 30%. We hope that future researchers in

under-represented clades will be able to help improve coverage to this level, by making

their time-calibrated trees available electronically or contributing them directly to the

Open Tree project.

Phylogenetic diversity

Previous studies of phylogenetic diversity have mainly focused on vertebrate animals,

and in particular mammals (class Mammalia) and birds (class Aves). For mammals, our

median estimate of PD was 57 billion years, which is within the range of 37-66 billion

years found across five previous studies [20, 50-53]. In birds, our median estimate is 96

billion years, broadly in line with figures of 74-90 billion years in four prior studies [50, 52-

54]. A full list by clade of comparisons to previous studies is provided in the supporting

information (S1 Table).

To our knowledge, two other studies have estimated PD across the entire tree of life. The

review by Diaz and Malhi [52] cited PD estimates made by one of us (Rosindell) using a

heuristic based on approximate clade age and species richness. In prokaryotes, Diaz and

Malhi estimated PD of 221 trillion years across the two domains of Bacteria and Archaea,

based on an estimated 740,000 operational taxonomic units (OTUs) [55-57]. Our tree

instead contains only around 79,000 "species" of bacteria and archaea, as labelled in the

Open Tree Taxonomy, resulting in the much smaller PD estimate from our tree of 10 trillion

years (**Fig 5**). For eukaryotes, the PD estimate in Diaz and Malhi is 124 trillion years, based

on an assumption of log-linear growth in species numbers over time. This assumption

does not account for extinctions and will therefore tend to overestimate the ages of

common ancestors, resulting in longer terminal branches and higher PD estimates. This

explains why our PD estimate for eukaryotes of 29 trillion years is low by comparison.

The second study to include a PD estimate for the entire tree, by Guo et al. [53], did so by
computing species-level evolutionary distinctiveness (ED) scores [20], which can be
summed to find PD for a clade. The ED scores were found by interpolating branch lengths
along the path from each individual species to the tree root [53], without computing a
single consistently-dated tree. Summing the scores across all species, this method
found the total PD of all life to be 31 trillion years, compared to our median estimate of
39.8 trillion years with our EQS-LS interpolation algorithm. We applied the ED
interpolation method to our fully-resolved tree, finding a median PD of 38.2 trillion years.
This was very similar to our estimate, which indicates that the original difference resulted
less from the method and more from the differing tree topologies. The prior study used
the tree from the OneZoom tree viewer [42], which uses the Open Tree synthetic tree as
a starting point but replaces certain clades of the tree with curated alternatives and
resolves all polytomies purely at random [42]. We used only the base Open Tree and,
where possible, resolve polytomies by imputing taxonomy-derived groups into the
phylogenetic backbone, producing a different topology.

Sources of uncertainty

In clades with few calibration dates, the choice of date interpolation algorithm will have
a major effect on PD. Algorithms that create longer pendant edges will produce larger PD
estimates, and vice versa. Here, we have used the EQS-LS algorithm because it produced
the closest reproductions of dates and pendant edge lengths in the trees we tested (**Fig
2**). A distribution of trees produced using our alternative birth model interpolation
algorithm is also available (see Data availability).

Having chosen the EQS-LS algorithm, our estimates of phylogenetic diversity
incorporated two dimensions of uncertainty. First, we explored topological uncertainty:
the distribution of plausible topologies for the tree of life created by the stochastic
resolution of polytomies in the Open Tree. In the Open Tree synthesis algorithm,
polytomies are created when taxonomic data is attached to the phylogenetic backbone.
In clades with high phylogenetic coverage such as the vertebrates, there are few
polytomies to resolve, and so we saw very low topological variation (**Fig 5**). In clades that

323 come mainly from taxonomic data, such as molluscs (phylum Mollusca) and red algae

324 (phylum Rhodophyta), we saw much broader distributions (**Fig 5**).

325 Second, we considered the temporal uncertainty resulting from variation across the

326 available input dates. There are 76,935 nodes in the synthetic tree for which at least one

327 published study has both estimated a date and been included in the Open Tree database.

328 The majority of these nodes are in vertebrate animals and vascular plants. Here, nodes

329 frequently had dates from multiple sources, and we were able to sample from the range

330 of sources to reflect uncertainty (**Fig 5**), or use a median to summarise the information in

331 multiple studies. Outside of these clades, to our knowledge most nodes in the tree of life

332 have not been dated in any published study. For the small proportion of nodes where we

333 did have a date, we usually had only a single source, and therefore we were not able to

334 reflect uncertainty through sampling. This is a limitation in our inference of temporal

335 uncertainty: if data are lacking this should imply higher uncertainty, not lower. Future

336 work will include searching for studies not yet included in the Open Tree database,

337 focusing on clades with low date coverage. Many source trees also included information

338 on the uncertainty around their date estimates, but this data is not currently available in

339 the Open Tree database.

340 There are further sources of uncertainty that could be considered in future. The crown

341 ages of large, ancient clades where the descendant tree structure is derived largely from

342 taxonomy are of particular importance to our method, because the crown ages will

343 determine the dates interpolated across the entire clade. For example, the ages of the

344 common ancestor of all Bacteria and of the largest fungus phylum Ascomycota are likely

345 to have a significant effect on PD estimates for the complete tree, but we currently have

346 only a single source for both these dates, so our analysis does not reflect this sensitivity.

347 We have also treated the set of extant species in the Open Tree Taxonomy as a fixed input.

348 While the taxonomy attempts to incorporate all formally-described species, more

349 species remain undescribed than have been described [58, 59]. Further, those

350 undescribed species are likely to be concentrated in certain clades of the tree, such as

351 Arthropoda [58, 60] and Fungi [61]. The incorporation of these "missing" species into the

tree would alter the topology of some regions, and the interpolated dates and
downstream analyses, in ways not reflected in our topological distributions. A PD
estimate for the Arthropoda phylum that truly included every arthropod species on Earth
would in all likelihood lie well beyond the bounds of our distribution, whereas our PD
distribution for Chordata is likely closer to the truth.

Finally, we emphasise that our work summarises information from studies included in
Open Tree's database, but this does not include all published data. We are aware of dated
trees not yet included in Open Tree, and where we are able to gain access to these trees
in electronic form we plan to curate them into the Open Tree database, reducing
uncertainty by expanding phylogenetic coverage and increasing the number of date
calibration nodes. We are grateful to researchers who makes trees available
electronically or inform us of regions of the tree that do not conform with the latest
published work; indeed, we hope that researchers will consider contributing trees
directly to the Open Tree project, to improve the resources available for all.

Conclusion

We have designed and implemented algorithms to produce plausible candidates for the
topology of the bifurcating tree of all life, and to interpolate dates across the tree, allowing
us to produce fully-dated complete trees. We have used the trees to compute a
distribution of phylogenetic diversity estimates for the whole tree of life. Our set of trees,
as well as our polytomy resolution and date interpolation algorithms, are freely available
(see Data availability); the input data we used and the seeds for the random number
generator are included, making the trees reproducible.

Our work has built on the Open Tree of Life, the most complete tree presently available,
deriving bifurcating trees that we think will be useful to new audiences. The source
studies and algorithms used by Open Tree and Chronosynth are made available under
the FAIR principles [62], and new time-calibrated trees can be curated into the Open Tree
database by anyone with a GitHub account [23, 63]. The synthetic tree is regularly
updated to incorporate newly-added phylogenetic and taxonomic information, and the

computational efficiency of our algorithms will allow our distribution of trees to be
updated in sync. We hope this work can spark new interest from potential contributors,
who can further improve the Open Tree project for the benefit of the whole community,
as well as enabling novel research that can take advantage of fully-dated complete trees.

Materials and methods

Overview of methods

We based our work on the Open Tree Synthetic Tree version 16.1 [27], which incorporates
2.3 million species, and the Open Tree Taxonomy 3.7.3 [64], which contains additional
metadata about each taxon. We first pruned the tree to remove subspecies and extinct
taxa, so that every leaf node represented an extant species. We then stochastically
resolved the topology of the tree to produce a distribution of bifurcating trees, each one
of which represented a plausible hypothesis for the evolutionary history of life. Finally, we
mapped dates from the trees in the Open Tree database to the internal nodes in our trees
[23, 33], and designed scalable algorithms to interpolate dates across all remaining
undated internal nodes. These steps are summarised in **Fig 6**.

Fig 6. Summary of our methods for producing a distribution of dated trees. The iterative process shown will generate a distribution of trees incorporating both topological and temporal uncertainty. To explore only topological uncertainty, we used the median of available date sources at each node instead of sampling a new source with each iteration. To explore only temporal uncertainty, we used a single fixed topology instead of sampling a new topology with each iteration. Green boxes represent data inputs; white boxes summarise algorithmic processes detailed in the Materials and methods section of this paper; and blue boxes represent intermediate or final outputs.

Initial pruning of the tree

Beginning with the labelled supertree (labelled_supertree_ottnames.tre) from the Open Tree of Life version 16.1 [27], we pruned any below-species-level nodes from the tree, including subspecies, varieties, formae, and leaf nodes labelled "no rank - terminal" in the taxonomy. If below-species nodes were present without a parent node representing their species, we kept one of them at random as a representative of that species. We removed nodes whose name contained the strings "uncultured" or "unidentified", and removed 15 nodes whose name contained the string "intergeneric hybrids". We pruned the birds clade (class Aves) to include only species present in the synthetic tree in

McTavish et al. [6], which is aligned to the Clements taxonomy [65], avoiding known issues with this clade in the Open Tree Taxonomy [6]. Similarly, we pruned the order of turtles (Testudines) to include only species present in the Turtles of the World checklist [66]. We removed the "extinct" flag that had erroneously been added to the species *Homo sapiens* in the latest version of the Open Tree Taxonomy. Finally, we removed any nodes labelled as extinct in the taxonomy and removed any of their ancestral nodes which may not themselves have been labelled extinct but had only extinct nodes below them.

These steps pruned 84,990 leaf nodes and 22,502 of their ancestors from the Open Tree. In addition, there were 1,718 leaf nodes that represented ranks above species, primarily empty genera; for each of these, we imputed a child node with the name "[Genus name]_sp._ott0000" as an unnamed representative species. The resulting tree had 2,294,776 leaf nodes, all now representing extant species, and 315,958 internal nodes.

Topology

Our aim was to create a set of trees that were plausible hypotheses for the common descent of all life: that is, trees that respected the taxonomic hierarchy and were fully bifurcating. The steps to achieve this are described here, and illustrated using a small example tree (**Fig 7**).

Fig 7. Illustration of our topology resolution algorithms. Top: A small phylogenetic tree illustrating how nodes in the Open Tree supertree that were absent from any underlying tree (squares, labelled with coloured lower case letters, with dashed-line branches above) were imputed into the phylogenetically-derived backbone (grey circles, with solid-line branches above). In random order, each coloured square node will be inserted into one of the branches indicated by the matching upper-case letters. For example, the *Laphyctes* genus is labelled with a lower-case "e", and will be inserted into one of the branches labelled with an upper-case "E", which are chosen to respect taxonomic hierarchy: they could be within any existing family in the backbone but not an existing genus. If the branch above the original parent node (Passeriformes in this example of *Laphyctes*) is chosen, the node to be moved would become sister to the entire descendant clade, with the original parent node remaining ancestral to both. Polytomy resolutions of this type were performed in random order, with newly-inserted groups added to the backbone as resolution progressed. As a final step, the polytomy in the *Rhectes* genus was resolved by choosing a topology at random. Bottom: one possible resolution, which will always be a bifurcating tree with taxonomic hierarchy respected and no existing monophyletic groups interrupted.

The Open Tree supertree algorithm first synthesises a bifurcating backbone tree
comprising all the taxa included in their curated database of phylogenetic trees [23, 24].
Then, taxa in the Open Tree Taxonomy that are not present in any tree are connected to
the crown node of their clade in the backbone, introducing polytomies. Working from our
pruned tree, we used the information in the annotations.json file included with the Open
Tree to label each node as derived either from the phylogenetic backbone or from the
taxonomy. We then located all polytomies in the supertree and categorised them into
three groups depending on their origin, resolving each category separately.

First, there were polytomies where taxonomy-derived groups had been attached to a
phylogenetically-derived backbone node (this case is represented in **Fig 7** by the taxa
labelled with lower-case b, c, d and e). There were 40,607 polytomous backbone nodes
in this category. We resolved these using an approach similar to tree completion tools
such as addTaxa [38], randtip [39], and pez [40], which insert taxonomy-derived species
into randomly-chosen branches in a phylogenetic tree, applying certain location
constraints. In our case, owing to the scale of the tree, location constraints were found
algorithmically based on taxonomic information. For each polytomy, we detached the
taxonomy-derived group, and searched below the polytomous node for candidate
insertion points that would respect taxonomic hierarchy and would not interrupt any
existing monophyly (for example, in **Fig 7**, candidate insertion points for the
Colluricinclidae family are labelled with an upper-case D). We additionally included the
possibility of the detached group becoming a sister to all the existing children of the
backbone node. An insertion point was then selected randomly with uniform probability
among the candidates. Successive resolutions of this type were performed in random
order, with newly-inserted groups added to the backbone as resolution progressed.

Second, we resolved polytomies resulting from non-monophyletic genera in the
supertree. In the phylogenetically-derived backbone tree, species from a genus can be
placed into groups that also include other genera, if there was phylogenetic backing for
those relationships. The supertree algorithm then takes taxonomy-derived species in the

genus and attaches them to the most recent common ancestor of all their congeneric backbone species, creating a polytomy at that common ancestor. (This is illustrated in **Fig 7** by the Mirafra genus, where Mirafra pulpa has been attached to the common ancestor of the backbone Mirafra species). To resolve the 5,064 polytomies of this type we followed the approach of addTaxa [38]: we detached the taxonomy-derived species from the common ancestor, and imputed a new parent node for it into a branch in one of the congeneric monophyletic groups, choosing with uniform probability among all branches in all groups. (In **Fig 7**, the new location for M. pulpa is chosen uniformly among the branches labelled upper-case A.) As before, successive resolutions were performed in random order, with new branches added to the set of choices for potential insertion as resolution progressed.

Third, after resolving the previous two categories, there remained 74,402 polytomies where there was no phylogenetic information within a taxonomic group (as in the Rhectes genus in **Fig 7**). We resolved each of these by choosing a bifurcating topology uniformly at random from all possible topologies.

As a final step, we removed all unary nodes, which were created by the supertree algorithm only to provide taxonomic representation, producing a fully-bifurcating tree.

Note on domains of life in the Open Tree

The current version of the Open Tree supertree has the root node of all life as the parent of the three domains Bacteria, Archaea, and Eukaryota. The presence of Bacteria and Archaea as sisters is derived from underlying phylogenetic trees, while Eukaryota is added from the taxonomy. We resolved this polytomy by making Eukaryota a sister to Archaea prior to any other polytomy resolution.

Dating

Assigning dates to the trees

We used the Chronosynth library to assign dates from Open Tree's underlying database of trees, called the phylesystem [23], to nodes in our bifurcating tree [6, 33]. Working with commit e060993 of the phylesystem (27 December 2025), we ran Chronosynth's combine_ages_from_sources function, which searches the database for trees with branch lengths in millions of years, and then maps nodes in those source trees to nodes in the Open Tree synthetic tree by matching common ancestors [6].

We found 334 source trees (**Fig 8**; a full list of citations is given in S1 Text), and from these we were able to map at least one date to 76,953 nodes. The clade with the most source trees was birds (class Aves), with 129 sources in total and individual nodes having up to 44 sources (the crown nodes of Passeriformes and Galloanserae). In mammals (class Mammalia) there were 49 sources across the clade, with 15 date sources for the crown node of all mammals. Most dated nodes, however, had only one date source (48,938 nodes) or two (13,842 nodes) (S3 Fig).

Clade	Species richness	Dated nodes	Date coverage	Date source trees	Phylogenetic coverage
All Life	2,294,775	76,953	3.4%	334	6.8%
Eukaryota	2,215,634	76,915	3.5%	333	6.9%
Metazoa	1,475,149	41,196	2.8%	285	5.7%
Chordata	82,903	29,735	35.9%	225	47.4%
Actinopteri	37,408	11,182	29.9%	31	36.9%
Squamata	11,221	4,113	36.7%	16	40.9%
Aves	10,931	8,762	80.2%	129	93.4%
Amphibia	9,835	1,937	19.7%	7	57.3%
Mammalia	7,964	2,993	37.6%	49	50.1%
Chondrichthyes	1,653	536	32.4%	6	39.1%
Testudines	326	171	52.6%	5	96.8%
Arthropoda	1,183,222	10,532	0.9%	57	3.3%
Pancrustacea	1,068,467	10,287	1.0%	54	3.6%
Insecta	984,720	9,615	1.0%	48	3.6%
Coleoptera	286,855	658	0.2%	10	2.8%
Lepidoptera	182,577	2,208	1.2%	9	3.5%
Diptera	166,902	237	0.1%	4	2.2%
Hymenoptera	142,384	5,832	4.1%	16	6.8%
Hemiptera	103,698	255	0.2%	6	2.3%
Odonata	6,314	104	1.6%	5	18.6%
Malacostraca	48,796	589	1.2%	6	3.2%
Copepoda	15,764	0	0.0%	0	4.0%
Branchiopoda	2,468	25	1.0%	3	1.9%
Chelicerata	96,446	209	0.2%	4	0.8%
Myriapoda	18,309	34	0.2%	2	0.2%
Mollusca	88,048	213	0.2%	13	1.1%
Platyhelminthes	25,209	20	0.1%	1	1.0%
Annelida	22,978	13	0.1%	10	2.1%
Nematoda	19,802	95	0.5%	3	2.0%
Cnidaria	15,435	190	1.2%	11	14.1%
Porifera	10,134	38	0.4%	3	7.8%
Echinodermata	8,715	47	0.5%	11	3.1%
Bryozoa	7,299	287	3.9%	1	5.0%
Tardigrada	1,473	3	0.2%	1	0.6%
Chloroplastida	426,729	35,203	8.2%	48	15.1%
Tracheophyta	390,260	34,814	8.9%	46	16.1%
Magnoliopsida	373,449	30,700	8.2%	41	15.5%
Polypodiopsida	13,239	3,619	27.3%	6	29.7%
Bryophyta	16,977	204	1.2%	3	5.3%
Marchantiophyta	9,587	151	1.6%	4	3.4%
Chlorophyta	6,869	23	0.3%	1	2.4%
Rhodophyta	9,274	8	0.1%	1	1.6%
Fungi	247,150	396	0.2%	4	1.4%
Ascomycota	163,301	187	0.1%	3	1.2%
Basidiomycota	78,410	201	0.3%	2	1.8%
Microsporidia	1,466	0	0.0%	0	0.8%
SAR	51,095	76	0.1%	2	1.6%
Stramenopiles	20,088	46	0.2%	2	2.5%
Alveolata	14,857	18	0.1%	2	1.7%
Rhizaria	16,150	10	0.1%	2	0.3%
Archaea	1,507	6	0.4%	1	8.8%
Bacteria	77,634	31	0.0%	2	3.9%

Fig 8: Breakdown of the calibration date coverage and phylogenetic coverage in our bifurcating
tree. Date coverage is the percentage of internal nodes in the clade with at least one date source.
Figures are for the original dates as mapped by Chronosynth, before checking for time consistency. In
the hierarchy on the left, the clades chosen are a non-exhaustive subset of their parent clade.

Our date interpolation algorithms required a dated root node. We used a root date of 4247

million years ago (Mya), found in the TimeTree of Kumar et al. [44], who justified this date

as lying between younger and older constraints on possible estimates for the origin of life.

Their younger bound of >3970 Mya came from a molecular clock study of the divergence

of archaea and eukaryotes [67], while the older bound of <4440 Mya was the earliest

plausible cessation of asteroid impacts on the early Earth that evaporated the oceans

and would have prevented life from developing [68].

Algorithms for interpolating missing dates

Our goal was to interpolate a date on every undated internal node in a set of sparsely-dated trees with no branch lengths. We will refer here to "dated" and "undated" nodes, by which we mean internal nodes. All leaf nodes represented extant species and were assumed to have a date of zero. In describing our algorithms, we will use "down" to refer to movement from the root of the tree to its descendants toward the leaves, and "up" for the reverse movement from leaves to parents to ancestors toward the root.

We divided the interpolation into two distinct problems: interpolation between a dated node and one or more dated descendant internal nodes; and interpolation when a dated node had only undated internal nodes and leaf nodes below it. In the former case, paths from the dated node to each of its dated descendants must be considered to avoid negative branch lengths (see S4 Fig, and the discussion in the supplementary material of Kuhn et al. [28]). In the latter case, negative branch lengths can be avoided more straightforwardly by working down the tree and ensuring each undated node is assigned a date more recent than its parent. In our partially-dated bifurcating tree, 96.5% of undated nodes fell into the latter category.

We tested five interpolation algorithms, all of which we have implemented with run times that scale linearly with the number of nodes in a tree, making it feasible to use them across the entire tree of life. The first algorithm we will call equal splits or EQS, following Kuhn et al. [28], commonly used in the software Phylocom where its implementation is called branch length adjustment or BLADJ [34]. The algorithm assigns equal branch lengths to each branch along an undated path between a dated node and its oldest descendant. By identifying the oldest descendant, negative branch lengths are always avoided. For this reason, we applied only this algorithm to date the 3.5% of undated nodes located along undated paths between two dated internal nodes.

We compared the five interpolation algorithms across the remaining 96.5% of undated nodes, i.e. those located in clades where all internal nodes were undated except for a dated crown node. In these cases, there will be multiple undated paths covering the same time period (from the dated node to the leaves). For the EQS algorithm, the choice

of which path is interpolated first will affect the result. In the BLADJ implementation, the

path with the most nodes along it is dated first (**Fig 9**); we called this variant EQS-L, for

"longest path". We tested a second version of this where, instead of choosing the path

with the most nodes, we first dated the path with the fewest nodes (**Fig 9**). We called this

EQS-S, for "shortest path".

Fig 9: Examples of our five interpolation algorithms. The algorithms are illustrated in a clade where
the only calibration date is 12 Mya on the crown node. The EQS-L and EQS-S algorithms space nodes
evenly along undated paths, with the red dotted lines indicating the paths interpolated first - the longer
path for EQS-L and the shorter path for EQS-S. The EQS-L algorithm produces pendant edges that are
longer than or equal to the EQS-S algorithm. The EQS-LS algorithm computes dates as a linear
combination of the two: 0.25*EQS-L + 0.75*EQS-S. The LnN and BM algorithms are inspired by birth
models, with logarithmically-spaced dates. Here we have a crown age of 12 Mya, and a clade of six
species. The LnN ages are (date of parent node) * (number of species in this clade) / (number of
species in parent's clade), applied recursively from crown to leaves down each lineage independently.
The BM ages are (date of crown node) * (logarithm of the number of terminal lineages in the clade -
logarithm of the number of lineages in the clade rootwards of this node in time) / (logarithm of the
number of terminal lineages in the clade), so that the number of lineages through time in the clade
plotted on a semilog plot is straight line.

Our third algorithm was a linear combination of the dates from these two approaches,

inspired by observing that the EQS-L approach leads to very long pendant edges relative

to a fully-dated tree, while the EQS-S approach produces extremely short pendant edges.

To find a suitable combination, we used the four published trees described in the section

below on quality assessment, which had molecular data for all species. We found that

assigning dates as 0.25*(EQS-L date) + 0.75*(EQS-S date) minimised error relative to the

published dates. We called this algorithm EQS-LS. Since both input algorithms produce

a consistent set of dates across the tree, and the EQS-S dates must always be younger or

equal to the EQS-L dates, any linear combination of the two must also produce a

consistent set of dates.

The fourth algorithm we tested was proposed by Purvis and Nee [35], based on a birth or

birth-death model where the age of a clade is in expectation proportional to the logarithm

of the number of species in the clade. This method, which we called LnN following Kuhn

et al. [28], computes a date for each undated child node by multiplying the date of its

parent by the ratio of the logarithm of the species counts in the child clade and the parent

clade. Given a dated root of the clade, this can be applied recursively down a tree to date

all nodes (**Fig 9**).

These four algorithms are deterministic, i.e. given a tree topology and a set of dated nodes

they will produce the same result on any run. Our fifth algorithm was, like the LnN

approach, based on a birth model with exponentially-distributed waiting times, but

added a stochastic element. Rather than working independently down each lineage of a

clade as the LnN algorithm does, we instead considered all speciation events in the clade

and assigned their time ordering at random: working down a clade, the probability of a

lineage being the site of the next speciation event is proportional to that lineage's fraction

of the total remaining speciation events. The dates for the speciation events are chosen

so that the number of lineages grows exponentially over time (**Fig 9**); that is, a lineages-

through-time semilog plot would show a straight line. Specifically, the sequence of dates

is (logarithm of the number of terminal lineages in the clade - logarithm of the number of

lineages in the clade rootwards of this node in time) / (logarithm of the number of terminal

lineages in the clade). We called this the BM algorithm, for birth model.

Implementation and time complexity of date interpolation algorithms

We implemented all algorithms using a dynamic programming approach with run times

that scaled linearly with the number of nodes (**Fig 1**). Here we describe the

implementation of the EQS-L algorithm, and then summarise the modifications required

by the other four algorithms.

The first step in the EQS-L algorithm was to traverse the tree and label each undated node

with the information required to interpolate missing dates: we require the date of the

oldest descendant of the node, and the number of nodes along the path to that date (for

an illustrative example of the procedure described here, see **Fig 10**, top). We begin with

leaf nodes, which we label with the tuple (0, 0), representing a date of 0 and a path length

to that date of 0. These tuples are reported to each node's parent. Then, if a parent is

undated, we look at the tuples it received from its children, and label it with the tuple that

contained the oldest date, incrementing the path length in the tuple by one to represent

the additional node now in the path. If multiple tuples are received with the same date,

the tuple with the longest path is retained (for example, in **Fig 10**, the common ancestor

of species M and P stores the path length to M). If the parent is dated, it is labelled with a

tuple containing its own date and a path length of 0. In either case, the parent reports its

tuple to its own parent, and this process continues recursively up to the tree root.

Fig 10. Illustration of the EQS-L algorithm for evenly interpolating missing dates between known dates. Top: Leaf nodes are shown as small grey circles, and are given a date of zero. Nodes with dates assigned are shown as blue-green circles, with the date given in red text above the branches. The yellow-brown squares are the nodes to be dated. Each node is shown with the tuple reported upwards, containing: (oldest date reported to this node, path length down the tree to that date). If the oldest date reported is a tie, the longer path length is used; for example, the common ancestor of M and P retains the path towards M. For the "shortest path" EQS-S version of the algorithm, the path to P would be retained. Bottom: Working down the tree from the dated root node, each undated node is given a date spaced evenly along the path between its parent and the oldest date below it, which is stored in its label.

Once all internal nodes are labelled we begin again, this time from the root, working down
the tree and assigning a date to every undated node. Since we require a dated root node
and work down the tree, a node's parent can be assumed to be dated. We therefore space
each undated node evenly along the path between the date of its parent and the oldest
date beneath it, using the oldest date and path length information recorded in the node's
label (**Fig 10**, bottom). This completes the dating of the tree in linear time, using one
postorder and one preorder traversal.

For the shortest path EQS-S algorithm, the procedure is the same except that when a
node has received multiple tuples with the same date, it retains the tuple with the
shortest path rather than the longest. The LnN algorithm uses similar recursive logic to
first label each node with the number of leaf nodes below it. It then uses the oldest-
descendant labels to identify when a dated node has only undated nodes and leaf nodes
below it, and in those cases, works down from the root assigning a date to each undated
child by multiplying the date of its parent node by the ratio of the logarithm of the leaf
counts in the child clade and the parent clade. The BM algorithm also identifies dated
nodes with only undated and leaf nodes below, then works down the clade, randomly
choosing among the lineages and assigning the expected date for each bifurcation given
the numbers of lineages.

Quality assessment of date interpolation algorithms

We assessed the interpolation algorithms by removing input dates at random from a
clade, interpolating all missing dates, and measuring the difference between the
interpolated dates and the input dates that we had removed. We did this first for four
published dated trees, each with molecular data for all species: the Jetz et al. tree of
6,645 bird species [69]; the tree of 4,988 frog species by Portik et al. [70]; the mammals
tree of Upham et al. with 4,064 species [71]; and the tree of 609 cartilaginous fish by Stein
et al. [72].

For each tree, we first removed a fixed proportion of dates - for example, 10% - at random
from the internal nodes of the tree, though never from the root node. We ran the first
interpolation algorithm over the tree, and computed the absolute difference in million

years between the interpolated dates and the original dates. We then divided the differences by the root age of the tree, to normalise across trees of different ages - average absolute differences can be expected to be larger if all the dates in the tree are larger (older). We then ran the other four algorithms on the same partly-dated tree. That completed one replicate for each algorithm.

We then restored all input dates and randomly chose a different 10% of dates to remove, again finding the interpolation errors for each algorithm, and repeated this until we had 10 replicates for each algorithm. The errors were then averaged across the ten replicates. This gave us one datapoint for each algorithm: the average interpolation error when we removed 10% of dates from one fully-dated tree. We did this for all four trees, and averaged the errors. Then we removed 25% of dates, repeating the entire process, and so on; we tested removing 10%, 25%, 33%, 50%, 60%, 70%, 80%, 90%, 93%, 96%, 98%, 99%, and 100% of the internal dates (excluding the root age).

At the same time, we computed for each replicate the mean and median pendant edge length in the tree, relative to the mean and median in the original fully-dated tree. We considered it desirable for the distribution of pendant edge lengths to remain close to the original distribution when interpolating dates, i.e. for the relative edge lengths to stay close to 1.

Finally, we repeated this analysis on seven clades of our bifurcating complete tree, chosen because they had a high proportion of their nodes dated including the crown node of the clade. The clades were: Aves (10,931 spp., 73% dated); Squamata (11,221 spp., 34% dated); Mammalia (7964 spp., 33% dated); Actinopteri (37,408 spp., 28% dated); Polypodiales (10,709 spp., 27% dated); Amphibia (9,835 spp., 19% dated); and Fagales (2,274 spp., 20% dated). Interpolation errors were measured relative to the crown age of the clade. Since we had no fully-dated baseline tree, pendant edge lengths were normalised relative to those resulting from the EQS-LS algorithm when no dates had been removed.

Date consistency across the tree

The dates we used came from 334 trees (a full list of citations is given in S1 Text). The dates and topologies in these trees were not necessarily consistent with one another. When the dates were assigned to the synthetic supertree, some nodes had dates younger than a descendant or older than an ancestor, which would imply negative branch lengths. The BLADJ implementation in Phylocom of the equal splits algorithm deals with inconsistent dates by working down the tree and removing any date older than an ancestral date [34]. This approach has linear time complexity, but by working in one direction from the root it will bias the results toward younger dates, and is likely to remove more dates than necessary to produce a consistent tree. For example, consider a node with a date of 10 Mya that has two older children dated 11 and 12 Mya. The BLADJ algorithm will remove the dates on the children because they are lower in the tree, but consistency could also be achieved by removing only the parent date of 10 Mya, retaining two of the original three dates rather than one.

We developed an alternative algorithm that removes the minimum number of dates necessary to make a tree consistent. We first traversed once through the tree and produced for each node a list of any older descendants and younger ancestors. Then we chose the node with the largest number of inconsistencies, removed the date from that node, then removed the node from the lists for all other nodes. We then chose the node with the most remaining inconsistencies, and so on until there were no inconsistencies left.

Our method has a worst-case time complexity of $O(n \times m)$, where n is the number of nodes and m is the number of dated nodes: we visit each node once (n nodes), and at each node we scan the list of all dated descendants (m nodes). On our sparsely-dated trees we have $m \ll n$, so the realised time complexity is very close to linear in the number of nodes (**Fig 1**).

Phylogenetic diversity distributions

For each fully-dated tree, we summed all branch lengths to compute phylogenetic diversity. We traversed the tree from leaves to root; at each node we stored the sum of desendant branch lengths, then added the length of the branch to the parent node and passed this sum on to the parent. This recursion computes and stores the PD of the clade below every node in the tree in linear time.

We then computed three PD distributions: one exploring temporal uncertainty, one exploring topological uncertainty and one exploring both topological and temporal uncertainty simultaneously.

First, temporal uncertainty in the tree was explored by choosing among the available date sources at each node. Here, we used a single fixed tree topology; we chose the topology that give us the median PD estimate from the distribution exploring topological uncertainty. We created a list of all date source trees. We randomly shuffled the list, then treated this list as a priority ordering of sources: when we assigned dates to nodes in the tree, we used the date from the highest-priority source in the list. We then made the dates time-consistent, interpolated all dates and computed PD. Then we removed all the dates, re-shuffled the list of sources, and repeated the process 501 times, producing a distribution of 501 PD estimates.

Second, we investigated topological uncertainty by running the stochastic polytomy resolution step repeatedly to produce 501 different dated trees. For each tree topology generated, we computed a PD estimate using the median of the available date sources at each node. We then removed all dates and assigned a single date source at each node using random prioritisation of sources as previously described. We repeated this to give us three different dated trees with the same topology but varying dates. These steps produced two further distributions, one of 501 PD estimates exploring topological variation only, and one of 1503 PD estimates exploring both topological and temporal variation simultaneously.

Benchmarking and software

We compared the run times of our EQS-L interpolation implementation and the BLADJ implementation in Phylocom [34] on increasingly large, partially-dated subtrees of our complete tree. The performance of BLADJ was measured using the software "hyperfine" [73] in the zsh shell on macOS Sequoia 15.1.1, running on an Apple M4 Pro Macbook Pro. The Phylocom software was compiled for Apple Silicon from source on the same machine. For benchmarking, our code read the same input files and wrote the same outputs as Phylocom, and was run in the CPython 3.13.0 implementation of Python using IPython's "%timeit" magic command to measure run times. Random numbers were generated using the default random number generator in Numpy 2.1.3 [74]. For tree structures and tree visualisation we used the ETE4 library [75].

Data availability

The Python code for all the algorithms described here is available at https://github.com/jdduke24/dated-complete-tree. Distributions of fully-dated complete trees using both the EQS-LS and birth model interpolation algorithms are available on Dryad. All input dated trees are available on Github in the Open Tree data store https://github.com/OpenTreeOfLife/phylesystem-1 and via the OpenTree data curation website https://tree.opentreeoflife.org/curator.

Funding

This work was supported by the Natural Environment Research Council [NE/S007415/1], and NSF ABI No. 1759846.

References

1. Darwin C. On the origin of species by means of natural selection, or preservation of favoured races in the struggle for life: London : John Murray, 1859; 1859.

2. Haeckel E. Generelle Morphologie der Organismen: Allgemeine Grundzüge der organischen Formen-Wissenschaft, mechanisch begründet durch die von Charles Darwin reformierte Descendenz-Theorie. Band 1: Allgemeine Anatomie. Band 2: Allgemeine Entwicklungsgeschichte: de Gruyter; 1866.

3. Sayers EW, Beck J, Bolton EE, Brister JR, Chan J, Connor R, et al. Database resources of the National Center for Biotechnology Information in 2025. Nucleic Acids Research. 2024;53(D1):D20.

4. Lewin HA, Robinson GE, Kress WJ, Baker WJ, Coddington J, Crandall KA, et al. Earth BioGenome Project: Sequencing life for the future of life. Proceedings of the National Academy of Sciences. 2018;115(17):4325-33. doi: doi:10.1073/pnas.1720115115.

5. Zuntini AR, Carruthers T, Maurin O, Bailey PC, Leempoel K, Brewer GE, et al. Phylogenomics and the rise of the angiosperms. Nature. 2024. doi: 10.1038/s41586-024-07324-0. PubMed PMID: WOS:001253160400001.

6. McTavish EJ, Gerbracht JA, Holder MT, Iliff MJ, Lepage D, Rasmussen PC, et al. A complete and dynamic tree of birds. Proceedings of the National Academy of Sciences. 2025;122(18):e2409658122. doi: 10.1073/pnas.2409658122.

7. Zhu Q, Mai U, Pfeiffer W, Janssen S, Asnicar F, Sanders JG, et al. Phylogenomics of 10,575 genomes reveals evolutionary proximity between domains Bacteria and Archaea. Nature Communications. 2019;10(1):5477. doi: 10.1038/s41467-019-13443-4.

8. Rabosky DL, Chang J, Title PO, Cowman PF, Sallan L, Friedman M, et al. An inverse latitudinal gradient in speciation rate for marine fishes. Nature. 2018;559(7714):392-5. doi: 10.1038/s41586-018-0273-1.

9. Testo W, Sundue M. A 4000-species dataset provides new insight into the evolution of ferns. Molecular Phylogenetics and Evolution. 2016;105:200-11. doi: https://doi.org/10.1016/j.ympev.2016.09.003.

10. Kawahara AY, Storer C, Carvalho APS, Plotkin DM, Condamine FL, Braga MP, et al. A global phylogeny of butterflies reveals their evolutionary history, ancestral hosts and biogeographic origins. Nature Ecology & Evolution. 2023;7(6):903-13. doi: 10.1038/s41559-023-02041-9.

11. Felsenstein J. Phylogenies and the Comparative Method. The American Naturalist. 1985;125(1):1-15.

12.   Grafen A, Hamilton WD. The phylogenetic regression. Philosophical Transactions
of the Royal Society of London B, Biological Sciences. 1989;326(1233):119-57. doi:
doi:10.1098/rstb.1989.0106.

13.   Martins EP, Hansen TF. Phylogenies and the Comparative Method: A General
Approach to Incorporating Phylogenetic Information into the Analysis of Interspecific
Data. The American Naturalist. 1997;149(4):646-67. doi: 10.1086/286013.

14.   Pagel M. Inferring the historical patterns of biological evolution. Nature.
1999;401(6756):877-84. doi: 10.1038/44766.

15.   Blomberg SP, Garland JR. T, Ives AR. Testing for phylogenetic signal in comparative
data: behavioural traits are more liable. Evolution. 2003;57(4):717-45. doi:
https://doi.org/10.1111/j.0014-3820.2003.tb00285.x.

16.   Swenson NG. Phylogenetic imputation of plant functional trait databases.
Ecography. 2014;37(2):105-10. doi: https://doi.org/10.1111/j.1600-0587.2013.00528.x.

17.   Penone C, Davidson AD, Shoemaker KT, Di Marco M, Rondinini C, Brooks TM, et al.
Imputation of missing data in life-history trait datasets: which approach performs the
best?   Methods   in   Ecology   and   Evolution.   2014;5(9):961-70.   doi:
https://doi.org/10.1111/2041-210X.12232.

18.   Moura MR, Ceron K, Guedes JJM, Chen-Zhao R, Sica YV, Hart J, et al. A phylogeny-
informed characterisation of global tetrapod traits addresses data gaps and biases. PLOS
Biology. 2024;22(7):e3002658. doi: 10.1371/journal.pbio.3002658.

19.   Faith DP. Conservation evaluation and phylogenetic diversity. Biological
Conservation. 1992;61(1):1-10. doi: https://doi.org/10.1016/0006-3207(92)91201-3.

20.   Isaac NJB, Turvey ST, Collen B, Waterman C, Baillie JEM. Mammals on the EDGE:
Conservation Priorities Based on Threat and Phylogeny. PLOS ONE. 2007;2(3):e296. doi:
10.1371/journal.pone.0000296.

21.   Hinchliff CE, Smith SA, Allman JF, Burleigh JG, Chaudhary R, Coghill LM, et al.
Synthesis of phylogeny and taxonomy into a comprehensive tree of life. Proceedings of
the National Academy of Sciences of the United States of America. 2015;112(41):12764-
9. doi: 10.1073/pnas.1423041112. PubMed PMID: WOS:000363130900055.

22.   McTavish EJ, Drew BT, Redelings B, Cranston KA. How and Why to Build a Unified
Tree   of   Life.   BioEssays.   2017;39(11):1700114.   doi:
https://doi.org/10.1002/bies.201700114.

23.   McTavish EJ, Hinchliff CE, Allman JF, Brown JW, Cranston KA, Holder MT, et al.
Phylesystem: a git-based data store for community-curated phylogenetic estimates.
Bioinformatics. 2015;31(17):2794-800. doi: 10.1093/bioinformatics/btv276.

24.   Redelings BD, Holder MT. A supertree pipeline for summarizing phylogenetic and
taxonomic information for millions of species. PeerJ. 2017;5:e3058. doi:
10.7717/peerj.3058.

25.   Rees JA, Cranston K. Automated assembly of a reference taxonomy for
phylogenetic data synthesis. Biodiversity Data Journal. 2017;5. doi:
10.3897/BDJ.5.e12581. PubMed PMID: WOS:000449744100001.

26.   Redelings BD, Holder MT. Taxonomic Supertree Construction with Incertae sedis
Taxa. In: Warnow T, editor. Bioinformatics and Phylogenetics: Seminal Contributions of
Bernard Moret. Cham: Springer International Publishing; 2019. p. 151-73.

27.   OpenTree. Open Tree of Life version 16.1 2025 [2026-01-12]. Available from:
https://tree.opentreeoflife.org/about/synthesis-release/v16.1.

28.   Kuhn TS, Mooers AØ, Thomas GH. A simple polytomy resolver for dated
phylogenies. Methods in Ecology and Evolution. 2011;2(5):427-36. doi:
https://doi.org/10.1111/j.2041-210X.2011.00103.x.

29.   Swenson NG. Phylogenetic resolution and quantifying the phylogenetic diversity
and dispersion of communities. PloS one. 2009;4(2):e4390.

30.   Thomas GH, Hartmann K, Jetz W, Joy JB, Mimoto A, Mooers AO. PASTIS: an R
package to facilitate phylogenetic assembly with soft taxonomic inferences. Methods in
Ecology and Evolution. 2013;4(11):1011-7. doi: https://doi.org/10.1111/2041-
210X.12117.

31.   Pearse WD, Chase MW, Crawley MJ, Dolphin K, Fay MF, Joseph JA, et al. Beyond
the EDGE with EDAM: Prioritising British Plant Species According to Evolutionary
Distinctiveness, and Accuracy and Magnitude of Decline. PLOS ONE.
2015;10(5):e0126524. doi: 10.1371/journal.pone.0126524.

32.   Sánchez Reyes LL, McTavish EJ, O'Meara B. DateLife: Leveraging Databases and
Analytical Tools to Reveal the Dated Tree of Life. Systematic Biology. 2024;73(2):470-85.
doi: 10.1093/sysbio/syae015.

33.   McTavish EJ, Gerbracht JA, Holder MT, Iliff MJ, Lepage D, Rasmussen P, et al. A
complete and dynamic tree of birds. bioRxiv. 2024:2024.05.20.595017. doi:
10.1101/2024.05.20.595017.

34.   Webb CO, Ackerly DD, Kembel SW. Phylocom: software for the analysis of
phylogenetic community structure and trait evolution. Bioinformatics. 2008;24(18):2098-
100. doi: 10.1093/bioinformatics/btn358.

35.   Purvis A. A composite estimate of primate phylogeny. Philosophical Transactions
of the Royal Society of London Series B: Biological Sciences. 1995;348(1326):405-21.
doi: doi:10.1098/rstb.1995.0078.

36.   Britton T, Anderson CL, Jacquet D, Lundqvist S, Bremer K. Estimating Divergence
Times in Large Phylogenetic Trees. Systematic Biology. 2007;56(5):741-52. doi:
10.1080/10635150701613783.

37.   Tamura K, Battistuzzi FU, Billing-Ross P, Murillo O, Filipski A, Kumar S. Estimating
divergence times in large molecular phylogenies. Proceedings of the National Academy
of Sciences. 2012;109(47):19333-8. doi: doi:10.1073/pnas.1213199109.

38.   Mast AR, Olde PM, Makinson RO, Jones E, Kubes A, Miller ET, et al. Paraphyly
changes understanding of timing and tempo of diversification in subtribe Hakeinae
(Proteaceae), a giant Australian plant radiation. American Journal of Botany.
2015;102(10):1634-46.

39.   Ramos-Gutiérrez I, Lima H, Vilela B, Molina-Venegas R. A generalized framework
to expand incomplete phylogenies using non-molecular phylogenetic information.
Global   Ecology   and   Biogeography.   2023;32(10):1707-16.   doi:
https://doi.org/10.1111/geb.13733.

40.   Pearse WD, Purvis A. phyloGenerator: an automated phylogeny generation tool for
ecologists.   Methods   in   Ecology   and   Evolution.   2013;4(7):692-8.   doi:
https://doi.org/10.1111/2041-210X.12055.

41.   OpenTree. Open Tree of Life reference taxonomy version 3.6 2023 [2025-06-02].
Available from: https://tree.opentreeoflife.org/about/taxonomy-version/ott3.6.

42.   Wong Y, Rosindell J. Dynamic visualisation of million-tip trees: The OneZoom
project.   Methods   in   Ecology   and   Evolution.   2022;13(2):303-13.   doi:
https://doi.org/10.1111/2041-210X.13766.

43.   Hedges SB, Dudley J, Kumar S. TimeTree: a public knowledge-base of divergence
times   among   organisms.   Bioinformatics.   2006;22(23):2971-2.   doi:
10.1093/bioinformatics/btl505.

44.   Kumar S, Suleski M, Craig JM, Kasprowicz AE, Sanderford M, Li M, et al. TimeTree
5: An Expanded Resource for Species Divergence Times. Molecular Biology and
Evolution. 2022;39(8). doi: 10.1093/molbev/msac174.

45.   Sayers EW, Cavanaugh M, Frisse L, Pruitt KD, Schneider VA, Underwood Beverly A,
et al. GenBank 2025 update. Nucleic Acids Research. 2024;53(D1):D56-D61. doi:
10.1093/nar/gkae1114.

46.   Blaxter M, Lewin HA, DiPalma F, Challis R, da Silva M, Durbin R, et al. The Earth
BioGenome Project Phase II: illuminating the eukaryotic tree of life. Frontiers in Science.
2025;3:1514835.

47.   Rabosky DL. No substitute for real data: A cautionary note on the use of
phylogenies from birth–death polytomy resolvers for downstream comparative analyses.
Evolution. 2015;69(12):3207-16. doi: https://doi.org/10.1111/evo.12817.

48.   Weedop KB, Mooers AØ, Tucker CM, Pearse WD. The effect of phylogenetic
uncertainty and imputation on EDGE Scores. Animal Conservation. 2019;22(6):527-36.
doi: https://doi.org/10.1111/acv.12495.

49.   Molina-Venegas R. How to get the most out of phylogenetic imputation without
abusing it. Methods in Ecology and Evolution. 2024;15(3):456-63. doi:
https://doi.org/10.1111/2041-210X.14198.

50.   Gumbs R, Scott O, Bates R, Böhm M, Forest F, Gray CL, et al. Global conservation
status of the jawed vertebrate Tree of Life. Nature Communications. 2024;15(1):1101.
doi: 10.1038/s41467-024-45119-z.

51.   Martyn I, Kuhn TS, Mooers AO, Moulton V, Spillner A. Computing evolutionary
distinctiveness indices in large scale analysis. Algorithms for Molecular Biology.
2012;7(1):6. doi: 10.1186/1748-7188-7-6.

52.   Díaz S, Malhi Y. Biodiversity: Concepts, Patterns, Trends, and Perspectives. Annual
Review of Environment and Resources. 2022;47(Volume 47, 2022):31-63. doi:
https://doi.org/10.1146/annurev-environ-120120-054300.

53.   Rosindell J, Guo J, McTavish EJ. Phylogenetic Diversity Across the Complete Tree
of Life. 2024.

54.   Jetz W, Thomas Gavin H, Joy Jeffrey B, Redding David W, Hartmann K, Mooers
Arne O. Global Distribution and Conservation of Evolutionary Distinctness in Birds.
Current Biology. 2014;24(9):919-30. doi: 10.1016/j.cub.2014.03.011.

55.   Konstantinidis KT, Ramette A, Tiedje JM. The bacterial species definition in the
genomic era. Philosophical Transactions of the Royal Society B: Biological Sciences.
2006;361(1475):1929-40. doi: doi:10.1098/rstb.2006.1920.

56.   Rosselló-Móra R, Amann R. Past and future species definitions for Bacteria and
Archaea. Systematic and Applied Microbiology. 2015;38(4):209-16. doi:
https://doi.org/10.1016/j.syapm.2015.02.001.

57.   Louca S, Mazel F, Doebeli M, Parfrey LW. A census-based estimate of Earth's
bacterial and archaeal diversity. PLOS Biology. 2019;17(2):e3000106. doi:
10.1371/journal.pbio.3000106.

58.   Mora C, Tittensor DP, Adl S, Simpson AGB, Worm B. How Many Species Are There
on Earth and in the Ocean? PLOS Biology. 2011;9(8):e1001127. doi:
10.1371/journal.pbio.1001127.

59.   Scheffers BR, Joppa LN, Pimm SL, Laurance WF. What we know and don't know
about Earth's missing biodiversity. Trends in Ecology & Evolution. 2012;27(9):501-10. doi:
https://doi.org/10.1016/j.tree.2012.05.008.

60.   Stork NE. How Many Species of Insects and Other Terrestrial Arthropods Are There
on Earth? Annual Review of Entomology. 2018;63(Volume 63, 2018):31-45. doi:
https://doi.org/10.1146/annurev-ento-020117-043348.

61.   Niskanen T, Lücking R, Dahlberg A, Gaya E, Suz LM, Mikryukov V, et al. Pushing the
Frontiers of Biodiversity Research: Unveiling the Global Diversity, Distribution, and
Conservation of Fungi. Annual Review of Environment and Resources. 2023;48(Volume
48, 2023):149-76. doi: https://doi.org/10.1146/annurev-environ-112621-090937.

62.   Wilkinson MD, Dumontier M, Aalbersberg IJ, Appleton G, Axton M, Baak A, et al.
The FAIR Guiding Principles for scientific data management and stewardship. Scientific
Data. 2016;3(1):160018. doi: 10.1038/sdata.2016.18.

63.   OpenTree. Open Tree of Life Study Curator 2025. Available from:
https://tree.opentreeoflife.org/curator.

64.   OpenTree. Open Tree of Life Taxonomy version 3.7.3 2025. Available from:
https://tree.opentreeoflife.org/about/taxonomy-version/ott3.7.3.

65.   Clements JF, Rasmussen PC, Schulenberg TS, Iliff MJ, Fredericks TA, Gerbracht JA,
et al. The eBird/Clements checklist of Birds of the World: v2023 2023 [cited 2025 2025-
11-07]. Available from: https://www.birds.cornell.edu/clementschecklist/download/.

66.   Turtle Taxonomy Working Group. Turtles of the World: Annotated Checklist and
Atlas of Taxonomy, Synonymy, Distribution, and Conservation Status. 10th ed: Chelonian
Research Foundation and Turtle Conservancy; 2025.

67.   Hedges SB, Chen H, Kumar S, Wang DYC, Thompson AS, Watanabe H. A genomic
timescale for the origin of eukaryotes. BMC Evolutionary Biology. 2001;1(1):4. doi:
10.1186/1471-2148-1-4.

68.   Sleep NH, Zahnle KJ, Kasting JF, Morowitz HJ. Annihilation of ecosystems by large
asteroid impacts on the early Earth. Nature. 1989;342(6246):139-42. doi:
10.1038/342139a0.

69.   Jetz W, Thomas GH, Joy JB, Hartmann K, Mooers AO. The global diversity of birds
in space and time. Nature. 2012;491(7424):444-8. doi: 10.1038/nature11631.

70.   Portik DM, Streicher JW, Wiens JJ. Frog phylogeny: A time-calibrated, species-level
tree based on hundreds of loci and 5,242 species. Molecular Phylogenetics and
Evolution. 2023;188:107907. doi: https://doi.org/10.1016/j.ympev.2023.107907.

71.   Upham NS, Esselstyn JA, Jetz W. Inferring the mammal tree: Species-level sets of
phylogenies for questions in ecology, evolution, and conservation. PLOS Biology.
2019;17(12):e3000494. doi: 10.1371/journal.pbio.3000494.

72. Stein RW, Mull CG, Kuhn TS, Aschliman NC, Davidson LNK, Joy JB, et al. Global
priorities for conserving the evolutionary history of sharks, rays and chimaeras. Nature
Ecology & Evolution. 2018;2(2):288-98. doi: 10.1038/s41559-017-0448-4.

73. Peter D. hyperfine. 1.16.1 ed2023.

74. Harris CR, Millman KJ, Van Der Walt SJ, Gommers R, Virtanen P, Cournapeau D, et
al. Array programming with NumPy. nature. 2020;585(7825):357-62.

75. Huerta-Cepas J, Serra F, Bork P. ETE 3: Reconstruction, Analysis, and Visualization
of Phylogenomic Data. Molecular Biology and Evolution. 2016;33(6):1635-8. doi:
10.1093/molbev/msw046.

76. Pybus OG, Harvey PH. Testing Macro-Evolutionary Models Using Incomplete
Molecular Phylogenies. Proceedings: Biological Sciences. 2000;267(1459):2267-72.

77. Tonini JFR, Beard KH, Ferreira RB, Jetz W, Pyron RA. Fully-sampled phylogenies of
squamates reveal evolutionary patterns in threat status. Biological Conservation.
2016;204:23-31. doi: https://doi.org/10.1016/j.biocon.2016.03.039.

78. Forest F, Moat J, Baloch E, Brummitt NA, Bachman SP, Ickert-Bond S, et al.
Gymnosperms on the EDGE. Scientific Reports. 2018;8(1):6053. doi: 10.1038/s41598-
018-24365-4.

Supporting information

1001 **S1 Fig: Full breakdown of date interpolation quality assessment on four fully-dated trees.**
1002 Columns from left to right: results for four fully-dated trees with molecular data for all included
1003 species: the mammals tree of Upham et al. [71]; the tree of 609 cartilaginous fish by Stein et al. [72];
1004 the tree of 4,988 frog species by Portik et al. [70]; and the Jetz et al. tree of 6,645 bird species [69]. We
1005 removed a proportion of dates at random from the tree (but never the root age), then interpolated
1006 dates across all undated internal nodes and compared trees with interpolated dates to the original
1007 trees. Top row: the average error in date reproduction tests (absolute difference in million years).
1008 Second row: the average error in date reproduction as a proportion of the root age of the tree. This
1009 normalises across trees of different ages; for example, sharks are older than mammals, so the
1010 absolute differences are expected to be larger. Third row: the mean pendant edge lengths for the same
1011 trees, divided by the mean for the fully-dated tree. Fourth row: Median pendant edge lengths, divided
1012 by the median for the fully-dated tree. Fifth row: Pybus and Harvey's gamma statistic characterising
1013 the distribution of branch lengths [76]. Positive values indicate younger dates and shorter pendant
1014 edges than expected from a birth model, and vice versa. All trees converge on 0 when 100% of dates
1015 are interpolated using birth model interpolation (red lines), as expected.

S2   **Fig:**   **As**   **in**

1018

S1 Fig but for seven subtrees of a partially-dated complete tree. Note the different x and y scales in each column.

1019
1020

1021

1022

1023 **S3 Fig: Distribution of the number of date sources at nodes with at least one source.** There were
1024 76,935 nodes with at least one date source. Around 82% (62,780) of these dated nodes had only one
1025 or two sources; 99% of nodes have seven or fewer sources. There are 25 nodes with 20 or more
1026 sources, all in the birds clade.

1027

1028

S4 Fig: Interpolation can result in negative branch lengths. The numbers shown next to the internal
nodes in this cladogram are dates in Mya; the node labelled A is to be dated. Assigning a date to A
using, for example, 12 x ln(4) / ln(5) = 10.3 Mya, as computed by the LnN algorithm explained in the
main text, would result in a negative branch length between A and the node dated 11. The EQS
algorithms find the oldest dated node leafwards of A, and place A midway in time between the node
dated 12 and the oldest dated node below it (11), guaranteeing no negative branch lengths.
1035

S1 Table: Comparisons of our median phylogenetic diversity estimates with estimates from previous studies.

1038

Clade	Study	Species richness	PD (Myr)	PD:spp ratio
All Life	This study	2,294,776	39,782,514	17.3
	Guo et al. (2025) [50]	2,235,076	31,000,000	13.9
	Diaz & Malhi (2022) [52]	2,850,746	345,367,000	121.1
Eukaryotes	This study	2,215,634	28,914,060	13.1
(Eukaryota)	Guo et al. (2025)	2,174,404	26,004,213	12.0
	Diaz & Malhi (2022)	2,110,866	124,417,000	58.9
Mammals	This study	7,965	56,828	7.1
(Mammalia)	Gumbs et al. (2024)	6,253	37,068	5.9
	Martyn et al. (2012) [51]	5,139	49,131	9.6
	Isaac et al. (2007) [20]	4,433	42,628	9.6
	Guo et al. (2025)	5,046	50,075	9.9
	Diaz & Malhi (2022)	5,045	66,000	13.1
Birds	This study	10,931	96,272	8.8
(Aves)	Gumbs et al. (2024)	10,988	89,747	8.2
	Jetz et al. (2014) [54]	9,993	77,149	7.7
	Guo et al. (2025)	10,002	74,177	7.4
	Diaz & Malhi (2022)	9,993	81,000	8.1
Ray-finned fish	This study	37,408	410,443	11.0
(Actinopterygii)	Gumbs et al. (2024)	32,760	329,935	10.1
	Guo et al. (2025)	34,432	348,288	10.1
	Diaz & Malhi (2022)	35,810	1,512,000	42.2
Amphibians	This study	9,835	126,956	12.9
(Amphibia)	Gumbs et al. (2024)	8,024	142,670	17.8
	Isaac et al. (2012)	4,331	57,980	13.4
	Guo et al. (2025)	8,688	95,735	11.0
Lepidosaurs	This study	11,221	176,652	15.7
(Lepidosauria)	Gumbs et al. (2024)	10,735	134,253	12.5
	Tonini et al. (2016) [77]	9,755	122,478	12.6
Gymnosperms	This study	2,014	38,360	19.0
(Acrogymnospermae)	Forest et al. (2018) [78]	1,090	12,671	11.6
Jawed vertebrates	This study	79,430	895,692	11.3
(Gnathostomata)	Gumbs et al. (2024)	70,426	782,102	11.1
Cartilaginous fish	This study	1,653	52,119	31.5
(Chondrichthyes)	Gumbs et al. (2024)	1,290	41,297	32.0
Crocodilians	This study	34	586	17.2
(Crocodilia)	Gumbs et al. (2024)	25	613	24.5
Turtles	This study	326	6,245	19.2
(Testudines)	Gumbs et al. (2024)	351	6,520	18.6

1039

S1 Text: References for studies in the Open Tree phylesystem that contain dated phylogenetic trees used as date sources in this work. Some studies contributed more than one tree. Listed in alphabetical order of first author surname.

1. Abalde, S., Tenorio, M. J., Afonso, C. M. L., & Zardoya, R. (2017). Mitogenomic phylogeny of cone snails endemic to Senegal. Molecular Phylogenetics and Evolution, 112, 79–87. https://doi.org/10.1016/j.ympev.2017.04.020

2. Abalde, S., Tenorio, M. J., Afonso, C. M. L., Uribe, J. E., Echeverry, A. M., & Zardoya, R. (2017). Phylogenetic relationships of cone snails endemic to Cabo Verde based on mitochondrial genomes. BMC Evolutionary Biology, 17(1). https://doi.org/10.1186/s12862-017-1069-x

3. Agarwal, I., Bauer, A. M., Gamble, T., Giri, V. B., Jablonski, D., Khandekar, A., Mohapatra, P. P., Masroor, R., Mishra, A., & Ramakrishnan, U. (2022). The evolutionary history of an accidental model organism, the leopard gecko Eublepharis macularius (Squamata: Eublepharidae). Molecular Phylogenetics and Evolution, 168, 107414. https://doi.org/10.1016/j.ympev.2022.107414

4. Aggerbeck, Marie, Jon Fjeldså, Les Christidis, Pierre-Henri Fabre, Knud Andreas Jønsson. 2014. Resolving deep lineage divergences in core corvoid passerine birds supports a proto-Papuan island origin. Molecular Phylogenetics and Evolution 70: 272-285.

5. Alencar, Laura R.V., Tiago B. Quental, Felipe G. Grazziotin, Michael L. Alfaro, Marcio Martins, Mericien Venzon, Hussam Zaher. 2016. Diversification in vipers: Phylogenetic relationships, time of divergence and shifts in speciation rates. Molecular Phylogenetics and Evolution 105: 50-62

6. Almeida, E. A. B., Bossert, S., Danforth, B. N., Porto, D. S., Freitas, F. V., Davis, C. C., Murray, E. A., Blaimer, B. B., Spasojevic, T., Ströher, P. R., Orr, M. C., Packer, L., Brady, S. G., Kuhlmann, M., Branstetter, M. G., & Pie, M. R. (2023). The evolutionary history of bees in time and space. Current Biology, 33(16), 3409-3422.e6. https://doi.org/10.1016/j.cub.2023.07.005

7. Alström, P., Cibois, A., Irestedt, M., Zuccon, D., Gelang, M., Fjeldså, J., Andersen, M. J., Moyle, R. G., Pasquet, E., & Olsson, U. (2018). Comprehensive molecular phylogeny of the grassbirds and allies (Locustellidae) reveals extensive non-monophyly of traditional genera, and a proposal for a new classification. Molecular Phylogenetics and Evolution, 127, 367–375. https://doi.org/10.1016/j.ympev.2018.03.029

8. Alström, P., Höhna, S., Gelang, M., Ericson, P. G., & Olsson, U. (2011). Non-monophyly and intricate morphological evolution within the avian family Cettiidae revealed by multilocus analysis of a taxonomically densely sampled dataset. BMC Evolutionary Biology, 11(1), 352. https://doi.org/10.1186/1471-2148-11-352

9. Alström, P., Mohammadi, Z., Enbody, E., Irestedt, M., Engelbrecht, D., Crochet, P.-A., Guillaumet, A., Rancilhac, L., Tieleman, B. I., Olsson, U., Donald, P. F., & Stervander, M. (2023). Systematics of the avian family Alaudidae using multilocus and genomic data. Avian Research, 100095. https://doi.org/10.1016/j.avrs.2023.100095

10. Alström, P., Rasmussen, P. C., Xia, C., Gelang, M., Liu, Y., Chen, G., Zhao, M., Hao, Y., Zhao, C., Zhao, J., Yao, C., Eaton, J. A., Hutchinson, R., Lei, F., & Olsson, U. (2018). Taxonomy of the White-browed Shortwing (Brachypteryx montana) complex on mainland Asia and Taiwan: an integrative approach supports recognition of three instead of one species. Avian Research, 9(1). https://doi.org/10.1186/s40657-018-0125-6

11. Alström, P., Rasmussen, P. C., Zhao, C., Xu, J., Dalvi, S., Cai, T., Guan, Y., Zhang, R., Kalyakin, M. V.,
Lei, F., & Olsson, U. (2016). Integrative taxonomy of the Plain-backed Thrush (Zoothera mollissima) complex
(Aves, Turdidae) reveals cryptic species, including a new species. Avian Research, 7(1).
https://doi.org/10.1186/s40657-016-0037-2

12. Amy E. Zanne, David C. Tank, William K. Cornwell, Jonathan M. Eastman, Stephen A. Smith,
Richard G. FitzJohn, Daniel J. McGlinn, Brian C. O'Meara, Angela T. Moles, Peter B. Reich, Dana L. Royer,
Douglas E. Soltis, Peter F. Stevens, Mark Westoby, Ian J. Wright, Lonnie Aarssen, Robert I. Bertin, Andre
Calaminus, Rafaël Govaerts, Frank Hemmings, Michelle R. Leishman, Jacek Oleksyn, Pamela S. Soltis,
Nathan G. Swenson, Laura Warman, Jeremy M. Beaulieu, 2013, 'Three keys to the radiation of angiosperms
into freezing environments', Nature, vol. 506, no. 7486, pp. 89-92

13. Andersen, Michael J., Jenna M. McCullough, William M. Mauck, Brian Tilston Smith, Robert G.
Moyle. 2017. A phylogeny of kingfishers reveals an Indomalayan origin and elevated rates of diversification
on oceanic islands. Journal of Biogeography

14. Anderson Ferreira Pinto Machado, Nina Rønsted, Sam Bruun-Lund, Rodrigo Augusto Santinelo
Pereira, Luciano Paganucci de Queiroz, 2018, 'Atlantic forests to the all Americas: Biogeographical history
and divergence times of Neotropical Ficus (Moraceae)', Molecular Phylogenetics and Evolution, vol. 122,
pp. 46-58

15. Andrew L. Hipp, Paul S. Manos, Antonio González-Rodríguez, Marlene Hahn, Matthew Kaproth,
John D. McVay, Susana Valencia Avalos, Jeannine Cavender-Bares, 2017, 'Sympatric parallel diversification
of major oak clades in the Americas and the origins of Mexican species diversity', New Phytologist, vol. 217,
no. 1, pp. 439-452

16. Andujar, C., Serrano J., & Gomez-zurita J. 2012. Winding up the molecular clock in the genus
Carabus (Coleoptera: Carabidae): assessment of methodological decisions on rate and node age
estimation. BMC Evolutionary Biology 12: 40.

17. Antonelli, A. 2009. Have giant lobelias evolved several times independently? Life form shifts and
historical biogeography of the cosmopolitan and highly diverse subfamily Loeblioideae (Campanulaceae).
BMC Biology 7: 82 - see http://www.biomedcentral.com/1741-7007/7/82

18. Arakaki M., Christin P., Nyffeler R., Lendel A., Eggli U., Ogburn R., Spriggs E.,

19. Moore M.J., & Edwards E.J. 2011. Contemporaneous and recent radiations of the world?s

20. major succulent plant lineages. PNAS, 108(20): 8379-8384.

21. Arbabi, Tayebeh, Javier Gonzalez, Michael Wink. 2014. A re-evaluation of phylogenetic
relationships within reed warblers (Aves: Acrocephalidae) based on eight molecular loci and ISSR profiles.
Molecular Phylogenetics and Evolution 78: 304-313.

22. Barker, F. K., K. J. Burns, J. Klicka, S. M. Lanyon, I. J. Lovette. 2013. Going to extremes: contrasting
rates of diversification in a recent radiation of New World passerine birds. Systematic Biology 62 (2): 298-
320.

23. Barker, F. Keith, Kevin J. Burns, John Klicka, Scott M. Lanyon, Irby J. Lovette. 2015. New insights into
New World biogeography: An integrated view from the phylogeny of blackbirds, cardinals, sparrows,
tanagers, warblers, and allies. The Auk 132 (2): 333-348.

24.   Batista, R., Olsson, U., Andermann, T., Aleixo, A., Ribas, C. C., & Antonelli, A. (2020).
Phylogenomics and biogeography of the world's thrushes (Aves,   Turdus   ): new evidence for a
more parsimonious evolutionary history. Proceedings of the Royal Society B: Biological Sciences,
287(1919), 20192400. https://doi.org/10.1098/rspb.2019.2400

25.   Beaulieu, Jeremy M., Brian C. O'Meara, Peter Crane, Michael J. Donoghue. 2015. Heterogeneous
rates of molecular evolution and diversification could explain the Triassic age estimate for Angiosperms.
Systematic Biology 64 (5): 869-878

26.   Bechteler, J., Peñaloza-Bojacá, G., Bell, D., Gordon Burleigh, J., McDaniel, S. F., Christine Davis, E.,
Sessa, E. B., Bippus, A., Christine Cargill, D., Chantanoarrapint, S., Draper, I., Endara, L., Forrest, L. L.,
Garilleti, R., Graham, S. W., Huttunen, S., Lazo, J. J., Lara, F., Larraín, J., ... Villarreal A., J. C. (2023).
Comprehensive phylogenomic time tree of bryophytes reveals deep relationships and uncovers gene
incongruences in the last 500 million years of diversification. American Journal of Botany, 110(11). Portico.
https://doi.org/10.1002/ajb2.16249

27.   Bendiksby, M., Schumacher, T., Gussarova, G., Nais, J., Mat-Salleh, K., Sofiyanti, N., ... & Barkman,
T. (2010). Elucidating the evolutionary history of the Southeast Asian, holoparasitic, giant-flowered
Rafflesiaceae: Pliocene vicariance, morphological convergence and character displacement. Molecular
Phylogenetics and Evolution, 57(2), 620-633.

28.   Bergh N.G., & Linder H.P. 2009. Cape diversification and repeated out-of-southern-Africa dispersal
in paper daisies. Molecular Phylogenetics and Evolution, 51(Special issue: origins and evolution of a
biodiversity hotspot, the biodiversity of the African Cape Floristic Region): 5 - 18.

29.   Bhak, Y., Jeon, Y., Jeon, S., Chung, O., Jho, S., Jun, J., Kim, H.-M., Cho, Y., Yoon, C., Lee, S., Kang, J.-
H., Lim, J.-D., An, J., Cho, Y. S., Ryu, D.-Y., & Bhak, J. (2017). Myotis rufoniger genome sequence and
analyses: M. rufoniger's genomic feature and the decreasing effective population size of Myotis bats. PLOS
ONE, 12(7), e0180418. https://doi.org/10.1371/journal.pone.0180418

30.   Bininda-Emonds, Olaf R. P., Marcel Cardillo, Kate E. Jones, Ross D. E. MacPhee, Robin M. D. Beck,
Richard Grenyer, Samantha A. Price, Rutger A. Vos, John L. Gittleman, Andy Purvis. 2007. The delayed rise
of present-day mammals. Nature 446 (7135): 507-512

31.   Blanchard, Benjamin D., Corrie S. Moreau. 2016. Defensive traits exhibit an evolutionary trade-off
and drive diversification in ants. Evolution

32.   Bodner M.R., & Maddison W. 2012. The biogeography and age of salticid spider radiations
(Araneae: Salticidae). Molecular phylogenetics and Evolution, .

33.   Boore J., Longhorn S., & Masta S. 2008. Arachnid relationships based on mitochondrial genomes:
asymmetric nucleotide and amino acid bias affects phylogenetic analyses. Molecular Phylogenetics and
Evolution, null.

34.   Brady, S. G., Sipes, S., Pearson, A., & Danforth, B. N. (2006). Recent and simultaneous origins of
eusociality in halictid bees. Proceedings of the Royal Society B: Biological Sciences, 273(1594), 1643–1649.
https://doi.org/10.1098/rspb.2006.3496

35.   Breinholt, Jessse, Marcos Pérez-Losada, Keith A. Crandall. 2009. The timing of the diversification
of the freshwater crayfishes. In: Decapod Crustacean Phylogenetics.Edited by Joel W. Martin , Keith A.
Crandall, Darryl L. Felder. CRC Press. pp. 343-355

36.   Bremer, B., & Eriksson, O. 2009. Time tree of Rubiaceae: Phylogeny and dating the family,
subfamilies and tribes. Internat. J. Plant Sci. 170: 766-793.

37.   Brian Tilston Smith, Robert W. Bryson, William M. Mauck, Jaime Chaves, Mark B. Robbins,
Alexandre Aleixo, John Klicka, 2018, 'Species delimitation and biogeography of the gnatcatchers and
gnatwrens (Aves: Polioptilidae)', Molecular Phylogenetics and Evolution, vol. 126, pp. 45-57

38.   Bridge, E. S., Jones, A. W., & Baker, A. J. (2005). A phylogenetic framework for the terns (Sternini)
inferred from mtDNA sequences: implications for taxonomy and plumage evolution. Molecular
Phylogenetics and Evolution, 35(2), 459–469. doi:10.1016/j.ympev.2004.12.010

39.   Brock, K. M., McTavish, E. J., & Edwards, D. L. (2020). Color Polymorphism is a Driver of
Diversification in the Lizard Family Lacertidae. https://doi.org/10.1101/2020.08.27.270207

40.   Brown, Joseph W., Joshua S. Rest, Jaime García-Moreno, Michael D. Sorenson, and David P.
Mindell. 2008. Strong mitochondrial DNA support for a Cretaceous origin of modern avian lineages. BMC
Biology 6: 6

41.   Brown, Joseph W., Robert B. Payne, David P. Mindell. 2007. Comment. Nuclear DNA does not
reconcile 'rocks' and 'clocks' in Neoaves: a comment on Ericson et al. Biology Letters 3 (3): 257-259.

42.   Bruno Frédérich, Laurie Sorenson, Francesco Santini, Graham J. Slater, Michael E. Alfaro, 2013,
'Iterative Ecological Radiation and Convergence during the Evolutionary History of Damselfishes
(Pomacentridae)', The American Naturalist, vol. 181, no. 1, pp. 94-113

43.   Buainain, N., Ferreira, M., Avendaño, J. E., Cadena, C. D., Faircloth, B. C., Brumfield, R. T., Cracraft,
J., & Ribas, C. C. (2022). Biogeography of a neotropical songbird radiation reveals similar diversification
dynamics between montane and lowland clades. Journal of Biogeography. Portico.
https://doi.org/10.1111/jbi.14379

44.   Bucher, M., Condamine, F. L., Luo, Y., Wang, M., & Bourgoin, T. (2023). Phylogeny and diversification
of planthoppers (Hemiptera: Fulgoromorpha) based on a comprehensive molecular dataset and large
taxon   sampling.   Molecular   Phylogenetics   and   Evolution,   186,   107862.
https://doi.org/10.1016/j.ympev.2023.107862

45.   Burns, Kevin J., Allison J. Shultz, Pascal O. Title, Nicholas A. Mason, F. Keith Barker, John Klicka,
Scott M. Lanyon, Irby J. Lovette. 2014. Phylogenetics and diversification of tanagers (Passeriformes:
Thraupidae), the largest radiation of Neotropical songbirds. Molecular Phylogenetics and Evolution 75: 41-
77.

46.   Cai, L., Cardoso, D., Tressel, L. G., Lee, C., Shrestha, B., Choi, I.-S., de Lima, H. C., de Queiroz, L.
P., Ruhlman, T. A., Jansen, R. K., & Wojciechowski, M. F. (2024). Well-resolved phylogeny reveals multiple
origins   of   keel   flowers   during   the   diversification   of   papilionoid   legumes.
https://doi.org/10.1101/2024.10.07.617045

47.   Cai, T., Wu, G., Sun, L., Zhang, Y., Peng, Z., Guo, Y., Liu, X., Pan, T., Chang, J., Sun, Z., & Zhang, B.
(2021). Biogeography and diversification of Old World buntings (Aves: Emberizidae): radiation in open
habitats. Journal of Avian Biology, 52(6). Portico. https://doi.org/10.1111/jav.02672

48.   Cardinal-McTeague, W. M., Sytsma, K. J., & Hall, J. C. (2016). Biogeography and diversification of
Brassicales: A 103 million year tale. Molecular Phylogenetics and Evolution, 99, 204–224.
doi:10.1016/j.ympev.2016.02.021

49. Cardinal-McTeague, W. M., Wurdack, K. J., Sigel, E. M., & Gillespie, L. J. (2019). Seed size evolution
and biogeography of Plukenetia (Euphorbiaceae), a pantropical genus with traditionally cultivated oilseed
species. BMC Evolutionary Biology, 19(1). doi:10.1186/s12862-018-1308-9

50. Cassemiro, F. A. S., Albert, J. S., Antonelli, A., Menegotto, A., Wüest, R. O., Cerezer, F., Coelho, M.
T. P., Reis, R. E., Tan, M., Tagliacollo, V., Bailly, D., da Silva, V. F. B., Frota, A., da Graça, W. J., Ré, R., Ramos,
T., Oliveira, A. G., Dias, M. S., Colwell, R. K., … Graham, C. H. (2023). Landscape dynamics and
diversification of the megadiverse South American freshwater fish fauna. Proceedings of the National
Academy of Sciences, 120(2). https://doi.org/10.1073/pnas.2211974120

51. Černý, D., & Natale, R. (2022). Comprehensive taxon sampling and vetted fossils help clarify the
time tree of shorebirds (Aves, Charadriiformes). Molecular Phylogenetics and Evolution, 177, 107620.
https://doi.org/10.1016/j.ympev.2022.107620

52. Chakrabarty, Prosanta, Matthew P. Davis, W. Leo Smith, Zachary H. Baldwin, John S. Sparks. 2011.
Is sexual selection driving diversification of the bioluminescent ponyfishes (Teleostei: Leiognathidae)?
Molecular Ecology 20 (13): 2818-2834.

53. Chan, Kin Onn , Rafe M. Brown. 2017. Did true frogs 'dispersify'?' Biology Letters 13 (8): 20170299

54. Chan, L.M., Choi D., Raselimanana A.P., Rakotondravony H.E., & Yoder A.D. 2012. Defining spatial
and temporal patterns of phylogeographic structure in Madagascar's iguanid lizards (genus Oplurus).
Molecular Ecology 21 (15): 3839-3851.

55. Chen, Wei-Jen, Sébastien Lavoué, and Richard L. Mayden. 2013. Evolutionary origin and early
biogeography of otophysan fishes (Ostariophysi: Teleostei). Evolution 67 (8): 2218–2239.

56. Chris J. Law, Graham J. Slater, Rita S. Mehta, 2017, 'Lineage Diversity and Size Disparity in
Musteloidea: Testing Patterns of Adaptive Radiation Using Molecular and Fossil-Based Methods',
Systematic Biology

57. Cibois, Alice, Jean-Claude Thibault, Céline Bonillo, Christopher E. Filardi, Dick Watling, Eric
Pasquet. 2014. Phylogeny and biogeography of the fruit doves (Aves: Columbidae). Molecular
Phylogenetics and Evolution 70: 442-453.

58. Claramunt, S., Sheard, C., Brown, J. W., Cortés-Ramírez, G., Cracraft, J., Su, M. M., Weeks, B. C., &
Tobias, J. A. (2025). A new time tree of birds reveals the interplay between dispersal, geographic range size,
and diversification. Current Biology, 35(16), 3883-3895.e4. https://doi.org/10.1016/j.cub.2025.07.004

59. Claramunt, Santiago, Joel Cracraft. 2015. A new time tree reveals Earth historys imprint on the
evolution of modern birds. Science Advances 1 (11): e1501005-e1501005

60. Corl, Ammon, Hans Ellegren. 2013. Sampling strategies for species trees: The effects on
phylogenetic inference of the number of genes, number of individuals, and whether loci are mitochondrial,
sex-linked, or autosomal. Molecular Phylogenetics and Evolution 67 (2): 358-366

61. Cushing, Paula E., Matthew R. Graham, Lorenzo Prendini, Jack O. Brookhart. 2015. A multilocus
molecular phylogeny of the endemic North American camel spider family Eremobatidae (Arachnida:
Solifugae). Molecular Phylogenetics and Evolution 92: 280-293

62.   Dahiana Arcila, R. Alexander Pyron, James C. Tyler, Guillermo Ortí, Ricardo Betancur-R., 2015, 'An
evaluation of fossil tip-dating versus node-age calibrations in tetraodontiform fishes (Teleostei:
Percomorphaceae)', Molecular Phylogenetics and Evolution, vol. 82, pp. 131-145

63.   Daniel L. Rabosky, Jonathan Chang, Pascal O. Title, Peter F. Cowman, Lauren Sallan, Matt
Friedman, Kristin Kaschner, Cristina Garilao, Thomas J. Near, Marta Coll, Michael E. Alfaro, 2018, 'An
inverse latitudinal gradient in speciation rate for marine fishes', Nature, vol. 559, no. 7714, pp. 392-395

64.   Daniela Campanella, Lily C. Hughes, Peter J. Unmack, Devin D. Bloom, Kyle R. Piller, Guillermo
Ortí, 2015, 'Multi-locus fossil-calibrated phylogeny of Atheriniformes (Teleostei, Ovalentaria)', Molecular
Phylogenetics and Evolution, vol. 86, pp. 8-23

65.   Dantas, Sidnei M., Jason D. Weckstein, John M. Bates, Niels K. Krabbe, Carlos Daniel Cadena, Mark
B. Robbins, Eugenio Valderrama, Alexandre Aleixo. 2016. Molecular systematics of the new world screech-
owls (Megascops: Aves, Strigidae): biogeographic and taxonomic implications. Molecular Phylogenetics
and Evolution 94: 626-634

66.   Delsuc, Frédéric, Gillian C. Gibb, Melanie Kuch, Guillaume Billet, Lionel Hautier, John Southon,
Jean-Marie Rouillard, Juan Carlos Fernicola, Sergio F. Vizcaíno, Ross D.E. MacPhee, Hendrik N. Poinar.
2016. The phylogenetic affinities of the extinct glyptodonts. Current Biology 26 (4): R155-R156

67.   Deng, W., Luo, X., Ho, S. Y. W., Liao, S., Wang, Z., & Che, Y. (2023). Inclusion of rare taxa from
Blattidae and Anaplectidae improves phylogenetic resolution in the cockroach superfamily Blattoidea.
Systematic Entomology, 48(1), 23–39. Portico. https://doi.org/10.1111/syen.12560

68.   Derryberry, E.P., Claramunt S., Derryberry G.E., Chesser R.T., Cracraft J., Aleixo A., Perez-eman J.,
Remsen J., and Brumfield R. 2011. Lineage diversification and morphological evolution in a large-scale
continental radiation: the Neotropical ovenbirds and woodcreepers (Aves: Furnariidae). Evolution 65 (10):
2973-2986.

69.   Dexter, Kyle, Jérôme Chave. 2016. Evolutionary patterns of range size, abundance and species
richness in Amazonian angiosperm trees. PeerJ 4: e2402

70.   Dool, S. E., Puechmaille, S. J., Foley, N. M., Allegrini, B., Bastian, A., Mutumi, G. L., Maluleke, T. G.,
Odendaal, L. J., Teeling, E. C., & Jacobs, D. S. (2016). Nuclear introns outperform mitochondrial DNA in
inter-specific phylogenetic reconstruction: Lessons from horseshoe bats (Rhinolophidae: Chiroptera).
Molecular Phylogenetics and Evolution, 97, 196–212. https://doi.org/10.1016/j.ympev.2016.01.003

71.   Dornburg, Alex, Jeffrey P Townsend, Matt Friedman, Thomas J Near. 2014. Phylogenetic
informativeness reconciles ray-finned fish molecular divergence times. BMC Evolutionary Biology 14(1):
169.

72.   Doré, M., Borowiec, M. L., Branstetter, M. G., Camacho, G. P., Fisher, B. L., Longino, J. T., Ward, P.
S., & Blaimer, B. B. (2025). Evolutionary history of ponerine ants highlights how the timing of dispersal
events shapes modern biodiversity. Nature Communications, 16(1). https://doi.org/10.1038/s41467-025-
63709-3

73.   Drew, Bryan T., Kenneth J. Sytsma. 2013. The South American radiation of Lepechinia (Lamiaceae):
phylogenetics, divergence times and evolution of dioecy. Botanical Journal of the Linnean Society 171 (1):
171-190.

74.   Drummond, C. S., Eastwood, R. J., Miotto, S. T., & Hughes, C. E. (2012). Multiple continental
radiations and correlates of diversification in Lupinus (Leguminosae): testing for key innovation with
incomplete taxon sampling. Systematic Biology 61(3), 443-460.

75.   Dufort, Matthew J. 2015. An augmented supermatrix phylogeny of the avian family Picidae reveals
uncertainty deep in the family tree. Molecular Phylogenetics and Evolution

76.   Dumont E.R., Davalos L.M., Goldberg A., Santana S.E., Rex K., & Voigt C.C. 2012. Morphological
innovation, diversification and invasion of a new adaptive zone. Proceedings of the Royal Society B:
Biological Sciences, 279: 1797-1805.

77.   Dupin J., & Smith S.D. 2018. Phylogenetics of Datureae (Solanaceae), including description of the
new genus Trompettia and re-circumscription of the tribe. Taxon, 67(2): 359-375.

78.   Economo, E. P., Narula, N., Friedman, N. R., Weiser, M. D., & Guénard, B. (2018). Macroecology
and macroevolution of the latitudinal diversity gradient in ants. Nature Communications, 9(1).
https://doi.org/10.1038/s41467-018-04218-4

79.   Ericson, P. G. P., Irestedt, M., Nylander, J. A. A., Christidis, L., Joseph, L., & Qu, Y. (2020). Parallel
Evolution of Bower-Building Behavior in Two Groups of Bowerbirds Suggested by Phylogenomics.
Systematic Biology, 69(5), 820–829. https://doi.org/10.1093/sysbio/syaa040

80.   Ericson, Per GP, Seraina Klopfstein, Martin Irestedt, Jacqueline MT Nguyen and Johan AA Nylander.
2014. Dating the diversification of the major lineages of Passeriformes (Aves). BMC Evolutionary Biology,
14(8).

81.   Fabre, Pierre-Henri, Lionel Hautier, Dimitar Dimitrov, Emmanuel J P Douzery. 2012. A glimpse on
the pattern of rodent diversification: a phylogenetic approach. BMC Evolutionary Biology 12 (1): 88.

82.   Fine, Paul V. A., Felipe Zapata, Douglas C. Daly. 2014. Investigating processes of neotropical rain
forest tree diversification by examining the evolution and historical biogeography of the Protieae
(Burseraceae). Evolution 68 (7): 1988-2004

83.   Foster, Charles S. P., Hervé Sauquet, Marlien van der Merwe, Hannah McPherson, Maurizio
Rossetto, Simon Y. W. Ho. 2016. Evaluating the impact of genomic data and priors on Bayesian estimates
of the angiosperm evolutionary timescale. Systematic Biology, p. syw086

84.   Francesco Santini, Luke J Harmon, Giorgio Carnevale, Michael E Alfaro, 2009, 'Did genome
duplication drive the origin of teleosts? A comparative study of diversification in ray-finned fishes', BMC
Evolutionary Biology, vol. 9, no. 1, p. 194

85.   Frank Portillo, William R. Branch, Werner Conradie, Mark-Oliver Rödel, Johannes Penner, Michael
F. Barej, Chifundera Kusamba, Wandege M. Muninga, Mwenebatu M. Aristote, Aaron M. Bauer, Jean-
François Trape, Zoltán T. Nagy, Piero Carlino, Olivier S.G. Pauwels, Michele Menegon, Marius Burger, Tomáš
Mazuch, Kate Jackson, Daniel F. Hughes, Mathias Behangana, Ange-Ghislain Zassi-Boulou, Eli
Greenbaum, 2018, 'Phylogeny and biogeography of the African burrowing snake subfamily Aparallactinae
(Squamata: Lamprophiidae)', Molecular Phylogenetics and Evolution, vol. 127, pp. 288-303

86.   Friedman, M., B. P. Keck, A. Dornburg, R. I. Eytan, C. H. Martin, C. D. Hulsey, P. C. Wainwright, T. J.
Near. 2013. Molecular and fossil evidence place the origin of cichlid fishes long after Gondwanan rifting.
Proceedings of the Royal Society B: Biological Sciences 280 (1770): 20131733

87.   Frolová, P., Horká, I., & Ďuriš, Z. (2022). Molecular phylogeny and historical biogeography of marine palaemonid shrimps (Palaemonidae: Palaemonella–Cuapetes group). Scientific Reports, 12(1). https://doi.org/10.1038/s41598-022-19372-5

88.   Fuchs, J., Johnson, J. A., & Mindell, D. P. (2015). Rapid diversification of falcons (Aves: Falconidae) due to expansion of open habitats in the Late Miocene. Molecular Phylogenetics and Evolution, 82, 166–182. https://doi.org/10.1016/j.ympev.2014.08.010

89.   Fuchs, J., Pasquet, E., Stuart, B. L., Woxvold, I. A., Duckworth, J. W., & Bowie, R. C. K. (2018). Phylogenetic affinities of the enigmatic Bare-faced Bulbul Pycnonotus hualon with description of a new genus. Ibis, 160(3), 659–665. Portico. https://doi.org/10.1111/ibi.12580

90.   Fusca, D. D., Dall'Acqua, M. N., Sánchez-Ramírez, S., & Cutter, A. D. (2025). Phylogenomic timetree-calibrated speciation clocks for Caenorhabditis nematodes reveal slow but disproportionate accumulation of post-zygotic reproductive isolation. PLOS Genetics, 21(9), e1011852. https://doi.org/10.1371/journal.pgen.1011852

91.   Garcia-R., J. C., & Trewick, S. A. (2015). Dispersal and speciation in purple swamphens (Rallidae:Porphyrio). The Auk, 132(1), 140–155. https://doi.org/10.1642/auk-14-114.1

92.   García-Girón, J., Múrria, C., Arnedo, M. A., Bonada, N., Cañedo-Argüelles, M., Derka, T., Fernández-Calero, J. M., Li, Z., Tierno de Figueroa, J. M., Xie, Z., & Heino, J. (2024). A time-calibrated 'Tree of Life' of aquatic insects for knitting historical patterns of evolution and measuring extant phylogenetic biodiversity across the world. Earth-Science Reviews, 252, 104767. https://doi.org/10.1016/j.earscirev.2024.104767

93.   García–R, Juan C., Gillian C. Gibb, Steve A. Trewick. 2014. Eocene diversification of crown group rails (Aves: Gruiformes: Rallidae). PLoS ONE 9 (10): e109635

94.   Gavryushkina, Alexandra, Tracy A. Heath, Daniel T. Ksepka, Tanja Stadler, David Welch, Alexei J. Drummond. 2016. Bayesian Total-Evidence Dating Reveals the Recent Crown Radiation of Penguins. Systematic Biology, p. syw060

95.   Gaynor, M. L., Fu, C., Gao, L., Lu, L., Soltis, D. E., & Soltis, P. S. (2023). Biogeography and ecological niche evolution in Diapensiaceae inferred from phylogenetic analysis. Journal of Systematics and Evolution, 58(5), 646–662. Portico. https://doi.org/10.1111/jse.12646

96.   Ghezelayagh, A., Harrington, R. C., Burress, E. D., Campbell, M. A., Buckner, J. C., Chakrabarty, P., Glass, J. R., McCraney, W. T., Unmack, P. J., Thacker, C. E., Alfaro, M. E., Friedman, S. T., Ludt, W. B., Cowman, P. F., Friedman, M., Price, S. A., Dornburg, A., Faircloth, B. C., Wainwright, P. C., & Near, T. J. (2022). Prolonged morphological expansion of spiny-rayed fishes following the end-Cretaceous. Nature Ecology & Evolution, 6(8), 1211–1220. https://doi.org/10.1038/s41559-022-01801-3

97.   Gibb, G. C., & Shepherd, L. D. (2022). Recent evolution of extreme sexual dimorphism in the huia (Heteralocha acutirostris; Callaeidae). Molecular Phylogenetics and Evolution, 175, 107575. https://doi.org/10.1016/j.ympev.2022.107575

98.   Gibb, Gillian C., Fabien L. Condamine, Melanie Kuch, Jacob Enk, Nadia Moraes-Barros, Mariella Superina, Hendrik N. Poinar, Frédéric Delsuc. 2015. Shotgun mitogenomics provides a reference phylogenetic framework and timescale for living Xenarthrans. Molecular Biology and Evolution 33 (3): 621-642

99.   Gibb, Gillian C., Martyn Kennedy, David Penny. 2013. Beyond phylogeny: pelecaniform and ciconiiform birds, and long-term niche stability. Molecular Phylogentics and Evolution 68 (2): 229-238.

100.   Gibb, Gillian C., Ryan England, Gerrit Hartig, P.A. (Trish) McLenachan, Briar L. Taylor Smith, Bennet J. McComish, Alan Cooper, David Penny. 2015. New Zealand passerines help clarify the diversification of major songbird lineages during the Oligocene. Genome Biology and Evolution 7 (11): 2983-2995.

101.   Grealy, Alicia, Matthew Phillips, Gifford Miller, M. Thomas P. Gilbert, Jean-Marie Rouillard, David Lambert, Michael Bunce, James Haile. 2017. Eggshell palaeogenomics: Palaeognath evolutionary history revealed through ancient nuclear and mitochondrial DNA from Madagascan elephant bird (Aepyornis sp.) eggshell. Molecular Phylogenetics and Evolution 109: 151-163

102.   Harrington, Richard C., Brant C. Faircloth, Ron I. Eytan, W. Leo Smith, Thomas J. Near, Michael E. Alfaro, Matt Friedman. 2016. Phylogenomic analysis of carangimorph fishes reveals flatfish asymmetry arose in a blink of the evolutionary eye. BMC Evolutionary Biology 16: 224

103.   Harris, Liam W., T. Jonathan Davies. 2016. A Complete Fossil-Calibrated Phylogeny of Seed Plant Families as a Tool for Comparative Analyses: Testing the 'Time for Speciation' Hypothesis. PLoS ONE 11 (10): e0162907

104.   Haynie, M. L., Tsuchiya, M. T. N., Ospina-Garcés, S. M., Arroyo-Cabrales, J., Medellín, R. A., Polaco, O. J., & Maldonado, J. E. (2016). Placement of the rediscoveredMyotis planiceps(Chiroptera: Vespertilionidae) within theMyotisphylogeny. Journal of Mammalogy, 97(3), 701–712. https://doi.org/10.1093/jmammal/gyv216

105.   He, Kai, Akio Shinohara, Xue-Long Jiang, Kevin L. Campbell. 2014. Multilocus phylogeny of talpine moles (Talpini, Talpidae, Eulipotyphla) and its implications for systematics. Molecular Phylogenetics and Evolution 70: 513-521.

106.   Hedges, S. Blair, Julie Marin, Michael Suleski, Madeline Paymer, Sudhir Kumar. 2015. Tree of life reveals clock-like speciation and diversification. Molecular Biology and Evolution 32 (4): 835-845

107.   Henríquez-Piskulich, P., Hugall, A. F., & Stuart-Fox, D. (2024). A supermatrix phylogeny of the world's bees (Hymenoptera: Anthophila). Molecular Phylogenetics and Evolution, 190, 107963. https://doi.org/10.1016/j.ympev.2023.107963

108.   Higdon J. W., Bininda-Emonds O. R. P., Beck R. M. D., Ferguson S. H. 2007. Phylogeny and divergence of the pinnipeds (Carnivora: Mammalia) assessed using a multigene dataset. BMC Evolutionary Biology 8: 216.

109.   Hipp, A. L., Manos, P. S., Hahn, M., Avishai, M., Bodénès, C., Cavender-Bares, J., … Valencia-Avalos, S. (2019). Genomic landscape of the global oak phylogeny. New Phytologist. doi:10.1111/nph.16162

110.   Hooper, Daniel M., Trevor D. Price. 2017. Chromosomal inversion differences correlate with range overlap in passerine birds. Nature Ecology & Evolution 1 (10): 1526-1534

111.   Hosner, P. A., Braun, E. L., & Kimball, R. T. (2016). Rapid and recent diversification of curassows, guans, and chachalacas (Galliformes: Cracidae) out of Mesoamerica: Phylogeny inferred from mitochondrial, intron, and ultraconserved element sequences. Molecular Phylogenetics and Evolution, 102, 320–330. https://doi.org/10.1016/j.ympev.2016.06.006

112.   Hubert N., Paradis E., Bruggemann H., & Planes S. 2011. Community assembly and diversification
in Indo-Pacific coral reef fishes. Ecology and Evolution, 1(3): 229-277.

113.   Imfeld, Barker, Brumfield, Mitochondrial genomes and thousands of ultraconserved elements
resolve the taxonomy and historical biogeography of the Euphonia and Chlorophonia finches
(Passeriformes: Fringillidae), The Auk, Volume 137, Issue 3, 1 July 2020, ukaa016,

114.   Imfeld, T. S., Barker, F. K., Vázquez-Miranda, H., Chaves, J. A., Escalante, P., Spellman, G. M., &
Klicka, J. (2024). Diversification and dispersal in the Americas revealed by new phylogenies of the wrens
and allies (Passeriformes: Certhioidea). Ornithology, 141(2). https://doi.org/10.1093/ornithology/ukae007

115.   Inoue, J. G., Miya, M., Lam, K., Tay, B.-H., Danks, J. A., Bell, J., Walker, T. I., & Venkatesh, B. (2010).
Evolutionary Origin and Phylogeny of the Modern Holocephalans (Chondrichthyes: Chimaeriformes): A
Mitogenomic   Perspective.   Molecular   Biology   and   Evolution,   27(11),   2576–2586.
https://doi.org/10.1093/molbev/msq147

116.   Irisarri, I., Mauro, D. S., Abascal, F., Ohler, A., Vences, M., & Zardoya, R. (2012). The origin of modern
frogs (Neobatrachia) was accompanied by acceleration in mitochondrial and nuclear substitution rates.
BMC Genomics, 13(1). doi:10.1186/1471-2164-13-626

117.   Islam, S., Peart, C., Kehlmaier, C., Sun, Y.-H., Lei, F., Dahl, A., Klemroth, S., Alexopoulou, D., del
Mar Delgado, M., Laiolo, P., Carlos Illera, J., Dirren, S., Hille, S., Lkhagvasuren, D., Töpfer, T., Kaiser, M.,
Gebauer, A., Martens, J., Paetzold, C., & Päckert, M. (2024). Museomics help resolving the phylogeny of
snowfinches (Aves, Passeridae, Montifringilla and allies). Molecular Phylogenetics and Evolution, 198,
108135. https://doi.org/10.1016/j.ympev.2024.108135

118.   James B. Pease, Joseph W. Brown, Joseph F. Walker, Cody E. Hinchliff, Stephen A. Smith, 2018,
'Quartet Sampling distinguishes lack of support from conflicting support in the green plant tree of life',
American Journal of Botany

119.   Janssens, S., Couvreur, T. L. P., Mertens, A., Dauby, G., Dagallier, L.-P., Vanden Abeele, S.,
Vandelook, F., Mascarello, M., Beeckman, H., Sosef, M., Droissart, V., van der Bank, M., Maurin, O.,
Hawthorne, W., Marshall, C., Réjou-Méchain, M., Beina, D., Baya, F., Merckx, V., ... Hardy, O. (2020). A large-
scale species level dated angiosperm phylogeny for evolutionary and ecological analyses. Biodiversity
Data Journal, 8. https://doi.org/10.3897/bdj.8.e39677

120.   Jarvis, E. D., S. Mirarab, A. J. Aberer, B. Li, P. Houde, C. Li, S. Y. W. Ho, B. C. Faircloth, B. Nabholz, J.
T. Howard, A. Suh, C. C. Weber, R. R. da Fonseca, J. Li, F. Zhang, H. Li, L. Zhou, N. Narula, L. Liu, G.
Ganapathy, B. Boussau, M. S. Bayzid, V. Zavidovych, S. Subramanian, T. Gabaldon, S. Capella-Gutierrez, J.
Huerta-Cepas, B. Rekepalli, K. Munch, M. Schierup, B. Lindow, W. C. Warren, D. Ray, R. E. Green, M. W.
Bruford, X. Zhan, A. Dixon, S. Li, N. Li, Y. Huang, E. P. Derryberry, M. F. Bertelsen, F. H. Sheldon, R. T.
Brumfield, C. V. Mello, P. V. Lovell, M. Wirthlin, M. P. C. Schneider, F. Prosdocimi, J. A. Samaniego, A. M. V.
Velazquez, A. Alfaro-Nunez, P. F. Campos, B. Petersen, T. Sicheritz-Ponten, A. Pas, T. Bailey, P. Scofield, M.
Bunce, D. M. Lambert, Q. Zhou, P. Perelman, A. C. Driskell, B. Shapiro, Z. Xiong, Y. Zeng, S. Liu, Z. Li, B. Liu,
K. Wu, J. Xiao, X. Yinqi, Q. Zheng, Y. Zhang, H. Yang, J. Wang, L. Smeds, F. E. Rheindt, M. Braun, J. Fjeldsa, L.
Orlando, F. K. Barker, K. A. Jonsson, W. Johnson, K.-P. Koepfli, S. O'Brien, D. Haussler, O. A. Ryder, C.
Rahbek, E. Willerslev, G. R. Graves, T. C. Glenn, J. McCormack, D. Burt, H. Ellegren, P. Alstrom, S. V.
Edwards, A. Stamatakis, D. P. Mindell, J. Cracraft, E. L. Braun, T. Warnow, W. Jun, M. T. P. Gilbert, G. Zhang.
2014. Whole-genome analyses resolve early branches in the tree of life of modern birds. Science 346
(6215): 1320-1331.

121.   Jeffrey L. Weinell, Aaron M. Bauer, 2018, 'Systematics and phylogeography of the widely distributed
African skink Trachylepis varia species complex', Molecular Phylogenetics and Evolution, vol. 120, pp. 103-
117

122.   Jeffrey L. Weinell, Rafe M. Brown, 2018, 'Discovery of an old, archipelago-wide, endemic radiation
of Philippine snakes', Molecular Phylogenetics and Evolution, vol. 119, pp. 144-150

123.   Jennifer L. Fessler, Mark W. Westneat, 2007, 'Molecular phylogenetics of the butterflyfishes
(Chaetodontidae): Taxonomy and biogeography of a global coral reef fish family', Molecular Phylogenetics
and Evolution, vol. 45, no. 1, pp. 50-68

124.   Jetz, W., G. H. Thomas, J. B. Joy, K. Hartmann, A. O. Mooers. 2012. The global diversity of birds in
space and time. Nature 491 (7424): 444-448

125.   Jha, A., Seneviratne, S., Prayag, H. S., & Vasudevan, K. (2021). Phylogeny identifies multiple
colonisation events and Miocene aridification as drivers of South Asian bulbul (Passeriformes:
Pycnonotidae)   diversification.   Organisms   Diversity   &   Evolution,   21(4),   783–794.
https://doi.org/10.1007/s13127-021-00506-y

126.   Joel P. Olfelt, William A. Freyman, 2014, ' Relationships of North American members of Rhodiola
(Crassulaceae) ', Botany, vol. 92, no. 12, pp. 901-910

127.   Johnson, Jeff A., Joseph W. Brown, Jérôme Fuchs, David P. Mindell, 2016, 'Multi-locus phylogenetic
inference among New World Vultures (Aves: Cathartidae)', Molecular Phylogenetics and Evolution, vol. 105,
pp. 193-199

128.   Johnson, K. P., Dietrich, C. H., Friedrich, F., Beutel, R. G., Wipfler, B., Peters, R. S., Allen, J. M.,
Petersen, M., Donath, A., Walden, K. K. O., Kozlov, A. M., Podsiadlowski, L., Mayer, C., Meusemann, K.,
Vasilikopoulos, A., Waterhouse, R. M., Cameron, S. L., Weirauch, C., Swanson, D. R., … Yoshizawa, K.
(2018). Phylogenomics and the evolution of hemipteroid insects. Proceedings of the National Academy of
Sciences, 115(50), 12775–12780. https://doi.org/10.1073/pnas.1815820115

129.   Johnson, M. T., FitzJohn, R. G., Smith, S. D., Rausher, M. D., & Otto, S. P. (2011). Loss of sexual
recombination and segregation is associated with increased diversification in evening primroses.
Evolution, 65(11), 3230-3240.

130.   Johnson, O., Howard, J. T., & Brumfield, R. T. (2021). Systematics of a Neotropical clade of dead-
leaf-foraging antwrens (Aves: Thamnophilidae; Epinecrophylla). Molecular Phylogenetics and Evolution,
154, 106962. https://doi.org/10.1016/j.ympev.2020.106962

131.   Jose Tavera, Arturo Acero P., Peter C. Wainwright, 2018, 'Multilocus phylogeny, divergence times,
and a major role for the benthic-to-pelagic axis in the diversification of grunts (Haemulidae)', Molecular
Phylogenetics and Evolution, vol. 121, pp. 212-223

132.   Jønsson, Knud Andreas, Pierre-Henri Fabre, Jonathan D. Kennedy, Ben G. Holt, Michael K.
Borregaard, Carsten Rahbek, Jon Fjeldså. 2016. A supermatrix phylogeny of corvoid passerine birds (Aves:
Corvides). Molecular Phylogenetics and Evolution 94: 87-94.

133.   Kathryn M. Everson, Voahangy Soarimalala, Steven M. Goodman, Link E. Olson, 2016, 'Multiple
Loci and Complete Taxonomic Sampling Resolve the Phylogeny and Biogeographic History of Tenrecs
(Mammalia: Tenrecidae) and Reveal Higher Speciation Rates in Madagascar's Humid Forests', Systematic
Biology, vol. 65, no. 5, pp. 890-909

134.   Kawahara, A. Y., Storer, C., Carvalho, A. P. S., Plotkin, D. M., Condamine, F. L., Braga, M. P., Ellis, E.
A., St Laurent, R. A., Li, X., Barve, V., Cai, L., Earl, C., Frandsen, P. B., Owens, H. L., Valencia-Montoya, W.
A., Aduse-Poku, K., Toussaint, E. F. A., Dexter, K. M., Doleck, T., ... Lohman, D. J. (2023). A global phylogeny
of butterflies reveals their evolutionary history, ancestral hosts and biogeographic origins. Nature Ecology
& Evolution. https://doi.org/10.1038/s41559-023-02041-9

135.   Kennedy, J. D., Marki, P. Z., Reeve, A. H., Blom, M. P. K., Prawiradilaga, D. M., Haryoko, T., Koane, B.,
Kamminga, P., Irestedt, M., Jønsson, K. A., & Sheard, C. (2022). Diversification and community assembly of
the world's largest tropical island. Global Ecology and Biogeography. Portico.
https://doi.org/10.1111/geb.13484

136.   Kimball, R. T., Oliveros, C. H., Wang, N., White, N. D., Barker, F. K., Field, D. J., ... Braun, E. L. (2019).
A Phylogenomic Supertree of Birds. Diversity, 11(7), 109. doi:10.3390/d11070109

137.   Kimberly L. Foster, Kyle R. Piller, 2018, 'Disentangling the drivers of diversification in an imperiled
group of freshwater fishes (Cyprinodontiformes: Goodeidae)', BMC Evolutionary Biology, vol. 18, no. 1

138.   Koepfli, Klaus-Peter, Kerry A Deere, Graham J Slater, Colleen Begg, Keith Begg, Lon Grassman,
Mauro Lucherini, Geraldine Veron, Robert K Wayne. 2008. Multigene phylogeny of the Mustelidae:
Resolving relationships, tempo and biogeographic history of a mammalian adaptive radiation. BMC Biology
6 (1): 10.

139.   K. M. Kozak, N. Wahlberg, A. F. E. Neild, K. K. Dasmahapatra, J. Mallet, C. D. Jiggins, 2015,
'Multilocus Species Trees Show the Recent Adaptive Radiation of the Mimetic Heliconius Butterflies',
Systematic Biology, vol. 64, no. 3, pp. 505-524

140.   Ksepka, Daniel T., Matthew J. Phillips. 2015. Avian diversification patterns across the K-Pg
boundary: influence of calibrations, datasets, and model misspecification. Annals of the Missouri
Botanical Garden 100 (4): 300-328

141.   Kuo L., Chen C., Shinohara W., Ebihara A., Hiroshi K., Hirotoshi S., Huang Y., & Chiou W. 2017. Not
only in the temperate zone: independent gametophytes of two vittarioid ferns (Pteridaceae, Polypodiales)
in East Asian subtropics. Journal of Plant Research, 130: 407-416.

142.   Lack J.B., & Van den bussche R.A. 2010. Identifying the Confounding Factors in Resolving
Phylogenetic Relationships in Vespertilionidae. Journal of Mammalogy, .

143.   Lapiedra, O., Sayol, F., Garcia-Porta, J., & Sol, D. (2021). Niche shifts after island colonization
spurred adaptive diversification and speciation in a cosmopolitan bird clade. Proceedings of the Royal
Society B: Biological Sciences, 288(1958), 20211022. https://doi.org/10.1098/rspb.2021.1022

144.   Larsen, P. A., Siles, L., Pedersen, S. C., & Kwiecinski, G. G. (2011). A new species of Micronycteris
(Chiroptera: Phyllostomidae) from Saint Vincent, Lesser Antilles. Mammalian Biology, 76(6), 687–700.
https://doi.org/10.1016/j.mambio.2011.01.006

145.   Lavrov, D. V., Diaz, M. C., Maldonado, M., Morrow, C. C., Perez, T., Pomponi, S. A., & Thacker, R. W.
(2023). Phylomitogenomics bolsters the high-level classification of Demospongiae (phylum Porifera). PLOS
ONE, 18(12), e0287281. https://doi.org/10.1371/journal.pone.0287281

146.   Lee, M. S., Soubrier, J., & Edgecombe, G. D. (2013). Rates of phenotypic and genomic evolution
during the Cambrian explosion. Current Biology 23 (19): 1889-1895.

147.	Leslie, A. B., J. M. Beaulieu, H. S. Rai, P. R. Crane, M. J. Donoghue, S. Mathews. 2012. Hemisphere-scale differences in conifer evolutionary dynamics. Proceedings of the National Academy of Sciences. 109 (40): 16217-16221.

148.	Licht, M., Schmuecker, K., Huelsken, T., Hanel, R., Bartsch, P., & Paeckert, M. (2012). Contribution to the molecular phylogenetic analysis of extant holocephalan fishes (Holocephali, Chimaeriformes). Organisms Diversity & Evolution, 12(4), 421–432. https://doi.org/10.1007/s13127-011-0071-1

149.	Lin, Qiang, Shaohua Fan, Yanhong Zhang, Meng Xu, Huixian Zhang, Yulan Yang, Alison P. Lee, Joost M. Woltering, Vydianathan Ravi, Helen M. Gunter, Wei Luo, Zexia Gao, Zhi Wei Lim, Geng Qin, Ralf F. Schneider, Xin Wang, Peiwen Xiong, Gang Li, Kai Wang, Jiumeng Min, Chi Zhang, Ying Qiu, Jie Bai, Weiming He, Chao Bian, Xinhui Zhang, Dai Shan, Hongyue Qu, Ying Sun, Qiang Gao, Liangmin Huang, Qiong Shi, Axel Meyer, Byrappa Venkatesh. 2016. The seahorse genome and the evolution of its specialized morphology. Nature 540 (7633): 395-399

150.	Lincoln Carneiro, Gustavo A. Bravo, Natalia Aristizábal, Andrés M. Cuervo, Alexandre Aleixo, 2018, 'Molecular systematics and biogeography of lowland antpittas (Aves, Grallariidae): The role of vicariance and dispersal in the diversification of a widespread Neotropical lineage', Molecular Phylogenetics and Evolution, vol. 120, pp. 375-389

151.	Liu, B., Alström, P., Olsson, U., Fjeldså, J., Quan, Q., Roselaar, K. C. S., Saitoh, T., Yao, C., Hao, Y., Wang, W., Qu, Y., & Lei, F. (2017). Explosive radiation and spatial expansion across the cold environments of the Old World in an avian family. Ecology and Evolution, 7(16), 6346–6357. Portico. https://doi.org/10.1002/ece3.3136

152.	Lohmann, L. G., Bell, C. D., Calió, M. F., & Winkworth, R. C. (2013). Pattern and timing of biogeographical history in the Neotropical tribe Bignonieae (Bignoniaceae). Botanical Journal of the Linnean Society, 171(1), 154-170.

153.	Magallón, Susana, Sandra Gómez-Acevedo, Luna L. Sánchez-Reyes, Tania Hernández-Hernández. 2015. A metacalibrated time-tree documents the early rise of flowering plant phylogenetic diversity. New Phytologist 207 (2): 437-453

154.	Mahler, D. L., T. Ingram, L. J. Revell, J. B. Losos. 2013. Exceptional Convergence on the Macroevolutionary Landscape in Island Lizard Radiations. Science 341 (6143): 292-295.

155.	K. Mao, R. I. Milne, L. Zhang, Y. Peng, J. Liu, P. Thomas, R. R. Mill, S. S. Renner. 2012. Distribution of living Cupressaceae reflects the breakup of Pangea. Proceedings of the National Academy of Sciences 109(20):7793-7798.

156.	Marianne Espeland, Jason P.W. Hall, Philip J. DeVries, David C. Lees, Mark Cornwall, Yu-Feng Hsu, Li-Wei Wu, Dana L. Campbell, Gerard Talavera, Roger Vila, Shayla Salzman, Sophie Ruehr, David J. Lohman, Naomi E. Pierce, 2015, 'Ancient Neotropical origin and recent recolonisation: Phylogeny, biogeography and diversification of the Riodinidae (Lepidoptera: Papilionoidea)', Molecular Phylogenetics and Evolution, vol. 93, pp. 296-306

157.	Marisol Sánchez-García, Patrick Brandon Matheny, 2016, 'Is the switch to an ectomycorrhizal state an evolutionary key innovation in mushroom-forming fungi? A case study in the Tricholomatineae (Agaricales)', Evolution, vol. 71, no. 1, pp. 51-65

158.   Marki, Petter Z., Knud A. Jønsson, Martin Irestedt, Jacqueline M.T. Nguyen, Carsten Rahbek, Jon
Fjeldså. 2017. Supermatrix phylogeny and biogeography of the Australasian Meliphagides radiation (Aves:
Passeriformes). Molecular Phylogenetics and Evolution 107: 516-529

159.   Martin Malmstrøm, Michael Matschiner, Ole K Tørresen, Bastiaan Star, Lars G Snipen, Thomas F
Hansen, Helle T Baalsrud, Alexander J Nederbragt, Reinhold Hanel, Walter Salzburger, Nils C Stenseth,
Kjetill S Jakobsen, Sissel Jentoft, 2016, 'Evolution of the immune system influences speciation rates in
teleost fishes', Nature Genetics

160.   Martin Schwentner, Thomas C.G. Bosch, 2015, 'Revisiting the age, evolutionary history and species
level diversity of the genus Hydra (Cnidaria: Hydrozoa)', Molecular Phylogenetics and Evolution, vol. 91, pp.
41-55

161.   McCord, C. L., Nash, C. M., Cooper, W. J., & Westneat, M. W. (2021). Phylogeny of the damselfishes
(Pomacentridae) and patterns of asymmetrical diversification in body size and feeding ecology. PLOS ONE,
16(10), e0258889. doi:10.1371/journal.pone.0258889

162.   McCord, Charlene L., Mark W. Westneat. 2016. Phylogenetic relationships and the evolution of
BMP4 in triggerfishes and filefishes (Balistoidea). Molecular Phylogenetics and Evolution 94: 397-409

163.   McCullough, J. M., Moyle, R. G., Smith, B. T., & Andersen, M. J. (2019). A Laurasian origin for a
pantropical bird radiation is supported by genomic and fossil data (Aves: Coraciiformes). Proceedings of
the Royal Society B: Biological Sciences, 286(1910), 20190122. https://doi.org/10.1098/rspb.2019.0122

164.   McCullough, J. M., Oliveros, C. H., Benz, B. W., Zenil-Ferguson, R., Cracraft, J., Moyle, R. G., &
Andersen, M. J. (2022). Wallacean and Melanesian Islands Promote Higher Rates of Diversification within
the Global Passerine Radiation Corvides. Systematic Biology. https://doi.org/10.1093/sysbio/syac044

165.   McGuire, Jimmy A., Christopher C. Witt, J.V. Remsen, Ammon Corl, Daniel L. Rabosky, Douglas L.
Altshuler, Robert Dudley. 2014. Molecular phylogenetics and the diversification of hummingbirds. Current
Biology 24 (8): 910-916.

166.   Mckenna, Duane D., Alexander L. Wild, Kojun Kanda, Charles L. Bellamy, Rolf G. Beutel, Michael
S. Caterino, Charles W. Farnum, David C. Hawks, Michael A. Ivie, Mary Liz Jameson, Richard A. B. Leschen,
Adriana E. Marvaldi, Joseph V. Mchugh, Alfred F. Newton, James A. Robertson, Margaret K. Thayer, Michael
F. Whiting, John F. Lawrence, Adam Ślipiński, David R. Maddison, Brian D. Farrell. 2015. The beetle tree of
life reveals that Coleoptera survived end-Permian mass extinction to diversify during the Cretaceous
terrestrial revolution. Systematic Entomology 40 (4): 835-880

167.   Michael E. Alfaro, Brant C. Faircloth, Richard C. Harrington, Laurie Sorenson, Matt Friedman,
Christine E. Thacker, Carl H. Oliveros, David Černý, Thomas J. Near, 2018, 'Explosive diversification of
marine fishes at the Cretaceous–Palaeogene boundary', Nature Ecology & Evolution, vol. 2, no. 4, pp. 688-
696

168.   Michelangeli, F. A., Dellinger, A. S., Goldenberg, R., Almeda, F., Mendoza-Cifuentes, H.,
Fernández-Fernández, D., Ulloa Ulloa, C., & Penneys, D. S. (2022). Phylogenetics and Taxonomy of the Tribe
Merianieae.   Systematics,   Evolution,   and   Ecology   of   Melastomataceae,   255–273.
https://doi.org/10.1007/978-3-030-99742-7_11

169.   Mikkelsen, E. K., & Weir, J. T. (2022). Phylogenomics Reveals that Mitochondrial Capture and
Nuclear Introgression Characterize Skua Species Proposed to be of Hybrid Origin. Systematic Biology,
72(1), 78–91. https://doi.org/10.1093/sysbio/syac078

170.   Millicent D. Sanciangco, Kent E. Carpenter, Ricardo Betancur-R., 2016, 'Phylogenetic placement
of enigmatic percomorph families (Teleostei: Percomorphaceae)', Molecular Phylogenetics and Evolution,
vol. 94, pp. 565-576

171.   Milá, B., Bruxaux, J., Friis, G., Sam, K., Ashari, H., & Thébaud, C. (2021). A new, undescribed species
of   Melanocharis   berrypecker from western New Guinea and the evolutionary history of the family
Melanocharitidae. Ibis, 163(4), 1310–1329. Portico. https://doi.org/10.1111/ibi.12981

172.   Mitchell, K. J., B. Llamas, J. Soubrier, N. J. Rawlence, T. H. Worthy, J. Wood, M. S. Y. Lee, A. Cooper.
2014. Ancient DNA reveals elephant birds and kiwi are sister taxa and clarifies ratite bird evolution. Science
344 (6186): 898-900.

173.   Mitchell, Kieren J., Jamie R. Wood, Bastien Llamas, Patricia A. McLenachan, Olga Kardailsky, R.
Paul Scofield, Trevor H. Worthy, Alan Cooper. 2016. Ancient mitochondrial genomes clarify the evolutionary
history of New Zealand's enigmatic acanthisittid wrens. Molecular Phylogenetics and Evolution 102: 295-
304

174.   Mongiardino Koch, N., Thompson, J. R., Hiley, A. S., McCowin, M. F., Armstrong, A. F., Coppard, S.
E., Aguilera, F., Bronstein, O., Kroh, A., Mooi, R., & Rouse, G. W. (2022). Phylogenomic analyses of echinoid
diversification   prompt   a   re-evaluation   of   their   fossil   record.   ELife,   11.   CLOCKSS.
https://doi.org/10.7554/elife.72460

175.   Moreau C.S., & Bell C. 2013. Testing the museum versus cradle tropical biological diversity
hypothesis: Phylogeny, diversification, and ancestral biogeographic range evolution of the ants. Evolution,
67(8): 2240-2257.

176.   Moyle, Robert G., Carl H. Oliveros, Michael J. Andersen, Peter A. Hosner, Brett W. Benz, Joseph D.
Manthey, Scott L. Travers, Rafe M. Brown, Brant C. Faircloth. 2016. Tectonic collision and uplift of Wallacea
triggered the global songbird radiation. Nature Communications 7: 12709

177.   Nash, J. A., Harrington, R. C., Zyskowski, K., Near, T. J., & Prum, R. O. (2024). Species status and
phylogenetic relationships of the enigmatic Negros Fruit Dove (Ptilinopus arcanus). Ibis, 166(3), 1023–
1040. Portico. https://doi.org/10.1111/ibi.13305

178.   Nattier, R., Robillard, T., Desutter-Grandcolas, L., Couloux, A., & Grandcolas, P. (2011). Older than
New Caledonia emergence? A molecular phylogenetic study of the eneopterine crickets (Orthoptera:
Grylloidea).   Journal   of   Biogeography,   38(11),   2195–2209.   https://doi.org/10.1111/j.1365-
2699.2011.02563.x

179.   Near, T. J., A. Dornburg, R. I. Eytan, B. P. Keck, W. L. Smith, K. L. Kuhn, J. A. Moore, S. A. Price, F. T.
Burbrink, M. Friedman, P. C. Wainwright, 2013, 'Phylogeny and tempo of diversification in the
superradiation of spiny-rayed fishes', Proceedings of the National Academy of Sciences, vol. 110, no. 31,
pp. 12738-12743

180.   Nelsen, M. P., Ree, R. H., & Moreau, C. S. (2018). Ant–plant interactions evolved through increasing
interdependence.   Proceedings   of   the   National   Academy   of   Sciences,   115(48),   12253–12258.
https://doi.org/10.1073/pnas.1719794115

181.   Nitta, J. H., Schuettpelz, E., Ramírez-Barahona, S., & Iwasaki, W. (2022). An open and continuously
updated fern tree of life. Frontiers in Plant Science, 13. https://doi.org/10.3389/fpls.2022.909768

182. Nyakatura, Katrin, Olaf RP Bininda-Emonds. 2012. Updating the evolutionary history of Carnivora (Mammalia): a new species-level supertree complete with divergence time estimates. BMC Biology 10 (1): 12

183. Ogawa, Lisa M., Paulo C. Pulgarin, Donald A. Vance, Jon Fjeldså, Marcel van Tuinen. 2015. Opposing demographic histories reveal rapid evolution in grebes (Aves: Podicipedidae). The Auk 132 (4): 771-786

184. Oliver, P. M., Heiniger, H., Hugall, A. F., Joseph, L., & Mitchell, K. J. (2020). Oligocene divergence of frogmouth birds (Podargidae) across Wallace's Line. Biology Letters, 16(5), 20200040. https://doi.org/10.1098/rsbl.2020.0040

185. Oliveros, C. H., Field, D. J., Ksepka, D. T., Barker, F. K., Aleixo, A., Andersen, M. J., … Faircloth, B. C. (2019). Earth history and the passerine superradiation. Proceedings of the National Academy of Sciences, 116(16), 7916–7925. doi:10.1073/pnas.1813206116

186. Orr, R. J. S., Di Martino, E., Ramsfjell, M. H., Gordon, D. P., Berning, B., Chowdhury, I., Craig, S., Cumming, R. L., Figuerola, B., Florence, W., Harmelin, J.-G., Hirose, M., Huang, D., Jain, S. S., Jenkins, H. L., Kotenko, O. N., Kuklinski, P., Lee, H. E., Madurell, T., … Liow, L. H. (2022). Paleozoic origins of cheilostome bryozoans and their parental care inferred by a new genome-skimmed phylogeny. Science Advances, 8(13). https://doi.org/10.1126/sciadv.abm7452

187. Ottenburghs, Jente, Hendrik-Jan Megens, Robert H.S. Kraus, Ole Madsen, Pim van Hooft, Sipke E. van Wieren, Richard P.M.A. Crooijmans, Ronald C. Ydenberg, Martien A.M. Groenen, Herbert H.T. Prins. 2016. A Tree of Geese: A Phylogenomic Perspective on the Evolutionary History of True Geese. Molecular Phylogenetics and Evolution 101: 303-313.

188. Parfrey, L. W., D. J. G. Lahr, A. H. Knoll, L. A. Katz. 2011. Estimating the timing of early eukaryotic diversification with multigene molecular clocks. Proceedings of the National Academy of Sciences 108 (33): 13624-13629.

189. Perry G. Beasley-Hall, Simon M. Tierney, Phillip Weinstein, Andrew D. Austin, 2018, 'A revised phylogeny of macropathine cave crickets (Orthoptera: Rhaphidophoridae) uncovers a paraphyletic Australian fauna', Molecular Phylogenetics and Evolution, vol. 126, pp. 153-161

190. Peter A. Hosner, Edward L. Braun, Rebecca T. Kimball, 2015, 'Land connectivity changes and global cooling shaped the colonization history and diversification of New World quail (Aves: Galliformes: Odontophoridae)', Journal of Biogeography, vol. 42, no. 10, pp. 1883-1895

191. Peterson, Rose D., et al. "Phylogenomics of Bony-Tongue Fishes (Osteoglossomorpha) Shed Light on the Craniofacial Evolution and Biogeography of the Weakly Electric Clade (Mormyridae)." Systematic Biology (2022).

192. Pinto-Ledezma, J. N., Jahn, A. E., Cueto, V. R., Diniz-Filho, J. A. F., & Villalobos, F. (2019). Drivers of Phylogenetic Assemblage Structure of the Furnariides, a Widespread Clade of Lowland Neotropical Birds. The American Naturalist, 193(2), E41–E56. https://doi.org/10.1086/700696

193. Price, Trevor D., Daniel M. Hooper, Caitlyn D. Buchanan, Ulf S. Johansson, D. Thomas Tietze, Per Alström, Urban Olsson, Mousumi Ghosh-Harihar, Farah Ishtiaq, Sandeep K. Gupta, Jochen Martens, Bettina Harr, Pratap Singh, Dhananjai Mohan. 2014. Niche filling slows the diversification of Himalayan songbirds. Nature 509: 222-225.

194.   Prum, Richard O., Jacob S. Berv, Alex Dornburg, Daniel J. Field, Jeffrey P. Townsend, Emily Moriarty
Lemmon, Alan R. Lemmon. 2015. A comprehensive phylogeny of birds (Aves) using targeted next-
generation DNA sequencing. Nature 526, (7574): 569-573

195.   Pyron, R. Alexander, Frank T. Burbrink. 2013. Early origin of viviparity and multiple reversions to
oviparity in squamate reptiles. Ecology Letters 17 (1): 13-21

196.   Qing, X., Zhang, Y. M., Sun, S., Ahmed, M., Lo, W.-S., Bert, W., Holovachov, O., & Li, H. (2024).
Phylogenomic Insights into the Evolution and Origin of Nematoda. Systematic Biology, 74(3), 349–358.
https://doi.org/10.1093/sysbio/syae073

197.   Righetti, N., Nicolini, F., Forni, G., & Luchetti, A. (2025). Towards a time-tree solution for
Branchiopoda diversification: a jackknife assessment of fossil age priors. Palaeoentomology, 8(3), 316–
328. https://doi.org/10.11646/palaeoentomology.8.3.8

1685

198.   Rainford, J. L., Hofreiter, M., Nicholson, D. B., & Mayhew, P. J. (2014). Phylogenetic Distribution of
Extant Richness Suggests Metamorphosis Is a Key Innovation Driving Diversification in Insects. PLoS ONE,
9(10), e109085. https://doi.org/10.1371/journal.pone.0109085

199.   Ralph S. Peters, Lars Krogmann, Christoph Mayer, Alexander Donath, Simon Gunkel, Karen
Meusemann, Alexey Kozlov, Lars Podsiadlowski, Malte Petersen, Robert Lanfear, Patricia A. Diez, John
Heraty, Karl M. Kjer, Seraina Klopfstein, Rudolf Meier, Carlo Polidori, Thomas Schmitt, Shanlin Liu, Xin Zhou,
Torsten Wappler, Jes Rust, Bernhard Misof, Oliver Niehuis, 2017, 'Evolutionary History of the Hymenoptera',
Current Biology

200.   Rapini, A. [et al. 2007], van den Berg, C., & Liede-Schumann, S. 2007.

201.   Diversification of Asclepiadoideae (Apocynaceae) in the New World.

202.   Ann. Missouri Bot. Gard. 94: 407-422.

203.   dos Reis, Mario, Yuttapong Thawornwattana, Konstantinos Angelis, Maximilian J. Telford, Philip C.J.
Donoghue, Ziheng Yang. 2015. Uncertainty in the timing of origin of animals and the limits of precision in
molecular timescales. Current Biology

204.   Ribas, C. C., A. Aleixo, A. C. R. Nogueira, C. Y. Miyaki, J. Cracraft, 2011. A palaeobiogeographic
model for biotic diversification within Amazonia over the past three million years. Proceedings of the Royal
Society B: Biological Sciences 279 (1729): 681-689.

205.   Ronquist, Fredrik, Seraina Klopfstein, Lars Vilhelmsen, Susanne Schulmeister, Debra L. Murray,
Alexandr P. Rasnitsyn. 2012. A Total-evidence approach to dating with fossils, applied to the early radiation
of the Hymenoptera. Systematic Biology, 61(6): 973-999.

206.   Roquet, C., Lavergne, S., & Thuiller, W. (2014). One Tree to Link Them All: A Phylogenetic Dataset
for   the   European   Tetrapoda.   PLoS   Currents.
doi:10.1371/currents.tol.5102670fff8aa5c918e78f5592790e48

207.   de Sá R.O., Streicher J.W., Selenkoyela R., Forlani M.C., Loader S.P., Greenbaum E., Richards S., &
Haddad C. 2012. Molecular phylogeny of microhylid frogs (Anura: Microhylidae) with emphasis on
relationships among New World genera. BMC Evolutionary Biology 12: 241 .

208.   Sangeet Lamichhaney, Jonas Berglund, Markus Sällman Almén, Khurram Maqbool, Manfred Grabherr, Alvaro Martinez-Barrio, Marta Promerová, Carl-Johan Rubin, Chao Wang, Neda Zamani, B. Rosemary Grant, Peter R. Grant, Matthew T. Webster, Leif Andersson, 2015, 'Evolution of Darwin's finches and their beaks revealed by genome sequencing', Nature, vol. 518, no. 7539, pp. 371-375

209.   Sann, M., Niehuis, O., Peters, R. S., Mayer, C., Kozlov, A., Podsiadlowski, L., Bank, S., Meusemann, K., Misof, B., Bleidorn, C., & Ohl, M. (2018). Phylogenomic analysis of Apoidea sheds new light on the sister group of bees. BMC Evolutionary Biology, 18(1). https://doi.org/10.1186/s12862-018-1155-8

210.   Särkinen, Tiina , Lynn Bohs, Richard G. Olmstead, Sandra Knapp. 2013. A phylogenetic framework for evolutionary study of the nightshades (Solanaceae): a dated 1000-tip tree. BMC Evolutionary Biology 13: 214.

211.   Satoshi Eguchi, Minoru N. Tamura, 2016, 'Evolutionary timescale of monocots determined by the fossilized birth-death model using a large number of fossil records', Evolution, vol. 70, no. 5, pp. 1136-1144

212.   Schenk, John J., Kevin C. Rowe, and Scott J. Steppan. 2013. Ecological opportunity and incumbency in the diversification of repeated continental colonizations by muroid rodents. Systematic Biology 62 (6): 837-864.

213.   Schield, D. R., Brown, C. E., Shakya, S. B., Calabrese, G. M., Safran, R. J., & Sheldon, F. H. (2024). Phylogeny and historical biogeography of the swallow family (Hirundinidae) inferred from comparisons of thousands of UCE loci. Molecular Phylogenetics and Evolution, 197, 108111. https://doi.org/10.1016/j.ympev.2024.108111

214.   A. C. Schneider, W. A. Freyman, C. M. Guilliams, Y. P. Springer, B. G. Baldwin, 2016, 'Pleistocene radiation of the serpentine-adapted genus Hesperolinon and other divergence times in Linaceae (Malpighiales)', American Journal of Botany, vol. 103, no. 2, pp. 221-232

215.   Schweizer, Manuel, Hadoram Shirihai. 2013. Phylogeny of the Oenanthe lugens complex (Aves, Muscicapidae: Saxicolinae): paraphyly of a morphologically cohesive group within a recent radiation of open-habitat chats. Molecular Phylogenetics and Evolution 69 (3): 450-461.

216.   Schweizer, Manuel, Timothy F. Wright, Joshua V. Peñalba, Erin E. Schirtzinger, Leo Joseph. 2015. Molecular phylogenetics suggests a New Guinean origin and frequent episodes of founder-event speciation in the nectarivorous lories and lorikeets (Aves: Psittaciformes). Molecular Phylogenetics and Evolution 90: 34-48.

217.   Scofield, R. Paul, Kieren J. Mitchell, Jamie R. Wood, Vanesa L. De Pietri, Scott Jarvie, Bastien Llamas, Alan Cooper. 2016. The Origin and Phylogenetic Relationships of the New Zealand Ravens. Molecular Phylogenetics and Evolution 106: 136–143

218.   Selvatti, Alexandre Pedro, Ana Galvão, Anieli Guirro Pereira, Luiz Pedreira Gonzaga, Claudia Augusta de Moraes Russo. 2016. An African origin of the Eurylaimides (Passeriformes) and the successful diversification of the ground-foraging pittas (Pittidae). Molecular Biology and Evolution, p. msw250

219.   Selvatti, Alexandre Pedro, Luiz Pedreira Gonzaga, Claudia Augusta de Moraes Russo. 2015. A Paleogene origin for crown passerines and the diversification of the Oscines in the New World. Molecular Phylogenetics and Evolution 88: 1-15.

220. Seán G Brady, Brian L Fisher, Ted R Schultz, Philip S Ward, 2014, 'The rise of army ants and their relatives: diversification of specialized predatory doryline ants', BMC Evolutionary Biology, vol. 14, no. 1, p. 93

221. Shang-Yin Vanson Liu, Bruno Frédérich, Sébastien Lavoué, Jonathan Chang, Mark V. Erdmann, Gusti Ngurah Mahardika, Paul H. Barber, 2018, 'Buccal venom gland associates with increased of diversification rate in the fang blenny fish Meiacanthus (Blenniidae; Teleostei)', Molecular Phylogenetics and Evolution, vol. 125, pp. 138-146

222. S. V. Shedko, I. L. Miroshnichenko, G. A. Nemkova, 2012, 'Phylogeny of salmonids (Salmoniformes: Salmonidae) and its molecular dating: Analysis of nuclear RAG1 gene', Russian Journal of Genetics, vol. 48, no. 5, pp. 575-579

223. Shen, X.-X., Steenwyk, J. L., LaBella, A. L., Opulente, D. A., Zhou, X., Kominek, J., Li, Y., Groenewald, M., Hittinger, C. T., & Rokas, A. (2020). Genome-scale phylogeny and contrasting modes of genome evolution in the fungal phylum Ascomycota. Science Advances, 6(45). https://doi.org/10.1126/sciadv.abd0079

224. Shi, Jeff J., Daniel L. Rabosky. 2015. Speciation dynamics during the global radiation of extant bats. Evolution 69 (6): 1528-1545

225. Shin, S., Clarke, D. J., Lemmon, A. R., Moriarty Lemmon, E., Aitken, A. L., Haddad, S., Farrell, B. D., Marvaldi, A. E., Oberprieler, R. G., & McKenna, D. D. (2017). Phylogenomic Data Yield New and Robust Insights into the Phylogeny and Evolution of Weevils. Molecular Biology and Evolution, 35(4), 823–836. https://doi.org/10.1093/molbev/msx324

226. Shipham, Ashlee, Daniel J. Schmidt, Leo Joseph, Jane M. Hughes. 2015. Phylogenetic analysis of the Australian rosella parrots (Platycercus) reveals discordance among molecules and plumage. Molecular Phylogenetics and Evolution 91: 150-159.

227. Sikes, D.S., & Venables C. 2013. Molecular phylogeny of the burying beetles (Coleoptera: Silphidae: Nicrophorinae). Molecular Phylogenetics and Evolution 69 (3): 552-565.

228. Skojec, C., Earl, C., Couch, C.D., Masonick, P.K. & Kawahara, A.Y. (2023). Phylogeny and divergence time estimation of Io moths and relatives (Lepidoptera: Saturniidae: Automeris).

229. Smith, B. T., Merwin, J., Provost, K. L., Thom, G., Brumfield, R. T., Ferreira, M., Mauck, W. M., Moyle, R. G., Wright, T. F., & Joseph, L. (2022). Phylogenomic Analysis of the Parrots of the World Distinguishes Artifactual from Biological Sources of Gene Tree Discordance. Systematic Biology. https://doi.org/10.1093/sysbio/syac055

230. Smith, SA. 2009. Taking into account phylogenetic and divergence-time uncertainty in a parametric biogeographical analysis of the Northern Hemisphere plant clade Caprifolieae. Journal of Biogeography.

231. Soares, André E. R. , Ben J. Novak, James Haile, Tim H. Heupink, Jon Fjeldså, M. Thomas P. Gilbert, Hendrik Poinar, George M. Church, Beth Shapiro. 2016. Complete mitochondrial genomes of living and extinct pigeons revise the timing of the columbiform radiation. BMC Evolutionary Biology 16: 230

232. Soares, P., Abrantes, D., Rito, T., Thomson, N., Radivojac, P., Li, B., Macaulay, V., Samuels, D. C., & Pereira, L. (2013). Evaluating Purifying Selection in the Mitochondrial DNA of Various Mammalian Species. PLoS ONE, 8(3), e58993. https://doi.org/10.1371/journal.pone.0058993

233.   L. Sorenson, F. Santini, M. E. Alfaro, 2014, 'The effect of habitat on modern shark diversification',
Journal of Evolutionary Biology, vol. 27, no. 8, pp. 1536-1548

234.   Springer, Mark S., Robert W. Meredith, John Gatesy, Christopher A. Emerling, Jong Park, Daniel L.
Rabosky, Tanja Stadler, Cynthia Steiner, Oliver A. Ryder, Jan E. Janečka, Colleen A. Fisher, William J. Murphy.
2012. Macroevolutionary dynamics and historical biogeography of primate diversification inferred from a
species supermatrix. PLoS ONE 7 (11): e49521.

235.   Steeman, M., Hebsgaard M., Fordyce R., Ho S., Rabosky D., Nielsen R., Rahbek C., Glenner H.,
Sørensen M., & Willerslev E. 2009. Radiation of Extant Cetaceans Driven by Restructuring of the Oceans.
Systematic Biology 58 (6): 573-585.

236.   Stein, R. W., Mull, C. G., Kuhn, T. S., Aschliman, N. C., Davidson, L. N. K., Joy, J. B., Smith, G. J.,
Dulvy, N. K., & Mooers, A. O. (2018). Global priorities for conserving the evolutionary history of sharks, rays
and chimaeras. Nature Ecology & Evolution, 2(2), 288–298. https://doi.org/10.1038/s41559-017-
0448-4

237.   Stein, R. Will, Joseph W. Brown, Arne Ø. Mooers. 2015. A molecular genetic time scale
demonstrates Cretaceous origins and multiple diversification rate shifts within the order Galliformes
(Aves). Molecular Phylogenetics and Evolution 92: 155-164

238.   Steppan, Scott J. , John J. Schenk. 2017. Muroid rodent phylogenetics: 900-species tree reveals
increasing diversification rates. PLOS ONE 12 (8): e0183070

239.   Steven Poe, Adrián Nieto-montes de oca, Omar Torres-carvajal, Kevin De Queiroz, Julián A.
Velasco, Brad Truett, Levi N. Gray, Mason J. Ryan, Gunther Köhler, Fernando Ayala-varela, Ian Latella, 2017,
'A Phylogenetic, Biogeographic, and Taxonomic study of all Extant Species of Anolis (Squamata; Iguanidae)',
Systematic Biology, vol. 66, no. 5, pp. 663-697

240.   Stiller, J., Feng, S., Chowdhury, AA. et al. Complexity of avian evolution revealed by family-level
genomes. Nature (2024). https://doi.org/10.1038/s41586-024-07323-1

241.   Stiller, J., Short, G., Hamilton, H., Saarman, N., Longo, S., Wainwright, P., Rouse, G. W., & Simison,
W. B. (2022). Phylogenomic analysis of Syngnathidae reveals novel relationships, origins of endemic
diversity and variable diversification rates. BMC Biology, 20(1). https://doi.org/10.1186/s12915-022-01271-
w

242.   Subramanian, S., G. Beans-Picon, S. K. Swaminathan, C. D. Millar, D. M. Lambert. 2013. Evidence
for a recent origin of penguins. Biology Letters 9 (6): 20130748-20130748.

243.   Suvorov, A., Kim, B. Y., Wang, J., Armstrong, E. E., Peede, D., D'Agostino, E. R. R., Price, D. K.,
Waddell, P. J., Lang, M., Courtier-Orgogozo, V., David, J. R., Petrov, D., Matute, D. R., Schrider, D. R., &
Comeault, A. A. (2022). Widespread introgression across a phylogeny of 155 Drosophila genomes. Current
Biology, 32(1), 111-123.e5. https://doi.org/10.1016/j.cub.2021.10.052

244.   Sweet, Andrew D., Kevin P. Johnson. 2015. Patterns of diversification in small New World ground
doves are consistent with major geologic events. The Auk 132 (1): 300-312.

245.   A. E. Syme, T. H. Oakley, 2011, 'Dispersal between Shallow and Abyssal Seas and Evolutionary Loss
and Regain of Compound Eyes in Cylindroleberidid Ostracods: Conflicting Conclusions from Different
Comparative Methods', Systematic Biology, vol. 61, no. 2, pp. 314-336

246.   Testo, Weston, Michael Sundue. 2016. A 4000-species dataset provides new insight into the
evolution of ferns. Molecular Phylogenetics and Evolution 105: 200-211

247.   Thomson, R. C., Spinks, P. Q., & Shaffer, H. B. (2021). A global phylogeny of turtles reveals a burst
of climate-associated diversification on continental margins. Proceedings of the National Academy of
Sciences, 118(7). https://doi.org/10.1073/pnas.2012215118

248.   Thornhill, Andrew H., Simon Y.W. Ho, Carsten Külheim, Michael D. Crisp. 2015. Interpreting the
modern distribution of Myrtaceae using a dated molecular phylogeny. Molecular Phylogenetics and
Evolution 93: 29-43

249.   Tierney, S. M., Sanjur, O., Grajales, G. G., Santos, L. M., Bermingham, E., & Wcislo, W. T. (2011).
Photic niche invasions: phylogenetic history of the dim-light foraging augochlorine bees (Halictidae).
Proceedings   of   the   Royal   Society   B: Biological   Sciences,   279(1729),   794–803.
https://doi.org/10.1098/rspb.2011.1355

250.   Toljagić O., Voje K.L., Matschiner M., Liow L., & Hansen T.F. 2017. Millions of Years Behind:
Slow Adaptation of Ruminants to Grasslands. Systematic Biology, .

251.   Torres, Chris R, Lisa M Ogawa, Mark AF Gillingham, Brittney Ferrari, Marcel van Tuinen. 2014. A
multi-locus inference of the evolutionary diversification of extant flamingos (Phoenicopteridae). BMC
Evolutionary Biology 14 (1): 36.

252.   Toussaint, Emmanuel F. A., Lars Hendrich, Helena Shaverdo, Michael Balke. 2015. Mosaic patterns
of diversification dynamics following the colonization of Melanesian islands. Scientific Reports 5: 16016

253.   Emmanuel F. A. Toussaint, Matthias Seidel, Emmanuel Arriaga-Varela, Jiří Hájek, David Král, Lukaš
Sekerka, Andrew E. Z. Short, Martin Fikáček, 2016, 'The peril of dating beetles', Systematic Entomology

254.   Tárcia Fernanda da Silva, Horacio Schneider, Iracilda Sampaio, Arturo Angulo, Marcelo Fulgêncio
Guedes Brito, Alexandre Clistenes de Alcântara Santos, Jonas de Andrade Santos, Alfredo Carvalho-Filho,
Simoni Santos, 2018, 'Phylogeny of the subfamily Stelliferinae suggests speciation in Ophioscion Gill,
1863 (Sciaenidae: Perciformes) in the western South Atlantic', Molecular Phylogenetics and Evolution, vol.
125, pp. 51-61

255.   Uit de Weerd, D. R., Robinson, D. G., & Rosenberg, G. (2016). Evolutionary and biogeographical
history of the land snail family Urocoptidae (Gastropoda: Pulmonata) across the Caribbean region. Journal
of Biogeography, 43(4), 763–777. doi:10.1111/jbi.12692

256.   Uit de weerd D.R., & Gittenberger E. 2013. Phylogeny of the land snail family Clausiliidae
(Gastropoda: Pulmonata). Molecular Phylogenetics and Evolution, 67(1): 201-216.

257.   Upham, N. S., Esselstyn, J. A., & Jetz, W. (2019). Inferring the mammal tree: Species-level sets of
phylogenies for questions in ecology, evolution, and conservation. PLOS Biology, 17(12), e3000494.
doi:10.1371/journal.pbio.3000494

258.   Upham, N.S. & B.D. Patterson. 2015. Evolution of the caviomorph rodents: a complete phylogeny
and timetree of living genera. Pp. 63–120 In: Biology of caviomorph rodents: diversity and evolution (A. I.
Vassallo & D. Antenucci, eds.). SAREM Series A, Buenos Aires.

259.   Uva, Vera , Martin Päckert, Alice Cibois, Luca Fumagalli, Alexandre Roulin. 2018. Comprehensive molecular phylogeny of barn owls and relatives (Family: Tytonidae), and their six major Pleistocene radiations. Molecular Phylogenetics and Evolution 125: 127-137.

260.   Uyeda, J. C., Pennell, M. W., Miller, E. T., Maia, R., & McClain, C. R. (2017). The Evolution of Energetic Scaling across the Vertebrate Tree of Life. The American Naturalist, 190(2), 185–199. doi:10.1086/692326

261.   Vaga, C. F., Quattrini, A. M., Galvão de Lossio e Seiblitz, I., Huang, D., Quek, Z. B. R., Stolarski, J., Cairns, S. D., & Kitahara, M. V. (2025). A global coral phylogeny reveals resilience and vulnerability through deep time. Nature. https://doi.org/10.1038/s41586-025-09615-6

262.   Valtueña, Francisco J., Chris D. Preston, and Joachim W. Kadereit. "Phylogeography of a Tertiary relict plant, Meconopsis cambrica (Papaveraceae), implies the existence of northern refugia for a temperate herb." Molecular Ecology 21.6 (2012): 1423-1437.

263.   Vasconcelos, T. N. C., Alcantara, S., Andrino, C. O., Forest, F., Reginato, M., Simon, M. F., & Pirani, J. R. (2020). Fast diversification through a mosaic of evolutionary histories characterizes the endemic flora of ancient Neotropical mountains. Proceedings of the Royal Society B: Biological Sciences, 287(1923), 20192933. doi:10.1098/rspb.2019.2933

264.   Vázquez-López, M., Ramírez-Barrera, S. M., Terrones-Ramírez, A. K., Robles-Bello, S. M., Nieto-Montes de Oca, A., Ruegg, K., & Hernández-Baños, B. E. (2024). Biogeographic factors contributing to the diversification of Euphoniinae (Aves, Passeriformes, Fringillidae): a phylogenetic and ancestral areas analysis. ZooKeys, 1188, 169–195. https://doi.org/10.3897/zookeys.1188.107047

265.   Vea, Isabelle M., David A. Grimaldi. 2016. Putting scales into evolutionary time: the divergence of major scale insect lineages (Hemiptera) predates the radiation of modern angiosperm hosts. Scientific Reports, 6: 23487 .

266.   Vianna, J. A., Fernandes, F. A. N., Frugone, M. J., Figueiró, H. V., Pertierra, L. R., Noll, D., ... Bowie, R. C. K. (2020). Genome-wide analyses reveal drivers of penguin diversification. Proceedings of the National Academy of Sciences, 117(36), 22303–22310. doi:10.1073/pnas.2006659117

267.   Wang, Ning, Rebecca T. Kimball, Edward L. Braun, Bin Liang, Zhengwang Zhang. 2016. Ancestral range reconstruction of Galliformes: the effects of topology and taxon sampling. Journal of Biogeography

268.   Wang, S., & Luo, H. (2025). Dating the Bacterial Tree of Life Based on Ancient Symbiosis. Systematic Biology, 74(4), 639–655. https://doi.org/10.1093/sysbio/syae071

269.   Wells, H. B. M., Crego, R. D., Opedal, Ø. H., Khasoha, L. M., Alston, J. M., Reed, C. G., Weiner, S., Kurukura, S., Hassan, A. A., Namoni, M., Ekadeli, J., Kimuyu, D. M., Young, T. P., Kartzinel, T. R., Palmer, T. M., Pringle, R. M., & Goheen, J. R. (2021). Experimental evidence that effects of megaherbivores on mesoherbivore space use are influenced by species' traits. Journal of Animal Ecology, 90(11), 2510–2522. Portico. https://doi.org/10.1111/1365-2656.13565

270.   Weston Testo, Benjamin Øllgaard, Ashley Field, Thaís Almeida, Michael Kessler, David Barrington, 2018, 'Phylogenetic systematics, morphological evolution, and natural groups in neotropical Phlegmariurus (Lycopodiaceae)', Molecular Phylogenetics and Evolution, vol. 125, pp. 1-13

271.   Wiemers, M., Chazot, N., Wheat, C., Schweiger, O., & Wahlberg, N. (2020). A complete time-
calibrated multi-gene phylogeny of the European butterflies. ZooKeys, 938, 97–124.
doi:10.3897/zookeys.938.50878

272.   Wildman, Derek E., et al. "Implications of natural selection in shaping 99.4% nonsynonymous DNA
identity between humans and chimpanzees: enlarging genus Homo." Proceedings of the national Academy
of Sciences 100.12 (2003): 7181-7188.

273.   Wright, April M., Kathleen M. Lyons, Matthew C. Brandley, David M. Hillis. 2015. Which came first:
The lizard or the egg? Robustness in phylogenetic reconstruction of ancestral states. Journal of
Experimental Zoology Part B: Molecular and Developmental Evolution 324 (6): 504-516

274.   Yao, X., Song, Y., Yang, J., Tan, Y., & Corlett, R. T. (2020). Phylogeny and biogeography of the hollies
(Ilex   L., Aquifoliaceae). Journal of Systematics and Evolution. doi:10.1111/jse.12567

275.   Yu, W., Wu, Y., & Yang, G. (2014). Early diversification trend and <scp>A</scp>sian origin for extent
bat   lineages.   Journal   of   Evolutionary   Biology,   27(10),   2204–2218.   Portico.
https://doi.org/10.1111/jeb.12477

276.   Yuchi Zheng, John J. Wiens, 2016, 'Combining phylogenomic and supermatrix approaches, and a
time-calibrated phylogeny for squamate reptiles (lizards and snakes) based on 52 genes and 4162 species',
Molecular Phylogenetics and Evolution, vol. 94, pp. 537-547

277.   Zhang, S.-Q., Che, L.-H., Li, Y., Dan Liang, Pang, H., Ślipiński, A., & Zhang, P. (2018). Evolutionary
history of Coleoptera revealed by extensive sampling of genes and species. Nature Communications, 9(1).
https://doi.org/10.1038/s41467-017-02644-4

278.   Zhang, Y. M., Delvare, G., Blaimer, B. B., Cruaud, A., Rasplus, J., Brady, S. G., & Gates, M. W. (2025).
Phasing in and out of phytophagy: Phylogeny and evolution of the family Eurytomidae (Hymenoptera:
Chalcidoidea)   based   on   Ultraconserved   Elements.   Systematic   Entomology.   Portico.
https://doi.org/10.1111/syen.12682

279.   Zhao, M., Thom, G., Faircloth, B. C., Andersen, M. J., Barker, F. K., Benz, B. W., Braun, M. J., Bravo,
G. A., Brumfield, R. T., Chesser, R. T., Derryberry, E. P., Glenn, T. C., Harvey, M. G., Hosner, P. A., Imfeld, T.
S., Joseph, L., Manthey, J. D., McCormack, J. E., McCullough, J. M., ... Smith, B. T. (2025). Efficient Inference
of   Macrophylogenies: Insights   from   the   Avian   Tree   of   Life.   Systematic   Biology.
https://doi.org/10.1093/sysbio/syaf080

280.   Zuntini, A.R., Carruthers, T., Maurin, O. et al. Phylogenomics and the rise of the angiosperms.
Nature (2024). https://doi.org/10.1038/s41586-024-07324-0