Plasma membrane nanoscale dynamics of *Arabidopsis* leucine-rich repeat receptor
kinase complexes

Michelle von Arx[1,2], Marie-Dominique Jolivet[1,2], David Biermann[1,2], Vaishali Gabani[1], Steven
S. Andrews[3], Cyril Zipfel[4,5], Julien Gronnier[1,2,4#]

Correspondence: julien.gronnier@tum.de

1. Plant Cell Biology, TUM School of Life Sciences, Technical University of Munich, Freising, Germany.
2. Center for Plant Molecular Biology (ZMBP), University of Tübingen, Germany.
3. Department of Bioengineering, University of Washington, United states of America.
4. Institute of Plant and Microbial Biology and Zurich-Basel Plant Science Center, University of Zürich,
Zürich, Switzerland.
5. The Sainsbury Laboratory, University of East Anglia, Norwich Research Park, Norwich, United
Kingdom.

Abstract
Plasma membrane-localized receptors operate as dynamic signaling complexes and
integrative networks[1-3], yet the spatial and temporal regulation of these interactions remain
largely unknown. Here, by analyzing the components of a minimal *Arabidopsis* leucine-rich
repeat receptor kinase network, we describe the differential diffusion and organization of
receptor complex components and unveil the nanoscale spatial and temporal logic
underlying the formation of receptor kinase complexes. The ligand-binding receptors FLS2
and BRI1, and the accessory receptor BIR3, are organized in plasma membrane
nanodomains, within which the co-receptor BAK1 diffuses and is spatially arrested upon
ligand perception. BAK1's spatial arrest relies on extracellular domain (ECD)-ECD
interactions but does not require receptor complex activation. Mathematical modelling,
single molecule imaging and bio-assays infer that accessory receptors maintain a dynamic
pool of co-receptors in the vicinity of ligand-binding receptors to promote ligand-induced
complex formation and signaling. We propose that ligand-induced receptor kinase complex
formation is a deterministic process defined by the relative nanoscale spatial positioning of
individual signaling and regulatory components.

Main
Plants and metazoan cell-surface receptors sense and relay a myriad of extracellular signals to
coordinate development and adjust to ever-changing environmental conditions[4,5]. In plants, the
repertoire of cell-surface receptor kinases has considerably expanded with over 400 members
encoded in the genome of *Arabidopsis thaliana* (hereafter Arabidopsis)[6]. Among them, leucine-
rich repeat receptor kinases (LRR-RKs) constitute a prominent class that perceive self, non-self

and modified-self molecules to regulate development, immunity and reproduction[7-9]. For

instance, the ligand-binding LRR-RK BRASSINOSTEROID-INSENSITIVE 1 (BRI1) functions as the

main receptor for plant steroid hormones brassinosteroids[7] and forms a complex with the co-

receptor BRI1-ASSOCIATED KINASE (BAK1, also called SOMATIC EMBRYOGENESIS RECEPTOR

KINASE 3, SERK3) to regulate growth[7,10]. BAK1 also serves as a co-receptor for other LRR-RKs, such

as FLAGELING SENSING 2 (FLS2) that sense the flg22 epitope derived from bacterial flagellin to

initiate immune signaling[9,11].

LRR-RKs have been proposed to function as a coordinated interaction network to process

extracellular signals[2]. In this framework, accessory receptor kinases regulate ligand-induced

association between ligand-binding receptors and their co-receptors. For instance, the LRR-RK

BAK1-INTERACTING RECEPTOR-LIKE KINASE 3 (BIR3) physically associates with BAK1[12,13] and has

been proposed to inhibit ligand-induced FLS2-BAK1 and BRI1-BAK1 complex formation[13-15].

Several studies have highlighted the spatial organization of LRR-RKs into plasma membrane

nanodomains[16-22]. For instance, the spatial separation of FLS2 and BRI1 has been proposed to

support signaling specificity[16] and alterations of the nanoscale organization of FLS2 and BAK1

correlate with defects in ligand-induced complex formation and signaling[17,23]. Yet, how the

dynamic associations of cell surface receptor are coordinated in space and time within the plasma

membrane remains unclear. Here, we studied a minimal LRR-RK network composed of the ligand-

binding receptors FLS2 and BRI1, their common co-receptor BAK1 and the accessory receptor

BIR3.

We analyzed the plasma membrane nanoscale organization of FLS2-GFP[24], BRI1-GFP[25] and BAK1-

GFP[26] by combining variable-angle total internal reflection microscopy (VA-TIRFM)[27,28] with

enhanced super-resolution radial fluctuations (eSRRF)[29] in hypocotyl epidermal cells of light

grown Arabidopsis seedlings. Consistent with previous observations made in Arabidopsis

cotyledons epidermal cells[16-18,30,31], we observed that FLS2-GFP and BRI1-GFP are organized into

well-defined nanodomains (Fig. 1a-b). In contrast, BAK1-GFP showed a more dispersed

distribution within the PM (Fig. 1a-b). To complement these observations, we used photochromic

reversion, a variant of single-particle tracking photoactivated localization microscopy (spt-PALM)

that enables long-term single molecule imaging[32]. We analyzed BRI1, FLS2 and BAK1 tagged with

the photoconvertible fluorescence protein mEOS3.2[33] expressed under the control of their native

promoter. We observed that FLS2-mEOS3.2 and BRI1-mEOS3.2 are overall static while BAK1 is

laterally diffusive and presents a higher diffusion coefficient (Fig. 1c-d; Extended Data Fig. 1a-b).

We observed that when transiently expressed in *Nicotiana benthamiana,* or expressed under the

control of the *ubiquitin 10* promoter[34] in Arabidopsis stable transgenic lines, FLS2-mEOS3.2

remained static and BAK1 remained diffusive (Fig. 1c-d; Extended Data Fig. 1a-d, 2a, 3c). We next

analyzed FLS2, BRI1 and BAK1 organization using nanoscale spatiotemporal indexing clustering

(NASTIC)[35]. We observed that FLS2 and BRI1 are more clustered than BAK1 (Fig. 1e-f). Together

these observations highlight the differential organization and diffusion of plant LRR-RK complex

components, features that contrast with the behavior of animal cell-surface receptors[36-39]. We

next investigated the effect of ligand perception on receptor dynamics. We imaged FLS2 and BRI1

within 10 min post ligand-treatment, a time-frame which corresponds to initiation of ligand-

induced complex formation and signaling[11,40,41]. In good agreement with the analysis of FLS2-GFP

diffusion in *N. benthamiana*[42], we observed that flg22 does not lead to changes in FLS2-mEOS3.2

diffusion in *N. benthamiana* and in Arabidopsis (Fig. 2c-d; Extended Data Fig. 2a-b). Further, flg22

treatment did change FLS2 clustering (Extended Data Fig. 2c-d). Similarly, we observed that BRI1-

mEOS3.2 nanoscale dynamics and clustering is unaffected by epi-brassinolide (eBL) treatment

(Fig. 2a-b; Extended Data Fig. 2e-f). To corroborate these observations, we used computational

analysis of spatial arrests (CASTA)[32] to analyze changes in FLS2 and BRI1 dynamics in long-term

single molecule imaging. We observed that flg22 and eBL do not lead to changes in FLS2 and BRI1

spatial arrest (Extended Data Fig. 2g-j). We conclude that ligand-induced complex formation and

early ligand-induced signaling is not associated with changes in ligand-binding receptor nanoscale

dynamics or organization.

We next examined the behavior of BAK1. We observed that BAK1-mEOS3.2 diffusion coefficient

is reduced upon eBL treatment (Fig. 2e-f). Further, we observed an increase in BAK1-mEOS3.2

clustering (Extended Data Fig. 3a-b) and in the frequency of its spatial arrest (Fig. 2i-j) events upon

eBL treatment. Similarly, we observed that flg22 treatment reduces BAK1-mEOS3.2 diffusion in

N. benthamiana (Extended Data Fig. 3c-d). However, in Arabidopsis hypocotyl cells, we did not

observe changes in BAK1 mobility behavior (Extended Data Fig. 3e-f). We presumed that the low

FLS2 expression in Arabidopsis hypocotyl cells (Extended Data Fig. 1a) combined with the inherent

stochasticity of BAK1-mEOS3.2 photoconversion impedes our ability to detect changes in BAK1

mobility behavior. To test this hypothesis, we crossed *bak1-4*/pBAK1::BAK1-mEOS3.2 with Col-

0/pFLS2::FLS2-GFP to increase FLS2 accumulation. We observed that introducing FLS2-GFP in

bak1-4/pBAK1::BAK1-mEOS3.2 leads to detectable flg22-induced decrease in BAK1-mEOS3.2

diffusion (Fig. 2g-h) and increase in BAK1 clustering (Extended Data Fig. 3g-h). Further, we

observed that similar to eBL, flg22 treatments lead to an increase in BAK1-mEOS3.2 spatial arrests

(Fig. 2i-l). Together these observations show that, while ligand binding RKs are static and

organized in clusters independently of exogenous ligand treatment, BAK1 clustering and spatial

arrestment increases upon ligand perception. Using dual-color VA-TIRFM we observed that FLS2-

GFP and BAK1-mCherry, as well as BRI1-GFP and BAK1-mCherry partially co-localize in control

conditions (Extended Data Fig. 4a-b and d-e). These observations are consistent with previous cell

surface-selective FLIM analyses of BRI1-GFP and BAK1-mCherry[20], and suggest that BAK1

navigates in the vicinity of ligand-binding receptor nanodomains. Further, we observed that loss

of the receptor kinase FERONIA (allele *fer*-4), which induces an increase in BRI1-BAK1

association[43], led to an increased co-localization between BRI1-GFP and BAK1-mCherry (Extended

Data Fig. 4f). Similarly, we observed that flg22 treatment leads to an increased co-localization
between FLS2-GFP and BAK1-mCherry (Extended Data Fig. 4c). In contrast to BRI1-BAK1, flg22-
induced FLS2-BAK1 complex formation was reduced in *fer-4*[44]. In agreement, we observed flg22
treatment does not lead to an increase in co-localization between FLS2-GFP and BAK1-mCherry
in *fer-4*. Together with single molecule tracking analyses, these observations suggest that BAK1
navigates within and is spatially arrested to pre-formed ligand-binding receptor nanodomains
upon ligand perception.

We next investigated the molecular events underlying ligand-induced BAK1 spatial arrestment.
We mutated two residues in the BAK1 ECD, Phenylalanine (F) 60 and F144, which are commonly
engaged in interaction with ligand-bound ligand-binding receptors, including BRI1, and are
required for ligand-induced complex formation[13,45-48]. We observed that the nanoscale dynamics
and organization of $BAK1^{F60A/F144A}$-mEOS3.2 is insensitive to eBL treatment (Fig. 2m; Extended
Data Fig. 5a-c), suggesting that ligand-induced BAK1 spatial arrestment relates to ligand-induced
complex formation. In agreement, using a BAK1 kinase-dead version of BAK1 ($BAK1^{D416N}$)[49,50], we
observed that BAK1 kinase activity is not required for eBL-induced BAK1 spatial arrestment and
clustering (Fig. 2n; Extended Data Fig. 5d-f). We conclude that ligand-induced BAK1 spatial
arrestment corresponds to ligand-induced complex formation and does not rely on receptor
complex activation and downstream signaling events.

We observed that a fraction of BAK1 is spatially arrested in absence of exogenous ligand
treatment (Fig. 2) and that mutations that abolish BAK1 association with a variety of ligand-
binding receptors (Fig. 2m) are not sufficient to abolish BAK1 clustering and spatial arrestment
(Extended Data Fig. 5g-i). We thus hypothesized that additional regulators control BAK1 spatial
arrestment. The accessory receptor kinase BIR3 physically associates with BAK1[13,14] and is in
proximity to FLS2 and BRI1[14]. Using VA-TIRFM-eSRRF we observed that, similar to FLS2 and BRI1
(Fig. 1), BIR3-GFP is organized into plasma membrane nanodomains (Extended data Fig. 6a).
Corroborating these observations, we observed that BIR3-mEOS3.2 is relatively static and forms
nanoclusters (Fig. 3a-d; Extended data Fig. 6b-c). Since BIR3 showed a slow and confined lateral
diffusion, we hypothesized that BAK1 transiently physically associates with BIR3 and conditions
BAK1 spatial arrestment. We therefore mutated BAK1 Aspartic acid (D) 122, which has been
shown to reduce BAK1-BIR3 ECD-ECD interactions *in vitro*[13]. We observed that substituting D122
with Alanine (A) led to an increase in BAK1 diffusion (Fig. 3e-f), as well as a reduction in BAK1
spatial arrestment (Fig. 3g-h) and clustering (Fig. 3i-j), indicating that BIR3 regulates BAK1
nanoscale dynamics. Corroborating these observations, we observed that loss of *BIR3* leads to an
increase in BAK1-mEOS3.2 diffusion and a decrease in BAK1 clustering (Extended Data Fig. 7a-e).
Loss of *BIR3* was not sufficient to significantly affect BAK1 spatial arrest (Extended Data Fig 7f-g),
suggesting the implication of additional factors; potentially the BIR3 paralogues BIR1 and/or
BIR2[51,52]. We conclude that the association of BAK1 to BIR3, and potentially other BIRs, contribute

to BAK1 clustering and spatial arrestment presumably in the vicinity of the ligand-binding
receptor kinases FLS2 and BRI1 (Fig. 3k).

We next set to reconstitute the plasma membrane nanoscale dynamics of ligand-induced
complex formation *in silico* using particle-based simulations (Fig. 4a)[53]. We used the diffusion
parameters extracted from single molecule imaging and known ECD-ECD binding kinetics[13,54] to
parametrize our model. These simulations closely recapitulated the nanoscale behavior of single
receptor molecules (Extended Data Fig. 8a-c). Interestingly, we observed that introducing BIR3 in
these simulations promotes the existence of a dynamic pool of BAK1 in the vicinity of FLS2
nanodomains (Fig. 4b-d) and flg22-induced complex formation (Fig. 4e), suggesting that BIR3 has
a positive role in regulating ligand-induced complex formation. In agreement, we observed that
loss of *BIR3* (*bir3-2* allele[14] and *bir3-c* CRISPR mutant (Extended Data Fig. 7)) led to a decrease in
flg22-induced FLS2-BAK1 complex formation in Arabidopsis seedlings (Fig. 4g; Extended Data
Figure 9a-c).

Corroborating these observations, loss of *BIR3* led to a decrease in flg22-induced MAPK
phosphorylation and flg22-induced ROS production (Fig. 4h; Extended Data Fig. 9d-e). These
results contrast with a previous report in Arabidopsis[14], but align with the effects of loss of BIRs
in wheat[55] and in *N. benthamiana*[56]. In addition, in line with previous observations in Arabidopsis
root and dark grown hypocotyls[14], and in rice[57], we observed that loss of *BIR3* led to reduced eBL-
responsiveness in light-grown Arabidopsis hypocotyls (Extended Data Fig. 9f, g). These results
together indicate that BIR3 functions as positive regulator of both flg22- and eBL-triggered
signaling. Altogether, these observations suggest that BIR3 attracts BAK1 in the vicinity of ligand-
binding receptors, providing a dynamic pool of BAK1 available for ligand-induced complex
formation with ligand-binding receptors. We next tested the effect of increasing the amount of
BIR3 *in silico*. These simulations predict a decrease in the number of BAK1 molecules in the
vicinity of FLS2 (Fig. 4c). Further, in agreement with the effect of BIR3 over-expression observed
in vivo[14], increasing the number of BIR3 molecules decreases FLS2-BAK1 complexes formed *in*
silico (Fig. 4f). In this system, BIR3 predominantly acted as a negative regulator keeping BAK1 away
from FLS2 nanodomains. Our data indicate a dual, concentration-dependent, role for the
accessory receptor kinase BIR3.

Discussion
177
How cell surface receptors dynamically assemble into signaling-competent complexes within the
plasma membrane remains a fundamental question. By combining super-resolution microscopy,
long-term single molecule imaging and mathematical modeling, our results suggest a nanoscale
organizational principle underlying the formation of Arabidopsis LRR-RK complexes. Ligand-
binding receptors are pre-organized into nanodomains, while the common co-receptor BAK1 is
diffusive and spatially arrested to ligand-binding receptors. The spatial and temporal

orchestration of plant LRR-RK and animal cell-surface receptor complexes thus differ. For instance,
compared to FLS2 or BRI1, the human epidermal growth factor receptor (EGFR) and type-I
interferon receptors are diffusive and undergo ligand-induced dimerization and spatial
stabilization[38,58]. Our experimental and *in silico* analyses converge in describing receptor kinases
as an organized and predictable system in which BIR3 maintains a dynamic pool of available BAK1
in the vicinity of ligand-binding receptors for ligand-induced complex formation and signaling (Fig.
4i). We thus propose that accessory receptors, such as BIR3, act as spatial regulators for co-
receptor availability. This reconciles previously conflicting genetic observations made in several
plant species[14,55-57] and highlights how subtle changes in accessory receptor abundance can tune
ligand responsiveness. Collectively, our findings suggest that ligand-induced complex formation
is a deterministic process defined by the relative spatial positioning of individual signaling and
regulatory components at the nanoscale rather than purely stochastic diffusion-driven
encounters. In this framework, signaling competence emerges from the pre-established spatial
architecture of the plasma membrane and may reflect a general organizational principle of
signaling systems.
199

Acknowledgments
We thank all members of the NanoSignaling Laboratory for fruitful discussions and
comments on the manuscript. We thank Niko Geldner for providing the Col-0/pBRI1::BRI1-
GFP, Sacco de Vries for providing the Col-0/pBRI1::BRI1-GFP/pBAK1::BAK1-mCherry and
Col-0/pBAK1::BAK1-mCherry lines, and Birgit Kemmerling for providing the *bir3-2* mutant.
Bio-imaging was performed at the ZMBP microscopy facility. We thank Sven zur Oven-
Krockhaus for maintaining the VA-TIRFM. We thank Birgit Kemmerling, Thorsten Nürnberger
and Kyle Bender for comments on the manuscript. This research was supported by the
Deutsche Forschungsgemeinschaft (DFG) grants (A09-SFB1101 and B01-TRR356) to J.G., the
European Molecular Biology Organization (postdoctoral fellowship EMBO LTF 438-2018, to
J.G.), the European Research Council under the European Union's Horizon 2020 research
and innovation programme grant agreement no. 773153 (project 'IMMUNO-PEPTALK' to
C.Z.), the University of Zürich (to C.Z.), the University of Tübingen (to J.G.) and the Technical
University of Munich (to J.G.).

Col-0/pFLS2::FLS2-GFP   Col-0/pBRI1::BRI1-GFP   *bak1-4*/pBAK1::BAK1-GFP

Localization density -   +

fls2/pUb::FLS2-mEOS3.2   Col-0/pBRI1::BRI1-mEOS3.2   *bak1-4*/pBAK1::BAK1-mEOS3.2

Molecule velocity -   +

fls2/pUb::FLS2-mEOS3.2   Col-0/pBRI1::BRI1-mEOS3.2   *bak1-4*/pBAK1::BAK1-mEOS3.2

Figures legends

Figure 1 | Differential organization and dynamics of LRR-RK complex components.

a, Representative eSRRF images of FLS2-GFP, BRI1-GFP and BAK1-GFP plasma membrane
organization in 5-day-old Arabidopsis hypocotyl. **b**, Quantification of the image wide spatial
clustering index (iSCI). Each data point represents the average value per cell (n), colors indicate
independent experiments. **c**, Representative images of FLS2-mEOS3.2, BRI1-mEOS3.2 and BAK1-
mEOS3.2 long-term single molecule trajectories in 5-day-old Arabidopsis hypocotyl. Trajectories
are colored based on velocity. **d**, Analysis of instantaneous diffusion coefficient (D). Each data
point represents the average log(D) value in $\mu m^2/s$ obtained per cell (n), colors indicate
independent experiments. **e**, Representative images of FLS2-mEOS3.2, BRI1-mEOS3.2 and BAK1-
mEOS3.2 trajectories nanoclustering in 5-day-old Arabidopsis hypocotyl. **f**, Quantification of
percentage of clustered trajectories. Each data point represents the percentage of clustered
trajectories per cell (n), colors indicate independent experiments. Conditions sharing a letter are
not significantly different in pairwise Wilcoxon with Bonferroni correction (p>0.05).

Col-0/pBRI1::BRI1-mEOS3.2

a Mock / eBL — Molecule velocity − +, 10 μm

b Log(D), a a, Mock eBL — 1 2

fls2/pUb::FLS2-mEOS3.2

c Mock / flg22 — 10 μm

d Log(D), a a, Mock flg22 — 1 2 3 4

bak1-4/pBAK1::BAK1-mEOS3.2

e Mock / eBL — Molecule velocity − +, 10 μm

f Log(D), a b, Mock eBL — 1 2 3 4 5 6 7

bak1-4/pBAK1::BAK1-mEOS3.2

g Mock / flg22 — 10 μm

h Log(D), a b, Mock flg22 — 1 2 3

bak1-4/pBAK1::BAK1-mEOS3.2

i Mock / eBL — Spatial arrest, 500 nm

j Spatial arrest, a b, Mock eBL — 1 2 3 4 5 6 7

bak1-4/pBAK1::BAK1-mEOS3.2

k Mock / flg22 — Spatial arrest, 500 nm

l Spatial arrest, a b, Mock flg22 — 1 2 3

m *bak1-4*/pBAK1::BAK1-mEOS3.2 — Log(D), a b / a a, BAK1^WT BAK1^F60A/F144A / Mock eBL, BAK1^F60A/F144A — 1 2 3 4

n *bak1-4*/pBAK1::BAK1-mEOS3.2 — a a / a b, BAK1^WT BAK1^D416N / Mock eBL, BAK1^D416N — 1 2 3 4

o BAK1 / Ligand-binding receptor, Ligand

Figure 2 | BAK1 is spatially arrested upon ligand perception.

a, Representative BRI1-mEOS3.2 trajectories, long-term single molecule imaging in presence of 1 µM eBL or corresponding mock control (EtOH) in Arabidopsis hypocotyl. **b**, Quantification of instantaneous diffusion coefficient (D) within 10 min of treatment. Each data point represents the average log(D) value obtained per cell (n), colors indicate independent experiments. **c**, Representative FLS2-mEOS3.2 trajectories, long-term single molecule imaging in presence of 1 µM flg22 or corresponding mock control (H_2O) in Arabidopsis hypocotyl. **d**, Quantification of instantaneous diffusion coefficient (D) within 10 min of treatment. Each data point represents the average log(D) value obtained per cell (n), colors indicate independent experiments. **e**, **g**, **i**, and **k**, Representative BAK1-mEOS3.2 trajectories, long-term single molecule imaging in presence of 1 µM eBL (**e** and **i**) or 1 µM flg22 (**g** and **k**) or the corresponding mock controls (EtOH or H_2O) in *bak1-4* (**e** and **i**) and *bak1-4*/pFLS2::FLS2-GFP (**g** and **k**) hypocotyls. **f**, **h**, **m** and **n**, Quantification of instantaneous diffusion coefficient (D) within 10 min of treatment. Each data point represents the average log(D) value obtained per cell (n), colors indicate independent experiments. **j** and **l**, Quantification of BAK1-mEOS3.2 spatial arrests within 10 min of treatment. Each data point represents the average number of spatial arrests per trajectory obtained per cell (n), colors indicate independent experiments. Conditions sharing a letter are not significantly different in pairwise Wilcoxon test with Bonferroni correction (p>0.05). **o**, Working model: BAK1 navigates within and is stabilized to pre-formed ligand-binding RK nanodomains upon ligand perception.

a *bak1-4*/pUb::BAK1-mEOS3.2   *bir3-2*/pUb::BIR3-mEOS3.2

Molecule velocity − ▬▬▬▬ +

b

Log(D)

a   b

	• 1
	• 2
	• 3

n=37   n=23

BAK1   BIR3

c *bak1-4*/pUb::BAK1-mEOS3.2   *bir3-2*/pUb::BIR3-mEOS3.2

Clusters

d

Clustered trajectories (%)

a   b

	• 1
	• 2
	• 3
	• 4

n=37   n=23

BAK1   BIR3

e *bak1-4*/pUb::BAK1-mEOS3.2   *bak1-4*/pUb::BAK1^D122A-mEOS3.2

f

Log(D)

a   b

	• 1
	• 2
	• 3

n=36   n=42

BAK1^WT   BAK1^D122A

g *bak1-4*/pUb::BAK1-mEOS3.2   *bak1-4*/pUb::BAK1^D122A-mEOS3.2

☐ Spatial arrest

h

Spatial arrest

a   b

	• 1
	• 2
	• 3

n=36   n=42

BAK1^WT   BAK1^D122A

i *bak1-4*/pUb::BAK1-mEOS3.2   *bak1-4*/pUb::BAK1^D122A-mEOS3.2

Clusters

j

Clustered trajectories (%)

a   b

	• 1
	• 2
	• 3

n=36   n=42

BAK1^WT   BAK1^D122A

k

BIR3
BAK1
Ligand-binding receptor

Figure 3 | BIRs regulates BAK1 plasma membrane nanoscale dynamics.
a and **e**, Representative BAK1-mEOS3.2 and BIR3-mESO3.2 (**a**) or BAK1-mEOS3.2 and BAK1^{D122A}-
mEOS3.2 (**e**) trajectories, long-term single molecule imaging, colors indicate velocity. **b** and **f**,
Quantification of instantaneous diffusion coefficient (D). Each data point represents the average
log(D) value obtained per cell (n), colors indicate independent experiments. **c** and **i**,
Representative images of BAK1-mEOS3.2. and BIR3-mEOS3.2 (**c**) or BAK1-mEOS3.2 and BAK1^{D122A}-
mEOS3.2 (**i**) trajectories nanoclustering. **d** and **j**, Quantification of clustered trajectories. Each data
point represents the percentage of clustered trajectories per cell (n), colors indicate independent
experiments. **g**, Representative images of BAK1-mEOS3.2 and BAK1^{D122A}-mEOS3.2 spatial arrest.
h, Quantification of BAK1-mEOS3.2 and BAK1^{D122A}-mEOS3.2 spatial arrests. Each data point
represents the average number of spatial arrests per trajectory obtained per cell (n), colors
indicate independent experiments. Conditions sharing a letter are not significantly different in
pairwise Wilcoxon with Bonferroni correction (p>0.05). **k**, Working model: pre-formed ligand-
binding RK nanodomains contain accessory RKs such as BIR3.
264

a Particle-based simulations

Molecules
FLS2
FLS2-flg22
BAK1
BIR3
BAK1-BIR3
FLS2-BAK1

Mesh ▽

2 µm

b BAK1

?

+/- BIR3
FLS2 nanodomain

c -BIR3 +BIR3 ++BIR3

Number of BAK1 in FLS2 nanodomains

Time (s)

d a   b

Time BAK1 remains in FLS2 nanodomains (s)

-BIR3   +BIR3   ++

e +BIR3
-BIR3

FLS2-flg22/BAK1 complexes

+ flg22

time (s)

f +BIR3
++BIR3

FLS2-flg22/BAK1 complexes

+ flg22

time (s)

g pBAK1::BAK1-mEOS3.2

bak1-4   bak1-4/bir3-c

-    +   -   +   flg22

170 kDa-   α-FLS2

100 kDa-   α-BAK1

IP α-FLS2

170 kDa-   α-FLS2

100 kDa-   α-BAK1

Input

h a   b

Total photon counts (×10⁶)

Col-0   fls2

i WT   bir3

BIR3
BAK1
Ligand-binding receptor

Ligand   Ligand

Figure 4 | BIR3 maintains a dynamic pool of BAK1 close to FLS2 for ligand-induced complex formation.

a, Snapshot of a particle-based simulation. **b,** Schematic depicting of the monitoring of BAK1 occurrence in FLS2 nanodomain in presence and absence of BIR3 in particle-based simulations. **c**, Number of BAK1 molecules present in FLS2 nanodomains over time. Data show mean ± se of 3 independent simulations. **d**, Duration of BAK1 occurrence in FLS2 nanodomains. Data points correspond to individual FLS2 nanodomain entry events, colors indicate independent simulations. **e-f,** Analysis of FLS2-flg22/BAK1 complex formation over time in presence of intermediate levels of BIR3 (+BIR3), no BIR3 (-BIR3) (**e**) or high levels of BIR3 (++BIR3) (**f**). The simulations were run 120 s prior to the addition of flg22 in the system. Data represent mean ± se of 10 independent simulations per scenario. **g**, Immunoprecipitation of FLS2 in *bak1-4*/pBAK1::BAK1-mEOS3.2 or *bir3-c*/*bak1-4*/pBAK1::BAK1-mEOS3.2 CRISPR line after treatment with 20 nM flg22 or corresponding mock solution (H_2O) for 2 min. Membranes were probed with anti-FLS2 or anti-BAK1. **h**, Reactive oxygen species production (ROS) in 4-week-old Arabidopsis leaf discs elicited with 100 nM flg22. Data points indicate total photon counts measured in individual leaf discs (n), colors indicate independent experiments. Conditions sharing a letter are not significantly different in pairwise Wilcoxon test with Bonferroni correction (p<0.05). **i**, Working model: In WT conditions BIR3 attracts and maintains a dynamic pool of BAK1 for efficient ligand-induced complex formation. In absence of accessory receptors, such as BIR3, BAK1 is not maintained in the vicinity of ligand-binding receptors, and the likelihood of ligand-induced complexes is reduced.

a *fls2*/pFLS2::FLS2-mEOS3.2   *fls2*/pUb::FLS2-mEOS3.2

10 µm

b *fls2*/FLS2-mEOS3.2

c *bak1-4*/pBAK1::BAK1-mEOS3.2   *bak1-4*/pUb::BAK1-mEOS3.2

10 µm

d *bak1-4*/BAK1-mEOS3.2

Extended Data Figure 1 | Analysis of FLS2 and BAK1 diffusion in Arabidopsis
a and **c**, Representative single molecule trajectories of FLS2-mEOS3-2 and BAK1-mEOS3.2, long-
term single molecule imaging in Arabidopsis hypocotyl epidermis, colors indicate velocity. **b** and
d, Quantification of instantaneous diffusion coefficient (D). Each data point represents the
average log(D) value obtained per cell (n). Conditions sharing a letter are not significantly different
in pairwise Wilcoxon test with Bonferroni correction (p<0.05).
292

Extended Data Figure 2 | Analysis of FLS2 and BRI1 diffusion and organization in presence and absence of exogenous ligand treatment.

a, Representative single molecule trajectories of FLS2-mEOS3.2 in presence of 1 μM flg22 or corresponding mock control (H$_2$O) in *N. benthamiana*. **b**, Quantification of instantaneous diffusion coefficient (D) within 10 min of treatment. Each data point represents the average log(D) value obtained per cell (n), colors indicate independent experiments. **c, e, g, i,** Representative images of FLS2-mEOS3.2 (**c, g**) or BRI1-mEOS3.2 (**e, i**) single molecule nanoclustering (**c, e**) or trajectories spatial arrest (**g, i**) in Arabidopsis hypocotyl in presence of 1 μM flg22 (**c, g**) or 1 μM eBL (**e, i**) or the corresponding mock controls (H$_2$O or EtOH). **d** and **f**, Quantification of FLS2-mEOS3.2 (**d**) or BRI1-mEOS3.2 (**f**) of clustered trajectories within 10 min of treatment. Each data point represents the percentage of clustered trajectories per cell (n), colors indicate independent experiments. Conditions sharing a letter are not significantly different in pairwise Wilcoxon with Bonferroni correction (p>0.05). **h** and **j,** Quantification of FLS2-mEOS3.2 (**h**) or BRI1-mEOS3.2 (**j**) spatial arrest within 10 min of treatment. Each data point represents the average number of spatial arrests per trajectory obtained per cell (n), colors indicate independent experiments

bak1-4/pBAK1::BAK1-mEOS3.2

pBAK1::BAK1-mEOS3.2 (*N. benthamiana*)

Molecule velocity - ▬▬▬ +

bak1-4/pBAK1::BAK1-mEOS3.2

bak1-4/pBAK1::BAK1-mEOS3.2/pFLS2::FLS2-GFP

Extended Data Figure 3 | Analysis of BAK1 diffusion and organization in presence and absence
of exogenous ligand treatment.
a and **g**, Representative images of BAK1-mEOS3.2 trajectories nanoclustering in Arabidopsis
hypocotyl in presence of 1 μM eBL (**a**) or 1 μM flg22 (**g**) or the corresponding mock controls (EtOH
or H_2O) in *bak1-4* (**a**) and *bak1-4*/pFLS2::FLS2-GFP (**g**). **b** and **h**, Quantification of clustered
trajectories within 10 min of treatment. Each data point represents the percentage of clustered
trajectories per cell (n), colors indicate independent experiments. **c** and **e**, Representative BAK1-
mEOS3.2 trajectories in presence of 1 μM flg22 or the corresponding mock control (H_2O)
expressed in *N. benthamiana* leaves (**c**) or stably expressed in Arabidopsis hypocotyl (**e**). **d** and **f,**
Quantification of instantaneous diffusion coefficient (D) within 10 min of treatment. Each data
point represents the average log(D) value obtained per cell (n), colors indicate independent
experiments. Conditions sharing a letter are not significantly different in pairwise Wilcoxon test
with Bonferroni correction (p<0.05).
322

pFLS2::FLS2-GFP/pBAK1::BAK1-mCherry

pBRI1::BRI1-GFP/pBAK1::BAK1-mCherry

Extended Data Figure 4 | Analysis of FLS2-BAK1 and BRI1-BAK1 co-localization.

a and **d**, Representative VA-TIRFM images of FLS2-GFP and BAK1-mCherry (**a**), and BRI1-GFP and BAK1-mCherry. **b** and **e**, Quantification of the co-localization between FLS2-GFP and BAK1-mCherry (**b**) and between BRI1-GFP and BAK1-mCherry (**e**). Each data point represents Pearson correlation coefficient for individual cells (n), lines connect raw and corresponding control rotated (90° mCherry) images. **c** and **f**, Quantification of the co-localization between FLS2-GFP and BAK1-mCherry (**c**) and between BRI1-GFP and BAK1-mCherry (**f**), in presence of 1 μM flg22 or the corresponding mock control (H_2O) (**c**) or control condition (f) in WT Col-0 or *fer-4*. Each data point represents the average Pearson correlation coefficient for individual cells (n), colors indicate independent experiments. Conditions which do not share a letter are significantly different in pairwise Wilcoxon test with Bonferroni p-value adjustment (p< 0,05).

bak1-4/pBAK1::BAK1^{F60A/F144A}-mEOS3.2

bak1-4/pBAK1::BAK1^{D416N}-mEOS3.2

bak1-4/pBAK1::BAK1-mEOS3.2

Extended Data Figure 5 | Analysis of the organization of BAK1 mutants in presence and absence of exogenous eBL treatment.

a, d and **g**, Representative single molecule trajectories clustering of BAK1^{F60AF144A}-mEOS3.2 (**a**), BAK1^{D416N}-mEOS3.2 (**d**) or BAK1^{WT}-mEOS3.2 and BAK1^{F60AF144A}-mEOS3.2 (**g**) in presence of 1 µM eBL (**a** and **d**) or the corresponding mock control (EtOH) (**a, d** and **g**). **b, e** and **h**, Quantification clustered trajectories within 10 min of treatment. Each data point represents the percentage of clustered trajectories per cell (n), colors indicate independent experiments. **c, f** and **i**, Quantification of BAK1^{F60AF144A}-mEOS3.2 (**c**), BAK1^{D416N}-mEOS3.2 (**f**) or BAK1^{WT}-mEOS3.2 and BAK1^{F60AF144A}-mEOS3.2 (**i**) spatial arrest within 10 min of treatment. Each data point represents the percentage of clustered trajectories per cell (n), colors indicate independent experiments. Conditions sharing a letter are not significantly different in pairwise Wilcoxon test with Bonferroni correction (p<0.05).

a BIR3-GFP PM-RFP

10 µm

b *bak1-4*/pUb::BAK1-mEOS3.2 *bir3-2*/pUb::BIR3-mEOS3.2

500 nm

☐ Spatial arrest

c

a   b

Spatial arrest

n=37   n=23

BAK1   BIR3

Extended Data Figure 6 | Analysis of BIR3 organization and diffusion.

a, Representative eSRRF images of BIR3-GFP control plasma membrane (PM) protein expressed in *N. benthamiana*. **b**, Representative single molecule trajectories of BAK1-mEOS3.2 and BIR3-mESO3.2, long term single molecule imaging in Arabidopsis hypocotyl. **c**, Quantification of spatial arrests. Each data point represents the average number of spatial arrests per trajectory obtained per cell (n), colors indicate independent experiments. Conditions sharing a letter are not significantly different in pairwise Wilcoxon test with Bonferroni correction (p<0.05).

a

gRNA

BIR3 (AT1G27190)

5′- ___________ -3′

bak1-4/pBAK1::BAK1-mEOS3.2

394-AAGC-TTGGTGGTTCGATTCCGACTCAGATCGTTGA

bir3-c/bak1-4/pBAK1::BAK1-mEOS3.2

394- AAGCTTTGGTGGTTCGATTCCGACTCAGATCGTTGA *

pBAK1::BAK1-mEOS3.2

b *bak1-4*   *bak1-4/bir3-c*

Molecule velocity - ▬ +

c
Log(D)

a   b

n=39   n=13

bak1-4   *bak1-4/bir3-c*

d *bak1-4*   *bak1-4/bir3-c*

Clusters

e
Clustered trajectories (%)

a   b

n=39   n=13

bak1-4   *bak1-4/bir3-c*

f *bak1-4*   *bak1-4/bir3-c*

▭ Spatial arrest

g
Spatial arrest

a   a

n=39   n=13

bak1-4   *bak1-4/bir3-c*

Extended Data Figure 7| Analysis of BAK1 diffusion and organization in a *BIR3* CRISPR mutant
a, Schematic describing the *bir3-c/bak1-4*/pBAK1::BAK1-mEOS3.2 CRISPR line. The number of base pairs from the start codon is indicated. Red letters represent insertions observed in gDNA sequencing. Asterisks indicate early stop codons. **b, d, f**. Representative trajectories of BAK1-mEOS3.2, long-term single molecule imaging in *bak1-4* or *bak1-4/bir3-c* Arabidopsis hypocotyls. **c**, Quantification of instantaneous diffusion coefficient (D). Each data point represents the average log(D) value obtained per cell (n), colors indicate independent experiments. **e**, Quantification of clustered trajectories. Each data point represents the percentage of clustered trajectories per cell (n), colors indicate independent experiments. **g**, Quantification of spatial arrests. Each data point represents the average number of spatial arrests per trajectory obtained per cell (n), colors indicate independent experiments. Conditions sharing a letter are not significantly different in pairwise Wilcoxon test with Bonferroni correction (p>0.05).

Extended Data Figure 8| Particle-based simulation recapitulates native protein behavior.
a, Quantification of the area of simulated FLS2 confinement zones (blue) and measured FLS2-
mEOS3.2 nanoclusters observed by spt-PALM and analysed by NASTIC (red). **b**, Quantification of
instantaneous diffusion coefficient (D) of simulated BAK1, FLS2 and BIR3 molecules. Conditions
sharing a letter are not significantly different in pairwise Wilcoxon test with Bonferroni correction
($p < 0.05$). **c**, Representative single molecule tracks of simulated BAK1, FLS2 and BIR3 molecules.
375

a
Col-0   *bir3-2*
− +   − +   flg22
170 kDa-   α-FLS2
70 kDa-   α-BAK1
IP α-FLS2
170 kDa-   α-FLS2
70 kDa-   α-BAK1
Input

b
Col-0   *bir3-2*
− +   − +   flg22
170 kDa-   α-FLS2
70 kDa-   α-BAK1
IP α-FLS2
170 kDa-   α-FLS2
70 kDa-   α-BAK1
Input

c
Col-0   *bir3-2*
− 20 100   − 20 100   nM flg22
170 kDa-   α-FLS2
70 kDa-   α-BAK1
IP α-FLS2
170 kDa-   α-FLS2
70 kDa-   α-BAK1
Input

d
flg22
Col-0   *bir3-2*
170 kDa-   α-FLS2
55 kDa-   α-pMAPK

e
Col-0
fls2
bir3-2
Photon counts (x10⁵)
Time (min)

f
Hypocotyl length (rel. to ctrl.)
a b a c a c
Mock
eBL
1 2 3
n=121 n=149 n=121 n=123 n=179 n=189
Col-0   *bak1-4*   *bir3-2*

g
Hypocotyl length (rel. to ctrl.)
a b a c a b
Mock
eBL
1 2 3
n=121 n=123 n=161 n=194 n=157 n=181
bak1-4   WT   *bir3c*
bak1-4/pBAK1::BAK1-mEOS3.2

Extended Data Figure 9| Analyses of eBL and flg22 responsiveness in *BIR3* loss-of-function
mutants.
a, **b**, and **c** Immunoprecipitation of FLS2 in Col-0 or *bir3-2* after treatment with 100 nM flg22 (a,
b, c) or corresponding mock solution (H$_2$O) for 5 min (**a, b, c**) or 20 nM flg22 and corresponding
mock solution (H$_2$O) for 2 min (**c**). Membranes were probed with anti-FLS2 or anti-BAK1. **c**d
Western blot analysis of MAPK phosphorylation in Arabidopsis seedlings treated for 5 min with
100 nM flg22. Membranes were probed with anti-FLS2 or anti-pERK42/44 (pMAPK) antibodies.
The same observations were made in 3 independent experiments. **e**, Time course analysis of ROS
production in 4-week-old Arabidopsis leaf discs treated with 100 nM flg22, values are mean ± se
of 4 independent experiments. **f** and **g**, Quantification of the hypocotyl length of 5-day-old
Arabidopsis seedlings grown on half MS medium containing 200 nM epi-Brassinolide (eBL) or
corresponding mock control solution (EtOH). Data points indicate measurements of the length of
individual hypocotyl relative to mock for each genotype, colors indicate independent experiments
Conditions sharing a letter are not significantly different in pairwise Wilcoxon test with Bonferroni
correction (p<0.05).
391

Material and methods

Plant materials and growth

Arabidopsis thaliana ecotype Columbia (Col-0) was used as WT control. Col-0/pFLS2::FLS2-GFP[59], *bak1-4*[11], Col-0/pBRI1::BRI1-GFP[25], *fls2c*[60], *bir3-2*[14], Col-0/pBRI1::BRI1-GFP/BAK1-mCherry[20], *bak1-4*/pBAK1::BAK1-GFP[26], Col-0/pBRI1::BRI1-mEOS3.2[61], were previously described. Col-0/pFLS2::FLS2-GFP[59], was crossed with Col-0/pBAK1::BAK1-mCherry[62] to obtain Col-0/pFLS2::FLS2-GFP/pBAK1::BAK1-mCherry. The following stable Arabopidopsis lines *fls2*/pUb::FLS2-mEOS3.2, *fls2*/pFLS2::FLS2-mEOS3.2, *bak1-4*/pBAK1::BAK1-mEOS3.2, *bak1-4*/pUb::BAK1-mEOS3.2, *bak1-4*/pBAK1::BAK1^{D416N}-mEOS3.2, *bak1-4*/pBAK1::BAK1$^{F60A/F144A}$-mEOS3.2, *bak1-4*/pUb::BAK1^{D122A}-mEOS3.2 and *bir3-2*/pUb::BIR3-mEOS3.2 were generated by the transforming corresponding knock-out backgrounds via floral dipping[63]. The transformants were selected on hygromycin (25 mg/L) to obtain homozygous mono insertional lines and analyzed by western blotting using anti-BAK1 or anti-FLS2 antibodies. Col-0/pFLS2::FLS2-GFP/pBAK1::BAK1-mCherry and Col-0/pBRI1::BRI1-GFP/BAK1-mCherry were crossed with *fer-4*[64] to obtain *fer-4*/pFLS2::FLS2-GFP/pBAK1::BAK1-mCherry and *fer-4*/pFLS2::FLS2-GFP/pBAK1::BAK1-mCherry. Col-0/pFLS2::FLS2-GFP was crossed with *bak1-4*/pBAK1::BAK1-mEOS3.2 to obtain *bak1-4*/pFLS2::FLS2-GFP/pBAK1::BAK1-mEOS3.2. *bir3-c/bak1-4*/pBAK1::BAK1-mEOS3.2 was obtained by transforming *bak1-4*/pBAK1::BAK1-mEOS3.2 with CRISPR-Cas9 constructs carrying a gRNA targeting *BIR3*. Transformants were selected based on seed coat fluorescence, sequenced and T3 Cas9i-negative homozygous *BIR3* mutants were selected for functional analyses. To analyse flg22-induced ROS production, plants were grown in individual pots at 21 °C with a 10-h light photoperiod in environmentally controlled growth rooms. For seedlings-based assays, seeds were surface-sterilized by incubating them in 0.1 % Tween20 in 70 % EtOH for 10 min, following 70 % EtOH for 10 min and 100% EtOH for 1 min. Seeds were stratified for 2 d in the dark at 4 °C and grown on half Murashige and Skoog (MS) media supplemented with vitamins, 1 % sucrose and 0.8 % agar at 22 °C and a 16-h light photoperiod.

Molecular cloning, plant transformation and CRISPR-Cas9-mediated mutagenesis

FLS2, BAK1 and BIR3 coding sequences and FLS2 and BAK1 promotors were obtained as synthetic fragments (Genewiz) and cloned into the level 1 Golden Gate plasmid pUC57_BB03 using Bpil[65,66]. The plasmid encoding for mEOS3.2 was previously described[32]. Nucleic acid fragments were assembled in the level 2 Golden Gate plasmid (BB24_LIIβ F 3-4)[66]. *BIR3* gRNA for CRISPR Cas-mediated editing was designed using CHOPCHOP (https://chopchop.cbu.uib.no/) and inserted into pAGM55261 using Golden Gate cloning[65]. All plasmids were verified by Sanger sequencing. The binary plasmids were transferred into

the *Agrobacterium tumefaciens* strain GV3101 for flower dip transformation of Arabidopsis
or transient transformation of *Nicotiana benthamiana* leaves.

Brassinosteroid-sensitivity assay
Brassinosteroid sensitivity assay was performed as previously described in[43]. Briefly, seeds
were surface-sterilized and individually placed in line on square Petri dishes containing half
MS 1 % sucrose, 0.8 % phytoagar, supplemented with 200 nM epi-brassinolide (Sigma,
Cat#E1641) or corresponding control solution (EtOH). The seeds were stratified at 4 °C for 2
d and then placed vertically in a growth chamber for 5 d. Hypocotyl length were measured
using Fiji[67].

Co-immunoprecipitation
Sample preparation and immunoprecipitation were performed as previously described[68].
Briefly, seedlings were grown on half MS media for 5 d and transferred to 6-well plates
containing liquid half MS to be cultivated for ten more days. Subsequently, seedlings from
10 wells were gathered and let rest for 1 h in liquid half MS. Seedlings were treated for 4 min
with either 20 nM or 100 nM flg22 or mock (distilled sterile water) and flash frozen in liquid
nitrogen. For protein extraction, tissues were grinded in liquid nitrogen and proteins were
isolated in 50 mM Tris-HCl pH 7.5, 150 mM NaCl, 10 % glycerol, 5 mM dithiothreitol, 1 %
protease inhibitor cocktail (Sigma-Aldrich), 2 mM Na2MoO4, 2.5 mM NaF, 1.5 mM activated
Na3VO4, 1 mM phenylmethanesulfonyl fluoride, and 0.5 % IGEPAL. For
immunoprecipitations, trueblot agarose beads (Rockland, 18-8816-33) were conjugated
with FLS2 antibodies (Agrisera, AS12 1857) and incubated with the crude protein extracts for
3 to 4 h at 4 °C. Subsequently, beads were washed three times with a washing buffer (50 mM
Tris-HCl pH 7.5, 150 mM NaCl, 1 mM phenylmethanesulfonyl fluoride, 0,1 % IGEPAL) before
adding 60 µL of a protein loading solution (6xSDS 10 % DTT) and incubating for 10 min at 90
°C. Analysis was carried out by SDS-PAGE and immunoblotting.

Immunoblotting
Protein samples were separated in 10 % bisacrylamide gels at 150 V for approximately 2
hours and transferred into activated PVDF membranes at 100 V for 90 min. Immunoblotting
was performed with antibodies diluted in blocking solution (5 % fat-free milk in TBS with 0.1
% [v/v] Tween-20). Antibodies used in this study were α-BAK1 (1:5000, Agrisera AS12 1858),
α-FLS2 (1:5000, Agrisera AS12 1857) or α-pMAPK (1:5000, pERK42/44 cell signal 9101S).
Blots were developed with Pierce ECL/ECL Femto Western Blotting Substrate (Thermo
Scientific). The following secondary antibody was used: anti-rabbit IgG (whole molecule)–
HRP (A0545, Sigma, dilution 1:10,000).

Variable angle total internal reflection microscopy (VA-TIRFM)

VA-TIRM imaging was performed as previously described in[32] Briefly, VA-TIRFM was operated on a custom-build TIRFM set-up[61] equipped with an 100x objective NA 1.49 (Zeiss, 421190-9800-000), four laser lines (405, 488, 561 and 642 nm), a polychromatic modulator (AOTF, AA OPTO-ELECTRONIC) and a sCMOS camera (Hamamatsu Photonics, ORCA-Flash4.0 V2). To ensure that the target laser power value always corresponds to the irradiance in the sample plane, the power of each laser line was calibrated before each experiment using a PM100D detector (Thorlabs). Five-day-old Arabidopsis seedlings or 3-week-old *N. benthamiana* leave samples (28 to 34 h post Agrobacterium infiltration) were delicately mounted between two coverslips (Epredia 24×50 mm #1) in liquid ½ MS medium and placed on the specimen stage without additional weight. Images were acquired at 20 Hz frame rate (50 ms), 6.66 Hz (150 ms) or 5 Hz (200 ms) for spt-PALM and photochromic reversion imaging conditions.

Single molecule imaging and analyses

Photochromic reversion and spt-PALM imaging were performed as previously described[32]. Briefly, for photochromic reversion, mEOS3.2 was photoconverted, excited and kept in a prolonged fluorescent state using 1000 µW 561nm laser power, 2-15 µW 405 nm laser power and 1-30 µW 488 nm laser power. The emitted light was collected using 568 LP Edge Basic Long-pass Filter, 584/40 ET Band-pass filters (AHF analysentechnik AG) and recorded from a 51.2 x 51.2 µm region of interest, 100 nm pixel size. Track reconstruction and analysis was carried out as previously described[32]. To ensure bona-fide single-molecule tracking we analyzed frames with relatively low molecule density (ca. 0.1 – 1 molecule per µm^2). We used the plugin TrackMate[69] in Fiji[67] to reconstruct single molecule trajectories. Single particles were segmented frame-by-frame by applying a Laplacian of Gaussian (LoG) filter and estimated particle size of 0.3 µm with a Quality threshold of 30. Individual single particles were localized with sub-pixel resolution using a built-in quadratic fitting scheme. Single-particle trajectories were reconstructed using a simple linear assignment problem[70] with a maximal linking distance of 0.4 µm and a 2 frames-gap-closing maximum. The coordinates of the single particle trajectories were then further analyzed using a custom-made python script to calculate the mean square displacement (MSD) and diffusion coefficient (D) in batch mode. Only tracks with a minimum of five localizations were kept for analysis. The MSD and diffusion coefficient was calculated based on the first four time points of each trajectory as previously described[71]. Single molecule trajectories clustering was performed using nastic[35] with the following threshold, time = 20 s, radius factor = 1.2, minimum trajectories = 3, size screen = 0.15 µm. Spatial arrests were analyzed using CASTA, with the default minimum track length of 25 timepoints[32] (https://pypi.org/project/casta/).

Enhanced super-resolution radial fluctuations (eSRRF)

Stable transgenic lines expressing BRI1-GFP, BAK1-GFP or FLS2-GFP were imaged by VA-TIRFM as described above at 5 Hz (200 ms) frame rate. Super resolved images were reconstructed using eSRRF[29] in Fiji[67]. Stream VA-TIRFM acquisitions were processed using the following parameters that were defined by iterative parameter sweep[29]: magnification = 6, radius = 3, sensitivity = 3 and frames = 100. The average (AVG) reconstructions of BRI1-GFP, BAK1-GFP and FLS2-GFP plasma membrane organization were segmented and analysed using Nanonet, a custom-made automated analysis pipeline (https://github.com/NanoSignalingLab/NanoNet), providing nanodomain density and image wide spatial clustering index (iSCI) based on predicted localization density.

Measurement of reactive oxygen species production

Arabidopsis plants were grown in individual pots at 21 °C with a 10-h light photoperiod in environmentally controlled growth rooms. Leaf discs (4 mm in diameter) were excised from 3- to 4-week-old plants and incubated in 96-well plates containing sterile distilled water overnight. The following day, water was replaced with a solution containing 0.5 µM L-012 and 20 µg/mL horseradish peroxidase (HRP, Sigma Aldrich) and 100 nM flg22 or control (water). Luminescence was measured using a microplate reader (Promega Glomax navigator) over a period of 60 min.

Particle based simulations

Computational simulations of single particles were performed using the Smoldyn software (version 2.75)[53], and a custom-made python script. We simulated a 2D space of 10 µm^2 with periodic boundaries conditions to avoid edge effects. We imposed a triangular mesh with an average compartment size corresponding to the size FLS2 nanoclusters obtained by NASTIC. To simulate FLS2, BAK1 and BIR3 molecular diffusion we provided as input data the instantaneous diffusion coefficient (D) measured by photochromic reversion. For each step and each molecule, a new displacement is randomly chosen from normally distributed possible displacements. The probability of molecule diffusion across compartment boundaries was defined as 0.00001 for FLS2, 0.00001 for BIR3 and 1 for BAK1. To simulate molecule binding kinetics, we used the Kon and Koff of flg22-bound FLS2 to BAK1[54] and the Kd of BAK1 to BIR3[13] converted into binding and unbinding radii. Simulations were run using a delta t of 0.0001s to ensure that the spatial resolution was smaller than the geometric features and the characteristic time scale of the system for each possible reaction. The initial location of each molecule was randomized for each simulation. To analyse the effect of BIR3 on BAK1 localization we monitored the number of BAK1 molecules within FLS2 nanodomains every 5 ms for 300 s. To simulate flg22 perception, FLS2 molecules were

542 converted to flg22-bound FLS2 molecules after 120 s and FLS2-flg22/BAK1 complexes were
543 measured every 50 ms for 300 s.

544
545

References

1   DeFalco, T. A. & Zipfel, C. Molecular mechanisms of early plant pattern-triggered immune
signaling. *Mol Cell* **81**, 3449–3467 (2021). https://doi.org/10.1016/j.molcel.2021.07.029
2   Smakowska-Luzan, E. *et al.* An extracellular network of Arabidopsis leucine-rich repeat
receptor kinases. *Nature* **553**, 342–346 (2018). https://doi.org/10.1038/nature25184
3   Özkan, E. *et al.* An extracellular interactome of immunoglobulin and LRR proteins reveals
receptor-ligand   networks.   *Cell*   **154**,   228–239   (2013).
https://doi.org/10.1016/j.cell.2013.06.006
4   Monaghan, J. & Zipfel, C. Plant pattern recognition receptor complexes at the plasma
membrane.   *Curr   Opin   Plant   Biol*   **15**,   349–357   (2012).
https://doi.org/10.1016/j.pbi.2012.05.006
5   Lemmon, M. A. & Schlessinger, J. Cell signaling by receptor tyrosine kinases. *Cell* **141**,
1117–1134 (2010). https://doi.org/10.1016/j.cell.2010.06.011
6   Shiu, S. H. & Bleecker, A. B. Receptor-like kinases from Arabidopsis form a monophyletic
gene family related to animal receptor kinases. *Proc Natl Acad Sci U S A* **98**, 10763–10768
(2001). https://doi.org/10.1073/pnas.181141598
7   Li, J. & Chory, J. A Putative Leucine-Rich Repeat Receptor Kinase Involved in
Brassinosteroid   Signal   Transduction.   *Cell*   **90**,   929–938   (1997).
https://doi.org/https://doi.org/10.1016/S0092-8674(00)80357-8
8   Jinn, T. L., Stone, J. M. & Walker, J. C. HAESA, an Arabidopsis leucine-rich repeat receptor
kinase, controls floral organ abscission. *Genes Dev* **14**, 108–117 (2000).
9   Gómez-Gómez, L., Bauer, Z. & Boller, T. Both the extracellular leucine-rich repeat domain
and the kinase activity of FSL2 are required for flagellin binding and signaling in
Arabidopsis. *Plant Cell* **13**, 1155–1163 (2001).
10 Nam, K. H. & Li, J. BRI1/BAK1, a receptor kinase pair mediating brassinosteroid signaling.
Cell **110**, 203–212 (2002). https://doi.org/10.1016/s0092-8674(02)00814-0
11 Chinchilla, D. *et al.* A flagellin-induced complex of the receptor FLS2 and BAK1 initiates
plant defence. *Nature* **448**, 497–500 (2007). https://doi.org/10.1038/nature05999
12 Ma, C. *et al.* Structural basis for BIR1-mediated negative regulation of plant immunity. *Cell*
Research **27**, 1521–1524 (2017). https://doi.org/10.1038/cr.2017.123
13 Hohmann, U., Nicolet, J., Moretti, A., Hothorn, L. A. & Hothorn, M. The SERK3 elongated
allele defines a role for BIR ectodomains in brassinosteroid signalling. *Nat Plants* **4**, 345–
351 (2018). https://doi.org/10.1038/s41477-018-0150-9
14 Imkampe, J. *et al.* The Arabidopsis Leucine-Rich Repeat Receptor Kinase BIR3 Negatively
Regulates BAK1 Receptor Complex Formation and Stabilizes BAK1. *Plant Cell* **29**, 2285–
2303 (2017). https://doi.org/10.1105/tpc.17.00376
15 Großeholz, R. *et al.* Specifying the role of BAK1-interacting receptor-like kinase 3 in
brassinosteroid   signaling.   *J   Integr   Plant   Biol*   **62**,   456–469   (2020).
https://doi.org/10.1111/jipb.12803

16 Bücherl, C. A. *et al.* Plant immune and growth receptors share common signalling
components but localise to distinct plasma membrane nanodomains. *Elife* **6** (2017).
https://doi.org/10.7554/eLife.25114

17 Gronnier, J. *et al.* Regulation of immune receptor kinase plasma membrane nanoscale
organization by a plant peptide hormone and its receptors. *eLife* **11**, e74162 (2022).
https://doi.org/10.7554/eLife.74162

18 McKenna, J. F. *et al.* The cell wall regulates dynamics and size of plasma-membrane
nanodomains in Arabidopsis. *Proceedings of the National Academy of Sciences* **116**,
12857–12862 (2019). https://doi.org/doi:10.1073/pnas.1819077116

19 Cui, Y. *et al.* Sterols regulate endocytic pathways during flg22-induced defense responses
in Arabidopsis. *Development* **145** (2018). https://doi.org/10.1242/dev.165688

20 Hutten, S. J. *et al.* Visualization of BRI1 and SERK3/BAK1 Nanoclusters in Arabidopsis
Roots. *PLOS ONE* **12**, e0169905 (2017). https://doi.org/10.1371/journal.pone.0169905

21 Ma, Z., Xie, Y., Chng, C.-P., Huang, C. & Miao, Y. Synthetic FLS2 receptor oligomer boosts
plant   innate   immunity.   *bioRxiv*,   2025.2001.2027.634983   (2025).
https://doi.org/10.1101/2025.01.27.634983

22 Jaillais, Y. *et al.* Guidelines for naming and studying plasma membrane domains in plants.
Nat Plants **10**, 1172–1183 (2024). https://doi.org/10.1038/s41477-024-01742-8

23 Liu, M. J. *et al.* Extracellular pectin-RALF phase separation mediates FERONIA global
signaling   function.   *Cell*   **187**,   312–330.e322   (2024).
https://doi.org/10.1016/j.cell.2023.11.038

24 Robatzek, S., Chinchilla, D. & Boller, T. Ligand-induced endocytosis of the pattern
recognition receptor FLS2 in Arabidopsis. *Genes Dev* **20**, 537–542 (2006).
https://doi.org/10.1101/gad.366506

25 Geldner, N., Hyman, D. L., Wang, X., Schumacher, K. & Chory, J. Endosomal signaling of
plant steroid receptor kinase BRI1. *Genes Dev* **21**, 1598–1602 (2007).
https://doi.org/10.1101/gad.1561307

26 Ntoukakis, V., Schwessinger, B., Segonzac, C. & Zipfel, C. Cautionary Notes on the Use of
C-Terminal BAK1 Fusion Proteins for Functional Studies. *The Plant Cell* **23**, 3871–3878
(2011). https://doi.org/10.1105/tpc.111.090779

27 Tokunaga, M., Imamoto, N. & Sakata-Sogawa, K. Highly inclined thin illumination enables
clear single-molecule imaging in cells. *Nat Methods* **5**, 159–161 (2008).
https://doi.org/10.1038/nmeth1171

28 Konopka, C. A. & Bednarek, S. Y. Variable-angle epifluorescence microscopy: a new way to
look at protein dynamics in the plant cell cortex. *The Plant Journal* **53**, 186–196 (2008).
https://doi.org/https://doi.org/10.1111/j.1365-313X.2007.03306.x

29 Laine, R. F. *et al.* High-fidelity 3D live-cell nanoscopy through data-driven enhanced super-
resolution   radial   fluctuation.   *Nature   Methods*   **20**,   1949–1956   (2023).
https://doi.org/10.1038/s41592-023-02057-w

30 Wang, L. *et al.* Spatiotemporal Dynamics of the BRI1 Receptor and its Regulation by
Membrane Microdomains in Living Arabidopsis Cells. *Molecular Plant* **8**, 1334–1349
(2015). https://doi.org/https://doi.org/10.1016/j.molp.2015.04.005

Tran, T. M. *et al.* The bacterial quorum sensing signal DSF hijacks Arabidopsis thaliana sterol biosynthesis to suppress plant innate immunity. *Life Sci Alliance* **3** (2020). https://doi.org/10.26508/lsa.202000720

von Arx, M., Xhelilaj, K., Schulz, P., zur Oven-Krockhaus, S. & Gronnier, J. Photochromic reversion enables long-term tracking of single molecules in living plants. *bioRxiv*, 2024.2004.2010.585335 (2024). https://doi.org/10.1101/2024.04.10.585335

Zhang, M. *et al.* Rational design of true monomeric and bright photoactivatable fluorescent proteins. *Nature Methods* **9**, 727–729 (2012). https://doi.org/10.1038/nmeth.2021

Grefen, C. *et al.* A ubiquitin-10 promoter-based vector set for fluorescent protein tagging facilitates temporal stability and native protein distribution in transient and stable expression studies. *Plant J* **64**, 355–365 (2010). https://doi.org/10.1111/j.1365-313X.2010.04322.x

Wallis, T. P. *et al.* Super-resolved trajectory-derived nanoclustering analysis using spatiotemporal indexing. *Nature Communications* **14**, 3353 (2023). https://doi.org/10.1038/s41467-023-38866-y

Low-Nam, S. T. *et al.* ErbB1 dimerization is promoted by domain co-confinement and stabilized by ligand binding. *Nat Struct Mol Biol* **18**, 1244–1249 (2011). https://doi.org/10.1038/nsmb.2135

Niwa, F. *et al.* Gephyrin-Independent GABAAR Mobility and Clustering during Plasticity. *PLOS ONE* **7**, e36148 (2012). https://doi.org/10.1371/journal.pone.0036148

Sako, Y., Minoghchi, S. & Yanagida, T. Single-molecule imaging of EGFR signalling on the surface of living cells. *Nature Cell Biology* **2**, 168–172 (2000). https://doi.org/10.1038/35004044

Wang, Y. *et al.* Regulation of EGFR nanocluster formation by ionic protein-lipid interaction. *Cell Research* **24**, 959–976 (2014). https://doi.org/10.1038/cr.2014.89

Schulze, B. *et al.* Rapid heteromerization and phosphorylation of ligand-activated plant transmembrane receptors and their associated kinase BAK1. *J Biol Chem* **285**, 9444–9451 (2010). https://doi.org/10.1074/jbc.M109.096842

Delesalle, C., Vert, G. & Fujita, S. The cell surface is the place to be for brassinosteroid perception and responses. *Nature Plants* **10**, 206–218 (2024). https://doi.org/10.1038/s41477-024-01621-2

Hurst, C. H. *et al.* S-acylation stabilizes ligand-induced receptor kinase complex formation during plant pattern-triggered immune signaling. *Current Biology* **33**, 1588–1596.e1586 (2023). https://doi.org/10.1016/j.cub.2023.02.065

Biermann, D. *et al.* A RALF-brassinosteroid signaling circuit regulates Arabidopsis hypocotyl cell shape. *Current Biology* **35**, 5002–5017.e5005 (2025). https://doi.org/10.1016/j.cub.2025.09.016

Stegmann, M. *et al.* The receptor kinase FER is a RALF-regulated scaffold controlling plant immune signaling. *Science* **355**, 287–289 (2017). https://doi.org/doi:10.1126/science.aal2541

Sun, Y. *et al.* Structural basis for flg22-induced activation of the Arabidopsis FLS2-BAK1 immune complex. *Science* **342**, 624–628 (2013). https://doi.org/10.1126/science.1243825

46 Santiago, J. *et al.* Mechanistic insight into a peptide hormone signaling complex mediating
floral organ abscission. *eLife* **5**, e15075 (2016). https://doi.org/10.7554/eLife.15075

47 Santiago, J., Henzler, C. & Hothorn, M. Molecular Mechanism for Plant Steroid Receptor
Activation by Somatic Embryogenesis Co-Receptor Kinases. *Science* **341**, 889–892 (2013).
https://doi.org/doi:10.1126/science.1242468

48 Sun, Y. *et al.* Structure reveals that BAK1 as a co-receptor recognizes the BRI1-bound
brassinolide. *Cell Research* **23**, 1326–1329 (2013). https://doi.org/10.1038/cr.2013.131

49 Schwessinger, B. *et al.* Phosphorylation-Dependent Differential Regulation of Plant
Growth, Cell Death, and Innate Immunity by the Regulatory Receptor-Like Kinase BAK1.
PLOS Genetics **7**, e1002046 (2011). https://doi.org/10.1371/journal.pgen.1002046

50 Wang, X. *et al.* Sequential Transphosphorylation of the BRI1/BAK1 Receptor Kinase
Complex Impacts Early Events in Brassinosteroid Signaling. *Developmental Cell* **15**, 220–
235 (2008). https://doi.org/10.1016/j.devcel.2008.06.011

51 Halter, T. *et al.* The leucine-rich repeat receptor kinase BIR2 is a negative regulator of BAK1
in   plant   immunity.   *Curr   Biol*   **24**,   134–143   (2014).
https://doi.org/10.1016/j.cub.2013.11.047

52 Guzmán-Benito, I. *et al.* The immune repressor BIR1 contributes to antiviral defense and
undergoes transcriptional and post-transcriptional regulation during viral infections. *New*
Phytol **224**, 421–438 (2019). https://doi.org/10.1111/nph.15931

53 Andrews, S. S. Smoldyn: particle-based simulation with rule-based modeling, improved
molecular interaction and a library interface. *Bioinformatics* **33**, 710–717 (2016).
https://doi.org/10.1093/bioinformatics/btw700

54 Okuda, S. *et al.* Molecular mechanism for the recognition of sequence-divergent CIF
peptides by the plant receptor kinases GSO1/SGN3 and GSO2. *Proceedings of the National*
Academy   of   Sciences   **117**,   2693–2703   (2020).
https://doi.org/doi:10.1073/pnas.1911553117

55 Sun, Y. *et al.* A Leucine-Rich Repeat Receptor-like Kinase TaBIR1 Contributes to Wheat
Resistance against Puccinia striiformis f. sp. tritici. *International Journal of Molecular*
Sciences **24**, 6438 (2023).

56 Liu, F. *et al.* BAK1 protects the receptor-like kinase BIR2 from SNIPER2a/b-mediated
degradation to promote pattern-triggered immunity in Nicotiana benthamiana. *The Plant*
Cell **35**, 3566–3584 (2023).

57 Guo, J. *et al.* OsBIR3 maintains the homeostasis of OsBRI1, OsREM4.1, and Brd2 protein
levels in brassinosteroid pathways in rice. *Plant Biotechnology Journal* **23**, 3024–3040
(2025). https://doi.org/https://doi.org/10.1111/pbi.70128

58 You, C. *et al.* Receptor dimer stabilization by hierarchical plasma membrane
microcompartments regulates cytokine signaling. *Science Advances* **2**, e1600452 (2016).
https://doi.org/doi:10.1126/sciadv.1600452

59 Göhre, V. *et al.* Plant Pattern-Recognition Receptor FLS2 Is Directed for Degradation by the
Bacterial   Ubiquitin   Ligase   AvrPtoB.   *Current   Biology*   **18**,   1824–1832   (2008).
https://doi.org/10.1016/j.cub.2008.10.063

60 Zipfel, C. *et al.* Bacterial disease resistance in Arabidopsis through flagellin perception.
Nature **428**, 764–767 (2004). https://doi.org/10.1038/nature02485

61 Rohr, L. *et al.* OneFlowTraX: a user-friendly software for super-resolution analysis of single-
molecule dynamics and nanoscale organization. *Frontiers in Plant Science* **Volume 15 -**
2024 (2024). https://doi.org/10.3389/fpls.2024.1358935

62 Bücherl, C. A. *et al.* Visualization of BRI1 and BAK1(SERK3) Membrane Receptor
Heterooligomers during Brassinosteroid Signaling *Plant Physiology* **162**, 1911–1925
(2013). https://doi.org/10.1104/pp.113.220152

63 Clough, S. J. & Bent, A. F. Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. *Plant J* **16**, 735–743 (1998).
https://doi.org/10.1046/j.1365-313x.1998.00343.x

64 Duan, Q., Kita, D., Li, C., Cheung, A. Y. & Wu, H. M. FERONIA receptor-like kinase regulates
RHO GTPase signaling of root hair development. *Proc Natl Acad Sci U S A* **107**, 17821–
17826 (2010). https://doi.org/10.1073/pnas.1005366107

65 Engler, C. *et al.* A Golden Gate Modular Cloning Toolbox for Plants. *ACS Synthetic Biology*
3, 839–843 (2014). https://doi.org/10.1021/sb4001504

66 Binder, A. *et al.* A modular plasmid assembly kit for multigene expression, gene silencing
and silencing rescue in plants. *PLoS One* **9**, e88218 (2014).
https://doi.org/10.1371/journal.pone.0088218

67 Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nature*
Methods **9**, 676–682 (2012). https://doi.org/10.1038/nmeth.2019

68 Kadota, Y. *et al.* Direct regulation of the NADPH oxidase RBOHD by the PRR-associated
kinase BIK1 during plant immunity. *Mol Cell* **54**, 43–55 (2014).
https://doi.org/10.1016/j.molcel.2014.02.021

69 Ershov, D. *et al.* TrackMate 7: integrating state-of-the-art segmentation algorithms into
tracking pipelines. *Nature Methods* **19**, 829–832 (2022). https://doi.org/10.1038/s41592-
022-01507-1

70 Jaqaman, K. *et al.* Robust single-particle tracking in live-cell time-lapse sequences. *Nature*
Methods **5**, 695–702 (2008). https://doi.org/10.1038/nmeth.1237

71 Saxton, M. J. & Jacobson, K. Single-particle tracking: applications to membrane dynamics.
Annu Rev Biophys Biomol Struct **26**, 373–399 (1997).
https://doi.org/10.1146/annurev.biophys.26.1.373

744