Digital Twin Brain simulation and manipulation of a functional brain network underlying

mental illness.

Authors:

Yunman Xia[1, 2†], Songjun Peng[3†], Juergen Dukart[4, 5], Chao Xie[6], Shitong Xiang[7, 8], Spase

Petkoski[9], Zilin Li[10], Joerg Hipp[11], Suresh Muthukumaraswamy[12], Anna Forsyth[12], Tianye Jia[2,

7, 8], Nilakshi Vaidya[1], Tristram, Lett[1], Liyi Qian[13], Xiao Chang[1, 2, 7, 8], Yuxiang Dai[1, 2], Tobias

Banaschewski[14], Gareth J. Barker[15], Arun L. W. Bokde[16], Rüdiger Brühl[17], Sylvane

Desrivières[18], Herta Flor[19, 20], Penny Gowland[21], Antoine Grigis[22], Andreas Heinz[23, 24], Hervé

Lemaître[22, 25], Frauke Nees[26], Dimitri Papadopoulos Orfanos[20], Luise Poustka[27], Michael N.

Smolka[24], Sarah Hohmann[14], Henrik Walter[28], Robert Whelan[29], Paul Wirsching[28], Zuo

Zhang[18, 30], Lauren Robinson[18, 31], Jeanne Winterer[30, 32], Yuning Zhang[33], Hedi Kebir[1], Ulrike

Schmidt[34, 35], Julia Sinclair[36], Yuchen Liu[7], Jiexiang Wang[7], Fei Dai[7], Longbin Zeng[7], Yubo

Hou[37], Huarui Wang[7], Leijun Ye[7], Chunhe Li[7, 38], Qibao Zheng[7], Andre Marquand[39],

Changsong Zhou[3], Viktor Jirsa[9], Jianfeng Feng[7, 8, 38, 40-42], Wenlian Lu[7, 8, 38, 43], Gunter

Schumann[1, 2, 44-46*], IMAGEN Consortium, STRATIFY Consortium, DTB Consortium, and

environMENTAL Consortium.

Affiliations:

1.   Centre for Population Neuroscience and Stratified Medicine (PONS), Department of

Psychiatry and Neuroscience, Charite-Universitaetsmedizin Berlin, Germany.

2.   PONS, Institute of Science and Technology for Brain-Inspired Intelligence, Fudan

University, Shanghai, China.

3.   Department of Physics, Hong Kong Baptist University, Hong Kong SAR, China

4. Institute of Neurosciences and Medicine, Brain & Behaviour (INM-7), Research Centre Juelich, Juelich, Germany.

5. Institute of Systems Neuroscience, Medical Faculty, Heinrich Heine University, Düsseldorf, Germany.

6. Department of Psychological and Cognitive Sciences, Tsinghua University, Beijing, China.

7. Institute of Science and Technology for Brain-Inspired Intelligence, Fudan University, Shanghai, China.

8. Key Laboratory of Computational Neuroscience and Brain-Inspired Intelligence, Fudan University, Ministry of Education, Shanghai, China.

9. Institut de Neurosciences des Systèmes (INS) UMR1106, Aix Marseille Université, Marseille, France.

10. Department of Psychiatry, University of Oxford, Oxford, United Kingdom.

11. Roche Pharma Research and Early Development, Neuroscience and Rare Diseases, Roche Innovation Center Basel, F. Hoffmann–La Roche Ltd., Basel, Switzerland.

12. School of Pharmacy, Faculty of Medical and Health Sciences, University of Auckland, Auckland, New Zealand.

13. Department of Neurosurgery, Huashan Hospital, Fudan University, Shanghai, China.

14. Department of Child and Adolescent Psychiatry and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany.

15. Department of Neuroimaging, Institute of Psychiatry, Psychology & Neuroscience,
King's College London, United Kingdom.

16. Discipline of Psychiatry, School of Medicine and Trinity College Institute of
Neuroscience, Trinity College Dublin, Dublin, Ireland.

17. Physikalisch-Technische Bundesanstalt (PTB), Braunschweig and Berlin, Germany.

18. Social, Genetic and Developmental Psychiatry Centre, Institute of Psychiatry,
Psychology & Neuroscience, King's College London, United Kingdom.

19. Institute of Cognitive and Clinical Neuroscience, Central Institute of Mental Health,
Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany.

20. Department of Psychology, School of Social Sciences, University of Mannheim,
Mannheim, Germany.

21. Sir Peter Mansfield Imaging Centre School of Physics and Astronomy, University of
Nottingham, University Park, Nottingham, United Kingdom.

22. NeuroSpin, CEA, Université Paris-Saclay, F-91191 Gif-sur-Yvette, France.

23. Department of Psychiatry and Psychotherapy, University of Tübingen, Germany;
German Center for Mental Health (DZPG), Site Tübingen, Germany.

24. Department of Psychiatry and Psychotherapy, Technische Universität Dresden, Dresden,
Germany.

25. Institut des Maladies Neurodégénératives, UMR 5293, CNRS, CEA, Université de
Bordeaux, 33076 Bordeaux, France.

26. Institute of Medical Psychology, Ludwig-Maximilians-Universität (LMU) in Munich,
Munich, Germany.

27. Department of Child and Adolescent Psychiatry, Center for Psychosocial Medicine,
University Hospital Heidelberg, Heidelberg, Germany.

28. Department of Psychiatry and Psychotherapy, Charité – Universitätsmedizin Berlin,
corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and
Berlin Institute of Health, Berlin, Germany.

29. School of Psychology and Global Brain Health Institute, Trinity College Dublin, Ireland.

30. School of Psychology, Institute for Mental Health, University of Birmingham,
Birmingham, United Kingdom.

31. Oxford Institute of Clinical Psychology Training and Research, University of Oxford,
Oxford, United Kingdom.

32. Department of Education and Psychology, Freie Universität Berlin, Berlin, Germany.

33. Psychology Department, University of Southampton, Southampton, United Kingdom.

34. Department of Psychological Medicine, Section for Eating Disorders, Institute of
Psychiatry, Psychology and Neuroscience, King's College London, London, United
Kingdom.

35. South London and Maudsley NHS Foundation Trust, London, United Kingdom.

36. Clinical and Experimental Sciences, Faculty of Medicine, University of Southampton,
Southampton, United Kingdom.

37. School of Mathematical Sciences, Fudan University, Shanghai, China.

38. Shanghai Center for Mathematical Sciences, Fudan University, Shanghai, China.

39. Donders Institute, Radboud University, Nijmegen, Netherlands.

40. School of Data Science, Fudan University, Shanghai, China.

41. Department of Computer Science, University of Warwick, Coventry, United Kingdom.

42. Zhangjiang Fudan International Innovation Centre, Shanghai, China.

43. Center for Applied Mathematics, Fudan University, Shanghai, China

44. PONS Centre, National Center for Neurologic Disorders, Huashan Hospital, Fudan
University, Shanghai, China.

45. German Centre for Mental Health (DZPG), Berlin-Brandenburg site, Germany.

46. Department of Psychiatry, Cambridge University, United Kingdom.

*Corresponding author:

Gunter Schumann

Email: gunter.schumann1961@gmail.com and gunter.schumann@charite.de

† These authors contributed equally to this work.

Abstract

Linking synaptic-level perturbations to distributed brain-network dynamics remains a central

challenge for understanding and treating mental illness. Although recent whole-brain models

can reproduce individual brain activity patterns, they largely function as descriptive

simulators rather than mechanistic, intervention-capable systems. Here we present an

intervention-capable digital twin of the human brain, integrating individual neuroanatomy

and task-evoked dynamics within a neuronal-scale framework. Individualised digital twin

brains recapitulate a participant-specific compact cortico-subcortical network phenotype

that captures transdiagnostic psychopathology across population and clinical cohorts. In

silico modulation of excitatory and inhibitory synaptic conductance produces bidirectional,

heterogeneous network responses across individuals. Population-scale simulations stratify

individuals and predict longitudinal symptom trajectories from DTB-derived response

profiles. Independent pharmacological functional MRI data further validate the predicted

baseline-dependent network responses in vivo. Together, these findings establish digital

brain models as experimental platforms for mechanistic perturbation, behavioural prediction

and stratification, providing a foundation for precision neuroscience and psychiatry.

Main

As human neuroimaging has reached individual-level precision[1,2], a central challenge has

emerged: how to transform rich multimodal measurements of brain structure and function

into mechanistic models that can be systematically perturbed to generate causal and

predictive insights into highly prevalent mental illness[3,4]. Digital twin models of the human

brain offer a powerful framework for addressing this challenge by integrating neuroanatomy,

physiology, and large-scale functional dynamics within a unified computational system[5-8].

Recent advances have demonstrated that Digital Twin Brain (DTB) framework can

approximate large-scale neuronal dynamics using the biologically constrained spiking neural

networks grounded in individual anatomy and empirical BOLD signals[9,10], establishing the

feasibility of personalised whole-brain simulation. However, existing digital twin brains

primarily function as descriptive simulators rather than enabling systematic mechanistic

perturbation and prediction.

This limitation is especially salient in psychiatry. Mental disorders are characterised by

distributed network alterations[11] and heterogeneous treatment response[12–15]. Neuroimaging

studies have identified reproducible functional connectivity abnormalities across disorders,

often implicating common circuits involved in reward processing, cognitive control, and

salience attribution[15–22]. Yet without the ability to perform controlled perturbations, these

findings remain difficult to translate into mechanistic insight or predictive models of

intervention effects[13,14]. At the same time, most psychiatric treatments act on specific

molecular targets - such as glutamatergic or GABAergic neurotransmission - whose effects

propagate across multiple spatial scales, from synapses to distributed functional brain

networks[23–27]. Bridging these levels of description remains a critical gap.

A promising strategy is to focus on behaviourally grounded, transdiagnostic brain

networks that provide low-dimensional interfaces between cognition, symptoms, and

neurobiology[18–20,28]. Converging evidence suggests that shared dimensions of

psychopathology are underpinned by common neural circuit disruptions that cut across

traditional diagnostic boundaries[17–20,28–30]. The neuropsychopathology (NP) factor, derived

from task-based functional connectivity, captures shared continuous variance across

internalising and externalising symptoms in the population cohort[28]. As such, it provides a

principled target for testing whether digital twin brains can move beyond simulation toward

mechanistic manipulation of behaviourally relevant circuits.

In parallel, a large body of neurochemical evidence implicates disruptions in cortical

excitation–inhibition (E/I) balance in the pathophysiology of mood and addiction disorders.

Magnetic resonance spectroscopy and postmortem studies report reductions in both

glutamatergic and GABAergic signalling across cortex[31–36], while pharmacological agents

targeting these systems - including ketamine and benzodiazepines - can induce rapid

changes in symptoms and functional connectivity[23,26,37–41]. Despite this, the relationship

between neurotransmitter-level perturbations and large-scale functional network

reconfiguration remains poorly understood. Empirical pharmaco-fMRI can quantify

individual-level effects of neurochemical interventions, but offers limited leverage

to mechanistically disentangle or systematically manipulate the underlying synaptic

processes. Existing whole-brain models, in turn, have not yet been configured to support

controlled, neurotransmitter-level perturbations to task-state network reconfiguration at the

individual level[8,42–44].

Here we build on the DTB framework[9,10] to address this unmet need. We tested whether

a neuronal-scale digital twin can reproduce a behaviourally relevant transdiagnostic brain

network (NP factor) and mechanistically manipulate it through abstract synaptic gain

control parameters inspired by E/I mechanisms. We further examined whether such in

silico manipulations yield predictions that generalise across tasks and populations, align

with empirical pharmacological effects, and forecast longitudinal symptom trajectories.

Together, this work marks a step change in digital brain modelling from passive emulation

of neural activity to intervention-capable, mechanistically closed-loop systems that

integrate simulation, perturbation, and prediction.

A compact cortico–subcortical network phenotype of psychopathology.

We first delineated the NP factor, a shared functional connectivity phenotype that captures

psychopathology across symptom domains and cohorts. As in our previous work[28],

conducted in 14 year-old adolescents of the IMAGEN study, we applied connectome-based

predictive modeling to the task-based functional connectomes derived from the monetary

incentive delay and stop-signal task in 19 year old young adults of IMAGEN follow-up 2

(n=1,050, age 18.4±0.7 years, Table S1). Four of six task conditions (stop success and failure

in SST; reward anticipation and positive feedback in MID) significantly predicted a range of

externalising and internalising symptoms at age 19 (Fig.1a; Table S2-3). Across these

predictive task conditions, we identified two transdiagnostic functional connectivity (FC)

profiles: one positive profile of 34 edges where stronger connectivity was associated with

higher symptom burden across domains, and one negative profile of 29 edges where weaker

connectivity was associated with higher burden. These edges were concentrated within and

between default mode, frontoparietal, somatomotor–frontal, limbic, and cerebellum systems

(Fig.1b). To quantify these functional connectivity profiles, we computed positive and

negative NP scores for each individual by summing the strength of edges in each profile.

To test whether the NP scores generalise to clinical populations, we re-computed them from

task-state FC in an independent case–control cohort (STRATIFY, n=434, age 21.8±2.0 years).

Only the negative NP score showed robust case-control differences, with lower scores in

patients than in healthy controls (Fig.1c; Fig.S1). Thus, the negative NP profile defines a

shared network phenotype of transdiagnostic psychopathology.

To characterise the neurobiological context of this negative NP circuit, we examined

whether negative NP-related regions exhibit a distinctive neurochemical signature in

empirical PET receptor maps. Using published PET-derived receptor density maps[45], we

found that NP-related regions showed higher receptor-map values for glutamatergic,

serotonergic, cholinergic, and cannabinoid systems than NP-unrelated regions (Fig.S2, Table

S4, all $P_{corrected} < 0.05$). These receptor systems are primarily involved in regulating cortical

excitatory drive and neuromodulatory control of network dynamics, and are known to

influence E/I balance at the circuit level through modulation of glutamatergic

transmission[46,47], GABA interneuron recruitment[48–50], recurrent coupling[51], and gain control[52]

mechanisms.

To enable computationally tractable digital twin brain (DTB) modeling, we restricted the

negative NP profile to cortico-subcortical edges, excluding the cerebellum and brainstem.

These regions exhibit distinct cytoarchitecture and neuronal dynamics[53] and are not currently

parameterized in our DTB framework (see Methods for details). Finally, we identified a

compact set of 12 task-state functional connections linking regions spanning prefrontal

control, default-mode, sensorimotor, and limbic systems (Fig.1d; Table S5). The summed

strength of these 12 edges remained significantly reduced in patients relative to controls

(Fig.1e; Fig.S1), and served as the final NP factor and target for subsequent DTB simulations

and in-silico E/I modulations.

Individual digital twin brains reveal a bidirectionally E/I-tunable NP factor.

We next asked whether this NP factor can be reconstructed and mechanistically manipulated

in individualised, biologically grounded models. To this end, we applied the previously

established DTB framework[9,10] to generate participant-specific task-state models (Methods;

Fig.2a; Fig.S3). Following a calibration experiment across neuronal resolutions (Fig.S4; Table

S8; more details in Methods and Supplementary Results), we implemented DTBs at three

complementary scales: billion-neuron voxel-wise models to establish biological fidelity and

simulation feasibility, 100-million-neuron models for controlled perturbational exploration,

and computationally tractable regional models for population-level statistical inference.

Across all scales, models were constructed using a common pipeline. The individual voxel-

wise grey and white matter anatomy were first transformed into a multi-population neuronal

network. The activity of each neuron in this network was modelled using the leaky integrate-

and-fire (LIF) model[54], equipped with excitatory (AMPA-mediated) and inhibitory (GABA-A-

mediated) synapses. We then estimated hyperparameters by assimilating empirical BOLD

signals in task-engaged regions (Methods; Table S6) to mimic external current inputs driving

the model from rest to task-states[9,10]. Finally, the voxel-wise neuronal activity (mean firing

rate of neuronal population) was converted into BOLD signals using the Balloon-Windkessel

model[55].

We first selected four participants as illustrative exemplars according to the behavioural

symptoms and negative NP scores, including participants with major depressive disorder

(MDD), alcohol use disorder (AUD), and healthy controls (HC01 and HC02) (demographics

information in Table S7, selection criteria in Methods), to demonstrate the feasibility and

mechanistic properties of neuronal-scale digital twin simulations. These examples are

presented for illustration only; all statistical inferences are based on group-level analyses that

account for site and sex effects (see Supplementary Results). Using a 1-billion-neuron scale

(Fig.S4; Table S8), their individualised DTBs accurately reproduced task-evoked dynamics

in MID and SST. Simulations achieved high voxel-wise correlations with empirical BOLD in

assimilated regions (mean r=0.9, in-sample correlations). Across the whole brain, mean

correlations of four individuals (mean r=0.30-0.44) were clearly above a reference

simulation obtained by a single random parameter shuffle (mean r=0) (Fig.2b-c; Fig.S5-6;

Table S9). These DTBs also produced whole-brain functional connectivity patterns that

closely matched empirical connectomes in both tasks (Fig.2d; Table S9).

Based on the simulated baseline NP configuration, we next examined whether NP circuit

dynamics could be systematically manipulated within the DTB framework. In recurrent

cortical networks, E/I balance emerges from the interaction of inhibitory interneuron

recruitment[56,57], recurrent coupling[51,57], and synaptic gain[58], and is governed by circuit

dynamics rather than receptor density alone. Consistent with this systems-level perspective,

we treated both AMPA- and GABA-A-mediated synaptic conductance as global dynamical

control parameters, enabling controlled perturbation of NP network activity within

individualised digital twins. Notably, publicly available PET datasets did not show

preferential expression of GABA receptor systems in NP-related regions, reinforcing the

interpretation that inhibitory modulation in the model reflects circuit-level dynamical control

rather than regionally localized receptor enrichment.

Accordingly, we varied the synaptic conductance parameters for AMPA and GABA-A

receptors in the 100-million-neuron DTBs of the same illustrative participants. Despite the

reduced resolution, DTBs preserved high fidelity in assimilated regions and acceptable

whole-brain correspondence (Table S10). Moreover, 100-million-neuron DTBs retained

sufficient subject-specific NP-factor phenotypes such that each individual's simulated NP

factor best matched their own empirical value, with significantly lower mean-squared error

for within-subject matches than any cross-subject pairing (permutation tests across 1,575

simulated–empirical pairings, Supplementary Results; Fig.S7), therefore supporting its use for

controlled perturbational experiments. Using these established DTB models, we performed

systematic perturbations by upregulating synaptic conductance for AMPA (0.0020-0.0052)

and GABA-A receptors (0.0015-0.0040) across a predefined parameter grid, guided by

evidence for reduced glutamatergic and GABAergic function in depression[24,25] and

addiction[27,34]. Parameter ranges were constrained by biologically plausible neuronal firing

rates[59], synchronization[60], and simulated BOLD activity. During GABA-A manipulations,

AMPA conductance was fixed at the value that maximized NP factor enhancement in prior

sweeps, allowing controlled assessment of E/I balance and avoiding excessive inhibition (see

more details in Methods).

Virtual modulation of AMPA- and GABA-A-mediated synaptic conductance produced

bidirectional, heterogeneous changes in NP connectivity across illustrative participants.

Specifically, in two patient brain models, both excitatory and inhibitory upregulation

increased NP-factor strength, and in illustrative healthy control models, the same

perturbations yielded divergent responses, with modest increases in one individual and clear

decreases in the other (Fig.2e-f; Tables S11-12). These examples demonstrate that identical

synaptic perturbations can lead to bidirectional changes in NP-factor strength across

individuals.

To test whether this bidirectional response generalises across cognitive contexts, we repeated

the same E/I perturbations in an alternative task (the Emotional Face Task) in the same

participants. Comparable bidirectional modulation effects were observed (Supplementary

Results; Fig.S8; Table S13), indicating that the E/I-sensitive response properties of the NP

circuit are not task-specific, but generalise across distinct cognitive contexts.

Population-scale in silico perturbations attenuate NP abnormalities and stratify individuals.

The single-subject analyses suggest that in silico E/I perturbation can shift NP network toward

the control distribution observed in STRATIFY (reflected by increased NP-factor strength),

although not uniformly across individuals. To further explore whether these effects generalise

across the cohort and relate to behavioural differences, we performed population-scale

simulations, which allow us to link individual susceptibility to excitatory and inhibitory

synaptic perturbations with behavioural symptoms. To enable large-cohort simulations, we

developed regional DTBs that convert regional MRI-derived structural features into spiking

neural networks comprising 3 million neurons (identified from calibration experiment, Fig.S4,

Table S14), with neuron counts allocated proportionally to regional grey matter volume to

preserve anatomical scaling while maintaining computational tractability (Methods). While

voxel-wise DTBs establish mechanistic feasibility at high-fidelity resolution, these regional

DTBs are used to test whether these principles scale to population-level inference.

We constructed individualised DTBs for 72 patients with MDD, 59 with AUD and 69 healthy

controls from STRATIFY, as well as 90 individuals with high externalising and internalising

symptoms from IMAGEN (Table S15). These participants from STRATIFY were included

because they possessed complete, high-quality multimodal imaging (T1-weighted, DTI, fMRI)

and behavioural data. These DTB models reproduced individual task-evoked BOLD signals

with moderate-to-high accuracy during in-sample fitting phase (Fig.3a), generated NP factors

that closely matched empirical values at the group level (Fig.S9), and mirrored case–control

group differences observed in vivo (Fig.3b; Fig.S9). As an additional validation beyond NP

circuit, simulated whole-brain FC also predicted task-related behavioural performances and,

on average, outperformed empirical FC (Supplementary Results; Fig.S10; Table S16),

suggesting that DTBs distill behaviourally relevant variance that is only partially captured by

conventional functional connectivity.

Motivated by these converging validations, we then applied the same AMPA and GABA-A

neuromodulation schemes used in the 100-million-neuron DTBs to all 290 participants. At

the group level, both excitatory and inhibitory upregulation shifted NP factors in patients

toward values observed in controls, in other words, after either AMPA or GABA-A

modulation, previously significant differences between patients, high-symptom individuals

and healthy controls were no longer significant (Fig.3c-d). Within each clinical group, NP

factors increased on average following both AMPA and GABA-A manipulations (Fig.3e; Fig.

S11).

Despite clear group-level increases, individual responses were heterogeneous and often

bidirectional (Fig.3f). Following AMPA modulation, both patients and controls showed

increased NP factor on group average, but the proportion of "increasers" (ΔNP > 0) was

substantially higher among patients than controls (Table S17). Baseline simulated NP factor

was strongly and inversely correlated with AMPA-induced change (Fig.3g), such that

individuals with lower baseline connectivity, typically patients, exhibited the largest gains,

whereas those with high baseline NP showed modest increases or decreases. GABA-A

modulation produced a similar, though weaker, negative association (Fig.3h). Importantly,

these baseline–ΔNP couplings significantly exceeded a null distribution generated by

permuting ΔNP values across individuals (10,000 permutations, two-sided $P<0.0001$; see

Methods), indicating that the relationship reflects structured baseline-dependent dynamics

rather than statistical regression to the mean.

To facilitate clinical interpretation of this response heterogeneity, we descriptively stratified

individuals based on the direction of their joint responses to AMPA and GABA-A

perturbations. Participants whose NP factors increased under both manipulations were

classified as "increased responders", whereas those with at least one decrease were classified

as "decreased responders". This stratification was used for visualization and group-level

comparison only; the underlying baseline–ΔNP relationships were treated as continuous

throughout statistical analyses. Increased responders were markedly more prevalent among

patients than among high-symptom individuals or healthy controls (Fig.3i) and did not differ

systematically by site or sex (Tables S18–19; Supplementary Results). Clinically, the

increased responders, characterised by lower simulated baseline NP factors, showed greater

symptom burden, including more panic attacks and negative affect, and reduced social

support, compared with the decreased responders after controlling for sex, site, and head

motion (Fig.3j; Table S20).

Together, these findings indicate that DTB-derived virtual responses offer an individualised,

mechanistically interpretable stratification, whereby different responses in NP circuit map

onto meaningful differences in clinical and behavioural profiles. A critical question, however,

is whether these virtual baseline-dependent, bidirectional response structures also emerge

under real pharmacological perturbations in vivo.

Pharmacological E/I perturbation in vivo mirrors digital twin predictions of NP

modulations.

To address this, we analysed pharmacological fMRI data from 27 healthy male participants

(27.3±6.2 years), who performed the MID task following ketamine (0.25 mg/kg), midazolam

(0.03 mg/kg), or placebo administration in a randomized, single-blinded, three-way

crossover design (Fig.4a). Ketamine, an NMDA receptor antagonist, reduces inhibitory

interneuron activity and indirectly enhances AMPA-mediated excitatory transmission[23,61],

whereas midazolam, a GABA-A receptor positive allosteric modulator, increases inhibitory

synaptic conductance and dampens network excitability[62,63]. Because pharmacological data

were only available for the MID task-fMRI, we focused on a MID-specific NP connectivity,

defined as the summed strength of the six MID edges that contributed to the NP circuit.

Placebo served as the empirical analogue of the DTB baseline simulation, whereas ketamine

and midazolam provided in vivo perturbations with predominant effects on excitatory-

leaning and inhibitory neurotransmission, respectively. We tested whether these

perturbations reproduce the baseline-dependent response structures predicted by AMPA and

GABA-A modulations in silico.

At the group level, neither ketamine nor midazolam induced a significant change in summed

NP-related MID connectivity relative to placebo (Fig.S12), suggesting that conventional

group-mean analyses would miss potential relationship between brain network configuration

and pharmacological response. However, individual-level analyses revealed a bidirectional,

baseline-dependent pattern closely resembling DTB simulations. Unsupervised k-means

clustering of combined ketamine- and midazolam-induced FC changes identified two

subgroups (Fig.S12). In Group 1 (n=19), both ketamine and midazolam significantly

increased NP-related connectivity after adjusting for head motion effects (Fig.4b-c), whereas

in Group 2 (n=8), both drugs significantly decreased it (Fig. 4d-e). Baseline NP-related

connectivity under placebo was significantly lower in the increased-response group than in

the decreased-response group (Fig.4f), and baseline FC was inversely related to drug-induced

change across individuals (Fig.4g-h). This inverse baseline–ΔFC relationship significantly

exceeded a permutation-derived null distribution generated by shuffling drug-induced

changes across individuals (10,000 permutations; midazolam: two-sided P=0.0056,

ketamine: two-sided P=0.069, showing a trend in the same direction; see Methods),

indicating that the bidirectional response is unlikely to reflect regression to the mean alone.

Thus, pharmacological E/I perturbation in vivo reproduces the bidirectional, baseline-

dependent modulation of NP circuit predicted in silico by DTBs.

To match the pharmacological dataset (MID only), we restricted virtual modulation to MID-

specific NP connectivity to place the model and pharmacological data in the same task

context. Importantly, a bidirectional, baseline-dependent response toward virtual

modulation of MID-specific NP connectivity was preserved (Supplementary Results; Fig.

S13), indicating that the core dynamic property of the NP circuit is robust to this task

restriction.

We then asked whether a quantitative mapping learned entirely in silico could generalise to

predict individual pharmacological responses. In DTBs, we first estimated a linear mapping

between simulated baseline MID FC and virtual modulation-induced changes in NP-related

connectivity. Applying this same mapping to empirical placebo MID FC, we obtained

subject-specific predictions of ketamine- and midazolam-induced FC changes. Predicted

and observed drug-induced changes closely matched at the individual level (Fig.4i-j) and

stratified participants with opposite response directions (AUC of ketamine=0.69; AUC of

midazolam=0.83; Fig.4k), further supporting the translational relevance of DTB-derived

virtual modulation.

Together, these findings show that DTBs not only recapitulate individual-level, baseline-

dependent response structure observed under pharmacological perturbation, but also

provide a mechanistic template for forecasting individual drug-induced circuit shifts from

baseline network configuration.

DTB-derived perturbational responses predicts future changes in symptoms.

We next asked whether DTB-derived perturbational responses carry prospective information

about longer-term symptom trajectories. Focusing on internalising symptoms (given the

limited prevalence of externalising symptoms at age 23), we first learn a linear mapping from

empirical NP connectivity at age 19 to concurrent internalising symptom scores (Fig.4l). We

then applied this mapping to simulated NP connectivity before and after virtual E/I

modulation to obtain subject-specific predicted symptom changes, termed the "behavioural

restoration" index.

We quantified observed symptom change over four years (Δbehaviour, ages 19–23) and

tested whether the DTB-derived "behavioural restoration" index predicted these longitudinal

changes. Across individuals, baseline symptoms and the restoration index for AMPA

modulation together explained ~23% of the variance in the symptom changes (Fig.4m). The

restoration index alone was significantly correlated with empirical symptoms changes

(Fig.S14), indicating that individuals whose NP circuit was more "normalisable" in silico

were less likely to deteriorate and more likely to remit. Importantly, the restoration index

contributed unique predictive value beyond baseline symptoms, significantly increasing

explained variance by ~5% relative to a symptom-only model (Fig.S14). However, the

behavioural restoration index for GABA-A modulation was not correlated with empirical

symptom changes (Fig. S14).

These findings suggest that DTB-derived perturbational response metrics capture not only

cross-sectional heterogeneity but also clinically relevant dynamical properties that forecast

longitudinal trajectories. Taken together with the pharmacological validation above, these

results position DTB models as a promising tool for predicting and stratifying individualised

responses to neurotransmitter-targeted interventions, with potential implications for the

development and personalization of future psychiatric treatments.

Discussion

We simulated a transdiagnostic, behaviourally relevant task-evoked functional network

phenotype (i.e., NP factor) within individualised neuronal-scale digital twin brain models.

By systematically perturbing AMPA- and GABA-A–mediated synaptic gain, we quantified

individual-specific, bidirectional network responses and linked these virtual response

profiles to both pharmacological fMRI effects and longitudinal symptom trajectories. These

results demonstrate that digital brain models can move beyond descriptive simulation of

neural activity to perform controlled in silico perturbations for mechanistic interrogation and

individualised counterfactual prediction. By introducing biologically interpretable synaptic

parameters, our framework allows precise manipulation of excitation–inhibition balance at

the neuronal scale while preserving each individual's anatomical and task-evoked

constraints, establishing a new class of perturbational and predictive digital brain models for

computational neuroscience and psychiatry.

The NP factor represents a compact functional connectivity pattern that relates to a broad

spectrum of internalising and externalising symptoms across heterogeneous populations. It

is characterised by reduced functional integration among limbic, default-mode,

frontoparietal control, and cingulo-opercular networks, which jointly support reward

processing, cognitive control, and attention allocation[64–67]. Disruptions within and between

these networks have been reported across mood, anxiety, and substance use disorders, as

well as in population-based samples spanning the full range of symptom severity[18–20,68]. In

this sense, the NP factor reflects a shared circuit phenotype rather than a disorder-specific

marker. Importantly, we treat NP as a continuous circuit dimension rather than a diagnostic

classifier. This makes it suitable as a perturbational readout: it is sufficiently low-dimensional

to allow explicit predictions about network reconfiguration, while remaining anchored in

identifiable large-scale functional systems.

A central contribution of the NP-DTB framework is that it renders a descriptive task-evoked

connectivity phenotype experimentally manipulable. Conventional neuroimaging analyses

typically relate connectivity measures to behavioural symptoms, but they do not permit

controlled intervention at the circuit level[69,70]. By contrast, DTBs establish an explicit link

between large-scale functional network organization and synaptic-scale parameters,

enabling in silico perturbation of defined excitatory and inhibitory mechanisms. In addition,

DTBs reproduce individualised, task-evoked functional connectivity patterns that are

challenging for the conventional whole-brain models, which often rely on coarse-grained

regional parcellations and focus primarily on resting-state dynamics[71,72]. This task-state

emphasis is critical, as many clinically relevant circuit abnormalities are context

dependent[20,73], and network configurations expressed during reward processing or inhibitory

control cannot be inferred from resting-state activity alone[74]. By assimilating task-state signals

and applying perturbations at the synaptic level, DTBs generate individual-specific

predictions of how functional circuits respond to mechanistically interpretable excitatory or

inhibitory modulation.

Using this framework, we observed a pronounced baseline-dependent heterogeneity in NP

connectivity response to virtual E/I perturbations. Individuals with lower baseline NP factor

scores tend to show larger increases following perturbation, whereas those with higher

baseline NP show modest increases or even decreases. Interestingly, both excitatory-leaning

(AMPA) and inhibitory-leaning (GABA-A) perturbations often shifted NP connectivity in the

same direction within a given individual model. This convergence does not contradict

excitation–inhibition balance; rather, it reflects the nonlinear, regime-dependent dynamics

of large-scale recurrent networks, in which multiple control parameters can move the system

toward a balanced operating regime depending on baseline state[75–77], consistent with prior

empirical and computational modeling studies[57,78–81]. Notably, the bidirectional convergence

of excitatory- and inhibitory-leaning perturbations at the network level does not imply

equivalence of their molecular mechanisms, but instead reflects compensatory and

nonlinear circuit dynamics in which E/I balance acts as a key control variable[57,79,80].

Two independent lines of evidence further support the biological plausibility of the causal

links between microscale neurotransmitter systems and large-scale functional networks

inferred by the DTB models. First, PET-derived receptor maps show that regions contributing

to the NP circuit are enriched in receptor systems involved in regulating excitation–inhibition

balance[46,49,51,52], providing a neurochemical context for why E/I perturbations plausibly

modulate this circuit, although these maps are group-level and cannot establish individual

causality. Second, pharmacological fMRI data provide an independent empirical

perturbation reference for examining how excitatory and inhibitory neurotransmitter

modulation affects NP-related connectivity. Although ketamine and midazolam did not

produce robust group-mean effects on NP-related connectivity, individual-level analyses

revealed bidirectional, baseline-dependent responses that closely mirror DTB predictions.

The convergence between in silico perturbations and pharmacological modulation in vivo

suggests that heterogeneity in NP network responses may reflect differences in baseline

circuit state, a property also evident in pharmacological responses, rather than an artifact of

the digital twin models.

Therefore, a key future implication of the DTB framework is that an individual's baseline

functional network configuration carries information about the direction and magnitude of

network change under perturbation. This is supported by our analyses showing a significant

inverse relationship between baseline NP connectivity and modulation-induced ΔNP in both

AMPA- and GABA-A–mediated virtual perturbations, as well as in pharmacological fMRI

data (Fig. 3g–h; Fig. 4g-k). Depending on their initial circuit state, the same intervention may

increase network integration in some individuals while decreasing it in others. In this work,

we illustrate the translational potential of this approach in two ways. First, we learned a

mapping from baseline NP connectivity to virtual perturbation responses in silico and

applied it out-of-sample to forecast drug-induced circuit changes in vivo. Second, by linking virtual NP network changes to behaviour, we define a "behavioural restoration" index that quantifies an individual's brain circuit susceptibility to intervention. This index predicts differences in future symptom trajectories beyond what baseline symptom severity alone can explain, providing a measurable, testable readout of circuit "normalisability" based directly on the model's simulated perturbation responses. Beyond prediction, the DTB framework also supports mechanistic stratification based on baseline brain network configuration rather than symptom severity alone. Individuals with similar symptom burden but distinct NP configurations can follow divergent trajectories over time, offering a mechanistic explanation for why symptom-based stratification often fails to anticipate treatment response. However, we should note that the excitatory and inhibitory perturbations implemented in the DTB represent controlled shifts in synaptic gain that move the system across distinct network operating regimes, rather than modeling reality drug actions. Thus, this predictive framework is intended for risk stratification and mechanistic insight, rather than deterministic individual-level prognosis.

Notably, model-empirical similarity increased with neuronal resolution in voxel-wise simulations, consistent with scaling properties of the previous studies[9,10]. Regional DTBs showed higher similarity in aggregated BOLD dynamics but lower correspondence in functional connectivity relative to voxel-wise models, likely reflecting reduced spatial granularity for capturing individual-level network structure. Nevertheless, regional models remained computationally tractable and retained sufficient fidelity to predict task performance and empirical group-level differences, supporting their use for population-scale perturbational analyses. Rather than serving as exact replicas of biological brains, DTBs

should be viewed as experimentally controllable models that enable causal hypothesis

testing in systems otherwise inaccessible to direct manipulation.

Several limitations should be noted. DTB inferences depend on simplifying modelling

assumptions, e.g., fixed excitatory-inhibitory neuron ratios, homogeneous synaptic

architecture, and simplified neuronal dynamics. Accordingly, AMPA- and GABA-A–labelled

parameters should be interpreted as phenomenological gain parameters rather than direct

mapping to drug mechanisms. The present DTB framework does not incorporate the full

range of neurotransmitter systems, nor cerebellar microcircuit dynamics, and

pharmacological validation was limited to a single task and cohort. Although PET receptor

maps provide biological plausibility, they do not capture individual neurochemical

variability. Despite these constraints, the convergent evidence suggests that a compact task-

evoked circuit phenotype can be linked to explicit synaptic control parameters in

individualised models, yielding baseline-dependent, and bidirectional predictions that align

with pharmacological perturbations and carry prognostic information. Together, these

findings motivate a shift from descriptive biomarkers toward perturbational models that

explain and predict individual differences in psychiatric trajectories.

References

1.   Dubois, J. & Adolphs, R. Building a Science of Individual Differences from fMRI.

Trends Cogn. Sci. **20**, 425–443 (2016).

2.   Bassett, D. S. & Sporns, O. Network neuroscience. *Nat. Neurosci.* **20**, 353–364

(2017).

3.   Sejnowski, T. J., Churchland, P. S. & Movshon, J. A. Putting big data to good use in

neuroscience. *Nat. Neurosci.* **17**, 1440–1441 (2014).

4.   Huys, Q. J. M., Maia, T. V. & Frank, M. J. Computational psychiatry as a bridge from

neuroscience to clinical applications. *Nat. Neurosci.* **19**, 404–413 (2016).

5.   Lee, J. H., Liu, Q. & Dadgar-Kiani, E. Solving brain circuit function and dysfunction

with computational modeling and optogenetic fMRI. *Science* **378**, 493–499 (2022).

6.   Jirsa, V. K., Sporns, O., Breakspear, M., Deco, G. & McIntosh, A. R. Towards The

Virtual Brain: network modeling of the intact and the damaged brain. *Arch. Ital. Biol.* **148**,

189–205 (2010).

7.   Deco, G. & Kringelbach, M. L. Great expectations: using whole-brain computational

connectomics for understanding neuropsychiatric disorders. *Neuron* **84**, 892–905 (2014).

8.   Shine, J. M. *et al.* Computational models link cellular mechanisms of

neuromodulation to large-scale neural dynamics. *Nat. Neurosci.* **24**, 765–776 (2021).

9.   Lu, W. *et al.* Imitating and exploring human brain's resting and task-performing states

via resembling brain computing: scaling and architecture. *Natl. Sci. Rev.* nwae080 (2024)

doi:10.1093/nsr/nwae080.

10.   Lu, W. *et al.* Simulation and assimilation of the digital human brain. *Nat. Comput.*

Sci. **4**, 890–898 (2024).

11.   Braun, U. *et al.* From Maps to Multi-dimensional Network Mechanisms of Mental

Disorders. *Neuron* **97**, 14–31 (2018).

12.   Global, regional, and national burden of 12 mental disorders in 204 countries and

territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019.

Lancet Psychiatry **9**, 137–150 (2022).

13.   Patel, V. *et al.* The Lancet Commission on global mental health and sustainable

development. *The Lancet* **392**, 1553–1598 (2018).

14.   McIntyre, R. S. *et al.* Treatment-resistant depression: definition, prevalence, detection,

management, and investigational interventions. *World Psychiatry Off. J. World Psychiatr.*

Assoc. WPA **22**, 394–412 (2023).

15.   Kotov, R. *et al.* The Hierarchical Taxonomy of Psychopathology (HiTOP): A

dimensional alternative to traditional nosologies. *J. Abnorm. Psychol.* **126**, 454–477 (2017).

16.   Insel, T. R. The NIMH Research Domain Criteria (RDoC) Project: precision medicine

for psychiatry. *Am. J. Psychiatry* **171**, 395–397 (2014).

17.   Caspi, A. *et al.* The p Factor: One General Psychopathology Factor in the Structure

of Psychiatric Disorders? *Clin. Psychol. Sci. J. Assoc. Psychol. Sci.* **2**, 119–137 (2014).

18.   Vanes, L. D. & Dolan, R. J. Transdiagnostic neuroimaging markers of psychiatric risk:

A narrative review. *NeuroImage Clin.* **30**, 102634 (2021).

19.   Goodkind, M. *et al.* Identification of a Common Neurobiological Substrate for Mental

Illness. *JAMA Psychiatry* **72**, 305–315 (2015).

20.   McTeague, L. M. *et al.* Identification of Common Neural Circuit Disruptions in

Cognitive Control Across Psychiatric Disorders. *Am. J. Psychiatry* **174**, 676–685 (2017).

21.   Zilverstand, A., Huang, A. S., Alia-Klein, N. & Goldstein, R. Z. Neuroimaging

Impaired Response Inhibition and Salience Attribution in Human Drug Addiction: A

Systematic Review. *Neuron* **98**, 886–903 (2018).

22.   Russo, S. J. & Nestler, E. J. The brain reward circuitry in mood disorders. *Nat. Rev.*

Neurosci. **14**, 609–625 (2013).

23.   Zanos, P. & Gould, T. D. Mechanisms of Ketamine Action as an Antidepressant. *Mol.*

Psychiatry **23**, 801 (2018).

24.   Duman, R. S., Sanacora, G. & Krystal, J. H. Altered connectivity in depression: GABA

and glutamate neurotransmitter deficits and reversal by novel treatments. *Neuron* **102**, 75–

90 (2019).

25.   Hu, Y.-T., Tan, Z.-L., Hirjak, D. & Northoff, G. Brain-wide changes in excitation-

inhibition balance of major depressive disorder: a systematic review of topographic patterns

of GABA- and glutamatergic alterations. *Mol. Psychiatry* **28**, (2023).

26.   Ren, Z. *et al.* Bidirectional Homeostatic Regulation of a Depression-Related Brain

State by Gamma-Aminobutyric Acidergic Deficits and Ketamine Treatment. *Biol. Psychiatry*

80, 457–468 (2016).

27.   Biria, M. *et al.* Cortical glutamate and GABA are related to compulsive behaviour in

individuals with obsessive compulsive disorder and healthy controls. *Nat. Commun.* **14**,

3324 (2023).

28.   Xie, C. *et al.* A shared neural basis underlying psychiatric comorbidity. *Nat. Med.* **29**,

1232–1242 (2023).

29.   McTeague, L. M., Goodkind, M. S. & Etkin, A. Transdiagnostic impairment of

cognitive control in mental illness. *J. Psychiatr. Res.* **83**, 37–46 (2016).

30.   Husain, M. & Roiser, J. P. Neuroscience of apathy and anhedonia: a transdiagnostic

approach. *Nat. Rev. Neurosci.* **19**, 470–484 (2018).

31.   Ritter, C. *et al.* Evaluation of Prefrontal γ-Aminobutyric Acid and Glutamate Levels

in Individuals With Major Depressive Disorder Using Proton Magnetic Resonance

Spectroscopy. *JAMA Psychiatry* **79**, 1209–1216 (2022).

32.   Marinkovic, K., Alderson Myers, A. B., Arienzo, D., Sereno, M. I. & Mason, G. F.

Cortical GABA levels are reduced in young adult binge drinkers: Association with recent

alcohol consumption and sex. *NeuroImage Clin.* **35**, 103091 (2022).

33.   Prévot, T. & Sibille, E. Altered GABA-mediated information processing and cognitive

dysfunctions in depression and other brain disorders. *Mol. Psychiatry* **26**, 151–167 (2021).

34.   Zhou, H. *et al.* Glutamate concentration of medial prefrontal cortex is inversely

associated with addictive behaviors: a translational study. *Transl. Psychiatry* **14**, 433 (2024).

35.   Godfrey, K. E. M., Gardner, A. C., Kwon, S., Chea, W. & Muthukumaraswamy, S. D.

Differences in excitatory and inhibitory neurotransmitter levels between depressed patients

and healthy controls: A systematic review and meta-analysis. *J. Psychiatr. Res.* **105**, 33–44

(2018).

36.   Sanacora, G., Treccani, G. & Popoli, M. Towards a glutamate hypothesis of

depression: An emerging frontier of neuropsychopharmacology for mood disorders.

Neuropharmacology **62**, 63–77 (2012).

37.   Grabski, M. *et al.* Adjunctive Ketamine With Relapse Prevention–Based

Psychological Therapy in the Treatment of Alcohol Use Disorder. *Am. J. Psychiatry* **179**,

152–162 (2022).

38.   Dubovsky, S. L. & Marshall, D. Benzodiazepines Remain Important Therapeutic

Options in Psychiatric Practice. *Psychother. Psychosom.* **91**, 307–334 (2022).

39.   Forsyth, A. *et al.* Comparison of local spectral modulation, and temporal correlation,

of simultaneously recorded EEG/fMRI signals during ketamine and midazolam sedation.

Psychopharmacology (Berl.) **235**, 3479–3493 (2018).

40.   Abdallah, C. G. *et al.* Ketamine Treatment and Global Brain Connectivity in Major

Depression. *Neuropsychopharmacology* **42**, 1210–1219 (2017).

41.   Kraguljac, N. V. *et al.* Ketamine modulates hippocampal neurochemistry and

functional connectivity: a combined magnetic resonance spectroscopy and resting-state

fMRI study in healthy volunteers. *Mol. Psychiatry* **22**, 562–569 (2017).

42.   Breakspear, M. Dynamic models of large-scale brain activity. *Nat. Neurosci.* **20**,

340–352 (2017).

43. Wang, H. *et al.* Virtual brain twins: from basic neuroscience to clinical use. *Natl. Sci.*

Rev. **11**, (2024).

44. Griffiths, J. D., Bastiaens, S. P. & Kaboodvand, N. Whole-Brain Modelling: Past,

Present, and Future. in *Computational Modelling of the Brain: Modelling Approaches to Cells,*

Circuits and Networks (eds. Giugliano, M., Negrello, M. & Linaro, D.) 313–355 (Springer

International Publishing, Cham, 2022). doi:10.1007/978-3-030-89439-9_13.

45. Hansen, J. Y. *et al.* Mapping neurotransmitter systems to the structural and functional

organization of the human neocortex. *Nat. Neurosci.* **25**, 1569–1581 (2022).

46. Brunel, N. Dynamics of sparsely connected networks of excitatory and inhibitory

spiking neurons. *J. Comput. Neurosci.* **8**, 183–208 (2000).

47. Swanson, C. J. *et al.* Metabotropic glutamate receptors as novel targets for anxiety

and stress disorders. *Nat. Rev. Drug Discov.* **4**, 131–144 (2005).

48. Puig, M. V., Watakabe, A., Ushimaru, M., Yamamori, T. & Kawaguchi, Y. Serotonin

Modulates Fast-Spiking Interneuron and Synchronous Activity in the Rat Prefrontal Cortex

through 5-HT1A and 5-HT2A Receptors. *J. Neurosci.* **30**, 2211–2222 (2010).

49. Celada, P., Puig, M. V. & Artigas, F. Serotonin modulation of cortical neurons and

networks. *Front. Integr. Neurosci.* **7**, 25 (2013).

50. William Moreau, A., Amar, M., Le Roux, N., Morel, N. & Fossier, P. Serotoninergic

Fine-Tuning of the Excitation–Inhibition Balance in Rat Visual Cortical Networks. *Cereb.*

Cortex **20**, 456–467 (2010).

51.   Hasselmo, M. E. The role of acetylcholine in learning and memory. *Curr. Opin.*

Neurobiol. **16**, 710–715 (2006).

52.   Wilson, R. I. & Nicoll, R. A. Endocannabinoid signaling in the brain. *Science* **296**,

678–682 (2002).

53.   Kandel, E. R., Koester, J. D., Mack, S. H. & Siegelbaum, S. A. *Principles of Neural*

Science, Sixth Edition. (McGraw Hill LLC, 2021).

54.   Abbott, L. F. Lapicque's introduction of the integrate-and-fire model neuron (1907).

Brain Res. Bull. **50**, 303–304 (1999).

55.   Friston, K. J., Mechelli, A., Turner, R. & Price, C. J. Nonlinear responses in fMRI: the

Balloon model, Volterra kernels, and other hemodynamics. *NeuroImage* **12**, 466–477 (2000).

56.   Isaacson, J. S. & Scanziani, M. How Inhibition Shapes Cortical Activity. *Neuron* **72**,

231–243 (2011).

57.   Deco, G. *et al.* How Local Excitation–Inhibition Ratio Impacts the Whole Brain

Dynamics. *J. Neurosci.* **34**, 7886–7898 (2014).

58.   Haider, B. & McCormick, D. A. Rapid neocortical dynamics: cellular and network

mechanisms. *Neuron* **62**, 171–189 (2009).

59.   Yu, Y. *et al.* A 3D atlas of functional human brain energetic connectome based on

neuropil distribution. *Cereb. Cortex* **33**, 3996–4012 (2023).

60.   Baroni, F. & Fulcher, B. Synchrony, oscillations, and phase relationships in collective

neuronal activity: A highly comparative overview of methods. *PLOS Comput. Biol.* **21**,

e1013597 (2025).

61.   Suzuki, A., Hara, H. & Kimura, H. Role of the AMPA receptor in antidepressant

effects of ketamine and potential of AMPA receptor potentiators as a novel antidepressant.

Neuropharmacology **222**, 109308 (2023).

62.   Dakwar, E. *et al.* A Single Ketamine Infusion Combined With Motivational

Enhancement Therapy for Alcohol Use Disorder: A Randomized Midazolam-Controlled Pilot

Trial. *Am. J. Psychiatry* **177**, 125–133 (2020).

63.   Riss, J., Cloyd, J., Gates, J. & Collins, S. Benzodiazepines in epilepsy: pharmacology

and pharmacokinetics. *Acta Neurol. Scand.* **118**, 69–86 (2008).

64.   Haber, S. N. & Knutson, B. The Reward Circuit: Linking Primate Anatomy and

Human Imaging. *Neuropsychopharmacology* **35**, 4–26 (2010).

65.   Buckner, R. L., Andrews-Hanna, J. R. & Schacter, D. L. The brain's default network:

anatomy, function, and relevance to disease. *Ann. N. Y. Acad. Sci.* **1124**, 1–38 (2008).

66.   Cole, M. W. *et al.* Multi-task connectivity reveals flexible hubs for adaptive task

control. *Nat. Neurosci.* **16**, 1348–1355 (2013).

67.   Power, J. D. *et al.* Functional network organization of the human brain. *Neuron* **72**,

665–678 (2011).

68.   Lynch, C. J. *et al.* Frontostriatal salience network expansion in individuals in

depression. *Nature* **633**, 624–633 (2024).

69.   Reid, A. T. *et al.* Advancing functional connectivity research from association to

causation. *Nat. Neurosci.* **22**, 1751–1760 (2019).

70.   Friston, K. J. Functional and Effective Connectivity: A Review. *Brain Connect.* **1**, 13–

36 (2011).

71.   Jirsa, V. *et al.* Personalised virtual brain models in epilepsy. *Lancet Neurol.* **22**, 443–

454 (2023).

72.   Wang, P. *et al.* Inversion of a large-scale circuit model reveals a cortical hierarchy in

the dynamic resting human brain. *Sci. Adv.* **5**, eaat7854 (2019).

73.   Stringaris, A. *et al.* The Brain's Response to Reward Anticipation and Depression in

Adolescence: Dimensionality, Specificity, and Longitudinal Predictions in a Community-

Based Sample. *Am. J. Psychiatry* **172**, 1215–1223 (2015).

74.   Cole, M. W., Ito, T., Cocuzza, C. & Sanchez-Romero, R. The Functional Relevance

of Task-State Functional Connectivity. *J. Neurosci.* **41**, 2684–2702 (2021).

75.   Ozeki, H., Finn, I. M., Schaffer, E. S., Miller, K. D. & Ferster, D. Inhibitory

stabilization of the cortical network underlies visual surround suppression. *Neuron* **62**, 578–

592 (2009).

76.   Murphy, B. K. & Miller, K. D. Balanced Amplification: A New Mechanism of

Selective Amplification of Neural Activity Patterns. *Neuron* **61**, 635–648 (2009).

77.   Joglekar, M. R., Mejias, J. F., Yang, G. R. & Wang, X.-J. Inter-areal Balanced

Amplification Enhances Signal Propagation in a Large-Scale Circuit Model of the Primate

Cortex. *Neuron* **98**, 222-234.e8 (2018).

78.   Murray, J. D. *et al.* Linking Microcircuit Dysfunction to Cognitive Impairment: Effects

of Disinhibition Associated with Schizophrenia in a Cortical Working Memory Model. *Cereb.*

Cortex **24**, 859–872 (2014).

79.   Yizhar, O. *et al.* Neocortical excitation/inhibition balance in information processing

and social dysfunction. *Nature* **477**, 171–178 (2011).

80.   Ahmadian, Y. & Miller, K. D. What is the dynamical regime of cerebral cortex?

Neuron **109**, 3373–3391 (2021).

81.   Bradley, C., Nydam, A. S., Dux, P. E. & Mattingley, J. B. State-dependent effects of

neural stimulation on brain function and cognition. *Nat. Rev. Neurosci.* **23**, 459–475 (2022).

Figure 1 Identification of a compact functional network phenotype of transdiagnostic psychopathology.

a, Task-based fMRI data and behavioural symptom scores from the IMAGEN cohort (n = 1,050) were analysed using connectome-based predictive modelling (CPM) to identify task-

evoked functional connectivity (FC) profiles associated with transdiagnostic

psychopathology. Predictive performances of task-specific FC derived from the Stop-Signal

Task (SST) and Monetary Incentive Delay (MID) were shown in a bubble plot, where bubble

size reflects Spearman's correlation between predicted and empirical symptom scores. The

mean P values across 1,000 repetitions were corrected for multiple comparisons across 36

models using false discovery rate (FDR) correction, corresponding to a minimum significant

effect size of Spearman's $\rho = 0.07$. Task-specific functional connectivity (FC) matrices were

then selected if they showed significant prediction performance for at least three behavioural

symptom domains. SST conditions included stop-success, stop-failure, and go-wrong; MID

conditions included positive feedback, negative feedback, and reward anticipation.

Behavioural measures comprised two externalising domains (attention-deficit/hyperactivity

disorder, ADHD, and conduct disorder, CD) and four internalising domains (eating disorder,

ED, specific phobia, SP, generalized anxiety disorder, GAD, and depression, DEP). A

transdiagnostic FC profile was identified that significantly predicted both externalising and

internalising symptoms. These transdiagnostic associated edges were divided into positive

(pink) or negative (blue) FC profiles based on the direction of their associations with

behavioural symptoms. **b,** Heatmaps show the canonical functional network affiliations of

positive and negative transdiagnostic FC profiles. Positively predictive connections were

primarily distributed across the default mode network (DMN), frontoparietal network (FPN),

somatomotor–frontal (SMF) network, and cerebellum, whereas negatively predictive

connections were concentrated in limbic, DMN, FPN, SMF, and cerebellar systems. **c,**

Negative NP scores (summed strength of the negative FC profile) were compared between

healthy controls (HC, n = 225) and patients (n = 209) from the STRATIFY cohort. Group

difference was assessed using an independent-samples t test with Bonferroni correction ($t =$

3.94, $P_{corrected}$ = 0.0003). Values represent residuals after regressing out sex, scanning site,

and head motion (mean framewise displacement). Violin plots show kernel density estimates

with embedded boxplots indicating the median and interquartile range (IQR); whiskers

extend to 1.5× IQR. **d,** After excluding cerebellar and brainstem connections, the negative

FC profile was reduced to a compact 12-edge NP circuit comprising cortico–subcortical

connections. Edges in the circos plot represent connection strength and negative associations

with psychopathology; corresponding brain regions are shown on cortical and subcortical

renderings. **e,** Individual summed 12-edge-NP factor scores were compared between HC

and patients from the STRATIFY cohort. Group difference was confirmed using an

independent-samples *t* test with Bonferroni correction (*t* = 3.27, $P_{corrected}$ = 0.0036). Scores

represent residuals after regressing out sex, scanning site, and head motion. Each dot denotes

one participant; black markers and vertical lines indicate group means and standard

deviations, respectively. Kernel density estimates illustrate the distributional shift in NP factor

scores between groups. **, $P_{corrected}$ < 0.01; ***, $P_{corrected}$ < 0.001.

Figure 2 Individualised digital twin brains (DTBs) reveal excitation/inhibition-sensitive NP

modulation.

a, Computational workflow for constructing individualised DTBs from multi-modal participant-specific MRI data. DTB model received assimilation hyperparameters as external current inputs and generated simulated neuronal firing rates, which were converted to BOLD signals using Balloon model. The simulation performance was assessed by Pearson's *r* between empirical and simulated BOLD signals. **b,** Radar plots showing simulation performances of 1-billion-neuron models for two patients (one MDD; one AUD) and two healthy controls (HC01; HC02). For each task, both the mean voxel-wise Pearson's correlations in assimilated voxels (assimilated) and the mean correlations across the whole voxels (whole). Task-specific FC matrices included stop-success and stop-failure for the SST, and reward anticipation and positive feedback for the MID task. **c,** Voxel-wise Pearson's correlations between empirical and simulated BOLD signals from the 1-billion-neuron DTB model for the MID and SST tasks of HC01. Bright yellow regions indicate higher similarity. **d,** Empirical and simulated FC matrices across four task conditions of SST and MID from the 1-billion-neuron DTB model of the HC01. **e–f,** In silico E/I modulation sweeps. Curves depict NP factor responses to systematic changes in AMPA (**e**) and GABA-A (**f**) synaptic conductance. For AMPA modulation, NP factor scores increased in patients and HC02 but decreased in HC01. For GABA-A modulation (with AMPA = 0.0044, the parameter yielding maximal NP factor improvement), the NP factor scores increased in two patients, while slightly decreased in HCs. The shaded areas indicate optimal E/I parameters for future perturbations (AMPA = 0.0044; GABA-A = 0.0040). **, $P_{corrected} < 0.01$.

Figure 3 Population-scale in silico perturbations and mechanistically grounded

stratification.

a, Regional mean Pearson's correlations between empirical and simulated BOLD signals

during SST and MID tasks from 3-million-neuron DTB models for 290 participants. For each

task, Violin plots show both the mean correlations within assimilated task-related regions

(assimilated) and the mean correlations across all regions (whole). The boxplots indicating

the median and IQR; whiskers extend to 1.5× IQR. **b-d,** Simulated NP factor scores under

baseline (**b**), AMPA-mediated (**c**), and GABA-A-mediated (**d**) virtual perturbations in healthy

controls (HC, n = 69), high-symptom individuals (n = 90), and patients (n = 131). Group

differences were assessed using ANOVA and post-doc tests with Bonferroni correction for

multiple comparisons (n.s., not significant; **, $P_{corrected} < 0.01$). Values are residuals after

regressing out sex, site, and head motion. Violin plots show kernel density estimates with

embedded boxplots indicating the mean and IQR; whiskers extend to 1.5× IQR. **e,** Kernel

density distributions of NP factor scores across healthy controls (HC, grey), high-symptom

individuals (yellow), and patients (orange) under three simulation conditions: (left) baseline,

(middle) AMPA modulation, and (right) GABA-A modulation. The x-axis represents the

simulated NP factor scores, and shaded areas denote the distribution density for each group.

Across the three panels, the degree of overlap between group distributions and the relative

shift of scores from negative to positive values correspond to the transition from baseline to

specific receptor manipulations. **f,** Individual changes in NP factor scores following the

AMPA and GABA-A perturbations. Dashed lines indicate zero change. Each dot represents

an individual subject. Individuals with ΔNP>0 (perturbed-baseline) under both perturbations

were classified as the increased-response group (n = 251, pink); others formed the decreased-

response group (n = 39, blue). **g-h,** Baseline NP factor scores were inversely correlated with

AMPA- (**g**, Pearson's $r = -0.86$, $P < 0.0001$) and GABA-A- (**h**, Pearson's $r = -0.25$, $P < 0.0001$)

induced changes, with a permutation test of 10,000 times. **i,** Distribution of response types

across groups (Chi-square test: χ^2 = 14.05, P = 0.007). Increased-response proportions:

patients, 124/131 (94.66%); high-symptom, 73/90 (81.11%); HC, 54/69 (78.26%). **j,** The

increased-response group exhibited lower baseline NP factors, and higher symptom burden,

including panic attacks, negative affect and reduced social support, than the decreased-

response group (Mann–Whitney U tests, controlling for sex, site, and head motion; *, $P<0.05$;

**, $P<0.01$; ****, $P<0.0001$). Boxplots represent the median and IQR; whiskers extend to

1.5× IQR.

Figure 4 Pharmacological fMRI validation and prospective symptom prediction.

a, Pharmacological fMRI experiment in 27 male healthy volunteers using a three-way randomized, controlled crossover design. **b-e,** Individual- and group-level changes in summed NP-related MID functional connectivity (summed MID FCs) following ketamine and midazolam administration in subgroup 1 (**b**, **c**, n = 19) and subgroup 2 (**d**, **e**, n = 8). Each dot represents one individual and is colored by drug condition: placebo (grey), ketamine

(gold), and midazolam (teal). Statistical significance was assessed using paired *t*-tests or

Wilcoxon signed-rank tests, as appropriate. Values represent residuals after regressing out

head motion. **f,** Subgroup 1, exhibiting drug-induced increased responses, showed lower

baseline summed MID FCs under placebo compared with subgroup 2 (Mann–Whitney U

tests, controlling for head motion; ***, *P*<0.001). **g-h,** Baseline summed MID FC was

inversely correlated with midazolam-induced change (**g**, Pearson's *r* = -0.52, 10,000

permutations, *P* = 0.0056), and showed a similar trend for ketamine-induced change (**h**,

Pearson's *r* = -0.35, 10,000 permutations, *P* = 0.069). **i-j,** Prediction of empirical drug-

induced FC changes from baseline summed MID FC using linear mappings between DTB-

simulated E/I perturbations and simulated baseline MID FC strength. Scatter plots show

predicted versus observed FC changes induced by ketamine (**i**, Pearson's *r* = 0.35, 1,000

permutations, *P* = 0.039) and midazolam (**j**, Pearson's *r* = 0.52, 1,000 permutations, *P* =

0.001). Each dot represents one individual; solid lines indicate linear regression fits with 95%

confidence intervals. **k,** Prediction of drug-induced response direction (ΔNP>0 or <0) from

baseline summed MID FCs using DTB-derived models (AMPA model AUC = 0.69; GABA-A

model AUC = 0.83). **l,** Linear association between empirical NP factor scores and

behavioural symptom severity at age 19 (n = 100, r^2 = 0.24, *F* = 2.3, *P* = 0.014). Each dot

represents one individual; solid lines indicate linear regression fits with 95% confidence

intervals. **m,** Prospective prediction of behavioural symptom changes over 4 years. Scatter

plot shows predicted versus empirical behavioural changes from age 19 to 23. The prediction

model explained a significant proportion of variance (n = 85, r^2 = 0.23, *F* = 4.80, permutation

test of 5,000 times, *P* = 0.0006). Each dot represents one individual; solid line indicates the

linear fit with 95% confidence intervals. All boxplots show the median and interquartile

range (IQR); whiskers extend to 1.5× IQR. *, *P* < 0.05; **, *P* < 0.01; ***, *P* < 0.001.

Methods

Participants and MRI dataset.

IMAGEN dataset

The discovery dataset used to idenitify brain phenotype of transdiagnostic symptoms was

derived from the IMAGEN cohort (https://www.imagen-project.org/), a longitudinal,

population-based study comprising behavioural, neuroimaging, environmental, and genetic

data from approximately 2,000 participants assessed at ages 14, 19, and 23[1,2]. IMAGEN

participants were recruited from eight research sites across the United Kingdom, Germany,

France, and Ireland, and were cognitively normal with no history of psychiatric diagnoses.

To reduce brain developmental confounds, we focused on early adulthood, specifically the

follow-up 2 timepoint (aged 19y) of the IMAGEN study. At this time point, task-fMRI data

were available for the Monetary Incentive Delay (MID, n = 1,390) and Stop Signal Task (SST,

n = 1,394). After excluding participants with excessive head motion (mean framewise

displacement > 0.5 mm) and ensuring matched multimode data, the final sample included

1,050 participants (mean age = 18.41 ± 0.67 years; female/male = 564/486).

The task-fMRI paradigms used in this study are summarized below. The MID task comprised

42 trials across three reward conditions (no-win, small-win, and big-win). Each trial consisted

of a reward cue (250 ms), a fixation period (4-4.5 s), a target response, and reward feedback

(1,450 ms). The SST included 480 go trials (respond to arrow direction) and 80 stop trials

(inhibit response when a stop signal appeared ~300 ms after the go cue). The Emotional Face

Task (EFT) presented angry, neutral, and happy human faces in a passive-viewing paradigm

without responses. Each emotional condition comprised four trials, with each trial lasting 18

s. More details were available in the original paper of IMAGEN study [2].

 STRATIFY dataset

We validated brain phenotype of transdiagnostic symptoms in independent samples from

the STRATIFY cohorts[3], which used identical MRI protocols and behavioural assessments to

IMAGEN. These datasets included patients with psychiatric diagnoses and demographically

matched healthy controls. After the same screening procedure with IMAGEN, the final

sample included 209 patients (age = 22.26 ± 2.21; female/male = 135/74) and 225 healthy

controls (age = 21.40 ± 1.72; female/male = 129/96). Specific diagnoses included alcohol

use disorder (AUD, female/male = 58/40), major depressive disorder (MDD, female/male =

75/29), psychosis (female/male = 2/4), and attention-deficit/hyperactivity disorder (ADHD,

female/male = 0/1) for STRATIFY.

Clinical diagnoses were determined based on established thresholds from standardized

assessment tools consistent with DSM-5 criteria. Individuals with AUD were defined by an

Alcohol Use Disorders Identification Test (AUDIT)[4] score greater than 15. MDD was defined

by a Patient Health Questionnaire-9 (PHQ-9)[5] score greater than 15. Psychosis was defined

using ICD-10 diagnosis, with inclusion requiring at least one episode of schizophreniform

or chronic schizophrenia. Healthy controls were screened using rigorous inclusion criteria

to minimize confounding factors. Participants were excluded if they had: a) any current or

past mental health disorder[6], b) first- or second-degree relatives with mental health diagnoses,

c) regular use of medication for serious physical health conditions, d) learning difficulties

(e.g., Wechsler Adult Intelligence Scale verbal comprehension score below the 10th

percentile), or e) any history of recreational drug use.

Pharmacological dataset

To validate the effects of in sillco neurotransmitter manipulation in vivo, we utilized

pharmacological task-fMRI data[7] from 30 healthy male participants (mean age

27.3 ± 6.2 years). Each participant underwent three separate scanning sessions in a

randomized, single-blind, placebo-controlled, three-way crossover design, receiving

ketamine, midazolam, or placebo. A minimum 48-hour interval was maintained between

sessions to prevent carryover effects. The experimental procedure included a 16-minute

resting-state fMRI scan, during which drug administration began at the 7-minute mark,

followed by task-fMRI acquisition during the MID task. Three participants were excluded

due to missing MID behavioural recording files (n = 2) and T1-weighted structural MRI data

(n = 1). The MID task used here comprised 90 trials across two conditions: no-win (45 trials)

and win (45 trials). Details on administration protocols are available in the original study[7].

All above studies received approval from the local ethics committee, and written informed

consent was obtained from all participants.

Behavioural symptoms assessments.

We assessed behavioural symptoms for each individual from IMAGEN and STRATIFY

cohorts using the Development and Well-Being Assessment[8] (DAWBA) and Strengths and

Difficulties Questionnaire[9] (SDQ). To characterise externalising and internalising symptoms,

we summed the scores of relevant items from DAWBA and SDQ[3]. The externalising

symptoms included ADHD (8 items) and conduct disorder (CD, 7 items), while the

internalising symptoms encompassed eating disorder (ED, 5 items), specific phobia (SP, 13

items), general anxiety disorder (GAD, 7 items), and depression (DEP, 8 items). A detailed

list of the specific items is provided in Table S2.

MRI acquisition and preprocessing.

The structural and functional MRI protocols for the IMAGEN and STRATIFY studies were

harmonised across sites and scanner manufacturers (Siemens: 6 sites, Philips: 2 sites, General

Electric: 1 site, and Bruker: 1 site), Standardized hardware for visual and auditory stimulus

presentation (Nordic Neurolabs, Norway) was used at all sites.

Structural T1-weighted images

T1-weighted images (T1-w) of the IMAGEN and STRATIFY cohorts were acquired using a

protocol based on those from the ADNI study (https://adni.loni.usc.edu/data-samples/adni-

data/neuroimaging/mri/mri-scanner-protocols/). Several parameters in these protocols

deliberately differ between scanner models, in order to produce consistent image contrast

and quality despite implementation differences between manufacturers; all scans were

however collected with a sagittal slice plane and voxel size = $1.1 \times 1.1 \times 1.1$ mm^3.

The structural T1-w images were preprocessed using fMRIPrep[10] (version 20.2.3), a robust

and standardized pipeline based on Nipype, supported by Advanced Normalization Tools

(ANTs, version 2.3.3), FMRIB Software Library (FSL, version 5.0.9), Analysis of Functional

NeuroImages (AFNI, version 20160207), and FreeSurfer (6.0.1) packages. Anatomical

preprocessing included bias field correction, skull-stripping, tissue segmentation and

nonlinear registration to standard Montreal Neurological Institute (MNI) space.

Diffusion-weighted images

Diffusion-weighted images (DWIs) of IMAGEN and STRATIFY cohorts were acquired using

a 2D diffusion weighted EPI sequence: TR = 15,000 ms; TE = 104 ms; FOV = 307×307

mm^2; number of slices = 60; scan time = 9 min 45 s; voxel size = $2.4 \times 2.4 \times 2.4$ mm^3; 36

optimal non-colinear diffusion-weighted directions with b = 1,300 s/mm^2.

The structural DWIs were preprocessed using the FSL and MRtrix3 package. Eddy current-

induced distortions and head motion were corrected using FSL's eddy tool, and the rotated

b-vectors were updated using eddy_rotated_bvecs. The preprocessed DWI data was then

converted to the MRtrix3 format (.mif) with the mrconvert tool. The mean b0 image was

extracted using the mrmath command (mean), followed by brain extraction with the bet tool

(fractional intensity threshold = 0.3) to generate a brain mask. The white matter fiber

orientation distributions (FODs) within each voxel were estimated using constrained

spherical deconvolution (CSD), with response function estimation performed using the

Tournier method. The anatomical T1-weighted image was registered to the native diffusion

space using FSL's flirt tool. A nonlinear transformation to MNI space was then computed

using FSL's fnirt, and the inverse warp was generated with invwarp. The resulting warp was

subsequently converted into an MRtrix-compatible format for fiber transformation. A five-

tissue-type segmentation was carried out with the 5ttgen command. The gray matter-white

matter interface (GMWMI) was extracted using the 5tt2gmwmi tool to serve as the seeding

region for tractography. Whole-brain fiber tracking was performed using the tckgen

command with the iFOD2 algorithm, seeding from the extracted GMWMI and applying

Anatomically Constrained Tractography (ACT). The generated fiber tracks were transformed

from diffusion space to MNI space and generate voxel-wise and regional-level structural

connectivity matrices.

Resting-state and task-state functional MRI images

The resting-state fMRI scans of IMAGEN and STRATIFY cohorts were acquired using a 2D

gradient echo EPI sequence with the following parameters: TR = 2,200 ms; TE = 30 ms; FA

= 75°; FOV = 220 × 220 mm^2; number of slices = 40; number of volumes = 164; and voxel

size = 3.4 × 3.4 × 2.4 mm^3. Task-state fMRI scans were conducted with identical scanning

parameters, except for the number of volumes, that is 191 volumes for the MID task, 349

volumes for SST task, and 202 volumes for EFT task .

Addtionally, the pharamacological MID task-state fMRI were acquired on a 3T scanner

(Siemens Skyra, Erlangen, Germany), using the EPI sequence: TR = 2,200 ms; TE = 27 ms;

FA = 79°; FOV = 215 × 215 mm^2; number of slices = 30; number of volumes = 410; and

voxel size = 3 × 3 × 3 mm^3.

All fMRI data were preprocessed using fMRIPrep (version 20.2.3), including motion

correction, slice timing correction, and susceptibility distortion correction using fMRIPrep's

fieldmap-less approach. Functional MRI images were co-registered to the T1-weighted image

using boundary-based registration and then normalized to MNI152 space. Confound

regressors were extracted to control for physiological and motion-related artifacts, including

framewise displacement, DVARS, and CompCor components. All transformations were

applied in a single interpolation step to minimize resampling effects. Then the preprocessed

data underwent temporal detrending to remove low-frequency drifts, followed by spatial

smoothing using a full-width at half maximum (FWHM) of 6 mm Gaussian kernel and a

band-pass filtering (0.01 ~ 0.1 Hz). Quality control was performed by excluding subjects

with incomplete demographic or scanning timepoints; excessive head motion (mean

framewise displacement > 0.5 mm); or unsuccessful spatial normalization.

Identification of a compact network phenotype of psychopathology in IMAGEN.

Followed by previous work identifing the neuropsychopathology (NP) factor in the IMAGEN

baseline at age 14[3], we constructed MID and SST task-specific functional connectivity (FC)

matrics using the CONN toolbox (version 16.h) and a well-established 268-node whole-

brain functional parcellation[11], applying weighted generalised linear models after regressing

out task condition regressors and nuisance variables. Subsequently, we employed a

Connectome-based Predictive Modeling (CPM) [12] with a 50-fold cross-validation to predict

each behavioural symptom scores (including ADHD conduct disorder, eating disorder,

specific phobia, general anxiety disorder, and depression) from each task-specific functional

connectomes (SST conditions included stop-success, stop-failure, and go-wrong; MID

conditions included positive feedback, negative feedback, and reward anticipation). The

model performance was assessed using the Spearman's correlation between predicted and

empirical behavioural symptoms scores. This procedure was repeated 1,000 times, and only

edges selected in over 95% of models were used for further analysis. The mean P values

across 1,000 repetitions were corrected for multiple comparisons across 36 predictive

models using false discovery rate (FDR) correction ($q < 0.05$). Task-specific functional

connectivity (FC) matrices were then selected if they showed statistically significant

prediction performance for at least three behavioural symptom domains, ensuring robustness

across multiple diagnostic categories.

From these selected task-specific FC matrices, we extracted edges that significantly predicted

both externalising and internalising symptoms. Edges that were consistently associated with

behavioural symptoms across all selected task conditions were defined as transdiagnostic

associated edges. To improve interpretability, we stratified these transdiagnostic associated

edges into positive or negative FC profiles based on the direction of their associations with

behavioural symptoms. The summed strengths of these two profiles were termed as positive

and negative NP scores, respectively.

Case-control comparison of NP factors in STRATIFY.

To test whether the NP scores generalise to clinical populations, we re-extracted these

transdiagnostic associated FC profiles in the independent STRATIFY dataset and re-

calculated NP scores for each individual. We then compared NP scores between patients

and healthy controls, as well as among depression and alcohol use disorder subgroups, using

independent-samples t tests while covarying for sex, site, and head motion (mean framewise

displacement). Multiple comparisons were corrected using the Bonferroni method.

Permutation testing of PET receptor maps enrichment in NP-related regions.

To assess the neurochemical context of the NP factor, we examined whether the NP-related

regions exhibited higher receptor map values relative to non-NP regions using published

whole-brain   PET-derived   receptor   density   maps[13]

(https://github.com/netneurolab/hansen_receptors/tree/main/data/PET_nifti_images). For a

given PET map, we first extracted regional expressive levels using the 268-node functional

parcellation[11], and then computed the empirical difference in mean values between NP (n =

32) and non-NP (n = 236) regions. Statistical significance was assessed using permutation

testing (10,000 times), in which region labels (NP-related vs NP-unrelated) were randomly

reassigned while preserving the original group sizes. For each permutation, the difference in

mean receptor values between the permuted groups was recomputed, yielding a null

distribution expected under no spatial specificity. Two-sided permutation P values were

calculated as the proportion of permuted differences whose absolute value exceeded the

empirical difference. Resulting P values were corrected for multiple comparisons across 39

receptor maps using FDR correction.

Multimode neuroimaging data for Digital Twin Brain (DTB) models.

To constructing individualised DTB models, we extracted voxel-wise multimodel

neuroimaging data, including grey matter volume, white matter structural connectivity, and

functional BOLD signals from resting-state and task-state fMRI.

Gray matter volume was estimated using voxel-based morphometry (VBM) in SPM12

(Matlab R2020b). T1-weighted images were segmented into tissue maps, normalized to MNI

space using DARTEL, modulated by Jacobian determinants, and smoothed with an 8 mm

FWHM Gaussian kernel at $3 \times 3 \times 3$ mm³ resolution.

White matter structural connectivity was derived from whole-brain tractography using the

iFOD2 algorithm in MRtrix3, with anatomical constraints (ACT). Tracking used 5 million

streamlines seeded from the GMWMI, with a step size of 0.2 mm, curvature threshold of

45°/step, and length range of 3–250 mm.

To extract BOLD signals, we first generated individualised brain mask by selecting voxels

that: a) overlapped with both the 268-node functional parcellation[11] and the MNI152

template, b) had existing white matter structural connections, and c) were located in the

cortex or subcortex. On average, ~12,000 voxels per subject were retained. The time series

from these voxels were then extracted as simulation targets for the DTB models.

Importantly, incorporating specific neuronal models for the cerebellum and brainstem would

substantially increase model complexity and computational cost, particularly given that the

DTB simulations are implemented at the scale of hundreds of millions to billions of neurons.

We therefore adopted a parsimonious modeling strategy focused on cortical and subcortical

circuits, while acknowledging that this exclusion represents a limitation and that future

extensions incorporating differentiated regional parameterization may enable more

comprehensive characterization of NP network dynamics.

Construction of individualised task-state DTB models.

Followed by the framework developed by Lu et al.[14,15], we firstly constructed a excitation-

inhibition balanced spiking neural network constrained by individual multimode

neuroimaging data. In this network, each voxel was modeled as a sub-unit comprising one

excitatory and one inhibitory neuronal population. Each voxel included both excitatory and

inhibitory neurons at a fixed ratio of 4:1[16]. The number of neurons within each voxel was

determined by its proportional grey matter volume. Within-voxel synaptic architecture

followed a 4:1:2 ratio of internal excitatory, internal inhibitory, and external excitatory

synapses, based on anatomical organization of the cat visual cortex[17]. Between-voxel

connections were weighted by row-normalized structural connectivity derived from

diffusion MRI, and exclusively excitatory[18,19]. Neuronal activity was simulated using the

leaky integrate-and-fire model[20], incorporating α-amino-3-hydroxy-5-methyl-4-

isoxazolepropionic-acid (AMPA; excitatory-leaning) and γ-aminobutyric-acid-A (GABA-A;

inhibitory-leaning) synaptic conductances to represent glutamatergic and GABAergic

signaling:

1086
$$C_i \frac{dV_i}{dt} = -g_{L,i}(V_i - V_L) + \sum_u I_{syn,i} + I_{bg,i} + I_{ext,i}, \ V_i < V_{th,i}$$

where C_i is the capacitance of the neuron membrane, $g_{L,i}$ is the leakage conductance, V_i is

the membrane potential of neuron i, V_L is the leakage voltage, $I_{syn,i}$ is the synaptic currents of

two synapse types (AMPA and GABA-A), $I_{bg,i}$ is the background current as noises, and $I_{ext,i}$ is

the external current input for the task. It should be noted that the $I_{ext,i}$ are the independent

parameters, which are estimated using empirical task-fMRI data. The background noise is

given by independent Ornstein–Uhlenbeck processes and described as in the original

paper[15].

Simulated neuronal activity was quantified as mean firing rate of neuronal population within

each voxel, and was transformed into simulated BOLD signals using the Balloon-Windkessel

model[21]. The model performance was quantified by Pearson's correlation between simulated

and empirical task-state BOLD signals.

To optimize the fit between simulated and empirical BOLD signals, we applied the

Hierarchical Mesoscale Data Assimilation (HMDA) method[15] to estimate voxel-wise, time-

dependent hyperparameters. This assimilation procedure iteratively alternated between

forward simulating neuronal activity and parameter updating to reduce the mismatch

between simulated and observed BOLD signals. At each time point for each assimilated

voxel, the parameters of synaptic conductances for AMPA and GABA-A were sampled from

gamma distributions, which were constrained by voxel-specific hyperparameters. These

synaptic conductances governed the simulated neuronal dynamics, which were transformed

into BOLD signals. Hyperparameters were then iteratively updated so that the resulting

simulated BOLD signals progressively converged toward the empirical fMRI data. Parameter

updating was implemented using a diffusion ensemble Kalman filter (EnKF), which estimates

latent states and parameters by propagating an ensemble of 30 parallel model

realizations. More methodological details of HMDA are describedin Lu et al.'s original

paper[15].

Resting-state digital twin brain models were first obtained by assimilating empirical resting-

state fMRI BOLD signals for all voxels using the HMDA framework. This procedure yielded

individualised resting-state DTBs with converged hyperparameters and stable neuronal

dynamics. To construct task-state DTB models, we next assimilated empirical task-state fMRI

BOLD signals from task-engaged regions to infer digital external input currents. These

inferred digital input currents were then injected into the corresponding voxels of the resting-

state DTB to induce task-related neuronal activity. For regions not directly engaged by the

task, model hyperparameters were fixed to the mean values estimated during the resting-

state assimilation, ensuring a stable baseline while allowing task-specific dynamics to

propagate through the network.

Task-relevant regions were identified based on prior literature and meta-analytic activation

maps from the Neurosynth database. For the MID task, regions where received assimilated

currents included the orbitofrontal cortex, dorsal anterior cingulate cortex, insula, amygdala,

dorsal striatum, and nucleus accumbens, associated with reward processing. For the SST task,

assimilated regions involved the orbitofrontal cortex, anterior and dorsolateral prefrontal

cortex, dorsal ACC, insula, and thalamus, related to executive control. For the EFT task,

assimilated regions included the amygdala, dorsal striatum, hippocampus, insula, thalamus,

anterior and posterior cingulate cortex, dorsolateral prefrontal cortex, and orbitofrontal

cortex, which are related to emotional recognition and processing. Primary visual were

included for all three tasks, and sensorimotor cortices for first two tasks. A full list of

assimilated regions is provided in Table S6.

Assessment of subject-specific information in DTB-assimilated hyperparameters.

To evaluate whether DTB-assimilated hyperparameters capture subject-specific neural

dynamics independent of empirical data, we analyzed EFT task-based fMRI data from an

independent subset of four male participants, matched for scanning site (Berlin) and age (one

MDD and one AUD, both aged 21 years, and two HCs aged 22 years). Each participant

completed four trials per emotional block (angry, neutral, and happy), yielding 12 trials per

subject.

We first independently estimated hyperparameters for each trial, without using any other

trial of the same participant. Each trial-specific hyperparameter was then injected into noise-

initialized DTB models to generate simulated BOLD time series of matched duration. Trial-

level activation maps were computed from simulated and empirical BOLD signals using

identical preprocessing and statistical modeling procedures. To quantify subject specificity,

we correlated activation maps from all simulated and empirical trials across subjects using

Pearson correlation coefficients, yielding a 48 × 48 similarity matrix (4 subjects × 12 trials),

and compared correlations between simulated and empirical trials from the same individual

(self–self) with those between trials from different individuals (self–other) using a Mann–

Whitney U test.

In addition, we tested whether simulated activation maps, generated without re-assimilation

of empirical data, were sufficiently distinctive to support individual identification. For each

participant, we created subject-specific empirical templates by averaging the first two trials

of each emotional condition. Hyperparameters estimated from these first-half trials were

used to simulate BOLD time series for the remaining trials without assimilation. We then

correlated activation maps from simulated second-half trials (4 subjects × 6 trials) with all

subject-specific templates, and aligned each simulated trial to the subject with the highest

correlation. The same identification procedure was applied to empirical activation maps

from the second-half trials as a benchmark.

Calibration experiment of DTB model scales.

The DTB framework allows flexible specification of total neuronal count across a wide range,

from approximately 1 million to 86 billion neurons[14,15]. To balance biological fidelity,

perturbational controllability, and computational feasibility, we performed calibration

experiments to determine optimal model resolutions for three complementary analytical

purposes: high-fidelity simulation, controlled perturbational exploration, and population-

level inference.

We first evaluated the influence of model scale using the DTB of an illustrative healthy

control participant (HC01), by varying neuronal counts from 10 million to 5 billion and

assessing simulation fidelity across resolutions. Model performance was quantified using the

mean voxel-wise Pearson's correlation between simulated and empirical BOLD signals, as

well as correlations between simulated and empirical whole-brain functional connectivity

matrices (268-node functional parcellation[11]). These experiments established the relationship

between neuronal scale and task-state simulation accuracy, providing an empirical basis for

selecting biologically realistic yet computationally tractable model resolutions.

Second, to identify the optimal resolution for the population-scale simulations, we

constructed reduced-scale DTBs ranging from 1 to 9 million neurons for the same participant.

In these models, individual MRI data were represented at the regional rather than voxel-wise

level. The number of neurons per region was assigned in proportion to the regional grey

matter volume. While voxel-wise grey matter volumes could in principle be used, the total

number of neurons in millions of neuron models would make it impractical to allocate

neurons to every individual voxel, as some voxels have a minimal ratio of grey matter volume.

By using regional-level allocation, we ensured that the neuron distribution faithfully reflected

grey matter proportions while remaining computationally tractable. Between-regional

connections were weighted by row-normalized regional structural connectivity derived from

diffusion MRI. The other details were consistent with the methodology used in our voxel-

wise simulation models. Model performance was quantified using the mean regional-level

Pearson's correlation between simulated and empirical BOLD signals, as well as correlations

between simulated and empirical whole-brain functional connectivity matrices (268-node functional parcellation[11]).

Together, these calibration procedures ensured that DTB simulations maintained high fidelity while enabling scalable individualised and population-level analyses.

DTB Simulations for four illustrative participants.

Following the calibration experiments described above (see Supplementary Results), high-fidelity 1-billion-neuron DTBs were constructed for four illustrative participants: two patients (one MDD and one AUD) and two healthy controls (HC01 and HC02; demographics information in Table S7). The patients were selected as those with the highest negative NP scores within their respective diagnostic groups, indicating the highest symptom burden. Among the healthy controls, HC01 and HC02 were chosen from the subset of participants with the lowest behavioural symptom summed scores in the IMAGEN cohort. HC01 had the lowest negative NP score, whereas HC02 had the highest negative NP score, illustrating the range of NP values even among individuals with minimal behavioural symptoms. All selected participants had complete multi-modal imaging data, including T1-weighted, DTI, and fMRI scans.

In addition to billion-neuron simulations, 100-million-neuron DTBs were constructed for the same participants to enable computationally efficient perturbational analyses while retaining individualised model characteristics. This intermediate resolution was selected by calibration results showing no clear performance inflection point across neuronal scales (Fig.S4; Table S8), suggesting that reduced-scale models could preserve essential network properties while improving controllability for systematic parameter exploration.

To evaluate whether 100-million-neuron DTBs retained subject-specific network phenotypes,

we extracted simulated NP factor scores from each individualised model and compared them

with empirical NP factors derived from all individuals in the IMAGEN and STRATIFY datasets.

Similarity was quantified using mean-squared error (MSE) between each pair of simulated

and empirical NP factors. For each illustrative participant, self-to-self MSE (simulated versus

own empirical NP factor) was contrasted against cross-participant MSE (simulated versus all

other empirical scores). Statistical significance was evaluated using permutation testing

across all 1,575 simulated - empirical pairings.

Systematic parameter sweeps of virtual AMPA and GABA-A perturbations.

Based on 100-million-neuron DTB models of four illustrative participants, we performed

systematic perturbations of AMPA- and GABA-A-mediated synaptic conductances. AMPA

conductance was varied from 0.0020 to 0.0052 in steps of 0.0004 (baseline = 0.0008), and

GABA-A conductance from 0.0015 to 0.0040 in steps of 0.0005 (baseline = 0.0015). The

baseline levels of AMPA and GABA-A in the model were determined across two key

dimensions: the mean firing rate of the neuronal population (<10 Hz), consistent with

physiological observations[22], and the synchrony measure, which represents the degree of

oscillatory behaviour. These metrics identified the optimal parameter window where the

model maintains robust, reasonable neuronal firing while avoiding excessive

synchronization[23], thereby ensuring a stable and biologically plausible baseline for the

model.

Crucially, the upper limit for the parameter of AMPA conductance was constrained by the

stability of the simulated BOLD signal: beyond a specific threshold, excessive neuronal firing

led to a cessation of dynamic fluctuations in the BOLD signal. We considered such states as

"over-firing" regimes, where the hemodynamic response reaches a non-biological plateau

and no longer reflect meaningful neural dynamics[24]. During GABA-A manipulations, AMPA

conductance was held fixed at the value that had produced the maximal NP factor

enhancement during prior AMPA sweeps, allowing controlled assessment of excitatory-

inhibitory balance, and preventing excessive inhibition that could silence neuronal activity

and abolish intrinsic BOLD fluctuations[25].

These synaptic conductance parameters were implemented as global control variables,

applied uniformly across the whole-brain model rather than specific regions. Importantly,

they were directly applied to DTB models that had already been calibrated to individual

empirical fMRI data; therefore, no additional data assimilation or parameter refitting was

required during virtual perturbations. The NP network responses were obtained by forward

simulation under systematically varied synaptic levels. This design allows direct assessment

of perturbational sensitivity around individualised baseline operating points. To ensure

reproducibility, each perturbation was repeated five times, and the mean NP factor scores

across repetitions were used for subsequent analyses. Finally, while multiple combinations

of synaptic parameters can give rise to similar network outputs, our goal was not to recover

exact physiological parameter values, but to use biologically interpretable synaptic

conductance parameters as control variables to probe regime-dependent network responses.

Evaluation of cross-task generalisability of virtual perturbations.

To test whether the effects of AMPA- and GABA-A–mediated perturbations on the NP factor

generalise across cognitive contexts, we conducted the same virtual perturbation to the DTB

models that simulate EFT task activity of the same four illustrative participants. Specifically,

we first calibrated DTB models at a 100-million-neuron scale to simulate BOLD signals of

EFT task-based fMRI data, and then applied the optimal perturbational parameters identified

in the parameter sweeps to these DTB models. For each participant, NP-related functional

connectivity was extracted from baseline and perturbed simulated EFT BOLD signals using

the same computational pipeline as the original MID and SST tasks, enabling direct

comparison of perturbation effects across cognitive tasks.

Population-scale virtual perturbations.

To assess whether virtual E/I modulations in DTB models generalise across the cohort, we

first constructed DTB models for 290 participants, including 72 patients with MDD (age =

22.33 ± 2.20, female/male = 48/24), 59 with AUD (age = 22.41 ± 2.02, female/male =

37/22), and 69 healthy controls (age = 21.45 ± 1.38, female/male = 38/31) from STRATIFY

dataset, as well as 90 subclinical participants with high behavioural symptom scores (sum of

six externalising and internalising symptoms ≥ 20) from IMAGEN (age = 18.42 ± 0.65,

female/male = 74/16). We then applied the optimal perturbational parameters identified in

the parameter sweeps to these DTB models.

For each participant, we calculated simulated NP factor scores derived from baseline and

perturbed BOLD signals. Group differences in empirical, simulated, and perturbed NP

factors were assessed using ANOVA analysis and post-doc tests, controlling for sex, site, and

head motion (mean framewise displacement). The paired-sample t tests were used to

compare simulated and empirical, and simulated and perturbed NP factors, among

subgroups. Multiple comparisons were corrected using the Holm-Bonferroni method.

Prediction of task performance from DTB simulations.

To assess the behavioural validity of digital twin brain simulations, we evaluated whether

simulated task-state functional connectivity metrics, derived from population-scale

simulations, significantly predicted individual differences in task performance during the

MID task.

Analyses were conducted in the 287 participants with 3-million-neuron DTB simulations

and MID task behavioural data. Individualised task-state functional connectomes were

extracted separately for the anticipation-hit and feedback-hit stages of the MID task.

Behavioural performance was quantified as mean response time (RT) under three incentive

conditions: big-win, small-win, and no-win. Prediction of behavioural performance was

performed using a connectome-based predictive modeling framework with 10-fold cross-

validation. For each fold, linear models were trained on task-state FC matrices from nine

folds to predict RT measures and evaluated on the held-out fold. Prediction accuracy was

quantified as the Spearman's correlation between predicted and observed RT values. The

entire cross-validation procedure was repeated 100 times to ensure stability, and prediction

accuracies were averaged across repetitions. Separate models were constructed for each

combination of task stage (anticipation-hit, feedback-hit) and behavioural measure (big-win,

small-win, no-win), resulting in six independent prediction models.

We also examined the prediction performances of empirical task-state FC for establishing a

benchmark. Prediction accuracies obtained using empirical FC and DTB-simulated FC were

compared using paired-sample *t* tests across the six model combinations.

Stratification of NP network responses to virtual perturbations.

In the population-level analysis, we quantified individual responses to virtual

neurotransmitter perturbations in the DTB models. For each participant, we computed the

change in simulated NP factor following AMPA and GABA-A modulation relative to baseline.

Participants were classified as "increasers" (ΔNP>0 following both modulations) or

"decreasers" (ΔNP ≤ 0 for any given perturbation). We compared the distribution of response

types across diagnostic groups (patients, high-symptom, and healthy controls), sex, and site

using Chi-square tests.

We also assessed the relationship between baseline simulated NP factor and the AMPA- or

GABA-A–induced changes using Pearson's correlation. Statistical significance was assessed

using permutation testing (10,000 permutations), in which ΔNP values were randomly

reassigned across individuals to generate a null distribution. For each permutation, the

correlation between simulated baseline NP and permuted ΔNP was recomputed. Two-sided

permutation P values were calculated as the proportion of null correlations whose absolute

value exceeded the observed correlation, directly testing whether modulation-induced

changes reflect systematic baseline dependence rather than symmetric fluctuations or

regression-to-the-mean effects.

Finally, we compared simulated baseline NP factors and behavioural symptom measures

using Mann–Whitney U tests, given unequal subgroup sizes and non-normal distributions.

All analyses controlled for sex, scanning site, and mean framewise displacement.

Behavioural measures were derived from the 69 entry items of the DAWBA and the SDQ.

Stratification of individual pharmacological responses.

To validate DTB-predicted effects of virtual AMPA and GABA-A modulations on the NP

factor, we analyzed an independent pharmacological fMRI dataset in which healthy

participants completed MID task-fMRI under placebo, ketamine, and midazolam in a

randomised cross-over design. For each participant and drug condition, we computed NP-

related MID FC strengths (six edges) and summed them as a "summed MID FC" metric.

We first tested group-level drug effects by comparing the summed MID FC between placebo

and drug conditions using paired-sample t tests. To further capture individual variability, we

computed the drug-induced change in summed MID FC relative to placebo for each

participant and applied k-means clustering (500 iterations) to identify two response

subgroups. Within each subgroup, paired-sample t tests, controlling for head motion, were

used to compare placebo and drug conditions. Wilcoxon signed-rank tests were applied

when normality or variance assumptions were violated.

We also compared summed MID FC under placebo between response subgroups using a

independent-samples t test. Finally, the relationship between baseline summed MID FC in

placebo and drug-induced FC change across all participants was assessed using Pearson's

correlations. Statistical significance was assessed using permutation testing (10,000

permutations), in which Δsummed MID FC values were randomly reassigned across

individuals to generate a null distribution. For each permutation, the correlation between

baseline summed MID FC and permuted Δsummed MID FC was recomputed. Two-sided

permutation P values were calculated as the proportion of null correlations whose absolute

value exceeded the observed correlation, directly testing whether drug-induced changes

reflect baseline dependence rather than regression-to-the-mean effects.

Prediction of pharmacological responses from DTB-derived virtual perturbations.

To test whether DTBs capture systematic relationships between baseline network

configuration and drug-induced NP-related connectivity changes, we first constructed a

linear regression model linking simulated baseline summed MID FC and FC changes

following virtual AMPA and GABA-A modulations. We then applied this regression model

to predict empirical FC changes under ketamine or midazolam from each participant's

baseline FC under placebo. Here, we chose a linear mapping to minimize overfitting and

maximize interpretability given the modest sample size. Prediction accuracy was quantified

by Pearson's correlation between predicted and observed FC changes, with significance

assessed via permutation tests (1,000 permutations of subject labels).

To evaluate whether DTB-based predictions can distinguish individuals with opposite drug

responses, participants were classified as "increasers" or "decreasers" based on the their

observed drug-induced change in summed MID FC. The predicted probabilities from the

DTB-derived regression model were then used to calculate the area under the receiver

operating characteristic curve (AUC) as a measure of how well the model correctly identifies

individuals in each response category.

Prediction of longitudinal symptom changes from DTB-derived virtual perturbations.

To test whether DTB-derived virtual E/I modulations predict longitudinal symptom

trajectories, we first estimated an empirical linear mapping between NP factors and

behavioural symptoms using baseline data. Specifically, we fitted a linear regression linking

NP factors at age 19 to summed internalising symptom scores at age 19 across four domains

(eating disorder, depression, generalized anxiety, and specific phobia). Externalising

symptoms (ADHD and conduct disorder) were excluded due to their low prevalence at

follow-up (age 23).

Baseline NP factors and NP factors under virtual AMPA and GABA-A modulation were

simulated for the full cohort using 3-million-neuron DTBs. From this cohort, we selected

individuals with imaging data at age 19 and symptom assessments at both ages 19 and 23.

The empirical NP–symptom mapping derived at baseline was then applied to the DTB-

simulated NP factors to generate predicted symptom scores at baseline and after virtual

modulation.

For each individual, we defined a DTB-derived "behavioural restoration" index as the

difference between predicted baseline and post-modulation symptom scores. Longitudinal

symptom change over four years (age 23 minus age 19) was subsequently modeled using

multiple linear regression, with baseline symptom scores and the DTB-derived "behavioural

restoration" index included as predictors. Prediction accuracy was quantified by Pearson's

correlation between predicted and observed behavioural symptom changes. Model fit was

quantified using the coefficient of determination (r^2). The incremental variance explained by

the DTB-derived index (Δr^2) was assessed by comparing full and reduced models using F

tests. Statistical significance of DTB-related effects was further evaluated using 5,000

permutation tests for both r^2 and the regression coefficient associated with the DTB index.

1380

Methods References

1.   Mascarell Maričić, L. *et al.* The IMAGEN study: a decade of imaging genetics in adolescents. *Mol. Psychiatry* 25, 2648–2671 (2020).

2.   Schumann, G. *et al.* The IMAGEN study: reinforcement-related behaviour in normal brain function and psychopathology. *Mol. Psychiatry* 15, 1128–1139 (2010).

3.   Xie, C. *et al.* A shared neural basis underlying psychiatric comorbidity. *Nat. Med.* 29, 1232–1242 (2023).

4.   Allen, J. P., Litten, R. Z., Fertig, J. B. & Babor, T. A review of research on the Alcohol Use Disorders Identification Test (AUDIT). *Alcohol. Clin. Exp. Res.* 21, 613–619 (1997).

5.   Kroenke, K., Spitzer, R. L. & Williams, J. B. The PHQ-9: validity of a brief depression severity measure. *J. Gen. Intern. Med.* 16, 606–613 (2001).

6.   Sheehan, D. V. *et al.* The Mini-International Neuropsychiatric Interview (M.I.N.I.): the development and validation of a structured diagnostic psychiatric interview for DSM-IV and ICD-10. *J. Clin. Psychiatry* 59 Suppl 20, 22-33;quiz 34-57 (1998).

7.   Forsyth, A. *et al.* Comparison of local spectral modulation, and temporal correlation, of simultaneously recorded EEG/fMRI signals during ketamine and midazolam sedation. *Psychopharmacology (Berl.)* 235, 3479–3493 (2018).

8.   Goodman, R., Ford, T., Richards, H., Gatward, R. & Meltzer, H. The Development and Well-Being Assessment: Description and initial validation of an integrated assessement of child and adolescent psychopathology. *J. Child Psychol. Psychiatry* https://doi.org/10.1017/S0021963099005909 (2000) doi:10.1017/S0021963099005909.

9.   Goodman, R. Psychometric properties of the strengths and difficulties questionnaire.

J. Am. Acad. Child Adolesc. Psychiatry 40, 1337–45 (2001).

10.   Esteban, O. *et al.* fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nat.*

Methods 16, 111–116 (2019).

11.   Shen, X., Tokoglu, F., Papademetris, X. & Constable, R. T. Groupwise whole-brain

parcellation from resting-state fMRI data for network node identification. *NeuroImage* 82,

403–415 (2013).

12.   Shen, X. *et al.* Using connectome-based predictive modeling to predict individual

behavior from brain connectivity. *Nat. Protoc.* 12, 506–518 (2017).

13.   Hansen, J. Y. *et al.* Mapping neurotransmitter systems to the structural and functional

organization of the human neocortex. *Nat. Neurosci.* 25, 1569–1581 (2022).

14.   Lu, W. *et al.* Imitating and exploring human brain's resting and task-performing states

via resembling brain computing: scaling and architecture. *Natl. Sci. Rev.* nwae080 (2024)

doi:10.1093/nsr/nwae080.

15.   Lu, W. *et al.* Simulation and assimilation of the digital human brain. *Nat. Comput.*

Sci. 4, 890–898 (2024).

16.   Brunel, N. Dynamics of sparsely connected networks of excitatory and inhibitory

spiking neurons. *J. Comput. Neurosci.* 8, 183–208 (2000).

17.   Binzegger, T., Douglas, R. J. & Martin, K. A. C. A Quantitative Map of the Circuit of

Cat Primary Visual Cortex. *J. Neurosci.* 24, 8441–8453 (2004).

18. Deco, G., Jirsa, V. K. & McIntosh, A. R. Emerging concepts for the dynamical

organization of resting-state activity in the brain. *Nat. Rev. Neurosci.* 12, 43–56 (2011).

19. Isaacson, J. S. & Scanziani, M. How Inhibition Shapes Cortical Activity. *Neuron* 72,

231–243 (2011).

20. Abbott, L. F. Lapicque's introduction of the integrate-and-fire model neuron (1907).

Brain Res. Bull. 50, 303–304 (1999).

21. Friston, K. J., Mechelli, A., Turner, R. & Price, C. J. Nonlinear responses in fMRI: the

Balloon model, Volterra kernels, and other hemodynamics. *NeuroImage* 12, 466–477 (2000).

22. Yu, Y. *et al.* A 3D atlas of functional human brain energetic connectome based on

neuropil distribution. *Cereb. Cortex* 33, 3996–4012 (2023).

23. Baroni, F. & Fulcher, B. Synchrony, oscillations, and phase relationships in collective

neuronal activity: A highly comparative overview of methods. *PLOS Comput. Biol.* 21,

e1013597 (2025).

24. Deco, G. *et al.* Resting-state functional connectivity emerges from structurally and

dynamically shaped slow linear fluctuations. *J. Neurosci. Off. J. Soc. Neurosci.* 33, 11239–

11252 (2013).

25. Li, J., Kells, P. A., Osgood, A. C., Gautam, S. H. & Shew, W. L. Collapse of complexity

of brain and body activity due to excessive inhibition and MeCP2 disruption. *Proc. Natl.*

Acad. Sci. 118, e2106378118 (2021).

1441

Acknowledgments

This work received support from the following sources: National Natural Science Foundation of China (32400935 to YX, W2541022 to GS, 82150710554 to GS), National Science and Technology Major Project (2025ZD0215100 to GS), 111 Project (B18015 to JF), Shanghai Municipal Science and Technology Major Project (2018SHZDZX01 to JF), the European Union-funded FP6 Integrated Project IMAGEN (Reinforcement-related behaviour in normal brain function and psychopathology) (LSHM-CT- 2007-037286 to GS), the Horizon 2020 funded ERC Advanced Grant 'STRATIFY' (Brain network based stratification of reinforcement-related disorders) (695313 to GS), Horizon Europe 'environMENTAL', grant no: 101057429 to GS, UK Research and Innovation (UKRI) Horizon Europe funding guarantee (10041392 and 10038599) to SD, Human Brain Project (HBP SGA 2, 785907, and HBP SGA 3, 945539) to GS. The German Center for Mental Health (DZPG), the Bundesministerium für Bildung und Forschung (BMBF grants 01GS08152; 01EV0711; Forschungsnetz AERIAL 01EE1406A, 01EE1406B; Forschungsnetz IMAC-Mind 01GL1745B), the Deutsche Forschungsgemeinschaft (DFG project numbers 458317126 [COPE] to GS, 186318919 [FOR 1617], 178833530 [SFB 940], 386691645 [NE 1383/14-1], 402170461 [TRR 265], 454245598 [IRTG 2773]), the Medical Research Foundation and Medical Research Council (grants MR/R00465X/1 and MR/S020306/1) to SD, the National Institutes of Health (NIH) funded ENIGMA-grants 5U54EB020403-05, 1R56AG058854-01 and U54 EB020403 as well as NIH R01DA049238, the National Institutes of Health, Science Foundation Ireland (16/ERCD/3797) to ALWB and RW.

Consortia

DTB Consortium

Yubin Bao, Boyu Chen, Siming Chen, Zhongyu Chen, Fei Dai, Weiyang Ding, Xin Du,

Jianfeng Feng, Yubo Hou, Mingda Ji, Peng Ji, Chong Li, Chunhe Li, Xiaoyi Li, Yuhao Liu,

Wenlian Lu, Zhihui Lv, Hengyuan Ma, Yang Qi, Edmund Rolls, He Wang, Huarui Wang,

Jiexiang Wang, Shouyan Wang, Ziyi Wang, Yunman Xia, Shitong Xiang, Chao Xie,

Xiangyang Xue, Leijun Ye, Longbin Zeng, Tianping Zeng, Chenfei Zhang, Jie Zhang, Nan

Zhang, Wenyong Zhang, Yicong Zhao & Qibao Zheng

IMAGEN Consortium

Tobias Banaschewski, Gareth J. Barker, Arun L.W. Bokde, Rüdiger Brühl, Sylvane Desrivières,

Herta Flor, Penny Gowland, Antoine Grigis, Andreas Heinz, Herve Lemaitre, Jean-Luc

Martinot, Marie-Laure Paillère Martinot, Eric Artiges, Frauke Nees, Dimitri Papadopoulos

Orfanos, Tomáš Paus, Luise Poustka, Michael N. Smolka, Sarah Hohmann, Nilakshi Vaidya,

Henrik Walter, Robert Whelan, Paul Wirsching & Gunter Schumann

STRATIFY Consortium

Nilakshi Vaidya, Zuo Zhang, Lauren Robinson, Jeanne Winterer, Yuning Zhang, Gareth J

Barker, Arun L.W. Bokde, Rüdiger Brühl, Hedi Kebir, Hervé Lemaître, Frauke Nees, Dimitri

Papadopoulos Orfanos, Ulrike Schmidt, Julia Sinclair, Robert Whelan, Henrik Walter,

Sylvane Desrivières & Gunter Schumann

EnvironMENTAL Consortium

Tobias Banaschewski, Sylvane Desrivières, Andreas Heinz, Frauke Nees, Nilakshi Vaidya,

Henrik Walter, Vince D. Calhoun & Gunter Schumann

Author contributions

Data availability

The data analysed in this study were obtained from the IMAGEN consortium and associated

cohorts. Due to ethical and data protection restrictions, the raw data are not publicly

available. Researchers may request access to the data through the official data access

procedures of the respective consortia. Processed data supporting the findings of this study

are available from the corresponding author upon reasonable request and subject to

consortium approval.

Code availability

The code used to perform the analyses in this study is publicly available at GitHub

(https://github.com/Shmily94/Digital_twin_brain-psychiatry-.git). Custom codes developed

in PyTorch for personal computing environments and in C++ for high-performance

computing (HPC) systems of the DTB platform is available via the DTB Consortium GitHub

repository   (https://github.com/DTB-consortium/Digital_twin_brain-open)   and   Zenodo

(https://doi.org/10.5281/zenodo.13995756).   Please   note   that   parts   of   the   HPC

implementation depend on specific hardware and software configurations. Further technical

information can be requested from the DTB Consortium via dtb.fudan@gmail.com.