Enhancing cGMP signaling with psilocybin reduces head twitch and restructures the synaptic proteome while maintaining antidepressant response

Gabriele Floris*[1], Sarah J. Jefferson*[1], Jocelyne Rondeau[1], Frank S. Menitti[2,3], Alex C. Kwan[4,5], Joao P. De Aquino[1,9], John H. Krystal[1,2,6,7], Christopher Pittenger[1,6,8,9,#], Alfred P. Kaye[1,9,10,11,#]

1. Yale University Department of Psychiatry, New Haven, CT

2. Freedom Biosciences, Tiburon, CA 94920

3. George & Anne Ryan Institute for Neuroscience, University of Rhode Island, Kingston, RI

4. Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY

5. Department of Psychiatry, Weill Cornell Medicine, New York, NY

6. Yale University Department of Psychology, New Haven, CT

7. Yale University Department of Neuroscience, New Haven, CT

8. Yale University Child Study Center, New Haven, CT

9. Yale University Center for Brain Mind Health, New Haven, CT

10. Yale University Department of Comparative Medicine, New Haven, CT

11. Yale University Wu Tsai Institute, New Haven, CT

* Equal Contributions

# These authors jointly supervised the work

Corresponding author: Alfred P. Kaye. Email: alfred.kaye@yale.edu

Abstract

New treatments for depression are needed that combine robust efficacy with improved scalability. Although psilocybin has demonstrated antidepressant effects in Phase 3 clinical trials, some of its psychedelic effects limit tolerability, necessitating administration in highly supervised clinical settings, and thus motivating development of serotonergic therapeutics that preserve antidepressant efficacy while reducing the acute psychedelic experience. We combined psilocybin with a phosphodiesterase-9 inhibitor (PDE9i), which raises cyclic GMP

levels, and observed substantial reduction in the mouse head twitch response (HTR) — a proxy for 5-HT$_2$A receptor-mediated psychedelic-like behavior in rodents — suggesting attenuation of acute psychedelic effects. Significantly, rescue of chronic stress-induced depressive-like behavior by psilocybin was maintained with the coadministration of PDE9i. Proteomic analysis of medial prefrontal cortex (mPFC) synaptosomes showed that the combination of PDE9i and psilocybin enhanced synaptogenesis pathways relative to psilocybin alone, while reducing pathways involved in G protein-coupled receptor (GPCR) signaling. Together, these results suggest that the combination of PDE9i and psilocybin may be a promising direction for psychedelic treatment, and point towards molecular pathways that dissociate acute psychedelic and antidepressant responses.

Introduction

Major Depressive Disorder affects over 300 million people worldwide and approximately one-third of them fail to respond to conventional antidepressants [1]. Psilocybin, a serotonin 2A receptor (5-HT$_2$AR) agonist, shows marked efficacy in refractory depression in phase 3 clinical trial (NCT05711940) [2]. Psilocybin administration is also accompanied by marked changes in perception mediated by 5-HT$_2$AR-dependent disruption of functional brain networks, including the default mode network [3]. The intensity of these acute psychedelic effects can limit broad clinical implementation, motivating efforts to dissociate therapeutic from acute psychedelic effects. Multiple strategies are being explored, including non-hallucinogenic analogs, biased agonists targeting specific 5-HT$_2$AR signaling pathways, and structure-based drug design approaches [4-6]. While these novel compounds show promise in preclinical studies, their clinical efficacy in patient populations remains unknown. Combination-based strategies offer an alternative approach that maintains psilocybin's therapeutic efficacy, while pharmacologically targeting mechanisms underlying its acute perceptual effects. Phosphodiesterase-9A (PDE9A) represents a compelling molecular target in this context. PDE9A is a high-affinity, cGMP-specific enzyme (Km ≈ 70 nM) with dense expression in hippocampus and prefrontal cortex, where it regulates cGMP pools

distinct from those controlled by other phosphodiesterases [7]. Selective PDE9 inhibitors, such as BAY 73-6691 and PF-04447943, enhance synaptic plasticity and produce robust pro-cognitive effects in rodent models [8]. Critically, PDE9 inhibition reverses mescaline-induced scratching behavior, a 5-HT$_2$AR-dependent response [9], providing direct evidence that cGMP modulation can influence behavioral outputs of 5-HT$_2$AR activation. Here, we investigated whether PDE9 inhibition modulates acute behavioral and molecular responses to psilocybin. We quantified head twitch response (HTR) as a behavioral readout of 5-HT$_2$AR engagement and predictor of acute perceptual effects, and assessed whether PDE9 inhibition interferes with psilocybin's antidepressant-like effects in a chronic stress paradigm. Finally, synaptic proteomic analysis was performed to characterize the molecular landscape underlying these combinatorial effects.

Methods

Animals and Housing

Male C57BL/6J mice (8–12 weeks old; The Jackson Laboratory) were group-housed under standard conditions (12:12 h light/dark cycle) with ad libitum food and water. All procedures were approved by the Yale University Institutional Animal Care and Use Committee.

Drugs

Psilocybin (item #14041, Cayman Chemical Company) was dissolved in saline and administered at 1 mg/kg, i.p. The PDE9A inhibitor PF-4181366 was synthesized and generously provided by Yuelian Xu at Chinglu Pharmaceutical Research LLC. It was analyzed by UPLC-MS at 98.95% purity. PF-4181366 was dissolved in 5% DMSO, 5% Tween-80 in saline at doses previously shown to increase cGMP levels in cortex and hippocampus (0.32, 1, 3.2, or 10 mg/kg, i.p.) [10].

The psilocybin dose selected here (1 mg/kg i.p.) is consistent with doses established to produce robust 5-HT$_2$AR-dependent HTR in rodents and to elicit neuroplastic responses in

preclinical depression models [11]; for reference, human therapeutic doses span approximately 0.14–0.36 mg/kg orally, a range shown to exhibit linear pharmacokinetics in healthy adults [12].

Head Twitch Response

We detected HTR using automated magnetometer-based system. Psilocybin (1 mg/kg) was administered 30 min after pretreatment with PF-4181366 (0.32, 1, 3.2, or 10 mg/kg) or vehicle and HTR was quantified immediately thereafter.

Locomotor Activity

Following 1h habituation, mice were placed individually into standard cages positioned within a 16-beam infrared photobeam grid (Med Associates Inc.). Locomotor activity was recorded for 60 min before and 60 min after injection of PF-4181366 (0.32, 1, 3.2, or 10 mg/kg) or saline.

Chronic Multimodal Stress

Chronic multimodal stress (CMMS) was adapted from Hesselgrave et al. [13]. Mice were single-housed and habituated to two-bottle choice (1.5% sucrose and water). Baseline sucrose preference was assessed on Days 2–3 using 1% sucrose during 14 h dark-phase sessions. Beginning Day 4, mice underwent 15 consecutive days of stress: 5 h daily restraint in ventilated 50 mL conical tubes under concurrent 2 Hz strobe lighting and 75 dB white noise. Session timing varied daily (8-11 AM). Sucrose preference was reassessed on Days 18–19 to evaluate stress-induced anhedonia.

Sucrose Preference Test

Sucrose preference was calculated as [sucrose intake / (sucrose + water intake)] × 100. A 1.5% solution was used during habituation; 1% was used for all testing sessions. Bottle positions were alternated to control for side preference.

Tail Suspension Test

On Day 17, mice were suspended by adhesive tape (1 cm from tail tip) to a horizontal metal bar. Behavior was video-recorded for 6 min, and immobility—defined as absence of active escape-oriented movements—was scored manually during minutes 1–5.

Pharmacological Treatment

Following TST, mice were stratified by sucrose preference and immobility into three groups: (1) vehicle–saline, (2) vehicle–psilocybin (1 mg/kg), and (3) PF-4181366 (10 mg/kg)–psilocybin (1 mg/kg). A 30 min interval separated pretreatment from psilocybin injection. PF-4181366 was administered intraperitoneally (i.p.) in 5% DMSO, 5% Tween-80, and saline; psilocybin was administered i.p. in sterile saline.

Post-Treatment Sucrose Preference

Four hours post-treatment, mice underwent a 14 h overnight sucrose preference test using 1% sucrose. Intake was measured by weight.

Forced Swim Test

On Day 21 (24 h post-treatment), mice were placed individually in cylindrical beakers filled with water (21 ± 1°C; 15 cm depth) for 6 min. Immobility—passive floating with minimal movements to maintain head above water—was scored during minutes 2–6.

Synaptosome isolation and LC-MS/MS

Mice were euthanized and the mPFC including ACC, PL and IL were dissected on ice. Crude synaptosomes were isolated as described in [14] and stored at -80 °C. LC-MS/MS data were acquired on a Thermo Scientific Orbitrap Fusion mass spectrometer coupled to a Waters M-Class UPLC system. Protein identification and label-free quantification (LFQ) were performed in Proteome Discoverer (Thermo Fisher Scientific, version 3.2.0.450) using standard processing and consensus workflows. Normalization between samples was based on total peptide amount and protein abundance ratios were calculated based on pairwise peptide ratios. Background-based t-tests were performed to identify significant differences in

protein abundances between conditions. Peptide- and protein-level FDR thresholds were set to 1% and only proteins with at least two unique peptides and those present in >50% of samples were considered for downstream analysis. Canonical pathway enrichment graphs and hierarchical clustering of pathways were generated through QIAGEN IPA (QIAGEN Inc., https://digitalinsights.qiagen.com/IPA).

Results

Pharmacological inhibition of PDE9 attenuates psilocybin-induced head-twitch responses without affecting locomotor activity.

Mice received psilocybin (1 mg/kg, i.p.) 30 min after vehicle or PDE9 inhibitor PF-4181366 pretreatment (0.32, 1, 3.2, or 10 mg/kg), and HTR was recorded for 30 min. ANOVA revealed a significant main effect of pretreatment on psilocybin-induced HTR ($F_{(4,23)}$ = 6.042, p = 0.0018). PF-4181366 at 3.2 mg/kg and 10 mg/kg significantly reduced HTR relative to vehicle (p = 0.0097 and p = 0.0025, respectively), whereas lower doses produced nonsignificant trends (Tukey's multiple comparisons test, both p = 0.0532). Expressed as percentage of vehicle control, PF-4181366 reduced psilocybin-induced HTR by approximately 47% at 0.32 and 1 mg/kg, 59% at 3.2 mg/kg, and 69% at 10 mg/kg, demonstrating robust, dose-dependent attenuation. To exclude nonspecific motor suppression, locomotor activity was assessed following PF-4181366 administration. Analysis revealed no significant effect of pretreatment on distance traveled ($F_{(4,24)}$ = 0.2975, p = 0.8766), with no differences between vehicle and any PF-4181366 dose (all adjusted p ≥ 0.96). Normalization to baseline yielded comparable results ($F_{(4,24)}$ = 0.8831, p = 0.4888). These findings indicate that PDE9 inhibition selectively attenuates psilocybin-induced head-twitch behavior without affecting general locomotor activity, supporting specific modulation of $5\text{-HT}_2\text{AR}$-mediated behavioral effects.

PDE9 inhibition does not interfere with the antidepressant-like effects of psilocybin in a chronic stress model.

Mice subjected to CMMS showed decreased sucrose preference across groups. Antidepressant-like behavior was assessed 24 h post-treatment using the forced swim test (FST). ANOVA revealed a significant treatment effect on immobility ($F_{(2,57)} = 9.855$, $p = 0.0002$). Psilocybin significantly reduced immobility relative to saline-treated stressed controls ($p = 0.0008$). Critically, co-administration of PDE9 inhibitor did not attenuate this effect—the combination group similarly exhibited reduced immobility compared with saline controls ($p = 0.0004$).

Sucrose preference was assessed at baseline, post-CMMS, and post-treatment. Repeated-measures ANOVA revealed a significant effect of time ($p < 0.0001$), reflecting stress-induced reduction, but no treatment effect ($p = 0.5987$) or interaction ($p = 0.7661$).

These results indicate that psilocybin significantly reduces antidepressant-like behavior following chronic stress, and PDE9 inhibitor co-administration does not interfere with this effect. Sucrose preference was not significantly modulated by treatment at the examined time point.

Synaptosomal proteomics reveal that psilocybin/PDE9 inhibitor combination enhances synaptogenesis-related pathway compared to psilocybin alone.

Proteomic analysis of mPFC synaptosomal fractions collected 24h post-treatment revealed that PDE9i–psilocybin significantly altered synaptic protein composition relative to psilocybin alone. Pathway analysis showed an overall pattern of pathway downregulation with the PDE9i–psilocybin combination, including GPCR and serotonin receptor signaling. Remarkably, synaptogenesis, a process that is believed to be crucial for the therapeutic effects of psychedelics, was upregulated in the combination group. Pathways that were similarly regulated by PDE9i-psilocybin and psilocybin alone included CREB signaling, S100 family signaling and the NF1-Ras pathway. These pathways may provide mechanistic insight into the downstream pathways involved in psilocybin's therapeutic efficacy.

Conclusions

The present findings demonstrate that pharmacological inhibition of PDE9 selectively attenuates the acute head-twitch response to psilocybin without compromising its antidepressant-like effects in a chronic stress model. This behavioral dissociation is mechanistically supported by synaptosomal proteomic analysis showing preserved synaptogenic signaling pathways in the combination group, consistent with maintained therapeutic potential. The ability to reduce acute 5-HT$_2$A receptor-mediated behavioral responses while preserving antidepressant efficacy suggests that PDE9 inhibition may represent a viable strategy for improving the clinical tolerability of psilocybin. Notably, the 24-hour post-treatment timepoint selected for synaptosomal proteomics aligns with the established onset of psilocybin-induced neuroplastic consolidation in humans, where neuroimaging studies report changes in brain network connectivity and clinical symptom relief emerging within 24 hours of dosing and persisting for at least one month—providing translational grounding for the molecular window captured here [15]. It is possible that the increase in cGMP following PDE9 inhibition functions to block psilocybin-induced hyperexcitability and thereby reduce head-twitch responses. Collectively, these results support the broader hypothesis that combination-based approaches targeting intracellular signaling cascades can modulate the acute subjective profile of psychedelics, improve their tolerability while preserving their therapeutic mechanisms, thus making them scalable.

A Pre-treatment   PSI 1 mg/kg   HTR

Veh/PDE9i T_0 30 min
−30 min

HTR BOX

B Head-Twitch Response
Psilocybin 1 mg/kg

HTR count (30 min)

VEH 0.32 1 3.2 10
Pre-treatment PF-4181366 (mg/kg)

C Baseline   PDE9i   Locomotion

−60 min T_0 60 min

Open Field   Open Field

D Post-Treatment
Locomotor activity

Total distance traveled (m/1h)

VEH 0.32 1 3.2 10
Pre-treatment PF-4181366 (mg/kg)

E SPT SPT CMMS (15 days) SPT TST SPT FST
                                              Drug Proteomic

1   2–3   4   18 19   20   21
            Time (d)   24 h   24 h

F Forced Swim

Immobility (sec)

Treatment

- Veh-Veh
- Veh-Psilo
- PDE9i-Psilo

G Sucrose Preference

Sucrose preference (%)

Baseline CRS Treatment

- Veh-Veh
- Veh-Psilo
- PDE9i-Psilo

H

Saline or   24 hrs   mPFC isolation   Synaptosome
7.8 mg/kg MDMA   enrichment

LC/MS/MS

I Psilo+PDE9i vs Psilo

Log2(Fold Change)

−log10(pvalue)

248 DEPs
91 up ●
157 down ●

J Psilo+PDE9i vs Psilo

Synaptogenesis Signaling Pathway
Oxytocin in Brain Signaling Pathway
Orexin Signaling Pathway
NF1 RAS Signaling Pathway
Serotonin Receptor Signaling
Adrenergic Receptor Signaling Pathway
cAMP-mediated signaling
Protein Kinase A Signaling
DHCR24 Signaling Pathway
GABAergic Receptor Signaling Pathway
Glutaminergic Receptor Signaling Pathway
CREB Signaling in Neurons
Acetylcholine Receptor Signaling Pathway
Opioid Signalling
G-Protein Coupled Receptor Signaling

z-score
−2.5   +2.5

−log(pvalue)

K

GPCR signaling
CREB signaling
Serotonin receptor signaling
S100 family pathway
G alpha (i) signaling
Adrenergic signaling
NF1 RAS pathway
Gap junction signaling
Ephrin receptor signaling
Oxytocin signaling
G beta gamma pathway

Psilo vs Sal
Psilo+PDE9i vs Sal
Psilo+PDE9i vs Psilo

Figure 1. PDE9 inhibition attenuates psilocybin-induced head-twitch responses without affecting locomotor activity or antidepressant efficacy, and preserves synaptogenic pathway engagement.

(A) Experimental timeline for head-twitch response (HTR) assessment. Mice received PF-4181366 or vehicle 30 minutes prior to psilocybin (1 mg/kg, i.p.) and HTR was recorded for 30 minutes in a dedicated HTR box. (B) Total HTR counts over 30 minutes showing dose-dependent attenuation following PF-4181366 pretreatment. Significant reductions were observed at 3.2 and 10 mg/kg compared to vehicle (**$p<0.01$; n=5–8/group). (C) Experimental timeline for locomotor activity assessment. Mice were placed in an open field 60 minutes before PF-4181366 treatment (baseline) and re-assessed for 60 minutes post-treatment. (D) Total distance traveled post-treatment showing no effect of PF-4181366 at any dose tested (n=5–7/group). (E) Experimental timeline for chronic multimodal stress (CMMS) and behavioral testing. Mice underwent SPT training (day 1), baseline SPT (days 2–3), 15 days of CMMS (days 4–18) with SPT on days 17–18, followed by TST (day 19), drug treatment with SPT (day 20), and FST (day 21; n=20/group). A separate cohort was used for synaptosomal proteomic analysis (n=6/group). (F) FST immobility time. Both psilocybin alone (Veh-Psilo) and in combination with PDE9 inhibitor (PDE9i-Psilo) significantly reduced immobility compared to vehicle-treated stressed controls (*** $p<0.001$). (G) Sucrose preference across baseline, post-stress (CRS), and post-treatment timepoints. (H) Schematic workflow for synaptosomal proteomic analysis. mPFC tissue was collected 24 h post-treatment, synaptosomal fractions were isolated, and proteins were quantified by LC/MS/MS. (I) Volcano plot of differentially expressed proteins (DEPs) in the psilocybin-PF-4181366 (Psilo+PDE9i) group versus Psilo alone (248 DEPs total; 91 upregulated, red; 157 downregulated, blue; n=6/group). (J) Ingenuity Pathway Analysis (IPA) of the Psilo+PDE9i vs. Psilo comparison showing significantly regulated canonical pathways ranked by −log(p-value). Bar color reflects predicted activation (orange) or inhibition (blue) based on z-score (scale: −2.5 to +2.5). (K) Heatmap comparing z-score-based pathway regulation across three contrasts (Psilo vs. Sal, Psilo+PDE9i vs. Sal, Psilo+PDE9i vs. Psilo). Data are

presented as mean ± SEM. HTR and locomotor data were analyzed by one-way ANOVA followed by Tukey's multiple comparisons test. Forced swim data were analyzed by one-way ANOVA with Dunnett's post hoc test. Sucrose preference was analyzed using two-way repeated-measures ANOVA.

Disclosures: This work was supported by a research contract from Freedom Biosciences (A.P.K., C.P., G.F.). Drs. Kaye, Krystal, and Pittenger are co-inventors on a patent application related to psychedelics. Dr. Kaye previously received contracted research funding from Transcend Therapeutics. Drs. Jefferson and Pittenger previously received research funding from Freedom Biosciences. Dr. De Aquino has received research support from Jazz Pharmaceuticals and Ananda Scientific and has served as a paid consultant for Boehringer Ingelheim. In the past three years, Dr. Pittenger has consulted for Biohaven Pharmaceuticals, Freedom Biosciences, Transcend Therapeutics, UCB BioPharma, Mind Therapeutics, Ceruvia Biosciences, F-Prime Capital Partners, and Madison Avenue Partners; has received research support from Biohaven Pharmaceuticals, Freedom Biosciences, and Transcend Therapeutics; owns equity in Alco Therapeutics, Mind Therapeutics, and Lucid/Care; receives royalties from Oxford University Press and UpToDate; and holds pending patents on pathogenic antibodies in pediatric OCD and on novel mechanisms of psychedelic drugs. Dr. Menniti has served as a consultant for Freedom Biosciences and is a co-inventor on patents claiming the use of PDE9 inhibitors in combination with psychedelic drugs. Dr. Krystal has served as a consultant for Aptinyx, Inc.; Biogen, Idec, MA; Bionomics, Limited (Australia); Boehringer Ingelheim International; Clearmind Medicine, Inc.; Cybin IRL (Ireland Limited Company); Enveric Biosciences; Epiodyne, Inc.; EpiVario, Inc.; Janssen Research & Development; Jazz Pharmaceuticals, Inc.; Otsuka America Pharmaceutical, Inc.; Perception Neuroscience, Inc.; Praxis Precision Medicines, Inc.; Spring Care, Inc.; Sunovion Pharmaceuticals, Inc. Dr. Krystal has served as a scientific advisory board member for: Biohaven Pharmaceuticals; BioXcel Therapeutics, Inc. (Clinical Advisory Board); Cerevel Therapeutics, LLC; Delix Therapeutics, Inc.; Eisai, Inc.; EpiVario, Inc.; Freedom Biosciences, Inc.; Jazz Pharmaceuticals, Inc.; Neumora Therapeutics, Inc.; Neurocrine Biosciences, Inc.; Novartis Pharmaceuticals Corporation; Praxis Precision Medicines, Inc.; PsychoGenics, Inc.; Tempero Bio, Inc.; Terran Biosciences, Inc. In the past 3 years, Dr. Krystal is an inventor on patents licensed by Yale University to Janssen Pharmaceuticals, Biohaven Pharmaceuticals, Spring Health, Freedom Biosciences, and Novartis Pharmaceuticals. Dr. Kwan has been a scientific advisor or consultant for Boehringer Ingelheim, Eli Lilly, Empyrean Neuroscience, Freedom Biosciences, Otsuka, and Xylo Bio. Dr. Kwan has received research support from Intra-Cellular Therapies. The remaining authors declare no conflict of interest.

Bibliography & References Cited

1. Friedrich MJ. Depression Is the Leading Cause of Disability Around the World. JAMA. 2017 Apr 18;317(15):1517. doi: 10.1001/jama.2017.3826. PMID: 28418490.
2. Goodwin GM, Aaronson ST, Alvarez O, Arden PC, Baker A, Bennett JC, Bird C, Blom RE, Brennan C, Brusch D, Burke L, Campbell-Coker K, Carhart-Harris R, Cattell J, Daniel A, DeBattista C, Dunlop BW, Eisen K, Feifel D, Forbes M, Haumann HM, Hellerstein DJ, Hoppe AI, Husain MI, Jelen LA, Kamphuis J, Kawasaki J, Kelly JR, Key RE, Kishon R, Knatz Peck S, Knight G, Koolen MHB, Lean M, Licht RW, Maples-Keller JL, Mars J, Marwood L, McElhiney MC, Miller TL, Mirow A, Mistry S,

Mletzko-Crowe T, Modlin LN, Nielsen RE, Nielson EM, Offerhaus SR, O'Keane V, Páleníček T, Printz D, Rademaker MC, van Reemst A, Reinholdt F, Repantis D, Rucker J, Rudow S, Ruffell S, Rush AJ, Schoevers RA, Seynaeve M, Shao S, Soares JC, Somers M, Stansfield SC, Sterling D, Strockis A, Tsai J, Visser L, Wahba M, Williams S, Young AH, Ywema P, Zisook S, Malievskaia E. Single-Dose Psilocybin for a Treatment-Resistant Episode of Major Depression. N Engl J Med. 2022 Nov 3;387(18):1637-1648. doi: 10.1056/NEJMoa2206443. PMID: 36322843.

3. Siegel JS, Subramanian S, Perry D, Kay BP, Gordon EM, Laumann TO, Reneau TR, Metcalf NV, Chacko RV, Gratton C, Horan C, Krimmel SR, Shimony JS, Schweiger JA, Wong DF, Bender DA, Scheidter KM, Whiting FI, Padawer-Curry JA, Shinohara RT, Chen Y, Moser J, Yacoub E, Nelson SM, Vizioli L, Fair DA, Lenze EJ, Carhart-Harris R, Raison CL, Raichle ME, Snyder AZ, Nicol GE, Dosenbach NUF. Psilocybin desynchronizes the human brain. Nature. 2024 Aug;632(8023):131-138. doi: 10.1038/s41586-024-07624-5. Epub 2024 Jul 17. PMID: 39020167; PMCID: PMC11291293.

4. Cameron LP, Tombari RJ, Lu J, Pell AJ, Hurley ZQ, Ehinger Y, Vargas MV, McCarroll MN, Taylor JC, Myers-Turnbull D, Liu T, Yaghoobi B, Laskowski LJ, Anderson EI, Zhang G, Viswanathan J, Brown BM, Tjia M, Dunlap LE, Rabow ZT, Fiehn O, Wulff H, McCorvy JD, Lein PJ, Kokel D, Ron D, Peters J, Zuo Y, Olson DE. A non-hallucinogenic psychedelic analogue with therapeutic potential. Nature. 2021 Jan;589(7842):474-479. doi: 10.1038/s41586-020-3008-z. Epub 2020 Dec 9. PMID: 33299186; PMCID: PMC7874389.

5. Wallach J, Cao AB, Calkins MM, Heim AJ, Lanham JK, Bonniwell EM, Hennessey JJ, Bock HA, Anderson EI, Sherwood AM, Morris H, de Klein R, Klein AK, Cuccurazzu B, Gamrat J, Fannana T, Zauhar R, Halberstadt AL, McCorvy JD. Identification of 5-HT$_{2A}$ receptor signaling pathways associated with psychedelic potential. Nat Commun. 2023 Dec 15;14(1):8221. doi: 10.1038/s41467-023-44016-1. PMID: 38102107; PMCID: PMC10724237.

6. Cao D, Yu J, Wang H, Luo Z, Liu X, He L, Qi J, Fan L, Tang L, Chen Z, Li J, Cheng J, Wang S. Structure-based discovery of nonhallucinogenic psychedelic analogs. Science. 2022 Jan 28;375(6579):403-411. doi: 10.1126/science.abl8615. Epub 2022 Jan 27. PMID: 35084960.

7. van Staveren WC, Glick J, Markerink-van Ittersum M, Shimizu M, Beavo JA, Steinbusch HW, de Vente J. Cloning and localization of the cGMP-specific phosphodiesterase type 9 in the rat brain. J Neurocytol. 2002 Sep-Nov;31(8-9):729-41. doi: 10.1023/a:1025704031210. PMID: 14501210.

8. Hutson PH, Finger EN, Magliaro BC, Smith SM, Converso A, Sanderson PE, Mullins D, Hyde LA, Eschle BK, Turnbull Z, Sloan H, Guzzi M, Zhang X, Wang A, Rindgen D, Mazzola R, Vivian JA, Eddins D, Uslaner JM, Bednar R, Gambone C, Le-Mair W, Marino MJ, Sachs N, Xu G, Parmentier-Batteur S. The selective phosphodiesterase 9 (PDE9) inhibitor PF-04447943 (6-[(3S,4S)-4-methyl-1-(pyrimidin-2-ylmethyl)pyrrolidin-3-yl]-1-(tetrahydro-2H-pyran-4-yl)-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one) enhances synaptic plasticity and cognitive function in rodents. Neuropharmacology. 2011 Sep;61(4):665-76. doi: 10.1016/j.neuropharm.2011.05.009. Epub 2011 May 19. PMID: 21619887.

9. Kleiman RJ, Chapin DS, Christoffersen C, Freeman J, Fonseca KR, Geoghegan KF, Grimwood S, Guanowsky V, Hajós M, Harms JF, Helal CJ, Hoffmann WE, Kocan GP, Majchrzak MJ, McGinnis D, McLean S, Menniti FS, Nelson F, Roof R, Schmidt AW, Seymour PA, Stephenson DT, Tingley FD, Vanase-Frawley M, Verhoest PR, Schmidt CJ. Phosphodiesterase 9A regulates central cGMP and modulates responses to cholinergic and monoaminergic perturbation in vivo. J Pharmacol Exp Ther. 2012 May;341(2):396-409. doi: 10.1124/jpet.111.191353. Epub 2012 Feb 10. PMID: 22328573.

10. Verhoest PR, Proulx-Lafrance C, Corman M, Chenard L, Helal CJ, Hou X, Kleiman R, Liu S, Marr E, Menniti FS, Schmidt CJ, Vanase-Frawley M, Schmidt AW, Williams

RD, Nelson FR, Fonseca KR, Liras S. Identification of a brain penetrant PDE9A inhibitor utilizing prospective design and chemical enablement as a rapid lead optimization strategy. J Med Chem. 2009 Dec 24;52(24):7946-9. doi: 10.1021/jm9015334. PMID: 19919087.

11. Shao LX, Liao C, Gregg I, Davoudian PA, Savalia NK, Delagarza K, Kwan AC. Psilocybin induces rapid and persistent growth of dendritic spines in frontal cortex in vivo. Neuron. 2021 Aug 18;109(16):2535-2544.e4. doi: 10.1016/j.neuron.2021.06.008. Epub 2021 Jul 5. PMID: 34228959; PMCID: PMC8376772.

12. Griffiths RR, Johnson MW, Richards WA, Richards BD, McCann U, Jesse R. Psilocybin occasioned mystical-type experiences: immediate and persisting dose-related effects. Psychopharmacology (Berl). 2011 Dec;218(4):649-65. doi: 10.1007/s00213-011-2358-5. Epub 2011 Jun 15. PMID: 21674151; PMCID: PMC3308357.

13. Hesselgrave N, Troppoli TA, Wulff AB, Cole AB, Thompson SM. Harnessing psilocybin: antidepressant-like behavioral and synaptic actions of psilocybin are independent of 5-HT2R activation in mice. Proc Natl Acad Sci U S A. 2021 Apr 27;118(17):e2022489118. doi: 10.1073/pnas.2022489118. PMID: 33850049; PMCID: PMC8092378.

14. Li N, Lee B, Liu RJ, Banasr M, Dwyer JM, Iwata M, Li XY, Aghajanian G, Duman RS. mTOR-dependent synapse formation underlies the rapid antidepressant effects of NMDA antagonists. Science. 2010 Aug 20;329(5994):959-64. doi: 10.1126/science.1190287. PMID: 20724638; PMCID: PMC3116441.

15. Carhart-Harris RL, Roseman L, Bolstridge M, Demetriou L, Pannekoek JN, Wall MB, Tanner M, Kaelen M, McGonigle J, Murphy K, Leech R, Curran HV, Nutt DJ. Psilocybin for treatment-resistant depression: fMRI-measured brain mechanisms. Sci Rep. 2017 Oct 13;7(1):13187. doi: 10.1038/s41598-017-13282-7. PMID: 29030624; PMCID: PMC5640601.