Biobank-scale genotyping of Robertsonian translocations reveals hidden structural variation on the human acrocentric chromosomes

Arang Rhie[1*], Juhyun Kim[1*], Francisco Rodriguez-Algarra[2,3], Steven J. Solar[1], Sergey Koren[1], Dmitry Antipov[1], Caralynn M. Wilczewski[4], George L. Maxwell[5], Jennifer Gerton[6], Justin Paschall[4], Tamara Potapova[6], Tyra G. Wolfsberg[7], Sumeeta Singh[7], Sandra O. del Castillo del Rio[2,3], Human Pangenome Reference Consortium, Clesson Turner[4], Vardhman K. Rakyan[2,3], Adam M. Phillippy[1]

[1]Genome Informatics Section, Center for Genomics and Data Science Research, NHGRI, NIH, Bethesda, MD, USA
[2]The Blizard Institute, School of Medicine and Dentistry, Queen Mary University of London, London, UK
[3]Centre for Epigenetics, Queen Mary University of London, London, UK
[4]Reverse Phenotyping Core, Center for Precision Health Research, NHGRI, NIH, Bethesda, MD, USA
[5]Women's Health Integrated Research Center, Inova Health System, Falls Church, VA, USA
[6]Stowers Institute for Medical Research, Kansas City, MO, USA
[7]Bioinformatics and Scientific Programming Core, Office of Scientific Core Facilities, NHGRI, NIH, Bethesda, MD, USA
*Contributed equally

Summary

Balanced Robertsonian translocation (ROB) is the most common chromosomal rearrangement in humans, with an estimated occurrence of 1 in 800 in newborn studies. Carriers are at increased risk of cancer and often diagnosed at fertility clinics after facing recurrent miscarriages, infertility, or aneuploid offspring. Genotyping carriers with DNA sequencing has been challenging because of gaps and misrepresentation of the translocation fusion site in the human reference genome. Only recently, telomere-to-telomere (T2T) human genomes successfully revealed sequences of the acrocentric short arms, including the most common ROB fusion site. A ROB results in loss of two ribosomal DNA (rDNA) arrays and its adjacent distal sequences, including the highly conserved distal junction (DJ). Here, we present a novel method to type ROB carriers directly from short sequencing reads by estimating DJ copy number. We demonstrate that our method successfully genotypes ROBs using a reference-free approach or alignments to either T2T-CHM13v2 or GRCh38. Applying the method to a cohort of healthy newborns and family members (n=4,172) as well as the UK Biobank (n=490,416), we find candidate ROBs at a frequency consistent with the previously reported 1 in 800 incidence (0.11-0.12%). In addition to ROB carriers, we report the frequency of one DJ loss (9, 2.8-3.4%) or gain (11+, 8.4-9.3%) from the two cohorts and the 1000 Genomes Project (n=3,202), and characterize the underlying structural variation in near-T2T genome assemblies from the Human Pangenome Reference Consortium. Importantly, our method provides the first sequencing-based diagnostic for Robertsonian chromosomes and can be applied to low-coverage sequencing data, enhancing its clinical applicability and enabling new studies of structural variation on the acrocentric chromosomes.

Keywords: Balanced Robertsonian translocation, Distal junction of the rDNA, Structural variation, acrocentric chromosomes, telomere-to-telomere genomes

INTRODUCTION

Robertsonian translocation is a type of chromosomal fusion involving acrocentric or telocentric chromosomes. In humans, Robertsonian chromosomes (ROBs) occur at a rate of 1 in 800 individuals[1], with the most common form of translocation found between chromosome 14 and either chromosome 13 or 21, accounting for 75% and 10% of cases, respectively[2,3]. If a person carries a ROB with two normal chromosomes to pair, it is considered a "balanced translocation" (hereafter shortened as "ROB carrier") as opposed to monosomy or trisomy. For example, one fused ROB chromosome between 14 and 21 is still able to pair with the unaffected copies of 14 and 21 during meiosis. However, such carriers are at increased risk of childhood leukemia, non-Hodgkin lymphoma and breast cancer[4]. The carriers are also at increased risk to produce gametes with an unbalanced karyotype, often leading to spontaneous miscarriage, infertility, and uniparental disomy or trisomy in the offspring[5]. The extra 21q copy from an inherited ROB chromosome has been reported to cause 2-4% of Down syndrome cases[6]. Similarly but less frequently, an additional 13q copy can be transmitted from a ROB chromosome, causing Patau syndrome[7].

The exact mechanism and breakpoint of the fusion site forming the most common type of ROB remained unknown until recently. In the past, ROB formation was classified as a centromeric fusion[8] or referred to as a fusion of unknown microsatellites[9]. The first complete telomere-to-telomere (T2T) human genome revealed the complex structure of the acrocentric p-arms, including regions with high sequence homology[10]. In a follow-up study, using human pangenome assemblies[11], frequent inter-chromosomal exchange was inferred within so-called pseudo-homolog regions (PHR)[12]. A particular PHR on the proximal short arms of Chrs. 13, 14, and 21 was suggested to mediate ROB translocations[13], which was later confirmed by the T2T sequencing and assembly of three ROB cell lines[14]. Such ROB formation results in the loss of one fusion site and sequences distal to it, which includes the rDNA gene array on each chromosome and the highly conserved distal junction (DJ) sequence of each rDNA array[12,14,15].

Until now, the most widely used method for identifying ROB carriers relies on karyotyping, fluorescence in situ hybridization (FISH)[16] or Hi-C[17]. However, these methods depend on live cells or intact tissues and are labor intensive. Thus, these methods may not be applicable for routine diagnostics or large-scale cohorts. Microarray or SNP based chips have also been developed to detect ROB carriers[18]. However, the lack of understanding of the exact ROB

breakpoint and gaps in the prior human reference genomes (i.e. both GRCh38 and hg19)
have posed a barrier to using whole genome sequencing (WGS) reads to detect ROB
carriers[19].

Recognizing that ROBs result in the predictable loss of acrocentric sequence along with the
increasing large-scale cohorts sequenced with WGS available, we sought to explore methods
to identify candidate ROB carriers from such datasets. Here, we show three ways to detect
ROBs from short-read sequencing data by using (1) T2T-CHM13/hs1 as an alternate
reference for read alignment, (2) pre-existing read alignments mapped to GRCh38, and (3) DJ
marker *k*-mers and their multiplicity directly obtained from sequencing reads. We
benchmarked these methods to reference genomes CHM13 and HG002 as controls and three
known ROB cell lines as cases[10,14]. Next, we applied the method on WGS in the Reverse
Phenotyping Core (RPC) cohort (n=4,172)[20,21] and the UK Biobank WGS cohort (n=490,444)[22].
We report that these methods identify potential ROB carriers in all cohorts at approximately
the expected frequency of 1 in 800. Finally, we typed all samples from the 1000 Genomes
Project (1KGP)[23] and further examined samples with atypical copies of the DJ in the recent,
near-T2T HPRC Release 2 assemblies[24]. We propose this method as a routine screening
approach for identifying potential ROB carriers in any whole-genome sequencing dataset.

RESULTS

Detecting ROBs by the loss of rDNA distal junctions

We first identified the most stable, highly conserved region to target for copy number
estimates. The most common form of ROB translocation is reported to have a breakpoint in
the proximal short arm, between the rDNA and centromere[14]. Therefore, ROB carriers are
expected to be missing two rDNA arrays and all sequences distal to these arrays (**Fig. 1A**).
Because of the difficulty in detecting the loss of rDNA units due to their highly variable copy
number, we targeted the region immediately distal of the rDNA array, the DJ, which had been
reported to be highly conserved in humans and expected to be in a single copy per
chromosome[15]. Surveying the copy number estimates from the Simons Genome Diversity
Project[10,25], we confirmed that most of the DJ region maintains a stable 10 copy pattern across
individuals regardless of ancestral background, suggesting the presence of a highly

conserved, single DJ per rDNA array (**Fig. 1B**). Thus, we hypothesized that two DJ copies will

be lost following most Robertsonian translocations, resulting in eight DJs in the ROB carriers.

Using this hypothesis, we developed a pipeline for read-mapping based copy number

estimates utilizing a masked T2T-CHM13 reference (**Fig. 1C**). Specifically, we masked all but

one copy of the DJ in the T2T-CHM13 reference and compared the coverage against the

background coverage collected from the rest of the non-heterochromatic autosomal regions.

In addition, we masked all but one copy of the 45S rDNAs and Robertsonian-associated

PHRs to estimate copy number for these regions as well (**Table S1 and Fig. S1-4**).

To expand the pipeline on large-scale cohorts already mapped to GRCh38, we sought to

look for the presence of the target DJ sequence in GRCh38. When mapping GRCh38 to each

target sequence extracted from CHM13, we identified nine DJ copies on GRCh38 chr21

organized as a repeat array with some inverted and incomplete copies interspersed and

surrounded with gaps (**Fig. 1D**). This is an arrangement seen in no T2T-assembled genome

since, and represents a clear mis-assembly in GRCh38. In addition, partial DJ sequences

were found on two unplaced sequences (chrUn_GL000195v1 and chrUn_GL000220v1) and

two unlocalized sequences (chr22_KI270733v1_random and chr17_GL000205v2_random),

with high sequence identity (average 96.2%) (**Tables S2-3**). Diverged pieces of DJs were

found on Chrs. 4 and 10, similar to those observed in CHM13, however these were excluded

from targeting due to their low sequence identity (<83.4%) and low coverage (<2.8%).

In addition to the mapping-based method, we also developed a reference-free approach to

reduce any potential mapping bias. To select target k-mers for the DJ, all k-mers (k=31) in

T2T-CHM13v2 were evaluated for their copy number multiplicity observed in Illumina reads

derived from CHM13. Among the k-mers found at least once in each of the five DJs (Chrs. 13,

14, 15, 21 and 22), target k-mers were selected from the ~10 copy peak to exclude highly

repetitive or variable regions (**Fig. 1E**). To estimate the copy number of the DJs for each new

sample, all k-mers are collected directly from the FASTQ files. The two-copy peak is then

extracted from the entire k-mer histogram and compared to the median multiplicity of the

target DJ k-mers. Since this method operates directly on FASTQ files, it eliminates the need

for a mapping process, thus reducing computational resource requirements.

To test the pipelines, we used Illumina reads from CHM13, HG002 (GM24385)—for which

10 DJs were confirmed[10,26]—and ROB cell lines GM03417, GM03786, and GM04890—for

which 8 DJs and ROBs were confirmed by T2T assemblies and FISH karyotyping by de Lima

et al.[14] (**Fig. 2**). We applied our mapping-based method using both GRCh38 and CHM13

reference genomes, utilizing BAM files aligned to each reference, as well as our *k*-mer-based

method using raw sequencing reads. All three methods consistently reported ~10 DJs for

both CHM13 and HG002 and ~8 DJs for the 3 known ROB cell lines, as expected (**Table 1,**

Fig. 2 and **Fig. S5**).

Fig. 1 | Target rDNA distal junction (DJ) for ROB carrier detection.

(A) Schematic overview of the most common karyotype of ROB carriers. Chr. 14 pseudo-homolog region (PHR)
has been identified in its inverted orientation compared to those on Chrs. 13 and 21, reported as the fusion site for
the most common forms of ROBs. When a ROB is formed through a fusion between two acrocentric chromosomes,
e.g. Chrs. 14 (lime) and 13 (red), one small segment carrying one PHR and two DJs gets lost. Legends follow as in
panel C. (B) Copy number estimates of the DJ shows conserved 10-20 copies across individuals from various
ancestral backgrounds in the Simons Genome Diversity Project. No other suitable regions on the short arms were
found with a stable copy number. Copy numbers for GRCh38 (at bottom) indicate this region is mis-assembled in
GRCh38. SGDP population labels: AFR, Africa; AMR, America; EAS, East Asia; OCN, Oceania; SAS, South Asia;
SIB, Siberia; WEA, Western Eurasia. (C) Masked regions of the T2T-CHM13v2.0 acrocentric p-arms for copy
number estimation of the DJ, rDNA, and PHR. Only one copy of each in Chr. 13 has been chosen as the target for
copy number estimates. (D) Identifying DJ target region for copy number estimates for sequencing read alignments
on GRCh38. Nine mis-assembled copies of the target were found on Chr. 21 surrounded by gaps, two copies on
unplaced contigs and two on unlocalized contigs of Chrs. 17 and 22. (E) Reference free DJ copy number estimates.
A *k*-mer found in all five DJs in T2T-CHM13v2.0 were selected as "DJ *k*-mers". Based on the *k*-mer copy number
count in Illumina reads (left), *k*-mers with ~10× coverage were selected as "DJ target *k*-mers". The *k*-mer
multiplicity on CHM13 Chr. 13 shows a stable 10× coverage across the region for the DJ target *k*-mers.

Fig. 2 | FISH karyotyping of the control cell lines.

(A) Control cell lines CHM13 and GM24385 (HG002). All 10 acrocentric haplotypes show the presence of rDNA (green) and DJ (magenta). (B) Previously reported ROB cell lines, which the breakpoint has been identified as the PHR in de Lima et al.[14] The ROB chromosomes are highlighted, showing the absence of the DJ.

Table 1. Pipeline confirms 8 DJs in 3 known ROB cell lines and two control samples, where the DJ was karyotyped before. Remarks are from Coriell. *Reported from de Lima et al.[14]

Sample	Sex	T2T-CHM13	GRCh38	k-mer	Remarks	Age
CHM13	XX	10.3	10.1	10.2	CHM13hTERT cell line, karyotypically normal	na
HG002	XY	10.1	10.7	10.1	GM24385 cell line, karyotypically normal	na
GM03417	XX	8.3	8.4	8.3	Clinical features of Down syndrome. Mosaic; 32% of cells are balanced 45,XX, t(14;21); *Cellular heterogeneity with regard to the number of copies of Chr. 21, with 3 copies of Chr. 21 being relatively rare (~10%)	15
GM03786	XX	8.0	8.1	8.0	Clinically normal	19
GM04890	XX	8.1	8.3	8.1	Clinically normal; has had 5 miscarriages	32

One in 800 ROB candidates in newborn family cohort

The Reverse Phenotyping Core (RPC) cohort includes 1,325 families with healthy newborns, parents, and siblings (where available), totalling 4,187 samples. Blood samples were collected at the time of birth. Illumina short read data aligned to hg19 from Wilczewski *et al.*[21] and Bodian *et al.*[20] were extracted and re-aligned to the masked version of T2T-CHM13v2.0 and evaluated for the copy number of the DJs, PHRs and rDNAs (**Fig. 3**). In total, we found 4 individuals with estimated 8 copies of the DJs (**Fig. 3A**). This includes two parents and one mother-newborn (daughter) pair. (**Table 2**). Counting the 3 out of 2,650 presumably unrelated parents, among which 2,612 were sequenced, yields an estimated ratio of 1 in 871-883 (0.11%), matching closely the expected 1 in 800 (0.125%) incidence[1]. Among the 2,612 parents, 2,277 (87.17%) appeared as having 10 estimated copies, comprising the majority of the individuals (**Fig. 3A**). Interestingly, regardless of with or without the offspring, we found 3.34-3.45% individuals with 9 DJs, and 9.26-9.27% with more than 10 DJs, with 13 being the most copies observed (**Fig. 3 and Table 3**). When comparing the number of samples classified in each DJ copy bin, no significant difference was observed between XX and XY individuals (χ^2 test of independence, *p*>0.5). We observed a few more 8 DJs in XX carriers (3 vs. 1), with one passed down from the mother to daughter, matching previous reports that ROBs are more frequently observed in XX samples and transmitted to XX offspring[27]. When comparing the distribution of DJ copy number between parents and offspring, we observed a slight shift toward higher copy numbers in the parental genomes (Mann–Whitney U test, p = 3.7×10^{-6}; **Fig. 3B and Fig. S6**).

Fig. 3 | DJ, PHR, and rDNA copy number estimation using T2T-CHM13, GRCh38, and *k*-mer based methods for the RPC cohort.

(A) Diploid copy number estimation of DJ, PHR, and rDNA in the RPC cohort using CHM13-based mapping approach. (B) Histogram of the probability density of DJ and PHR copy numbers across samples, stratified by parents and newborns. The area under each group's histogram sums to one. A two-sided Mann–Whitney U test was performed using raw counts to compare parents and newborns, and the rank-biserial correlation (RBC) is reported as the effect size. (C) Each point represents a sample from the RPC cohort. No obvious correlations were

observed between the DJ, PHR, and rDNA copy numbers. (D) Comparison between DJ counts derived from mapping-based estimates (CHM13 and GRCh38) and the *k*-mer–based method using subsampled samples (n = 92), showing strong correlation. Density levels are shown with four contour levels.

Table 2. ROB candidates from the RPC cohort. Estimated num. of DJs, PHRs, and rDNAs are shown below with the categorized copy counts binned. ROB candidates are highlighted with a green background.

Family	Sex	DJ	Bin	PHR	Bin	rDNA	Relationship
A	XX	8.16	8	3.87	4	386.6	Mother
	XY	10.09	10	4.85	5	433.8	Father
	XY	8.88	9	4.71	5	415.6	Child
B	XX	10.2	10	6.00	6	338.3	Mother
	XY	8.04	8	4.04	4	229.6	Father
	XX	9.10	9	5.40	5	357.7	Child-A
	XX	8.95	9	5.31	5	334.8	Child-B
C	XX	8.05	8	4.77	5	254.7	Mother
	XY	9.04	9	4.85	5	325.6	Father
	XX	8.04	8	5.80	6	316.2	Child-A
	XX	9.15	9	4.90	5	424.9	Child-B

Table 3. Copy number estimates of the DJs across the RPC cohort (n=4,187) and UK Biobank WGS cohort (n=490,369). Regardless of the cohorts, estimated ROBs with 8 DJs had the smallest fraction of 0.11-0.12%. The largest fraction of samples were observed with 10 DJs (87.17-88.70%), followed by 11 DJs (7.81-9.00%), 9 DJs (2.81-3.45%), and more than 11 DJs (0.26-0.56%), indicating loss of one or gains of one or more DJs are relatively common.

Cohort	Num. of DJs	Num. of samples	%	XX	%	XY	%
RPC	<8	0	0.00%	0	0.00%	0	0.00%
	8	**4**	**0.10%**	**3**	**0.14%**	**1**	**0.05%**
	9	140	3.34%	81	3.82%	59	2.86%
	10	3,659	87.39%	1,848	87.05%	1,811	87.74%
	11	373	8.91%	186	8.76%	187	9.06%
	>11	11	0.26%	5	0.24%	6	0.29%
	Total	4,187	100.00%	2,123	100.00%	2,064	100.00%
RPC (Parents only)	<8	0	0.00%	0	0.00%	0	0.00%
	8	**3**	**0.11%**	**2**	**0.15%**	**1**	**0.08%**
	9	90	3.45%	51	3.91%	39	2.98%
	10	2,277	87.17%	1,135	87.04%	1,142	87.31%
	11	235	9.00%	113	8.67%	122	9.33%
	>11	7	0.27%	3	0.23%	4	0.31%
	Total	2,612	100.00%	1,304	100.00%	1,308	100.00%
UK Biobank	<8	9	0.002%	6	0.002%	3	0.00%
	8	**583**	**0.12%**	**312**	**0.12%**	**271**	**0.12%**
	9	13,778	2.81%	7,460	2.80%	6,318	2.82%
	10	434,951	88.70%	235,927	88.70%	199,024	88.70%
	11	38,291	7.81%	20,791	7.82%	17,500	7.80%
	>11	2,757	0.56%	1,500	0.56%	1,257	0.56%
	Total	490,369	100.00%	265,996	100.00%	224,373	100.00%

However, when comparing the number of PHRs, we observed a broader polymorphic copy number distribution compared to the DJs (**Fig. 3A**). Initially, we expected that the majority of individuals would carry six copies of the PHRs as observed in CHM13 and HG002, and five copies for those with 8 DJs due to the ROB as reported in Guarracino et al.[12] and de Lima et

al.[14]. A little over half the samples were observed as 6 copies (52.3% in all and 56.6% among
parents), still forming the largest frequency, followed by 5, 4 and 7 copies (39.3-35.6%, 6.2-
5.5% and 2.0-2.3%, respectively for all and among parents (**Table 4**). Among the parental
genomes, the portion of 6 PHR copies observed was slightly higher in samples carrying XX
over XY (57.9% vs. 55.2%). For 5 and 4 copies, however, the observed frequency was slightly
higher in XY carriers over XX (36.5% vs. 34.7% and 5.8% vs. 5.1%). Unlike the DJ, the
distribution of PHR copy numbers was not significantly different between parents and their
offsprings (Mann–Whitney U test, *p*>0.5. **Fig. 3B**). The copy number of the PHRs in the 8 DJ
samples also varied from 4 to 6 (**Table 2**). Distribution of the PHR copy number also showed
no significant relationship with the DJ gains or losses and showed a very weak correlation
when restricted to samples with 8 to 10 DJ copies (Pearson correlation, *r*=0.01, *p*=0.39),
indicating there could be other polymorphic variations of the PHRs that are yet to be
discovered (**Fig. 3C**).

Next, we examined the changes in rDNA copy number associated with DJ loss or gain. The
rDNA copy number varied widely among samples, ranging from 137 to 738 copies (**Fig. 3A**).
Unlike the DJ or PHR distribution, we observed a unimodal peak with a wide deviation
(mean=352, sd=70.2) as previously reported for the UK Biobank cohort[28]. Again, the rDNA
copy number distribution showed a very weak positive correlation for samples with 8 to 10 DJ
copies (DJ < 10.4; Pearson correlation: r = 0.09, p = 8.1 × 10□□). However, for samples with
10 or more copies, copy number ranges were similar and less correlated (DJ >= 10.4,
Pearson r = 0.12, p = 1.4 × 10□²). (**Fig. 3C**). No significant correlation was observed between
PHR copy number and rDNA copy numbers (p > 0.05) (**Fig. 3C**).

Cohort	Num. of PHRs	Num. of samples	%	XX	%	XY	%
RPC	<4	3	0.07%	0	0.00%	3	0.15%
	4	261	6.23%	126	5.93%	135	6.54%
	5	1,645	39.29%	828	39.00%	817	39.58%
	6	2,189	52.28%	1,127	53.09%	1,062	51.45%
	7	85	2.03%	39	1.84%	46	2.23%
	>7	4	0.10%	3	0.14%	1	0.05%
	Total	4,187	100.00%	2,123	100.00%	2,064	100.00%
RPC (Parents only)	<4	1	0.04%	0	0.00%	1	0.08%
	4	143	5.47%	67	5.14%	76	5.81%
	5	931	35.64%	453	34.74%	478	36.54%
	6	1,477	56.55%	755	57.90%	722	55.20%
	7	59	2.26%	28	2.15%	31	2.37%
	>7	1	0.04%	1	0.08%	0	0.00%
	Total	2,612	1	1,304	100.00%	1,308	100.00%
UK Biobank	<4	234	0.05%	128	0.05%	106	0.05%
	4	33,964	6.93%	18,195	6.84%	15,769	7.03%
	5	185,625	37.85%	100,601	37.82%	85,024	37.89%
	6	262,723	53.58%	142,803	53.69%	119,920	53.45%
	7	7,631	1.56%	4,164	1.57%	3,467	1.55%
	>7	197	0.04%	104	0.04%	93	0.04%
	Total	490,374	100.00%	265,995	100.00%	224,379	100.00%

We then evaluated whether our two alternative approaches—using GRCh38 as the reference or using marker k-mers—yielded high correlation with the CHM13 mapping–based approach. We selected samples representing each classified DJ copy number category, including all samples with 8 DJ copies and all samples with more than 12 DJ copies (n = 92 total). Among the GRCh38-based modes, we selected the "Fast-precise" mode (more details in **"Computing resources and prediction model"** section and **Methods**), which showed the

highest correlation with CHM13-based estimates and the lowest residual sum of squares
(RSS) when fitting DJ copy numbers derived from GRCh38 alignments (**Fig. S7 and Table**
S4). All three approaches showed similar distributions and strong correlations (Pearson
correlation r² and p-values shown in **Fig. 3D**), demonstrating the robustness and broad
applicability of these methods across different data types.

DJ and PHR copy number distribution in UK Biobank

To confirm the distribution of the DJ copy numbers and the PHRs in an independent cohort,
we applied our analysis to the UK Biobank (n=490,444), for which CRAM files aligned to
GRCh38 are available (Byrska-Bishop *et al.*, Methods[23]). To enable DJ and PHR copy number
estimation in this large cohort, we applied the mapping-based approach of our pipeline to
reduce computational demands ("pilot Fast mode" in **Methods and Fig. S8**). Among the
samples, we were able to calculate DJ and PHR counts for 490,369 and 490,374 samples,
respectively. The distribution pattern and percent samples assigned to each discrete DJ and
PHR count replicated what we observed in the RPC cohort, both when considering all
individuals and when restricted to unrelated individuals (**Fig. 4A**). In total, we identified 583
samples with 8 DJ copies, corresponding to a predicted ROB frequency of 1 in 841 (0.12%),
again closely matching the expected 1 in 800. Unlike the RPC cohort, we observed some
individuals with 7 DJ counts. More individuals with 3 PHR counts were observed, presumably
due to the larger size of the cohort, with a frequency of 0.05% regardless of the sex
chromosome composition, and the lowest PHR count observed 2, in a single sample. Again,
we did not observe a statistically significant difference in the frequency of each discrete copy
number bin between sex (χ^2 test of independence, *p*>0.5, **Table 3**). Unlike seen in the RPC
cohort, we did not observe an increased number of 8 DJ carriers in XX individuals. We also
confirmed that DJ counts were significantly (albeit slightly) higher not only in the parent group
compared to the offspring (**Fig. 4B**), but also in older groups compared to younger groups
(**Fig. S9**).

Fig. 4 | DJ and PHR counts in the UK Biobank.

(A) DJ and PHR count distribution among all and unrelated white British (WB) samples. (B) Density distribution

between the parent group and the offspring group. The RBC was calculated using the paired Wilcoxon Rank Sum

test. (C) DJ copies as a function for PHR copy numbers. Left, for all DJs. Right, inset of the samples classified as 8

or lower. (D) DJ (left) and PHR (right) counts in monozygotic twins. The twins share the same copy number

(Pearson's r test). (E) DJ counts in siblings (except monozygotic twins) with one of the siblings determined to have

8 DJs. No particular inheritance patterns were found. (F) Inheritance pattern of all 8 DJ carriers found with father-

offspring or mother-offspring relationships. No 8 DJ offspring with both parents were found in the cohort. (G)

Inheritance pattern of the DJ in offspring among the available trios, with at least one member of the trio having 9 or

fewer estimated DJs. No trio with 9 DJs in both parents was observed.

With the increased number of samples carrying 8 DJ copies, we were able to more precisely

evaluate the relationship between PHR and DJ copy number. The overall distribution mirrored

that observed in the RPC cohort, with a broader PHR copy number distribution among

individuals with 10 DJ copies, again presumably due to the larger sample size (**Fig. 4C**). A

higher proportion of individuals with 8 or fewer DJ copies had 5 PHR copies, as we initially

hypothesized (54.4%, compared with 37.8% in individuals with 9 or more DJs). However,

substantial individuals carried 3, 4, 6 or 7 copies of the PHRs (45.6%), suggesting other

structural variations may be present altering the PHR counts. For example, prior HPRC

assemblies showed 34.5% of the Chr. 14 haplotypes lacked the PHR (n=9 out of 26

haplotypes; de Lima *et al.*[14], Extended Data Fig. 9], supporting our observation of variable

PHR copy numbers in the population.

We further confirmed that monozygotic twins have identical DJ and PHR copy numbers (**Fig.**

4D), whereas dizygotic twins or siblings show differences in DJ copy number inheritance

pattern (**Fig. 4E**), confirming the precision of our method. Among samples carrying 8 DJs, 9

samples had either the maternal or paternal genome available. Among the 3 paternal samples

with 8 DJs, we found 1 case where the offspring inherited the same pattern (**Fig. 4F left**).

Among the 6 maternal samples, 1 offspring had inherited 8 DJs (**Fig. 4F right**). The copy

number of the offspring matched one of the parents in most cases, or were at 10 copies in

most cases when one of the parents had ≥11 copies (**Fig. 4G** and **Table S5**). Three samples

were found to have 9 DJs while both parental genomes had 10 DJs, suggesting a possible *de*

novo loss of the DJ in the offspring. No cases were observed in which both parents had 9 DJs

among the available trios.

Uncovering structural variation within the DJ of near-T2T genomes

For detailed inspection of structural gains or losses in DJ copy number, we analyzed 1KGP

samples both with and without near-T2T assemblies generated by the HPRC. The 1KGP

samples have recently been re-sequenced using Illumina short reads (30×, n=3,202 among
which n=2,504 are unrelated)[23]. A subset of the samples (n=226) have been long-read
sequenced by the HPRC to build a reference pangenome from *de novo* assembled
haplotypes. The cell lines were selected to broadly sample global genomic variation and were
restricted to lines with a low passage count and normal karyotype to avoid artifacts from cell
culture. These lines were assembled with Verkko[29] using long reads, Hi-C and parental data
when available[24].

We downloaded the Illumina BAM files aligned to GRCh38 from IGSR[30] and counted the DJ
copy numbers using the GRCh38 and *k*-mer based methods. To assess compatibility between
methods and across independent cohorts, we applied previously derived linear models for
each GRCh38 mode, using subsampled RPC data to transform DJ counts obtained from
GRCh38 alignment into the CHM13-based distribution. This model was subsequently used to
transform DJ counts in an independent 1KGP cohort. Among the evaluated approaches, the
"Fast-precise" mode demonstrated the lowest RSS (**Table S4**). Moreover, it was the only
method for which the transformed 1KGP distribution was statistically indistinguishable from
the original RPC distribution generated using the masked T2T-CHM13 method (**Fig. 5A and
Fig. S10**). Therefore, we chose to use the "fast-precise transformed" DJ counts for further
analyses (**Fig. 5A**). We obtained a similar distribution as found in the RPC or UK Biobank
samples, albeit with a slightly higher fraction of individuals binned in 8, 9 or 11+ DJ copies
(**Fig. 5A**, **Table S6 and Fig. S11**). Here, we also observed a pattern in which the parents have
a distribution shifted slightly higher than the offspring, consistent with our findings from the
RPC cohort and UK Biobank samples (**Fig. 5A and 5B**).

Seven individuals in the 1KGP dataset were classified as having 8 DJs, suggesting possible
ROB carriers (**Table S7**). All seven samples were identified within pedigreed families, allowing
us to confirm the inheritance pattern. Among the four families, one family showed a potential
ROB was inherited in the father-son relationship (HG01204-HG01206), and in the mother-
daughter relationship (HG00651-HG00652). Two potential ROB carriers were found only in
the parental genomes and not observed in their offspring. Interestingly, one offspring had 8.1-
8.2 copies, while neither parent carried 8 DJ copies, indicative of a possible *de novo* ROB.
Alternatively, given the parents are both classified as 9 copies, it is also possible that this child
inherited two DJ-less acrocentric chromosomes. Unfortunately, these samples were not
selected by the HPRC and so never underwent karyotyping. However, one sample, HG01204,

was initially selected for sequencing by the HPRC and underwent G-banding for quality

control. This revealed a karyotypic aberration involving a fusion between Chrs. 13 and 21 (**Fig.**

S12), which is an extremely rare type of a ROB chromosomal fusion with only a few case

reports available to date (for example, Chen *et al.*[31]). HPRC ultimately excluded this sample

due to its abnormal karyotype, but this example highlights the ability of our method to correctly

identify a true ROB from a large biobank.

Since no 8 DJ cases were sequenced by the HPRC, we evaluated all remaining cases of

abnormal DJ counts. Six samples with 9 predicted DJs and ten samples with 11 predicted DJs

were included in the HPRCr2 assemblies, allowing us to further inspect the sequence-level

structure of the DJs and the surrounding region. To confirm the entirety of the DJ structure, we

divided the DJ target into 3 regions—two palindromic arms residing close to the CER repeat

and the region flanking to the rDNA (DJarms and DJflank in **Fig. 5C**). We searched for the DJ,

rDNAs and other satellite repeats commonly found on the short arms of the acrocentric

chromosomes (**Fig. 5D and Table S8**). Next, we investigated the assembly graph to confirm

that the distal sequences were unambiguously assembled and connected to the rDNA nodes

(**Table S9**). HiFi or ONT reads were mapped to the junction to validate the loss of DJs.

Fig. 5 | Polymorphism of the rDNA DJs and surrounding sequence on the short arms of the acrocentric chromosomes.

(A) Distribution of DJ copy numbers in the RPC cohort, calculated using masked CHM13 as the ground truth, and in the 1KGP cohort, transformed from GRCh38 alignments (Fast-precise mode). (B) Distribution of DJ copy numbers across 1KGP samples (n = 3,202), with samples assembled by HPRC highlighted in orange. The normalized density plot (top) shows a slight shift of parental samples toward higher values among those with ≥10 copies. The right marginal plot displays the distribution of samples with assemblies. (C) Structure of the DJ and its functional components. The self-dotplot on top shows the palindromic arms inside the DJ, marked as DJ arm. The region flanking to the rDNA (DJ flank) shows a relatively unique pattern, with the beginning showing diverged

inverted repeat shared at the beginning of the DJ (next to the CER). (D) Distal sequences and repeats as observed
in CHM13. All DJ sequences are intact. Chr. 13 does not have the sequence below the bSat unlike the other
sequences. (E) Ideograms of cases with loss of DJs. HG00621 had an unplaced sequence containing the loss of
DJ. Sequences lost include the beginning of the rDNA array down to the bSat, resulting in loss of sequences
including the entire DJ, CER, ACRO and HSat3(A4). The rDNA array started at the large subunit (S28). HG01891
lost sequences including the HSat1A, with a rare rDNA inversion found at the beginning of the rDNA array.
HG01981 had only a partial loss of the DJ flank. Sequences larger than 1 Mbp are reduced and re-scaled for clarity.
(F) FISH with probes for the DJs and rDNAs on the 2 cases found to have 9 DJs. HG00621 exhibited the same DJ
loss pattern as the paternal genome, while HG01891 shared the pattern observed in the maternal genome. (G)
Ideograms of cases with gain of DJs. Six samples shared the same duplicated form of DJ, sharing the same
breakpoint at the DJ flank. The HSat1A block size varied, however the overall structure of the duplicated copy
(green box) remained identical. The 2 samples at the bottom showed a unique structure, with one having the DJ
palindromes duplicated and the inverted repeat part of the DJ flank. The last case shows a sequence within the
rDNA array, with one rDNA array (left) having the PJ intact. This time, the additional DJ has only one of the DJ
arms present with the DJ flank. (H) FISH for the two samples with 11 DJs. Because the DJs are too close to each
other, and highly packed inside the heterochromatin, the duplicated DJs aren't clearly distinguishable. Still, one
haplotype (red arrow) exhibits stronger intensity compared to the other. The other haplotype of the same
chromosome was assembled with a single DJ flanking the rDNA array.

Among the 6 samples with 9 DJ copies, 3 samples successfully assembled the junction with
the loss of DJ (**Fig. 5D** and **Fig. S13**). In the first case, HG00621, the DJ was lost entirely,
including the HSat3 (A4) distal to the DJ and the beginning of the rDNA array. The rDNA array
started at the 28S subunit rather than the regular 5' external transcribed spacer. FISH painting
of the DJs for this family showed the loss of DJ was also observed in the paternal genome,
HG00619 (**Fig. 5F**). In the second case, HG01891, we found the DJ was also lost entirely,
including part of HSat3 (**Fig. 5D**). The beginning of the rDNA array was found inverted with a
gap at the inversion breakpoint within the array. A local re-assembly of the inverted rDNA
junction was confirmed with ONT UL reads, suggesting this is a true inversion of the rDNA
(**Fig. S14**). The chromosome painting showed the absence of DJ was also present in the
maternal genome on Chr. 15 (**Fig. 5F**). The third case, HG01981, had DJ counts of 9.1-9.4,
indicating slightly higher counts for the samples grouped as 9 DJs (**Fig. 5G**). We found the DJ
was mostly intact except for the sequence flanking to the rDNA array on Chr. 21. Given that all
3 cases are showing non-overlapping breakpoints, we conclude that loss of 1 DJ is less
common, however still transmissible within families.

When investigating the 8 samples with extra DJ junction assembled, we found 7 samples
assembled the entire distal sequence to the telomere, including the duplicated DJ (**Fig. 5G**
and **Fig. S15**). Within those samples, we found the second DJ was partially duplicated with
the breakpoint residing internal to the flanking sequence, including distal sequences. Six
samples shared a similar segmentally duplicated block spanning sequences down to the large
beta satellite block. The DJ arm was truncated, with only the beginning 34 kb part being
duplicated. This truncated DJ flank sequence is shared downstream the DJ, as an inverted
repeat, with long-noncoding transcripts and pseudogenes. Chromosome painting of the
duplicated DJ in NA20752 (GM20752) and NA20805 (GM20805) was not visible, likely due to
their proximity at this scale (**Fig. 5H**). Arguably, one haplotype of Chr. 21 showed stronger
intensity for the DJ probes, suggestive for a possible duplication. The second case, found in
HG03654, had a block duplicated but starting with a smaller portion of the DJ flank down to
the ACRO repeat (**Fig. 5G**). The last case, HG00320, was partially assembled, surrounded by
rDNA sequences. In this sample, the partial DJ contained only one arm of the DJ palindromes,
with the flanking sequence intact to the rDNA. Given that we see 6 cases with overlapping
segmentally duplicated breakpoints ending at the same DJ flank sequence, we conclude this
form is the more common form present among the extra DJ carriers.

Computing resources and prediction model

Considering that current and past large-scale sequencing studies predominantly rely on
short reads, utilizing the GRCh38 reference genome in most cases, we developed our method
into a tool "DJCounter" to enable accurate detection of ROBs and determine the DJ copy
numbers, all while maintaining reasonable computational demands. Here we summarize the
various filtering and normalization strategies that were tested (**Table S4**).

Initially, we used the most conservative approach, normalizing DJ counts based on read
depth from alignments to masked T2T-CHM13 ("High-resolution" mode). However, this
approach proved too computationally intensive to apply to large cohorts such as the UK
Biobank dataset. To address this limitation, we tested an alternative "Fast" mode that utilizes
raw read counts directly from the BAM index file. While doing so, we refined the DJ target to
include the DJ flank sequences that were missing in the "pilot-Fast" mode. Although
computationally efficient, this method does not filter out secondary, supplementary or
duplicated read alignments, resulting in inconsistent results (batch effects) depending on how

the BAM file was generated. We next tested two additional modes that filter out the
aforementioned reads prior to estimating DJ counts. The first option ("Fast-precise" mode)
uses all filtered reads mapped to the autosomes and normalizes by the total size of the
autosomes. The second option ("Fast-refine" mode) calculates the median filtered read depth
across all autosomes and uses the selected autosome size for normalization.

433
Figure 6 | Computational resources required for DJ estimation using GRCh38 alignment–based and *k*-mer
approaches, and DJ classification using a GMM trained on 1KGP distribution.

(A-C) Computational resources required for DJ calculation across methods, including total CPU time in hour,
maximum memory usage (MaxMem, GB) and Elapsed time (wall-clock hours). (D) DJ copy number estimation
using different approaches, methods, and sequencing coverages. HG002 short read data were used and
subsampled to varying depths.(E) Gaussian mixture model (GMM) fitting on the training set subsampled from

1KGP (n = 2,402). (F) Zoomed-in view of the GMM fit. (G) DJ classification using the trained GMM on the
independent test subset of 1KGP (n = 800). (H) Transformed DJ copy numbers with corresponding classification
probabilities for each DJ category. (I) Recommendations for using DJCounter based on input format and sample
size.

We benchmarked the computational requirements of the three modes for GRCh38 and the
k-mer based method across different sequencing depths. We subsampled the Illumina
NovaSeq reads for HG002[32] to 5✕, 10✕, 20✕, 30✕, 40✕, and 50✕. As expected, the "fast" mode
was the most computationally efficient option, requiring substantially less CPU time and
maximum peak memory (**Fig. 6A-C and S16**). The "fast-precise" mode utilized CPU time
close to the "fast" mode, making it suitable for the next best option. The *k*-mer based method
was faster than the "fast-refine" mode, utilizing the given 120 GB of memory. The *k*-mer based
method operates on Meryl[33], which can trade shorter CPU wall time for higher memory. The
"fast-precise" and "fast-refine" methods show higher DJ count estimates (+0.7), however
provide robust results across different sequencing coverage over 5✕ (**Fig 6D**). The *k*-mer
based method reports closer to the expected 10 copies over 10✕. Nevertheless, these results
indicate that both mapping and *k*-mer based methods are applicable in clinical settings even
for shallow sequencing data generated.

A limitation of this approach is that it requires a large cohort to accurately estimate the
distribution and identify ROB samples based on that distribution. Therefore, we developed a
framework that operates at a single-sample level, enabling calculation of DJ copy number
directly from GRCh38 alignments and identifies ROB candidates using a reference density
derived from a large cohort. We constructed a Gaussian mixture model (GMM) to transform
DJ copy number estimates using subsamples from the 1KGP cohort (training set, n = 2,402)
and validated the model on the remaining samples (test set, n = 800). This framework
identifies candidate ROB samples and classifies each sample into a discrete copy number bin,
even when only a single sample is available. Due to the uneven density across peaks, we
fixed the mean of each component and constrained peak positions to align between the RPC
and 1KGP cohorts (**Fig S17)**. An additional intermediate component between 10 and 11
copies was included to reflect the partially duplicated DJ we observed (**Fig. 5F**) as a separate
category (**Fig. 6E-F**). Using this model, we could assign a discrete integer bin for each sample
in both the 1KGP test set (**Fig. 6G**) and the RPC cohort (**Fig. S18**), along with posterior
classification probabilities. The classification probabilities indicated high confidence (>0.5) for

all samples tested, with the majority of the test set showing strong support (>0.9), including
ROB samples (**Fig. 6H**).

For optimal performance, we recommend using DJCounter under different scenarios (**Fig.**
6I). For studies with large sample sizes (n > 1,000), where the cohort-specific distribution can
be reliably estimated and reads are aligned to GRCh38, we recommend running the "fast"
mode and inspecting the overall distribution to identify potential ROB carriers. However,
applying a strict cutoff of 8 copies may be challenging, as the fast mode includes duplicates
and supplementary reads, which can shift the overall distribution. In contrast, for smaller
cohorts (n < 1,000), where the distribution cannot be robustly inferred, the "fast-precise" mode
is preferred. In this setting, the classification approach should be applied to DJ estimates
generated using the fast-precise mode, with duplicate reads properly removed or marked in
the alignment files. The *k*-mer based method is reference-free and does not require the
alignment step. Therefore, it is recommended when rapid diagnosis is needed without
alignments.

DISCUSSION

Here we used recently completed T2T and near-T2T assemblies to understand the structure
and conservation of the short arms of human acrocentrics chromosomes, particularly
surrounding the rDNA arrays. Using this information, we developed a new method for
genotyping the rDNA distal junction and other features of the short arms to predict the
presence of Robertsonian chromosomes. Importantly, our method can be successfully applied
to short-read sequencing datasets, enabling its application to large-scale biobanks, as
demonstrated here using the RPC, UK Biobank, and 1KGP datasets.

Given the concordance between the observed frequency and the previously reported ROB
prevalence, as well as the observed inheritance patterns, we propose that a DJ copy number
of 8 or less is a reliable marker for screening ROB carriers. We also looked into the frequency
distribution of the PHR on the proximal short arms of Chrs. 13, 14, and 21, where one copy is
expected to be lost after ROB formation. However, a wider range of PHR copy numbers was
observed, indicating the polymorphic nature of this sequence, and PHR copy number was not
significantly associated with DJ copy number. Thus, in contrast to the DJ, the copy number of

the PHRs does not appear to be a reliable marker of ROBs. Prior work points to the SST1
macrosatellite and an enrichment of PRDM9 motifs within the ROB-associated PHR as
possible recombination hotspots, both explaining the higher prevalence of 13;14 and 14;21
ROBs, as well as the high degree of structural variation[14]. Cataloging structural variants and
their implications across the acrocentric short arms is an area of future work.

The DJ and the structure of the sequences distal to the DJ on the acrocentric short arms are
nearly impossible to genotype without near complete genome assemblies built from long
reads. The near-T2T HPRCr2 assemblies allowed us to understand the loss and gain of the
DJs, revealing previously uncharacterized structural variation in this region. The 9 DJ cases
found in the 1KGP set were sporadic, with the DJ-less junction sequence not found in the
RPC cohort. The gain of DJs was more frequently observed, with blocks of neighboring
sequences duplicated alongside. The recurrent duplication found in 6 out of 8 cases across
the 1KGP populations indicate that this is the most common form of duplication, regardless of
ancestral background.

Originally, we expected the copy number of the DJs to be found as 10 in most samples, with
only a handful of 8 DJs, as an indication for ROBs. However, our genotyping results from the
two large cohorts showed a greater variability in the copy number, especially in the 9 or 11
range. Using the learned frequency of the 9 DJs we observed, we estimate a rough false
positive rate for having two DJ-less haplotypes could be up to one third of the 8 DJ cases
(**Methods**). Alternatively, it is also possible to miss a true ROB carrier when the individual
carries the duplicated DJ in one of the non-ROB chromosomes, leading to a rough false
negative rate of 7.2%. Copy number analysis alone is limited to distinguish a true ROB carrier
from these alternative scenarios. Therefore, we suggest using our tool as a screening method
to identify potential ROB carriers, followed by karyotype confirmation. This approach will be
particularly useful for screening large cohorts to prioritize samples for confirmatory cytogenetic
testing.

Our method for DJ counting and ROB prediction could have immediate clinical application,
where whole-genome sequencing is commonly performed during routine carrier screening, but
where Robertsonian chromosomes are not currently diagnosed[34–36]. A positive result returned
by a sequencing-based method could be followed by a karyotype for confirmation, providing
more sensitive screening. Furthermore, ROB carriers are typically phenotypically normal at

the time of birth and most carriers go undiagnosed throughout their lives; however, aside from the fertility complications, little is known about any additional health implications. Also uncertain are the implications of DJ gain or loss, which have not been characterized prior to this study. The DJ includes important regulatory elements that have been shown with nucleolar function[37]. Thus, disruption of the DJ through either duplication or deletion may affect the transcription and function of the adjacent rDNA array. Future studies, aided by the methods and results presented here, are now needed to understand the clinical implications of structural variation on the short arms of the acrocentric chromosomes.

Limitations

To validate the four 8 DJ individuals we identified in the RPC cohort are ROB carriers, we attempted to contact them to request participation. Despite our efforts, all individuals were unable to be recontacted or declined to participate. Additionally, due to the extensive review processes required to re-contact ROB candidates within the UK Biobank, we instead genotyped the DJ copy numbers in the widely studied 1KGP lymphoblastoid cell lines. The structural variation in the DJ region may contain variants specific to the cell line as reported in the 1KGP dataset[23].

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources should be directed to and will be fulfilled by the lead contact, Arang Rhie rhiea@nih.gov and Adam M. Phillippy adam.phillippy@nih.gov.

Materials availability

This study did not generate new unique reagents or sequencing data.

Data and code availability

DJCounter is available on GitHub (https://github.com/marbl/DJCounter). The initial DJ counting method for the masked T2T-CHM13 version is available on GitHub (https://github.com/arangrhie/T2T-Ref). The acrocentric haplotypes presented in this study

560 from the Verkko assemblies are available on Zenodo
561 (https://doi.org/10.5281/zenodo.18895340) along with the acrocentric satellite annotation.

ACKNOWLEDGMENTS

A.R., J.K., S.J.S., S.K., D.A., A.M.P. were supported by NHGRI ZIA HG200398. J.K. was
supported by a grant of the Korea Health Technology R&D Project through the Korea Health
Industry Development Institute (KHIDI), funded by the Ministry of Health & Welfare, Republic
of Korea (grant number: RS-2022-KH131838). F.R-A and V.K.R. were supported by grants
from the Biotechnology and Biological Sciences Research Council (BB/R00675X/1). F.R-A
was additionally supported by a Barts Charity seed grant (G-002983). S.O.C.R was supported
by a grant from the Medical Research Council (MR/W007045/1). C.T. and C.M.W. were
supported by NHGRI ZIA HG200418-02. T.G.W. and S.S. were supported by NIH grant ZIC
HG200345. Members of the Reverse Phenotyping Core are as follows: Felicia Akinwande,
MBA, MSN, RN, Leslie G. Biesecker, MD, Jennifer Johnston, PhD, Justin Paschall, MS,
Sumeeta Singh, MS, Clesson Turner, MD, Caralynn Wilczewski, PhD, ScM, and Tyra
Wolfsberg, PhD. J.G. and T.P. are supported by the Stowers Institute for Medical Research
and by NCI R01CA266339. T.P. is supported by NCI R50CA305001. This research was
supported in part by the Intramural Research Program of the National Institutes of Health
(NIH). The contributions of the NIH author(s) are considered Works of the United States
Government. The findings and conclusions presented in this paper are those of the author(s)
and do not necessarily reflect the views of the NIH or the U.S. Department of Health and
Human Services. This work utilized the computational resources of the NIH HPC Biowulf
cluster (https://hpc.nih.gov). During the preparation of this work the authors used ChatGPT5.3
and Gemini to rephrase text for fluency. After using this tool, the authors reviewed and edited
the content as needed and take full responsibility for the content of the published article.

AUTHOR CONTRIBUTIONS

A.R. and J.K. conceived, developed DJCounter, conducted analysis for the RPC and 1KGP
cohorts, and drafted the manuscript. C.M.W., G.L.M., J.P., T.G.W., S.S. and C.T. conceived
analysis and administered the RPC dataset. F.R.-A., S.O.C.R and V.R. performed analysis on
the UK Biobank cohort. A.R., S.J.S. and A.M.P. defined the target regions. A.R., S.J.S. S.K.

and D.A. generated the HPRC Verkko assemblies and performed manual inspection to
validate the DJ variation. T.P. and J.G. conducted the FISH karyotyping. C.M.W. and G.L.M.
re-contacted participants for validation. A.M.P. supervised the project and provided critical
revisions. All authors reviewed and approved the final manuscript.

Declaration of Interests

595 The authors declare no competing interests.

REFERENCES

1. Hamerton, J.L., Canning, N., Ray, M., and Smith, S. (1975). A cytogenetic survey of 14,069
newborn infants. Clin. Genet. *8*, 223–243. https://doi.org/10.1111/j.1399-
0004.1975.tb01498.x.

2. Therman, E., Susman, B., and Denniston, C. (1989). The nonrandom participation of
human acrocentric chromosomes in Robertsonian translocations. Ann. Hum. Genet. *53*,
49–65. https://doi.org/10.1111/j.1469-1809.1989.tb01121.x.

3. Scriven, P.N., Flinter, F.A., Braude, P.R., and Ogilvie, C.M. (2001). Robertsonian
translocations--reproductive risks and indications for preimplantation genetic diagnosis.
Hum. Reprod. *16*, 2267–2273. https://doi.org/10.1093/humrep/16.11.2267.

4. Schoemaker, M.J., Jones, M.E., Higgins, C.D., Wright, A.F., and Swerdlow, A.J. (2019).
Mortality and Cancer Incidence in Carriers of Balanced Robertsonian Translocations: A
National Cohort Study. Am. J. Epidemiol. *188*, 500–508. https://doi.org/10.1093/aje/kwy266.

5. Kolgeci, S., Kolgeci, J., Azemi, M., Shala, R., Dakas, A., and Sopjani, M. (2013).
Reproductive risk of the silent carrier of Robertsonian translocation. Med. Arch. *67*, 56–59.
https://doi.org/10.5455/medarh.2013.67.56-59.

6. Dallaire, L., and Fraser, F.C. (1964). TWO UNUSUAL CASES OF FAMILIAL MONGOLISM.
Can. J. Genet. Cytol. J. Can. Genet. Cytol. *6*, 540–547. https://doi.org/10.1139/g64-069.

7. Hook, E.B. (1981). Unbalanced Robertsonian translocations associated with Down's
syndrome or Patau's syndrome: chromosome subtype, proportion inherited, mutation rates,
and sex ratio. Hum. Genet. *59*, 235–239. https://doi.org/10.1007/BF00283671.

8. Wilch, E.S., and Morton, C.C. (2018). Historical and Clinical Perspectives on Chromosomal
Translocations. Adv. Exp. Med. Biol. *1044*, 1–14. https://doi.org/10.1007/978-981-13-0593-
1_1.

9. Gravholt, C.H., Friedrich, U., Caprani, M., and Jørgensen, A.L. (1992). Breakpoints in
Robertsonian translocations are localized to satellite III DNA by fluorescence in situ
hybridization. Genomics *14*, 924–930. https://doi.org/10.1016/s0888-7543(05)80113-2.

10.   Nurk, S., Koren, S., Rhie, A., Rautiainen, M., Bzikadze, A.V., Mikheenko, A., Vollger,
624   M.R., Altemose, N., Uralsky, L., Gershman, A., et al. (2022). The complete sequence of a
625   human genome. Science *376*, 44–53. https://doi.org/10.1126/science.abj6987.

626   11.   Liao, W.-W., Asri, M., Ebler, J., Doerr, D., Haukness, M., Hickey, G., Lu, S., Lucas,
627   J.K., Monlong, J., Abel, H.J., et al. (2023). A draft human pangenome reference. Nature
628   *617*, 312–324. https://doi.org/10.1038/s41586-023-05896-x.

629   12.   Guarracino, A., Buonaiuto, S., de Lima, L.G., Potapova, T., Rhie, A., Koren, S.,
630   Rubinstein, B., Fischer, C., Gerton, J.L., Phillippy, A.M., et al. (2023). Recombination
631   between heterologous human acrocentric chromosomes. Nature *617*, 335–343.
632   https://doi.org/10.1038/s41586-023-05976-y.

633   13.   Gerton, J.L. (2024). A working model for the formation of Robertsonian chromosomes.
634   J. Cell Sci. *137*, jcs261912. https://doi.org/10.1242/jcs.261912.

635   14.   de Lima, L.G., Guarracino, A., Koren, S., Potapova, T., McKinney, S., Rhie, A., Solar,
636   S.J., Seidel, C., Fagen, B.L., Walenz, B.P., et al. (2025). The formation and propagation of
637   human Robertsonian chromosomes. Nature 647, 952–961. https://doi.org/10.1038/s41586-
638   025-09540-8.

639   15.   McStay, B. (2023). The p-Arms of Human Acrocentric Chromosomes Play by a
640   Different Set of Rules. Annu. Rev. Genomics Hum. Genet. *24*, 63–83.
641   https://doi.org/10.1146/annurev-genom-101122-081642.

642   16.   Sullivan, B.A., Jenkins, L.S., Karson, E.M., Leana-Cox, J., and Schwartz, S. (1996).
643   Evidence for structural heterogeneity from molecular cytogenetic analysis of dicentric
644   Robertsonian translocations. Am. J. Hum. Genet. *59*, 167–175.

645   17.   Álvarez-González, L., and Ruiz-Herrera, A. (2025). Leveraging Hi-C Data to Detect
646   Chromosomal Reorganizations. Methods Mol. Biol. *2968*, 213–230.
647   https://doi.org/10.1007/978-1-0716-4750-9_12.

648   18.   Huang, S., Juneau, K., Bogard, P.E., Davies, K.A., Wang, E.T., Kingsley, C.B., Struble,
649   C.A., Oliphant, A., Zahn, J.M., and Nicolaides, K.H. (2016). Identifying Robertsonian
650   Translocation Carriers by Microarray-Based DNA Analysis. Fetal Diagn. Ther. *40*, 59–62.
651   https://doi.org/10.1159/000441945.

652   19.   Hochstenbach, R., van Binsbergen, E., Schuring-Blom, H., Buijs, A., and Ploos van
653   Amstel, H.K. (2019). A survey of undetected, clinically relevant chromosome abnormalities
654   when replacing postnatal karyotyping by Whole Genome Sequencing. Eur. J. Med. Genet.
655   *62*, 103543. https://doi.org/10.1016/j.ejmg.2018.09.010.

656   20.   Bodian, D.L., Klein, E., Iyer, R.K., Wong, W.S.W., Kothiyal, P., Stauffer, D.,
657   Huddleston, K.C., Gaither, A.D., Remsburg, I., Khromykh, A., et al. (2016). Utility of whole-
658   genome sequencing for detection of newborn screening disorders in a population cohort of
659   1,696 neonates. Genet. Med. *18*, 221–230. https://doi.org/10.1038/gim.2015.111.

660   21.   Wilczewski, C.M., Obasohan, J., Paschall, J.E., Zhang, S., Singh, S., Maxwell, G.L.,
661   Similuk, M., Wolfsberg, T.G., Turner, C., Biesecker, L.G., et al. (2023). Genotype first:

Clinical genomics research through a reverse phenotyping approach. Am. J. Hum. Genet.
110, 3–12. https://doi.org/10.1016/j.ajhg.2022.12.004.

22.   UK Biobank Whole-Genome Sequencing Consortium (2025). Whole-genome
sequencing of 490,640 UK Biobank participants. Nature *645*, 692–701.
https://doi.org/10.1038/s41586-025-09272-9.

23.   Byrska-Bishop, M., Evani, U.S., Zhao, X., Basile, A.O., Abel, H.J., Regier, A.A.,
Corvelo, A., Clarke, W.E., Musunuri, R., Nagulapalli, K., et al. (2022). High-coverage whole-
genome sequencing of the expanded 1000 Genomes Project cohort including 602 trios.
Cell *185*, 3426-3440.e19. https://doi.org/10.1016/j.cell.2022.08.004.

24.   HPRC   Data   Release   2   Hum.   Pangenome   Ref.   Consort.
https://humanpangenome.org/hprc-data-release-2/.

25.   Mallick, S., Li, H., Lipson, M., Mathieson, I., Gymrek, M., Racimo, F., Zhao, M.,
Chennagiri, N., Nordenfelt, S., Tandon, A., et al. (2016). The Simons Genome Diversity
Project: 300 genomes from 142 diverse populations. Nature *538*, 201–206.
https://doi.org/10.1038/nature18964.

26.   Hansen, N.F., Dwarshuis, N., Ji, H.J., Rhie, A., Loucks, H., Logsdon, G.A., Vollger,
M.R., Storer, J.M., Kim, J., Adam, E., et al. (2025). A complete diploid human genome
benchmark for personalized genomics. BioRxiv Prepr. Serv. Biol., 2025.09.21.677443.
https://doi.org/10.1101/2025.09.21.677443.

27.   Pardo-Manuel de Villena, F., and Sapienza, C. (2001). Transmission ratio distortion in
offspring of heterozygous female carriers of Robertsonian translocations. Hum. Genet. *108*,
31–36. https://doi.org/10.1007/s004390000437.

28.   Rodriguez-Algarra, F., Evans, D.M., and Rakyan, V.K. (2024). Ribosomal DNA copy
number variation associates with hematological profiles and renal function in the UK
Biobank. Cell Genomics *4*, 100562. https://doi.org/10.1016/j.xgen.2024.100562.

29.   Antipov, D., Rautiainen, M., Nurk, S., Walenz, B.P., Solar, S.J., Phillippy, A.M., and
Koren, S. (2025). Verkko2 integrates proximity-ligation data with long-read De Bruijn graphs
for efficient telomere-to-telomere genome assembly, phasing, and scaffolding. Genome
Res. *35*, 1583–1594. https://doi.org/10.1101/gr.280383.124.

30.   30x GRCh38 | IGSR data collection https://www.internationalgenome.org/data-
portal/data-collection/1000genomes_30x.

31.   Chen, C.-P., Chern, S.-R., Wu, P.-C., Tsai, F.-J., Lee, C.-C., Town, D.-D., Chen, W.-L.,
Chen, L.-F., Lee, M.-S., Pan, C.-W., et al. (2009). Unbalanced and balanced acrocentric
rearrangements involving chromosomes other than chromosome 21 at amniocentesis.
Taiwan. J. Obstet. Gynecol. *48*, 389–399. https://doi.org/10.1016/S1028-4559(09)60329-6.

32.   Baid, G., Nattestad, M., Kolesnikov, A., Goel, S., Yang, H., Chang, P.-C., and Carroll,
A. (2020). An Extensive Sequence Dataset of Gold-Standard Samples for Benchmarking
and Development. Preprint at bioRxiv, https://doi.org/10.1101/2020.12.11.422022
https://doi.org/10.1101/2020.12.11.422022.

33.   Rhie, A., Walenz, B.P., Koren, S., and Phillippy, A.M. (2020). Merqury: reference-free
quality, completeness, and phasing assessment for genome assemblies. Genome Biol. *21*,
245. https://doi.org/10.1186/s13059-020-02134-9.

34.   Lu, W., Zhou, J., Rao, H., Yuan, H., Huang, S., Liu, Y., and Yang, B. (2024). A
Retrospective Analysis of Robertsonian Translocations from a Single Center in China.
Reprod. Sci. *31*, 851–856. https://doi.org/10.1007/s43032-023-01398-3.

35.   Beyer, C.E., and Willats, E. (2017). Natural selection between day 3 and day 5/6 PGD
embryos in couples with reciprocal or Robertsonian translocations. J. Assist. Reprod. Genet.
34, 1483–1492. https://doi.org/10.1007/s10815-017-1009-0.

36.   Zhang, S., Lei, C., Wu, J., Zhou, J., Sun, H., Fu, J., Sun, Y., Sun, X., Lu, D., and
Zhang, Y. (2017). The establishment and application of preimplantation genetic haplotyping
in embryo diagnosis for reciprocal and Robertsonian translocation carriers. BMC Med.
Genomics *10*, 60. https://doi.org/10.1186/s12920-017-0294-x.

37.   Vetter, P.H., Krauss, S., Steiner, O., Kropp, P., Möller, W.D., Moises, H.W., and Köller,
O. (1999). Vascular dementia versus dementia of Alzheimer's type: do they have
differential effects on caregivers' burden? J. Gerontol. B. Psychol. Sci. Soc. Sci. *54*, S93-98.
https://doi.org/10.1093/geronb/54b.2.s93.

Methods

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Oligonucleotides and Probes		
DJ OligoMix - 5' Biotin L	LC Sciences, LLC	Custom 100bp oligo paint corresponding to CHM13 chr13:5424523-5770548
CenSat D14Z1/D22Z1	Cytocell	LPE 014
Human whole chromosome paint (chr15)	Applied Spectral Imaging	FPRPR0167
45S rDNA probe	Empire genomics	RP11-450E20
Experimental models: Cell lines		
CHM13	Magee-Womens Hospital (Pittsburgh) IRB MWH-20-054	CHM13
HG002	Coriell Institute for Medical Research	GM24385
GM03417	Coriell Institute for Medical Research	GM03417
GM03786	Coriell Institute for Medical Research	GM03786
GM04890	Coriell Institute for Medical Research	GM04890
HG01204	Coriell Institute for Medical Research	HG01204
HG00619	Coriell Institute for Medical Research	HG00619
HG00620	Coriell Institute for Medical Research	HG00620
HG01981	Coriell Institute for Medical Research	HG01981
HG01890	Coriell Institute for Medical Research	HG01890
HG01889	Coriell Institute for Medical Research	HG01889
GM02752	Coriell Institute for Medical Research	GM02752
GM20805	Coriell Institute for Medical Research	GM20805
Software and algorithms		
Bedtools (v2.31.1)	Aaron R. Quinlan *et al.*	https://bedtools.readthedocs.io/en/latest/
BWA-MEM (v0.7.17)	Heng Li *et al.*	https://github.com/lh3/bwa
compleasm (v0.2.7)	Neng Huang *et al.*	https://github.com/huangnengCSU/compleasm
GenomeScope (v2.0)	Rhyker Ranallo-Benavidez *et al.*	https://github.com/schatzlab/genomescope
IGV (v2.18.2)	Robinson *et al.*	https://igv.org/

mashmap (v3.1.1)	Chirag Jain *et al.*	https://github.com/marbl/MashMap
merqury (v1.4.1)	Arang Rhie *et al.*	https://github.com/marbl/merqury
meryl (v1.4.1)	Arang Rhie *et al.*	https://github.com/marbl/merqury
Minimap2 (v2.30)	Heng Li	https://lh3.github.io/minimap2/
ModDotPlot (v0.9.0)	Sweeten *et al.*	https://github.com/marbl/ModDotPlot
paftools	Heng Li	https://lh3.github.io/minimap2/
Picard (v3.3.0)		https://github.com/lh3/bwa
Rustybam (v0.1.33)	Mitchell R Vollger *et al.*	https://www.vollgerlab.com/rustybam/
Samblaster (v0.1.26)	Gregory G. Faust *et al.*	https://github.com/GregoryFaust/samblaster
Samtools (v1.21)	Heng Li *et al.*	https://www.htslib.org/
scipy (v1.13.1)	Virtanen, P., *et al.*	https://scipy.org/
seqtk (v1.4)	Heng Li	https://github.com/lh3/seqtk
statsmodels (v0.14.5)	Seabold and Skipper *et al.*	https://www.statsmodels.org/stable/index.html
Verkko (v2.2.1)	Dmitry Antipov *et al.*	https://github.com/marbl/verkko
Yak (yak-0.1 (r56))	Heng Li	https://github.com/lh3/yak

METHOD DETAILS

Samples and data used

Control data

For CHM13, PCR-free Illumina Hi-Seq 2500 paired-end reads from Nurk *et al.*[1] were used. The reads are available on SRA: SRX1009644.

For HG002, PCR-free Illumina NovaSeq paired-end reads from Baid *et al.*[2] were used. The reads are available under google storage: https://console.cloud.google.com/storage/browser/brain-genomics-public/research/sequencing/fastq/novaseq/wgs_pcr_free/50x.

For the three previously identified ROB samples, PCR-free Illumina NovaSeq X paired-end reads from de Lima *et al.*[3] were used. The reads are available under dbGap: phs003920.v1.p1.

Tested cohorts

Reverse Phenotyping Core cohort

The original Reverse Phenotyping Core (RPC) cohort data included in the present analysis consisted of BAM files per sample, originally collected as part of previously published newborn trio genomic sequencing study[4] aligned to the hg19 reference genome. We therefore extracted all paired-end reads from the BAM files using samtools fastq and realigned them to either CHM13v2.0 or GRCh38.

```
samtools sort -@ $cpu -n -o ${name}_nameSort.bam ${inBam}
samtools fastq -@ $cpu ${name}_nameSort.bam \
    -1 $tmp/${name}_1.fq \
    -2 $tmp/${name}_2.fq
```

UK Biobank

To obtain DJ and PHR count estimates on the UK Biobank, GRCh38 alignment files in `cram` format stored in subfolders of the `/Bulk/GATK and GraphTyper WGS/Whole genome GATK CRAM files and indices [500k release]/` path were accessed through the UK Biobank Research Analysis Platform (UKB RAP, https://www.ukbiobank.ac.uk/use-our-data/research-analysis-platform). The Fast copy number estimation method described below was then executed via the UKB RAP application called swiss-army-knife (SAK) version 4.5.0, with local and remote scripts communicating through the `dxpy` python library.

1KGP

The 1000 Genomes Project dataset 30x Illumina NovaSeq 6000 paired-end reads were obtained from Byrska-Bishop *et al.*[5]. BAM files aligned to GRCh38 were downloaded and processed locally following instructions on IGSR for the 2504 unrelated and the 698 samples: https://www.internationalgenome.org/data-portal/data-collection/1000genomes_30x.

Target regions for DJ, PHR and rDNAs

Here we describe 3 features for copy number estimates of the distal junction (DJ), pseudo-homolog region (PHR) and the ribosomal DNA (rDNA).

Target regions in T2T-CHM13 and masked T2T-CHM13 reference

We used CHM13v2.0 as the reference genome, with the revised Cambridge Reference Sequence (rCRS) included for mitochondrial sequences. Centromeric Satellite (CenSat) repeat annotations from Altemose *et al*.[6] were used to define target regions of interest. For identifying internal repeat structure, we used ModDotPlot v0.9.0[7] to visualize sequence identity and its orientation.

The DJ region spans ~341 kbp upstream of the rDNA arrays in the short arms of the acrocentric chromosomes. We defined the DJ excluding the CER repeat as it was identified in the CenSat annotation from Nurk *et al*.[1] and Altemose *et al*. 2023[6]. After comparing the DJs on T2T-CHM13v2.0 to each other, we found minor variations on each unit, however the majority stayed concordant (**Fig. S1-2**). The DJs were all similar in terms of length (341,898-346,124 bps) with high identity (>95%), except for the bSat between the DJ arm and DJ flank (**Fig. 5C**) and the variation found within the second DJ arm as reported before[8].

For the PHR, we used the region as presented in Guarracino *et al*.[9] and refined the boundaries to exclude the inverted repeat. The PHR on Chr. 13 is the longest, spanning 1,174,544 bps, while Chr. 14 and Chr. 21 have a shorter form, spanning 1,124,403 and 1,126,601 bps, respectively. We chose the longest form on Chr. 13 as the target for the PHR. The PHR consists of two regions outside the highly repetitive satellite repeats, which we refer to here as PHR arm1 and PHR arm2 (**Fig. S3**).

For the rDNA, we compared the first rDNA units in each chromosomal array (**Fig. S4**). Among them, the unit on Chr. 13 had the longest form and was therefore chosen.

The final target regions for the DJ, PHR arm1 and the first rDNA unit are on Chr. 13. We masked the rest of the units present elsewhere in the reference using bedtools maskfasta (v 2.31.1). Although not used in this study, we additionally masked all 5S rDNA units except the last copy on Chr. 1.

To account for differences in sex chromosome complement, we applied the sex chromosome complement reference (SCC) approach to ensure proper read mapping on the sex chromosomes. Specifically, we used a pseudoautosomal region (PAR)-masked Y reference for XY individuals and a Y-masked reference for XX individuals. **Table S1** contains the exact coordinates of the target and masked region used on T2T-CHM13v2.0.

Target regions on GRCh38

To establish target coordinates on GRCh38, we extracted the corresponding target sequences from CHM13 and aligned the GRCh38 reference genome (https://rcs.bu.edu/examples/bioinformatics/gatk/ref/) to each target sequence using Minimap2 (v2.30)[10] with the -ax asm20 option. The resulting alignments were converted to PAF format using paftools.js (sam2paf). Regions shorter than 10 kbp were excluded from further analysis (**Tables S2-4**).

```
minimap2 -ax asm20 {target}.fasta Homo_sapiens_assembly38.fasta.gz
```

The PHR regions were defined as described above, resulting in a single location found on chr21:9722154-9796322 (+). Additional inspection was performed to refine the DJ region in GRCh38.

Initially, we evaluated the palindromic region of the DJ (ct_13_9) for counting the DJs. Before applying this approach to the larger cohort, we selected 50 samples from RPC and aligned the reads to GRCh38. Paired-end short reads were aligned to the GRCh38 reference using the same pipeline that we applied to RPC samples aligned to CHM13 (see "Read alignment on each reference" below). Next, we inspected the base-level sequencing depth across the DJ region by lifting over the alignments from GRCh38 to CHM13. This was done using the target and query coordinates in the PAF file, scaling the query's start and end positions to match the full DJ coordinates.

Per–base-pair depth was calculated using samtools depth. Following manual inspection, a subregion within the original DJ target in CHM13 (chr13:5,566,000–5,600,000) was excluded due to copy number variable alignment behavior in GRCh38. The refined DJ target coordinates were subsequently lifted back to GRCh38 to generate a final BED file for copy number estimation (**Table S2**). We found that using this target produced the expected distribution compared to the samples distribution on T2T-CHM13, with the majority of samples exhibiting approximately 10 DJ copies. This BED file was used to assess DJ copy number variation across UK Biobank samples.

However, we later found that extending the DJ target region to include upstream and downstream of the initial ct_13_9 sequences encompassing the full DJ and its flanking regions resulted in a stronger correlation with the CHM13-based mapping results. Regions on chromosomes 4 and 10 with low sequence identity (<83.4%) and low coverage (<2.8%) were

excluded from analysis. Therefore, this expanded region was selected as the final region of interest for DJ copy number estimation.

Read alignment on each reference

CHM13

Pilot testing on the control samples

We used Illumina short reads from the 3 identified ROB samples from de Lima *et al*[3], and mapped them to the masked reference. As a control, we used Illumina short reads from CHM13 and HG002, which were previously identified to have 10 DJs from FiSH experimental data (Nurk *et al* and Supplementary Figure XVIII.59 in Yoo *et al*.)[1,11].

RPC

We utilized the read coverage on chromosome X and Y in the original GRCh37 alignments to infer the sample's biological sex. First, we confirmed the BAM file is sorted based on genomic coordinates. Next, we used samtools idxstats to calculate the number of mapped reads to each sex chromosome and collected the size of the sex chromosomes. The coverage for chromosomes X and Y was determined by dividing the total number of mapped reads by the total chromosome length for each respective chromosome. To infer the sample's sex, we calculated the coverage ratio between chromosome X and chromosome Y (covX/covY). Samples with a ratio greater than 4 were classified as XX, while those with a ratio of 4 or less were classified as XY. Once the presence of Y was confirmed, the proper masked reference has been set to be used in the downstream analysis.

Short-read paired-end data from both the control samples and the RPC were aligned to the corresponding SCC reference using BWA (v0.7.17)[12] and post-processed with samtools (v1.21)[13] using fixmate, sort and markdup. Secondary alignments were removed with samtools view -F0x100 option.

```
bwa       mem       -t       $cpu       $ref       $r1       $r2
samtools     fixmate     -m     -@$cpu     $out.sam     $out.fix.bam
samtools  sort  -@$cpu  -O  bam  -o  $out.bam  -T  $out.tmp $tmp/$out.fix.bam
samtools markdup -r -@$cpu $out.bam $out.dedup.bam
```

GRCh38

To match with the method used in the UKbiobank and 1KGP, the paired-end reads from the six control samples and the 92 subsampled RPC samples were aligned to the reference genome using BWA-MEM (v0.7.17)[12]. The resulting SAM files were processed with SAMBLASTER (v0.1.26)[14] to add mate tags and sort reads by query name, after which duplicate reads were marked using Picard *MarkDuplicates* (v3.3.0).

Copy number estimate

Mapping based approaches

T2T-CHM13v2.0

Copy number estimates for target regions were calculated as the ratio of median sequencing depth in the target region to that of the autosomal background, excluding acrocentric short arms. Median depth for each target and background region was computed using samtools (v1.23)[13] and bedtools (v2.31.1)[15] with the following command:

```
samtools depth -@ $cpu -b $bed $bam | \
  awk '{print $1,$2-1,$2,$3}' OFS='\t' - | \
  bedtools map -a $bed -b - -c 4 -o median > $prefix.coverage_results.bed
```

For a given target T, the copy number (CN_T) was calculated as:

$$CN_T = 2 \times \frac{Median(Depth_T)}{Median(Depth_B)}$$

where $Depth_T$ represents the depth per base-pair within the target region and $Depth_B$ represents the depth across the autosomal region.

GRCh38

   Challenges in the normalization process arose from collapsed, misassembled, and fragmented target sequences in the acrocentric short arms, in particular the DJ regions in GRCh38, as well as from potential PCR-duplicate–like biases observed in reads mapped to

these regions. To reduce computational bottlenecks and minimize the impact on the target copy number estimation, we evaluated four different approaches for calculating the target and background sequencing depth.

Starting from the pilot-Fast mode, we developed four approaches and benchmarked the 92 RPC samples and compared correlations of the distribution to the result from T2T-CHM13 (**Fig. S7**).

For the pilot Fast mode applied to the UK Biobank cohort, we used samtools (v1.20)[13] to collect the number of reads mapped to the target and all regions. Specifically, we used `samtools view -c` aligned to the target (RC_T, **Table S2** for the DJ target) and `samtools idxstats` to collect reads aligned to the autosomes by using num. of mapped reads – num. of unmapped reads. In this mode, no reads are filtered prior to calculation. Unlike the RPC samples, we learned `samtools depth` was creating a bottleneck and was too costly to run on all 490k samples on DNAnexus (1 hour vs. 2 minutes using samtools idxstats). Albeit a little shifted, we found a good correlation to the T2T-CHM13 based result for the 92 subsamples (Pearson's correlation, r=9.81, p=57x10[-66]) between the two methods (**Fig. S8**), and decided to move forward using this approach. The script was deployed on DNAnexus and collected copy counts of the DJ, rDNA, and PHR.

Copy number for each target was estimated using read count normalization. Read counts (RC) mapped to each target were normalized to autosomal coverage, which was calculated by subtracting unmapped read counts (RU_A) from the total reads mapped to autosomes (RC_A) and dividing by the total autosomal length (Len_A). The normalized target RC was then used to infer copy number.

$$CN_{GRCh38-pilot\ Fast} = 2 \times \frac{SUM(RC_T)/Len_T}{SUM(RC_A - RU_A)/Len_A}$$

After a series of evaluations, we updated our method to include the DJ flanking region in the target as it yielded better copy number estimates. The following four modes (**Table S5**) use the target DJ region as shown in **Table S3**.

The Fast mode estimates copy number using the number of reads aligned to the target region (RC_T) calculated by `samtools view -c`, normalized by the original target length in CHM13v2.0 (Len_T). Autosomal coverage is calculated by using the median of autosomal read

coverage, defined as the total number of reads mapped to each autosome (RC_C) divided by the length of the autosomal (Len_C) reported by using `samtools idxstat`. In this mode, no reads are filtered prior to calculation.

$$CN_{GRCh38-Fast} = 2 \times \frac{SUM(RC_T)/Len_T}{Median(RC_C/Len_C)}$$

In the Fast-precise mode, reads marked as duplicates, as well as secondary and supplementary alignments, were excluded from analysis. Target coverage was calculated as the total filtered read counts aligned to the target region (FRC_T) (as reported by `samtools view -c -F 3332`) divided by the target length (Len_T). Background coverage was calculated as the total number of filtered reads aligned to the autosomes (FRC_A) divided by the total autosomal length (Len_A).

$$CN_{GRCh38-Fast-precise} = 2 \times \frac{SUM(FRC_T)/Len_T}{FRC_A/Len_A}$$

The Fast-refine mode uses the same approach to calculate target coverage. However, background coverage is defined as the median coverage across autosomal regions rather than coverage calculated from the entire autosome.

$$CN_{GRCh38-Fast-refine} = 2 \times \frac{SUM(FRC_T)/Len_T}{Median(FRC_C/Len_C)}$$

The High-resolution mode calculates coverage at per-base resolution, consistent with the approach used for CHM13v2.0. The depth for each target region was computed using `samtools depth` ($Depth_T$) and normalized by the target length (Len_T). For background coverage, we used `samtools coverage` to obtain coverage metrics for each autosome (Cov_C) and defined the background as the median autosomal coverage.

$$CN_{GRCh38-High-resolution} = 2 \times \frac{SUM(Depth_T)/Len_T}{Median(Cov_C)}$$

Reference-free k-mer based approach

For identifying the DJ k-mers, 31-mers were collected with Meryl (v1.4.1)[16] for each of the five DJ copies in T2T-CHM13v2.0. The DJ sequence was extracted with `samtools faidx`, and `meryl count k=31` was used to build the `dj.$chr.meryl` for each chromosome. Only *k-*

mers present once in each chromosome were selected and intersected, to ensure we use *k*-mers present once in each DJ and in all 5 DJs. This yielded 178,731 potential target *k*-mers in `dj.meryl`.

```
meryl intersect [ equal-to 1 dj.chr13.meryl ] [ equal-to 1 dj.chr14.meryl ]
output dj.chr13_chr14.1.meryl
meryl intersect [ equal-to 1 dj.chr15.meryl ] [ equal-to 1 dj.chr21.meryl ]
output dj.chr15_chr21.1.meryl
meryl intersect [ intersect dj.chr13_chr14.1.meryl dj.chr15_chr21.1.meryl ]
[ equal-to 1 dj.chr22.meryl ] output dj.meryl
```

For identifying the DJ target k-mers, the DJ *k*-mers were filtered one more time after projecting the k-mer multiplicity as found in the short reads. The 31-mer meryl DB was built with `meryl count k=31`, and the histogram was generated with `meryl histogram`. The 2 copy peak was inferred using GenomeScope 2.0[17] with the following command line:

```
Rscript $tools/genomescope2.0/genomescope.R -i chm13.k31.hist -o
chm13.k31.genomescope2 --fitted_hist --testing -k 31
```

The predicted haploid coverage (`kcov`) resulted in 49.9. The inspected 10 copy peak in CHM13 is therefore ~50 x 10; resulting in ~500x. This matched the peak distribution of the DJ *k*-mers when projected to the read spectrum (**Fig. 1E**). The final selected DJ target *k*-mers were chosen falling between the range of 475-525x (±0.5 copy), to allow room for small variations or sequencing biases. This resulted in 52,227 distinct DJ target *k*-mers in `dj.target.meryl`.

```
meryl   greater-than   475   [   less-than   525   dj.chm13_count.meryl   ]\
  output dj.target.meryl
```

For copy number estimate the DJs in a given sample, we first collect 31-mers from the samples' short reads. The 31-mer database is then queried to collect the k-mer multiplicity histogram of the DJ target *k*-mers. The median of the DJ target k-mers found in the reads is used as the *k*-mer frequency of the DJs, and the "2-copy peak" is inferred from the histogram of all 31-mers in the reads. The 2-copy peak is obtained using a script in Merqury (`kmerHistToPloidyDepth.jar`), which differentiates the slope of the *k*-mer histogram and reports the estimated ploidy and its peak and boundary by collecting the local maxima[16].

```
# Collect 31-mers from reads
meryl count k=31 output $sample.k31.meryl $sample.fq.gz
# Intersect DJ target k-mers
meryl intersect threads=24 $sample.k31.meryl dj.target.meryl output
$sample.k31.DJ.meryl
# Histogram of the DJ kmers found in $sample
meryl histogram $sample.k31.DJ.meryl > ${sample}_DJ.hist
# Median
count=`cat ${sample}_DJ.hist | awk '{if (NR>1) { count+=$NF;} } END {print
count/2}'`
med=`cat ${sample}_DJ.hist | \
  awk -v cnt=$count '{cnt_sum+=$NF; if (NR>1 && cnt_sum > cnt) {print $(NF-
1); exit;} }'`
# 2-copy peak
java -jar -Xmx256m $MERQURY/eval/kmerHistToPloidyDepth.jar $sample.k31.hist
| tail -n1 | awk '{print $2}'
```

$$CN_{k-mer} = 2 \times \frac{Median(DJ\ target\ k-mer\ counts)}{2-copy\ peak}$$

Chromosome spreads and Fluorescent In-Situ Hybridization (FISH) for rDNA and DJ

Human lymphoblastoid cell lines (LCL) and fibroblast cell lines were according to Coriell protocols. CHM13 immortalized using human telomerase reverse transcriptase (hTERT) were propagated in DMEM:F12 with L-glutamine supplemented with 1x non-essential amino acids (Gibco), 1x Insulin-Transferrin-Selenium (Gibco), 1mM sodium pyruvate (Gibco), and 10% FBS. For preparation of chromosome spreads, cultures were blocked in mitosis by the addition of Karyomax colcemid solution (0.1 µg/ml, Thermo Fisher) for 6-7h. Adherent fibroblast cells were collected by trypsinization. Cells were incubated in hypotonic 0.4% KCl solution for 12 min and prefixed by addition of methanol:acetic acid (3:1) fixative solution (1% total volume). Pre-fixed cells were spun down and then fixed in Methanol:Acetic acid (3:1). Chromosome spreads were dropped on a glass slide and incubated at 65°C overnight. Before hybridization, slides were treated with 0.1mg/ml RNAse A (Qiagen) in 2xSSC for 45 minutes at 37°C and dehydrated in a 70%, 80%, and 100% ethanol series for 2 minutes each. Slides

were denatured in 70% deionized formamide/2X SSC solution pre-heated to 72°C for 1.2-1.5 min. Denaturation was stopped by immersing slides in 70%, 80%, and 100% ethanol series chilled to -20°C. Probes used in combinations were denatured in a hybridization buffer (Empire genomics) by heating to 80°C for 7 minutes before applying to denatured slides. Specimens were hybridized to the probes under a glass coverslip or HybriSlip hybridization cover (GRACE Biolabs) sealed with the rubber cement or Cytobond (SciGene) in a humidified chamber at 37°C for 48-72hours. After hybridization, slides were washed in 50% formamide/2X SSC 3 times for 5 minutes per wash at 45°C, then in 1x SSC solution at 45°C for 5 minutes twice and at room temperature once. For biotin detection, slides were incubated with streptavidin conjugated to Cy5 (Thermo) for 2-3 hours in PBS containing 0.1% Triton X-100 and 5% bovine serum albumin (BSA), and then washed 3 times for 5 minutes with PBS/0.1% Triton X-100. Slides were rinsed in water, air-dried in the dark, and mounted in Vectashield containing DAPI (Vector Laboratories). Z-stack confocal images were acquired on the Nikon TiE microscope equipped with 100x objective NA 1.45, Yokogawa CSU-W1 spinning disk, and Flash 4.0 sCMOS camera (Hamamatsu). Image processing was performed in FIJI.

Pattern of DJs in the HPRC samples

HPRC Verkko assemblies

In parallel to the HPRCr2 assemblies, a Verkko v2.2.1[18] assembly of each genome was made using automated scripts to download data from AWS, build parental k-mer databases, and run Verkko with both Hi-C and Trio data in parallel. Standardized post-processing and QC was performed with the same pipeline which used Yak (https://github.com/lh3/yak) to measure QV (k=31) and switch error rate, compleasm[19] for gene completeness, and computed T2T contig and scaffold statistics using seqtk to identify the telomeric and gap sequences in the assemblies. For each sample where a trio and Hi-C was available, the assembly with the most T2T contigs was selected. In total, there were 59 trios and 159 Hi-C assemblies.

Chromosome assignment and orientation

Mashmap v3.1.1[20] was used to map the assembly to CHM13v2.0. The contigs were assigned to chromosomes if they had mappings over 99% identity covering more than 10% of the reference chromosome in a given orientation (forward or reverse-complement). If a contig

matched over 10% of more than one chromosome, multiple assignments were noted and the longest one was used. The XY pseudo-autosomal regions (PAR) and distal regions of the acrocentric were excluded from the covered fraction count as they recombine between individuals and cannot be accurately assigned via reference alignment. Therefore, these regions could only be named as a chromosome if they were assembled together with a unique region of the chromosome. Across all samples and assembly types (verkko-hi-c or verkko-trio+hi-c), there were 23,094 contigs totaling 2.11 Tbp with an average size of 91.58 Mbp which could be oriented and assigned. There were 338 contigs with ambiguous assignments with an average size of 5.7 Mbp and a total of 1.936 Gbp while 75,565 contigs could not be assigned with an average size of 0.12 Mbp totaling 8.73 Gbp.

In parallel, the homopolymer-compressed Verkko gfa were aligned using Mashmap (v3.1.1)[20] to a compressed version of CHM13v2.0 to assign the individual nodes to a chromosome. A two pass assignment was performed. First, nodes with mappings over 99% identity and 5 Mbp matches to a chromosome were assigned to that chromosome. Second, if a connected component had no nodes assigned by the first criteria, nodes over 500 kbp were assigned. This identified connected components in the assembly graph belonging to each chromosome.

Repeat annotation of the acrocentric chromosomes

The acrocentric chromosomal short arms carry human satellite repeats (HSat) at and around the centromere, some specific to the acrocentric region forming a particular pattern[1,21]. To obtain an overview of the sequence content and confirm sequences belonging to the acrocentric short arms, particularly the distal and proximal sequences from the rDNA, satellite sequences in the assemblies were annotated based on a set of targeted sequence (Table S8).

In brief, the target repeat sequences were selected based on the CenSatv2.1 annotation available for T2T-CHM13v2.0 (https://s3-us-west-2.amazonaws.com/human-pangenomics/T2T/CHM13/assemblies/annotation/chm13v2.0_censat_v2.1.bed). Most were selected from chr13 or chrY, except for HSat1A, HSat3_A5, and HSat3_B3. For these three HSat repeats, the original consensus generated for classifying the satellite array subgroups were used as described by Altemose *et al.*[22]. The consensus sequences are available as fasta files on https://github.com/altemose/HSatReview/tree/main. The rDNA reference KY962518 was rotated to begin upstream of the 45S TSS, to include the 45S promoter region.

The target sequences were found using minimap2 (v2.28) in each assembly. Because most mappers are designed to find one best match for each query in the reference, using the assembly as the reference would yield only one result for each mapped target satellite. Thus, the reference and query sequences were flipped, mapping the assembly (query) to the targeted repeat sequence (reference) instead. This approach allows a more sensitive search of the target sequence in the assembly. Once all the alignments are collected in a PAF file, the reference and query coordinates were inverted with rustybam (v0.1.33)[23] and converted to BED format. Alignment blocks within 500 bp were merged with BEDTools (v2.31.1)[15] and formatted with designated colors and filtered to be over 2 kbp or 7–8 kbp to exclude excessive alignment matches to LINE elements.

Below are the command lines used:

```
minimap2 -x asm20 --eqx --MD -t $cpus -c $target.fa $asm_fa >>
${sample}_to_sat.paf
rustybam invert ${sample}_to_sat.paf > sat_to_${sample}.paf
cat sat_to_${sample}.paf |\
  awk -v OFS='\t' '{if ($10<$11) {idy=100*$10/$11;} \
    else {idy=100*$11/$10;} print $6,$8,$9,$1,idy,$5}' |\
  sort -k1,1V -k2,2n - > sat_to_${sample}.bed
```

For each target satellite, merging and filtering was applied along with the color assignment. Telomere sequences and gaps were annotated with Seqtk (v1.4, https://github.com/lh3/seqtk) using tel -d5000 and gap -l 1 parameters.

```
awk -v chr_hap=$chr_hap -v sat=$sat '$1==chr_hap && $4==sat'
${out}/sat_to_${sample}.bed | \
 bedtools merge -s -d 500 -c 4,5,6 -o distinct,median,distinct -i - |\
 awk -v len=$len '$3-$2>len' |\
 awk -v chr=$chr -v col=$col -v OFS="\t" '{print chr, $2, $3, $4, $5, $6, $2,
$3, col}' >> $OUT
```

The final BED file was then concatenated and sorted for manual inspection and visualization on IGV (v2.18.2)[24].

Benchmarking and optimizing DJCounter for GRCh38

Subsampling and benchmarking computing resources

 HG002 reads aligned to GRCh38 were subsampled to the desired sequencing depths (1×, 5×, 10×, 20×, 30×, 40×, and 50×) using samtools (v1.21)[13] to evaluate the stability and computational requirements of each method. We ran DJCounter on the subsampled BAM files using the GRCh38-based modes and additionally extracted reads from the BAM files for analysis with the k-mer–based method.

 To benchmark computational requirements of each DJ estimation mode, we extracted runtime and resource utilization metrics from SLURM job accounting records using sacct. For each job, we summarized: (i) Elapsed time as the maximum wall-clock duration across steps (hours); (ii) Total CPU time as the sum of CPU time across steps (hours); (iii) maxMem as the maximum resident memory usage (MaxRSS, GB); (iv) maxVmem as the maximum virtual memory usage (MaxVMSize, GB); (v) %CPU as the average CPU utilization relative to a single fully utilized core (100%); and (vi) AvgCPU as the mean CPU time per step (hours).

Linear transformation between GRCh38- and CHM13-derived DJ copy numbers

To determine which GRCh38 mapping mode exhibited the least cohort bias and to harmonize DJ copy number estimates derived from GRCh38 alignments with those obtained from CHM13, we fitted an ordinary least squares (OLS) linear regression model using the *statsmodels* package (v0.14.5)[25]. Specifically, DJ copy numbers calculated from CHM13 alignments of 92 RPC subsamples were modeled as a function of DJ copy numbers derived from GRCh38 alignments of the matched samples:

$$DJ_{CHM13} = \beta_0 + \beta_1 \times DJ_{GRCh38} + \epsilon$$

where β_0 represents the intercept, β_1 the slope, and ϵ the residual error term.

We fitted this model separately using DJ copy numbers generated from different GRCh38 modes (Fast, Fast-precise, Fast-refine, and High-resolution) and selected the Fast-precise mode because it yielded the lowest residual sum of squares (RSS) among all modes and

showed no significant difference between the CHM13-aligned RPC DJ distribution and the transformed 1KGP DJ distribution.

To harmonize DJ copy numbers across cohorts, we applied the fitted Fast-precise regression model to transform GRCh38-derived DJ estimates to the CHM13 scale using the `predict` function in *statsmodels*. The transformed values were subsequently used for GMM fitting and classification analyses

GMM fitting and classification

We used the transformed DJ copy numbers to split the 1KGP samples into a training set and a test set at a ratio of 0.75 and 0.25, respectively. When applying a standard GMM without fixed means, the model failed to identify peaks in sparsely populated regions and exhibited over-dispersion or component collapse, particularly for rare categories such as ROB samples with 8 copies. To address this issue, we performed GMM fitting with fixed component means and constrained standard deviations ($0.1 \leq \sigma \leq 0.2$).

Candidate peak locations were identified from the DJ copy number distributions of the RPC cohort aligned to CHM13 and the transformed 1KGP (GRCh38-aligned) cohort using the `find_peak` function from the *scipy.signal* module (v1.13.1)[26]. Peaks that were consistent across cohorts were selected as $7(k)$ fixed component means (u): [8, 9.1, 10.1, 10.7, 11.1, 11.9, 13].

To quantify uncertainty in the estimated mixture parameters, we performed nonparametric bootstrap resampling on the training set. Using the DJ copy numbers from the training data, we generated B=1,000 bootstrap datasets by sampling observations with replacement. For each bootstrap replicate b, we fit a one-dimensional GMM with fixed means and constrained standard deviations using an expectation–maximization (EM) algorithm. For each replicate, we estimated the mixture weights (ψ_{kb}) and standard deviations (σ_{kb}) for each k. Model fitting was performed with a maximum of 500 iterations, and convergence was defined as a change in log-likelihood of less than 1×10e-10 times between successive iterations. Final parameter (ψ_k and σ_k) estimates were obtained by averaging the bootstrap estimates of weights and standard deviations.

For classification, we used the fitted GMM parameters to assign samples from the test set to discrete DJ copy number categories. For each sample, the unnormalized posterior density for each component was computed as $\psi_k \cdot norm.pdf(x_i, u_k, \sigma_k)$ and normalized across components to obtain posterior membership probabilities. Each sample was assigned to the category with the highest posterior probability (maximum a posteriori classification).

False positive and false negative rate

False positive rate. We assume ROB carriers have 8 DJ copies based on observations from prior studies. However, as shown in our study, we found 1 DJ-less haplotypes (9 DJs) across different cohorts at a frequency of 3.1%. Assuming the 10 acrocentric chromosomes are independent, this gives us a rough estimate that each acrocentric chromosome can lack a DJ with probability of $1 - (1 - 0.031)^{1/10} \approx 0.31\%$. As such, the probability of a deletion of DJs is 0.0031, the probability that exactly 2 of them have a deletion is given by

$$P(FP\ ROB) = (0.0031)^2 \times (1 - 0.0031)^8 \times C(10,2)$$

This corresponds to approximately 0.04% of all samples. This could consist up to ⅓ of detected 0.12% 8 DJ cases.

False negative rate. Given the high frequency we observed for one or more DJ gains (~8.9%), it is also possible that a ROB carrier can be misclassified as having 9 copies when one of the non-fused acrocentric haplotype has the duplicated DJ present. Similarly as above, we can roughly estimate the probability for having a duplicated DJ in each acrocentric chromosome as $1 - (1 - 0.089)^{1/10} \approx 0.0093$. Thus, the chance that a ROB carrier will have duplicated DJ in at least one of the other acrocentric chromosomes and thus more than 8 DJ can be estimated as

P(FN ROB | ROB) = $1 - (1 - 0.0093)^8 \approx 0.07$

As a result, we expect that roughly 7% of ROBs can be missed by using 8 DJs as the sole marker for identifying ROB carriers.

Copy number analysis alone cannot distinguish between a true ROB carrier and this alternative scenario. To determine whether samples with 8 copies carry a genuine fusion chromosome, additional analysis such as karyotyping or Hi-C are required. Given that we did not observe any offspring with both parents having 9 DJs in the UK Biobank dataset, we estimate the chances might be even lower. We believe a more dedicated study on familial participants with identified 8 DJ offspring and 9 DJ parents may answer this question.

Methods References

1. Nurk, S., Koren, S., Rhie, A., Rautiainen, M., Bzikadze, A.V., Mikheenko, A., Vollger, M.R., Altemose, N., Uralsky, L., Gershman, A., et al. (2022). The complete sequence of a human genome. Science *376*, 44–53. https://doi.org/10.1126/science.abj6987.

2. Baid, G., Nattestad, M., Kolesnikov, A., Goel, S., Yang, H., Chang, P.-C., and Carroll, A. (2020). An Extensive Sequence Dataset of Gold-Standard Samples for Benchmarking and Development. Preprint at bioRxiv, https://doi.org/10.1101/2020.12.11.422022 https://doi.org/10.1101/2020.12.11.422022.

3. de Lima, L.G., Guarracino, A., Koren, S., Potapova, T., McKinney, S., Rhie, A., Solar, S.J., Seidel, C., Fagen, B.L., Walenz, B.P., et al. (2025). The formation and propagation of human Robertsonian chromosomes. Nature *647*, 952–961. https://doi.org/10.1038/s41586-025-09540-8.

4. Bodian, D.L., Klein, E., Iyer, R.K., Wong, W.S.W., Kothiyal, P., Stauffer, D., Huddleston, K.C., Gaither, A.D., Remsburg, I., Khromykh, A., et al. (2016). Utility of whole-genome sequencing for detection of newborn screening disorders in a population cohort of 1,696 neonates. Genet. Med. *18*, 221–230. https://doi.org/10.1038/gim.2015.111.

5. Byrska-Bishop, M., Evani, U.S., Zhao, X., Basile, A.O., Abel, H.J., Regier, A.A., Corvelo, A., Clarke, W.E., Musunuri, R., Nagulapalli, K., et al. (2022). High-coverage whole-genome sequencing of the expanded 1000 Genomes Project cohort including 602 trios. Cell *185*, 3426-3440.e19. https://doi.org/10.1016/j.cell.2022.08.004.

6. Altemose, N., Logsdon, G.A., Bzikadze, A.V., Sidhwani, P., Langley, S.A., Caldas, G.V., Hoyt, S.J., Uralsky, L., Ryabov, F.D., Shew, C.J., et al. (2022). Complete genomic and epigenetic maps of human centromeres. Science *376*, eabl4178. https://doi.org/10.1126/science.abl4178.

7. Sweeten, A.P., Schatz, M.C., and Phillippy, A.M. (2024). ModDotPlot-rapid and interactive visualization of tandem repeats. Bioinformatics *40*, btae493. https://doi.org/10.1093/bioinformatics/btae493.

8. van Sluis, M., Gailín, M.Ó., McCarter, J.G.W., Mangan, H., Grob, A., and McStay, B. (2019). Human NORs, comprising rDNA arrays and functionally conserved distal elements, are located within dynamic chromosomal regions. Genes Dev. *33*, 1688–1701. https://doi.org/10.1101/gad.331892.119.

9. Guarracino, A., Buonaiuto, S., de Lima, L.G., Potapova, T., Rhie, A., Koren, S., Rubinstein, B., Fischer, C., Gerton, J.L., Phillippy, A.M., et al. (2023). Recombination between heterologous human acrocentric chromosomes. Nature *617*, 335–343. https://doi.org/10.1038/s41586-023-05976-y.

10. Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics *34*, 3094–3100. https://doi.org/10.1093/bioinformatics/bty191.

11. Yoo, D., Rhie, A., Hebbar, P., Antonacci, F., Logsdon, G.A., Solar, S.J., Antipov, D., Pickett, B.D., Safonova, Y., Montinaro, F., et al. (2025). Complete sequencing of ape genomes. Nature *641*, 401–418. https://doi.org/10.1038/s41586-025-08816-3.

12. Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics *25*, 1754–1760. https://doi.org/10.1093/bioinformatics/btp324.

13. Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079. https://doi.org/10.1093/bioinformatics/btp352.

14. Faust, G.G., and Hall, I.M. (2014). SAMBLASTER: fast duplicate marking and structural variant read extraction. Bioinformatics *30*, 2503–2505. https://doi.org/10.1093/bioinformatics/btu314.

15. Quinlan, A.R., and Hall, I.M. (2010). BEDTools: a flexible suite of utilities for comparing genomic features. Bioinformatics *26*, 841–842. https://doi.org/10.1093/bioinformatics/btq033.

16. Rhie, A., Walenz, B.P., Koren, S., and Phillippy, A.M. (2020). Merqury: reference-free quality, completeness, and phasing assessment for genome assemblies. Genome Biol. *21*, 245. https://doi.org/10.1186/s13059-020-02134-9.

17. Ranallo-Benavidez, T.R., Jaron, K.S., and Schatz, M.C. (2020). GenomeScope 2.0 and Smudgeplot for reference-free profiling of polyploid genomes. Nat. Commun. *11*, 1432. https://doi.org/10.1038/s41467-020-14998-3.

18. Antipov, D., Rautiainen, M., Nurk, S., Walenz, B.P., Solar, S.J., Phillippy, A.M., and Koren, S. (2025). Verkko2 integrates proximity-ligation data with long-read De Bruijn graphs for efficient telomere-to-telomere genome assembly, phasing, and scaffolding. Genome Res. *35*, 1583–1594. https://doi.org/10.1101/gr.280383.124.

19. Huang, N., and Li, H. (2023). compleasm: a faster and more accurate reimplementation of BUSCO. Bioinformatics *39*, btad595. https://doi.org/10.1093/bioinformatics/btad595.

20. Kille, B., Garrison, E., Treangen, T.J., and Phillippy, A.M. (2023). Minmers are a generalization of minimizers that enable unbiased local Jaccard estimation. Bioinformatics *39*, btad512. https://doi.org/10.1093/bioinformatics/btad512.

21. Altemose, N., Logsdon, G.A., Bzikadze, A.V., Sidhwani, P., Langley, S.A., Caldas, G.V., Hoyt, S.J., Uralsky, L., Ryabov, F.D., Shew, C.J., et al. (2022). Complete genomic and epigenetic maps of human centromeres. Science *376*, eabl4178. https://doi.org/10.1126/science.abl4178.

22. Altemose, N. (2022). A classical revival: Human satellite DNAs enter the genomics era. Semin. Cell Dev. Biol. *128*, 2–14. https://doi.org/10.1016/j.semcdb.2022.04.012.

23. Vollger, M.R. (2026). rustybam: a composable toolkit for alignment analysis and visualization with SafFire. BioRxiv Prepr. Serv. Biol., 2026.02.16.706142. https://doi.org/10.64898/2026.02.16.706142.

24. Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26. https://doi.org/10.1038/nbt.1754.

25. Seabold, S., and Perktold, J. (2010). Statsmodels: Econometric and Statistical Modeling with Python. SciPy 2010. https://doi.org/10.25080/Majora-92bf1922-011.

26. Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0: fundamental algorithms for scientific computing in Python. Nat. Methods *17*, 261–272. https://doi.org/10.1038/s41592-019-0686-2.