The Stochastic Pacemaker: Cumulative Behavioral Noise Drives Morphological Plasticity in Pea Aphids

Xiaomi Liu[1*,] Kristopher Murdza[1], Yuxu Feng[1], Leila Lin[1], Elizabeth I. Croyle[1], Jennifer A. Brisson[1*]

1. Department of Biology, University of Rochester, Rochester, NY 14627, USA

Corresponding author: Xiaomi Liu (liuxiaomi1996@gmail.com), Jennifer A. Brisson (Jennifer.brisson@rochester.edu)

Abstract

Phenotypic variation within a single genotype under the same environment (intragenetic

variation), the biologically meaningful part of V_{error}, is frequently treated as a statistical

nuisance rather than a biological reality, yet it represents an evolutionary driver of

fitness that remains poorly integrated into evolutionary theory. The mechanism that

translates such stochasticity into deterministic developmental phenotypic outcomes is

not well understood. Here, we test a cumulative stochasticity model using wing

polyphenism of the pea aphid (*Acyrthosiphon pisum*), where asexual mothers produce

winged instead of wingless offspring in response to tactile cues. The model predicts that

stochastic variation in maternal locomotor behavior alters the rate of tactile cue

accumulation and thereby influences the probability of producing winged offspring. We

demonstrate that maternal locomotor activity acts as a "stochastic pacemaker", where

an individual's movement determines the rate at which it actively constructs its

environment and accumulates environmental cues. Our results reveal that genotypes

differ significantly in both mobility and the temporal pattern of wing induction, with

behavioral variation explaining approximately 20% of the total phenotypic variance

across genotypes. Crucially, we show that maternal mobility increases progressively

during crowding, accompanied by significant temporal heteroscedasticity. This

expanding variance and increase in mean suggest that initial, trivial stochasticity is

magnified into systematic behavioral divergence through the integration of

environmental signals. By demonstrating that total accumulated locomotor activity

predicts offspring phenotype, we provide a mechanistic bridge between transient

behavioral noise and stable morphological shifts. More broadly, our work reveals that

V_{error} is a dynamic product of behavioral history, suggesting a fundamental role for

individual-level niche construction in generating macro-phenotypic diversity.

Introduction

Understanding the source of phenotypic variation has long been a key focus in evolutionary biology because this variation is the target of natural selection. From a quantitative genetic view, phenotype variance can be partitioned as $V_P = V_G + V_E + V_{GxE} + V_{error}$, where each component represents a distinct source of variation (Lynch and Walsh, 1998; Tonsor et al., 2013; Westneat et al., 2015). Genetic variance (V_G) captures differences among genotypes within a single environment, whereas environmental variance (V_E) reflects phenotypic plasticity, the differences generated when genetically identical individuals experience different environments (Whitman and Agrawal, 2009). Genotype-by-environment interactions ($V_{G×E}$) represent heritable differences in plasticity itself, providing another target for evolutionary change (Via and Lande, 1985; West-Eberhard, 2003a; Whitman and Agrawal, 2009). The final component, the residual term, V_{error}, captures phenotypic differences among genetically identical individuals raised in nominally identical environments (Scheiner, 1993; Tonsor et al., 2013; Viney and Reece, 2013). Despite each component's potential to impact evolutionary dynamics, they differ markedly in how well their underlying mechanisms are understood.

Of particular interest here is the understudied role of V_{error}. This variation arises from several sources, such as measurement error, unmeasured or unmeasurable microenvironmental heterogeneity, and true stochastic developmental processes, like bet-hedging and stochastic gene expression (Kærn et al., 2005; Raj and Oudenaarden, 2008; Slatkin, 1974; Starrfelt and Kokko, 2012; Tonsor et al., 2013). The biologically meaningful portion of this residual variance, here referred to as intra-genotypic variation, can have a genetic basis and fitness consequences (Cai et al., 2008; Feinberg and Irizarry, 2010; Fraser et al., 2004; Kærn et al., 2005; Viney and Reece, 2013; Westneat et al., 2015). Yet despite its potential importance for evolution, intra-genotypic variation is frequently treated as a statistical nuisance rather than a biological reality, and has yet to be properly integrated in evolutionary theory (Cleasby and Nakagawa, 2011; Westneat et al., 2015). Characterizing this 'residual' component is essential for capturing the full spectrum of variation available to selection and for shedding light on understudied, interesting biological processes that shape phenotypic expression (Viney and Reece, 2013; Westneat et al., 2015).

Behavior provides a compelling candidate mechanism for generating such variation. As

the most flexible response to environmental change, behavior serves as a rapid and

dynamic interface between individuals and their environments (Duckworth, 2009; West-

Eberhard, 2003b). Behavioral traits are highly variable, even among clonal individuals

(Bierbach et al., 2017; Schuett et al., 2011), hinting that behavioral stochasticity may be

a primary driver of residual phenotypic variance (V_{error}). Furthermore, through niche

construction, organisms, especially through behavior, can modify their own micro-

environments, thereby altering the selective pressures and expression of non-

behavioral traits (Duckworth, 2009; Odling-Smee et al., 2003, 1996). Despite long-

standing theoretical predictions that behavior can modify the environment and thereby

influence the expression and evolution of other traits (Bateson, 2004; Duckworth, 2009;

West-Eberhard, 2003b), empirical studies that quantitatively link stochastic behavioral

noise to non-behavioral variance remain comparatively limited. Identifying such

connections would provide a critical, and currently missing perspective on how behavior

contributes to the generation and evolution of phenotypic diversity.

Aphids provide an exceptional opportunity to empirically study the connection between

behavioral variation to non-behavioral phenotypic outcomes. Their cyclical

parthenogenesis enables the production of genetically identical individuals, allowing

intra-genotypic variation to be isolated. Aphids also exhibit a well-characterized wing

polyphenism, in which asexual females give birth to clonal daughters that are either

winged or wingless depending on environmental conditions (Braendle et al., 2006;

Brisson, 2010; Deem et al., 2024; Sutherland, 1969). Importantly, tactile stimulation

generated during crowding is the key cue that induces more winged relative to wingless

offspring production (Johnson, 1965; Müller et al., 2001; Sutherland, 1969; Sutherland

and Mittler, 1971). Because tactile stimulation arises from physical interactions among

individuals, variation in movement directly modulates cue exposure. We propose that

individual locomotor behavior acts as an active, sensory-driven pace maker. Through

their own movement, individuals, even clonal individuals, actively modulate the

frequency of tactile stimuli they encounter, essentially "constructing" the environmental

signal they receive. Thus, behavior is not only a consequence of an organism's state,

but also the causal driver of the developmental trajectory.

In this study, we used the pea aphid wing polyphenism to examine how both inter- and intra-genotypic maternal locomotor behavioral variation shape offspring phenotype compositions. We test the hypothesis that intra-genotypic variation (V_{error}) is not an untraceable statistical constant, but a cumulative product of behavioral history. We propose a cumulative stochasticity model, where behavior acts as a "stochastic pacemaker": if locomotor activity facilitates tactile stimulation, then even minor initial differences in movement should undergo a process of linear accumulation over time. Our data demonstrate that genotypes differ significantly in both mobility and wing induction dynamics, with their correlation explaining about 20% of the total phenotypic variance. Crucially, within specific genotypes, individual mobility predicts winged offspring proportions, identifying behavior as a significant biological driver of V_{error}. By quantifying the temporal increase in locomotor and its variance over time, we show that mobility increases progressively during crowding, magnifying initial stochasticity into systematic behavioral divergence. This provides a mechanistic bridge between stochastic behavioral noise and deterministic morphological outcomes. More broadly, our work addresses a critical gap in evolutionary theory by demonstrating how both inter- and intra-genotypic variation arises through behavioral integration and scales to generate macro-phenotypic diversity.

Materials and Methods

Aphid Stocks: Pea aphid (*Acyrthosiphon pisum*) stocks were reared on *Vicia faba* seedlings (Improved Long Pod, Harris Seeds) by transferring four first instars on fresh seedlings every 11-13 days. Each plant was grown in a single plastic plant pot (S17648, Fisher Scientific) and enclosed in a cylindrical cage cut from clear PETG tubes (VISIPAK) sealed with fine mesh netting (Noseeum netting, 117 inches, Barre Army Navy Store). Cages were maintained in an incubator under long-day conditions (16L:8D), with a relative humidity of 30 ± 10% and a temperature of 19 ± 3.5°C. These environmental conditions enable the pea aphid to remain in an asexual life stage and reproduce viviparously. Stocks were reared under the described conditions for multiple generations to eliminate transgenerational density effects before the experiment.

Recording Setup: Four aphid genotypes were used: ROC1, SSC3, 375, and 663. Lines ROC1 and 375 show a pink color morph, while lines SSC3 and 663 show a green color morph (Figure 1A). To minimize stimuli from outside the recording, fresh adult aphids were collected directly from plants and immediately used for behavioral recording. Due to the limited ability of existing tracking software to reliably differentiate between visually similar moving individuals, particularly when animals frequently interact or obstruct one another, individual identity error is unavoidable(Dell et al., 2014). Here, we leveraged the natural color variation of pea aphids to ensure 100% identification accuracy. During the recording, one focal aphid of one color morph, either pink or green, was paired with 11 background aphids of the alternate color. Together, 12 aphids were placed in a 35 mm × 10 mm Petri dish. Six such Petri dishes were simultaneously recorded on a 5000K 70% brightness video light (RALENO, PLV-S116) for 7hrs using a mobile phone camera (Figure 1B). For the recording, line ROC1 was paired with line SSC3 and is hereafter referred to as pair 1 pink and pair 1 green, respectively. Similarly, lines 375 and 663 were paired and are referred to as pair 2 pink and pair 2 green. Following the recording, the focal aphids were individually transferred onto fresh seedlings and allowed to reproduce for 24 hours. The winged offspring percentage was scored upon adulthood. This experimental design enabled the accurate collection of behavioral and phenotypic data for each focal aphid, eliminating the risk of mismatches between behaviors and phenotypes due to limitations in tracking accuracy. We also collected an additional 3-hour recording for ROC1 (pair1 pink) for the reasons listed in the later context. All recordings started around 9 a.m..

Video Analysis: To estimate the amount of tactile stimulation aphids received during the recording period, we directly counted interactions. Two observers independently quantified interaction to account for human error. Due to time and resource limitations, we collected only interaction data for the pair 1pink recording.

To estimate aphid activity levels, we extracted each aphid's movement data for downstream analysis. One frame was extracted every two seconds from the original recording with FFmpeg (Tomar, 2006). Frames were then cleaned and processed using customized scripts and were subsequently split into pink and green channels using the ColorDeconvolution2 plugin (Ruifrok and Johnston, 2001) in ImageJ (Schneider et al.,

2012). Frames from the color channel containing focal aphids were further tidied up with

customized scripts and used to track aphid movement using TrackMate (Ershov et al.,

2021; Tinevez et al., 2017). The tracking results were exported as data tables

containing X and Y coordinates for aphid centroid positions in each frame. All positions

for a recording were plotted to determine the boundary for each Petri dish (Figure 1C).

Boundaries were assigned to different Petri dishes and corrected for false detection

using customized scripts. After false detection correction, each Petri dish was corrected

for camera distortion by resetting x boundaries to range from 0 to 400 (pixels) for both X

and Y coordinates. Frames from the color channel that contain background aphids were

tracked with StarDist (Schmidt et al., 2018). The trajectory results were exported as data

tables with trajectory distance for each identified track.

Profiling temporal dynamics in response to crowding: To test the effect of different

crowding durations on offspring wing percentage, we crowded aphids for varying

durations: 0 hours (no crowding), 4, 7, 10, or 24 hours. All treatments started around 8

a.m. During the crowding, 12 healthy adult aphids were placed in a 35 mm Petri dish

and then stored in an additional 90 mm Petri dish. A sheet of damped Waltman paper

was placed inside both the 35 mm dish and the larger Petri dish to prevent the aphids

from severe dehydration. After crowding, aphids were transferred individually onto fresh

seedlings and allowed to reproduce for 24 hours. For the 0hr group, aphids were

collected directly from the plants and placed on new seedlings for 24hrs to reproduce.

The offspring wing percentage was scored upon adulthood. Following the initial data

collection, based on the results, we selectively collected additional data at 1, 2, and 3hrs

for pair 1 pink and at 36 hours for pair 2 green.

Data analysis: We only included records in which treated aphids produced more than

two offspring. For both focal and background aphids, we measured activity level by

calculating the total distance the aphid moved. For the focal aphids, the distance was

calculated by summing up the geometric distance traveled between every frame. For

the background aphids, the total travel distance was calculated by summing up the

distance for all detected trajectories. Activity data, interaction number, and offspring

wing percentage data were all tested for normality with the Shapiro test. Because some

data were not normally distributed (Table S1), we used nonparametric tests or rank-

transformed data when needed. When multiple comparisons were performed, we

adjusted p-values with the Benjamini-Hochberg method (Benjamini and Hochberg,

1995).

All analyses were done in R 4.3.1.

Results

Locomotor activity strongly predicts tactile stimulation rate

To lay the foundation for downstream hypotheses and analyses, we first tested the

expectation that greater mobility would increase the likelihood of physical interaction

between aphids, thereby enhancing tactile stimulation. We measured interaction

number as a direct indicator of tactile stimulation, and quantified activity level using total

travel distance as a proxy for mobility. As expected, we observed a tight correlation

between the number of interactions and travel distance (Figure S1, Table S1, Spearman

Rank Correlation Test, Observer 1: correlation coefficient ρ = 0.883, adjusted p-value =

1.101×10^{-12}; Observer 2: correlation coefficient ρ = 0.893, adjusted p-value = 4.948 ×

10^{-13}). This result suggests that more active aphids experience greater tactile

stimulation, providing a basis for subsequent analyses examining the relationship

between activity and plasticity response.

Intra-genotypic variation: individual behavioral differences impact phenotypic variation

To investigate the source of V_{error}, we hypothesized that individual-level activity, which

varies even among clonal individuals, introduces microenvironmental heterogeneity that

contributes to final phenotype variation. We measured the final offspring wing

percentage phenotype and individual locomotion, and performed correlation tests within

each genotype. Significant correlations between offspring wing percentage and activity

level were detected in two out of the four tested genotypes (Figure 2, Kendall Rank

Correlation Test, pair 1 green adjusted p-value = 0.024, pair 2 pink adjusted p-value =

0.024). We then estimated how much of the phenotypic variance could be explained by

behavior. We calculated Kendall's tau^2 as a nonparametric proxy of association. For pair

1 (pink) and pair 2 (green) where significant correlations were detected, the tau^2 values

were 0.115 and 0.097, respectively. We also applied a linear regression model,

recognizing that the data did not follow a normal distribution; thus, interpretation should

be cautious. We obtained R^2 values of 0.248 and 0.177 for pair 1 pink and pair 2 green,

respectively. The higher R^2 and lower tau^2 suggest a generally linear yet noisy

relationship with tied ranks, consistent with the nature of our data (Figure 2). Taken

together, these results indicate that intra-genotypic individual-level activity accounts for

a considerable yet incomplete portion of final phenotype variation, helping to identify

part of the source of the V_{error} term.

Inter-genotypic variation: behavioral divergence across genotypes contributes to
228
phenotypic variation

To assess inter-genotypic variation in the wing percentage phenotype, we collected

offspring wing percentage data post-crowding and conducted a nonparametric statistical

test. We found that pair 1 pink and green have a higher propensity to produce winged

offspring compared to pair 2 pink and green (Figure 3A, Table S1, Kruskal-Wallis chi-

squared = 105.9, p-value = 8.3 x 10^{-23}, see detailed Dunn test result in Table S1),

supporting the idea that there are genetic differences underlying this wing induction

phenotype. The four tested genotypes also showed non-homogeneous levels of data

dispersion (Figure 3A, Table S1, Fligner-Killeen test, p-value = 2.797 × 10^{-6}), suggesting

that the variation level for the phenotype itself also has a genetic basis.

To assess intergenotypic variation in behavior, we calculated the total geometric

distance traveled by aphids, both as focal and background individuals, and conducted

nonparametric statistical tests. When acting as the single focal aphid, both pair 1 pink

and pair 1 green showed significantly higher activity levels compared to the two

genotypes from pair 2 (Figure 3B, Kruskal-Wallis chi-squared = 20.800, p-value = 1.158

x 10^{-4}, see detailed Dunn test results in Table S1). When comparing the activity level of

the background aphids, we found some similarity in behavior patterns, where pair 1 pink

moved significantly more than all other genotypes (Figure 3C, Kruskal-Wallis chi-

squared = 95.107, p-value < 1.751 x 10^{-20}, see detailed Dunn test results in Table S1).

However, pair 1 green now showed a similar activity level to pair 2 pink (Figure 3C; see

detailed Dunn test results in Table S1). Pair 2 green showed the lowest activity level

(Figure 3C). The movement of focal and background aphids cannot be directly

compared because they were collected using different tracking methods. Additionally,

background aphid travel distance represents the summed activity of all eleven

individuals within a Petri dish, reflecting summary statistics to some extent and resulting

in a narrower data dispersion. For both focal and background travel distance, pair 1 pink

exhibited greater variation than the other three genotypes (Fig. 3B, 3C, Fligner-Killeen

test, focal aphids distance p-value = 0.015, background aphids distance p-value = 1.093

$\times 10^{-13}$), suggesting increased individual-level behavioral variation in this genotype.

Together, our results show rich inter-genotypic variation in wing phenotype and

behavior.

We next examined how locomotion changed over time during the recording period.

Mean travel distance was calculated for each dish (1-6) and recording date using 5-min

windows. Time was standardized (mean-centered and scaled by its standard deviation)

to match the magnitude of speed and to improve numeric stability, yielding

approximately 2500 – 3500 observations per genotype. Mean travel distance was

analyzed using a linear mixed-effect model, where time, genotype, and interaction terms

were included as fixed effects. Individual identity was modeled with both random

intercept and slope to account for repeated measurements and individual trajectory.

Residual variance was allowed to vary as a function of time to test for

heteroscedasticity. As shown in Figure 3D, time had a strong positive impact on

locomotion. Genotype-specific slopes were estimated as follows: pair 1 pink (β = 4.10,

95% CI 3.23 - 4.98), pair 1 green (β = 2.84, 95% CI 1.88 - 3.80), pair 2 pink (β = 3.32,

95% CI 2.51 - 4.13), and pair 2 pink (β = 2.69, 95% CI 1.81 - 3.56), with all estimated

slopes significantly greater than 0. This result indicates a progressive increase in

locomotion throughout the assay. The time × genotype interaction was significant for

some genotypes (Table S1), demonstrating genotype-dependent differences in the

magnitude of temporal response relative to the reference genotype (pair 1 pink). Thus,

genotypes differed not only in baseline mobility but also in the rate of behavioral change

over time. Including individual-specific random slopes significantly improved model fit

compared to a random-intercept-only model (likelihood ratio test, p < 0.0001; Table S1),

indicating substantial among-individual variation in temporal trajectories. Furthermore,

allowing residual variance to scale with time significantly improved model fit relative to a

homoscedastic model (likelihood ratio test, p < 0.0001; Table S1, Figure 3D),

demonstrating an increase in within-genotype locomotor variability over time. Taken

together, these results revealed a strong temporal increase in locomotion behavior, as

well as inter- and intra-genotypic variation in movement dynamics.

After observing substantial variation within and among genotypes, we next examined

whether the relationship between activity and phenotype holds when all genotypes are

considered simultaneously. To do this, we modeled offspring wing percentage and

activity data from the four genotypes combined. Given that our data were enriched for

0% and 100%, we converted the percentages to proportions and employed a

generalized additive model with a quasibinomial family. The analysis revealed a

significant positive linear relationship between offspring wing percentage and focal

aphid movement (Figure 3E, p-value = 9.685×10^{-7}, effective degrees of freedom =

1.000). A similar significant but non-linear relationship was detected with the

background aphid movement data (Figure 3F, p-value = 0, effective degree of freedom

= 2.016). Additionally, the focal aphid movement explains 17.1% deviance, while the

background aphid movement explains 29.5% deviance (Table S1). Taken together, our

results suggest a general pattern: more active individuals produce a higher proportion of

winged offspring, implying that inter-genotypic behavioral variance can account for

approximately 20% of inter-genotypic phenotypic variance.

Genotype-specific Behavior-Sensitive Window

As we observed a general significant correlation between behavior and phenotype

across genotypes, and in two out of four genotypes, we next asked why we failed to

detect such a correlation in pair 1 pink, which nonetheless showed substantial variation

in both behavior and phenotype. We hypothesized that different genotypes exhibit

distinct dynamics in generating the plasticity response, thus affecting the behavior-

sensitive window. To test this, we measured wing percentage data across varying time

(0, 4, 7,10, or 24hrs) spent crowded to capture temporal dynamics. Our results revealed

drastically different responses among genotypes (Figure 4A). Pair 1 pink responded

rapidly to crowding, reaching a plateau after only 4 h, whereas pair 1 green and pair 3

pink reached their maximal responses after approximately 7 h and 10 h, respectively. In contrast, pair 2 green exhibited a general lack of response. To refine our temporal resolution, we collected additional early time points (1 h, 2 h, and 3 h) for pair 1 pink and an extended 36 h time point for pair 2 green to maximize signal detection, while maintaining the response shape. The maximum response at the plateau also differed among genotypes, with pair 1 pink and pair 1 green showing a high propensity for producing winged offspring, whereas pair 2 pink showed a moderate to low propensity (Figure 4A). Overall, these results demonstrate that time is a strong positive effector of the final phenotype in three out of four tested genotypes, supporting our hypothesis (Figure 4A, Kruskal-Wallis Test, adjusted p-value pair 1 pink: 5.544×10^{-11}, pair 1 green: 1.401×10^{-4}, pair 2 pink: 6.932×10^{-5}). The early saturation of the response in pair 1 pink explains the absence of a detectable behavior-phenotype correlation under the 7 h recording setting, as the phenotype was already at its plateau. Contrastingly, the lack of a correlation in pair 2 green likely comes from its limited overall responsiveness to crowding. As for pair 1 green and pair 2 pink, the 7 h window corresponds to the dynamic range in which variation in activity generates variation in tactile stimulation, resembling either shorter or longer crowding times, thus affecting the final phenotype output. Taken together, these results indicate that genotypes differ in their plasticity response dynamics, revealing inherent sensitive windows in which behavior-phenotype correlations can be detected.

Based on the unique behavior-sensitive window hypothesis, we predicted that a significant correlation between phenotype and behavior would be detected at an earlier time window for pair 1 pink. To test this, we repeated the recording at 3 h for pair 1 pink. The analysis detected a significant correlation between movement and offspring wing percentage, supporting our hypothesis (Figure 4B, Kendall Rank Correlation Test, p-value = 0.016). Similar to pair 1 green and pair 2 pink, 7h recording data, the tau2 is 0.093, suggesting a similar amount of variation is explained by intra-genotypic individual-level behavioral differences.

Discussion

A central contribution of this study is the empirical demonstration that maternal behavior serves as a "stochastic pacemaker", explaining a substantial portion of offspring wing plasticity both within and between pea aphid genotypes. At the intra-genotypic level, we show that seemingly stochastic differences in maternal movement create heterogeneous microenvironments by modulating active niche construction, altering the rate at which individuals encounter and accumulate tactile cues, which in turn alter the likelihood of producing winged offspring. In this framework, variance previously classified as stochastic residual error (V_{error}) is not random noise, but rather an integrated consequence of each mother's behavioral history. Behavioral stochasticity, therefore, becomes biologically consequential through cumulative signal acquisition. At the inter-genotypic level, the strong correlation between maternal behavior and offspring phenotype across genotypes indicates that heritable differences shape how mothers perceive, integrate, and respond to environmental cues. Importantly, genotypes differed not only in mean mobility and wing plasticity but also in the temporal dynamics of behavioral change under crowding. These differences suggest that selection can act not only on the magnitude of wing plasticity but also in the temporal dynamics of signal integration. Together, linking transient behavioral variability to stable morphological outcomes, our results provide empirical support for the cumulative stochasticity model and position behavioral variance as an evolvable component of developmental plasticity.

Under the Cumulative Stochasticity Model, maternal behavior modulates morph determination through iterative amplification of tactile cues. Tactile stimulation generated by movement promotes additional movement, gradually magnifying small initial differences in activity into substantial divergence in cue exposure over time. Behavioral responses can also potentially integrate other environmental inputs. For example, alarm pheromone ((E)-β-farnesene) induces "pseudo-crowding" by rapidly increasing walking behavior, which in turn elevates the proportion of winged offspring (Kunert et al., 2005). Similarly, intensified foraging behavior is observed under starvation, another nutrition-associated condition that promotes offspring wing production (Baz et al., 2025). Such behaviorally mediated cue amplification is widespread across insects. Because behavior is typically the earliest response to environmental change and is itself one of

the most plastic trait (Dingemanse and Wolf, 2013; Duckworth, 2009; Gosling, 2001; Sih et al., 2004; Wolf et al., 2008; Wolf and Weissing, 2012), it can act as a primary modifier that strengthens or prolongs initial cues before any downstream physiological development pathway are involved. Locust phase polyphenism provides a classic example of behavior reinforcement: solitary individuals increase aggregation behavior within hours of crowding, and this behavior then reinforces the gregarizing stimuli (Bouaïchi et al., 1995; Ellis, 1963; Gillett, 1973; Roessingh and Simpson, 1994; Rogers et al., 2003). Consistent with this general pattern, our data (Figure 3D) shows progressively increasing maternal activity as crowding proceeds. This temporal escalation supports the interpretation that behavior does not merely respond to crowding but actively intensifies cumulative signal exposure, thereby strengthening its contribution to morph determination.

Our findings integrate a model linking behavior, signaling dynamics, and offspring phenotypic plasticity. In this framework, mothers perceive tactile stimulation during crowding and transduce this information into signaling pathways, including glutamate, dopamine, ecdysone, and insulin/insulin-like growth factor signaling, that regulate wing plasticity (Brisson et al., 2026; Deem et al., 2024; Grantham et al., 2020; Liu and Brisson, 2023; Vellichirammal et al., 2017; Yuan et al., 2025, 2023). Behavioral variation potentially influences this system in two ways. First, differences in accumulated movement alter the intensity and frequency of tactile stimulation (Figure S1), thereby likely generating variation in signaling-molecule concentrations within and across genotypes. To evaluate how broadly such behavioral effects may apply, we examined multiple genotypes to increase the generality of our intra- and intergenotypic conclusions. Consistent with our intra-genotypic conclusion, another study using a different aphid genotype also reported a positive association between intra-genotypic behavioral activity and winged offspring production in a 2-individual, 2-h crowding setup (Yuan et al., 2025). Second, variation in the timing of the behaviorally sensitive window may interact with genotype-specific signaling thresholds to determine developmental outcomes, a hallmark of polyphenic systems (Nijhout, 2003, 1999; Pfennig, 2021). The interplay between behavioral modulation of cue exposure, genetically determined

thresholds, and the temporal dynamics of signal production likely all contribute to the
observed variation in winged-offspring production.

Despite considerable residual variance (V_{error}) being explained by inter- and intra-
genotypic behavioral variation, a large proportion of the variance remains unexplained.
One source of developmental stochasticity for V_{error} can be bet hedging, a strategy
favored when environmental cues are unreliable predictors of future conditions (Slatkin,
1974; Starrfelt and Kokko, 2012; Tonsor et al., 2013). Pea aphids, like many other
systems, employ a mixed plasticity and bet-hedging strategy: mothers stochastically
produce both morphs regardless of environmental conditions, ensuring that some
offspring match unpredictable environments and thereby increasing overall survival of
the species (Grantham et al., 2016). Such intrinsic stochasticity can blur the quantitative
relationship between cue intensity and phenotypic outcome.

Finally, our results provide empirical support for the theoretical expectation that
behavioral variation can shape, and in some cases even lead, the evolution of other
traits (Duckworth, 2009; Odling-Smee et al., 2003, 1996; West-Eberhard, 2003b). In our
system, maternal locomotion does not merely respond to environmental conditions; it
modulates the effective strength and duration of crowding cues experienced by the
individual. In doing so, behavior alters the local developmental environment that
determines offspring morphology. This aligns with niche construction theory, which
emphasizes that organisms actively modify their environments and thereby alter the
selective pressures they experience (Odling-Smee et al., 2003, 1996). Under this
framework, variation in movement rate translates directly into variation in cumulative
signal exposure, thereby influencing the probability of wing induction. As we show, even
clonal individuals present phenotypic variation rooted in maternal behavior history,
demonstrating that behavioral variation can generate structured developmental
divergence without genetic differences. More broadly, our findings suggest that
individual-level micro-environmental heterogeneity may represent an underappreciated
mechanism through which behavior contributes to macro phenotypic diversity.

Figures and table

Figure1. Experimental setup and visualization of behavioral recordings.

(A) Representative photo of the four pea aphid genotypes used in the study. From left to right: pair 1 pink,
pair 1 green, pair 2 pink, and pair 2 green. (B) Representative single frame from a behavioral recording
showing aphids within the observation arena. (C) Overlay of all detected aphid centroid positions across
the full recording period for one example recording, with 1 position sample every 2s, illustrating spatial
coverage of aphid movement within the arena.

Figure 2. Intra-genotypic variation in maternal locomotor behavior correlates with
offspring wing phenotype in some genotypes.

Scatter plots show the relationship between focal aphid locomotor activity (x-axis) and the percentage of
winged offspring (y-axis). Each panel shows a single pea aphid genotype, with each point representing
one focal individual.

Figure 3. Inter-genotypic variation in maternal locomotor behavior contributes to the offspring wing phenotype.

(A) Offspring winged percentage produced by focal aphids from each of four genotypes. (B) Locomotor activity of *focal* aphids from each of four genotypes. (C) Locomotor activity of aphids from the same genotypes when serving as *background* individuals. D) Temporal profile of locomotor activity, showing per-frame movement averaged across 5 min time windows (y-axis) as a function of time during recording (x-axis). Colors represent aphid color morphs; each panel represents one of the four genotypes. (E) Association between *focal* aphid locomotor activity and *focal* aphid offspring wing proportion. (F) Association between *background* aphid locomotor activity and *focal* aphid offspring wing proportion. For panels E and F, the raw data points are shown as small grey points. Data were grouped into 25 bins along the x-axis (distance moved), with uneven bin widths to yield approximately equal sample sizes per

bin (5–7 points). Black points represent the mean value for each bin, with the light gray lines indicating

the binomial standard error. GAM model fits are shown as light grey dashed lines.

Figure 4. Genotypes exhibit distinct temporal patterns of wing induction in response to

crowding.

Relationship between crowding duration (x-axis) and offspring wing percentage (y-axis). Each small panel

corresponds to each of the four genotypes.

Figure S1. Tight correlation between aphid locomotor activity and interaction frequency.

Interaction number (y-axis) is plotted as a function of focal aphids locomotor activity (x-axis). Colors and

point shapes represent two independent observers who quantified the aphid-aphid interactions.

Table S1. Detailed statistical analysis table.

Acknowledgements

This research is supported by the National Institute of General Medical Sciences within the National Institutes of Health under award number R35GM144001 to JAB. We thank Arron L. Forbes for helpful discussions and assistance in designing the interaction-counting criteria used in this study.

Reference

Bateson, P., 2004. The Active Role of Behaviour in Evolution. Biology & Philosophy 19, 283–298. https://doi.org/10.1023/B:BIPH.0000024468.12161.83

Baz, E.-S., Huygens, C., Calevro, F., Callaerts, P., 2025. Analysis of the impact of extrinsic and intrinsic factors on aphid locomotion using a novel machine-learning based system to track insect behavior. Entomologia Generalis 45.

Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal statistical society: series B (Methodological) 57, 289–300.

Bierbach, D., Laskowski, K.L., Wolf, M., 2017. Behavioural individuality in clonal fish arises despite near-identical rearing conditions. Nat Commun 8, 15361. https://doi.org/10.1038/ncomms15361

Bouaïchi, A., Roessingh, P., Simpson, S.J., 1995. An analysis of the behavioural effects of crowding and re-isolation on solitary-reared adult desert locusts (Schistocerca gregaria) and their offspring. Physiological Entomology 20, 199–208. https://doi.org/10.1111/j.1365-3032.1995.tb00002.x

Braendle, C., Davis, G.K., Brisson, J.A., Stern, D.L., 2006. Wing dimorphism in aphids. Heredity 97, 192–199. https://doi.org/10.1038/sj.hdy.6800863

Brisson, J.A., 2010. Aphid wing dimorphisms: linking environmental and genetic control of trait variation. Philosophical Transactions of the Royal Society B: Biological Sciences 365, 605–616. https://doi.org/10.1098/rstb.2009.0255

Brisson, J.A., Deem, K.D., Liu, X., 2026. Aphid wing plasticity: hormonal and epigenetic control mechanisms. Current Opinion in Insect Science 75, 101492. https://doi.org/10.1016/j.cois.2026.101492

496 Cai, L., Dalal, C.K., Elowitz, M.B., 2008. Frequency-modulated nuclear localization
497   bursts coordinate gene regulation. Nature 455, 485–490.
498   https://doi.org/10.1038/nature07292

499 Cleasby, I.R., Nakagawa, S., 2011. Neglected biological patterns in the residuals. Behav
500   Ecol Sociobiol 65, 2361–2372. https://doi.org/10.1007/s00265-011-1254-7

501 Deem, K.D., Gregory, L.E., Liu, X., Ziabari, O.S., Brisson, J.A., 2024. Evolution and
502   molecular mechanisms of wing plasticity in aphids. Current Opinion in Insect
503   Science 61, 101142. https://doi.org/10.1016/j.cois.2023.101142

504 Dell, A.I., Bender, J.A., Branson, K., Couzin, I.D., Polavieja, G.G. de, Noldus, L.P.J.J.,
505   Pérez-Escudero, A., Perona, P., Straw, A.D., Wikelski, M., Brose, U., 2014.
506   Automated image-based tracking and its application in ecology. Trends in
507   Ecology & Evolution 29, 417–428. https://doi.org/10.1016/j.tree.2014.05.004

508 Dingemanse, N.J., Wolf, M., 2013. Between-individual differences in behavioural
509   plasticity within populations: causes and consequences. Animal Behaviour,
510   Special Issue: Behavioural Plasticity and Evolution 85, 1031–1039.
511   https://doi.org/10.1016/j.anbehav.2012.12.032

512 Duckworth, R.A., 2009. The role of behavior in evolution: a search for mechanism. Evol
513   Ecol 23, 513–531. https://doi.org/10.1007/s10682-008-9252-6

514 Ellis, P.E., 1963. Changes in the social aggregation of locust hoppers with changes in
515   rearing conditions. Animal Behaviour 11, 152–160. https://doi.org/10.1016/0003-
516   3472(63)90023-X

517 Ershov, D., Phan, M.-S., Pylvänäinen, J.W., Rigaud, S.U., Le Blanc, L., Charles-Orszag,
518   A., Conway, J.R., Laine, R.F., Roy, N.H., Bonazzi, D., 2021. Bringing TrackMate
519   into the era of machine-learning and deep-learning. BioRxiv 2021–09.

520 Feinberg, A.P., Irizarry, R.A., 2010. Stochastic epigenetic variation as a driving force of
521   development, evolutionary adaptation, and disease. Proceedings of the National
522   Academy of Sciences 107, 1757–1764. https://doi.org/10.1073/pnas.0906183107

Fraser, H.B., Hirsh, A.E., Giaever, G., Kumm, J., Eisen, M.B., 2004. Noise Minimization
in Eukaryotic Gene Expression. PLOS Biology 2, e137.
https://doi.org/10.1371/journal.pbio.0020137

Gillett, S.D., 1973. Social determinants of aggregation behaviour in adults of the desert
locust. Animal Behaviour 21, 599–606. https://doi.org/10.1016/S0003-
3472(73)80022-3

Gosling, S.D., 2001. From mice to men: What can we learn about personality from
animal research? Psychological Bulletin 127, 45–86.
https://doi.org/10.1037/0033-2909.127.1.45

Grantham, M.E., Antonio, C.J., O'Neil, B.R., Zhan, Y.X., Brisson, J.A., 2016. A case for a
joint strategy of diversified bet hedging and plasticity in the pea aphid wing
polyphenism. Biology Letters 12, 20160654.
https://doi.org/10.1098/rsbl.2016.0654

Grantham, M.E., Shingleton, A.W., Dudley, E., Brisson, J.A., 2020. Expression profiling
of winged- and wingless-destined pea aphid embryos implicates insulin/insulin
growth factor signaling in morph differences. Evolution & Development 22, 257–
268. https://doi.org/10.1111/ede.12326

Johnson, B., 1965. Wing Polymorphism in Aphids Ii. Interaction Between Aphids.
Entomologia Experimentalis et Applicata 8, 49–64. https://doi.org/10.1111/j.1570-
7458.1965.tb02342.x

Kærn, M., Elston, T.C., Blake, W.J., Collins, J.J., 2005. Stochasticity in gene expression:
from theories to phenotypes. Nature Reviews Genetics 6, 451–464.
https://doi.org/10.1038/nrg1615

Kunert, G., Otto, S., Röse, U.S.R., Gershenzon, J., Weisser, W.W., 2005. Alarm
pheromone mediates production of winged dispersal morphs in aphids. Ecology
Letters 8, 596–603. https://doi.org/10.1111/j.1461-0248.2005.00754.x

Liu, X., Brisson, J.A., 2023. Dopamine mediates the pea aphid wing plasticity. Biol. Lett.
19, 20230024. https://doi.org/10.1098/rsbl.2023.0024

551 Lynch, M., Walsh, B., 1998. Genetics and analysis of quantitative traits. Sinauer

552   Sunderland, MA.

553 Müller, C.B., Williams, I.S., Hardie, J., 2001. The role of nutrition, crowding and

554   interspecific interactions in the development of winged aphids. Ecological

555   Entomology 26, 330–340. https://doi.org/10.1046/j.1365-2311.2001.00321.x

556 Nijhout, H.F., 2003. Development and evolution of adaptive polyphenisms. Evolution &

557   Development 5, 9–18. https://doi.org/10.1046/j.1525-142X.2003.03003.x

558 Nijhout, H.F., 1999. Control Mechanisms of Polyphenic Development in Insects: In

559   polyphenic development, environmental factors alter some aspects of

560   development in an orderly and predictable way. BioScience 49, 181–192.

561   https://doi.org/10.2307/1313508

562 Odling-Smee, F.J., Lala, K.N., Feldman, M., 2003. Niche Construction: The Neglected

563   Process in Evolution. Princeton University Press, Princeton, UNITED STATES.

564 Odling-Smee, F.J., Laland, K.N., Feldman, M.W., 1996. Niche Construction. The

565   American Naturalist 147, 641–648.

566 Pfennig, D.W., 2021. Phenotypic plasticity & evolution: causes, consequences,

567   controversies. Taylor & Francis.

568 Raj, A., Oudenaarden, A. van, 2008. Nature, Nurture, or Chance: Stochastic Gene

569   Expression and Its Consequences. Cell 135, 216–226.

570   https://doi.org/10.1016/j.cell.2008.09.050

571 Roessingh, P., Simpson, S.J., 1994. The time-course of behavioural phase change in

572   nymphs of the desert locust, Schistocerca gregaria. Physiological Entomology

573   19, 191–197. https://doi.org/10.1111/j.1365-3032.1994.tb01042.x

574 Rogers, S.M., Matheson, T., Despland, E., Dodgson, T., Burrows, M., Simpson, S.J.,

575   2003. Mechanosensory-induced behavioural gregarization in the desert locust

576   Schistocerca gregaria. Journal of Experimental Biology 206, 3991–4002.

577 Ruifrok, A.C., Johnston, D.A., 2001. Quantification of histochemical staining by color

578   deconvolution. Anal Quant Cytol Histol 23, 291–299.

Scheiner, S.M., 1993. Genetics and Evolution of Phenotypic Plasticity. Annual Review of
Ecology, Evolution, and Systematics 24, 35–68.
https://doi.org/10.1146/annurev.es.24.110193.000343

Schmidt, U., Weigert, M., Broaddus, C., Myers, G., 2018. Cell Detection with Star-
Convex Polygons, in: Frangi, A.F., Schnabel, J.A., Davatzikos, C., Alberola-
López, C., Fichtinger, G. (Eds.), Medical Image Computing and Computer
Assisted Intervention – MICCAI 2018, Lecture Notes in Computer Science.
Springer International Publishing, Cham, pp. 265–273.
https://doi.org/10.1007/978-3-030-00934-2_30

Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of
image analysis. Nature methods 9, 671–675.

Schuett, W., Dall, S.R.X., Baeumer, J., Kloesener, M.H., Nakagawa, S., Beinlich, F.,
Eggers, T., 2011. Personality variation in a clonal insect: The pea aphid,
Acyrthosiphon pisum. Developmental Psychobiology 53, 631–640.
https://doi.org/10.1002/dev.20538

Sih, A., Bell, A.M., Johnson, J.C., Ziemba, R.E., 2004. Behavioral Syndromes: An
Integrative Overview. The Quarterly Review of Biology 79, 241–277.
https://doi.org/10.1086/422893

Slatkin, M., 1974. Hedging one's evolutionary bets. Nature 250, 704–705.
https://doi.org/10.1038/250704b0

Starrfelt, J., Kokko, H., 2012. Bet-hedging—a triple trade-off between means, variances
and correlations. Biological Reviews 87, 742–755. https://doi.org/10.1111/j.1469-
185X.2012.00225.x

Sutherland, O.R.W., 1969. The role of crowding in the production of winged forms by
two strains of the pea aphid, *Acyrthosiphon pisum*. Journal of Insect Physiology
15, 1385–1410. https://doi.org/10.1016/0022-1910(69)90199-1

Sutherland, O.R.W., Mittler, T.E., 1971. Influence of diet composition and crowding on
wing production by the aphid *Myzus persicae*. Journal of Insect Physiology 17,
321–328. https://doi.org/10.1016/0022-1910(71)90216-2

Tinevez, J.-Y., Perry, N., Schindelin, J., Hoopes, G.M., Reynolds, G.D., Laplantine, E.,
Bednarek, S.Y., Shorte, S.L., Eliceiri, K.W., 2017. TrackMate: An open and
extensible platform for single-particle tracking. Methods 115, 80–90.

Tomar, S., 2006. Converting video formats with FFmpeg. Linux journal 2006, 10.

Tonsor, S.J., Elnaccash, T.W., Scheiner, S.M., 2013. DEVELOPMENTAL INSTABILITY
IS GENETICALLY CORRELATED WITH PHENOTYPIC PLASTICITY,
CONSTRAINING HERITABILITY, AND FITNESS. Evol 67, 2923–2935.
https://doi.org/10.1111/evo.12175

Vellichirammal, N.N., Gupta, P., Hall, T.A., Brisson, J.A., 2017. Ecdysone signaling
underlies the pea aphid transgenerational wing polyphenism. Proceedings of the
National Academy of Sciences 114, 1419–1423.
https://doi.org/10.1073/pnas.1617640114

Via, S., Lande, R., 1985. Genotype-Environment Interaction and the Evolution of
Phenotypic Plasticity. Evolution 39, 505–522. https://doi.org/10.1111/j.1558-
5646.1985.tb00391.x

Viney, M., Reece, S.E., 2013. Adaptive noise. Proceedings of the Royal Society B:
Biological Sciences 280, 20131104. https://doi.org/10.1098/rspb.2013.1104

West-Eberhard, M.J., 2003a. Developmental plasticity and evolution. Oxford University
Press.

West-Eberhard, M.J., 2003b. Developmental plasticity and evolution. Oxford University
Press.

Westneat, D.F., Wright, J., Dingemanse, N.J., 2015. The biology hidden inside residual
within-individual phenotypic variation. Biological Reviews 90, 729–743.
https://doi.org/10.1111/brv.12131

Whitman, D.W., Agrawal, A.A., 2009. What is phenotypic plasticity and why is it
important. Phenotypic plasticity of insects: Mechanisms and consequences 1–63.

Wolf, M., van Doorn, G.S., Weissing, F.J., 2008. Evolutionary emergence of responsive

and unresponsive personalities. Proceedings of the National Academy of

Sciences 105, 15825–15830. https://doi.org/10.1073/pnas.0805473105

Wolf, M., Weissing, F.J., 2012. Animal personalities: consequences for ecology and

evolution. Trends in Ecology & Evolution 27, 452–461.

https://doi.org/10.1016/j.tree.2012.05.001

Yuan, Y., Wang, Y., Ye, W., Xie, L., Yuan, E., Guo, H., Wang, S., Dong, F., Zhu-Salzman,

K., Ge, F., Sun, Y., 2025. The maternal vGluT2 and embryonic mGluR3 signaling

relay system controls offspring wing dimorphism in pea aphid. iScience 28.

https://doi.org/10.1016/j.isci.2025.112591

Yuan, Y., Wang, Y., Ye, W., Yuan, E., Di, J., Chen, X., Xing, Y., Sun, Y., Ge, F., 2023.

Functional evaluation of the insulin/insulin-like growth factor signaling pathway in

determination of wing polyphenism in pea aphid. Insect Science 30, 816–828.

https://doi.org/10.1111/1744-7917.13121