Title: Coordinated IL-2 and TGF-β Signaling via a Novel Fusion Protein Selectively Expands and

Activates Regulatory T Cells

One Sentence Summary

CUE-401 coordinates IL-2 and TGF-β signaling to expand and activate regulatory T cells and

restrain autoimmune responses.

Authors

Nicholas M. Jackson[1], Javier A. Carrero[1], Kristin Yeung[2], Evelyn Needham[2], Emily Christie[2], Natalie

Yakobian[1], Challen Pretorious[1], Stella Hoft[1], Kirsten Foley[1], Eric Ford[1], Simon Low[2], Ronald Seidel[2],

Anish Suri[2], Natasha M. Girgis[2], Steven N. Quayle[2], Richard J. DiPaolo[1*]

[1]Department of Molecular Microbiology and Immunology, Saint Louis University School of Medicine,

Saint Louis, Missouri

[2]CUE Biopharma, Boston, Massachusetts

Corresponding author, richard.dipaolo@health.slu.edu

Abstract

Safely and selectively re-establishing immune tolerance remains a central challenge in the treatment of

autoimmune and inflammatory diseases. Regulatory T cells (Tregs) are essential mediators of immune

homeostasis and peripheral tolerance. Interleukin-2 (IL-2) and transforming growth factor–β (TGF-β)

synergize to promote Treg activation, differentiation, and stability, but coordinating selective in vivo

delivery of these signals has limited potential therapeutic applications. Here, we describe CUE-401, a

bifunctional biologic consisting of attenuated IL-2 and attenuated, receptor-masked TGF-β3 to promote

co-signaling on target cells. In vitro, CUE-401 induced naïve CD4$^+$ T cells to convert to Foxp3$^+$ induced

Tregs (iTregs), expanded endogenous Tregs, and suppressed inflammatory cytokine production by

memory and effector CD4$^+$ T cells. In mice, a single dose of CUE-401 markedly increased the frequency

of CD4$^+$Foxp3$^+$ Tregs exhibiting a stable and activated phenotype, with minimal activation of

CD4+Foxp3− conventional T cells (Tconv) relative to an IL-2-only comparator. In a

Treg-deficiency-driven adoptive transfer model of autoimmunity, early administration of CUE-401

activated Tregs, reduced gastric infiltration by autoreactive T cells, and provided long-term protection

from autoimmune gastritis in the absence of continued treatment. Together, these findings support

CUE-401 as a dual IL-2/TGF-β3 biologic that promotes immune tolerance through coordinated induction,

expansion, and activation of Tregs while limiting activation of autoreactive CD4+ T cells.

Introduction

Autoimmune diseases arise from the breakdown of peripheral immune tolerance, resulting in persistent

immune-mediated damage to self-tissues. The discovery of CD4+CD25+ regulatory T cells (Tregs)

marked a paradigm shift in understanding immune tolerance (*1*). Tregs maintain tolerance by suppressing

autoreactive effector T cell responses while preserving protective immunity through multiple

mechanisms, including consumption of interleukin-2 (IL-2), expression of inhibitory receptors (e.g.

CTLA-4), and production of immunoregulatory cytokines, such as transforming growth factor–β (TGF-β)

(*2*). The essential role of Tregs in immune homeostasis is underscored by the severe autoimmune

pathology observed in humans and mice with mutations in the transcription factor Foxp3, which is

required for Treg lineage commitment and stability (*3-6*).

Tregs are classically defined by expression of CD4, the high-affinity IL-2 receptor α-chain (CD25), and

Foxp3. This population includes thymus-derived natural Tregs (nTregs) and peripherally induced Tregs

(pTregs), which arise from conventional CD4+ T cells in response to antigenic stimulation and cytokine

cues (*7, 8*). In vitro, naïve CD4+ T cells stimulated with TCR signals plus TGF-β, and IL-2 differentiate

into Foxp3+ induced Tregs (iTregs) with potent suppressive activity (*9*). Precise modulation of IL-2–

dependent STAT5 activation together with TGF-β–mediated Foxp3 induction is also critical for

maintaining stable, suppressive Tregs (*9-11*). Accordingly, this cytokine combination has been widely

used to generate suppressive iTregs ex vivo for adoptive cell therapies that are effective in preclinical

models of autoimmunity and graft-versus-host disease (*12-14*).

The immunosuppressive capacity of Tregs has motivated their therapeutic use in transplantation and

immune-mediated diseases. Several different IL-2 variants (muteins) have been engineered to activate and

expand Tregs and are being tested for effectiveness in treating inflammatory diseases (*15-17*). Key

challenges include a limited therapeutic window due to off-target stimulation of non-Treg populations, the

short half-life of IL-2, and the need for precise dosing to achieve selective Treg engagement. Moreover,

Treg induction, stability, and specialization in vivo often require additional cues, including TGF-β.

Coordinating IL-2 and TGF-β signaling in vivo is challenging due to short in vivo half-lives of cytokines,

tolerability considerations, and the need to co-engage receptors on the same target cell(s). CUE-401, a

novel Fc-based IL-2/TGF-β3 fusion protein designed for coordinated delivery of IL-2 and TGF-β signals

in vivo, was designed to address these challenges. CUE-401 includes an attenuated IL-2 mutein (F42A,

H16A) with reduced affinity for CD25 and CD122 (*18-20*), and a novel receptor-masked attenuated TGF-

β3 variant, linked via flexible linkers to an Fc scaffold to promote IL-2/TGF-β3 co-signaling and extend

half-life. All amino acid sequences are human-derived, and both cytokines cross-react with mouse

receptors (*21, 22*). Here, we evaluated the biological activities of CUE-401 in vitro and in vivo for its

ability to enhance Tregs, suppress effector T cell responses, and mitigate autoimmune pathology.

Results

CUE-401 delivers IL-2 and TGF-β signals to T cells

CUE-401 is a bispecific fusion protein composed of affinity attenuated variants of IL-2 and TGF-β3 (Fig.

1A). The IL-2 mutein (F42A and H16A) reduces affinity to IL-2Rα (CD25) and IL-2Rβ (CD122) (*18*). The

TGF-β3 component is engineered as a monomeric, receptor masked variant to enhance manufacturability

and attenuate receptor affinity. The TGF-β3 moiety in CUE-401 carries a C77S mutation that prevents

formation of the interchain disulfide bond while maintaining its ability to assemble the natural signaling

complex. The masking component consists of a modified TGFβR2 extracellular domain lacking the

N-terminal 25 amino acids required for TGFβR1 interaction and incorporating an D118A mutation to

reduce affinity for TGF-β3 relative to the wild-type receptor (*23, 24*). This masked configuration preserves

biological activity but enables CUE-401 to be manufactured at scale in soluble form and subsequently

activated upon engagement at the cell surface. Both cytokine domains are connected via flexible linkers to

a heterodimeric Fc scaffold to promote co-signaling, enhance molecular stability, extend serum half-life.

The IL-2 activity of the CUE-401 molecule was quantified using CTLL-2 proliferation assays. Three

conditions were tested: 1. wild-type human IL-2 (Proleukin), 2. CUE-401, and 3. A version of CUE-401

that contains the IL-2 mutein (F42A and H16A) but is lacking a TGF-β3 (referred to as CUE-401ΔTGFβ;

Fig. 1B). Both CUE-401 and CUE-401ΔTGFβ induced dose-dependent proliferation that was attenuated

compared to Proleukin (Fig. 1B). TGF-β3 activity of CUE-401 was demonstrated by SMAD2/3

phosphorylation in murine splenocytes treated with CUE-401 or recombinant TGF-β3, but not with rIL-2

alone, or in cells that were pretreated with a selective inhibitor of TGF-βR signaling, SB431542 (Fig. 1C).

CUE-401 induces and expands Tregs and suppresses effector T cell function in vitro

After confirming that both cytokine components of CUE-401 were biologically active, we determined

whether CUE-401 could induce de novo Foxp3 expression in CD4[+] T cells. CD4[+]CD8[-]Foxp3.GFP[-] thymic

T cells were sorted from thymi of BALB/c.Foxp3.eGFP mice and stimulated with anti-CD3/anti-CD28 in

the presence of recombinant human IL-2 (rhIL-2), rhIL-2 plus recombinant human TGF-β3 (rhTGF-β3), or

CUE-401 (Fig. 2). IL-2 alone induced minimal Foxp3 expression (~1%), whereas both rhIL-2 + rhTGF-β3,

and CUE-401 induced Foxp3 expression in >75% of CD4[+] T cells (Fig. 2A). CUE-401 induced Foxp3

expression in a dose-dependent manner, with an EC_{50} of ~18 nM (Fig. 2B). These data demonstrate that

CUE-401 is effective at co-delivering IL-2/TGF-β signals required for Foxp3[+] iTreg induction from naïve

CD4 T cells.

To assess whether CUE-401 expands endogenous CD4[+]Foxp3[+] Tregs, CD4[+]Foxp3.eGFP[+] Tregs sorted

from the spleens of BALB/c.Foxp3.eGFP mice were stimulated with anti-CD3/anti-CD28 in the presence

of either rhIL-2 (control) or CUE-401. As expected, Tregs maintained Foxp3 expression when cultured

with rhIL-2 (Fig. 2C). Similarly, Tregs stimulated in the presence of CUE-401 also maintained Foxp3

expression and expanded in a dose-dependent manner (Fig. 2D). These results demonstrate that CUE-401

effectively promotes Treg maintenance and expansion in vitro.

To determine the effects of CUE-401 on effector/memory T cells, CD4[+]CD44[hi]Foxp3.eGFP[−] cells sorted

from spleens of BALB/c.Foxp3.eGFP mice were stimulated with anti-CD3/anti-CD28 in the presence of

either rhIL-2, rhIL-2 + rhTGF-β3, or CUE-401 (Fig. 2E). Treatment with rhIL-2 resulted in minimal

induction of Foxp3 expression (~1%). However, stimulation of these purified effector/memory T cells in

the presence of rhIL-2 + rhTGF-β3 or CUE-401 induced Foxp3 expression in ~10% of effector/memory

CD4[+] T cells (Fig. 2E-F). Next, we assessed cytokine secretion by effector/memory CD4[+] T cells from

these same cultures. In the presence of IL-2, effector/memory T cells secreted IFN-γ, IL-4, TNF, and IL-

17A. However, CUE-401 significantly decreased the secretion of IFN-γ, IL-4 and TNF from these

stimulated effector cells compared to rhIL-2 alone (Fig. 2G). Together, these findings establish three key

biological activities of CUE-401 in vitro: (1) induction of iTregs from naïve CD4[+] T cells, (2) maintenance

of Foxp3 expression and expansion of existing Tregs, and (3) induction of Foxp3 expression in a subset

and reduced proinflammatory cytokine production by effector/memory CD4[+] T cells.

CUE-401 administration induces robust expansion of stable regulatory T cells in mice

To evaluate the biological activities of CUE-401 in vivo, BALB/c mice received a single dose of CUE-401

or vehicle (PBS). By day 6, Treg frequencies in the spleen of CUE-401 treated mice increased to ~61% of

CD4[+] T cells compared to ~20% in controls (Fig. 3A-B). To characterize the phenotype and stability of

Tregs expanded by CUE-401 in vivo, we assessed methylation of the Treg-specific demethylated region

(TSDR), a CpG-rich enhancer within the *Foxp3* locus (CNS2) (*25*). Splenocytes from vehicle and CUE-

401 treated mice were sorted to be CD4$^+$Foxp3.eGFP$^+$ and CD4$^+$Foxp3.eGFP$^-$ (Fig. 3C). Tregs from CUE-

401-treated mice exhibited comparable demethylation of the TSDR to that observed in natural Tregs in

control mice, whereas the TSDR in Tconv cells from CUE-401 treated mice was similarly methylated as in

natural Tconv cells from untreated animals (Fig. 3D). Furthermore, Tregs from both groups maintained

stable Foxp3 expression after restimulation for 6 days in vitro in the absence of further treatment (Fig. 3E).

These findings demonstrate that CUE-401 expands stable, lineage-committed Tregs in vivo.

Expanded regulatory T cells exhibit an activated phenotype following CUE-401 administration

After establishing that CUE-401 administration induced a robust expansion of Tregs, we next compared

Treg phenotypes between treated and control mice 6 days after treatment. As shown above, CUE-401

significantly increased the frequency of CD4$^+$Foxp3$^+$ Tregs compared with PBS-treated controls (Fig. 4A).

These Tregs were then analyzed for expression of markers of Treg activation and suppressor functions (*26,*

27), including Foxp3, GITR, CD25, CTLA-4, CD103, CD73, CD38, CD134, CD137, and Ki67. Flow

cytometry and UMAP analysis identified enrichment of activated Treg clusters in CUE-401–treated mice,

with increased frequencies of GITR^{hi}/CD25^{hi} and CTLA-4^{hi}/CD103^{hi} Tregs, and higher expression of

CD25, CTLA-4, and GITR overall (Fig. 4A–C).

To gain deeper insight into the effects of administering CUE-401 on Tregs, we compared gene expression

profiles on Tregs from CUE-401 treated and control mice (Fig. 4D). Bulk RNA-seq of purified Tregs

revealed upregulation of several genes associated with Treg activation and suppressive function in CUE-

401 treated mice, including *Itgae* (CD103), *Nrp1* (neuropilin-1), *Il2ra* (CD25), *Il2rb* (IL-2Rβ), *Lag3* (LAG-

3), *Tnfrsf18* (GITR), *Icosl* (ICOSL), and *Klrg1* (KLRG1) (*28-32*). Together, these data demonstrate that

CUE-401 not only expands a stable Treg population in vivo but also promotes a transcriptional program

associated with enhanced activation and suppressive capacity (*33-36*).

Combined IL-2 and TGF-β signaling responses in Tregs are distinct from IL-2 alone

To dissect the contribution of TGF-β3 and to compare the biological activity of CUE-401 to a biologic that delivers only IL-2, we compared CUE-401 to CUE-401ΔTGF-β3, which contain the identical variant of IL-2. In vivo, both molecules induced significant expansion of splenic Tregs, although Treg frequencies were higher in CUE-401-treated mice (Fig. 5A, B). CUE-401 also increased the proportion of GITR^{hi} CD38^{hi} Tregs more than CUE-401ΔTGF-β3 did (Fig. 5C, D). Furthermore, the IL-2–only version increased CD25 and CD103 expression on Tconv, whereas the TGF-β3-containing CUE-401 did not (Fig. 5E–F).

Transcriptomic analyses of sorted CD4⁺Foxp3.GFP⁺ Tregs and CD4⁺Foxp3.GFP⁻ Tconv from PBS, CUE-401ΔTGF-β3, and CUE-401 treated mice also revealed different biological activities of IL-2 in the presence or absence of TGF-β3 signaling. Both CUE-401ΔTGF-β3 and CUE-401 increased the expression of core Treg-associated genes, including: *Ctla4*, *Foxp3*, *Il2ra*, and *Lag3*, relative to PBS-treated controls (Fig. 5G). However, several of the IL-2 induced genes induced in Tregs by the IL-2 only version (CUE-401ΔTGF-β3) were not induced following treatment with CUE-401 (i.e. IL-2 and TGF-β3). Furthermore, a subset of transcripts known to be regulated by TGF-β, including *Alpk1*, *Amot1*, *Cyp4f18*, *Lgmn*, and *Trim47*, were specifically induced in Tregs treated with CUE-401 and not with CUE-401ΔTGF-β3. Together, these findings demonstrate that co-delivery of IL-2 and TGF-β3 by CUE-401 promotes a Treg-specific transcriptional state that is distinct from, and not simply an amplification of, the program induced by IL-2 alone.

Conversely, within the CD4⁺Foxp3[−] Tconv population, treatment with CUE-401ΔTGF-β3 induced a transcriptional program characteristic of IL-2–driven cellular activation, including *Ccl4, Eomes, Fut7, GzmA, Gzmb, Lgals1, Lgals3, Ly6a, Morrbid, Nfil3, S100a6*, which were not increased in Tconv of CUE-401–treated mice (Fig. 5H). Together, these data show that the IL-2 component of CUE-401 can activate Tconv, but that this IL-2 induced Tconv activation does not occur following co-delivery with TGF-β3.

Thus, CUE-401 elicits distinct responses in Tregs compared to IL-2 alone and exhibits greater selectivity

for Treg responses compared to Tconv.

Early administration of CUE-401 ameliorates autoimmune gastritis.

To evaluate the therapeutic effects of CUE-401 in autoimmunity, we used a well-established mouse model

of Treg-deficiency–driven autoimmune gastritis, in which $CD25^+$ cells (Tregs) are depleted from BALB/c

splenocytes and transferred into athymic BALB/c *nu/nu* recipients (*1*). Severe Treg loss in the transferred

population reliably induces autoimmune gastritis in 100% of recipients. To facilitate tracking of antigen-

specific autoreactive Tconv cells in this system, a small number of $CD4^+$ TCR-transgenic Tconv cells

specific for the gastric autoantigen H^+/K^+ ATPase (*37*) and expressing the Thy1.1 congenic marker were

mixed with CD25-depleted BALB/c splenocytes $(Thy1.2^+)$ prior to transfer. Figure 6A shows a

representative flow cytometry plot of CD4-gated T cells before transfer, illustrating the CD25-depleted

BALB/c splenocytes $(Thy1.2^+)$ mixed with autoreactive transgenic T cells $(Thy1.1^+)$.

Recipient BALB/c *nu/nu* mice were dosed on days 1 and 14 with PBS or CUE-401. At day 62, the frequency

of autoreactive H^+/K^+ ATPase-specific $CD4^+Thy1.1^+$ T cells infiltrating the gastric mucosa was

significantly lower in CUE-401–treated mice compared to PBS treated mice (approximately 10% versus

44%; $P = 0.0005$; Fig. 6B, C). Stomachs were also processed for histopathological analysis to assess disease

severity, including inflammation, glandular atrophy, and hyperplasia, three hallmark features of

autoimmune gastritis (*38*). As expected, control mice developed moderate-to-severe autoimmune gastritis,

with a median disease score of 6 out of 12. In contrast, disease severity was significantly reduced in CUE-

401–treated mice, with a median pathology score of 3 out of 12, with three mice exhibiting no detectable

disease (Fig. 6D, E). Together, these results demonstrate that early and transient administration of CUE-

401 conferred durable suppression of auto-reactive T cells, as well as protection against autoimmune

gastritis–associated pathology in a well-established model of Treg deficiency–driven autoimmunity.

CUE-401 treatment induces a distinct regulatory T cell phenotype in a disease setting

To investigate the mechanism(s) by which the administration of CUE-401 suppressed autoimmune gastritis in this model we analyzed the immune cells in PBS and CUE-401 mice using flow cytometry and single cell RNA sequencing on cells isolated from gastric lymph nodes and spleens seven days after cell transfer (six days after a single administration of PBS or CUE-401). Treg frequency in stomach draining lymph nodes was significantly increased in CUE-401–treated mice (~25%) compared with PBS-treated controls (~10%) (Fig. 7A, B). To further compare Treg phenotypes, we gated on CD4$^+$Foxp3$^+$ Tregs and performed high-dimensional flow cytometry analysis, applying unsupervised dimensionality reduction with Uniform Manifold Approximation and Projection (UMAP) of expression of Foxp3, GITR, CD25, CTLA-4, CD103, CD73, CD38, CD134, CD137, and Ki67. UMAP projections revealed distinct clustering of Tregs from PBS versus CUE-401 treated mice, indicating differential protein expression profiles between treatment groups (Fig. 7C). Among these, CD25 expression was higher on Tregs from CUE-401 treated mice compared with PBS controls (Fig. 7D). These data demonstrate that CUE-401 both increases the frequency of Tregs and enhances expression of activation-associated receptors such as CD25 in this disease model.

To further define the effects of CUE-401 on the Treg phenotype and transcriptional state, we performed single-cell RNA sequencing on cells isolated from gastric lymph nodes and spleens seven days after cell transfer (six days after a single administration of PBS or CUE-401). Foxp3$^+$ Tregs formed a unique UMAP cluster in each treatment condition (Fig. 7F). Foxp3$^+$ cells were isolated for downstream analysis, and gene set enrichment revealed that Tregs from CUE-401–treated mice were significantly enriched for higher expression of IL-2/STAT5 signaling signature transcripts compared with Tregs from PBS-treated mice (Fig. 7G).

Consistent with this observation, a heatmap highlighting differentially expressed genes demonstrated increased expression of transcripts associated with Treg activation, suppressive function, and survival in CUE-401–treated mice, including *Il2ra* (CD25), *Tnfrsf4* (OX40), *Tnfrsf9* (4-1BB), *Tnfsf11* (RANKL),

Il1rl1 (ST2), and *Bcl2l1* (BCL-xL) relative to PBS-treated mice (Fig. 7H). Together, these data demonstrate

that administration of CUE-401 following transfer of Treg-depleted splenocytes increases Treg abundance

and induces a distinct transcriptional program in Tregs that is associated with activation and suppressive

activities, providing mechanistic insight into the early immunological effects of CUE-401 that contribute

to long-term protection from autoimmune disease.

Discussion

CUE-401 is a novel bifunctional Fc–IL-2/TGF-β3 fusion protein that enables coordinated cytokine delivery

to selectively promote Treg induction, expansion, and activation. In vitro, CUE-401 efficiently generated

induced Tregs (iTregs), expanded endogenous Tregs, and enhanced Foxp3 expression while suppressing

cytokine production in effector and memory T cells, which are effects consistent with integrated IL-2 and

TGF-β3 signaling. In vivo, CUE-401 increased Treg frequency and activation while limiting Tconv

activation relative to IL-2 alone, and ameliorated disease in a Treg-deficiency–driven gastritis model.

Transcriptomic analyses further indicate that CUE-401 elicits a Treg program that is qualitatively distinct

from that induced by IL-2 monotherapy, and while IL-2 alone activates Tconv, the IL-2 and TGF-β3

together in CUE-401 did not activate Tconv, speaking to an opportunity for enhanced Treg selectivity with

this approach. While additional studies will be required to more fully assess these responses in additional

chronic disease settings, the current data supports robust short-term Treg reprogramming in vivo and long-

term prevention of autoimmunity.

Prior studies established that iTregs generated ex vivo with IL-2 and TGF-β can suppress autoimmunity

following adoptive transfer (*33-36, 39, 40*). However, these approaches required cell manufacturing and

did not directly address whether coordinated cytokine delivery can be achieved pharmacologically in vivo.

Our findings demonstrate that CUE-401 enables such coordination, promoting Treg expansion and

activation without broadly stimulating conventional T cells. Notably, CUE-401 induced an activated Treg

phenotype while minimizing activation of Foxp3⁻ Tconv, an effect not observed with an IL-2–only variant,

supporting the premise that linked IL-2 and TGF-β signaling preferentially biases responses toward the

Treg lineage. Future studies dissecting receptor-proximal signaling dynamics and cell-type–specific

cytokine engagement will further refine this model (*41*).

At the molecular level, Tregs treated with CUE-401 exhibited durable Foxp3 expression, demethylation of

the Foxp3 TSDR, and sustained Foxp3 maintenance upon restimulation, features associated with lineage

stability. These properties are particularly important given the reported instability of some iTreg

populations under inflammatory conditions (*42-45*). Although overall Treg frequencies normalized to

baseline levels after ~14 days (data not shown), CUE-401 expanded cells maintained a stable Foxp3$^+$

phenotype in both homeostatic and inflammatory settings, consistent with physiological regulation of the

compartment rather than loss of lineage identity. Longer-term lineage-tracing studies will be valuable to

definitively distinguish de novo induction from selective expansion in vivo.

Mechanistically, CUE-401 drove coordinated activation of CD25/IL-2/STAT5 pathways while avoiding

induction of effector-like programs. Studies with a TGF-β–deficient variant (CUEΔTGF-β3) established

that TGF-β3 is required for selective Treg activation and for constraining off-target CD4$^+$ Tconv responses,

supporting a TGF-β–dependent mechanism of action. Together, these data indicate that physically linked

cytokine co-delivery can reshape IL-2 responsiveness toward a regulatory-biased state. Whether this

reflects altered cytokine receptor geometry, local concentration effects, or cooperative signaling remains

an important question for future mechanistic studies.

Collectively, our findings position CUE-401 as a modular strategy for in vivo Treg programming that may

overcome key limitations of current IL-2–based therapies. By promoting expansion and activation of stable,

lineage-committed Tregs while minimizing effector T cell activation, CUE-401 represents a promising

therapeutic approach for autoimmune and inflammatory diseases. Evaluation in additional disease models

and human systems will be important next steps toward clinical translation.

Materials and Methods

Study design

The objective of this research was to define the activities of a new biologic that co-engages IL-2 and TGF-β pathways on regulatory T cells. This was accomplished by experimentally testing the function of the biologic, named CUE-401, in vitro and in vivo. In vitro, testing was performed to address the induction of iTregs from naïve $CD4^+$ T cells, maintenance of Foxp3 expression and expansion of existing Tregs, and induction of Foxp3 expression with reduced proinflammatory cytokine production in a subset of effector/memory $CD4^+$ T cells. CUE-401 was also administered to mice. Expression of Foxp3 and other activation related proteins were measured by flow cytometry and compared by uniform manifold approximation and projection analysis. Stability of in vivo Tregs was assessed by measuring the methylation state of CpG sites in the Treg specific demethylated region, and $Foxp3^+$ cells were restimulated to verify maintenance of Foxp3 expression. CUE-401's ability to affect disease severity was measured through tracking autoreactive T cells, pathological scores of disease in the stomachs of an established mouse model of autoimmunity. Single cell RNA sequencing was done to address potential mechanisms of action for disease.

CUE-401 design and production

CUE-401 was produced by transient transfection of ExpiCHO-S cells using the Gibco ExpiCHO Expression System Max Titer Protocol. Each transfection is performed with two plasmids at a 1:1 molar ratio encoding (1) IL2 mutein linked to human IgG1Fc (knob) linked to monomeric TGFß3 and (2) human IgG1Fc (hole) linked to truncated TGFßR2, respectively. Upon harvest, conditioned media was loaded onto a mAb Select SuRe protein A column using an AKTA Pure 25 system for affinity-based purification. After a 10-column volume (cv) wash with 20mM HEPES pH 7.5 + 1M NaCl, CUE-401 was eluted using 50mM glycine pH 2.8 directly into 10% 1M Tris pH 9.0 neutralization solution. Eluant was pooled, concentrated and loaded

onto a HiLoad 26/600 Superdex 200 for final step purification by size exclusion chromatography in 1x PBS

pH 7.4. Main protein elution peak was collected, pooled, concentrated, filtered using a 2um syringe filter.

Protein concentration was measured by absorbance at 280nm (A_{280}) using a NanoDrop system. Protein

integrity, size, solution state, and purity were confirmed by liquid chromatography-mass spectrometry,

SDS-PAGE, and analytical size exclusion chromatography. Final purified protein was aliquoted and stored

at -80°C until use.

Mice

Female BALB/c Thy1.1+ mice were bred in house and BALB/c *nu/nu* (4–8 wk old) were purchased from

the Charles River Laboratory and housed under specific pathogen-free conditions. TCR transgenic mice

were crossed with BALB/c Thy1.1/1.1 expressing mice as previously described (*37*). TCR Transgenic

Thy1.1/1.1 expressing mice that develop autoimmune gastritis were crossed with BALB/c Thy1.1/1.1

Foxp3.eGFP reporter mice (C.Cg-*Foxp3^{tm2Tch}*/J) to generate TCR transgenic Thy1.1/1.1 Foxp3.eGFP

reporter mice (*42*). All animals were maintained in the Department of Comparative Medicine, Saint Louis

University School of Medicine in accordance with institutional guidelines (Association for Assessment and

Accreditation of Laboratory Animal Care, International accreditation number 000656).

CUE-401 Dosing

329 CUE-401 was formulated in sterile PBS, pH 7.4. Mice received intraperitoneal (i.p.) injections of CUE-401

330 at 2 mg/kg (100 μg/mL; 10 mL/kg; typical injection volume ~0.2–0.3 mL per 20–30 g mouse) or PBS

331 (vehicle) on the indicated schedule. For the adoptive transfer model, dosing occurred on days +1 and +14

332 post-transfer. For pharmacodynamic studies of Treg expansion, a single dose was administered, and tissues

333 were analyzed on day 6.

CTLL-2 proliferation assays

CTLL-2 cells were cultured in RPMI 1640 medium (Thermo Fisher Scientific, Waltham, MA, USA)

supplemented with 10% heat-inactivated fetal bovine serum (Thermo Fisher Scientific, Waltham, MA,

USA) and were starved of IL-2 for 22–24 h by washing cells free of cytokine and incubating them at

1×10^5 cells/mL in IL-2–free medium at 37°C with 5% CO_2. Cell viability and morphology were assessed

before and after starvation via acridine orange/propidium iodide staining using a Cellaca MX automated

cell counter (Nexcelom Bioscience, Lawrence, MA, USA). On the day of assay, 5×10^3 viable CTLL-2

cells were seeded in 96-well tissue-culture-treated plates (Corning/Costar, Corning, NY, USA) in 100 μL

medium, with outer wells filled with DPBS (Thermo Fisher Scientific, Waltham, MA, USA) to minimize

edge effects. Test molecules and recombinant rhIL-2 were prepared at 2× concentration and added in 100

μL to achieve a final volume of 200 μL per well, with all conditions plated in triplicate. After 72 h

incubation at 37°C, cell viability was quantified using CellTiter-Glo® reagent (Promega, Madison, WI,

USA): plates were equilibrated to room temperature, 100 μL of suspension was transferred to white

flat-bottom 96-well assay plates (Corning/Costar, Corning, NY, USA), an equal volume of CellTiter-Glo

was added via reverse pipetting, plates were agitated and incubated for 10 min, and luminescence was

measured using a Synergy Neo2 multimode plate reader (BioTek Instruments, Winooski, VT, USA).

pSMAD Detection

Murine splenocytes and human PBMCs were used to evaluate pSMAD2 signaling in response to

CUE-401 stimulation and TGF-β pathway inhibition. Fresh spleens from BALB/c mice were processed

through a 0.7-μm strainer to obtain single-cell suspensions, followed by ACK lysis, washing in PBS, and

counting on a Cellaca MX automated cell counter. Cells were aliquoted at 5×10^7 per condition and rested

for 2 h at 37°C, with subsets pre-treated for 90 min with 10 μM TGF-β inhibitor. PBMCs were thawed

and washed similarly. Cells were treated for 30 min at 37°C with IL-2, TGF-β3, CUE-401, or

combinations thereof, and then washed and lysed with RIPA buffer (Thermo Fisher Scientific, Waltham,

MA, USA) supplemented with Halt Protease/Phosphatase Inhibitor Cocktail (Thermo Fisher Scientific,

Waltham, MA, USA). Lysates were homogenized by pipetting, clarified by centrifugation at 10,000 rpm

for 10 min at 4°C, and protein content was quantified using a BCA assay. Samples were mixed with LDS

loading buffer (Thermo Fisher Scientific, Waltham, MA, USA), heated, and resolved on Bolt 4–12%

Bis-Tris Plus polyacrylamide gels (Thermo Fisher Scientific, Waltham, MA, USA) alongside SeeBlue

Plus Prestained Standards (Thermo Fisher Scientific, Waltham, MA, USA). Following electrophoresis,

proteins were transferred to PVDF membranes using the iBlot 2 Dry Blotting System (Thermo Fisher

Scientific, Waltham, MA, USA), then blocked for 2 h at room temperature in SuperBlock buffer (Thermo

Fisher Scientific, Waltham, MA, USA). Membranes were incubated overnight at 4°C with primary

monoclonal antibodies against phospho-SMAD2 (Cell Signaling Technology, Danvers, MA, USA) or

total SMAD2/3 (Cell Signaling Technology, Danvers, MA, USA). After washing, membranes were

incubated with HRP-conjugated goat anti-rabbit IgG (Cell Signaling Technology, Danvers, MA, USA)

for 1 h at room temperature, washed again, and developed using SuperSignal West Femto or West Pico

chemiluminescent substrates (Thermo Fisher Scientific, Waltham, MA, USA). Western blots were

imaged using a Syngene G:BOX darkroom imaging system (Syngene, Frederick, MD, USA). Band

intensities were analyzed to quantify pSMAD2 induction and inhibition across treatment conditions.

Isolation of Immune Cells

Isolation of immune cells from stomach tissue has been described previously (*38, 43*). Briefly, stomachs

were harvested from mice, gastric lymph nodes were removed, and the stomachs were dissected along the

lesser curvature. Stomachs were rinsed with cold PBS then injected with 10 mL of cold MACS buffer (1 x

PBS,0.5% BSA (Fisher Scientific Waltham, MA), 2 mM EDTA (Promega, Fitchburg, Wisconsin)) using a

27-gauge needle, to inflate the tissue and free infiltrating immune cells. After injection, the stomach was

diced and collected with the 10mL of MACS buffer and transferred to a 50 mL conical tube. The dispersed

stomach tissue was mixed thoroughly then passed through a 40-micron filter, washed with 40 mL of MACS

buffer and used for analysis. Immune cells from the spleen and gastric/pancreatic lymph node were

processed by dispersing on a 40-micron filter and washed with 15 mL of MACS buffer. Spleen cells were

spun down at 2000 RPM for 5 minutes after initial processing. Spleen cells were resuspended in 5 mL of

1x ACK lysis buffer (Thermo Fisher Scientific, Waltham, MA) for 10 seconds then washed with 45 mL of

MACS buffer and spun again at 2000 RPM for 5 minutes. Cells were resuspended and filtered again through

a 40-micron filter before being counted and utilized for experiments.

T cell purification, iTreg differentiation, and nTreg expansion

Inducible Tregs (iTreg) were generated as previously described (*9*). Briefly, CD4$^+$CD8$^-$ thymocytes or

CD4$^+$Foxp3.eGFP$^+$ splenocytes were sorted from 6-8-wk-old BALB/c mice and stimulated with plate-

bound anti-CD3 (1 mg/ml) and anti-CD28 (2 mg/ml; BD Biosciences) in 24-well plates (2.5x10^5 cells/well)

in 2 ml RPMI media (modified RPMI 1640 (Corning, Tewksbury, Massachusetts) supplemented with 10%

fetal bovine serum (FBS; Biotechne, Minneapolis, Minnesota), 2 mM glutamine (Sigma-Aldrich), 100 U/ml

penicillin,100 mg/ml streptomycin (Sigma-Aldrich), 10 mM HEPES (Sigma-Aldrich), β-mercaptoethanol

(55mM), 1 mM essential amino acids (Gibco, Billings, Montana), and 1 mM sodium pyruvate (Sigma-

Aldrich) supplemented with recombinant human IL-2(rhIL-2) (National Institutes of Health, Bethesda,

Maryland) (100 U/ml). Recombinant human TGF-β3 (rhTGF-β3) (Biotechne) (5 ng/ml) was added to

induce iTregs. Cells were removed from plate-bound antibodies after 48 hours and added to new wells with

fresh media supplemented with 100 U/ml recombinant human IL-2. Cells were used after 7 days in culture.

Endogenous Treg expansion was assessed by isolating FACS-purified splenic CD4$^+$Foxp3.eGFP$^+$ Tregs.

Tregs were stimulated with plate-bound anti-CD3 & anti-CD28 and expanded with IL-2 (100 U/mL) or

CUE-401 as indicated for 5–7 days. Foxp3 maintenance and cell counts were assessed by flow cytometry.

Effector/Memory CD4$^+$ T-Cell Restimulation and Cytokine Quantification

Splenic CD4$^+$CD44^{hi} Foxp3.eGFP$^-$ T cells were FACS-sorted and stimulated with plate-bound anti-CD3 &

anti-CD28 in the presence of rhIL-2 (100 U/mL), rhIL-2 + rhTGF-β3 (5 ng/mL), or CUE-401. For cytokine

secretion, cells were cultured at 1×10^6 cells/mL and restimulated with 25ng/mL phorbol myristate acetate

(PMA) (Sigma-Aldrich, St. Louis, Missouri), and 500 ng/mL ionomycin (Sigma-Aldrich), for 72 hours at

37°C for 72 h. Supernatants were collected and cytokine secretion was analyzed using a Th1, Th2, Th17

and IL-13 flex set (BD Biosciences) cytometric bead array kit and analyzed by flow cytometry following

the kit protocol and analyzed by flow cytometry.

Treg Stability: TSDR (CNS2) Methylation Analysis

$CD4^+Foxp3^+$ and $CD4^+Foxp3^-$ cells were sorted from spleens of PBS- or CUE-401-treated mice 6 days

post-dose. Genomic DNA was isolated (DNeasy Blood & Tissue Kit, Qiagen), bisulfite-converted (EZ

DNA Methylation-Gold Kit, Zymo Research), and the Foxp3 TSDR (CNS2) region was PCR-amplified

using validated primers spanning CpG sites within the enhancer. Amplicons were analyzed by

pyrosequencing [or Sanger sequencing of cloned products] to quantify methylation at individual CpGs.

Percent demethylation was calculated per CpG and averaged across the TSDR. Males and female mice

were used for TSDR analyses. Methylation state was normalized in female mice to account for the X-linked

Foxp3 locus inactivation in females.

Bulk RNA-seq (Tregs and Tconv)

$CD4^+Foxp3.eGFP^+$ regulatory T cells (Tregs) and $CD4^+Foxp3.eGFP^-$ conventional T cells (Tconv) were

FACS-sorted from spleens of PBS-, CUE-401–, or CUE-401ΔTGF-β3–treated BALB/c mice 6 days after

treatment (≥95% purity). Sorted cells were pelleted and immediately lysed in lysis buffer supplied with the

PureLink™ RNA Micro Kit (Invitrogen, Thermo Fisher Scientific), and total RNA was extracted according

to the manufacturer's protocol. RNA was eluted in RNase-free water and quantified using a Qubit RNA HS

Assay (Thermo Fisher). RNA integrity was assessed using the Agilent Bioanalyzer RNA Pico Kit, and only

samples with RIN ≥7.0 were used for library preparation.

Libraries were generated using the Takara-Clontech SMARTer construction method for Illumina with

poly-A mRNA enrichment, following manufacturer instructions. Indexed libraries were pooled and

sequenced on an Illumina NovaSeq x Pluse (300 cycles) platform (paired-end 2×100 bp), targeting ≥30

million read pairs per sample.

Samples were processed and sequenced at the Genome Technology Access Center @ McDonnell Genome

Institute in St. Louis, Missouri. RNA-seq reads were then aligned to the Ensembl GRCm39.113 primary

assembly with STAR version 2.7.11b. Gene counts were derived from the number of uniquely aligned

fragments by Subread:featureCount version 2.0.8. Differential gene expression analysis was performed

with EdgeR using quasi-likelihood F-test and was adjusted using the Benjamini-Hochberg method. Genes

with a false discovery rate (FDR) < 0.05 were considered significantly differentially expressed and adjusted

P value was $p < 0.05$. Level counts were obtained with featureCount. Visualization of the heatmaps was

generated using the R package pheatmap. Level counts were obtained with featureCounts. Differential gene

expression analyses were performed using DESeq2. Significance thresholds were set at an adjusted

449   Cell purification and injection for adoptive transfer model of autoimmune gastritis

Naive TxA23 T cells were isolated from a single-cell suspension of thymii from TxA23.Thy-1.1

Foxp3.eGFP mice. Thymocytes were lysed with ACK lysis buffer (BioSource) and washed with MACS

buffer prior to staining with CD4 BV421 (BD Biosciences) and CD8 PE (BD Biosciences). Cells were

placed in FACS sort buffer (2% FBS (Biotechne, Minneapolis, Minnesota), 100 U/ml penicillin, 100 mg/ml

streptomycin, in complete RPMI without phenol red (Sigma-Aldrich), and then purified using the BD

FACSAria Fusion cell sorter (BD Biosciences) to obtain CD4$^+$CD8$^-$ GFP$^-$ Cells. BALB/c Splenocytes were

from spleens of healthy Thy-1.2 expressing BALB/c mice, RBC were removed with ACK lysis buffer and

then purified as follows. To deplete CD25$^+$ cells, cells were incubated for 10 min with anti-CD25-PE

(PC61), washed in MACS buffer, and incubated for an additional 10 min with anti-PE MACS microbeads

(Miltenyi Biotec, Bergisch Gladbach) then subjected to autoMACS purification (Miltenyi Biotec). The

negative fraction was used as CD25-depleted spleen cells. A typical depletion resulted in <1% of the CD4$^+$

cells expressing CD25. Prior to injection, all cells were washed twice in PBS. Naive TxA23.Thy-1.1 T cells

that were isolated from thymus, were mixed into the BALB/c Thy1.2 splenocyte population such that an

i.p. injection of 0.3 ml/mouse resulted in the transfer of 100,000 TxA23.Thy-1.1 T cells and 20×10^6 spleen

cells. Mice were monitored weekly and euthanized at ~ day 62 for tissue analyses.

Histopathology

BALB/c *nu/nu* mice were euthanized 62 days after receiving an adoptive transfer of splenocytes and naïve CD4$^+$ T cells from autoreactive mice. The stomach opened, washed with cold PBS, and fixed in 4% neutral buffered formalin. Fixed tissues were embedded in paraffin, stained with hematoxylin and eosin (H&E) by Saint Louis University Research Histology Facility (SLU ASBRH) and disease pathology was scored similarly to previously published assessments (*38*). Briefly multiple pathological features, including inflammation, glandular atrophy, and epithelial hyperplasia were graded on a scale of 1–4, depending on the extent of mononuclear cell infiltration and parietal and chief cell destruction. All stomach sections were scored blindly by at least two individuals. General descriptions for scores are as follows: 0, healthy tissue with no abnormalities 0.5, scattered lymphocytes throughout submucosa and muscularis with one or two small dense blankets of lymphocytes; 1, two to four small dense clusters of lymphocytes in the submucosa/mucosa; 1.5, two or three areas with intermediate infiltration spanning one-third of mucosa; 2, big nodules of lymphocytic accumulation spanning half to all of mucosa; 2.5, big nodules of lymphocytic accumulation spanning half to all of mucosa beginning to show evidence of parietal and chief cell destruction (<25%); 3.0, heavy lymphocytic infiltration throughout mucosa, parietal and chief cell loss (25–75%), and replacement by foamy cells; and 4.0, total parietal and chief cell loss, no mucosal architecture, and many foamy cells.

Flow Cytometry

Immune cells were isolated from gastric mucosa, gastric/pancreatic or inguinal lymph nodes, and spleens following the protocol previously described and cell surface staining was performed according to standard procedures (*38, 43*). Immune cell subsets and proteins expressed on Tregs were identified using the following antibodies: CD4 BV421, CD8 BV510, CD90.2 BV786, CD4 BV711, CD38 BV711, CD137 BV650 CD152 (CTLA4), CD103 PE-Cy7, CD8 PE, CD19 PE (BD Biosciences, Franklin Lakes, New Jersey), CD73 PE/Dazzle594, CD90.1, CD134 PerCP/Cyanine 5.5, Ki67 BV605 (BioLegend, San Diego, California), CD25 APC-eFluor780, Foxp3 APC, CD62L eFlour450, CD44 APC-eFlour780 (Thermo

Fisher Scientific). Cells were washed with 200 uL of PBS and stained in only PBS at 4°C. Cells were

fixed and permeabilized using eBioscience Foxp3/Transcription Factor Staining Buffer Set (Thermo

Fisher Scientific). Cells were incubated overnight with the intracellular antibodies and then washed and

analyzed by flow cytometry. Flow cytometry was performed on a BD LSRII, a BD Fortessa (BD

Biosciences), or a Cytek Aurora (Cytek Biosciences, Fremont California), and analyzed using FlowJo

(FlowJo, Ashland, OR).

Single-Cell RNA-seq

Cells from gastric lymph nodes and spleens (day 7 post transfer; day 6 post-treatment) were prepared as

above, RBC depleted as needed and filtered through 40μm strainers. Viability (>85%) was confirmed by

trypan blue. Cells were loaded on a 10x Genomics Chromium Controller (10X Genomics, Pleasanton,

CA) targeting up to 10,000 cells per library using the Next GEM Single Cell 5′ Kit v2 (10x #1000263).

cDNA amplification and library prep followed manufacturer protocols. PCR reactions were conducted in

the T100 Thermal Cycler (BioRad, Hercules, CA). Libraries were QC'd on an Agilent 2100 Bioanalyzer

(Agilent, Santa Clara, CA) and sequenced on an Illumina NovaSeq 6000 (Illumina, San Diego, CA).

Raw data were processed through CellRanger 6.0.2 pipeline (10x Genomics). Conditional repeats were

combined using cellranger aggr. Analysis was conducted using Seurat. (*44*) Each dataset was filtered to

exclude cells with low gene expression (<500/cell), high gene counts (outliers), and high percent

mitochondrial transcript load (>10%/cell). Datasets were merged, normalized, scaled, and dimensionally

reduced. Layers were joined and clusters were generated using a resolution of 2. Cluster identities were

established by attaining distinguishing genes per cluster and relating them to known profiles for immune

cells present in spleen and gastric lymph nodes. T cells clusters were subset and pathway analysis was

conducted using the R package enrichR and the Msig database for transcripts. Gene ontology, pathway

enrichment, and signature analyses (e.g., IL2/STAT5 signaling) were conducted

using fgsea and enrichR with MSigDB gene sets. Visualization of feature plots was performed using the

ggplot2 package in combination with the scCustomize package. Heatmaps were generated using Seurat's

DoHeatMap function. Statistical analyses were performed in Prism 10.0.0 (GraphPad Software). *P* values

less than 0.05 were considered significant.

Statistical Analysis

In vitro data are expressed as means of individual determination ± SD. In vivo data are expressed as median

of individual determination ± 95% confidence interval. Statistical analysis was performed using GraphPad

Prism 10 (GraphPad Software, San Diego, CA). Dose titrations of CUE-401 on nTregs utilized Kruskal-

Wallis test. Cytokine analysis utilized Wilcoxon matched-pairs signed rank test. In vitro treatment

comparisons between rhIL-2, rhIL-2 + TGF-β, and CUE-401 utilized Ordinary one-way ANOVA with

Tukey's multiple comparisons test. Histopathology score significance was performed using the Mann–

Whitney *U* test. Differential expression analysis between CUE-401 and PBS treated Tregs in scRNA-seq

data was performed using the Wilcoxon rank-sum test implemented in Seurat. Differentially expressed

genes were identified using the FindMarkers function in the Seurat package with default statistical

parameters, a log fold-change threshold of 0.6, and a minimum expression fraction of 0.1. Genes with an

adjusted p-value < 0.05 were considered differentially expressed. For bulk RNA sequencing, P-values were

adjusted for multiple comparisons using the Benjamini–Hochberg method to control false discovery rate.

Figures:

Fig. 1. CUE-401 promotes IL-2 and TGF-β signaling activity.

(**A**) Model of CUE-401 design. (**B**) CTLL-2 proliferation with recombinant human IL-2 (Proleukin),

CUE-401, or CUE-401ΔTGF-β3. (**C**) Immunoblot detecting phosphorylated SMAD-2 (pSMAD-2) and

 total SMAD-2/3 after 30 minutes of stimulation with rhIL-2, rhTGF-β3, CUE-401, or CUE-401 +

SB43152.

Fig. 2. CUE-401 promotes iTreg induction, nTreg expansion, and Teff suppression

(**A**). Representative flow cytometric analysis showing the percentage of CD4$^+$ gated cells expressing

Foxp3 in vitro following stimulation with rhIL-2, rhIL-2 + rhTGF-β3, or CUE-401. (**B**). Dose-response

curve displaying Foxp3 expression with increasing doses of CUE-401. (**C**). Representative flow

cytometric analysis showing the percentage of CD4$^+$ gated cells expressing Foxp3 after six days in vitro

stimulation of Tregs with rhIL-2 or CUE-401. (**D**). Number of Foxp3 expressing cells at various

concentrations of CUE-401. (**E**). Flow cytometric analysis displaying the percentage of CD4$^+$ gated cells

expressing Foxp3 after six days in vitro simulation of Teff with rhIL-2, rhIL-2 + rhTGF-β3, or CUE-401.

(**F**) Percentages of CD4$^+$Foxp3$^+$ Tregs in rhIL-2, rhIL-2 + rhTGF-β3 and CUE-401 treated samples. (**G**).

Amounts of IFN-γ, IL-4, TNF, and IL-17A secreted from CD4$^+$CD44^{hi} Foxp3.GFP$^-$ T cells with rhIL-2 or

CUE-401 treatment. Flow cytometry plots are representative of individual sorted samples analyzed from

thymus (A) or spleen (D,F,G) of 4-8 healthy BALB/c mice. EC$_{50}$ calculated by taking the concentration

halfway between the top and bottom value (approximately 20-80%). Bars in each graph represent the

mean +/-SD. *P*-values calculated using Kruskal-Wallis test (D), ordinary one-way ANOVA with Tukey's

multiple comparisons test (F), or Wilcoxon matched-pairs signed rank test (G) with $*p = 0.0332$, $**p =$

0.0021, $***p = 0.0002$, and $****p < 0.0001$.

Fig. 3. CUE-401 treatment increases proportions of stable Foxp3+ Tregs in vivo

(**A**) Flow cytometric analysis showing percentages of CD4$^+$ gated cells expressing Foxp3 six days after

treatment of BALB/c mice. (**B**) Combined data for the percentages of CD4$^+$Foxp3$^+$ Tregs in PBS and

CUE-401 treated mice. (**C**) Flow cytometric analysis of CD4$^+$Foxp3$^+$ expression prior to and after

performing FACS to purify CD4$^+$Foxp3$^+$GFP$^+$ Tregs. (**D**) Results of qPCR quantifying the relative

percentage of DNA demethylation at key CpG sites within the Treg-specific demethylated region (TSDR)

of the *Foxp3* locus of CD4$^+$Foxp3.eGFP$^-$ and CD4$^+$Foxp3.eGFP$^+$ cells sorted from PBS and CUE-401

treated mice. (**E**) Representative flow cytometric plots showing Foxp3 expression of purified T cells from

PBS and CUE-401 treated mice after 6 days of restimulation in vitro in the absence of cytokines. For all

graphs, controls were PBS treated age-matched mice. Data in (A-B) was generated from spleens of 2-5

mice in 7 independent experiments. Plots in (C) show the average proportion of Foxp3$^+$ T cells pre and

post sorting from 8 mice in 2 independent experiments. Bars represent the median +/- 95% C.I. TSDR

analysis in (D) includes 12 mice from 2 independent experiments. Female mice methylation status was

normalized for methylation state on both x chromosomes. *P* values were calculated utilizing the Mann-

Whitney *U* test. ****$p < 0.001$.

Fig. 4. CUE-401 increases expression of transcripts and proteins associated with Treg activation.

(**A**). Flow cytometric analysis shows the percentage of CD4$^+$Foxp3$^+$ expressing cells following treatment

of mice with PBS or CUE-401. CD4$^+$Foxp3$^+$ Tregs were identified, and UMAP projections were

generated using OMIQ software to cluster cells based on protein expression. The feature plot shows

CD25 expression within the Foxp3$^+$ populations. (**B**). Cells were gated on the CD4$^+$Foxp3$^+$ populations to

assess surface expression of GITR, CD25, CTLA-4 and CD103. (**C**). Histograms display the expression

of GITR, CD25, CTLA-4, and CD103 within the Treg compartment of PBS-treated controls compared to

CUE-401–treated mice. (**D**). Heatmap from bulk RNA sequencing of sorted Tregs, highlighting

differentially expressed genes between PBS and CUE-401 treated mice. Genes included had a log$_2$ fold

change > 2 and adjusted p-value < 0.05, using the Benjamini–Hochberg correction to control false

discovery rate (FDR). For all flow cytometry plots, data are representative of individual spleens from a

single experiment with 3 mice per treatment group. PBS-treated, age-matched mice served as controls in

all analyses.

Fig. 5. The combined effects of IL-2 and TGF-β3 in CUE-401 differ from IL-2 treatment alone.

(**A**). Representative flow cytometric analysis showing the percentage of CD4$^+$ gated cells expressing

Foxp3 six days after treatment with PBS (left), CUE-401ΔTGF-β3 (middle), or CUE-401 (right) in vivo.

(**B**). Combined data showing the percentages of CD4$^+$Foxp3$^+$ Tregs in PBS, CUE-401ΔTGF-β3, or CUE-

401 treated mice. (**C**). Representative flow cytometric analysis comparing GITR by CD38 expression in

CD4$^+$Foxp3$^+$ Tregs following the indicated treatments. (**D**). Combined data showing the percentage of

GITR$^+$CD38$^+$ Tregs following PBS, CUE-401ΔTGF-β3, or CUE-401 treatment. (**E**). Representative flow

cytometric analysis comparing CD25 by CD103 of CD4$^+$Foxp3$^-$ gated cells. (**F**). Combined data showing

the percentage of CD25$^+$CD4$^+$Foxp3$^-$ cells and CD103$^+$CD4$^+$Foxp3$^-$ cells six days after treatment. Data

shown are representative of two independent experiments (N = 9 per group). Bars represent the median ±

95% confidence interval. (**G**, **H**). Heatmaps of gene expression profiles in sorted CD4$^+$Foxp3.eGFP$^+$

Treg cells or CD4$^+$Foxp3.eGFP$^-$ Tconv cells, derived from a subset of control or treated mice (N = 2,3 per

group, respectively). *P* values for flow results were determined utilizing an ordinary one-way ANOVA

with Tukey's multiple comparisons test with *p = 0.0332, **p = 0.0021, *** *p* = 0.0002, and ****p <

0.0001.

Fig. 6. CUE-401 treatment inhibits Teff responses and decreases autoimmune disease severity.

(**A**). Flow cytometric analysis of Thy1.2$^+$CD4$^+$ splenocytes and autoreactive Thy1.1$^+$CD4$^+$ splenocytes

prior to injection into BALB/c *nu/nu* mice. (**B**). Flow cytometric analysis of Thy1.1 and Thy1.2

expression in CD4$^+$ T cells from stomachs of BALB/c *nu/nu* mice ~62 days after transfer. (**C**) Combined

analysis of Thy1.1 expressing autoreactive cells in the gastric mucosa of nude mice ~62 days after

transfer and subsequent treatment with PBS or CUE-401. Quantified data are presented as mean ± SEM.

Experiments were conducted independently three times, with 2–5 mice per group per experiment. (**D**).

Representative images from four independent experiments, comparing tissues from healthy untreated

BALB/c mice, PBS-treated BALB/c *nu/nu* controls, and CUE-401–treated BALB/c *nu/nu* mice. (**E**).

Histological disease scores were determined based on multiple pathological features, including

inflammation, glandular atrophy, and epithelial hyperplasia. The final score represents the combined

values from these categories. Graph bars represent the median +/- 95% C.I. Experiments contained 3-5

aged matched female mice. *P*-values were determined utilizing either Unpaired Student's T test (C) or

Mann-Whitney *U* test (D) with **p = 0.0040 and ***p = 0.0005. Scale bars in micrographs represent 180

μM.

Fig. 7. CUE-401 induces activated Tregs with a distinct transcriptional program.

(**A**). Representative flow cytometric analysis of CD4 gated Foxp3$^+$ cells in PBS and CUE-401 treated

BALB/c nude mice 7 days after splenocyte transfer. (**B**). Cumulative data showing the percentages of

CD4$^+$Foxp3$^+$ Tregs in PBS and CUE-401 treated mice 6 days after a single treatment. (**C**). CD4$^+$Foxp3$^+$

Tregs were identified in each treatment condition, and UMAP projections generated distinct clusters of

cells based on protein expression. (**D**). Representative flow cytometric analysis of CD4$^+$Foxp3$^+$ gated,

CD25 expressing cells. (**E**). Combined data showing the percentages of CD25 expressing Tregs in PBS

and CUE-401 treated mice. (**F**). Single cell RNA sequencing clustering analysis of T cells. T cell

populations were subset and Foxp3 expression identified the Treg populations in PBS and CUE-401

treated mice. (**G**). Foxp3 expressing T cells were re-clustered and signatures were generated based on IL-

2/STAT5 signaling transcripts that were identified through gene set enrichment analysis

(**H**). Heatmap that highlights differentially expressed genes in single Foxp3$^+$ cells between PBS and CUE-

401 treatment. Differential expression analysis between CUE-401 and PBS Tregs used the Wilcoxon

rank-sum test implemented in Seurat. P-values were adjusted for multiple comparisons using the

Benjamini–Hochberg method to control false discovery rate (FDR). Representative images in (A) from 3

independent experiments with 3-5 age matched female mice per experiment. *P*-values were determined

utilizing Mann-Whitney *U* test with $p < 0.05$ as significant for the flow cytometry analyses. Single cell

analysis in (F-H) generated cells of 8 total mice in 2 independent experiments (4 treated and 4 untreated).

Acknowledgments:

We thank Saint Louis University Flow Cytometry Core (Joy Eslick, and Jacqueline Spencer) and the

Advanced Research Histology Facility (Caroline Murphy, Vasiliki Grammatopoulou) for assistance with

data collection. We thank Saint Louis University Genomics and Bioinformatics Core Facility (Dr.

Michelle Brennan) for assistance with processing and analysis of single cell and bulk RNA sequencing.

We thank Saint Louis University Department of Comparative Medicine (Lisa Tempel, Tawanika

Williams, Kelsey Ernst) for assistance in animal care and handling

Funding:

This work was funded in part by a sponsored research agreement between CUE Biopharma and Saint

Louis University.

Author contributions:

Conceptualization: NMJ, RS, NMG, SNQ, AS, CAP, SGH, EF, RJD

Methodology: NMJ, NMG, SNQ, RJD, SL, EC, KY, EN

Investigation: NMJ

Visualization: NMJ, KF, JAC, RJD

Formal Analysis: NMJ, NY, CAP, SGH, KF, EF

Software: SGH, KF

Funding acquisition: AS, NMG, SNQ, RJD

Writing – original draft: NMJ, JAC, RJD

Writing – review & editing: NMJ, JAC, NMG, SNQ, RJD

All authors approved the final manuscript.

Competing interests: AS, SL, NMG, SNQ, RS, KY, EN, and EC are employees of CUE Biopharma. All

other authors declare that they have no competing interests.

Data and materials availability: All data necessary to understand and evaluate the conclusions of the

paper are provided in the manuscript and supplementary materials. The CUE-401 biologic can be

available upon request under a material transfer agreement with CUE Biopharma. Sequencing

data will be made publicly available upon acceptance of this publication.

References:

1. S. Sakaguchi, N. Sakaguchi, M. Asano, M. Itoh, M. Toda, Immunologic self-tolerance maintained by activated T cells expressing IL-2 receptor alpha-chains (CD25). Breakdown of a single mechanism of self-tolerance causes various autoimmune diseases. *J Immunol* **155**, 1151-1164 (1995).

2. L. Wang *et al.*, Regulatory T cells in homeostasis and disease: molecular mechanisms and therapeutic potential. *Signal Transduct Target Ther* **10**, 345 (2025).

3. C. L. Bennett *et al.*, The immune dysregulation, polyendocrinopathy, enteropathy, X-linked syndrome (IPEX) is caused by mutations of FOXP3. *Nat Genet* **27**, 20-21 (2001).

4. S. Hori, T. Nomura, S. Sakaguchi, Control of regulatory T cell development by the transcription factor Foxp3. *Science* **299**, 1057-1061 (2003).

5. R. Khattri, T. Cox, S. A. Yasayko, F. Ramsdell, An essential role for Scurfin in CD4+CD25+ T regulatory cells. *Nat Immunol* **4**, 337-342 (2003).

6. J. D. Fontenot, M. A. Gavin, A. Y. Rudensky, Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol* **4**, 330-336 (2003).

7. E. M. Shevach, A. M. Thornton, tTregs, pTregs, and iTregs: similarities and differences. *Immunol Rev* **259**, 88-102 (2014).

8. A. Jones, D. Hawiger, Peripherally Induced Regulatory T Cells: Recruited Protectors of the Central Nervous System against Autoimmune Neuroinflammation. *Front Immunol* **8**, 532 (2017).

9. T. S. Davidson, R. J. DiPaolo, J. Andersson, E. M. Shevach, Cutting Edge: IL-2 is essential for TGF-beta-mediated induction of Foxp3+ T regulatory cells. *J Immunol* **178**, 4022-4026 (2007).

10. W. Chen *et al.*, Conversion of peripheral CD4+CD25- naive T cells to CD4+CD25+ regulatory T cells by TGF-beta induction of transcription factor Foxp3. *J Exp Med* **198**, 1875-1886 (2003).

11. T. L. Nguyen, N. T. Makhlouf, B. A. Anthony, R. M. Teague, R. J. DiPaolo, In vitro induced regulatory T cells are unique from endogenous regulatory T cells and effective at suppressing late stages of ongoing autoimmunity. *PLoS One* **9**, e104698 (2014).

12. S. E. Weber *et al.*, Adaptive islet-specific regulatory CD4 T cells control autoimmune diabetes and mediate the disappearance of pathogenic Th1 cells in vivo. *J Immunol* **176**, 4730-4739 (2006).

13. R. J. DiPaolo *et al.*, Autoantigen-specific TGFbeta-induced Foxp3+ regulatory T cells prevent autoimmunity by inhibiting dendritic cells from activating autoreactive T cells. *J Immunol* **179**, 4685-4693 (2007).

14. S. G. Zheng *et al.*, CD4+ and CD8+ regulatory T cells generated ex vivo with IL-2 and TGF-beta suppress a stimulatory graft-versus-host disease with a lupus-like syndrome. *J Immunol* **172**, 1531-1539 (2004).

15. M. E. Raeber, D. Sahin, U. Karakus, O. Boyman, A systematic review of interleukin-2-based immunotherapies in clinical trials for cancer and autoimmune diseases. *EBioMedicine* **90**, 104539 (2023).

16. R. Zhang *et al.*, Low-dose IL-2 therapy in autoimmune diseases: An update review. *Int Rev Immunol* **43**, 113-137 (2024).

17. L. Khoryati *et al.*, An IL-2 mutein engineered to promote expansion of regulatory T cells arrests ongoing autoimmunity in mice. *Sci Immunol* **5**, (2020).

18. S. N. Quayle *et al.*, CUE-101, a Novel E7-pHLA-IL2-Fc Fusion Protein, Enhances Tumor Antigen-Specific T-Cell Activation for the Treatment of HPV16-Driven Malignancies. *Clin Cancer Res* **26**, 1953-1964 (2020).

19. D. J. Stauber, E. W. Debler, P. A. Horton, K. A. Smith, I. A. Wilson, Crystal structure of the IL-2 signaling complex: paradigm for a heterotrimeric cytokine receptor. *Proc Natl Acad Sci U S A* **103**, 2788-2793 (2006).

20. K. M. Heaton, G. Ju, E. A. Grimm, Human interleukin 2 analogues that preferentially bind the intermediate-affinity interleukin 2 receptor lead to reduced secondary cytokine secretion: implications for the use of these interleukin 2 analogues in cancer immunotherapy. *Cancer Res* **53**, 2597-2602 (1993).

21. T. R. Mosmann *et al.*, Species-specificity of T cell stimulating activities of IL 2 and BSF-1 (IL 4): comparison of normal and recombinant, mouse and human IL 2 and BSF-1 (IL 4). *J Immunol* **138**, 1813-1816 (1987).

22. K. Abnaof *et al.*, TGF-beta stimulation in human and murine cells reveals commonly affected biological processes and pathways at transcription level. *BMC Syst Biol* **8**, 55 (2014).

23. J. E. Zuniga *et al.*, Assembly of TbetaRI:TbetaRII:TGFbeta ternary complex in vitro with receptor extracellular domains is cooperative and isoform-dependent. *J Mol Biol* **354**, 1052-1068 (2005).

24. J. Groppe *et al.*, Cooperative assembly of TGF-beta superfamily signaling complexes is mediated by two
disparate mechanisms and distinct modes of receptor binding. *Mol Cell* **29**, 157-168 (2008).
25. X. Li, Y. Liang, M. LeBlanc, C. Benner, Y. Zheng, Function of a Foxp3 cis-element in protecting
regulatory T cell identity. *Cell* **158**, 734-748 (2014).
26. A. S. Wegrzyn, A. E. Kedzierska, A. Obojski, Identification and classification of distinct surface markers
of T regulatory cells. *Front Immunol* **13**, 1055805 (2022).
27. C. Gootjes, J. J. Zwaginga, B. O. Roep, T. Nikolic, Defining Human Regulatory T Cells beyond FOXP3:
The Need to Combine Phenotype with Function. *Cells* **13**, (2024).
28. S. Dikiy, A. Y. Rudensky, Principles of regulatory T cell function. *Immunity* **56**, 240-255 (2023).
29. J. D. Fontenot, J. P. Rasmussen, M. A. Gavin, A. Y. Rudensky, A function for interleukin 2 in Foxp3-
expressing regulatory T cells. *Nat Immunol* **6**, 1142-1151 (2005).
30. A. Arvey *et al.*, Inflammation-induced repression of chromatin bound by the transcription factor Foxp3 in
regulatory T cells. *Nat Immunol* **15**, 580-587 (2014).
31. J. A. Hill *et al.*, Foxp3 transcription-factor-dependent and -independent regulation of the regulatory T cell
transcriptional signature. *Immunity* **27**, 786-800 (2007).
32. N. Sugimoto *et al.*, Foxp3-dependent and -independent molecules specific for CD25+CD4+ natural
regulatory T cells revealed by DNA microarray analysis. *Int Immunol* **18**, 1197-1209 (2006).
33. M. Ahn, J. Dostal, P. Hegde, D. J. Lee, CTLA-4 expression on tregs is needed for suppression of
autoimmune uveitis. *Sci Rep* **15**, 17745 (2025).
34. M. G. Petrillo *et al.*, GITR+ regulatory T cells in the treatment of autoimmune diseases. *Autoimmun Rev*
14, 117-126 (2015).
35. S. Tagkareli *et al.*, CD103 integrin identifies a high IL-10-producing FoxP3(+) regulatory T-cell population
suppressing allergic airway inflammation. *Allergy* **77**, 1150-1164 (2022).
36. K. E. Earle *et al.*, In vitro expanded human CD4+CD25+ regulatory T cells suppress effector T cell
proliferation. *Clin Immunol* **115**, 3-9 (2005).
37. R. S. McHugh, E. M. Shevach, D. H. Margulies, K. Natarajan, A T cell receptor transgenic model of
severe, spontaneous organ-specific autoimmunity. *Eur J Immunol* **31**, 2094-2103 (2001).
38. R. J. DiPaolo, D. D. Glass, K. E. Bijwaard, E. M. Shevach, CD4+CD25+ T cells prevent the development
of organ-specific autoimmune disease by inhibiting the differentiation of autoreactive effector T cells. *J*
Immunol **175**, 7135-7142 (2005).
39. S. Gaggero *et al.*, IL-2 is inactivated by the acidic pH environment of tumors enabling engineering of a pH-
selective mutein. *Sci Immunol* **7**, eade5686 (2022).
40. W. J. Leonard, J. X. Lin, Strategies to therapeutically modulate cytokine action. *Nat Rev Drug Discov* **22**,
827-854 (2023).
41. D. A. Horwitz *et al.*, Nanoparticles loaded with IL-2 and TGF-beta promote transplantation tolerance to
alloantigen. *Front Immunol* **15**, 1429335 (2024).
42. W. Lin *et al.*, Regulatory T cell development in the absence of functional Foxp3. *Nat Immunol* **8**, 359-368
(2007).
43. F. Alderuccio, B. H. Toh, P. A. Gleeson, I. R. van Driel, A novel method for isolating mononuclear cells
from the stomachs of mice with experimental autoimmune gastritis. *Autoimmunity* **21**, 215-221 (1995).
44. Y. Hao *et al.*, Dictionary learning for integrative, multimodal and scalable single-cell analysis. *Nat*
Biotechnol **42**, 293-304 (2024).
45. D. Haribhai *et al.*, Regulatory T cells dynamically control the primary immune response to foreign antigen.
J Immunol **178**, 2961-2972 (2007).
785