A proteostasis clock underlies the timing of bacterial dormancy and antibiotic tolerance

Fa-Zhan Wang[1], Yi-Wen Zhang[1,2], and Jia-Feng Liu[1,3]*

[1]Center for Infection Biology, School of Basic Medical Sciences, Tsinghua University, Beijing 100084, China.

[2]Peking University-Tsinghua University-National Institute of Biological Sciences Joint Graduate Program, Academy

for Advanced Interdisciplinary Studies, Peking University, Beijing 100871, China.

[3]Tsinghua-Peking Center for Life Sciences, Beijing 100084, China.

*Corresponding authors: Email: jfliu@mail.tsinghua.edu.cn

8   ABSTRACT

Dormancy, characterized by an extended lag time before resuming growth, protects bacteria from antibiotic

stress and facilitates the emergence of resistance. However, a common mechanism that precisely regulates

bacterial lag time remains poorly defined. Here we show that the formation and disassembly dynamics of

protein aggregates function as a proteostasis clock that governs the lag time distribution. By tracking the

aggregate and regrowth dynamics of thousands of cells, we observed that dormant bacteria from diverse

genetic and environmental contexts converge to delayed aggregate disassembly as a common rate-limiting

step for growth resumption. Mechanistically, the chromosomal replication initiator DnaA is sequestered in

aggregates and spatially separated from the nucleoid, thereby preventing DNA replication until proteostasis

restoration enables aggregate disassembly. We modulated bacterial lag time distribution by genetically and

chemically targeting the proteostasis-replication axis. Remarkably, lag extension scales linearly with the

disruption level of proteostasis across diverse backgrounds, indicating a universal coupling between

dormancy and proteostasis. This proteostasis collapse renders dormant bacteria, including clinical isolates,

more sensitive to additional proteotoxic stress. These findings reveal a timekeeping mechanism in cellular

dormancy, highlighting proteostasis as a shared target for combating antibiotic tolerance.

24 INTRODUCTION

The timing of replication initiation is critical for cells to balance stress tolerance and proliferation[1]. Dormant

bacteria exhibit an extended lag time before resuming growth, even upon transition to nutrient-rich

environments[2,3]. This growth delay confers protection against antibiotics targeting actively proliferating cells,

a phenomenon termed antibiotic tolerance[2-5]. *In vitro* evolution experiments demonstrate that lag duration

can be optimized to match the duration of antibiotic treatment, thereby coordinating improved survival and

timely regrowth[2]. Increasing evidence establishes lag time as a plastic, heritable, and evolvable trait that

facilitates bacterial adaptation to fluctuating environments. Antibiotic tolerance conferred by lag extension

impedes the clearance of pathogens during treatment and facilitates the subsequent evolution of

resistance[3,6]. Consequently, a better understanding of dormancy regulation is essential for developing

strategies to eliminate antibiotic persisters.

Despite numerous mutations extending lag time have been identified in toxin-antitoxin systems[7], aminoacyl-

tRNA synthetases[2,3,5], transcriptional regulators[8], and metabolic enzymes[9], it has been a long-standing

challenge to determine whether there is a common mechanism underlying these different dormant states.

Recent evidence has established a correlation between exit from dormancy and disassembly of protein

aggregates[4,10-12]. Nevertheless, a causal link between protein aggregation and dormancy remains poorly
defined, and the specific identity of the aggregate components that govern the lag time distribution has yet
to be elucidated[11,12]. We previously reported that the sequestration of seryl-tRNA synthetase (SerRS) into
biomolecular condensates regulates compartmentalized translation and bacterial dormancy[13]. However,
direct evidence linking the aggregation dynamics of specific proteins to the regulation of the dormancy-
proliferation transition remains lacking. More broadly, elucidating the logic of dormancy timing would provide
insights into the design principle of timekeeping in biological systems.

To characterize the molecular basis determining the lag time distribution, we integrated automated single-
cell lag-time measurements with high-resolution fluorescence microscopy and quantitative omics profiling.
Using long-lag mutants of *Escherichia coli*, we observed that increased aggregation of specific proteins,
such as the phenylalanyl-tRNA synthetase α subunit (PheS), overwhelmed cellular proteostasis capacity,
leading to dose-dependent delays in aggregate disassembly and corresponding increases in lag time. We
showed that this aggregate burden delayed replication initiation by stimulating proteases to sequester the
chromosomal replication initiation protein DnaA away from the nucleoid. The introduction of aggregates or
proteases was sufficient to extend lag time, whereas increasing soluble DnaA levels accelerated dormancy
exit, demonstrating a direct proteostasis-replication axis. Across diverse genetic mutants, environmental
stresses, and clinical isolates, lag time increased in proportion to the degree of proteostasis disruption, and
in turn, accelerating aggregate disassembly through scavenging hydroxyl radicals facilitated dormancy exit.
These findings reveal that lag time can be understood as the timescale of cellular recovery from disrupted
proteostasis to a replication-competent state. These findings establish cellular proteostasis as a regulating
clock that operates in dormant cells.

RESULTS

Long-lag mutants exhibit delayed aggregate disassembly and disrupted proteostasis
To elucidate the regulatory mechanisms governing bacterial lag time, we first performed evolution
experiments to identify genetic factors that shape the lag time distribution. Based on the observation that
cyclic ampicillin exposure selects for extended lag time[2,3], we evolved *E. coli* KLY under cyclic treatment
with the carbapenem ertapenem (ETP). This antibiotic was specifically chosen for the lack of ETP-resistant
carbapenemase genes in the KLY strain[2], thus preventing the rapid emergence of resistance and enabling
long-term tracking of dormancy evolution. Using ScanLag[14] to quantify single-cell lag time, we observed
that, following cyclic ETP treatments, both the mean and variance of colony appearance times increased
across three parallel batch cultures (Extended Data Fig. 1a,b). Single colonies exhibiting different lag times
were isolated from the evolved lines and analysed by whole-genome sequencing (WGS). Mutations were
identified in several aminoacyl-tRNA synthetases (*pheT*, *alaS*, *leuS*) and the cysteine tRNA (*cysT*), each
conferring a specific increase in lag time (Fig. 1a and Supplementary Table 1). Notably, two independent
mutations were mapped to the phenylalanyl-tRNA synthetase β subunit (*pheT*), both of which produced
pronounced lag extension. Restoring the wild-type (WT) allele of *pheT* fully reverted this phenotype
(Extended Data Fig. 1c). In line with prior reports implicating multiple aminoacyl-tRNA synthetases (e.g.,
metG and *gltX*) in lag regulation[2,3,5,15], these results hinted at a conserved yet unresolved link between this
gene family and dormancy control.

Increasing evidence suggests that protein aggregation facilitates bacterial dormancy, whereas aggregate
disassembly is critical for cellular resuscitation[4,11,12]. To examine whether the identified mutants exhibited
changed aggregate dynamics, we stained WT and two *pheT* mutants with Proteostat, a fluorescent dye that
specifically binds to insoluble protein aggregates[16,17]. Consistent with previous studies[4], we observed the
formation of insoluble protein aggregates in stationary-phase (S-phase) bacteria, visible as dark foci in
brightfield microscopy and associated with elevated Proteostat fluorescence (Fig. 1b and Extended Data
Fig. 1d). However, quantitative analysis revealed that the proportion of cells containing protein aggregates
(Fig. 1c) and the intensity of Proteostat fluorescence (Extended Data Fig. 1e) were indistinguishable
between stationary-phase WT and *pheT* mutants, despite their markedly different lag times (Fig. 1a).
Isolation and quantification of insoluble protein fractions[18] further confirmed comparable levels of
aggregated proteins in these strains (Extended Data Fig. 1f). These results indicated that the aggregate
amount in stationary-phase cells was insufficient to account for the extended lag time of *pheT* mutants.
However, after nutrient replenishment, we observed that WT cells exhibited rapid protein disaggregation
within two hours, whereas aggregates in *pheT* mutants persisted for more than six hours until cell division
(Fig. 1b,c and Extended Data Fig. 1d,e). Thus, despite the similar aggregate amount, aggregates
disassemble more slowly in dormant mutants upon cell resuscitation.

We next determined whether the delayed aggregate disassembly in dormant mutants is merely an
epiphenomenon of extended lag time or instead reflects a compromised proteostasis capacity to clear
protein aggregates. It is well established that cells with disrupted proteostasis typically exhibit reduced
ability to cope with additional proteotoxic stress[19-21]. We compared the sensitivity of WT and *pheT* cells to
multiple proteotoxic stressors, including heat shock, oxidative stress, and puromycin, all is known to
promote protein aggregation[4,22,23]. Remarkably, killing assays revealed that ETP-tolerant *pheT* mutants
were more susceptible to heat shock and hydrogen peroxide than WT cells (Fig. 1d,e). Furthermore,
puromycin, an aminoacyl-tRNA analogue that induces misfolded protein accumulation[23,24], caused more
pronounced growth inhibition in *pheT* mutants than in WT cells, an effect that was not observed with other
antibiotics or non-proteotoxic stressors (Fig. 1f and Extended Data Fig. 2). These results indicate that long
lag time confers ETP tolerance but concurrently compromises cellular proteostasis, characterized by
delayed aggregate disassembly and increased vulnerability to proteotoxic stress.

Increased PheS aggregation extends lag time in *pheT* mutants
To investigate the molecular basis for the delayed aggregate disassembly in *pheT* mutants, we compared
the composition of insoluble aggregates isolated from stationary-phase WT and *pheT* cells. Label-free
quantitative proteomics identified phenylalanyl-tRNA synthetase α subunit (PheS) as a top-enriched protein
within the aggregates of *pheT* mutants compared to WT cells (Fig. 2a,b). In *E. coli*, phenylalanyl-tRNA
synthetase functions as a heterotetramer of two α (PheS) and two β (PheT) subunits, both of which are
essential for bacterial viability[25]. Quantitative analysis confirmed that total protein levels of PheS did not
increase in both *pheT* mutants, indicating that its increased aggregation results from reduced solubility
(Extended Data Fig. 3a,b). To validate these proteomic data, we tagged PheS with ECFP and monitored
its spatiotemporal localization during dormancy. We observed the formation of PheS-ECFP aggregates in
stationary-phase cells that colocalized with Proteostat signals (Fig. 2c and Extended Data Fig. 3c). Upon
nutrient replenishment, these aggregates disassembled rapidly in WT cells but persisted longer in *pheT*

mutants (Fig. 2c and Extended Data Fig. 3c). These results show that in dormant bacteria, *pheT* mutations
cause increased accumulation and delayed clearance of PheS aggregates.

To establish a causal relationship between PheS aggregation and dormancy, we examined how titrating
PheS levels modulated aggregate dynamics and lag time. Specifically, bacteria carrying an IPTG-inducible
PheS-ECFP system were grown for 48h into stationary phase under graded concentrations of IPTG.
Quantitative analysis of PheS-ECFP and Proteostat fluorescence showed that increased PheS induction
elevated the total aggregate burden in *pheT* mutants (Extended Data Fig. 3d). Crucially, upon nutrient
replenishment, this elevated PheS aggregation decreased the aggregate disassembly rates in a dose-
dependent manner (Fig. 2d) and led to corresponding extensions of lag times (Fig. 2e and Extended Data
Fig. 3e,f). At higher levels of PheS induction, the extended lag phase progressed into a loss of culturability,
characterized by a remarkable reduction in colony-forming units (CFUs) (Fig. 2f). Increased PheS
expression had no detectable effect on aggregate dynamics or lag time in WT cells (Fig. 2d–f and Extended
Data Fig. 3d), underscoring the role of wild-type PheT in maintaining PheS solubility. Consistent with
previous results, this PheS-induced extension of lag time further enhanced ETP tolerance but compromised
the capacity of *pheT* mutants to cope with proteotoxic stress (Extended Data Fig. 3g,h). Collectively, these
results show that modulating the degree of PheS aggregation is sufficient to tune cellular proteostasis and
lag time distribution in *pheT* mutants.

Aggregate-stimulated proteases sequester DnaA to delay replication initiation
We next sought to elucidate why increased PheS aggregation suffices to delay the transition from dormancy
to replication initiation. Given that chaperones and proteases have been reported to facilitate protein
disaggregation and cell resuscitation[4,12,26], we tested whether their overexpression could modulate bacterial
exit from dormancy. We found that overexpressing chaperones (DnaK, ClpB) failed to accelerate growth
resumption, whereas overexpressing proteases (HslVU, Lon) unexpectedly further delayed dormancy exit
(Extended Data Fig. 4a). Microscopy showed that the protease HslU formed aggregates colocalized with
PheS-ECFP and exhibited synchronized delayed disaggregation in *pheT* mutants after nutrient
replenishment (Extended Data Fig. 4b). However, if the primary role of proteases was to clear protein
aggregates, their overexpression would be expected to accelerate, rather than delay, the exit from
dormancy. A recent study showed that depletion of the protease Lon abolished the extended lag time in a
metG mutant[5], supporting the function of proteases in promoting dormancy. Indeed, beyond their role in
aggregate clearance, accumulating evidence suggests that proteases also actively target and degrade
native proteins involved in cell-cycle regulation[1,26,27]. In *Caulobacter crescentus*, proteases are activated by
misfolded proteins to degrade the chromosomal replication initiator DnaA and induce cell-cycle arrest[1].
Given the conserved role of DnaA in replication initiation and its specificity as a protease substrate in *E.*
coli[28,29], we next explored whether the aggregate-stimulated proteases might regulate DnaA function during
cell-cycle progression.

We constructed chromosomal fluorescent fusions of HslU and DnaA and confirmed that this dual-labelling
strategy did not perturb the lag time distribution of WT or *pheT^D1* mutant (Extended Data Fig. 5a).
Remarkably, we observed that mCherry-DnaA formed aggregates that colocalized with HslU-ECFP in
stationary-phase WT and *pheT^D1* cells (Fig. 3a,b). Upon nutrient replenishment, these aggregates dissolved
rapidly in WT cells; in contrast, DnaA persisted in a sequestered state within aggregates for over six hours

in the dormant mutant (Fig. 3a,b). As previous research has established that protein aggregates typically
localize to nucleoid-free regions owing to macromolecular crowding effects in the nucleoid[30,31], we used
DAPI staining to examine the spatiotemporal relationship between DnaA and the bacterial nucleoid. We
found that DnaA aggregates maintained a spatial separation from the nucleoid throughout the lag phase,
rendering the initiator inaccessible to the chromosomal origin of replication (*oriC*) (Fig. 3a,b). As a result,
the integration of DnaA into protein aggregates directly prevents DNA replication until restored proteostasis
enables aggregate disassembly.

Modulating lag time distribution by targeting the proteostasis-replication axis
To validate this proposed mechanism of dormancy, we reasoned that increasing the abundance of protein
aggregates or proteases would further exacerbate DnaA sequestration and delay growth resumption.
ScanLag analysis demonstrated that either elevated expression of PheS or HslVU markedly extended lag
times in the dormant mutants (Fig. 3c,d). In addition, promoting protein aggregation by the antibiotic
puromycin also produced similar lag time extensions (Fig. 3e), indicating that this proteostasis-replication
axis can integrate different proteotoxic stresses to determine the timing of dormancy. Furthermore,
increased HslVU expression acted synergistically with the *pheT* mutations to cause a dose-dependent
reduction in colony-forming capacity despite 3 days of incubation (Fig. 3f). These observations indicate that
modulating the level of aggregates and proteases could establish a dormancy continuum spanning from
lag time extension to an unculturable state[32].

Our findings also raise the possibility that releasing DnaA from aggregate-mediated sequestration may
facilitate the exit from dormancy. Guided by prior work showing that fluorescent tag positioning may
modulate the localization and activity of DnaA[33,34], we compared the phenotypes of variants with mCherry
fused to the amino-terminus, carboxy-terminus, or internal regions of DnaA. Notably, we found that, in
stationary-phase cells, the carboxy-terminal DnaA-mCherry exhibited significantly lower aggregation levels
compared to other variants (Extended Data Fig. 5b,c). As previously reported[29], overexpressing native
DnaA caused pronounced growth defects (Extended Data Fig. 5d) owing to hyper-replication initiation.
Strikingly, however, we found that overexpression of the soluble DnaA variant not only accelerated growth
resumption in *pheT[D1]* cells but also completely abolished the extended dormancy induced by elevated PheS
and HslVU levels (Fig. 3g and Extended Data Fig. 5d). This enhanced DnaA solubility is likely attributable
to the structural masking of its carboxy-terminal domain, supported by the experiments that fusing a
solubility-enhancing Maltose-binding protein (MBP) tag exclusively to the carboxy-terminus produced
similar effects (Extended Data Fig. 5e,f). Thus, these findings indicate that bypassing the sequestration of
DnaA within protein aggregates can directly promote dormancy exit, underlining a decisive role of DnaA in
gating the transition from dormancy to proliferation.

Nevertheless, we noticed that overexpressing soluble DnaA failed to accelerate dormancy exit in the *pheT[D2]*
mutant (Extended Data Fig. 5g), implying the existence of additional protease targets beyond DnaA in this
specific background. Given that *E. coli* encodes multiple proteases, including HslVU, Lon, ClpP (complexed
with ClpX or ClpA), and FtsH[26], we systematically examined how expressing each protease affected
dormancy exit of different mutants. We found that, while both *pheT* mutants shared a susceptibility to HslVU,
they exhibited specific sensitivity to other proteases: *pheT[D1]* bacteria were more sensitive to Lon, whereas
pheT[D2] bacteria showed greater sensitivity to ClpA and ClpX (Fig. 3h). These results imply that bacteria

may rely on multiple, partly redundant proteolytic pathways to coordinate stress responses with DNA
replication. Altogether, these results reveal a proteostasis-replication axis that can be targeted to precisely
modulate the lag time distribution.

Long-lag phenotypes are associated with disrupted proteostasis across diverse backgrounds
We next addressed whether the regulatory mechanism identified in *pheT* mutants could be generalized to
other genetically distinct dormant mutants. We observed that a cysteine tRNA mutant (*cysT*[D1]) exhibited
similarly delayed aggregate disassembly and increased puromycin sensitivity (Extended Data Fig. 6a–d).
Quantitative proteomic analysis of the insoluble fractions revealed mutation-specific heterogeneity in
aggregate composition across *pheT*[D1], *pheT*[D2], and *cysT*[D1] cells (Extended Data Fig. 6e,f). Notably, the
ferrichrome-transporting outer-membrane protein FhuA[35] showed significantly elevated aggregation levels
across all three dormant mutants (Extended Data Fig. 6e,f). We subsequently demonstrated that inducible
FhuA expression alone was sufficient to facilitate dormancy in these mutants (Extended Data Fig. 6g,h).
This effect was independent of its canonical role in ferrichrome transport, as a transport-defective FhuA
Δ236-248 variant[36] produced a comparable extension of lag time (Extended Data Fig. 6i). Moreover,
overexpression of the proteases HslVU and Lon further delayed dormancy exit in *cysT*[D1] cells (Extended
Data Fig. 6j,k). Together, these results indicate that the disruption of proteostasis may represent a shared
physiological state of distinct long-lag mutants.

These results prompted us to investigate whether the proteostasis-governed dormancy driven by genetic
mutations also extends to aggregates induced by environmental stresses. We first confirmed that both heat
shock (49 or 55 °C) and puromycin can induce the formation of protein aggregates in WT cells (Extended
Data Fig. 7a). Proteostat staining and ScanLag analysis revealed that exposure to 55 °C for 1 hour resulted
in delayed aggregate disassembly and a corresponding extension of lag time in WT cells (Fig. 4a,b). In
contrast, treatment at 49 °C for 1 hour, although sufficient to induce aggregates in exponentially growing
cells (Extended Data Fig. 7a), had no detectable effect on aggregate disassembly or lag time (Fig. 4a,b).
These data imply that the bacterial protein quality-control network can buffer low levels of proteotoxic stress
to support timely regrowth, and lag time is extended only when cumulative stress overwhelms cellular
proteostasis capacity to clear aggregates. Supporting this view, at the concentration below the minimum
inhibitory concentration (MIC), puromycin permitted bacterial growth but extended lag times in a dose-
dependent manner (Extended Data Fig. 7b). Following treatment with puromycin above the MIC for varying
durations, transfer to antibiotic-free medium revealed a post-antibiotic effect where the lag time increased
with the duration of preceding exposure (Fig. 4a–c and Extended Data Fig. 7c). These results suggest a
cellular memory of accumulated proteotoxic stress. Similar to dormant mutants, stress-induced long-lag
WT cells also showed persistent sequestration of HslU and DnaA within aggregates during dormancy
(Extended Data Fig. 7d), indicating the conservation of the proteostasis-replication axis across diverse
conditions.

Lag-time extension scales with the degree of proteostasis disruption across stresses
To establish a quantitative relationship between proteostasis disruption and lag-time extension across
various stresses, we sought a functional indicator that quantifies the degree of proteostasis disruption under
different conditions. Since the formation of protein aggregates mainly represents a cellular response to

proteotoxic stress rather than a definitive hallmark of proteostasis collapse (Fig. 4a,b and Extended Data
Fig. 7a), we tried to employ the bacterial vulnerability to an exogenous proteotoxic challenge, puromycin,
as a functional indicator of proteostasis disruption based on the following observations.

First, we have confirmed that puromycin can promote protein aggregation, delay aggregate disassembly,
and extend lag time in a dose-dependent manner (Fig. 4a–c and Extended Data Fig. 7a–c). Accordingly,
progressive disruption of proteostasis would compromise the cellular capacity to handle puromycin-induced
aggregation. Second, we previously found that dormant mutants contained a similar aggregate amount
relative to WT cells, but exhibited markedly increased puromycin sensitivity (Fig. 1f and Extended Data Fig.
1f), indicating that puromycin sensitivity reflects proteostasis disruption more accurately than the aggregate
amount. Third, treatment at 49 °C for 1 hour failed to delay aggregate disassembly, consistent with its
unaffected sensitivity to puromycin (Fig. 4a,b and Extended Data Fig. 7e). For comparison, the 55°C
treatment slowed aggregate disassembly and increased the sensitivity to puromycin at the same time (Fig.
4a,b and Extended Data Fig. 7e). These observations support puromycin sensitivity as a sensitive indicator
of proteostasis disruption in bacteria.

On this basis, we quantified the level of proteostasis disruption following various stresses by measuring
their puromycin sensitivity, calculated as the $\log_{10}$ ratio of CFU on antibiotic-free relative to puromycin plates.
Strikingly, we observed that lag time increased in proportion to the degree of proteostasis disruption,
regardless of stress type (Fig. 4d). In addition to canonical proteotoxic stresses, prolonged exposure to the
antibiotic azithromycin or chloramphenicol similarly compromised proteostasis and extended lag times (Fig.
4d). This scaling relation suggests a universal coupling between proteostasis disruption and lag extension
of bacteria. The quantitative relationship also extended to the dormant mutants, including the previously
described[3] long-lag mutant $metG^T$ (Fig. 4e). Thus, in bacteria, progressive disruption of proteostasis serves
as a tunable timer that governs lag time distribution across diverse contexts. As observed in dormant
mutants, stress-induced dormancy conferred increased ETP tolerance while simultaneously sensitizing
bacteria to puromycin (Fig. 4f). From a clinical perspective, this disrupted proteostasis could represent an
Achilles' heel of dormant bacteria, offering a common target for therapeutic exploitation.

Restoring proteostasis accelerates dormancy exit
Once antibiotic stress is removed, rapid growth resumption should confer a significant fitness advantage.
We therefore performed evolution experiments to identify genetic factors capable of shortening the lag
times of *pheT* mutants. Serial passaging of *pheT* mutants in antibiotic-free LB medium resulted in the rapid
evolution of shortened lag times within seven cycles (Extended Data Fig. 8a,b). Reverse mutants were
subsequently isolated and analysed by WGS (Extended Data Fig. 8c and Supplementary Table 2).
Intriguingly, while these reverse mutations mapped to diverse gene families, restored proteostasis emerges
as a convergent hallmark among reverse mutants, manifested as accelerated protein disaggregation and
reduced sensitivity to proteotoxic stress (Extended Data Fig. 8d,e). These results underscore proteostasis
as a regulatory hub shaping lag time distribution.

As delayed aggregate disassembly extends lag time, we reasoned that accelerating aggregate disassembly
might facilitate dormancy exit. Increasing evidence indicates that reactive oxygen species (ROS),
particularly hydrogen peroxide, can form spontaneously at aggregate interfaces and establish a self-
reinforcing feedback loop that promotes aggregation[37,38]. Using the H_2O_2-specific fluorogenic probe Peroxy

Orange 1 (PO-1)[39,40], we confirmed that H_2O_2 is enriched in aggregates compared to the cytoplasm (Fig.
5a,b). Notably, $pheT^{D1}$ cells maintained higher H_2O_2 levels than WT cells throughout dormancy (Fig. 5a,b).
Consistently, $pheT^{D1}$ cells exhibited elevated sensitivity to Fe^{2+} treatment, which reacts with H_2O_2 via the
Fenton reaction to generate more hydroxyl radicals[41] (Extended Data Fig. 9a). To determine if the increased
H_2O_2 level contributes to delayed aggregate disassembly, we supplemented the medium with catalase to
scavenge H_2O_2[42] and monitored disaggregation dynamics. Remarkably, catalase treatment significantly
reduced intracellular H_2O_2 levels in $pheT^{D1}$ cells (Fig. 5a,b) and accelerated aggregate disassembly (Fig.
5c,d). Accordingly, we observed that the addition of catalases directly facilitated the exit from dormancy
(Fig. 5e). Similar promoting effects were obtained with the addition of thiourea, another well-characterized
hydroxyl radical scavenger[43] (Extended Data Fig. 9b,c). Beyond dormant mutants, catalase treatment also
accelerated bacterial growth resumption from puromycin-induced dormancy (Extended Data Fig. 9d). Thus,
scavenging hydroxyl radicals serves as an effective strategy to accelerate aggregate disassembly and
promote bacterial dormancy exit.

Proteostasis-governed dormancy is observed in clinical isolates
Finally, we extended our analysis to other bacterial species, including the clinical pathogenic isolate
Klebsiella pneumoniae S837. In this strain, prolonged starvation resulted in a pronounced extension of lag
time (Fig. 6a). Through Proteostat staining, we observed the formation of protein aggregates in stationary-
phase *K. pneumoniae* cells, and upon nutrient replenishment, aggregates disassembled more slowly in the
long-lag clinical isolate S837 compared to the laboratory strain ATCC 43816 (Fig. 6b). These observations
confirmed a temporal coupling between dormancy exit and aggregate disassembly in *K. pneumoniae*.
Moreover, we found that the introduction of misfolded proteins with puromycin alone was sufficient to extend
lag times in both *K. pneumoniae* and the Gram-positive bacterium *Bacillus subtilis* (Fig. 6c), indicating that
this proteotoxicity-induced dormancy is not restricted to a single bacterial species. From a clinical
perspective, extended lag time in *K. pneumoniae* S837 was associated with increased ETP tolerance (Fig.
6d); however, this dormant state also sensitized the pathogen to the proteotoxic stressor puromycin (Fig.
6d). These findings suggest that the proteostasis-governed dormancy is a common strategy across
bacterial species and implicate the proteostasis network as a potential target for therapeutic intervention
against antibiotic persisters.

DISCUSSION

The accumulation of protein aggregates is typically linked to cytotoxicity and aging-related diseases in
eukaryotes[19-21]. The disassembly of such aggregates is often viewed as too slow to be relevant on
physiological timescales[44]. However, our work proposes that the kinetics of aggregate disassembly can be
repurposed as a functional timing device in bacteria for fine-tuning lag time to match the duration of stress
(Fig. 6e). Conceptually, this proteostasis-governed timing device and canonical circadian clocks driven by
transcription-translation feedback loops[45] represent two distinct evolutionary solutions to the problem of
temporal control. Circadian clocks provide a stable, cell-autonomous mechanism to synchronize internal
physiology with the 24-hour light-dark cycle, whereas the proteostasis clock integrates the history of
proteotoxic stress and encodes the timescale from disrupted proteostasis to a replication-capable state.
From an evolutionary perspective, while dedicated oscillators are favoured in environments with regular

cycles, a highly tunable timer based on the proteostasis network offers a more economical and flexible solution for bacteria adapting to fluctuating and unpredictable environments.

While protein aggregation has been implicated as a bacterial strategy to facilitate dormancy and antibiotic tolerance[46], the mechanistic link between aggregates and cell-cycle progression has remained elusive. Our results suggest that protein aggregation is not simply a passive process of deposition but actively regulates the spatiotemporal dynamics of the replication initiator DnaA, thereby gating the transition from dormancy to replication. Within this framework, increased aggregation engages proteases to sequester DnaA away from the nucleoid, which delays replication initiation until aggregates are disassembled. Guided by this, bacterial lag time can be precisely tuned by genetically or chemically targeting components of the proteostasis-replication axis. While DnaA is identified here as a critical client, other cell-cycle regulators are probably integrated into this timing device, yet its full regulatory architecture remains to be elucidated. Future *in vitro* reconstitution of the minimal components will be essential to unveil the operating principles of this timekeeping module.

Our work decouples the disruption of proteostasis from the mere presence of aggregates. We show that protein aggregation *per se* represents cellular responses to proteotoxic stresses, rather than a definitive indicator of dormancy. Instead, lag time distribution is determined by the disassembly rate of aggregates, a kinetic parameter that reflects the cellular proteostasis capacity. In bacteria, proteostasis is maintained by an intricate network coordinating protein synthesis, folding, trafficking, and degradation[47]. This systemic nature may explain why long-lag phenotypes are associated with such diverse genetic mutations and environmental factors: these perturbations may represent distinct ways that convergently shift the global proteostasis network away from a replication-competent state. In this view, lag time can be understood as a manifestation of how far the proteostasis network has been driven away from the replication state.

Across *E. coli* and clinical *K. pneumoniae* isolates, starvation-induced extension of lag time confers ertapenem tolerance while simultaneously causing collateral sensitivity to proteotoxic stress, revealing a universal trade-off between dormancy and proteostasis. This trade-off highlights proteostasis pathways as promising targets for the clinical elimination of antibiotic persisters, providing two directions for combating dormant bacteria. One approach is to promote protein disaggregation to accelerate dormancy exit, thereby sensitizing bacteria to antibiotics that target actively proliferating cells. An alternative strategy is to exacerbate proteotoxic stress, which would extend lag time and ultimately drive cells toward a loss of culturability.

Beyond bacteria, protein aggregates have been observed in diverse dormant systems, including yeast[48], oocytes[49], stem cells[17], and cancer cells[50,51]. Despite their well-established function in protein storage, our work hints at an active role of aggregates in regulating cell-cycle progression. The evolutionary conservation of the proteostasis network and DNA replication raises the possibility that regulated proteostasis may represent a broader design principle for timekeeping in cellular dormancy.

METHODS

Bacterial strains, plasmids, and growth conditions

All strains used in this study are indicated in Supplementary Table 3, plasmids in Supplementary Table 4, and primers in Supplementary Table 5. *K. pneumoniae, B. subtilis, E. coli* KLY, and derivative strains were grown at 37 °C in Luria-Bertani (LB) broth (Miller formulation; 10 g/L NaCl, 10 g/L tryptone, and 5 g/L yeast extract) with 220 rpm shaking or on solid LB medium prepared by supplementing the broth with 15 g/L agar, unless otherwise stated.

Polymerase chain reaction (PCR) amplification was performed using PrimeSTAR Max DNA Polymerase (Takara) according to the manufacturer's instructions. For plasmid construction, the gene PCR product was amplified from *E. coli* KLY and inserted into the pET28a/pET28a-mCherry backbone using the ClonExpress II One Step Cloning Kit (Vazyme). Cells were plated on LB plates containing kanamycin (100 µg/mL) for selection. Positive clones were validated by Sanger sequencing.

To label HslU and DnaA with fluorescent proteins on the chromosome, we used a two-step λ Red recombineering. At the first step, the *kan::sacB* cassette was amplified and transformed into the electrocompetent *E. coli* KLY cells with induced Red protein with IPTG (pSC101-RED-tet). Cells were plated on LB plates containing kanamycin (100 µg/mL) for selection. At the second step, the *hslU/dnaA-ECFP/mCherry* translational fusion fragment was amplified, transformed, and recombined into the bacterial genome of the electrocompetent cells with *kan::sacB* cassette at the targeting site and induced Red protein with IPTG, then the cells were counter-selected on sucrose plates. Positive clones were validated by Sanger sequencing. To restore the WT alleles of *pheT* mutants, P1 phage (ATCC 25404-B1) was used to conduct general transduction[52], and primer design was using Dup-In[53].

Evolutionary protocol

For antibiotic evolution, an overnight culture was diluted 1:100 into 1 mL LB medium supplemented with 1 µg/mL ertapenem and incubated for 4 or 12 hours alternately at 37 °C with shaking. After killing, the cultures were washed twice with 0.9% NaCl to remove antibiotics, resuspended in 1 mL fresh LB medium, and then grown overnight at 37 °C with shaking.

For reverse evolution, an overnight culture of dormant mutant was diluted 1:100 into 1 mL antibiotic-free LB medium and grown overnight at 37 °C with shaking.

ScanLag analysis

Bacteria were serially diluted to approximately 200 CFU per plate and plated on solid LB medium. The plates were then placed in a ScanLag setup[14] at 37 °C. This setup automatically monitors colony growth every 20 minutes and extracts the distribution of appearance times. The colony detection threshold size is set at 100 pixels.

Whole genome sequencing

Genomic DNA was extracted using a TIANamp Bacteria DNA Kit (TIANGEN; Beijing; China). Sequencing was performed by Sinobiocore using Illumina-HiSeq with 150 bp paired-end at an average coverage of 100-200. Single nucleotide polymorphisms (SNPs) and indels were identified by bwa version 0.7.18.

Cell staining and microscopy

For Proteostat staining, bacteria were processed following the manufacturer's instructions (Enzo Life
Sciences; enz-51035-K100): Briefly, bacteria were fixed with 4% PFA for 30 minutes at room temperature
and permeabilized in Assay Buffer with 0.5% Triton X-100 for 30 minutes on ice. Cells were then washed
twice with PBS and stained with Proteostat dye for 30 minutes at room temperature. The stained cells were
washed twice with PBS before imaging.
For DAPI staining, fixed cells were stained with DAPI (MCE, HY-D0814) at a concentration of 1 µg/mL in
PBS for 10 minutes at room temperature. The stained cells were washed twice with PBS before imaging.
For Peroxy Orange 1 (PO-1) staining, bacteria were stained with PO-1 (MCE, HY-103469) at a
concentration of 10 µM in PBS for 1 hour at 37 °C. The stained cells were washed twice with PBS before
imaging.
Brightfield and fluorescence images were captured using a FV3000 confocal microscope (Olympus) with a
100 × objective. Appropriate filter sets were chosen for each fluorophore based on their excitation and
emission spectra. All images were analyzed using Fiji.
416

Insoluble protein isolation
The isolation of insoluble proteins was modified from Tomoyasu[16]. Briefly, cells (1 mL, 2×10^9 bacteria)
were collected and incubated in 40 µL buffer A (10 mM potassium phosphate buffer, pH 6.5, 1 mM EDTA,
20% (w/v) sucrose, 1 mg/mL lysozyme) for 30 minutes on ice. Cells were then mixed with 360 µL buffer B
(10 mM potassium phosphate buffer, pH 6.5, 1 mM EDTA) and lysed by sonication while cooling. Intact
cells were removed by centrifugation (2000 g, 20 minutes, 4 °C). The insoluble cell fraction was isolated by
centrifugation (15000 g, 20 minutes, 4 °C). The pellet fractions were resuspended in 400 µL buffer B with
2% NP40 to dissolve membrane proteins and isolated by centrifugation (15000 g, 30 minutes, 4 °C). NP40-
insoluble pellets were washed twice with 400 µL buffer B and resuspended in 100 µL buffer B by brief
sonication. The concentration of insoluble protein is determined using the BCA protein assay kit
(Beyotime).
428

429 **Killing assays**
430 For ertapenem and H_2O_2 killing, bacteria were diluted 1:100 into LB medium supplemented with 1 µg/mL
431 ertapenem or 10 mM H_2O_2 and incubated for the indicated time at 37 °C with shaking. For heat killing,
432 bacteria were diluted 1:100 into LB medium and incubated for 30 minutes at 55 °C. The CFUs were
433 evaluated by plating before and after stress exposure.
434

435 **Culturability tests**
436 Tenfold dilutions of bacteria were prepared in PBS and subsequently plated on solid LB medium
437 supplemented with the indicated concentrations of puromycin or IPTG. The CFUs were calculated after 72
438 hours of incubation at 37 °C.
439

440 **Growth curves**
441 Bacteria were diluted to 2×10^5 cells/mL in a 96-well plate of LB medium. Growth measurements were
442 performed by incubating the plate in a SPARK multimode microplate reader (Tecan) at 37 °C with
443 shaking. OD_{600} was measured every 10 minutes.
444

Quantitative proteomics

The total protein of bacteria is extracted using the Bacterial Protein Extraction Kit (Sangon). The insoluble protein is isolated as previously mentioned. The proteins were separated by SDS-PAGE, respectively. The gel bands of interest were excised from the gel, reduced with 5 mM of DTT, and alkylated with 11 mM iodoacetamide, which was followed by in-gel digestion with sequencing grade modified trypsin at 37 °C overnight. The peptides were extracted twice with 0.1% trifluoroacetic acid in 50% acetonitrile aqueous solution for 30 min and then dried in a speedvac. Peptides were redissolved in 20 µL 0.1% trifluoroacetic acid, and 6 µL of extracted peptides were analyzed by Thermo Scientific Orbitrap Exploris 480.

For LC-MS/MS analysis, the peptides were separated by an 85-min gradient elution at a flow rate of 0.30 µl/min with a Thermo-Dionex Ultimate 3000 HPLC system, which was directly interfaced with a Thermo Scientific Orbitrap Exploris 480 mass spectrometer. The analytical column was a home-made fused silica capillary column (75 µm ID, 350 mm length) packed with C-18 resin (1.9 µm, Dr.Maisch GmbH). Mobile phase consisted of 0.1% formic acid, and mobile phase B consisted of 80% acetonitrile and 0.1% formic acid. Orbitrap Exploris 480 mass spectrometer was operated in the data-dependent acquisition mode using Xcalibur 4.5.445.18 software, and there was a single full-scan mass spectrum in the orbitrap (350-1600 m/z, 60,000 resolution) followed by 2 seconds data-dependent MS/MS scans in an Ion Routing Multipole at 30 normalized collision energy (HCD).

The MS data were analyzed using the MaxQuant algorithm (v.2.4.9.0). Spectra were searched against the *E. coli* K12 protein sequences in the Uniprot database. The global false discovery rate (FDR) cutoff for peptide and protein identification was set to 0.01. The label-free quantification (LFQ) algorithm in MaxQuant was used to compare samples.

Proteomics statistical analysis was performed using the Perseus algorithm (v.2.0.11.0). LFQ intensities for the protein group table were uploaded to Perseus. Protein groups were filtered to retain only proteins with at least 2 valid values in at least one group. Missing values were replaced from a normal distribution (width = 0.3, down shift =1.8). To compare protein intensities, a *t*-test was performed with the correction for multiple hypotheses by FDR-adjusted $p < 0.05$.

Statistical analysis

Statistical analyses were performed in GraphPad Prism using tests described in the Figure Legends.

Reporting summary

Further information on research design is available in the file Reporting Summary.

Data availability

All data needed to evaluate the conclusions in the paper are present in the file Combined Manuscript and the file Supplementary Information.

ACKNOWLEDGMENTS

We thank Jingjing Wang at the Cell Biology Facility, Center of Biomedical Analysis, Tsinghua University, for help with confocal microscopy. We thank Meng Han in Protein Chemistry and proteomics Facility at Technology Center for Protein Sciences, Tsinghua University, for protein MS analysis. We thank Nathalie Balaban's lab at Racah Institute of Physics, The Hebrew University of Jerusalem, for providing the *E. coli* KLY strain. This work was supported by the National Key Research and Development Program of China (2022YFC2303202), the Tsinghua-Peking Joint Center for Life Sciences (20111770319), and the Tsinghua University Dushi Program (20251080040) to J. L.

AUTHOR CONTRIBUTIONS

J. L. mainly designed experiments, acquired funding, and revised the paper. F. W. designed and performed experiments, analyzed data, and wrote the paper. Y. Z. performed the genome editing experiments, helped with the analysis of clinical isolates, and revised the paper.

COMPETING INTERESTS

The authors declare no competing interests.

MATERIALS & CORRESPONDENCE

Further information and requests for resources and reagents should be directed to, and will be fulfilled by, the lead contact, Jia-Feng Liu (jfliu@mail.tsinghua.edu.cn).

SUPPLEMENTARY INFORMATION

The file Supplementary Information includes Extended Data Figs. 1–9 and Supplementary Tables 1–5.

 FIGURE TITLES AND LEGENDS

Fig. 1. Delayed aggregate disassembly and increased vulnerability to proteotoxic stress in long-lag mutants. a, Appearance time distribution for WT and identified dormant mutants measured by ScanLag. **b**, Representative brightfield and fluorescence images of WT and *pheT^D1* cells stained with Proteostat. Cells from the stationary phase (S phase) were re-cultured in fresh LB medium and sampled at the indicated time points. Arrows indicate Proteostat-positive aggregates. Scale bars, 2 μm. **c**, Quantification of the fraction of WT, *pheT^D1*, and *pheT^D2* cells containing Proteostat-positive aggregates upon re-culturing in fresh LB medium. Data are presented as mean ± SD (n = 4). **d**, Survival of WT, *pheT^D1*, and *pheT^D2* cells after 1 μg/mL ertapenem treatment for 4 h or 12 h. Data are presented as mean ± SD (n = 3). **e**, Survival of WT, *pheT^D1*, and *pheT^D2* cells after oxidative stress (10 mM H_2O_2, 30 minutes) or heat shock (55 °C, 30 minutes). Asterisk indicates values below detection limit (< 10^{-7}). Data are presented as mean ± SD (n = 3). **f**, Growth fraction of WT, *pheT^D1*, and *pheT^D2* cells on solid medium containing the indicated concentrations of puromycin relative to drug-free plates. Data are presented as mean ± SD (n = 3).

Fig. 2. Increased PheS aggregation delays aggregate disassembly and extends lag time. a,b, Volcano plots showing MS analysis of the insoluble proteome in *pheT^D1^* and *pheT^D2^* relative to WT at the stationary phase. Student's *t*-test with the correction to multiple hypotheses by FDR adjusted $p < 0.05$. n = 3 biological replicates. **c,** Representative brightfield and fluorescence images of WT and *pheT^D1^* cells expressing ECFP-tagged PheS and stained with Proteostat. Cells from the stationary phase were re-cultured in fresh LB medium and stained with Proteostat at the indicated time points. Scale bars, 2 μm. **d,** Quantification of the fraction of WT and *pheT^D1^* cells containing Proteostat-positive aggregates during recovery in fresh LB medium. Cells with the inducible expression of PheS-ECFP were grown for 48 hours at the indicated concentrations of IPTG, and then transferred to fresh LB medium for recovery. Data are presented as mean ± SD (n = 4). **e,** Appearance time distribution for WT and *pheT^D1^* cells after the inducible expression of PheS-ECFP for 48 hours at the indicated concentrations of IPTG. **f,** CFU of WT, *pheT^D1^*, and *pheT^D2^* bacteria after the inducible expression of PheS-ECFP for 48 hours at the indicated concentrations of IPTG. Data are presented as mean ± SD (n = 3).

Fig. 3. Sequestration of the replication initiator DnaA within protein aggregates delays replication initiation. a, Fluorescence images of dual-labelled strains (chromosomally labelled HslU-ECFP and mCherry-DnaA) stained with DAPI upon re-culturing. Cells from the stationary phase were re-cultured in fresh LB medium and sampled at the indicated time points. Scale bars, 2 µm. **b**, Quantification of the fraction of WT and *pheT^{D1}* cells containing HslU-ECFP foci or mCherry-DnaA foci upon re-culturing. Data are presented as mean ± SD (n = 4). **c–e**, Appearance time distribution for WT, *pheT^{D1}*, and *pheT^{D2}* cells when incubated with the induced expression of PheS-ECFP (0.15 mM IPTG), HslVU-mCherry (0.25 mM IPTG), or puromycin (16 µg/mL). **f**, Growth fraction of WT, *pheT^{D1}*, and *pheT^{D2}* cells when incubated with the inducible expression of HslVU-mCherry at the indicated concentrations of IPTG relative to IPTG-free plates. Data are presented as mean ± SD (n = 3). **g**, Appearance time distribution for *pheT^{D1}* cells when incubated with the anhydrotetracycline-induced PheS (1 ng/mL anhydrotetracycline), arabinose-induced HslVU-ECFP (100 mM arabinose), or IPTG-inducible soluble DnaA-mCherry (0.25 mM IPTG). **h**, Growth fraction of WT, *pheT^{D1}*, and *pheT^{D2}* cells when incubated with the inducible expression of proteases (0.25 mM IPTG) relative to IPTG-free plates. Data are presented as mean ± SD (n = 3).

Fig. 4. Lag time scales with proteostasis disruption across stresses. a, Quantification of the fraction of stress-treated WT cells containing Proteostat-positive aggregates upon re-culturing. WT cells were treated with the indicated stress, and upon stress removal, bacteria were re-cultured with fresh LB medium and stained with Proteostat at the indicated time points. Data are presented as mean ± SD (n = 4). **b**, Appearance time distribution for WT after treatment with the indicated stress. **c**, Appearance time distribution for WT after exposure to 256 μg/mL puromycin for the indicated durations. **d**, Mean appearance time plotted against the disruption level of proteostasis after treatment with the indicated stress. Each point represents the mean of n = 3 biological replicates for each stress condition. The disruption level of proteostasis is quantified as the $\log_{10}$ ratio of CFU on antibiotic-free versus 64 μg/mL puromycin plates. From left to right: Heat: 49 °C for 1 h, 24 h, or 48 h; 55 °C for 0.5 h or 1 h; Chloramphenicol (Chl): 500 μg/mL for 48 h, 72 h, 96 h; Streptomycin (Str): 50 μg/mL for 2 h; Azithromycin (Azi): 100 μg/mL for 24 h, 48 h; H_2O_2: 5 mM or 10 mM for 0.5 h. The solid line denotes linear regression (R^2 = 0.97, P < 0.0001). **e**, Appearance time plotted against the disruption level of proteostasis for the identified dormant mutants. The disruption level of proteostasis is quantified as the $\log_{10}$ ratio of CFU on antibiotic-free versus 64 μg/mL puromycin plates. The solid line denotes linear regression (R^2 = 0.71, P = 0.017). Data are presented as mean ± SD (n = 3). **f**, Growth fraction of stress-treated WT cells on puromycin plates (64 μg/mL) relative to drug-free plates and survival after 4 h of 1 μg/mL ertapenem treatment. Data are presented as mean ± SD (n = 3).

Fig. 5. Restoration of proteostasis promotes exit from dormancy. a, Representative brightfield and
fluorescence images of WT and *pheT*[D1] bacteria stained with Peroxy Orange 1 (PO-1). Cells from the
stationary phase were re-cultured in LB or LB supplemented with 0.1 mg/mL catalase and sampled at the
indicated time points. Arrows indicate PO-1-positive aggregates. Scale bars, 2 μm. **b**, PO-1 fluorescence
intensity distribution for WT and *pheT*[D1] cells at the indicated time points during recovery. **c**, Representative
fluorescence images of *pheT*[D1] bacteria stained with Proteostat after re-culturing in LB or LB supplemented
with 0.1 mg/mL catalase. Scale bars, 2 μm. **d**, Quantification of the fraction of *pheT*[D1] cells containing
Proteostat-positive aggregates upon re-culturing in LB or LB supplemented with 0.1 mg/mL catalase. Data
are presented as mean ± SD (n = 4). **e**, Growth curves of *pheT*[D1] bacteria were monitored by measuring
optical density at 600 nm (OD_{600}) over time in LB or LB supplemented with 0.1 mg/mL catalase. Data are
presented as mean ± SD (n = 3).

Fig. 6. Proteostasis-governed dormancy in clinical isolates. a, Appearance time distribution for clinical isolate *K. pneumoniae* S837 at the stationary phase (S phase) or exponential phase (E phase), and laboratory strain ATCC 43816 at the S phase. **b**, Representative brightfield and fluorescence images of *K. pneumoniae* ATCC 43816 and clinical *K. pneumoniae* S837 stained with Proteostat. Bacteria from the stationary phase were re-cultured in fresh LB medium and sampled at the indicated time points. Arrows indicate Proteostat-positive aggregates. Scale bars, 2 μm. **c**, Appearance time distribution for *K. pneumoniae* ATCC 43816 (bottom) and *B. subtilis* PY79 (upper) after 256 μg/mL puromycin exposure for the indicated durations. **d**, Growth fraction of S/E-phase *K. pneumoniae* S837 on puromycin plates (64 μg/mL) relative to drug-free plates and survival after 3 h of 1 μg/mL ertapenem treatment. Data are presented as mean ± SD (n = 3). **e**, Formation and disassembly of protein aggregates function as a timing device for bacterial dormancy. Upon entry into dormancy, protein aggregates, such as insoluble PheS, sequester the protease HslVU and the replication initiator DnaA away from the bacterial nucleoid, thereby preventing replication initiation. After nutrient replenishment, bacterial replication initiation remains inhibited until aggregates are disassembled, allowing DnaA to reassociate with the chromosomal origin of replication (*oriC*). The rate of aggregate disassembly thus determines the duration of bacterial dormancy.

REFERENCES

Jonas, K., Liu, J., Chien, P. & Laub, M. T. Proteotoxic stress induces a cell-cycle arrest by stimulating Lon to degrade the replication initiator DnaA. *Cell* **154**, 623-636 (2013). https://doi.org:10.1016/j.cell.2013.06.034

Fridman, O., Goldberg, A., Ronin, I., Shoresh, N. & Balaban, N. Q. Optimization of lag time underlies antibiotic tolerance in evolved bacterial populations. *Nature* **513**, 418-421 (2014). https://doi.org:10.1038/nature13469

Levin-Reisman, I. *et al.* Antibiotic tolerance facilitates the evolution of resistance. *Science* **355**, 826-830 (2017). https://doi.org:10.1126/science.aaj2191

Pu, Y. *et al.* ATP-Dependent Dynamic Protein Aggregation Regulates Bacterial Dormancy Depth Critical for Antibiotic Tolerance. *Mol Cell* **73**, 143-156.e144 (2019). https://doi.org:10.1016/j.molcel.2018.10.022

Blattman, S. B. *et al.* Identification and genetic dissection of convergent persister cell states. *Nature* **636**, 438-446 (2024). https://doi.org:10.1038/s41586-024-08124-2

Liu, J., Gefen, O., Ronin, I., Bar-Meir, M. & Balaban, N. Q. Effect of tolerance on the evolution of antibiotic resistance under drug combinations. *Science* **367**, 200-204 (2020). https://doi.org:10.1126/science.aay3041

Balaban, N. Q., Merrin, J., Chait, R., Kowalik, L. & Leibler, S. Bacterial persistence as a phenotypic switch. *Science* **305**, 1622-1625 (2004). https://doi.org:10.1126/science.1099390

Hansen, S., Lewis, K. & Vulić, M. Role of global regulators and nucleotide metabolism in antibiotic tolerance in *Escherichia coli*. *Antimicrob Agents Chemother* **52**, 2718-2726 (2008). https://doi.org:10.1128/aac.00144-08

Girgis, H. S., Harris, K. & Tavazoie, S. Large mutational target size for rapid emergence of bacterial persistence. *Proc Natl Acad Sci U S A* **109**, 12740-12745 (2012). https://doi.org:10.1073/pnas.1205124109

Yu, J., Liu, Y., Yin, H. & Chang, Z. Regrowth-delay body as a bacterial subcellular structure marking multidrug-tolerant persisters. *Cell Discov* **5**, 8 (2019). https://doi.org:10.1038/s41421-019-0080-3

Dewachter, L. *et al.* The Dynamic Transition of Persistence toward the Viable but Nonculturable State during Stationary Phase Is Driven by Protein Aggregation. *mBio* **12**, e0070321 (2021). https://doi.org:10.1128/mBio.00703-21

Bollen, C. *et al.* Composition and liquid-to-solid maturation of protein aggregates contribute to bacterial dormancy development and recovery. *Nat Commun* **16**, 1046 (2025). https://doi.org:10.1038/s41467-025-56387-8

Zhang, Z., Li, D., Zheng, B. & Liu, J. F. DEAD-box ATPase-marked condensates coordinate compartmentalized translation and antibiotic persistence. *Sci Adv* **12**, eady1930 (2026). https://doi.org:10.1126/sciadv.ady1930

Levin-Reisman, I. *et al.* Automated imaging with ScanLag reveals previously undetectable bacterial growth phenotypes. *Nat Methods* **7**, 737-739 (2010). https://doi.org:10.1038/nmeth.1485

Kaspy, I. *et al.* HipA-mediated antibiotic persistence via phosphorylation of the glutamyl-tRNA-synthetase. *Nat Commun* **4**, 3001 (2013). https://doi.org:10.1038/ncomms4001

Shen, D. *et al.* Novel cell- and tissue-based assays for detecting misfolded and aggregated protein accumulation within aggresomes and inclusion bodies. *Cell Biochem Biophys* **60**, 173-185 (2011). https://doi.org:10.1007/s12013-010-9138-4

17 Leeman, D. S. *et al.* Lysosome activation clears aggregates and enhances quiescent neural stem
cell activation during aging. *Science* **359**, 1277-1283 (2018).
https://doi.org:10.1126/science.aag3048
18 Tomoyasu, T., Mogk, A., Langen, H., Goloubinoff, P. & Bukau, B. Genetic dissection of the roles
of chaperones and proteases in protein folding and degradation in the *Escherichia coli* cytosol.
Mol Microbiol **40**, 397-413 (2001). https://doi.org:10.1046/j.1365-2958.2001.02383.x
19 Gidalevitz, T., Ben-Zvi, A., Ho, K. H., Brignull, H. R. & Morimoto, R. I. Progressive disruption of
cellular protein folding in models of polyglutamine diseases. *Science* **311**, 1471-1474 (2006).
https://doi.org:10.1126/science.1124514
20 Labbadia, J. & Morimoto, R. I. The biology of proteostasis in aging and disease. *Annu Rev*
Biochem **84**, 435-464 (2015). https://doi.org:10.1146/annurev-biochem-060614-033955
21 Hipp, M. S. & Hartl, F. U. Interplay of Proteostasis Capacity and Protein Aggregation: Implications
for Cellular Function and Disease. *J Mol Biol* **436**, 168615 (2024).
https://doi.org:10.1016/j.jmb.2024.168615
22 Mirzaei, H. & Regnier, F. Protein:protein aggregation induced by protein oxidation. *J Chromatogr*
B Analyt Technol Biomed Life Sci **873**, 8-14 (2008).
https://doi.org:10.1016/j.jchromb.2008.04.025
23 Brøndsted, L., Andersen, M. T., Parker, M., Jørgensen, K. & Ingmer, H. The HtrA protease of
Campylobacter jejuni is required for heat and oxygen tolerance and for optimal interaction with
human epithelial cells. *Appl Environ Microbiol* **71**, 3205-3212 (2005).
https://doi.org:10.1128/aem.71.6.3205-3212.2005
24 Servant, P. *et al.* The ClpPX protease is required for radioresistance and regulates cell division
after gamma-irradiation in *Deinococcus radiodurans*. *Mol Microbiol* **66**, 1231-1239 (2007).
https://doi.org:10.1111/j.1365-2958.2007.06003.x
25 Mermershtain, I. *et al.* Idiosyncrasy and identity in the prokaryotic Phe-system: crystal structure
of *E. coli* phenylalanyl-tRNA synthetase complexed with phenylalanine and AMP. *Protein Sci* **20**,
160-167 (2011). https://doi.org:10.1002/pro.549
26 Mahmoud, S. A. & Chien, P. Regulated Proteolysis in Bacteria. *Annu Rev Biochem* **87**, 677-696
(2018). https://doi.org:10.1146/annurev-biochem-062917-012848
27 Joshi, K. K. & Chien, P. Regulated Proteolysis in Bacteria: *Caulobacter*. *Annu Rev Genet* **50**, 423-
445 (2016). https://doi.org:10.1146/annurev-genet-120215-035235
28 Gross, M. H. & Konieczny, I. Polyphosphate induces the proteolysis of ADP-bound fraction of
initiator to inhibit DNA replication initiation upon stress in *Escherichia coli*. *Nucleic Acids Res* **48**,
5457-5466 (2020). https://doi.org:10.1093/nar/gkaa217
29 Gbolahan, A. *et al.* A novel membrane stress response that blocks chromosomal replication by
targeting the DnaA initiator via the ClpP protease. *J Bacteriol* **207**, e0015125 (2025).
https://doi.org:10.1128/jb.00151-25
30 Winkler, J. *et al.* Quantitative and spatio-temporal features of protein aggregation in *Escherichia*
coli and consequences on protein quality control and cellular ageing. *Embo j* **29**, 910-923 (2010).
https://doi.org:10.1038/emboj.2009.412
31 Coquel, A. S. *et al.* Localization of protein aggregation in *Escherichia coli* is governed by diffusion
and nucleoid macromolecular crowding effect. *PLoS Comput Biol* **9**, e1003038 (2013).
https://doi.org:10.1371/journal.pcbi.1003038
32 Ayrapetyan, M., Williams, T. C. & Oliver, J. D. Bridging the gap between viable but non-
culturable and antibiotic persistent bacteria. *Trends Microbiol* **23**, 7-13 (2015).
https://doi.org:10.1016/j.tim.2014.09.004

33 Nozaki, S., Niki, H. & Ogawa, T. Replication initiator DnaA of *Escherichia coli* changes its assembly
form on the replication origin during the cell cycle. *J Bacteriol* **191**, 4807-4814 (2009).
https://doi.org/10.1128/jb.00435-09
34 Boeneman, K. *et al. Escherichia coli* DnaA forms helical structures along the longitudinal cell axis
distinct from MreB filaments. *Mol Microbiol* **72**, 645-657 (2009). https://doi.org/10.1111/j.1365-
2958.2009.06674.x
35 Ferguson, A. D., Hofmann, E., Coulton, J. W., Diederichs, K. & Welte, W. Siderophore-mediated
iron transport: crystal structure of FhuA with bound lipopolysaccharide. *Science* **282**, 2215-2220
(1998). https://doi.org/10.1126/science.282.5397.2215
36 Killmann, H., Herrmann, C., Wolff, H. & Braun, V. Identification of a new site for ferrichrome
transport by comparison of the FhuA proteins of *Escherichia coli*, *Salmonella paratyphi* B,
Salmonella typhimurium, and *Pantoea agglomerans*. J Bacteriol **180**, 3845-3852 (1998).
https://doi.org/10.1128/jb.180.15.3845-3852.1998
37 Dai, Y. *et al.* Interface of biomolecular condensates modulates redox reactions. *Chem* **9**, 1594-
1609 (2023). https://doi.org/10.1016/j.chempr.2023.04.001
38 Chen, M. W. *et al.* Transition-State-Dependent Spontaneous Generation of Reactive Oxygen
Species by Aβ Assemblies Encodes a Self-Regulated Positive Feedback Loop for Aggregate
Formation. *J Am Chem Soc* **147**, 8267-8279 (2025). https://doi.org/10.1021/jacs.4c15532
39 Dickinson, B. C., Huynh, C. & Chang, C. J. A palette of fluorescent probes with varying emission
colors for imaging hydrogen peroxide signaling in living cells. *J Am Chem Soc* **132**, 5906-5915
(2010). https://doi.org/10.1021/ja1014103
40 Lippert, A. R., Van de Bittner, G. C. & Chang, C. J. Boronate oxidation as a bioorthogonal reaction
approach for studying the chemistry of hydrogen peroxide in living systems. *Acc Chem Res* **44**,
793-804 (2011). https://doi.org/10.1021/ar200126t
41 Py, B. & Barras, F. Building Fe-S proteins: bacterial strategies. *Nat Rev Microbiol* **8**, 436-446
(2010). https://doi.org/10.1038/nrmicro2356
42 Alfonso-Prieto, M., Biarnés, X., Vidossich, P. & Rovira, C. The molecular mechanism of the
catalase reaction. *J Am Chem Soc* **131**, 11751-11761 (2009). https://doi.org/10.1021/ja9018572
43 Kohanski, M. A., Dwyer, D. J., Hayete, B., Lawrence, C. A. & Collins, J. J. A common mechanism of
cellular death induced by bactericidal antibiotics. *Cell* **130**, 797-810 (2007).
https://doi.org/10.1016/j.cell.2007.06.049
44 Alberti, S. & Hyman, A. A. Biomolecular condensates at the nexus of cellular stress, protein
aggregation disease and ageing. *Nature Reviews Molecular Cell Biology* **22**, 196-213 (2021).
https://doi.org/10.1038/s41580-020-00326-6
45 Zhang, E. E. & Kay, S. A. Clocks not winding down: unravelling circadian networks. *Nat Rev Mol
Cell Biol* **11**, 764-776 (2010). https://doi.org/10.1038/nrm2995
46 Bollen, C., Dewachter, L. & Michiels, J. Protein Aggregation as a Bacterial Strategy to Survive
Antibiotic Treatment. *Front Mol Biosci* **8**, 669664 (2021).
https://doi.org/10.3389/fmolb.2021.669664
47 Schramm, F. D., Schroeder, K. & Jonas, K. Protein aggregation in bacteria. *FEMS Microbiol Rev*
44, 54-72 (2020). https://doi.org/10.1093/femsre/fuz026
48 Saad, S. *et al.* Reversible protein aggregation is a protective mechanism to ensure cell cycle
restart after stress. *Nat Cell Biol* **19**, 1202-1213 (2017). https://doi.org/10.1038/ncb3600
49 Zaffagnini, G. *et al.* Mouse oocytes sequester aggregated proteins in degradative super-
organelles. *Cell* **187**, 1109-1126.e1121 (2024). https://doi.org/10.1016/j.cell.2024.01.031
50 Audas, T. E. *et al.* Adaptation to Stressors by Systemic Protein Amyloidogenesis. *Dev Cell* **39**,
155-168 (2016). https://doi.org/10.1016/j.devcel.2016.09.002

51   Rosswag de Souza, S., Böke, E. & Zaffagnini, G. Proteostasis in cellular dormancy: lessons from
yeast to oocytes. *Trends Biochem Sci* **50**, 646-662 (2025).
https://doi.org:10.1016/j.tibs.2025.05.004
52   Thomason, L. C., Costantino, N. & Court, D. L. *E. coli* genome manipulation by P1 transduction.
Curr Protoc Mol Biol **Chapter 1**, 1.17.11-11.17.18 (2007).
https://doi.org:10.1002/0471142727.mb0117s79
53   Näsvall, J., Knöppel, A. & Andersson, D. I. Duplication-Insertion Recombineering: a fast and scar-
free method for efficient transfer of multiple mutations in bacteria. *Nucleic Acids Res* **45**, e33
(2017). https://doi.org:10.1093/nar/gkw1078

742