Cognitive reappraisal of food and emotion cues involves common and unique neural

contributions

Julia M. Laing-Young[1,2], Cary R. Savage[1,2], Cara Tomaso[3], Maital Neta [1,2], Timothy D. Nelson [1,2]
& Douglas H. Schultz [1,2]
[1]Center for Brain, Biology and Behavior, University of Nebraska-Lincoln, Lincoln, NE
[2]Department of Psychology, University of Nebraska-Lincoln, Lincoln, NE
[3]Department of Pediatrics, Yale Child Study Center, Yale School of Medicine, New Haven, CT

Abstract

Obesity is a growing public health concern with more than 40% of adults meeting criteria for obesity in the United States. Although many treatments seek to lower individual's weight, few treatments have focused on cognitive strategies to change the way individuals think about food, therefore, decreasing consumption of non-nutrient-dense foods. Cognitive reappraisal is one strategy that involves changing the way one thinks about a situation and can be used to downregulate responses to those stimuli. Leveraging this intuitive, cost-effective strategy to decrease one's desire to eat unhealthy food and therefore, decrease overeating, could improve physical and mental health. The present study identified brain regions that are differentially activated when using cognitive reappraisal to downregulate responses to food (FR) versus when using the same strategy to downregulate negative emotions (ER). We collected functional magnetic resonance imaging (fMRI) data in 63 undergraduate students while participants completed both tasks. There was increased reappraisal-related activation in widespread regions across both tasks, including in expected subcortical (i.e., striatum) and cortical areas (i.e., visual, frontoparietal). We also found domain-specific activity, with greater insula activation in the FR than the ER task and greater hippocampal activation in the ER than the FR task. These results reveal domain-general and domain-specific effects of cognitive reappraisal in FR and ER tasks that inform future work examining eating behavior. Taken together, a better explication of the overlapping and discrete processes of food regulation, as it compares to other applications of this regulatory strategy can inform new intervention targets.

Introduction

Obesity is a growing public health concern. In the United States, more than 40% of adults meet criteria for clinical obesity (body mass index [BMI] of 30 or higher), and an additional 30% of adults are classified as overweight (National Center for Health Statistics, Centers for Disease Control and Prevention, 2024). Obesity is a complex and difficult to treat chronic disease associated with significant health consequences, including diabetes, cardiovascular disease, and even mortality (Koliaki et al., 2023). The obesity epidemic is further fueled by the consequences of economic, social, and technological advances in recent decades which promote increased calorie consumption and reduced physical activity, and place increased demands on self-regulation. For example, palatable foods with high caloric density are easily available in fast food restaurants and prepackaged items (Pi-Sunyer, 2002; Catenacci et al., 2009), the increase in popularity of food delivery services (e.g., DoorDash, UberEats), and Americans are more likely than Europeans to use automobiles for their primary form of transportation, resulting in less walking (Buehler & Hamre, 2014; Buehler & Pucher, 2023). To decrease obesity prevalence, interventions are needed at the individual level, for example, by decreasing individual-level food cravings (i.e., the desire to eat high caloric density, readily available foods).

The ability to regulate food cravings is associated with reduced risk for obesity through lower caloric consumption (Potenza & Grolo, 2014; Sun & Kober, 2021). These reductions in caloric consumption demonstrate the importance of food regulation for physical wellbeing. Cognitive reappraisal is a key regulation strategy used to down-regulate negative emotions and food cravings and is defined as a cognitive strategy that involves deliberately changing how one interprets the meaning or salience of a given stimulus (McRae, Ciesielski, & Gross, 2012a). In the context of negative emotions, cognitive reappraisal may be used to help an individual find

less negative or even identify positive interpretations of an emotional stimulus (e.g., viewing a bruised body as a scene from a movie). In the context of food regulation, the individual might focus on the health consequences of consuming unhealthy foods to decrease their desire for a meal or snack.

Recent functional magnetic resonance imaging (fMRI) research has investigated the neural processes involved in regulatory behavior, specifically food regulation (FR) and emotion regulation (ER). These studies have identified several brain regions that are involved when participants reappraise food stimuli relative to when they passively view food stimuli or are allowed to crave them. This core set of brain regions, including the anterior insula, inferior and middle frontal gyrus, supplementary motor cortex and parietal cortex – components of the cingulo-opercular and frontoparietal networks (Hausman et al., 2021) – have been broadly implicated in cognitive control (Vincent et al., 2008; Zanto et al., 2013). Other work has also linked these regions with dietary self-control (Han et al., 2018). Given that, it is not necessarily clear the extent to which these effects apply to regulation abilities in general rather than being specific to food regulation.

The use of cognitive reappraisal to downregulate responses to negative emotional images is associated with activation in various regions of the prefrontal cortex (PFC; e.g., dorsolateral, ventrolateral, and dorsomedial PFC, the pre-supplemental motor area, and the anterior cingulate cortex; Chambers et al., 2009; Buhle et al., 2014), which are also regions of the cingulo-opercular and frontoparietal networks (Friedman & Robbins, 2021; Sokolowski et al., 2021). A meta-analysis (Zilverstand et al., 2017) found clinical populations consistently demonstrate reduced recruitment of the dorsolateral and ventrolateral PFC when downregulating negative emotions, suggesting there may be a core deficit in this reappraisal process. As noted in the

context of food regulation, these brain regions are broadly implicated in domain-general cognitive control and may not be specific to emotion regulation.

Despite it being unclear whether these regulatory processes involve overlapping or distinct neural processes, food dysregulation and emotion dysregulation are similar in that they are both associated with clinical presentations. The inability to control food cravings is associated with overeating and obesity (Giuliani et al., 2013), and the inability to regulate negative emotions is a core component of mood disorders (Farb et al., 2012). Further, food dysregulation is a common feature of depression, and individuals with difficulty regulating their emotions may use food (i.e., binge-eating or food restriction) as an attempt to cope (Dingermans, Danner, & Parks, 2017). This example illustrates a potential mechanistic link between food and emotion dysregulation. Although FR and ER have been studied independently (Ochsner & Gross, 2005; Pruessner et al., 2020; Wardle, 1988; Kleiman et al., 2016; Davidson & Jones, 2019), minimal research has directly compared their underlying neural mechanisms using a within subjects design. A recent between-subjects meta-analysis (Gerosa et al., 2024) found a large overlap between brain regions involved in downregulating food cravings and emotional stimuli (e.g., dorsolateral and ventrolateral PFC), as described above. There were also domain-specific effects for food cravings, notably activation in the right insula and left inferior frontal gyrus. However, differences in task design across ER and FR limited the interpretation of these comparisons.

To address this gap in the literature, we examined FR and ER using a similar task design in the same set of participants. The aim of this study was to identify the brain regions associated with cognitive reappraisal during both FR and ER, referred to as domain-general hereafter, and brain regions uniquely associated with only one of these tasks, referred to as domain-specific.

We hypothesized that domain-general cognitive reappraisal would recruit activation in cognitive control regions (e.g., dorsolateral PFC) that are affiliated with fronto-parietal and cingulo-opercular networks. Additionally, we hypothesized domain-specific effects in reward-related brain regions, such as the insula, striatum, and orbitofrontal cortex, for the FR task relative to the ER task. Identifying overlapping and unique mechanisms for these regulatory tasks has the potential to inform future intervention strategies, which could in turn improve mental and physical health throughout the lifespan. For example, if FR and ER have distinct mechanisms, this research could provide insight into food dysregulation and emotion dysregulation requiring domain-specific interventions to target health outcomes.

Methods

Participants

Sixty-eight undergraduate students were enrolled from the University of Nebraska-Lincoln's psychology research participant pool and received course credit for participation. Eligibility criteria included being 18 or 19 years old, right-handed, and no MRI contraindications (e.g., metallic implants, claustrophobia, pregnancy). Of the 68 participants enrolled that completed the scanning procedures, 63 were included in the analyses (M_{age} = 19.13*; M_{BMI}* = 24.13; 75% female; 85.9% white, 6.3% Asian American, 4.7% Black, 3.1% did not report; 10.7% Hispanic/Latinx*)*. Neuroimaging data for two participants were unusable due to high motion (>33% of datapoints during tasks were censored), and three participants were excluded due to incomplete data (i.e., missing behavioral ratings on tasks).

Procedures

Participants who met inclusion criteria attended a multi-hour laboratory visit at the University of Nebraska-Lincoln's Center for Brain, Biology, and Behavior (CB3). At the start of

the session, participants confirmed that they had not consumed any food or caffeinated beverages during the four hours prior to their scheduled appointment. Participants then completed the informed consent process with a trained research assistant, as well as having their height and weight measured. Prior to starting the fMRI session, participants were trained on fMRI tasks to ensure they were appropriately using reappraisal strategies. All study procedures were approved by the Institutional Review Board at the University of Nebraska-Lincoln (UNL).

fMRI Tasks

Food Regulation Task: Participants completed a food regulation task adapted from Yokum and Stice (2013), in which they were shown 48 appetitive food images (Figure 1). The task was presented using E-Prime software (version 2.0) (Psychology Software Tools, Inc.). Images were selected from the Open Library of Affective Foods (OLAF; Miccoli et al., 2016), supplemented with images of high calorie foods used in previously published work (Ness et al., 2014). All images were rated ≥6 on a 1-9 craving scale in this previous online pilot study. During the *IMAGINE* condition (24 trials), which was cued with the word "LOOK," participants were instructed to imagine themselves eating the food, including what the food might smell and taste like. During the *REAPPRAISAL* condition (24 trials), which was cued with the word "DECREASE," participants were trained to regulate their responses by thinking about the long-term health consequences of eating the food (e.g., "Eating fried food may lead to future unwanted weight gain"). Participants were encouraged to consider health consequences that are most relevant to their personal health goals. This strategy was selected because of its face validity with appetitive executive control abilities and because it is very similar to other effective strategies used to decrease appetitive responses in earlier work (Yokum & Stice, 2013).

During training, participants verbalized their strategies to the *IMAGINE* condition and to the *REAPPRAISAL* condition to ensure they could quickly and effectively think of negative health consequences without using a non-cognitive strategy (e.g., attention deployment strategies such as looking away from the image). If participants' responses while training revealed other strategies, the research assistant provided corrective feedback until they could successfully use the reappraisal strategy. For each trial, the condition instruction (i.e., "LOOK" or "DECREASE") appeared at the center of the screen for 4 seconds. Then a food image was presented for 7 seconds, followed by a prompt that appeared for 4 seconds asking participants to press a button rating their desire for the food on a 5-point scale (1 = not at all to 5 = very much). A variable intertrial interval (2-4 seconds) separated each trial, during which time participants were instructed to relax and clear their minds. The entire task lasted 12 minutes, and the order in which the trials were presented was pseudo-randomized such that no more than two sequential trials were of the same condition. Conditions were also counterbalanced, and "LOOK" and "DECREASE" stimuli were randomly assigned for each participant.

Emotion Regulation Task: Participants completed an emotion regulation task adapted from McRae et al., 2012 (Figure 1) in which they were shown a total of 60 unique images. The task was presented with E-Prime software (version 2.0) (Psychology Software Tools, Inc.). Images, which were selected from the International Affective Picture System (IAPS; Lang et al., 2008), included 40 negative and highly arousing images taken from prior research (Neta et al., 2022). Each trial began with an instruction screen lasting 4 seconds with either "LOOK" or "DECREASE" written on a green or blue background, respectively. This was followed by a presentation of an image for 7 seconds against the same-colored background as the instruction screen. For the "LOOK" instruction trials, half of the images had a negative valence ("Look

Negative", 20 trials) and half of the images had neutral valence ("Look Neutral", 20 trials); participants were instructed to respond naturally and allow whatever feelings may arise. For the "DECREASE" instruction trials (*REAPPRAISE*; 20 trials), all the images were negative, and participants were instructed to use cognitive reappraisal to reinterpret the content to feel less negative (see more below). Next, a rating screen appeared for 4 seconds and participants were asked to indicate the degree of negative emotion they felt at the end of the image presentation (i.e., after reappraisal or look.) Participants were asked, "How bad do you feel?" on a 5-point scale (1 = not bad at all; 5 = very bad). A variable intertrial interval (2-4 seconds) separated each trial, during which time participants were instructed to relax and clear their minds. The entire task lasted approximately 16 minutes, and as in the FR task, trials were presented pseudo-randomly, and the negative images that were assigned to "LOOK" or "DECREASE" were counterbalanced across participants.

Prior to task onset, participants were given instructions and shown example images not used during the task itself. Participants completed a set of three practice trials (two "DECREASE" and one "LOOK" trial). For the "DECREASE" trials, the experimenter suggested three types of re-interpretations: It is not real (i.e., it's just a scene from a movie, they are pretending), things will improve with time, and things are not as they appear to be. During the training, participants were asked to explain how they reappraised each scene. If participant responses during training revealed that they were not using a reappraisal strategy, the experimenter offered corrective feedback and re-directed the participant to use one of the three re-interpretation strategies provided above.

Figure 1:

Food Regulation Task:

Emotion Regulation Task

Figure 1: Food Regulation Task (adapted from Yokum and Stice (2013)): During the imagine condition (cued with LOOK), participants were instructed to imagine themselves eating the food that appeared in the subsequent image. During the reappraisal condition (cued with DECREASE), participants were instructed to think about the long-term negative health consequences of eating the food that appeared in the subsequent image. After the image disappeared, they rated how much they wanted to eat the food on a 1 (not at all) to 5 (very much) scale. **Emotion Regulation Task** (adapted from McRae et al., 2012): When the "LOOK" cue appeared, participants were instructed to participants were instructed to respond naturally and allow whatever feelings may arise. During the reappraisal condition (cued with DECREASE), participants were instructed to think about three types of re-interpretation: It is not real (i.e., it's just a scene from a movie, they are pretending), things will improve with time, and things are not as they appear to be. After the image disappeared, they rated how much they wanted to eat the food on a 1 (not at all) to 5 (very much) scale.

fMRI image acquisition

fMRI data were collected with a 3.0-T Skyra Siemens scanner with a 64-channel head coil. Prior to functional imaging, high-resolution anatomical scans were acquired, T1 weighted

multi-echo magnetization prepared rapid gradient echo sequence (MP-RAGE; TR = 2530 msec, TE1 = 1.69 msec, TE2 = 3.55 msec, TE3 = 5.41 msec, TE4 = 7.27 msec, matrix size = 256 x 256 mm^2, field of view = 256 mm, slice thickness = 1mm, number of sliced = 176, and TA = 6.03 min). Functional scans used a multi-band factor = 8, accelerated echo planar imaging (EPI) sequence with the following parameters: repetition time (TR) = 1000 msec, time of echo (TE) = 29.80 msec, flip angel, 60 degrees, matrix size = 210 x 210 mm^2, field of view = 210 mm, slice thickness = 2.5 mm, number of sliced = 51, and voxel size = 2.5 mm^3. Throughout the imaging protocol, participants' heads were secured with foam pads and participants wore foam earplugs. Participants provided ratings for their food desire (FR task) and for how badly they felt after viewing the images (ER task) using a Celeritas Fiber Optic Five-Button Response System placed on their right side (Psychological Software Tools, Inc.). Lastly, movement during the tasks was monitored using Framewise Integrated Real-Time MRI Monitoring (FIRMM) software (Dosenbach et al., 2017).

MRI preprocessing

Functional imaging preprocessing was performed using Analysis of Functional Neuroimages (AFNI; Cox, 1996). Anatomical images were segmented using FreeSurfer (Fischl et al., 2004; Desikan et al., 2006). Processing was completed using afni_proc.py and involved despiking, slice time correction, standardization of anatomical images to Montreal Neurological Institute (MNI) template (152_2009), volume registration to the minimum outlier fraction, spatial smoothing using a 4 mm full width at half maximum Gaussian filter, the time course of each voxel scaled to have a mean of 100, and a general linear model (GLM) using the six motion estimates from volume registration as regressors of no interest. We also used up to third-order

polynominals to model baseline and drift. Pairs of volumes where the Euclidean norm of the

motion derivatives exceeded 0.3 were censored and eliminated from further analysis.

To compare the effects of the regulate and look/imagine strategies, blood oxygenation

level-dependent (BOLD) responses for each condition were contrasted. We used the "BLOCK"

basis function to model response functions at the start of each task trial. Specifically, the

"BLOCK" basis function had a duration of 4 seconds for the instruction prompt (i.e., LOOK or

DECREASE cue) and 7 seconds for the images.

Primary data analysis

AFNI's 3dMVM (Chen et al., 2014) was used to conduct a multivariate modeling

analysis with task (FR, ER) and condition (regulate vs. imagine/look) as within-subject variables.

The spatial smoothness of participants' residuals was estimated with the "-acf" option (Cox et al.,

2017). Mean values for the sample were 0.7151, 2.0648, 6.2374, respectively. Next, a cluster-

based approach for multiple comparisons correction was used to identify which regions showed a

significant main effect or interaction. Specifically, AFNI's 3dClustSim was used to generate ten

thousand random maps using the smoothness parameters and thresholded at a voxel-wise $p <$

0.001. The largest surviving cluster from each of these simulations was recorded, and this

distribution was used to estimate the probability of a false positive. Based on these estimations,

we selected a voxel-wise threshold of $p < 0.001$ and a minimum cluster size of eight contiguous

voxels, resulting in a corrected two-tailed alpha of $p < 0.05$. This resulted in a map for the main

effect for task (FR vs. ER), the main effect for condition (regulate vs. imagine/look), and the

interaction.

Finally, post-hoc tests were conducted to interrogate the interaction effects. Using the

previously identified interaction clusters, the means from the look condition were subtracted

from the regulate condition for each task. We categorized each cluster as showing a greater differential activation FR or ER by comparing the absolute value of the 'look – regulate' for each task and assigning it to a respective task based on which mean was furthest away from zero. The absolute value of 'look – regulate' activations for each task were also correlated with a difference score of 'look – regulate' behavioral ratings (i.e., regulation success).

Results

Behavioral results

Participants were instructed, on some trials, to use cognitive reappraisal while viewing food or emotional stimuli, and on other trials, to look at the images allowing themselves to feel what they would naturally feel. As expected, ratings of food cravings on decrease trials were lower than look trials in the FR task ($t = 14.23$, $p = <0.001$, Figure 2) and ratings of negative affect on decrease trials were lower than look trials in the ER task ($t = 14.10$, $p = <0.001$, Figure 2).

Figure 2: Ratings of food cravings for the FR task and ratings of negative affect for the ER task. Ratings on "LOOK" trials were higher than "DECREASE" (Regulate) trials for both FR (*Mlook* = 3.81, *Mdecrease* = 2.21) and ER (*Mlook* = 3.88, *Mdecrase* = 2.34) tasks, with higher ratings corresponding to higher desire that participants had to eating the food (FR) or feeling more negative while viewing the images (ER). *** = p <.001

We operationalized regulation success in each task as the difference in ratings for look minus regulate trials (FR: *M* =1.59, *SD* = 0.90, *range* = 0.08, -3.3; ER: *M* =1.53, *SD* = 0.85, *range* = 0.5, -3.3). There was a strong positive relationship for regulation success between the FR task and ER task (rho= 0.69, rho^2 =0.47, p = <0.001, Figure 3).

Figure 3: There was a strong positive relationship between performance on emotion regulation and food regulation tasks (rho = 0.69, rho^2 =0.47, *p* = <0.001). Difference score was calculated by subtracting look - regulate condition.

fMRI results

Domain-general regulation

We conducted a 2 (task: FR, ER) x 2 (condition: regulate, imagine/look) ANOVA on the imaging data. There was a main effect for condition, revealing greater activation for regulate than look trials in portions of the right insula, right inferior and middle frontal gyrus, and bilateral inferior parietal lobule, consistent with previous work (Carnell et al., 2012; Kumar et al., 2016; Li et al., 2021) (Figure 4 and Table 1).

Figure 4: Domain-general (main effect) for food regulation and emotion regulation tasks.

		MNI Coordinates		
Region	Volume (mm^3)	x	y	z
Regulate > Look				
Rolandic operculum	31797	51	-5	1
SMA	19656	4	-27	53
Inferior parietal	9406	49	29	45
Angular gyrus	8063	51	61	35
Cerebellum	5969	-28	77	-33
Inferior frontal gyrus	5281	-47	-32	-5
Middle cingulate cortex	3609	2	34	33
Inferior temporal gyrus	2438	-52	-6	-30
Inferior parietal lobule	2313	-54	56	39
Middle frontal gyrus	1719	-42	-23	45
Superior frontal gyrus	938	-19	-49	37
Middle temporal gyrus	578	-64	12	-11
Postcentral gyrus	516	51	15	52
Hippocampus	359	-29	23	-11
Lingual gyrus	359	16	100	-7
Middle temporal gyrus	281	-64	32	-1
Calcarine gyrus	203	12	75	10
Precuneus	172	11	40	4

Middle temporal gyrus	141	-51	15	-11
Caudate nucleus	125	14	-9	15

Look > Regulate

Left rolandic operculum	256	43	2	9
Right postcentral gyrus	130	-61	7	17
Right insula lobe	123	-40	-4	6
Right lingual gyrus	118	-7	82	-3
Right inferior parietal lobule	113	-37	48	44
Left SMA	74	6	8	55
Right cerebellum	59	-23	53	-23
Left middle occipital gyrus	45	32	79	17
Left precentral gyrus	28	60	-3	35
Left superior parietal lobule	27	345	51	61
Right precentral gyrus	26	-56	-7	35
Right cerebellum	23	-24	52	-50
Right cerebellum	18	-13	69	-47
Right caudate nucleus	17	-5	-16	14
Left superior occipital gyrus	15	26	66	35
Left precuneus	15	14	60	61
Right SMA	15	-14	3	64
Left cerebellum cortex	13	21	66	-47
Right superior occipital gyrus	13	-28	78	24
Left SMA	13	9	-3	44

Right supra-marginal gyrus	13	-60	26	45
Left fusiform gyrus	8	38	44	-8
Right middle occipital gyrus	8	-32	73	38

Table 1: Significant clusters associated with domain-general regulation (main effect) for food regulation and emotion regulation tasks. Note: MNI coordinates are listed as x = RL, y = AP, z = IS (center of mass)

Domain-specific regulation

In addition to the main effect of condition, there was also a significant task x condition interaction. Post-hoc tests of the interaction were conducted by subtracting the imagine/look condition from the regulate condition for each task (Figure 5 and Table 2). Interaction clusters that showed the greatest differential activation for FR > ER included the posterior insula and the right inferior frontal gyrus (Hollmann et al., 2012; Gerosa et al., 2023), whereas the reverse (ER > FR) was evident in the left posterior cingulate cortex and right superior frontal gyrus, consistent with previous research (Goldin et al., 2008; Grecucci et al., 2013) (Figure 4).

z = 0

z = 5

z = 10

z = 15

z = 20

Figure 5: Regions that show greater differential regulation for domain-specific emotion regulation (yellow), domain-specific food regulation (blue), and the domain-general regulation main effect with no interaction (green).

		MNI Coordinates		
Region	Volume (mm^3)	x	y	z
Emotion > Food				
Inferior occipital gyrus	4578	35	84	-6
Calcarine gyrus	2484	-7	65	11
Cerebellum	1609	33	50	-18
Superior frontal gyrus	406	-23	-34	47
Calcarine gyrus	328	12	59	7
Middle orbital gyrus	250	-38	-45	-10
Superior frontal gyrus	219	-25	-15	51
Hippocampus	203	23	26	-7
Superior frontal gyrus	188	-21	-27	58
Hippocampus	156	-22	26	-6
Lingual gyrus	156	-18	96	-2
Cerebellum	141	14	45	-48
Food > Emotion				
Middle cingulate cortex	1641	1	25	30
Cerebellum	703	-26	40	-18
Rolandic operculum	547	41	5	10
Inferior frontal gyrus	500	-49	-41	8
Inferior occipital gyrus	469	-32	92	-4
Cuneus	453	10	74	38

Middle temporal gyrus	406	-48	64	23
Insula lobe	359	-40	1	5
Middle temporal gyrus	203	-59	54	3
Postcentral gyrus	188	-64	13	29
Postcentral gyrus	172	-53	1	30
Cuneus	172	-13	68	38
Inferior frontal gyrus	156	45	-25	-6
Cerebellum	141	-28	78	-36
Cerebellum	141	-21	87	-30
Inferior occipital gyrus	141	-46	77	-3
Posterior cingulate cortex	140	7	43	26
Insula lobe	125	41	-1	-8
Inferior frontal gyrus	125	-52	-16	24
Angular gyrus	125	38	68	49

Table 2: Domain-specific significant clusters associated with the task x condition interaction. Note: MNI coordinates are listed as x = RL, y = AP, z = IS (center of mass)

Brain-behavior relationships

We examined the relationship between regulation success and domain-specific brain activation by correlating the difference scores of look > regulate related activity in the interaction clusters with our measure of regulation success (i.e., the difference in ratings between look and regulate for each respective task). There were no statistically significant relationships between regulation success and domain-specific brain activation. We also examined the relationship

between regulation success and domain-general brain activation by correlating the difference scores of look > regulate related activity in regions showing a main effect for regulation, excluding voxels showing an interaction (green regions in Figure 5). There were no statistically significant relationships between regulation success and domain-general brain activation.

Discussion

The present study used a within-subjects design to identify overlapping and distinct neural mechanisms for cognitive reappraisal in the context of FR and ER. There was domain-general activation in the right insula, right inferior and middle frontal gyrus, and bilateral inferior parietal lobule. There was domain-specific activation for the FR task in the posterior insula and for the ER task in the left posterior cingulate cortex and right superior frontal gyrus. Although previous studies have investigated the use of cognitive reappraisal in FR and ER separately and the present results are consistent with these findings, the present study makes novel contributions by identifying overlapping and unique regulatory mechanisms across these tasks using a within-subjects design. This research provides important insight into domain-general and domain-specific effects of regulation that could theoretically inform interventions for emotion and food dysregulation.

In both the FR and ER tasks, participants had decreased ratings in the "regulate" condition relative to the "look" condition, demonstrating successful ability to use the cognitive reappraisal strategy. Additionally, there was a strong relationship between regulation success on both tasks such that participants who were able to effectively regulate emotion cues were also able to effectively regulate food cues. However, there was no significant relationship between regulation success and brain activation (i.e., activation did not vary as a function of how successfully participants performed the respective tasks). Due to task performance on one task

explaining about 47% of the variance in performance on the second task, it's important to consider both domain-general and domain-specific cognitive control since these results indicate that, while there may be some overlap between regulatory cognitive processes (i.e., domain-general), there are also differences (i.e., domain-specific). This set of findings provides further evidence that a core set of brain regions are associated with cognitive control across domains, and there are also distinct brain regions involved in regulation within specific task domains.

Our results are consistent with previous between-subjects studies of cognitive reappraisal that show overlapping neural activation for FR and ER. The findings that the right inferior and middle frontal gyrus and the bilateral inferior parietal lobe are involved in domain-general regulation are consistent with previous findings that brain networks involving the PFC (e.g., fronto-parietal and cingulo-opercular networks) are critical for regulatory behavior such as inhibition and attentional control (Chambers et al., 2009; Friedman & Robbins, 2021; Menon & Esposito, 2021; Zanto & Gazzaley, 2014), and that the parietal cortex is involved in initiating regulatory behavior (Chen et al., 2022). Activation in control brain networks appear to be critical in regulatory processes, regardless of whether participants are regulating responses to food or negative emotions. This finding suggests that there is an overlapping mechanism supporting regulation that is required across various domains or instances of reappraisal.

The evidence for an underlying regulation mechanism has potentially important implications for intervention. Indeed, cognitive control is critical to being able to regulate one's food cravings and negative emotions. This study highlights similarities in these two regulatory processes suggesting that one intervention could potentially target and improve both food- and emotion-related outcomes. For example, current research has shown that 20-30 minutes of moderate-intensity aerobic exercise, which increases functional connectivity among control-

related brain regions (e.g., left dorsolateral PFC, posterior cingulate cortex, and the right inferior parietal cortex), may improve cognitive control across the lifespan (Themanson, Pontifex & Hillman, 2008, Hillman et al., 2009, Weng et al., 2017). In the present study, the dorsolateral PFC and posterior cingulate cortex showed domain-specific activation for FR and ER, respectively. Given the role of cognitive control in regulatory behavior, and the overlap of activation found in studies of aerobic exercise and of FR and ER, aerobic exercise is one area of potential future research for intervention to improve domain-general regulation.

Above and beyond the overlapping neural mechanism, differences were also observed between these two tasks, suggesting some aspects of regulation are domain-specific. Consistent with our findings, previous neuroimaging studies have described activation in brain regions associated with reward processing and the sensory perception of food-regulated stimuli, such as the insula, during the cognitive reappraisal of food (Frank et al., 2013; Kumar et al., 2016). We also observed greater differential activation in the left posterior cingulate cortex for emotion regulation, consistent with literature showing the involvement of the posterior cingulate cortex to the processing of emotions (Rolls, 2019), autobiographical memory (Auger & Maguire, 2013; Leech & Sharp, 2014), altering responses based on task demands (Pearson et al., 2011), and self-regulatory behavior (Kirlic et al., 2022).

Some limitations of the study should be noted. First, average participant BMI fell within the healthy weight range, which makes it unclear if there would be unique regulatory behavior responses (particularly for FR) for participants that fell in the underweight, overweight, or obese BMI categorizations. Second, although participants were instructed to either passively look at the images or use a cognitive reappraisal technique while viewing images in both tasks, the instructions for the FR task specifically instructed participants to reappraise by focusing on the

health consequences of eating the desirable food, whereas the ER task instructions were more open-ended about how to apply reappraisal. The greater specificity in the FR instructions may have influenced the differences in brain mechanisms between the two tasks. A future study could either narrow the ER instructions to a single approach or broaden the FR task instructions to cognitively reappraise the food stimuli to make it less appetizing (rather than relating it to health consequences of eating the food). Third, although we did not find a statistically significant relationship between regulatory success and brain activation, the literature suggests that these behaviors are related to success during the ER task. One possible explanation for the lack of replicating the behavior-brain relationship is due to our limited sample size. Lastly, participants were undergraduate students enrolled in a psychology course, so it is unclear to what extent these findings would generalize to other samples.

The current study builds on the existing literature by identifying overlapping brain regions that are associated with cognitive reappraisal during FR and ER tasks, and brain regions that are differentially associated with these two regulatory processes. The existing literature has previously examined these regulatory processes separately, and the current study results expand on these findings in a within-subjects context, paving the way for future research identifying and testing interventions for related mental and physical health outcomes. Future research should determine the extent to which these domain-general and domain-specific mechanisms are also evident in clinically relevant populations (e.g., individuals suffering from food and/or emotion dysregulation, such as those who are overweight or obese, or those presenting with mental health problems).

Author CRediT Statement

J.L., C.S., M.N., T.N., and D.S. contributed to the conceptualization of the project; T.N. acquired funding and resources for the project, and oversaw the investigation process and data collection; J.L. and C.T. contributed to the data curation; J.L. performed the formal analysis, visualization of data, and took the lead in writing the manuscript; J.L., T.N., and D.S. contributed to the methodology; D.S. provided supervision to all aspects of manuscript preparation (i.e., data analysis interpretation and revisions for original draft prior to sending to other co-authors); M.N. and D.S. provided software (i.e., E-Prime tasks) for the project; C.S., M.N., C.T., T.N., and D.S. reviewed and edited the manuscript.

Data Availability Statement

The data for this project cannot be shared publicly due to participants not consenting to having their data shared at the time of study participation. Please contact the corresponding author, Douglas Schultz, with additional questions on data availability.

Acknowledgements

This research was supported by the National Institute of General Medical Sciences (U54GM115458) and the National Institute of Diabetes and Digestive and Kidney Diseases (R01DK125651, F31DK122636) of the National Institutes of Health (NIH). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Declaration of Interest

Declarations of interest: none

References

Assem, M., Glasser, M. F., Van Essen, D. C., & Duncan, J. (2020). A Domain-General

    Cognitive Core Defined in Multimodally Parcellated Human Cortex. *Cerebral Cortex*,

    30(8), 4361–4380. https://doi.org/10.1093/cercor/bhaa023

Auger, S. D., & Maguire, E. A. (2013). Assessing the mechanism of response in the retrosplenial

    cortex of good and poor navigators. *Cortex*, *49*(10), 2904–2913.

    https://doi.org/10.1016/j.cortex.2013.08.002

Botvinick, M. M., Cohen, J. D., & Carter, C. S. (2004). Conflict monitoring and anterior

    cingulate cortex: An update. *Trends in Cognitive Sciences*, *8*(12), 539–546.

    https://doi.org/10.1016/j.tics.2004.10.003

Buhle, J. T., Silvers, J. A., Wager, T. D., Lopez, R., Onyemekwu, C., Kober, H., Weber, J., &

    Ochsner, K. N. (2014). Cognitive reappraisal of emotion: A meta-analysis of human

    neuroimaging studies. *Cerebral Cortex (New York, N.Y.: 1991)*, *24*(11), 2981–2990.

    https://doi.org/10.1093/cercor/bht154

Cameron, E. A., Kwiatkowski, K. J., Lee, B.-H., Hamaker, B. R., Koropatkin, N. M., & Martens,

    E. C. (2014). Multifunctional nutrient-binding proteins adapt human symbiotic bacteria

    for glycan competition in the gut by separately promoting enhanced sensing and

    catalysis. *mBio*, *5*(5), e01441-01414. https://doi.org/10.1128/mBio.01441-14

Chambers, C. D., Garavan, H., & Bellgrove, M. A. (2009). Insights into the neural basis of

    response inhibition from cognitive and clinical neuroscience. *Neuroscience &*

    Biobehavioral Reviews, *33*(5), 631–646. https://doi.org/10.1016/j.neubiorev.2008.08.016

Chen, G., Adleman, N. E., Saad, Z. S., Leibenluft, E., & Cox, R. W. (2014). Applications of

    multivariate modeling to neuroimaging group analysis: A comprehensive alternative to

univariate general linear model. *NeuroImage*, *99*, 571–588.

https://doi.org/10.1016/j.neuroimage.2014.06.027

Chen, X., Zirnsak, M., Vega, G. M., Govil, E., Lomber, S. G., & Moore, T. (2020). Parietal

Cortex Regulates Visual Salience and Salience-Driven Behavior. *Neuron*, *106*(1), 177-

187.e4. https://doi.org/10.1016/j.neuron.2020.01.016

Cole, M. W., Braver, T. S., & Meiran, N. (2017). The task novelty paradox: Flexible control of

inflexible neural pathways during rapid instructed task learning. *Neuroscience &*

Biobehavioral Reviews, *81*, 4–15. https://doi.org/10.1016/j.neubiorev.2017.02.009

Cole, M. W., Reynolds, J. R., Power, J. D., Repovs, G., Anticevic, A., & Braver, T. S. (2013).

Multi-task connectivity reveals flexible hubs for adaptive task control. *Nature*

Neuroscience, *16*(9), 1348–1355. https://doi.org/10.1038/nn.3470

Cox, R. W. (1996). AFNI: Software for analysis and visualization of functional magnetic

resonance neuroimages. *Computers and Biomedical Research, an International Journal*,

29(3), 162–173. https://doi.org/10.1006/cbmr.1996.0014

Dalley, J. W., Everitt, B. J., & Robbins, T. W. (2011). Impulsivity, Compulsivity, and Top-Down

Cognitive Control. *Neuron*, *69*(4), 680–694. https://doi.org/10.1016/j.neuron.2011.01.020

Daniel, R., & Pollmann, S. (2014). A universal role of the ventral striatum in reward-based

learning: Evidence from human studies. *Neurobiology of Learning and Memory*, *114*, 90–

100. https://doi.org/10.1016/j.nlm.2014.05.002

Davidson, T. L., Jones, S., Roy, M., & Stevenson, R. J. (2019). The Cognitive Control of Eating

and Body Weight: It's More Than What You "Think." *Frontiers in Psychology*, *10*, 62.

https://doi.org/10.3389/fpsyg.2019.00062

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner, R.

L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. (2006). An

automated labeling system for subdividing the human cerebral cortex on MRI scans into

gyral based regions of interest. *NeuroImage*, *31*(3), 968–980.

https://doi.org/10.1016/j.neuroimage.2006.01.021

Diamond, A. (2013). Executive functions. *Annual Review of Psychology*, *64*, 135–168.

https://doi.org/10.1146/annurev-psych-113011-143750

Dingemans, A., Danner, U., & Parks, M. (2017). Emotion Regulation in Binge Eating Disorder:

A Review. *Nutrients*, *9*(11), 1274. https://doi.org/10.3390/nu9111274

Dosenbach, N. U. F., Koller, J. M., Earl, E. A., Miranda-Dominguez, O., Klein, R. L., Van, A.

N., Snyder, A. Z., Nagel, B. J., Nigg, J. T., Nguyen, A. L., Wesevich, V., Greene, D. J.,

& Fair, D. A. (2017). Real-time motion analytics during brain MRI improve data quality

and reduce costs. *NeuroImage*, *161*, 80–93.

https://doi.org/10.1016/j.neuroimage.2017.08.025

Farb, N. A. S., Anderson, A. K., & Segal, Z. V. (2012). The Mindful Brain and Emotion

Regulation in Mood Disorders. *The Canadian Journal of Psychiatry*, *57*(2), 70–77.

https://doi.org/10.1177/070674371205700203

Fischl, B. (2012). FreeSurfer. *NeuroImage*, *62*(2), 774–781.

https://doi.org/10.1016/j.neuroimage.2012.01.021

Fischl, B., van der Kouwe, A., Destrieux, C., Halgren, E., Ségonne, F., Salat, D. H., Busa, E.,

Seidman, L. J., Goldstein, J., Kennedy, D., Caviness, V., Makris, N., Rosen, B., & Dale,

A. M. (2004). Automatically parcellating the human cerebral cortex. *Cerebral Cortex

(New York, N.Y.: 1991)*, *14*(1), 11–22. https://doi.org/10.1093/cercor/bhg087

Frank, S., Kullmann, S., & Veit, R. (2013). Food related processes in the insular cortex.

    Frontiers in Human Neuroscience, *7*. https://doi.org/10.3389/fnhum.2013.00499

Friedman, N. P., & Robbins, T. W. (2022). The role of prefrontal cortex in cognitive control and

    executive function. *Neuropsychopharmacology*, *47*(1), 72–89.

    https://doi.org/10.1038/s41386-021-01132-0

Gerosa, M., Canessa, N., Morawetz, C., & Mattavelli, G. (2024). Cognitive reappraisal of food

    craving and emotions: A coordinate-based meta-analysis of fMRI studies. *Social*

    Cognitive and Affective Neuroscience, *19*(1), nsad077.

    https://doi.org/10.1093/scan/nsad077

Giuliani, N. R., & Berkman, E. T. (2015). Craving is an affective state and its regulation can be

    understood in terms of the extended process model of emotion regulation. *Psychological*

    Inquiry, *26*(1), 48–53. https://doi.org/10.1080/1047840X.2015.955072

Giuliani, N. R., & Pfeifer, J. H. (2015). Age-related changes in reappraisal of appetitive cravings

    during adolescence. *NeuroImage*, *108*, 173–181.

    https://doi.org/10.1016/j.neuroimage.2014.12.037

Goschke, T. (2014). Dysfunctions of decision-making and cognitive control as transdiagnostic

    mechanisms of mental disorders: Advances, gaps, and needs in current research.

    International Journal of Methods in Psychiatric Research, *23*(S1), 41–57.

    https://doi.org/10.1002/mpr.1410

Grahek, I., Shenhav, A., Musslick, S., Krebs, R. M., & Koster, E. H. W. (2019). Motivation and

    cognitive control in depression. *Neuroscience & Biobehavioral Reviews*, *102*, 371–381.

    https://doi.org/10.1016/j.neubiorev.2019.04.011

Grecucci, A., Giorgetta, C., Van'T Wout, M., Bonini, N., & Sanfey, A. G. (2013). Reappraising the Ultimatum: An fMRI Study of Emotion Regulation and Decision Making. *Cerebral Cortex*, *23*(2), 399–410. https://doi.org/10.1093/cercor/bhs028

Gross, J. J. (2001). Emotion regulation in adulthood: Timing is everything. *Current Directions in Psychological Science*, *10*(6), 214–219. https://doi.org/10.1111/1467-8721.00152

Gross, J. J. (2015). Emotion regulation: Current status and future prospects. *Psychological Inquiry*, *26*(1), 1–26. https://doi.org/10.1080/1047840X.2014.940781

Hall, K. D., Farooqi, I. S., Friedman, J. M., Klein, S., Loos, R. J. F., Mangelsdorf, D. J., O'Rahilly, S., Ravussin, E., Redman, L. M., Ryan, D. H., Speakman, J. R., & Tobias, D. K. (2022). The energy balance model of obesity: Beyond calories in, calories out. *The American Journal of Clinical Nutrition*, *115*(5), 1243–1254. https://doi.org/10.1093/ajcn/nqac031

Han, W., Tellez, L. A., Perkins, M. H., Perez, I. O., Qu, T., Ferreira, J., Ferreira, T. L., Quinn, D., Liu, Z.-W., Gao, X.-B., Kaelberer, M. M., Bohórquez, D. V., Shammah-Lagnado, S. J., de Lartigue, G., & de Araujo, I. E. (2018). A Neural Circuit for Gut-Induced Reward. *Cell*, *175*(3), 665-678.e23. https://doi.org/10.1016/j.cell.2018.08.049

Heatherton, T. F. (2011). Neuroscience of Self and Self-Regulation. *Annual Review of Psychology*, *62*(1), 363–390. https://doi.org/10.1146/annurev.psych.121208.131616

Heatherton, T. F., & Wagner, D. D. (2011). Cognitive Neuroscience of Self-Regulation Failure. *Trends in Cognitive Sciences*, *15*(3), 132–139. https://doi.org/10.1016/j.tics.2010.12.005

Hillman, C. H., Pontifex, M. B., Raine, L. B., Castelli, D. M., Hall, E. E., & Kramer, A. F. (2009). The effect of acute treadmill walking on cognitive control and academic

achievement in preadolescent children. *Neuroscience*, *159*(3), 1044–1054.

https://doi.org/10.1016/j.neuroscience.2009.01.057

Hollmann, M., Hellrung, L., Pleger, B., Schlögl, H., Kabisch, S., Stumvoll, M., Villringer, A., &
Horstmann, A. (2012). Neural correlates of the volitional regulation of the desire for
food. *International Journal of Obesity*, *36*(5), 648–655.
https://doi.org/10.1038/ijo.2011.125

Kirlic, N., Cohen, Z. P., Tsuchiyagaito, A., Misaki, M., McDermott, T. J., Aupperle, R. L.,
Stewart, J. L., Singh, M. K., Paulus, M. P., & Bodurka, J. (2022). Self-regulation of the
posterior cingulate cortex with real-time fMRI neurofeedback augmented mindfulness
training in healthy adolescents: A nonrandomized feasibility study. *Cognitive, Affective,
& Behavioral Neuroscience*, *22*(4), 849–867. https://doi.org/10.3758/s13415-022-00991-
4

Kleiman, T., Trope, Y., & Amodio, D. M. (2016). Cognitive control modulates attention to food
cues: Support for the control readiness model of self-control. *Brain and Cognition*, *110*,
94–101. https://doi.org/10.1016/j.bandc.2016.04.006

Kober, H., Mende-Siedlecki, P., Kross, E. F., Weber, J., Mischel, W., Hart, C. L., & Ochsner, K.
N. (2010). Prefrontal–striatal pathway underlies cognitive regulation of craving.
Proceedings of the National Academy of Sciences, *107*(33), 14811–14816.
https://doi.org/10.1073/pnas.1007779107

Kumar, S., Grundeis, F., Brand, C., Hwang, H.-J., Mehnert, J., & Pleger, B. (2016). Differences
in Insula and Pre-/Frontal Responses during Reappraisal of Food in Lean and Obese
Humans. *Frontiers in Human Neuroscience*, *10*.
https://doi.org/10.3389/fnhum.2016.00233

Lang, P. J., Bradley, M. M., Cuthbert, B. N., & Center for the Study of Emotion and Attention. (2020). *International Affective Picture System* [Dataset]. https://doi.org/10.1037/t66667-000

Leech, R., & Sharp, D. J. (2014). The role of the posterior cingulate cortex in cognition and disease. *Brain*, *137*(1), 12–32. https://doi.org/10.1093/brain/awt162

Lesh, T. A., Niendam, T. A., Minzenberg, M. J., & Carter, C. S. (2011). Cognitive Control Deficits in Schizophrenia: Mechanisms and Meaning. *Neuropsychopharmacology*, *36*(1), 316–338. https://doi.org/10.1038/npp.2010.156

Lewis, M., & Haviland-Jones, J. M. (Eds.). (2000). *Handbook of emotions, 2nd ed* (pp. xvi, 720). The Guilford Press.

Lewis, R. G., Florio, E., Punzo, D., & Borrelli, E. (2021). The Brain's Reward System in Health and Disease. *Advances in Experimental Medicine and Biology*, *1344*, 57–69. https://doi.org/10.1007/978-3-030-81147-1_4

Marche, K., Martel, A.-C., & Apicella, P. (2017). Differences between Dorsal and Ventral Striatum in the Sensitivity of Tonically Active Neurons to Rewarding Events. *Frontiers in Systems Neuroscience*, *11*, 52. https://doi.org/10.3389/fnsys.2017.00052

Matthews, M., Webb, T. L., Shafir, R., Snow, M., & Sheppes, G. (2021). Identifying the determinants of emotion regulation choice: A systematic review with meta-analysis. *Cognition and Emotion*, *35*(6), 1056–1084. https://doi.org/10.1080/02699931.2021.1945538

McRae, K., Ciesielski, B., & Gross, J. J. (2012). Unpacking cognitive reappraisal: Goals, tactics, and outcomes. *Emotion (Washington, D.C.)*, *12*(2), 250–255. https://doi.org/10.1037/a0026351

McRae, K., & Gross, J. J. (2020). Emotion regulation. *Emotion (Washington, D.C.)*, *20*(1), 1–9.

> https://doi.org/10.1037/emo0000703

McRae, K., Gross, J. J., Weber, J., Robertson, E. R., Sokol-Hessner, P., Ray, R. D., Gabrieli, J.

> D. E., & Ochsner, K. N. (2012). The development of emotion regulation: An fMRI study

> of cognitive reappraisal in children, adolescents and young adults. *Social Cognitive and*

> *Affective Neuroscience*, *7*(1), 11–22. https://doi.org/10.1093/scan/nsr093

Menefee, D. S., Ledoux, T., & Johnston, C. A. (2022). The Importance of Emotional Regulation

> in Mental Health. *American Journal of Lifestyle Medicine*, *16*(1), 28–31.

> https://doi.org/10.1177/15598276211049771

Menon, V., & D'Esposito, M. (2022). The role of PFC networks in cognitive control and

> executive function. *Neuropsychopharmacology*, *47*(1), 90–103.

> https://doi.org/10.1038/s41386-021-01152-w

Miccoli, L., Delgado, R., Guerra, P., Versace, F., Rodríguez-Ruiz, S., & Fernández-Santaella, M.

> C. (2016). Affective Pictures and the Open Library of Affective Foods (OLAF): Tools to

> Investigate Emotions toward Food in Adults. *PLoS ONE*, *11*(8), e0158991.

> https://doi.org/10.1371/journal.pone.0158991

Milyavsky, M., Webber, D., Fernandez, J. R., Kruglanski, A. W., Goldenberg, A., Suri, G., &

> Gross, J. J. (2019). To reappraise or not to reappraise? Emotion regulation choice and

> cognitive energetics. *Emotion*, *19*(6), 964–981. https://doi.org/10.1037/emo0000498

Moccia, L., Pettorruso, M., De Crescenzo, F., De Risio, L., di Nuzzo, L., Martinotti, G., Bifone,

> A., Janiri, L., & Di Nicola, M. (2017). Neural correlates of cognitive control in gambling

> disorder: A systematic review of fMRI studies. *Neuroscience & Biobehavioral Reviews*,

> *78*, 104–116. https://doi.org/10.1016/j.neubiorev.2017.04.025

Munakata, Y., Herd, S. A., Chatham, C. H., Depue, B. E., Banich, M. T., & O'Reilly, R. C. (2011). A unified framework for inhibitory control. *Trends in Cognitive Sciences*, *15*(10), 453–459. https://doi.org/10.1016/j.tics.2011.07.011

Ness, A., Bruce, J., Bruce, A., Aupperle, R., Lepping, R., Martin, L., . . . Savage, C. R. (2014). Pre-surgical cortical activation to food pictures is associated with weight loss following bariatric surgery. Surgery for Obesity and Related Diseases, 10(6), 1188–1195. https://doi.org/10.1016/j.soard.2014.06.005

Ochsner, K. N., & Gross, J. J. (2005). The cognitive control of emotion. *Trends in Cognitive Sciences*, *9*(5), 242–249. https://doi.org/10.1016/j.tics.2005.03.010

Park, J.-H. (2022). Effects of acute moderate-intensity aerobic exercise on executive function and prefrontal cortex activity in community-dwelling older adults: A single-blind, randomized controlled trial. *Geriatrics & Gerontology International*, *22*(3), 227–232. https://doi.org/10.1111/ggi.14352

Pearson, J. M., Heilbronner, S. R., Barack, D. L., Hayden, B. Y., & Platt, M. L. (2011). Posterior cingulate cortex: Adapting behavior to a changing world. *Trends in Cognitive Sciences*, *15*(4), 143–151. https://doi.org/10.1016/j.tics.2011.02.002

PhD, R. S. L., & PhD, S. F. (1984). *Stress, Appraisal, and Coping*. Springer Publishing Company.

Potenza, M. N., & Grilo, C. M. (2014). How Relevant is Food Craving to Obesity and Its Treatment? *Frontiers in Psychiatry*, *5*. https://doi.org/10.3389/fpsyt.2014.00164

Prefrontal–striatal pathway underlies cognitive regulation of craving | PNAS. (n.d.). Retrieved July 27, 2024, from https://www.pnas.org/doi/full/10.1073/pnas.1007779107

Pruessner, L., Barnow, S., Holt, D. V., Joormann, J., & Schulze, K. (2020). A cognitive control

framework for understanding emotion regulation flexibility. *Emotion*, *20*(1), 21–29.

https://doi.org/10.1037/emo0000658

Ravichandran, S., Bhatt, R. R., Pandit, B., Osadchiy, V., Alaverdyan, A., Vora, P., Stains, J.,

Naliboff, B., Mayer, E. A., & Gupta, A. (2021). Alterations in reward network functional

connectivity are associated with increased food addiction in obese individuals. *Scientific

Reports*, *11*(1), 3386. https://doi.org/10.1038/s41598-021-83116-0

Reader, S. W., Lopez, R. B., & Denny, B. T. (2018). Cognitive reappraisal of low-calorie food

predicts real-world craving and consumption of high- and low-calorie foods in daily life.

Appetite, *131*, 44–52. https://doi.org/10.1016/j.appet.2018.08.036

Rolls, E. T. (2019). The cingulate cortex and limbic systems for emotion, action, and memory.

Brain Structure and Function, *224*(9), 3001–3018. https://doi.org/10.1007/s00429-019-

01945-2

Sander, D., & Nummenmaa, L. (2021). Reward and emotion: An affective neuroscience

approach. *Current Opinion in Behavioral Sciences*, *39*, 161–167.

https://doi.org/10.1016/j.cobeha.2021.03.016

Scharmüller, W., Übel, S., Ebner, F., & Schienle, A. (2012). Appetite regulation during food cue

exposure: A comparison of normal-weight and obese women. *Neuroscience Letters*,

518(2), 106–110. https://doi.org/10.1016/j.neulet.2012.04.063

Scherer, K. R. (2009). Emotions are emergent processes: They require a dynamic computational

architecture. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

364(1535), 3459–3474. https://doi.org/10.1098/rstb.2009.0141

Schimmelpfennig, J., Topczewski, J., Zajkowski, W., & Jankowiak-Siuda, K. (2023). The role of the salience network in cognitive and affective deficits. *Frontiers in Human Neuroscience*, *17*. https://doi.org/10.3389/fnhum.2023.1133367

Schultz, D. H., Ito, T., & Cole, M. W. (2022). Global connectivity fingerprints predict the domain generality of multiple-demand regions. *Cerebral Cortex*, *32*(20), 4464–4479. https://doi.org/10.1093/cercor/bhab495

Sheppes, G., Scheibe, S., Suri, G., & Gross, J. J. (2011). Emotion-Regulation Choice. *Psychological Science*, *22*(11), 1391–1396. https://doi.org/10.1177/0956797611418350

Small, D. M., Jones-Gotman, M., & Dagher, A. (2003). Feeding-induced dopamine release in dorsal striatum correlates with meal pleasantness ratings in healthy human volunteers. *NeuroImage*, *19*(4), 1709–1715. https://doi.org/10.1016/s1053-8119(03)00253-2

Stice, E., Yokum, S., Burger, K., Rohde, P., Shaw, H., & Gau, J. M. (2015). A Pilot Randomized Trial of a Cognitive Reappraisal Obesity Prevention Program. *Physiology & Behavior*, *0*, 124–132. https://doi.org/10.1016/j.physbeh.2014.10.022

Sun, W., & Kober, H. (2020). Regulating food craving: From mechanisms to interventions. *Physiology & Behavior*, *222*, 112878. https://doi.org/10.1016/j.physbeh.2020.112878

Themanson, J. R., Pontifex, M. B., & Hillman, C. H. (2008). Fitness and action monitoring: Evidence for improved cognitive flexibility in young adults. *Neuroscience*, *157*(2), 319–328. https://doi.org/10.1016/j.neuroscience.2008.09.014

Vanveen, V., & Carter, C. (2002). The anterior cingulate as a conflict monitor: fMRI and ERP studies. *Physiology & Behavior*, *77*(4–5), 477–482. https://doi.org/10.1016/S0031-9384(02)00930-7

Villalobos, D., Pacios, J., & Vázquez, C. (2021). Cognitive Control, Cognitive Biases and

    Emotion Regulation in Depression: A New Proposal for an Integrative Interplay Model.

    Frontiers in Psychology, *12*, 628416. https://doi.org/10.3389/fpsyg.2021.628416

Wardle, J., & Beales, S. (1988). Control and loss of control over eating: An experimental

    investigation. *Journal of Abnormal Psychology*, *97*(1), 35–40.

    https://doi.org/10.1037/0021-843X.97.1.35

Weng, T. B., Pierce, G. L., Darling, W. G., Falk, D., Magnotta, V. A., & Voss, M. W. (n.d.). The

    Acute Effects of Aerobic Exercise on the Functional Connectivity of Human Brain

    Networks. *Brain Plasticity*, *2*(2), 171–190. https://doi.org/10.3233/BPL-160039

Wilms, R., Lanwehr, R., & Kastenmüller, A. (2020). Emotion Regulation in Everyday Life: The

    Role of Goals and Situational Factors. *Frontiers in Psychology*, *11*, 877.

    https://doi.org/10.3389/fpsyg.2020.00877

Yanagisawa, H., Dan, I., Tsuzuki, D., Kato, M., Okamoto, M., Kyutoku, Y., & Soya, H. (2010).

    Acute moderate exercise elicits increased dorsolateral prefrontal activation and improves

    cognitive performance with Stroop test. *NeuroImage*, *50*(4), 1702–1710.

    https://doi.org/10.1016/j.neuroimage.2009.12.023

Yokum, S., & Stice, E. (2013). Cognitive regulation of food craving: Effects of three cognitive

    reappraisal strategies on neural response to palatable foods. *International Journal of*

    Obesity (2005), *37*(12), 1565–1570. https://doi.org/10.1038/ijo.2013.39

Young, G., & Suri, G. (2020). Emotion regulation choice: A broad examination of external

    factors. *Cognition and Emotion*, *34*(2), 242–261.

    https://doi.org/10.1080/02699931.2019.1611544

Zanto, T. P., & Gazzaley, A. (2013). Fronto-parietal network: Flexible hub of cognitive control. *Trends in Cognitive Sciences*, *17*(12), 10.1016/j.tics.2013.10.001. https://doi.org/10.1016/j.tics.2013.10.001

Zhu, D., Wang, X., Zhao, E., Nozari, N., Notebaert, W., & Braem, S. (2025). Cognitive control is task specific: Further evidence against the idea of domain-general conflict adaptation. *Journal of Experimental Psychology: Learning, Memory, and Cognition*. https://doi.org/10.1037/xlm0001480

Zilverstand, A., Parvaz, M. A., & Goldstein, R. Z. (2017). Neuroimaging cognitive reappraisal in clinical populations to define neural targets for enhancing emotion regulation. A systematic review. *NeuroImage*, *151*, 105–116. https://doi.org/10.1016/j.neuroimage.2016.06.009