Feeding and reproduction of a tropical coastal copepod across warming and copper gradients

Running title: Thermal modulation of copper effects

Ngoc-Anh Vu[1,a], Minh-Hoang Le[2,a], Tuan-Anh Hoang Lu[1], Hoang Viet Luu[1], Nam X. Doan[2], Kiem N.

Truong[1], Khuong V. Dinh[2,3,*]

[1]Department of Ecology, University of Science, Vietnam National University, Hanoi, 334 Nguyen Trai,

Thanh Xuan, Hanoi, Vietnam

[2]Department of Aquaculture, School of Fisheries and Life Sciences, Nha Trang University, No 2 Nguyen

Dinh Chieu, Nha Trang, Vietnam

[3]Department of Biosciences, University of Oslo, Blindernveien 31, 0371, Oslo, Norway

[a] Co-first authors

Corresponding author: Khuong V. Dinh (van.k.dinh@ibv.uio.no)

Abstract

Tropical coastal ecosystems in Southeast Asia are facing rapid warming and increasing pollution. Shallow coastal waters now frequently exceed 34 °C during marine heatwaves, potentially pushing tropical ectotherms beyond their thermal optimum while they are simultaneously exposed to copper (Cu) contamination, especially from aquaculture and shipping activities. However, how warming alters Cu toxicity in dominant tropical zooplankton remains poorly understood. We examined the effects of Cu (0 - 40 µg L^{-1}) across a realistic temperature gradient (26 - 35 °C) on the calanoid copepod *Pseudodiaptomus annandalei*, a dominant grazer of coastal plankton communities. Adult survival, cumulative faecal pellet production (as a proxy for energy intake), and cumulative nauplii production were quantified over seven days. No significant effects of temperature or Cu on adult survival were detected, likely reflecting age-dependent variability among wild-collected individuals. In contrast, temperature strongly structured feeding and reproductive performance, which peaked at 29 - 32 °C and declined at 35 °C. Cu exposure alone had no significant effects at 26 - 32 °C due to high variability in responses. At 35 °C, however, cumulative nauplii production decreased significantly at 30 µg Cu L^{-1} but increased at 20 µg Cu L^{-1}, while faecal pellet production was reduced in Cu-exposed copepods. These findings indicate that warming can modify contaminant effects in tropical zooplankton and highlight the importance of incorporating realistic thermal regimes and natural population variability into ecological risk assessments under climate change.

Keywords: Energy allocation trade-offs, multiple stressors, reproduction, tropical coastal ecosystems, temperature-dependent toxicity

1. Introduction

Tropical coastal ecosystems are increasingly exposed to multiple, overlapping anthropogenic stressors (Halpern et al., 2025, Barlow et al., 2018, Dinh, 2019). Rapid expansion of aquaculture, urbanization, and industrialization in Southeast Asia has resulted in substantial pollutant inputs to estuarine and nearshore environments (Crain et al., 2008, Halpern et al., 2025, Dinh, 2019). Among these pollutants, copper (Cu) is of particular concern due to its widespread use in antifouling paints, algaecides, and aquaculture practices (U.S. Environmental Protection Agency, 2009, Ebenezer et al., 2014). Although Cu rapidly adsorbs to particles and sediments, recurrent inputs from aquaculture and shipping activities may result in chronic exposure for planktonic organisms.

Tropical marine ecosystems are increasingly affected by climate warming and extreme temperature events (Lee et al., 2023, Smale et al., 2019, Gruber et al., 2021, Yao and Wang, 2021). Marine heatwaves (MHWs) have become more frequent, longer-lasting, and more intense over the past decades (Frölicher et al., 2018, Smale et al., 2019). In shallow coastal waters of Southeast Asia, temperatures of 34 °C or higher are frequently recorded during warm seasons (Doan et al., 2018). Tropical marine ectotherms typically live close to their upper thermal limits (Tewksbury et al., 2008, Deutsch et al., 2008). Consequently, even small increases above their thermal optimum may result in disproportionate declines in survival, growth, and reproduction (Nguyen et al., 2011, Doan et al., 2019).

Importantly, warming and pollutants can interact in non-additive manners, resulting in an intensifying or alleviating combined effects (Crain et al., 2008, Dinh et al., 2022). Understanding how warming interacts with contaminants has become a major challenge in aquatic ecology and ecotoxicology in the past two decades (Sokolova and Lannig, 2008, Noyes et al., 2009, Jackson et al., 2016, Dinh et al., 2022). Most studies have focused on temperate species, and are limited to two stressor levels (e.g., the absence *vs*. presence), which do not allow for testing for the non-linear ecological responses (Dinh et al., 2022, Orr et al., 2024), even ecological surprises (e.g., mass mortality Darling and Côté, 2008, Garrabou et al., 2009), especially at sub- and or supra-optimal thermal conditions (Dinh et al., 2022, Orr et al., 2024). Verheyen and Stoks (, 2023) showed that pesticide toxicity was minimal near optimal temperatures but increased at both lower and higher thermal extremes. In addition, most laboratory experiments rely on highly standardized cohorts of organisms, which may not fully capture the demographic variability typical of natural populations (Rohr et al., 2016). Differences in age, reproductive stage, and physiological condition can influence organismal responses to environmental

stressors, potentially altering the magnitude or detectability of contaminant effects under realistic ecological conditions (Muyssen and Janssen, 2007, Dinh et al., 2022, Schmid et al., 2023, Rohr et al., 2016). This may complicate extrapolation from laboratory assays to natural populations (Rohr et al., 2016, Dinh et al., 2022).

Despite growing recognition of temperature-dependent toxicity, few studies have evaluated contaminant effects across a realistic thermal gradient (but see e.g., Verheyen and Stoks, 2023). This gap is particularly relevant for short-lived planktonic organisms such as copepods, which form the foundation of coastal food webs (Chew et al., 2012). Calanoid copepods of the genus *Pseudodiaptomus* are dominant components of tropical plankton communities and play a critical role in transferring primary production to higher trophic levels (Truong et al., 2022, Truong et al., 2014, Chew et al., 2012). Because of their short generation time (5–12 days depending on temperature; Doan et al., 2019, Nguyen et al., 2020), changes in survival or reproduction can rapidly alter population dynamics and ecosystem functioning.

In this study, we investigated the sensitivity of *Pseudodiaptomus annandalei* to a gradient of temperatures (26 - 35 °C) and copper concentrations (0 - 40 µg L^{-1}) over seven days. By quantifying adult survival, cumulative faecal pellet production (as a proxy for food intake and energy assimilation, Isla et al., 2008), and cumulative nauplii production, we aimed to evaluate how warming and Cu exposure affect energy allocation on survival and reproduction, two of the most critical parameters for population growth. Specifically, we addressed three hypotheses (H1-H3): **H1)** Increasing temperature beyond the optimal range reduces feeding activity and reproductive output of *P. annandalei*, with stronger effects on reproduction than on survival. **H2)** Exposure to elevated copper concentrations reduces survival, feeding, and reproductive performance of *P. annandalei*, particularly at higher Cu levels. **H3)** Combined exposure to elevated temperature and copper results in stronger negative effects than either stressor alone, reflecting increased physiological costs under heat stress. By linking feeding activity and reproduction within an energy allocation framework, this study provides insights into how warming and pollution may interact in tropical coastal ecosystems.

2. Materials and methods

2.1 Pseudodiaptomus annandalei

Pseudodiaptomus annandalei is a dominant calanoid copepod occurring along the central coast of Vietnam and Southeast Asian regions (Truong et al., 2022, Truong et al., 2014, Chew et al., 2012). Adult males and females of *P. annandalei* were collected from an aquaculture pond in the Cam Ranh Centre for Tropical Marine

Research and Aquaculture in the summer, using a zooplankton net (mesh size = 200 μm). The sampling site was located at 11°49'26.3"B, 109°07'23.9"E. At the time of sampling, the water temperature and salinity of the pond were 26 °C and 29 PSU, respectively.

2.2 Copper

Copper stock solutions were prepared from $CuSO_4$ (purity >99%, Sigma-Aldrich, Søborg, Denmark) following a method similar to that described by Dinh et al. (, 2021, 2020). Copper concentrations in the control treatment (hereafter referred to as the control) were below the detection limit (1 μg L^{-1}, ICP-MS).

2.3 Experimental design and setup

To test the sensitivity of *P. annandalei* to a range of temperatures and copper exposure, adult copepods were subjected to a fully orthogonal experimental design comprising 20 treatments: 5 copper concentrations (control, 10, 20, 30, and 40 μg L^{-1}) × 4 temperatures (26, 29, 32, and 35°C) (Figure 1). The Cu concentrations varies from ecologically relevant to high levels, but such a range of Cu levels has been found in the coastal waters in Vietnam and worldwide (Jonathan et al., 2011, Teasdale et al., 2003, Marcussen et al., 2014). These temperatures were chosen to cover typical temperature ranges in the shallow coastal water in the South of Vietnam, where the study species is abundant (Doan et al., 2018). Each treatment was replicated ten times, resulting in a total of 100 experimental units (50-ml Falcon tubes).

Each replicate consisted of one egg-carrying female and one male *P. annandalei* placed in a 50 mL Falcon tube containing 20 mL of seawater with the designated copper concentration. The tubes were placed in temperature-controlled water baths maintained using aquarium heaters. All copepods were fed with *Isochrysis galbana* at a concentration of 30,000–33,000 cells L^{-1} (equivalent to 800–850 μg C L^{-1}; Berggreen et al., 1988; Doan et al., 2018). Throughout the experiment, the salinity, photoperiod, and dissolved oxygen concentration were maintained at 30 PSU, 12 h light : 12 h dark, and 5–6 mg L^{-1}, respectively. The contents of each tube were replaced daily with freshly prepared solutions to maintain consistent experimental conditions. Over seven consecutive days, adult survival, cumulative nauplii, and cumulative faecal pellets were recorded daily.

To determine the survival of adult *P. annandalei*, cumulative nauplii production, and cumulative faecal pellets, the contents in each bottle were filtered every day (mesh size = 25 μm). The dead copepods were removed from the tube, and their sex was identified. Alive males and females were then returned to the tube. The remaining contents, containing nauplii and pellets, were transferred to 12-well plates, fixed with Lugol

4%. Numbers of nauplii and faecal pellets were counted using a stereo-microscope (SZ51, Olympus, Japan).

The number of alive females and males, and the total number of nauplii and faecal pellets of each tube in seven

days, were used for statistical analysis.

2.4 Statistical analyses

Statistical analyses were conducted in R (version 4.4.2). The survival of males and females were analysed

using generalized linear mixed models (GLMMs) with a binomial error distribution and logit link function.

Temperature, copper, and their interaction were fitted as fixed effects. The significance of fixed effects was

evaluated using likelihood ratio tests. Cumulative nauplii production and cumulative faecal pellets were

analysed using a generalized linear model with a negative binomial error distribution as their counts showed

overdispersion under a Poisson model (dispersion ~15). A log link function to ensure positive fitted values

appropriate for cumulative count data (Truong et al., 2022). Temperature, copper, and their interaction were

fixed factors. Model significance was assessed using likelihood ratio chi-square tests (Truong et al., 2022).

134 3. Results and discussion

3.1 Effects of temperature on P. annandalei

Survival did not differ among treatments during this short-term exposure (Table 1, Figures 2 and 3). In our

previous studies, temperatures of 34 - 35 °C caused reductions in survival when the exposure duration lasted

for an entire one or two generations of *P. incisus* a congeneric species (Dinh et al., 2021), suggesting a species

specific vulnerability and also the short-term tolerance. In a short-term exposure, copepods may prioritize

maintenance metabolism to preserve survival under acute heat stress (Sokolova, 2013). Such prioritization

often comes at energetic costs, which are often manifested in energy and resources limitation (Metcalfe, 2025,

Sokolova, 2013, Saiz et al., 2022). The reduction in feeding activity at 35 °C suggests (see below) that energy

intake did not fully match the elevated metabolic demand associated with warming (Metcalfe, 2025, Sokolova,

2013, Saiz et al., 2022). Under these conditions, available energy is preferentially allocated to essential

maintenance processes, such as upregulation of heat shock protein, and coping with oxidative damage (Harada

et al., 2019, Low et al., 2018), while energetically expensive processes are constrained.

Acute heat stress significantly reduced the cumulative nauplii and faecal pellet production of *P.*

annandalei. Highest cumulative nauplii produced by *P. annandalei* was found at 32 °C, which is approximately

11% higher than those at 29 °C, and 26% higher than the lowest ones at 26 °C and 35 °C (Table 1, Fig. 4).

Similarly, the cumulative number of pellets per individual was highest at 32 °C, approximately 10% higher

than at 35 °C and 16% higher than those at 26 °C and 29 °C (Table 1, Figure 5). Our results support H1 as the

unimodal pattern in feeding and nauplii production suggests that 29 - 32 °C approximates the optimal thermal

range under the present experimental conditions. As faecal pellet production scales with ingestion and carbon

assimilation, it provides a proxy for energy intake (Isla et al., 2008, Dinh et al., 2019), suggesting a higher

energy acquisition than at both sup- or supra-temperatures. Temperature of 35 °C likely approaches the upper

performance boundary, which is in agreement with general findings of the vulnerability of tropical ectotherms

to temperature extremes (Tewksbury et al., 2008, Deutsch et al., 2008, Grønning et al., 2019). A decline in

faecal pellet production at 35 °C relative to the peak at 32 °C suggests that feeding activity may begin to

decrease as temperatures approach the upper thermal boundary, a pattern has been observed in previous studies

(Doan et al., 2018, Doan et al., 2019, Saiz et al., 2022). This creates a double energetic constraint: increased

expenditure combined with reduced intake efficiency, that may partly contribute to reduced reproduction

(Sokolova and Lannig, 2008, Holmstrup et al., 2010, Jackson et al., 2016, Dinh et al., 2022).

3.2 Temperature dependent Cu-toxicity in P. annandalei

Contrary to H2, Cu exposure did not significantly affect the survival of male or female *P. annandalei* during

the seven-day experiment (Table 1, Figures 2 and 3). Survival varied considerably among temperature × Cu

treatments, and in some cases the lowest survival occurred in the control while the highest survival was

observed at the highest Cu concentration (Figures 2 and 3). Such patterns suggest that variability in the

physiological condition or age structure may have influenced mortality during the experiment (Saiz et al.,

2015, Truong et al., 2020). Age-specific mortality is common in field-collected copepods and can obscure

treatment effects when individuals of different physiological states are used. This variability likely masked

potential effects of temperature and Cu exposure on survival. In our previous studies, we found the strong

effects of both temperature extreme and Cu on the survival of *P. annandalei* and *P. incisus* at comparable

temperatures and Cu concentrations (Dinh et al., 2021, Dinh et al., 2020).

High variabilities of treatment-specific responses of *P. annandalei* were also observed for cumulative

nauplii and faecal pellet production, especially at temperatures of 26 and 29 °C, that may be explained by

similar age-specific performance (Saiz et al., 2015, Truong et al., 2020). However, under extreme temperature,

the interactive effects of temperature and Cu on these parameters emerged, but only partially supporting H3.

At 32°C, Cu-exposed *P. annandalei* had lower cumulative nauplii and faecal pellet production, while at 35 °C,

the effects of Cu on reproductive output were highly complicated and concentration-dependent, generating

interactive effects of Temperature and Cu (Table 1, Figures 4 and 5). Specifically, at 35 °C, cumulative nauplii

production was reduced by 40% at 30 µg Cu L^{-1} compared to the corresponding value at 32 °C, and , and

approximately 24% lower than the control treatment at one at 35 °C, yet at 20 µg Cu L^{-1} it was higher than in

the control treatment at 35 °C and comparable to that observed at 32 °C (Figure 4, Table 1).

For faecal pellet production, *P. anndanalei* showed an average number lower at 35 °C than at 32 °C,

yet this difference was largely driven by reduced pellet production in the control treatment while in Cu-exposed

copepods, pellet production remained similar between the two temperatures (Figure 5, Table 1). Cu-induced

energetic constrains may be even more severe under the combined stressor exposures (Sokolova and Lannig

2008, Sokolova, 2013, Holmstrup et al. 2010). At 35 °C, Cu could further increase oxidative damage and

demands additional ATP for detoxification processes such as metallothionein synthesis and antioxidant

(Harada et al., 2019). Maintenance and detoxification costs could rise simultaneously (Pörtner, 2010, Sokolova

et al., 2012, Lushchak, 2011) that could result in the energy demand exceeding the supply, reproduction is

compromised before survival. This energetic trade-off is consistent with our previous findings, where

combined Cu and MHW exposures reduced fecundity (Dinh et al., 2021, Dinh et al., 2020). We are cautious

of a possibility that variability in male survival influenced reproductive output in some treatments (Kiørboe,

2007, Gusmao and McKinnon, 2009). For example, the loss of males may reduce mating success and

consequently limit nauplii production. However, this mechanism cannot be confirmed with the present data

and requires further investigation.

Importantly, our results also suggest that Cu did not shift the thermal optimum but narrowed

performance breadth by steepening the decline at the upper thermal limit. This supports the principle

articulated in recent thermal performance curve-based ecotoxicology studies: contaminant toxicity is minimal

near the thermal optimum but increases toward thermal extremes (Verheyen and Stoks, 2023). In tropical

systems, where organisms already live close to their upper thermal limits (Deutsch et al., 2008), even small

temperature increases may amplify contaminant effects. Reproduction emerged as the most sensitive endpoint.

Embryonic development is energetically demanding and particularly vulnerable to oxidative stress from

extreme temperatures and pollutants, increasing embryonic mortality, lowering hatching success and resulting in reduced nauplii production (Grønning et al., 2019, Dinh et al., 2021).

3.4 Implications for ecological risk assessments and conclusions

Our findings have several implications for understanding and managing the impacts of multiple environmental stressors in tropical coastal ecosystems. In particular, they provide empirical support for the growing recognition that contaminant effects cannot be evaluated independently of environmental temperature, a central issue in ecotoxicology under global change (Moe et al., 2013, Sokolova and Lannig, 2008, Holmstrup et al., 2010, Dinh et al., 2022, Noyes et al., 2025). The influence of Cu on reproductive output became evident near the upper thermal boundary, suggesting that contaminant toxicity may only emerge under particular environmental contexts, such as during marine heatwaves or other extreme events. Consequently, ecological risk assessments based on single-temperature toxicity tests may underestimate the vulnerability of tropical marine organisms under future warming scenarios. The copepods of the genus *Pseudodiaptomus* play a central role in tropical coastal food webs by transferring primary production from phytoplankton to higher trophic levels, including fish larvae and juvenile stages (Chew et al., 2012). Changes in feeding activity and reproductive output can therefore propagate through the food web and influence energy transfer in coastal ecosystems (Chew et al., 2012, Turner, 2004).

Finally, the relatively high variability observed among treatments likely reflects the natural heterogeneity of wild populations (Blanda et al., 2015, Grønning et al., 2019). Field-collected copepods may differ in age, reproductive stage, and physiological condition, all of which can influence their responses to environmental stressors (Muyssen and Janssen, 2007, Dinh et al., 2022, Schmid et al., 2023, Rohr et al., 2016). While such variability can obscure treatment effects in short-term laboratory experiments, it may better represent the demographic structure of natural populations. In this context, our results suggest that environmentally realistic population variability may mask stressor effects that could become more pronounced under extreme environmental conditions, further challenging the ecological risk assessments. Therefore, combining controlled laboratory studies with standardised life stages and physiological conditions, with ecologically realistic experiments, and field observations will be necessary to improve predictions of how multiple stressors influence tropical marine ecosystems under ongoing global change (Dinh et al., 2022).

Funding

We thank the support of The Ministry of Education and Training, project number B2024-TSN-15.

Acknowledgments

We thank Nha Trang University for supporting facilities to conduct this study and to implement the project B2024-TSN-15.

Conflict of interest

None.

References

BARLOW, J., FRANCA, F., GARDNER, T. A., HICKS, C. C., LENNOX, G. D., BERENGUER, E., CASTELLO, L., ECONOMO, E. P., FERREIRA, J., GUENARD, B., GONTIJO LEAL, C., ISAAC, V., LEES, A. C., PARR, C. L., WILSON, S. K., YOUNG, P. J. & GRAHAM, N. A. J. 2018. The future of hyperdiverse tropical ecosystems. *Nature,* 559**,** 517–526.

BLANDA, E., DRILLET, G., HUANG, C.-C., HWANG, J.-S., JAKOBSEN, H. H., RAYNER, T. A., SU, H.-M., WU, C.-H. & HANSEN, B. W. 2015. Trophic interactions and productivity of copepods as live feed from tropical Taiwanese outdoor aquaculture ponds. *Aquaculture,* 445**,** 11–21.

CHEW, L. L., CHONG, V. C., TANAKA, K. & SASEKUMAR, A. 2012. Phytoplankton fuel the energy flow from zooplankton to small nekton in turbid mangrove waters. *Marine Ecology Progress Series,* 469**,** 7–24.

CRAIN, C. M., KROEKER, K. & HALPERN, B. S. 2008. Interactive and cumulative effects of multiple human stressors in marine systems. *Ecology Letters,* 11**,** 1304–1315.

DARLING, E. S. & CÔTÉ, I. M. 2008. Quantifying the evidence for ecological synergies. *Ecology Letters,* 11**,** 1278–1286.

DEUTSCH, C. A., TEWKSBURY, J. J., HUEY, R. B., SHELDON, K. S., GHALAMBOR, C. K.,

HAAK, D. C. & MARTIN, P. R. 2008. Impacts of climate warming on terrestrial

ectotherms across latitude. *Proceedings of the National Academy of Sciences of the United*

States of America, 105**,** 6668–6672.

DINH, K. V. 2019. Vietnam's fish kill remains unexamined. *Science,* 365**,** 333–333.

DINH, K. V., DOAN, K. L. U., DOAN, N. X., PHAM, H. Q., LE, T. H. O., LE, M.-H., VU, M. T.

T., DAHMS, H.-U. & TRUONG, K. N. 2021. Parental exposures increase the vulnerability

of copepod offspring to copper and a simulated marine heatwave. *Environmental Pollution,*

287**,** 117603.

DINH, K. V., KONESTABO, H. S., BORGÅ, K., HYLLAND, K., MACAULAY, S. J.,

JACKSON, M. C., VERHEYEN, J. & STOKS, R. 2022. Interactive effects of warming and

pollutants on marine and freshwater invertebrates. *Current Pollution Reports,* 8**,** 341–359.

DINH, K. V., NGUYEN, T. T. Q., VO, T.-M.-C., BUI, B. T., DAO, T.-S., TRAN, D. M., DOAN,

N. X., TRUONG, S. H. T., WISZ, S. M., NIELSEN, T. G., VU, T. T. M. & LE, M.-H. 2020.

Interactive effects of extreme temperature and a widespread coastal metal contaminant

reduce the fitness of a common tropical copepod across generations. *Marine Pollution*

Bulletin, 159**,** 111509.

DINH, K. V., OLSEN, M. W., ALTIN, D., VISMANN, B. & NIELSEN, T. G. 2019. Impact of

temperature and pyrene exposure on the functional response of males and females of the

copepod *Calanus finmarchicus*. *Environmental Science and Pollution Research,* 26**,** 29327–

29333.

DOAN, N. X., VU, M. T. T., NGUYEN, H. T., TRAN, H. T. N., PHAM, H. Q. & DINH, K. V.

2018. Temperature- and sex-specific grazing rate of a tropical copepod *Pseudodiaptomus*

annandalei to food availability: implications for live feed in aquaculture. *Aquaculture*

Research, 49**,** 3864–3873.

DOAN, X. N., VU, M. T. T., PHAM, H. Q., WISZ, M. S., NIELSEN, T. G. & DINH, K. V. 2019.

Extreme temperature impairs growth and productivity in a common tropical marine

copepod. *Scientific Reports,* 9**,** 4550.

EBENEZER, V., LIM, W. A. & KI, J.-S. 2014. Effects of the algicides CuSO4 and NaOCl on

various physiological parameters in the harmful dinoflagellate *Cochlodinium polykrikoides*.

Journal of Applied Phycology, 26**,** 2357–2365.

FRÖLICHER, T. L., FISCHER, E. M. & GRUBER, N. 2018. Marine heatwaves under global

warming. *Nature,* 560**,** 360–364.

GARRABOU, J., COMA, R., BENSOUSSAN, N., BALLY, M., CHEVALDONNÉ, P.,

CIGLIANO, M., DIAZ, D., HARMELIN, J. G., GAMBI, M. C., KERSTING, D. K.,

LEDOUX, J. B., LEJEUSNE, C., LINARES, C., MARSCHAL, C., PÉREZ, T., RIBES, M.,

ROMANO, J. C., SERRANO, E., TEIXIDO, N., TORRENTS, O., ZABALA, M.,

ZUBERER, F. & CERRANO, C. 2009. Mass mortality in Northwestern Mediterranean

rocky benthic communities: effects of the 2003 heat wave. *Global Change Biology,* 15**,**

1090–1103.

GRUBER, N., BOYD, P. W., FRÖLICHER, T. L. & VOGT, M. 2021. Biogeochemical extremes

and compound events in the ocean. *Nature,* 600**,** 395–407.

GRØNNING, J. B., DOAN, N. X., DINH, T. N., DINH, K. V. & NIELSEN, T. G. 2019. Ecology

of *Pseudodiaptomus annandalei* in tropical aquaculture ponds with emphasis on the

limitation of production. *Journal of Plankton Research,* 41**,** 741–758.

GUSMAO, L. F. M. & MCKINNON, A. D. 2009. Sex ratios, intersexuality and sex change in

copepods. *Journal of Plankton Research,* 31**,** 1101–1117.

HALPERN, B. S., FRAZIER, M., O'HARA, C. C., VARGAS-FONSECA, O. A. & LOMBARD,

A. T. 2025. Cumulative impacts to global marine ecosystems projected to more than double

by mid-century. *Science,* 389**,** 1216–1219.

HARADA, A. E., HEALY, T. M. & BURTON, R. S. 2019. Variation in thermal tolerance and its

relationship to mitochondrial function across populations of *Tigriopus californicus*.

Frontiers in Physiology, 10.

HOLMSTRUP, M., BINDESBOL, A.-M., OOSTINGH, G. J., DUSCHL, A., SCHEIL, V.,

KOEHLER, H.-R., LOUREIRO, S., SOARES, A. M. V. M., FERREIRA, A. L. G.,

KIENLE, C., GERHARDT, A., LASKOWSKI, R., KRAMARZ, P. E., BAYLEY, M.,

SVENDSEN, C. & SPURGEON, D. J. 2010. Interactions between effects of environmental

chemicals and natural stressors: A review. *Science of The Total Environment,* 408**,** 3746–

3762.

ISLA, J. A., LENGFELLNER, K. & SOMMER, U. 2008. Physiological response of the copepod

Pseudocalanus sp in the Baltic Sea at different thermal scenarios. *Global Change Biology,*

14**,** 895–906.

JACKSON, M. C., LOEWEN, C. J. G., VINEBROOKE, R. D. & CHIMIMBA, C. T. 2016. Net

effects of multiple stressors in freshwater ecosystems: a meta-analysis. *Global Change*

Biology, 22**,** 180–189.

JONATHAN, M. P., ROY, P. D., THANGADURAI, N., SRINIVASALU, S., RODRIGUEZ-

ESPINOSA, P. F., SARKAR, S. K., LAKSHUMANAN, C., NAVARRETE-LOPEZ, M. &

MUNOZ-SEVILLA, N. P. 2011. Metal concentrations in water and sediments from tourist

beaches of Acapulco, Mexico. *Marine Pollution Bulletin,* 62**,** 845–850.

KIØRBOE, T. 2007. Mate finding, mating, and population dynamics in a planktonic copepod

Oithona davisae: There are too few males. *Limnology and Oceanography,* 52**,** 1511–1522.

LEE, H., CALVIN, K., DASGUPTA, D., KRINNER, G., MUKHERJI, A., THORNE, P., TRISOS,

C., ROMERO, J., ALDUNCE, P. & BARRET, K. 2023. IPCC, 2023: Climate change 2023:

Synthesis report, summary for policymakers. Contribution of working groups i, II and III to

the sixth assessment report of the intergovernmental panel on climate change [core writing

team, h. Lee and j. Romero (eds.)]. IPCC, geneva, Switzerland.

LOW, J. S., CHEW, L. L., NG, C. C., GOH, H. C., LEHETTE, P. & CHONG, V. C. 2018. Heat

shock response and metabolic stress in the tropical estuarine copepod *Pseudodiaptomus*

annandalei converge at its upper thermal optimum. *Journal of Thermal Biology,* 74**,** 14–22.

LUSHCHAK, V. I. 2011. Environmentally induced oxidative stress in aquatic animals. *Aquatic*

toxicology, 101**,** 13–30.

MARCUSSEN, H., LOJMAND, H., DALSGAARD, A., HAI, D. M. & HOLM, P. E. 2014. Copper

use and accumulation in catfish culture in the Mekong Delta, Vietnam. *Journal of*

Environmental Science and Health Part A - Toxic/Hazardous Substances & Environmental

Engineering, 49**,** 187–192.

METCALFE, N. B. 2025. How can we best measure the energetic trade-offs faced by animals?

Journal of Experimental Biology, 228**,** jeb250303.

MOE, S. J., DE SCHAMPHELAERE, K., CLEMENTS, W. H., SORENSEN, M. T., VAN DEN

BRINK, P. J. & LIESS, M. 2013. Combined and interactive effects of global climate change

and toxicants on populations and communities. *Environmental Toxicology and Chemistry,*

32**,** 49–61.

MUYSSEN, B. & JANSSEN, C. 2007. Age and exposure duration as a factor influencing Cu and

Zn toxicity toward Daphnia magna. *Ecotoxicology and Environmental Safety,* 68**,** 436–442.

NGUYEN, K. D. T., MORLEY, S. A., LAI, C. H., CLARK, M. S., TAN, K. S., BATES, A. E. &

PECK, L. S. 2011. Upper temperature limits of tropical marine ectotherms: global warming

implications. *Plos One,* 6**,** e29340.

NGUYEN, T. T., LE, M. H., DOAN, N. X., PHAM, H. Q., VU, M. T. T. & DINH, K. V. 2020.

Artificial light pollution increases the sensitivity of tropical zooplankton to extreme

warming. *Environmental Technology & Innovation,* 20**,** 101179.

NOYES, P. D., MCELWEE, M. K., MILLER, H. D., CLARK, B. W., VAN TIEM, L. A.,

WALCOTT, K. C., ERWIN, K. N. & LEVIN, E. D. 2009. The toxicology of climate

change: Environmental contaminants in a warming world. *Environment International,* 35**,** 971–986.

NOYES, P. D., MIRANDA, D., OLIVEIRA DE CARVALHO, G., PERFETTI-BOLAÑO, A., GUIDA, Y., BARBOSA MACHADO TORRES, F., PAULO MACHADO TORRES, J., MIGLIORANZA, K. S., HATJE, V. & BARRA, R. O. 2025. Climate change drives persistent organic pollutant dynamics in marine environments. *Communications Earth & Environment,* 6**,** 363.

ORR, J. A., MACAULAY, S. J., MORDENTE, A., BURGESS, B., ALBINI, D., HUNN, J. G., RESTREPO-SULEZ, K., WILSON, R., SCHECHNER, A., ROBERTSON, A. M., LEE, B., STUPARYK, B. R., SINGH, D., O'LOUGHLIN, I., PIGGOTT, J. J., ZHU, J., DINH, K. V., ARCHER, L. C., PENK, M., VU, M. T. T., JUVIGNY-KHENAFOU, N. P. D., ZHANG, P., SANDERS, P., SCHÄFER, R. B., VINEBROOKE, R. D., HILT, S., REED, T. & JACKSON, M. C. 2024. Studying interactions among anthropogenic stressors in freshwater ecosystems: A systematic review of 2396 multiple-stressor experiments. *Ecology Letters,* 27**,** e14463.

PÖRTNER, H. O. 2010. Oxygen- and capacity-limitation of thermal tolerance: a matrix for integrating climate-related stressor effects in marine ecosystems. *Journal of Experimental Biology,* 213**,** 881–893.

ROHR, J. R., SALICE, C. J. & NISBET, R. M. 2016. The pros and cons of ecological risk assessment based on data from different levels of biological organization. *Critical reviews in toxicology,* 46**,** 756–784.

SAIZ, E., CALBET, A., GRIFFELL, K., BERSANO, J. G. F., ISARI, S., SOLE, M., PETERS, J. & ALCARAZ, M. 2015. Ageing and caloric restriction in a marine planktonic copepod. *Scientific Reports,* 5**,** 14962.

SAIZ, E., GRIFFELL, K., OLIVARES, M., SOLÉ, M., THEODOROU, I. & CALBET, A. 2022.

Reduction in thermal stress of marine copepods after physiological acclimation. *Journal of*

Plankton Research, 44**,** 427–442.

SCHMID, S., RUNDBERGET, J. T., SONG, Y. & TOLLEFSEN, K. E. 2023. Age and

synchronization of *Daphnia magna* affect sensitivity to teflubenzuron in acute standardized

toxicity tests. *Environmental Toxicology and Chemistry,* 42**,** 1806–1815.

SMALE, D. A., WERNBERG, T., OLIVER, E. C. J., THOMSEN, M., HARVEY, B. P., STRAUB,

S. C., BURROWS, M. T., ALEXANDER, L. V., BENTHUYSEN, J. A., DONAT, M. G.,

FENG, M., HOBDAY, A. J., HOLBROOK, N. J., PERKINS-KIRKPATRICK, S. E.,

SCANNELL, H. A., SEN GUPTA, A., PAYNE, B. L. & MOORE, P. J. 2019. Marine

heatwaves threaten global biodiversity and the provision of ecosystem services. *Nature*

Climate Change, 9**,** 306–312.

SOKOLOVA, I. M. 2013. Energy-limited tolerance to stress as a conceptual framework to integrate

the effects of multiple stressors. *Integrative and Comparative Biology,* 53**,** 597–608.

SOKOLOVA, I. M., FREDERICH, M., BAGWE, R., LANNIG, G. & SUKHOTIN, A. A. 2012.

Energy homeostasis as an integrative tool for assessing limits of environmental stress

tolerance in aquatic invertebrates. *Marine environmental research,* 79**,** 1–15.

SOKOLOVA, I. M. & LANNIG, G. 2008. Interactive effects of metal pollution and temperature on

metabolism in aquatic ectotherms: implications of global climate change. *Climate Research,*

37**,** 181–201.

TEASDALE, P. R., APTE, S. C., FORD, P. W., BATLEY, G. E. & KOEHNKEN, L. 2003.

Geochemical cycling and speciation of copper in waters and sediments of Macquarie

Harbour, Western Tasmania. *Estuarine Coastal and Shelf Science,* 57**,** 475–487.

TEWKSBURY, J. J., HUEY, R. B. & DEUTSCH, C. A. 2008. Putting the heat on tropical animals.

Science, 320**,** 1296–1297.

TRUONG, K. N., VU, N.-A., DOAN, N. X., BUI, C. V., LE, M.-H., VU, M. T. T. & DINH, K. V.

2022. Transgenerational exposure to marine heatwaves ameliorates the lethal effect on

tropical copepods regardless of predation stress. *Ecology and Evolution,* 12**,** e9149.

TRUONG, K. N., VU, N.-A., DOAN, N. X., LE, M.-H., VU, M. T. T. & DINH, K. V. 2020.

Predator cues increase negative effects of a simulated marine heatwave on tropical

zooplankton. *Journal of Experimental Marine Biology and Ecology,* 530-531**,** 151415.

TRUONG, T. S. H., NGUYEN, C., LAM, N. N. & JENSEN, K. T. 2014. Seasonal and spatial

distribution of mesozooplankton in a tropical estuary, Nha Phu, South Central Viet Nam.

Biologia, 69**,** 80–91.

TURNER, J. T. 2004. The importance of small planktonic copepods and their roles in pelagic

marine food webs. *Zoological Studies,* 43**,** 255–266.

U.S. ENVIRONMENTAL PROTECTION AGENCY 2009. Reregistration eligibility decision

(RED) for coppers. *In:* U.S. ENVIRONMENTAL PROTECTION AGENCY (ed.).

Washington, DC. .

VERHEYEN, J. & STOKS, R. 2023. Thermal performance curves in a polluted world: too cold and

too hot temperatures synergistically increase pesticide toxicity. *Environmental Science &*

Technology, 57**,** 3270–3279.

YAO, Y. & WANG, C. 2021. Variations in summer marine heatwaves in the South China Sea.

Journal of Geophysical Research: Oceans, 126**,** e2021JC017792.

Table 1. The results of the statistical analyses testing effects of heat stress (Temp) and copper concentration

(Cu) on survival of males and females, cumulative nauplii, and cumulative fecal pellets of *Pseudodiaptomus*

annandalei. Significant *P* values are signed in bold.

Effects	Male survival			Female survival			Cumulative nauplii production			Cumulative faecal pellet production		
	df1, df2	F	*P*	df1, df2	*F*	*P*	df	χ^2	*P*	df	χ^2	*P*
Temp	3, 180	2.493	0.075	3, 180	0.226	0.878	3	11.806	**0.008**	3	13.679	**0.003**
Cu	4, 180	0.389	0.610	4, 180	1.637	0.167	4	6.153	0.188	4	3.794	0.434
Temp × Cu	12, 180	1.366	0.184	12, 180	1.254	0.249	12	29.650	**0.003**	12	47.788	**<0.001**

Figure legends

Figure 1. Experimental design used to test the effects of temperature (26, 29, 32, and 35 °C) and copper concentrations (0, 10, 20, 30 and 40 µg L⁻¹) on the tropical copepod *Pseudodiaptomus annandalei*.

Figure 2. Effects of temperature and copper on male survival of *Pseudodiaptomus annandalei*. Data are visualized as mean ± SEs (n = 10 replications).

Figure 3. Effects of temperature and copper on female survival of *Pseudodiaptomus annandalei*. Data are visualised as mean ± SEs (n = 10 replications).

Figure 4. Effects of temperature and copper on cumulative nauplii of *Pseudodiaptomus annandalei*. Data are visualized as mean ± SEs (n = 10 replications).

Figure 5. Effects of temperature and copper on cumulative fecal pellets of *Pseudodiaptomus annandalei*. Data are visualized as mean ± SEs (n = 10 replications).

Figure 1

Figure 2.

456   **Figure 3**.

460   **Figure 4**.

465   **Figure 5**.