Thermoregulatory Constraints on Regenerative Competence: Evolutionary Trade-Offs Between Metabolic Homeostasis and Tissue Repair

Daniel Pelaez[1,2*], Chloe Moulin[1,2], Jaelyn Chang[3], Kendall W. Knechtel[1]

[1]Bascom Palmer Eye Institute, Department of Ophthalmology, University of Miami Miller School of Medicine, Miami, FL, 33136, USA

[2]Department of Cell and Systems Biology, University of Miami Miller School of Medicine, Miami, FL, 33136, USA

[3]University of Miami Miller School of Medicine, Miami, FL, 33136, USA

*corresponding author

Correspondence: Dr. D. Pelaez, Bascom Palmer Eye Institute, Department of Ophthalmology, University of Miami Miller School of Medicine, 1638 NW 10th Ave, Miami, FL 33136, USA, Tel 305- 482-4850, Fax 305-547-3658. E-mail: dpelaez@miami.edu

Keywords: Thermoregulation, Regeneration, Endothermy, Central Nervous System, Evolution, Calcium, Endo/Sarcoplasmic Reticulum, SERCA

Abstract

Regenerative capacity varies widely across the animal kingdom, yet adult mammals and birds exhibit limited ability to regenerate tissues such as the central nervous system (CNS). The evolutionary basis for this loss of regenerative competence remains unclear. Here we advance a *Thermoregulatory Theory of Regenerative Scope*, positing that the emergence and refinement of endothermic thermoregulation imposed energetic and molecular constraints that progressively limited regenerative potential in highly metabolic tissues. Comparative biology observations reveal that robust regeneration is retained in ectothermic vertebrates and in specific mammalian contexts where thermoregulation is developmentally immature or physiologically attenuated, including neonatal rodents and poikilothermic species. Notably, the developmental acquisition of endothermic capacity in mammals coincides with the decline of regenerative competence. We propose that this trade-off arises from both energetic competition and molecular interference between thermogenic pathways and injury-response programs. Central to this framework is calcium-dependent signaling associated with thermogenesis. Futile Ca^{2+} cycling across the sarco/endoplasmic reticulum through SERCA pumps contributes to heat generation but can also promote sustained intracellular Ca^{2+} elevations that activate inflammatory and fibrotic signaling pathways incompatible with functional regeneration. Proteomic analyses identifying NNAT and TRIM59 as candidate SERCA-associated regulators in the mammalian CNS suggest a molecular crosstalk linking thermogenic Ca^{2+} handling with regenerative failure. Together, these observations support a model in which the evolution of endothermy favored metabolic responsiveness and immune competence at the expense of regenerative capacity.

Introduction

Within the breadth and scope of more than 4.5 billion years of evolutionary refinement by natural selection, the molecular mechanisms underlying most biological processes can be found and studied. This extensive evolutionary molecular repertoire offers a unique opportunity to elucidate potential solutions to intractable challenges in human health, such as the negligible regenerative capacity of the central nervous system (CNS). Regeneration – conceptualized as the ideal outcome of a healing process – varies widely across species throughout the animal kingdom, ranging from what is categorized as biological 'immortality' in some non-vertebrates (e.g. *Turritopsis dorhnii* and *Hydra vulgaris*) [1, 2], to full regenerative competence of limbs and CNS structures in fish and amphibians[3-6]. Unlike other vertebrates, however, most species within the mammaliaform and avian clades undergo regeneration failure following injury to the CNS [7, 8]

The questions of why and how some vertebrate species retain robust regenerative competence while others, particularly mammals and birds, lost this capacity have eluded evolutionary biologists and biomedical researchers for decades [9, 10]. Answers to these questions can unlock critical insights for regenerative medicine and human health.

Despite the general inability to regenerate, there exist instances within the mammalian and avian clades in which robust CNS regeneration can still be observed. Immediately after its development, and approximately up to the first postnatal week in rats and mice, as well as in the chicken while *in ovo*, the optic nerve can undergo spontaneous regeneration when injured [11-14]. The naked mole rat (*Heterocephalus glaber*), while possessing a regressed visual system that senses light presumably only for circadian and seasonal rhythm entrainment [15], undergoes spontaneous regeneration of the axons in its optic nerve following crush injury[16]. The naked mole rat has long been a model organism for mammalian aging and cancer research due to its

unusually long lifespan for a rodent, negligible cellular senescence[17, 18], and extremely low incidence of neoplastic disease[19]. Recently, other rodent species – the brush-furred rat (*Lophuromys*), and the spiny mouse (*Acomys*) – have garnered attention as regenerative models for mammals [20, 21], including for the spiny mouse's increased capacity for retinal regeneration compared to the traditional laboratory mouse (*Mus musculus*) [22, 23]. Understanding what drives the enhanced regenerative potential in altricial mammals and birds, the naked mole rat and *Acomys*, and whether there is any mechanistic overlap with the robust CNS regenerative competence seen in fish and amphibians would significantly advance our approach to promoting enhanced healing outcomes in humans. These instances establish a framework in which elucidating conserved mechanisms governing regenerative competence can be performed through comparative biology across clades, where large differences exist across all domains of molecular biology, between species of the mammalian clade, where these differences are lessened, and within individuals of a particular species (Figure 1).

Figure 1: Schematic of the comparative biology framework, molecular biology domains in which differences are expected in each comparator group, and regenerative segment intersection in which conserved mechanisms underlying enhanced regenerative competence can be elucidated

Thermoregulation and Regeneration

Upon closer analysis, a notable conserved mechanism that emerges between the regenerative segments in the above-highlighted examples relates to the maturity and complexity (or lack

thereof) of the molecular mechanisms responsible for internal body heat generation (endothermy) and thermoregulation. While this is clear for fully ectothermic vertebrate species such as reptiles, amphibians, and fish, the link between whole-body endothermy and regenerative competence becomes much more nuanced in birds and mammals due to the myriad mechanisms underlying endothermic and homeothermic control, as well as the difference in developmental timing and maturity of these mechanisms between species (and even within individuals) of these clades (Figure 2A).

Within the mammalian clade, only the naked mole rat is recognized as a true poikilotherm (i.e. an organism that cannot regulate its own body temperature except by behavioral means) with weak endothermic capacity, adding to the potential argument for a counterbalance between the molecular mechanisms for thermoregulation and regenerative competence. Moreover, while generally recognized as endothermic homeotherms, the thermoregulatory capacity of several members of the *Acomys* and *Lophuromys* families has been shown to be highly dependent on their environment [24-27] indicating a significant degree of poikilothermy, and inadequate or underdeveloped homeothermic mechanisms in these rodent species. A recent study in *Lophuromys melanonyx* [26] demonstrated a decreased thermoregulatory mechanism in the alpine species of these brush-furred rats resulting in a lower heat insulation capacity and suggested a need for solar radiation to support full homeothermy in these diurnal animals. In both *Acomys Russatus* as well as *Acomys Cahirinus*, significant variations in thermal regulation and metabolic rate adaptative capacity to varying environmental conditions have been recorded between populations of animals captured from different habitats [24, 27], showing a degree of poikilothermy which has not yet been studied in detail.

In mice and chicken, a central developmental timing difference in thermoregulatory capacity relates to the precocial and altricial state of these species at birth or hatching, respectively. Embryonically, both species are ectothermic and poikilothermic, relying on external heat sources (namely placental or *in ovo* incubation) without which the embryo is non-viable. As precocial birds, however, chicken transition to an endothermic phenotype upon hatching [28-30], whereas mice and most mammals are altricial at birth, and continue to require external thermal support for several days postnatally [31-33]. Laboratory mouse pups (*Mus musculus*) remain fully poikilothermic for the first week of life and do not develop endothermic and homeothermic regulation until around postnatal day 15 [34]. In rats, full physiological thermoregulatory abilities through both chemical and physical mechanisms is achieved by 14 to 18 days after birth

Figure 2: *(A) Phylogenetic tree of species across the animal kingdom with disparate nervous regeneration and thermoregulatory capacities. (B) Schematic depicting the temporal counterbalance between the acquisition of thermoregulatory maturity and the regenerative capacity in different vertebrate groups.*

[35]. The remarkable temporal overlap between the appearance and maturation of endothermic and thermoregulatory mechanisms and the decline in regenerative capacity within birds and mammals is highly conspicuous yet remains purely circumstantial (Figure 2B). While other researchers have highlighted this observation in other organ systems and offered mechanistic hypotheses [36], whether these are valid and a causative relationship exists remains to be proven.

Thermoregulatory Theory of Regenerative Scope

Teleologically, the loss of regenerative capacity appears counterintuitive to survival, as regeneration enhances resilience to injury. Arguably however, the evolution of endothermy itself is equally as counterintuitive as one of the most energetically expensive biological processes in vertebrates, equivalent to injury response and healing processes, and surpassed only by growth during development [37, 38]. Several hypotheses explain the evolution of endothermy, with the aerobic scope hypothesis – positing that endothermy enhances locomotion and metabolic capacity – being the most widely accepted [38, 39]. A proposed byproduct of endothermy is the development of larger brains and increased neuronal density, which can confer cognitive advantages that offset the loss of regenerative competence [40]. Hypotheses linking thermoregulation to regenerative scope, however, remain largely observational and not testable.

To begin exploring these argumentatively, we consider a proof by contradiction: Could mammals and birds have evolved and thrived without endothermy? Ectothermic organisms, reliant on environmental heat, face increased predation risk during thermoregulation, necessitating robust healing mechanisms to survive non-lethal injurious encounters. In contrast, endothermy enables night foraging and hiding, habitation of resource-diverse environments, and reduced predation, potentially relieving evolutionary pressure to maintain full regenerative competence. This is further supported by the fact that endothermic species produce fewer offspring compared to ectotherms, as the endothermic survival adaptation can prioritize quality over quantity [41]. Consistent with this, the naked mole rat has one of the highest reproductive rates for a mammal and is recognized as a unique exception to the 'one-half rule' for their remarkably large litter sizes [42].

Under a more axiomatic proof, we accept endothermy to be fundamentally predicated on sustained high metabolic flux and make the assumption that an enhanced healing capacity (up to and including full tissue regeneration) confers improved somatic maintenance and thereby extends lifespan. *Ipso facto*, species with lower baseline metabolic rates and reduced thermogenic demands would be predicted to exhibit both slower aging trajectories and greater longevity. This hypothesis is consistent with empirical observations in long-lived mammals characterized by comparatively low metabolic intensity and low core temperatures. The longest-lived mammal, the bowhead whale, demonstrates exceptionally low mass-specific metabolic rates and an average deep body temperature lower than that of most non-hibernating eutherian mammals [43, 44]. Similarly, long-lived terrestrial mammals such as the two-toed sloth and the naked mole-rat exhibit reduced metabolic rates and significantly longer lifespans relative to phylogenetically comparable species. Collectively, these data support the working model that diminished thermoregulatory metabolic demand may facilitate extended lifespans through enhanced healing competences.

How, then, could thermoregulatory complexity interfere with regenerative competence? Building upon our conceptual framework, our theory delineates two interdependent mechanistic dimensions that can collectively explain this inverse functional relationship:

Energetic resource constraints: heat generation and maintenance in endothermic homeotherms is arguably the most metabolically demanding process in vertebrate biology, rivaled only by healing, growth, and regeneration. The high energetic cost of homeothermy may limit the availability of bioenergetic resources for functional healing (regeneration), particularly in the most metabolic demanding tissues of a 'warm-blooded' organism, such as the myocardium or CNS [45]. This competition for energy can explain why regenerative processes in the most metabolically demanding tissues are curtailed in mammals as their endothermic capacity expands [36]. Studies

in laboratory mice have suggested that maturation of the central nervous system is probably the main factor responsible for the development of thermogenesis [34]. In the myocardium, the temporal correlation between the loss of regenerative capacity and the development of endothermy has been linked to changes in thyroid hormone levels around the same developmental period [36].

Molecular crosstalk interference: The molecular mechanisms underlying thermogenesis directly interfere with injury response and regenerative pathways in mammals. Heat is generated when any amount of free energy is not converted into downstream chemical or kinetic energy (work) but rather dissipated into the surroundings. Physiologically, thermogenesis results from the decoupling of the release of chemical energy (stored in ATP and other bioenergetic molecules) from the establishment of the chemiosmotic gradients (work) that drive myriad cell biological processes. This decoupling and loss of chemiosmotic control can disrupt the tightly orchestrated processes required for regeneration, including cell cycle reentry and transient proliferation, cellular reprogramming, and/or tissue remodeling [46-48].

Thermoregulatory and Regenerative Mechanism Crosstalk

Endothermy is an umbrella term which encompasses a multitude of physical and biochemical mechanisms for internal heat generation in different species, and even within an individual. In broad mechanical terms, endothermic mechanisms are categorized into shivering thermogenesis and non-shivering thermogenesis (NST). Biochemically, however, endothermic mechanisms can be categorized into two different types: *i.* those that involve the flux of protons (H^+), and *ii.* those that involve the flux of charged ions — most notably free calcium ions (Ca^{2+}) — when uncoupled from the chemical energy expenditure intended to drive their compartmentalization in the cell. The most well-known example of NST involving proton movement occurs in the mitochondria of brown adipose tissues (BAT), where the permeation of the inner mitochondrial membrane (IMM)

to protons (H^+) by uncoupling proteins (UCP1/2/3) results in futile proton shuttling at the expense of NADH and $FADH_2$ oxidation by complexes of the electron transport chain (ETC) and dissipation of the excess decoupled energy as heat. In shivering thermogenesis, rapid, small, involuntary muscle contractions in response to cold temperatures are triggered by calcium ion (Ca^{2+}) release from sarcoplasmic reticulum (SR) stores and their subsequent binding to troponin to enable the establishment of the myosin-actin cross-bridge. Detachment of the myosin head from actin filaments requires ATP hydrolysis to prepare the contractile unit for the next interaction. Unlike in large coordinated and voluntary muscle contractions, during shivering thermogenesis, restriction of calcium reuptake to the SR through the Sarcoplasmic/Endoplasmic Reticulum Calcium ATPase (SERCA) by negative regulators like sarcolipin (SLN) [49] and phospholamban (PLN) [50] uncouples ATP hydrolysis from Ca^{2+} flux [51], resulting in multiple subsequent, and uncoordinated small muscle contractions and heat generation. Shivering thermogenesis thus results in significant heat generation through ATP hydrolysis at two sites: the myosin head uncoupled from force-generating muscle movements (work), and at SERCA channels uncoupled from calcium reuptake (work) to the SR/ER stores.

Calcium is a ubiquitous second messenger, and calcium ion fluxes underlie a vast number of biological processes, making the elucidation of molecular crosstalk interference between thermoregulation and regeneration a difficult task. Central to the discussion, however, sustained elevation of intracellular Ca^{2+} is a central proximal signal linking tissue injury to inflammatory amplification and fibrotic or gliotic remodeling. Cytosolic Ca^{2+} overload arising from plasma membrane influx or endoplasmic reticulum (ER) release through IP_3 (IP_3R) and ryanodine receptors (RyR) activates multiple kinase pathways converging on transcriptional programs governed by NF-kappaB that drive expression of pro-inflammatory mediators [52-54]. In parallel,

Ca²⁺ mobilization promotes assembly and activation of the NLRP3 inflammasome, facilitating caspase 1-dependent maturation of IL-1β and IL-18 and propagating sterile inflammation [55, 56]. Persistent Ca²⁺ signaling also potentiates profibrotic pathways by augmenting transforming growth factor beta (TGF-β)/SMAD signaling, reinforcing extracellular matrix deposition [55]. In the adult mammalian CNS, pervasive intracellular Ca²⁺ overload has been demonstrated as an injury-induced mediator of mitochondrial oxidation and structural remodeling [57]. Moreover, Ca²⁺ waves in astrocytes enhance inflammatory and STAT3-dependent transcriptional responses, promoting reactive gliosis characterized by GFAP upregulation and glial scar formation [58-60]. Conversely, in the regenerative competent *xenopus laevis* [61], Ca²⁺ transients following optic nerve transection are short-lived, and calcium dynamics return to homeostatic levels within minutes. Inhibition of SERCA2 — but not Na²⁺ channels — in this model extends calcium permanence and slows clearance and abrogates the regenerative outcome and functional recovery of sight compared to the SERCA2-unihibited controls [manuscript in preparation]. In a recent review, the strategies that promote axonal regeneration in the adult mammalian CNS have been shown to modulated intracellular calcium dynamics via one of three mechanisms: *i*. restriction of Ca²⁺ release from the ER via IP₃R and RyR; *ii*. sequestration or chelation of free Ca²⁺ ions, or *iii*. Stimulation of Ca²⁺ reuptake to ER stores via modulation of SERCA negative regulators, with strategies that engage more than one of these mechanisms exerting a synergistic effect on axonal regeneration [manuscript in preparation]. Collectively, these data position dysregulated and sustained intracellular free Ca²⁺ as a nodal integrator of inflammatory transcriptional activation and maladaptive fibrotic or gliotic remodeling across organ systems in mammals.

Calcium dynamics are subject to stringent spatial and molecular regulation across tissues within an individual and between species, with distinct repertoires of sarco/endoplasmic reticulum

Ca²⁺-ATPase (SERCA) modulators shaping both metabolic output and cellular fate decisions [62, 63]. The central nervous system (CNS), which by mass is the most metabolically active organ system in mammals and a major contributor to endothermic heat production [45], exemplifies this specialization. In excitable tissues, futile Ca²⁺ cycling across the endoplasmic reticulum (ER) constitutes a thermogenic mechanism that elevates basal metabolic rate. Notably, the liver, despite being highly metabolic and one of the few mammalian organs retaining partial regenerative capacity, is devoid of functional ryanodine receptors (RyR) [64, 65], suggesting that attenuation of rapid ER Ca²⁺ release may be permissive for regenerative competence and reinforcing a mechanistic link between Ca²⁺ cycling dynamics and healing potential. Unlike striated muscle, canonical SERCA regulators such as phospholamban (PLN), myoregulin (MRLN), and sarcolipin (SLN) have not been identified in the mammalian CNS, implying the presence of distinct neural modulators. Neuronatin (NNAT), a neural tissue ER-associated proteolipid, has been linked to neurogenesis and systemic metabolic regulation [63, 66] positioning it as a candidate integrator of developmental and thermogenic programs in these tissues. A homology-based screen of SERCA-interacting proteins identified the transmembrane domain of the Tripartite Motif Containing 59 (TRIM59) protein as structurally and molecularly analogous to SLN (Figure 3A-C), raising the possibility that TRIM59 modulates ER Ca²⁺ handling in neural tissue. Intriguingly, NNAT lacks clear orthologs in regeneration-competent species, and the transmembrane domain of TRIM59 is significantly divergent in these organisms (Figure 3D). Finally, protein docking modeling shows a strong interaction between human SERCA2 and the transmembrane-domain of the human TRIM59 (Figure 3E), whereas there were no predicted interactions between these proteins from *xenopus laevis* (not shown).

Figure 3: *Structural (A) and amino acid sequence (B) homology between sarcolipin (SLN) and the transmembrane-domain (TM) of the tripartite motif containing 59 (TRIM59) protein in humans and mice. (C) Atomic positional alignment between the TM-TRIM59 and SLN from humans. (D) Sequence alignments of neuronatin (NNAT) and TRIM59 across species with divergent regenerative capacity: Mus musculus, Acomys russatus, Heterocephalus glaber, Xenopus laevis, Danio rerio, and Ambystoma mexicanum. (E)Protein docking models for predicted interactions between human SERCA2 and TRIM59-TM.*

Immunoprecipitation (IP) and co-immunoprecipitation (co-IP) followed by liquid chromatography and mass spectrometry (LC-MS/MS) studies of SERCA2 from mammalian optic

nerves corroborate this finding, showing that only NNAT and TRIM59 co-precipitate with SERCA2 in this tissue (Figure 4A). Additionally, the assays identified differences in SERCA2 and TRIM59 abundance between samples collected at developmental timepoints when the mouse optic nerve retains robust endogenous regenerative competence or undergoes regenerative failure after injury (Figure 4B), revealing a divergent regulatory framework for SERCA2 in these disparate regenerative paradigms (Figure 4C). Given that TRIM59 is intricately implicated in immune regulation and inflammatory signaling [67-69], these evolutionary modifications could suggest that ER Ca^{2+}-dependent thermogenesis may have been co-opted to fine-tune immune and inflammatory competence in endotherms. However, the same Ca^{2+}-driven inflammatory modulation, while evolutionarily beneficial, could impose a trade-off by promoting gliotic and growth suppressive responses that constrain regenerative scope, thereby advancing a thermoregulatory theory in which the evolution of endothermy mechanistically intersects with, and potentially hinders, tissue regeneration.

Figure 4: *SERCA2 binding partners in the mammalian CNS. (A) SERCA2 co-IP and LC-MS/MS quantification of canonical SERCA2 regulators and TRIM59 in adult mouse optic nerves. (B) Mass Spectrometry quantification of SERCA2, TRIM59, and NNAT from mouse optic nerve tissue at two different developmental times. (C) Relative abundance of NNAT or TRIM59 to SERCA2 in mice optic nerves during the first 9 postnatal days, or at 3-5 months of age.*

Discussion

Here we advance a *Thermoregulatory Theory of Regenerative Scope*, proposing that the evolutionary acquisition and elaboration of composite homeothermy imposed energetic and molecular constraints that progressively limited regenerative competence in highly metabolic tissues, with a particular focus on the adult mammalian central nervous system (CNS). Our framework integrates comparative physiology, developmental timing, and calcium-dependent signaling pathways to suggest that thermogenic specialization and regeneration are not merely coincidental traits, but mechanistically and evolutionarily intertwined.

A central tenet of this theory is that the maturation of endothermic capacity, specifically mechanisms involving futile calcium cycling across the sarco/endoplasmic reticulum (SR/ER), directly interferes with the cellular processes required for functional regeneration. In mammals, shivering thermogenesis depends, in part, on uncoupling ATP hydrolysis from effective Ca^{2+} reuptake through SERCA via negative regulators such as sarcolipin and phospholamban. We hypothesize that removing or attenuating negative regulation of SERCA in non-regenerating tissues would re-couple ATP hydrolysis to rapid Ca^{2+} clearance, reduce cytosolic Ca^{2+} permanence after injury, and thereby enhance regenerative capacity. This prediction is consistent with experimental paradigms demonstrating that accelerated Ca^{2+} sequestration and restriction of ER Ca^{2+} release promote axonal regeneration, whereas prolonged intracellular Ca^{2+} elevation amplifies inflammatory transcriptional programs and fibrotic or gliotic remodeling [52, 55, 57].

Tissue-specific differences further support this mechanistic axis. Hepatocytes retain robust regenerative capacity despite their high metabolic demands. Notably, the liver lacks functional ryanodine receptors (RyR) [64, 65], potentially restricting rapid ER Ca^{2+} release and buffering against sustained Ca^{2+}-driven inflammatory amplification. In contrast, cardiomyocytes and CNS

neurons — cells that exhibit profound regenerative failure in adult mammals — are embedded in tightly regulated Ca^{2+} cycling architectures integral to excitation-contraction coupling, action potential propagation and thermogenic output. Hirose and colleagues [36] previously associated the postnatal acquisition of endothermic physiology in cardiomyocytes with loss of regenerative competence, linking thyroid hormone-dependent transitions to polyploidization and cell cycle exit. Thyroid hormones are established regulators of Ca^{2+} cycling proteins and ion transport complexes [70-72], including uncoupled ATPase activity by SERCA [73] providing a plausible endocrine bridge between systemic thermogenic maturation and intracellular calcium homeostasis. Another thyroid-specific calcium modulating hormone — calcitonin — is known to be highest during the same perinatal transition stages and decreases in serum concentration with age [74, 75]. Calcitonin has been shown essential for peripheral nerve regeneration[76, 77]. Extending this logic to the CNS, our data identifying NNAT and TRIM59 as candidate SERCA-associated interactors in mammalian optic nerve suggest that neural thermogenic specialization may likewise alter ER Ca^{2+} handling in ways that bias injury responses toward inflammation and gliosis rather than regeneration.

From an evolutionary perspective, the loss of regenerative capacity appears maladaptive. However, the energetic burden of endothermy suggests that the trade-off may have conferred alternative survival advantages. Enhanced immune and inflammatory competence offers a compelling candidate. Sustained and rapidly mobilized Ca^{2+} flux is central not only to thermogenesis but also to activation of NF-κB–dependent transcription, inflammasome assembly, cytokine maturation, and reactive gliosis. The same ER Ca^{2+} dynamics that permit rapid thermogenic adaptation may therefore potentiate host defense in injury-prone, high-activity endotherms. In this view, regenerative suppression is not a passive loss but a systems-level

consequence of prioritizing inflammatory efficiency and metabolic responsiveness. The co-option of ER Ca^{2+}-dependent futile cycling to support immune surveillance and inflammatory amplification could have been selectively advantageous in endothermic lineages expanding to new and variable ecological niches.

This theory generates experimentally tractable hypotheses. First, genetic or pharmacologic re-coupling of SERCA activity in adult mammalian CNS tissue would be predicted to reduce injury-induced Ca^{2+} persistence, dampen inflammatory transcriptional activation, and enhance structural healing. Second, comparative analyses across species with variable thermogenic capacities should reveal correlations between the SERCA regulatory framework, RyR and IP_3R expression, metabolic hormone signaling intensity, and regenerative outcomes. Third, developmental transitions in thermoregulation, such as the postnatal maturation of endothermy in altricial rodents, should coincide with measurable shifts in ER Ca^{2+} kinetics and inflammatory moiety expression profiles that precede regenerative decline.

While circumstantial at present, the convergence of evolutionary physiology, endocrine regulation, calcium signaling biology, and injury response supports a coherent and testable model. By reframing regenerative failure not as an isolated deficiency but as a thermoregulatory byproduct of metabolic evolution, the *Thermoregulatory Theory of Regenerative Scope* provides a unifying lens through which disparate observations — ranging from neonatal mammalian plasticity to naked mole rat resilience and ectothermic regeneration — can be interpreted. Further mechanistic dissection of calcium-dependent thermogenic pathways in the CNS and other high-energy tissues will determine whether endothermy's evolutionary success indeed came at the cost of regenerative potential.

References:

1. Velasco-Munoz, V., et al., *Regenerative characteristics of the immortal jellyfish, Turritopsis dohrnii, and their potential implications for human aging.* Rev Esp Geriatr Gerontol, 2025. **60**(3): p. 101607.
2. Morris Little, H., et al., *A molecular, spatial and regulatory atlas of the Hydra vulgaris nervous system.* Development, 2025. **152**(20).
3. Zhong, J., et al., *Multi-species atlas resolves an axolotl limb development and regeneration paradox.* Nat Commun, 2023. **14**(1): p. 6346.
4. Wei, X., et al., *Single-cell Stereo-seq reveals induced progenitor cells involved in axolotl brain regeneration.* Science, 2022. **377**(6610): p. eabp9444.
5. Phipps, L.S., et al., *Model systems for regeneration: Xenopus.* Development, 2020. **147**(6).
6. Namikawa, K., S. Pose-Mendez, and R.W. Koster, *Genetic modeling of degenerative diseases and mechanisms of neuronal regeneration in the zebrafish cerebellum.* Cell Mol Life Sci, 2024. **82**(1): p. 26.
7. Varadarajan, S.G., et al., *Central nervous system regeneration.* Cell, 2022. **185**(1): p. 77-94.
8. Silver, J., M.E. Schwab, and P.G. Popovich, *Central nervous system regenerative failure: role of oligodendrocytes, astrocytes, and microglia.* Cold Spring Harb Perspect Biol, 2014. **7**(3): p. a020602.
9. Galis, F., G.P. Wagner, and E.L. Jockusch, *Why is limb regeneration possible in amphibians but not in reptiles, birds, and mammals?* Evol Dev, 2003. **5**(2): p. 208-20.
10. Alibardi, L., *Organ regeneration evolved in fish and amphibians in relation to metamorphosis: Speculations on a post-embryonic developmental process lost in amniotes after the water to land transition.* Ann Anat, 2019. **222**: p. 114-119.
11. Palazzo, I., et al., *NF-kappaB signaling regulates the formation of proliferating Muller glia-derived progenitor cells in the avian retina.* Development, 2020. **147**(10).
12. Goldberg, J.L., et al., *Amacrine-signaled loss of intrinsic axon growth ability by retinal ganglion cells.* Science, 2002. **296**(5574): p. 1860-4.
13. Yun, M.H., *Changes in Regenerative Capacity through Lifespan.* Int J Mol Sci, 2015. **16**(10): p. 25392-432.
14. L. Belecky-Adams, T., et al., *Chapter 8 - The Chick as a Model for Retina Development and Regeneration*, in *Animal Models in Eye Research*, P.A. Tsonis, Editor. 2008, Academic Press: London. p. 102-119.
15. Crish, S.D., C.M. Dengler-Crish, and K.C. Catania, *Central visual system of the naked mole-rat (Heterocephalus glaber).* Anat Rec A Discov Mol Cell Evol Biol, 2006. **288**(2): p. 205-12.
16. Park, K.K., et al., *Retinal ganglion cell survival and axon regeneration after optic nerve injury in naked mole-rats.* J Comp Neurol, 2017. **525**(2): p. 380-388.

17. Buffenstein, R., *Negligible senescence in the longest living rodent, the naked mole-rat: insights from a successfully aging species.* J Comp Physiol B, 2008. **178**(4): p. 439-45.

18. Buffenstein, R. and J.U. Jarvis, *The naked mole rat--a new record for the oldest living rodent.* Sci Aging Knowledge Environ, 2002. **2002**(21): p. pe7.

19. Delaney, M.A., et al., *Initial Case Reports of Cancer in Naked Mole-rats (Heterocephalus glaber).* Vet Pathol, 2016. **53**(3): p. 691-6.

20. Riddell, B., et al., *Complex tissue regeneration in Lophuromys reveals a phylogenetic signal for enhanced regenerative ability in deomyine rodents.* Proc Natl Acad Sci U S A, 2025. **122**(1): p. e2420726122.

21. Maden, M. and J.A. Varholick, *Model systems for regeneration: the spiny mouse, Acomys cahirinus.* Development, 2020. **147**(4).

22. Bills, J., et al., *Spiny mice display a pro-regenerative response to acute retinal damage.* Investigative Ophthalmology & Visual Science, 2023. **64**(8): p. 1625-1625.

23. Biswal, M.R., et al., *Retinal Pigment Epithelium (RPE) injury response in African spiny mice (Acomys) following laser injury.* Investigative Ophthalmology & Visual Science, 2024. **65**(7): p. 3461-3461.

24. Scantlebury, M., et al., *Differential energy costs of winter acclimatized common spiny mice Acomys cahirinus from two adjacent habitats.* Comp Biochem Physiol A Mol Integr Physiol, 2004. **137**(2): p. 419-23.

25. Levy, O., T. Dayan, and N. Kronfeld-Schor, *Adaptive thermoregulation in golden spiny mice: the influence of season and food availability on body temperature.* Physiol Biochem Zool, 2011. **84**(2): p. 175-84.

26. Ivlev, Y.F. and L.A. Lavrenchenko, *A decrease in heat insulation of the black-clawed brush-furred rat (Lophuromys melanonyx, Petter) during adaptation to high altitudes.* Dokl Biol Sci, 2016. **466**: p. 36-41.

27. Haim, A. and A. Borut, *Heat production and dissipation in golden spiny mice,Acomys russatus from two extreme habitats.* Journal of comparative physiology, 1981. **142**(4): p. 445-450.

28. Seebacher, F., T.S. Schwartz, and M.B. Thompson, *Transition from ectothermy to endothermy: the development of metabolic capacity in a bird (Gallus gallus).* Proc Biol Sci, 2006. **273**(1586): p. 565-70.

29. Price, E.R. and E.M. Dzialowski, *Development of endothermy in birds: patterns and mechanisms.* J Comp Physiol B, 2018. **188**(3): p. 373-391.

30. Rippamonti, J. and E.M. Dzialowski, *Thyroid hormone manipulation influences development of endothermy and hatching in white leghorn chickens (Gallus gallus).* J Therm Biol, 2023. **114**: p. 103582.

31. Zepeda, J.A., et al., *Individual differences in early body mass affect thermogenic performance and sibling interactions in litter huddles of the house mouse.* Dev Psychobiol, 2018. **60**(7): p. 825-835.

32.   Alberts, J.R., *Huddling by rat pups: multisensory control of contact behavior.* J Comp Physiol Psychol, 1978. **92**(2): p. 220-30.

33.   Michener, G.R. and T. Dic Charge, *Chapter 16 - Development of Thermoregulation in Altricial Rodents*, in *Exploring Animal Behavior in Laboratory and Field*, B.J. Ploger and K. Yasukawa, Editors. 2003, Academic Press: San Diego. p. 187-197.

34.   Lagerspetz, K.Y.H., *Postnatal development of thermoregulation in laboratory mice.* Helgoländer wissenschaftliche Meeresuntersuchungen, 1966. **14**(1): p. 559-571.

35.   Malik, S.S. and J.E. Fewell, *Thermoregulation in rats during early postnatal maturation: importance of nitric oxide.* Am J Physiol Regul Integr Comp Physiol, 2003. **285**(6): p. R1366-72.

36.   Hirose, K., et al., *Evidence for hormonal control of heart regenerative capacity during endothermy acquisition.* Science, 2019. **364**(6436): p. 184-188.

37.   Hulbert, A.J. and P.L. Else, *Membranes as possible pacemakers of metabolism.* J Theor Biol, 1999. **199**(3): p. 257-74.

38.   Clarke, A. and H.O. Portner, *Temperature, metabolic power and the evolution of endothermy.* Biol Rev Camb Philos Soc, 2010. **85**(4): p. 703-27.

39.   Hayes, J.P. and T. Garland, Jr., *The Evolution of Endothermy: Testing the Aerobic Capacity Model.* Evolution, 1995. **49**(5): p. 836-847.

40.   Osvath, M., et al., *Thought for food: the endothermic brain hypothesis.* Trends Cogn Sci, 2024. **28**(11): p. 998-1010.

41.   Labocha, M.K. and J.P. Hayes, *Endotherm ☆*, in *Encyclopedia of Ecology (Second Edition)*, B. Fath, Editor. 2019, Elsevier: Oxford. p. 368-374.

42.   Sherman, P.W., S. Braude, and J.U.M. Jarvis, *Litter Sizes and Mammary Numbers of Naked Mole-Rats: Breaking the One-Half Rule.* Journal of Mammalogy, 1999. **80**(3): p. 720-733.

43.   George, J.C., *Growth, Morphology And Energetics Of Bowhead Whales (Balaena Mysticetus)*. 2009.

44.   George, J.C. *Energetics and Temperature Regulation*. North Slope Borough Wildlife Management - Bowhead Whale Anatomy & Physiology Studies [Official Website of the North Slope Borough] March 5, 2026]; Available from: https://www.north-slope.org/departments/wildlife-management/studies-research-projects/bowhead-whales/bowhead-whale-anatomy-physiology-studies/energetics-temperature-regulation.

45.   Karbowski, J., *Global and regional brain metabolic scaling and its functional consequences.* BMC Biol, 2007. **5**: p. 18.

46.   Lin, T.Y., et al., *Fibroblast dedifferentiation as a determinant of successful regeneration.* Dev Cell, 2021. **56**(10): p. 1541-1551 e6.

47.   Jopling, C., et al., *Zebrafish heart regeneration occurs by cardiomyocyte dedifferentiation and proliferation.* Nature, 2010. **464**(7288): p. 606-9.

48.   Fang, J., et al., *Cellular polyploidy in organ homeostasis and regeneration.* Protein Cell, 2023. **14**(8): p. 560-578.

49.   Bal, N.C. and M. Periasamy, *Uncoupling of sarcoendoplasmic reticulum calcium ATPase pump activity by sarcolipin as the basis for muscle non-shivering thermogenesis.* Philos Trans R Soc Lond B Biol Sci, 2020. **375**(1793): p. 20190135.

50.   Bhupathy, P., G.J. Babu, and M. Periasamy, *Sarcolipin and phospholamban as regulators of cardiac sarcoplasmic reticulum Ca2+ ATPase.* J Mol Cell Cardiol, 2007. **42**(5): p. 903-11.

51.   de Meis, L., *Uncoupled ATPase activity and heat production by the sarcoplasmic reticulum Ca2+-ATPase. Regulation by ADP.* J Biol Chem, 2001. **276**(27): p. 25078-87.

52.   Suetomi, T., et al., *Inflammation and NLRP3 Inflammasome Activation Initiated in Response to Pressure Overload by Ca(2+)/Calmodulin-Dependent Protein Kinase II delta Signaling in Cardiomyocytes Are Essential for Adverse Cardiac Remodeling.* Circulation, 2018. **138**(22): p. 2530-2544.

53.   Berry, C.T., M.J. May, and B.D. Freedman, *STIM- and Orai-mediated calcium entry controls NF-kappaB activity and function in lymphocytes.* Cell Calcium, 2018. **74**: p. 131-143.

54.   Valdes, J.A., et al., *NF-kappaB activation by depolarization of skeletal muscle cells depends on ryanodine and IP3 receptor-mediated calcium signals.* Am J Physiol Cell Physiol, 2007. **292**(5): p. C1960-70.

55.   Wang, W., et al., *Inflammasome-independent NLRP3 augments TGF-beta signaling in kidney epithelium.* J Immunol, 2013. **190**(3): p. 1239-49.

56.   Zhang, W.J., et al., *Inflammasomes and Fibrosis.* Front Immunol, 2021. **12**: p. 643149.

57.   Breckwoldt, M.O., et al., *Multiparametric optical analysis of mitochondrial redox signals during neuronal physiology and pathology in vivo.* Nat Med, 2014. **20**(5): p. 555-60.

58.   Sofroniew, M.V., *Molecular dissection of reactive astrogliosis and glial scar formation.* Trends Neurosci, 2009. **32**(12): p. 638-47.

59.   Luo, J., *TGF-beta as a Key Modulator of Astrocyte Reactivity: Disease Relevance and Therapeutic Implications.* Biomedicines, 2022. **10**(5).

60.   Du, S., et al., *Calcium influx and activation of calpain I mediate acute reactive gliosis in injured spinal cord.* Exp Neurol, 1999. **157**(1): p. 96-105.

61.   Moulin, C., et al., *Novel laser model of optic nerve transection provides valuable insights about the dynamics of optic nerve regeneration.* Sci Rep, 2024. **14**(1): p. 27412.

62.   Periasamy, M., et al., *Role of SERCA Pump in Muscle Thermogenesis and Metabolism.* Compr Physiol, 2017. **7**(3): p. 879-890.

63.   Lin, H.H., et al., *Neuronatin promotes neural lineage in ESCs via Ca(2+) signaling.* Stem Cells, 2010. **28**(11): p. 1950-60.

64.   Amaya, M.J. and M.H. Nathanson, *Calcium signaling in the liver.* Compr Physiol, 2013. **3**(1): p. 515-39.

65.   Giannini, G., et al., *The ryanodine receptor/calcium channel genes are widely and differentially expressed in murine brain and peripheral tissues.* J Cell Biol, 1995. **128**(5): p. 893-904.

66. Braun, J.L., et al., *Neuronatin regulates whole-body metabolism: is thermogenesis involved?* FASEB Bioadv, 2020. **2**(10): p. 579-586.

67. Jin, Z., et al., *The role of TRIM59 in immunity and immune-related diseases.* Int Rev Immunol, 2024. **43**(1): p. 33-40.

68. An, Y., et al., *TRIM59 expression is regulated by Sp1 and Nrf1 in LPS-activated macrophages through JNK signaling pathway.* Cell Signal, 2020. **67**: p. 109522.

69. Jin, Z., et al., *TRIM59 Protects Mice From Sepsis by Regulating Inflammation and Phagocytosis in Macrophages.* Front Immunol, 2020. **11**: p. 263.

70. Carr, A.N. and E.G. Kranias, *Thyroid hormone regulation of calcium cycling proteins.* Thyroid, 2002. **12**(6): p. 453-7.

71. Little, A.G. and F. Seebacher, *The evolution of endothermy is explained by thyroid hormone-mediated responses to cold in early vertebrates.* J Exp Biol, 2014. **217**(Pt 10): p. 1642-8.

72. Garel, J.M., P. Besnard, and C. Rebut-Bonneton, *C cell activity during the prenatal and postnatal periods in the rat.* Endocrinology, 1981. **109**(5): p. 1573-7.

73. Arruda, A.P., et al., *Hyperthyroidism increases the uncoupled ATPase activity and heat production by the sarcoplasmic reticulum Ca2+-ATPase.* Biochem J, 2003. **375**(Pt 3): p. 753-60.

74. Deftos, L.J., et al., *Influence of age and sex on plasma calcitonin in human beings.* N Engl J Med, 1980. **302**(24): p. 1351-3.

75. Parthemore, J.G. and L.J. Deftos, *Calcitonin secretion in normal human subjects.* J Clin Endocrinol Metab, 1978. **47**(1): p. 184-8.

76. Yan, J.G., et al., *Calcitonin pump improves nerve regeneration after transection injury and repair.* Muscle Nerve, 2015. **51**(2): p. 229-34.

77. Toth, C.C., et al., *Locally synthesized calcitonin gene-related Peptide has a critical role in peripheral nerve regeneration.* J Neuropathol Exp Neurol, 2009. **68**(3): p. 326-37.