Title: Basal ganglia output dynamically controls skilled forelimb kinematics in real time

Authors: Shaolin Ruan[1], Henry H. Yin[1,2]

Affiliations: [1]Department of Neurobiology, Duke University School of Medicine, Durham, NC,

6 27708, USA.,[2]Department of Psychology and Neuroscience, Duke University, Durham, NC,

7 27708, USA.

Corresponding Author: Henry H. Yin

Email: hy43@duke.edu

Keywords: basal ganglia, substantia nigra pars reticulata, motor control, kinematics, reaching

Abstract

The substantia nigra pars reticulata (SNr), the principal output nucleus of the basal ganglia, is traditionally viewed as a binary gate that permits actions through disinhibition. However, this framework fails to explain the fluid, high-dimensional control required for skilled behavior. Using high-resolution 3D kinematic tracking and automated behavioral classification, we show that SNr population activity paradoxically increases during skilled forelimb reaching and scales with motif-specific kinematics. Systematic optogenetic and chemogenetic perturbations reveal that the SNr bidirectionally controls movement vigor and online kinematics. Remarkably, a transient 12.5 ms pause is sufficient to abort a motor program, while a 12.5 ms burst enhances velocity and reshapes trajectories without altering sequence identity. Calcium imaging at single-neuron resolution further confirms that endogenous SNr dynamics represent real-time 3D kinematics. These findings demonstrate that SNr output does not merely gate movement initiation but continuously regulates the stability and kinematic evolution of skilled actions, expanding the functional framework of basal ganglia output from binary selection to the real-time dynamical control of motor execution.

Highlights

- SNr activity paradoxically increases and scales with motor motif kinematics
- Bidirectional SNr control of action vigor for skilled forelimb reaching
- Brief SNr pauses or bursts reshape online motor execution
- The SNr acts as a continuous controller for real-time motor execution

Introduction

The execution of skilled forelimb movements exemplifies the sophistication of mammalian motor control, requiring the brain to select the appropriate action while precisely calibrating its vigor and controlling its kinematics in real-time. While this orchestration involves a distributed network of brain regions[1–8], the basal ganglia (BG) are essential[9,10,3,11–16], with the substantia nigra pars reticulata (SNr) serving as the primary output nucleus[17,18]. The SNr sends inhibitory projections to the thalamus and brainstem[19–24], but the functional role of this output remains unclear.

Classic models of BG function emphasize a gating mechanism. Inhibitory striatal projections to the SNr can produce a pause in the tonically active, inhibitory SNr neurons to permit movement via disinhibition[25,26]. While this model provides a coherent framework for action selection[27–31], it struggles to reconcile the binary nature of gating with the continuous, high-dimensional control required for the execution of skilled motor sequences. Recent evidence suggests that the

striatum, the primary input nuclei of the BG[32], exerts significant influence over movement

vigor[33–35] and online kinematics[36,37,11], yet it remains unclear how the BG shapes movement at

the output stage. Does the SNr merely relay a binary go/no-go signal, as assumed in traditional

models, or does it contribute to the vigor and the real-time kinematic control of movement.

This gap is further accentuated by a paradox in the literature on the SNr: contrary to the

canonical disinhibition model[25,26], some recent observations reveal that many SNr neurons

increase their firing during behavior[16,19,38–42], often outnumbering those that decrease[16,41,42]. This

paradoxical recruitment suggests that the traditional model is incomplete at best. Importantly, it

remains to be determined whether this active recruitment is a mere byproduct of circuit

dynamics or a functional requirement for the execution of skilled behavior.

To address these questions, we developed a high-resolution 3D tracking paradigm and a GRU-

based classifier to decompose the reaching sequence into a series of stereotyped motor motifs.

This high-

vigor, and online kinematics at millisecond temporal resolution. In this study, we provide

evidence that fundamentally challenges the canonical disinhibition model. We reveal a

recruitment of the SNr population that serves as a continuous, real-time drive for execution of

skilled forelimb movements. By momentarily pausing and augmenting SNr activity during

specific behavioral windows corresponding to individual action primitives, we demonstrate that

SNr can continuously shapes the spatial and temporal parameters of movement. Together,

these findings redefine the SNr not as a simple binary gate, but as a dynamic orchestrator of the

fine-grained kinematics in skilled behavior.

Results

Paradoxical increase in SNr population activity underlying stereotyped forelimb reaching

action sequence.

To investigate the contribution of basal ganglia output to the execution of skilled forelimb

movements, we developed a high-resolution water-reaching paradigm for freely moving mice.

This setup leveraged two orthogonal, synchronized high-speed cameras to permit the 3D

kinematic reconstruction of forelimb trajectories (Figure 1A). To achieve unbiased, high-

throughput behavioral analysis, we implemented a Gated Recurrent Unit (GRU)-based

classifier[43] that utilized augmented body-part coordinates to automatically segment reaching

sequences into discrete motor motifs. These segmented labels and their associated kinematics

served as the foundation for all subsequent motif-specific analyses (Figure S1A).

The water-reaching sequence can be consistently decomposed into six distinct, stereotyped

action motifs: Lift, Reach, Open, Grasp, Retract, and Drink (Figure 1B and Video S1). Each

motif was defined by a unique and reproducible kinematic signature (Figure S1B–M). For

instance, the Open and Grasp phases were characterized by minimum proximity to the water

spout (Figure S1F) and a maximum increase in inter-digit distance (Figure S1G). Furthermore,

inter-digit velocity peaked specifically during these dexterous phases, contrasting with the more

ballistic profiles observed during the Reach and Retract segments (Figure S1K, M). The

progression through this motor program was highly organized and reliable, as evidenced by the

structured behavioral ethograms (Figure 1C) and the high probability along the superdiagonal in

the motif transition matrix (Figure 1D). Consistent with this modular organization, 3D trajectory

reconstructions revealed a remarkably stereotyped execution of the reaching sequence across

trials (Figure 1E and Video S2).

To investigate the recruitment of the SNr during skilled forelimb tasks at the population level, we

expressed the genetically encoded calcium indicator jGCaMP7f in the SNr and performed fiber

photometry during the water-reaching task (Figure 1F). Contrary to the classical disinhibition

model, which predicts a pause in SNr output during movement, we observed a robust increase

in both ipsilateral and contralateral SNr population activity during the reaching sequence (Figure

1G). To determine if this activity was coupled to the initiation of the motor program, we stratified

trials by reaction time (fastest 30% vs. slowest 30%). In both hemispheres, the peak of SNr

calcium dynamics tracked the temporal shift in behavior, peaking significantly earlier in fast-

reaction trials compared to slow-reaction trials (Figure 1H).

We next examined whether these population dynamics encoded specific kinematic parameters

of the forelimb. We performed a correlation analysis between SNr activity and three key

kinematic features: motif trajectory length, mean velocity, and peak velocity. SNr activity was

positively correlated with these motif specific kinematics parameters across several motifs, from

the Reach through the Retract motif. Notably, these correlations seem to be more pronounced

in the contralateral hemisphere (Figure 1I), suggesting that the encoding of skilled movements

may be lateralized. Representative correlation plots (Figure 1J) further illustrate that SNr activity

does not merely signal movement onset, but rather scales with the physical execution of the

motor motif.

Collectively, these data demonstrate that, at least at the population level as measured by

photometry recording, an increase, rather than a pause, in SNr neurons is associated with the

execution of stereotyped reaching, scaling with the kinematics of the action sequence.

SNr activation is required for action selection and vigor control of dexterous reaching behavior.

The observed recruitment of the SNr population during reaching suggests that these dynamics are not merely correlative, but are functionally required for motor execution. To test this hypothesis, we employed optogenetic inhibition using the inhibitory opsin GtACR2, delivering photo-stimulation (473 nm, 4 s) at water delivery to silence SNr activity throughout the reaching sequence (Figure 2A).

In striking contrast to classical disinhibition models, which would predict that silencing the SNr might facilitate movement, optogenetic inhibition significantly suppressed reaching, as shown in the reduced success probability (Figure 2B). While the time to success remained unchanged in the sparse subset of successful trials (Figure 2C), the overall behavioral output was severely stunted (Figure 2D-H and Video S3, 4). Motif ethograms (Figure 2D) and quantification of motif initiation-related parameters (Figure 2E–H). revealed a global suppression across all motor motifs. Specifically, SNr inhibition led to a significant decrease in motif occurrence probability (Figure 2E), an increase in latency (Figure 2F), and a reduction in both motif count (Figure 2G) and total duration per trial (Figure 2H).

Beyond action selection, we examined whether the SNr also modulates motor vigor by analyzing forelimb velocity. We found that silencing the SNr resulted in a marked reduction in forelimb velocity across the Lift, Reach, Open, Grasp, and Retract motifs (Figure 2I and Video S4). Quantifying these effects revealed significant decreases in both mean and peak velocities for nearly all motor motifs (Figure 2J–K), suggesting a role of SNr activation in maintaining forelimb vigor.

To further validate the specificity of these effects and demonstrate that reaching behavior is not just being interrupted by the laser inhibition but is functionally titrated by the level of SNr activation, we performed a dose-response analysis by varying the intensity of SNr photoinhibition (1.25 mW, 2.5 mW, 5 mW, and 10 mW; Figure S2A). We observed a robust, intensity-dependent suppression of reaching: higher laser power resulted in lower reach probabilities during the stimulation period (Figure S2B) and a correspondingly higher rebound reaching probability immediately following laser offset (Figure S2C). This titration was further reflected in the latency to reach, which scaled linearly with inhibition intensity (Figure S2D). Moreover, analysis of motif transition dynamics revealed that increasing inhibition intensity progressively suppressed reaching sequence (Figure S2E), characterized by a reduction in the

probability of forward-sequence transitions (along the superdiagonal) and a reciprocal increase

in the probability of no-reach states (leftmost column; Figure S2F). Finally, by focusing on lower

intensities (1.25 mW and 2.5 mW) where movement was not entirely abolished, we confirmed

that motor vigor is also titratable. Greater SNr inhibition led to a gradual, intensity-dependent

reduction in both mean and peak forelimb velocities (Figure S2G–I).

One potential concern with focal optogenetic inhibition in an interconnected GABAergic nucleus

like the SNr is that silencing a localized population could paradoxically activate distant SNr

neurons via the relief of local inhibitory axonal collaterals[44]. To control for this and achieve a

more global silencing of the SNr, we employed an alternative approach using chemogenetics.

We expressed the inhibitory DREADD hM4D(Gi) in the SNr and administered the potent agonist

Deschloroclozapine (DCZ) to inhibit the population more broadly (Figure 2L). While

chemogenetic inhibition yielded a less pronounced effect on action selection (Figure 2M–S and

Video S3-4), compared to high-intensity photoinhibition, its impact on motor vigor was even

more robust (Figure 2T-V). Specifically, we observed no change in overall success probability

(Figure 2M), but a significant increase in the time to success (Figure 2N). While motif probability

and latency remained largely unaffected (Figure 2O–Q), we observed an increase in the number

of motifs per trial accompanied by a significant reduction in the distance traveled for each motif

(Figure 2R, S). This indicates that the individual motor motifs were hypometric, necessitating a

higher number of reaching attempts to successfully reach water. Consistently, chemogenetic

silencing resulted in a marked reduction in forelimb velocity across the Lift, Reach, Open,

Grasp, and Retract motifs (Figure 2T–V). These effects on action selection and kinematic vigor

closely mirrored those seen with low-intensity photoinhibition (Figure S2), reflecting a more

moderate but widespread neural suppression. Ultimately, these chemogenetic results confirm

that the suppression of skilled reaching stems from the inhibition of SNr output rather than

unintended collateral activation.

Collectively, these data demonstrate that rather than facilitating movement through a pause of

inhibition, as suggested by classical models, the activation of the SNr is required for action

selection and vigor control of dexterous reaching behavior.

Post-inhibitory rebound reaches exhibit enhanced vigor

The observed recruitment of the SNr population during reaching, coupled with the suppressive

effects of its inhibition, suggests that SNr activation is a key driver of dexterous motor execution.

Given that higher photoinhibition intensities triggered rebound reach robustly upon laser offset

(Figure S2C), and since prolonged photoinhibition is frequently associated with a compensatory
burst of rebound neural activity, we hypothesized that this post-inhibitory rebound neural activity
might actively promote subsequent reach. To test this, we silenced the SNr using GtACR2 (473
nm, 5 mW) for randomly interleaved durations of 1 s, 2 s, 3 s, or 4 s (Figure S3A). This protocol
reliably suppressed reaching during the laser window and triggered immediate rebound
reaching upon offset, regardless of the prior inhibition duration (Figure S3B–D). This temporal
precision allowed us to isolate and characterize the specific kinematic signatures of motor
behavior driven by a post-inhibitory state. While rebound reaches maintained a normal motif
progression (Figure S3E) and achieved success probabilities comparable to control trials
(Figure S3F), their latency to success was significantly reduced in rebound trials (Figure S3G), a
result that stands in direct opposition to the increased latency observed during SNr inhibition
(Figure 2N). Further motif ethogram analysis revealed that these rebound reaches were more
vigorous than their control counterparts (Figure S3H). Moreover, we observed a significant
increase in forelimb velocity during the initial phases of the sequence, specifically during the Lift,
Reach, and Open motifs (Figure S3I–K). Collectively, these results demonstrate that the neural
rebound following SNr inhibition drives rebound reaching behavior with enhanced kinematic
signatures. This potentiation of motor vigor provides further causal evidence that SNr activation,
rather than inhibiting movement, acts as a potent promoter of skilled forelimb reaching.

SNr activation is a critical, frequency-dependent driver of the kinematic vigor for
dexterous reaching.
To directly test the hypothesis that SNr activation promotes rather than suppresses skilled motor
execution, we employed parallel gain-of-function strategies. We performed focal optogenetic
activation using ChR2 (473 nm, 4 s starting at onset of water delivery) and, to account for
potential artifacts of local collateral-mediated inhibition, we utilized a more global chemogenetic
approach expressing the excitatory DREADD hM3D(Gq) in SNr (Figure 3A, L).

Overall, we observed consistent promoting effects across both optogenetic and chemogenetic
activation (Video S3-4). Neither focal nor global activation impaired behavioral performance with
success probability unaffected by activation (Figure 3B, M), and motif occurrence probabilities
and latencies were largely unaffected (Figure 3E, F and P, Q). These findings contradict a
recent report suggesting that SNr activation impairs reaching[16], instead pointing toward an
opposite role in skilled forelimb movement. Consistent with this, we observed a significant
reduction in the latency to success (Figure 3C, N) and a pronounced leftward shift in the
behavioral ethograms (Figure 3D, O), indicating that SNr activation accelerates the progression

of the motor program. Detailed kinematic quantification revealed that this enhancement was
characterized by motif-specific modulations of duration, distance travelled, and velocity. During
the ballistic phases of the sequence (Lift, Reach, and Retract), SNr activation significantly
increased movement velocity (Figure I-K and T-V) while simultaneously shortening motif
duration (Figure 3G, R). As a result, the total distance traveled during these segments remained
relatively stable (Figure 3H, S). In contrast, during the more dexterous middle phases (Open
and Grasp), the increase in velocity (Figure 3G, R) was not accompanied by a reduction in
duration (Figure I–K and T–V). Consequently, these motifs exhibited a significant increase in
trajectory length (Figure 3H, S), driven by the sustained increase in velocity.

To further validate the specificity of these effects and demonstrate that reaching behavior is
functionally titrated by the level of SNr activation, we performed a dose-response analysis by
varying the frequency of SNr photoactivation (20 Hz and 40 Hz; Figure S4A). Consistent with
our main findings, neither reach probability nor the latency to reach were affected by either
frequency (Figure S4B–D). Intriguingly, we even observed a slight but significant improvement
on success probability during 20 Hz stimulation (Figure S4E). Both activation frequencies
produced a comparable reduction in the latency to success (Figure S4F) and a pronounced
leftward shift in the behavioral ethograms (Figure S4H), further establishing that SNr activation
accelerates motor program progression. Finally, detailed kinematic quantification confirmed that
movement vigor can be titrated. Greater SNr activation frequencies led to a gradual, frequency-
dependent increase in both mean and peak forelimb velocities across motifs, from the Reach
through the Retract segments (Figure S4I–K).

Collectively, these data provide causal evidence that SNr activation, contrary to classical
disinhibition theory and recent findings[16], acts as a motif-specific driver of motor vigor, where
activity levels causally scale with the kinematic vigor of skilled forelimb reaching.

SNr is not involved in the trial-to-trial refinement and reinforcement of forelimb dexterous
reaching behavior.
While our results establish that SNr activity actively modulates the execution of reaching
through action selection and vigor control, it remains unclear how it affects future behavior, e.g.
reinforcement of the motor program. Given that reaching is a learned behavior driven by
exploration and reinforcement[45] and that the basal ganglia output is involvement in reinforcing[46]
and punishment[47], we tested whether SNr optogenetic perturbations leave a functional
reinforcement or punishment signal on subsequent motor performance. To investigate this, we

performed a trial-history analysis comparing control trials immediately following a
photostimulation trial (Post Stim) to those following a control trial (Post Ctr). We reasoned that if
the SNr were involved in the refinement and reinforcement of the motor program, the behavioral
profile of the Post Stim trial would deviate from that of the Post Ctr trial. This analysis was
applied to both the optogenetic inhibition (Figure S5A) and activation (Figure S6A) datasets.
Across both datasets, we found that the reaching profiles for Post Ctr and Post Stim trials
remained virtually identical (Figures S5 and S6). In the inhibition dataset, all metrics, including
reach probability and latency, success probability and latency, motif transition matrix, motif
ethogram, and 3D kinematics, were statistically indistinguishable between trial types (Figure
S5B–J). Similarly, in the activation dataset, the motor program exhibited remarkable stability;
with the exception of a minor decrease in reach latency (Figure S6C), all other primary features
remained unchanged (Figure S6B–J). Collectively, the lack of carry-over effects suggests that
the SNr does not play a role in the reinforcement and refinement of the forelimb motor program.

A momentary pause in SNr activity abolishes the ongoing reaching sequence, revealing
its obligatory role in real-time motor execution.
Having demonstrated that SNr activity is necessary for the overall selection and vigor of
reaching, we next asked whether this activity is required for the continuous, moment-by-moment
control of the motor sequence. To address this, we applied transient pulses of photoinhibition,
either 100 ms (Figure 4A) or 12.5 ms (Figure 4I), timed to coincide with reaching motifs. All trials
where a motif (from Lift through Retract) coincided with the pulse onset were included in the
analysis.

Consistent with our long-term inhibition results, even a momentary pause in SNr activity was
sufficient to disrupt the progression of the reaching sequence (Figure 4B-D and J-L; Video S5,
6). Motif transition analysis revealed a reduction in forward-sequence transitions along the
superdiagonal, coupled with a reciprocal increase in no-reach states (leftmost column; Figure
4B, J). This behavioral disruption was further evidenced by a significant reduction in both motif
duration and the distance traveled for motifs targeted by the laser pulse (Figure 4C, D and K, L).
Furthermore, laser-onset-aligned 3D trajectory (Figure 4E, M), XYZ coordinate profiles (Figure
4F, N), and distance to water spout (Figure 4G, O) clearly illustrated that the reaching
movement was abolished upon laser onset, with the duration of the behavioral disruption and
subsequent recovery scaling with the length of the inhibitory pulse. Kinematic analysis revealed
a complex velocity profile following the pause: a brief, transient increase in velocity was followed
by a sustained decrease (Figure 4H, P). We reasoned that the initial velocity spike was not a

direct consequence of SNr silencing, as its duration remained constant regardless of the pulse

length. In contrast, the duration of the subsequent velocity reduction scaled precisely with the

inhibition period, confirming that the pause of SNr activity directly reduced motor velocity in real-

time. Strikingly, while the 100 ms pause induced a more profound disruption across all

measured parameters (Figure 4B–H)., even the brief 12.5 ms inhibition was sufficient to abolish

the ongoing motor program (Figure 4J–L).

To investigate whether this moment-by-moment control exhibits motif-specific sensitivity, trials

were grouped post-hoc based on the specific action motif (Lift, Reach, Open, Grasp, or Retract)

coincident with the 12.5 ms pulse onset (Figure S7A). We found that the requirement for SNr

drive was non-uniform across the motor program. The ongoing action sequence was abolished

during the outbound phases (Lift, Reach, and Open), followed by a re-initiation of the reach

upon recovery, while the inbound phases (Grasp and Retract) were notably less affected in their

sequential progression (Figure S7B). Correspondingly, laser-onset-aligned 3D trajectories

(Figure S7C), coordinate profiles (Figure S7D), and distance-to-target metrics (Figure S7E)

revealed a profound vulnerability from the Lift through the Grasp motifs, whereas the Retract

motif remained relatively more resilient to the transient pause in SNr activity.

These findings establish that the SNr provides a continuous, motif-specific drive essential for the

moment-by-moment execution of skilled reaching. Our results demonstrate that basal ganglia

output is not merely a transient trigger for movement initiation or a tonic modulator of baseline

vigor; rather, it is required to maintain the structural integrity and kinematic flow of the motor

sequence in real-time.

Transient SNr bursts alter the kinematics of ongoing reaching in real-time

While we have established that SNr is required for the moment-by-moment execution of skilled

forelimb reaching, it is unknown if this signal serves an instructive role in real-time. To probe it,

we applied transient pulses of photoactivation, either 300 ms (Figure 5A) or 12.5 ms (Figure 5I),

timed to coincide with reaching motifs. All trials where a motif (from Lift through Retract)

coincided with the pulse onset were included in the analysis.

Consistent with our long-term activation results, we found that a momentary burst in SNr activity

did not impair the structural progression of the reaching sequence (Figure 5B, J and Video 5, 6).

However, laser-onset-aligned 3D trajectory (Figure 5C, K), XYZ coordinate profiles (Figure 5D,

L), and distance to water spout (Figure 5E, M) illustrated that the reaching movement was

significantly altered upon laser onset. The altered kinematics was characterized by a distinct

spatial protrusion in the trajectory (Video 5, 6), with the duration of the altered kinematics
scaling with the length of the excitatory burst (Figure 5C-E and K-M). Another striking evidence
of this real-time control was observed in the velocity profiles. While the 300 ms burst induced a
substantially longer period of velocity enhancement, the momentary 12.5 ms burst, a duration
representing only a fraction of a single motor motif, was nonetheless sufficient to drive an
immediate and significant increase in forelimb velocity (Figure 5F, N). Detailed quantification
across motifs confirmed that these excitatory bursts significantly enhanced both mean and peak
forelimb velocities for most motifs expect Lift (Figure 5G, H and O, P).

Similarly, we grouped trials post-hoc based on the specific action motif (Lift, Reach, Open,
Grasp, or Retract) coincident with the 12.5 ms pulse onset (Figure S8A). In contrast to the
behavioral abolishment effect observed during SNr transient inhibition (Figure S7), the structural
progression of the action sequence remained largely uninterrupted during brief activation bursts
(Figure S8B). However, laser-onset-aligned 3D trajectories (Figure S8C), coordinate profiles
(Figure S8D), and velocity metrics (Figure S8E) revealed an altered reaching kinematics from
the Reach through the Retract motifs, whereas the Lift showed no significant deviation in its
spatial trajectory or velocity profile (Figure S8C–E).

Collectively, these results reveal that rather than disrupting the progression of the forelimb
sequence, a momentary burst of SNr activity preserves the structural integrity of the reach while
reshaping its detailed kinematics. This demonstrates that SNr output provides a high-resolution,
instructive signal that modulates the spatial and velocity profiles of reaching behavior on a
moment-by-moment basis.

Endogenous SNr population dynamics continuously represent real-time 3D kinematics
Having established that SNr activity is necessary and sufficient to drive 3D kinematics, we next
investigated how endogenous SNr population dynamics encode these precise, moment-by-
moment movements. To achieve this, we expressed the genetically encoded calcium indicator
jGCaMP7f in the SNr and performed in vivo miniscope imaging to monitor calcium dynamics in
GABAergic projection neurons at single-cell resolution during the reaching behavior (Figure 6A).
Consistent with our fiber photometry results (Figure 1F-H), the population-averaged calcium
signal revealed a robust increase in peri-reach activity (Figure 6B). At the single-cell level, we
observed a heterogeneous mix of movement-aligned responses (Figure 6C). While movement-
inhibited neurons slightly outnumbered movement-excited neurons during the initial "Lift" phase,
this balance shifted as the movement progressed, with movement-excited neurons dominating

the subsequent reach, grasp, retract, and drink motifs (Figure 6D). To determine how individual

neurons, modulate distinct behavioral motifs, we sorted the population based on modulation

magnitude during the Retract phase. We observed that population activity evolved progressively

and smoothly throughout the forelimb movement sequence (from Lift to Retract). In contrast, we

noted a sudden, distinct shift in population dynamics when transitioning into the orofacial "Drink"

motif, suggesting distinct encoding strategies for forelimb versus consummatory movements

(Figure 6E). To quantify the kinematic information embedded within this network, we trained

linear regression models to predict real-time 3D kinematics using the single-cell population

activity. Strikingly, the population activity robustly and continuously predicted real-time 3D

trajectories (X, Y, and Z coordinates) as well as movement velocity (Figure 6F, G). Furthermore,

individual neurons exhibited distributed and diverse functional weights across these different

kinematic measurements (Figure 6H). Taken together, these data demonstrate that intrinsic in

vivo SNr population activity contains rich, predictive representations of movement, acting as a

continuous controller for moment-by-moment 3D kinematics.

Discussion

Our findings establish the substantia nigra pars reticulata (SNr) as a key component of a real-

time, high-dimensional controller for skilled behavior. By integrating high-resolution 3D tracking

with precisely timed perturbations and calcium imaging, we move beyond the categorical gating

model to reveal a more dynamic role for basal ganglia output. We demonstrate that SNr

dynamics do not merely permit movement initiation but actively dictate its kinematic flow. The

paradoxical recruitment of this inhibitory output during reaching, coupled with its requirement for

the moment-by-moment maintenance of motor trajectories, identifies the SNr as a critical

instructive signal. By concurrently governing action selection, kinematic vigor, and real-time

motor form, the SNr ensures the structural integrity and fluid execution of the complex action

sequences that define motor expertise.

For decades the action selection model has dominated the study of basal ganglia function[27–

29,31]. In this framework, the direct and indirect pathways are thought to exert opposing influences

on the tonically active SNr to either permit or suppress motor output through downstream

disinhibition[25,26]. While the discovery of co-activation between these pathways[48] during

movement led to a refined model[18,27,49,50], where the direct pathway promotes desired actions

and the indirect pathway suppresses competitors, recent evidence suggests a far more complex

architecture. Specifically, the observation that the vast majority of SNr neurons show congruent,

increased activity across disparate motor modalities[51], reaching and licking, is difficult to

reconcile with the refined action selection mode, which predicts that distinct motor behaviors

result from opposite modulation of the firing activity of same neurons. Our study extends these

observations by providing causal evidence that the SNr serves as more than a categorical

switch. Through bidirectional optogenetic and chemogenetic manipulations, we demonstrate

that the net activity of the SNr is engaged in both the selection of action and the titration of its

kinematics (Figure 2, 3 and Figure S2-S4). Most importantly, by using transient optogenetics to

perturb millisecond-scale dynamics, we reveal a critical role for the SNr in the continuous, online

control of motor form (Figure 4, 5 and Figure S7, S8). Together, these results suggest that the

SNr does not simply select and gate motor behavior, but actively orchestrates the detailed

physical parameters of its moment-by-moment execution.

According to the dominant model, the SNr facilitates movement through a pause in its tonic

firing, thereby disinhibiting downstream motor centers[25,26]. However, we demonstrate that this

mechanism cannot fully account for action selection, because SNr population activity increases,

rather than decreases, during skilled forelimb movement (Figure 1 and Figure 6). As this finding

contradicts the canonical disinhibition model, we examined alternative circuit mechanisms that

might reconcile our findings with the traditional framework. First, we addressed whether the

observed population increase is a non-functional byproduct, masking a smaller, inhibited subset

of neurons that actually drives the behavior. Second, we considered whether SNr activation

might locally inhibit neighboring projection neurons via inhibitory axonal collaterals[44], thereby

inducing net downstream disinhibition. By using chemogenetics to modulate the SNr output

globally and presumably minimizing local collateral competition, we observed behavioral effects

comparable to our focal optogenetic results (Figure 2 and Figure 3). This suggests that the

collective activity of the SNr, rather than a specific sub-population or modulation by axonal

collaterals, plays a key role in shaping the skilled forelimb movement. In addition, we

investigated whether this effect arose from post-stimulation compensation, specifically post-

inhibitory rebound or post-excitatory depression, as suggested by previous work[52]. To address

this, we used both chemogenetics (Figure 2 and Figure 3) and transient optogenetic

perturbations (Figures 4 and Figure 5) to minimize compensatory SNr network dynamics.

Through these complementary approaches, we demonstrate that the observed behavioral

changes are a direct consequence of SNr drive, rather than an artifact of post-stimulation

compensation. Finally, it remains possible that SNr neurons target inhibitory interneurons in

downstream nuclei, providing an additional layer of nested disinhibition[53]. While our functional

data cannot rule out this possibility, future studies employing cell-type-specific tracing and
recording in downstream targets will be essential to map these functional connections.

Despite the long-standing popularity of the gating model, direct in vivo evidence for functional
disinhibition has remained surprisingly elusive. While studies have identified movement-related
firing decreases in the SNr and corresponding firing increases in its postsynaptic targets[25,26],
recent physiological surveys have found only a limited number of putative monosynaptically
connected SNr-target pairs that exhibit clear inhibitory effects[16]. More tellingly, the latencies of
movement-related activity changes in the SNr and its postsynaptic targets are often
indistinguishable[16]. This lack of temporal offset is difficult to reconcile with a sequential
disinhibition model, which would necessitate a clear delay between the release of the inhibition
and the onset of downstream activity. Moreover, recent evidence suggests that the SNr has only
a modulatory influence in vivo[54], rather than showing decisive disinhibition as posited by the
traditional model.

One way to reconcile the paradoxical finding that SNr excitation increases vigor, whereas even
a 12.5 ms pause can abort an ongoing reach, is to interpret its output in dynamical rather than
purely level-based terms. The striatonigral pathway has been proposed to transform transient,
velocity-related striatal signals into sustained motor commands from the SNr output[19,33]. In such
integrator-
deviations from baseline decay following drive. Tonic SNr firing sets an operating point, whereas
intrinsic leak reflects network stability, potentially shaped by recurrent inhibition within SNr and
its connections with the rest of the BG. Our observation that global excitation accelerates
reaching without impairing action selection (Figure 3, 5 and Figure S4), whereas inhibition
collapses velocity and truncates motif progression (Figure 2, 4 and Figure S2, S3), is consistent
with the possibility that SNr manipulations alter this dynamical stability. Global SNr excitation
may shift the balance of channel-specific competition and recurrent inhibitory feedback,
effectively reducing leak. In an integrator, reduced leak would allow inputs to accumulate more
quickly, increasing movement velocity despite elevated absolute inhibition. Conversely, brief
pauses would transiently destabilize the accumulated motor command, causing it to decay
toward baseline and resulting in rapid termination of the ongoing movement. The known
intranigral inhibitory microcircuit, which has been proposed to implement forms of feedback
gain[44], provides a plausible substrate for such stability modulation. Although this account
remains speculative at present, the traditional model of BG fails to account for our observations.
Any comprehensive model of basal ganglia function must therefore account for the critical role

of SNr activity, as it dynamically stabilizes, accelerates, or terminates ongoing motor commands
on a millisecond timescale.

Methods

Mice

All experimental procedures were conducted in accordance with the guidelines and with the
approval of the Institutional Animal Care and Use Committee (IACUC) at Duke University
(Protocol # 162-22-09). Adult mice (3–10 months of age) were maintained on a 12-hour light/dark
cycle with ad libitum access to food and water until the start of behavioral training. Following
surgical procedures, mice were singly housed and provided with environmental enrichment. This
study utilized both male and female VGAT-Cre and PV-Cre transgenic lines. For behavioral
experiments, mice were placed under a water restriction protocol, with their body weight
monitored daily and maintained at 85–90% of their baseline weight.

Stereotaxic injection and surgical procedures

Mice were initially anesthetized in an induction chamber with 2.0–3.0% isoflurane and
subsequently secured in a stereotaxic frame (David Kopf Instruments, Tujunga, CA). Subjects
received intraperitoneal injections of meloxicam (2 mg/kg) and local administration of bupivacaine
for analgesia, with anesthesia maintained at 1.0–1.5% isoflurane throughout the procedure.
Unilateral or bilateral craniotomies were drilled above the substantia nigra pars reticulata (SNr) at
AP -3.7 mm and ML ±1.5 mm relative to bregma. Viral vectors were infused via a microinjector
(Nanoject 3000, Drummond Scientific) at a rate of 1 nL/s at DV -4.4 mm relative to bregma.
Following injection, the pipette remained at the target coordinate for 10 minutes to ensure
adequate viral diffusion and absorption. For chemogenetic experiments, 50 nL of AAV2-hSyn-
DIO-hM4D(Gi)-mCherry (Addgene #44362) or AAV2-hSyn-DIO-hM3D(Gq)-mCherry (Addgene
#44361) was infused bilaterally into the SNr of VGAT-Cre or PV-Cre mice. For optogenetic
experiments, 100 nL of AAV1-hSyn1-SIO-stGtACR2-FusionRed (Addgene #105677) or AAV5-
EF1a-DIO-hChR2(E123T/T159C)-EYFP (Addgene #35509) was injected bilaterally into the SNr
of VGAT-Cre or PV-Cre mice. Two custom-made optic fibers (105 µm core diameter, >85%
transmittance, 5 mm length below the ferrule) were then implanted above the SNr at AP -3.7 mm,
ML ±1.5 mm, and DV -4.4 mm relative to bregma. For fiber photometry, 100 nL of AAV9-Syn-
FLEX-jGCaMP7f (Addgene #104488) was infused bilaterally into the SNr, followed by the
implantation of an optic fiber (400 µm core diameter, 5 mm length below the ferrule; RWD) using
the same coordinates as the optogenetic protocols. Fibers were secured to the skull using dental
acrylic. For miniscope calcium imaging, 100 nL of AAV9-Syn-FLEX-jGCaMP7f (Addgene

#104488) was injected unilaterally into the left SNr. A GRIN lens (6 mm x 0.5 mm; Inscopix) was
implanted above the left SNr (AP -3.7 mm, ML -1.5 mm, DV -4.4 mm) and secured with dental
acrylic. A baseplate for miniscope attachment was affixed to the GRIN lens 2–3 weeks post-
surgery following verification of viral expression. All mice were allowed to recover for a minimum
of three weeks before the commencement of behavioral experiments.

Histology

Following the conclusion of all behavioral experiments, mice were sacrificed to confirm viral
expression and verify the precise anatomical placement of optic fibers and GRIN lenses. Mice
were anesthetized with 3.0% isoflurane until immobile and transcardially perfused with 0.1 M
phosphate-buffered saline (PBS), followed by 4% paraformaldehyde (PFA) in PBS. Brains were
carefully dissected and post-fixed in 4% PFA at 4 °C. After 24 hours, the tissue was transferred to
a 30% sucrose solution for cryoprotection. Coronal sections (60 µm) were obtained using a
cryostat (Leica CM 1850). Brain slices were subsequently mounted and coverslipped using
Fluoroshield with DAPI (Abcam #Ab104139) to visualize cell nuclei. Fluorescent images were
acquired using a ZEISS microscope (Axio Zoom.V16, RRID:SCR_016980).

Water-reaching task

Water-restricted mice were trained in a custom-built, transparent acrylic chamber (10 cm L × 10
cm W × 15 cm H) featuring a vertical slit (8.5 mm W × 40 mm H) on the front wall. The dimensions
of the slit were designed to permit reaching movements using only the right forelimb while
preventing direct access to the waterspout for licking. Outside the chamber, a blunted 21G needle
was used as a waterspout to deliver ~5 µL water droplets via a solenoid valve (NResearch). Each
trial was initiated with an auditory cue, and the spatial position of the spout was fixed for each
subject using a three-axis manual micromanipulator. Mice were provided a 7 s response window
to reach before the spout was retracted, with inter-trial intervals (ITI) randomized between 2 and
15 s. To capture the kinetics of the forelimb reaching sequence, two high-speed cameras (DMK
33UX287; The Imaging Source) were positioned perpendicularly and synchronized to record
behavior at 500 Hz. An infrared (IR) light source was mounted near the cameras to ensure high-
contrast illumination for kinematic tracking.

Behavioral classification and kinematics quantification

To achieve high-resolution tracking of forelimb kinematics, two deep neural networks were trained
independently using DeepLabCut (https://github.com/DeepLabCut) to digitize specific anatomical
landmarks (digits 2–4, paw, mouth, and nose) across both camera views. Three-dimensional (3D)
coordinates were subsequently reconstructed using the DeepLabCut 3D support pipeline

following camera calibration with a checkerboard pattern[55]. These reconstructed coordinates were
utilized to compute pairwise inter-bodypart coordinates and were further augmented with derived
kinematic features, including instantaneous velocity and acceleration. The centroid of the three
digits and the paw was used to represent the 3D coordinates of forelimb. To facilitate unbiased,
high-throughput behavioral analysis, we implemented a Gated Recurrent Unit (GRU)-based
classifier. This model utilized the augmented feature sets (coordinates, velocity, and acceleration)
as inputs to automatically segment reaching sequences into seven discrete motor motifs: Lift,
Reach, Open, Grasp, Open, Retract, and Drink. The GRU network consisted of a single hidden
layer with 256 units, and was trained using a cross-entropy loss function on a large-scale,
manually annotated dataset comprising 1,237,503 frames. Each identified motif is characterized
by a unique and reproducible kinematic profile (Figure S1). To evaluate motor performance, motif-
specific metrics were first calculated for each occurrence within a trial and then averaged to
produce a single representative value per trial. The following kinematic and temporal parameters
were defined. Motif number: the total count of motif occurrence per trial; Motif latency: the time
elapsed from the water delivery to the first occurrence of a specific motif; Motif duration: the time
spent in a single occurrence of a specific motif; Motif total duration: the cumulative time spent in
a specific motif per trial; Motif distance travelled: the cumulative length of 3d trajectory traveled
by the forelimb during a specific motif; Motif mean/peak velocity: the average or maximum velocity
of the forelimb during a specific motif. We further defined trial-wide metrics to quantify global
behavior. Latency to reach: the time from water delivery to the first Lift-Reach sequence; Latency
to success: time from first Lift-Reach sequence to the first Drink motif with a duration > 600 ms.
Every single parameter from each trial was then averaged to get a single numerical quantification
for that session. Session-level quantifications included: Reach probability (percentage of trials
containing at least one Lift-Reach sequence); Success probability (percentage of reaching trials
resulting in a Drink motif > 600 ms); and Motif probability (percentage of trials containing at least
one occurrence of a specific motif).

Optogenetic experiments

528
Optogenetics were performed using a 473 nm blue laser (Shanghai Laser & Optics Century Co.,
Ltd.) delivered bilaterally, unless otherwise specified, during randomized trials. For control trials,
we assigned pseudo-laser start and end times that matched the exact temporal distribution of
stimulation trials within the same session to ensure rigorous quantification and comparison. All
motif-specific metrics were analyzed across three distinct windows: Within-Laser, After-Laser, and
Whole-trial as indicated in the figures. We conducted three sets of optogenetic inhibition
experiments: 1) for fixed duration and varying intensity, a 4 s laser was delivered at water onset

in 33% of trials at intensities of 10 mW, 5 mW, 2.5 mW, or 1.25 mW; 2) for fixed intensity and
varying duration, a 5 mW laser was delivered at water onset for 1 s, 2 s, 3 s, or 4 s in 33% of
trials; and 3) for momentary pulse inhibition, brief pulses of 12.5 ms or 100 ms were delivered in
60% of trials at optimized, mouse-specific latencies relative to water delivery to intercept specific
behavioral motifs. We also performed two sets of optogenetic activation experiments using a 10
mw laser: 1) for fixed duration and varying frequency, a 4 s laser pulse (2 ms pulse width at 40
Hz or 20 Hz) was delivered at water onset in 25% of trials; and 2) for momentary pulse excitation,
a laser of 12.5 ms (constant) or of 300 ms (40 Hz, 2 ms pulse width) were delivered in 50% of
trials at calibrated latencies to target reaching motifs.

Chemogenetic experiments
Chemogenetic manipulations were performed using Deschloroclozapine[56] (DCZ; Hello Bio
#HB9126), a potent and highly selective actuator for hM3D(Gq) and hM4D(Gi) DREADD
receptors. Each experimental session began with 75 baseline trials to establish pre-treatment
performance levels. To ensure a non-invasive administration and minimize stress-induced
behavioral artifacts, mice were allowed to voluntarily consume 100 µL of water containing either
saline (control) or DCZ (0.2 µg/g body weight). Following a 6-minute absorption period to allow
for rapid ligand uptake, an additional 75 trials were recorded to assess the effects of SNr activation
or inhibition on forelimb skilled movement. Control and DCZ sessions were performed on
consecutive days, with the saline session serving as a within-subject control for the subsequent
chemogenetic manipulation.

Fiber photometry
Following a three-week recovery period for viral expression of jGCaMP7f, cell-type-specific
calcium activity in the SNr was recorded during the water-reaching task. Fluorescent signals were
acquired using a multi-channel fiber photometry system (RWD Life Science). Two light-emitting
diodes (LEDs) were used: a 470 nm LED to excite the calcium-dependent jGCaMP7f signal and
a 410 nm LED as an isosbestic control to account for movement artifacts and photo-bleaching.
Light was delivered through a dichroic mirror and a multimode fiber-optic patch cord (400 µm
core, 0.5 NA), with power at the fiber tip maintained between 40 mW to minimize phototoxicity.
The emitted fluorescence was collected through the same patch cord, filtered, and detected by a
femtowatt photoreceiver. The isosbestic 410 nm signal was fitted to the 470 nm signal to calculate
the change in fluorescence ($\Delta F/F$), defined as ($F_{recorded}$ - F_{fitted}/ F_{fitted}). The resulting $\Delta F/F$ traces
were Z-score normalized and temporally aligned to the behavioral metadata. The mean and peak
calcium activity were calculated for each occurrence of the motor motifs (Lift, Reach, Open,

Grasp, Open, and Retract) to evaluate the correlation between SNr population activity and
kinematics of specific motor motifs.

Miniscope Imaging
Following a three-week recovery period to ensure robust viral expression of jGCaMP7f, calcium
dynamics in SNr projection neurons were recorded at single-cell resolution during the free-moving
water-reaching task. Imaging was performed using a custom-built miniature fluorescence
microscope paired with custom data acquisition software. To minimize stress and tethering-related
behavioral artifacts, mice were extensively habituated to the behavioral arena while wearing a
weight-matched dummy miniscope for several days prior to the start of recording experiments. To
mitigate photobleaching of the calcium indicator and prevent potential phototoxicity, recording
sessions were strictly capped at 60 trials per animal. Raw calcium imaging videos were
preprocessed and analyzed using the MiniAn Python package[57]. The standard processing
pipeline included motion correction to remove movement artifacts, background subtraction, and
constrained non-negative matrix factorization (CNMF) to extract the spatial footprints and
calculate temporal calcium traces of individual neurons. For trial-based analysis, the baseline
activity for each trial was defined as the mean fluorescence signal during a predefined window (
-2.0 s to -0.4 s relative to water delivery onset). To quantify the magnitude of behavioral
modulation, a ΔZscore was calculated for each neuron by subtracting the baseline mean Zscore
from the mean Zscore recorded during specific action motifs. The statistical significance of this
neuronal modulation was evaluated using a Wilcoxon signed-rank test. Finally, to evaluate the
relationship between neural activity and continuous behavior, a linear regression model was
trained to predict moment-by-moment 3D kinematic variables (e.g., XYZ spatial coordinates and
forelimb velocity) utilizing the concurrently recorded SNr population activity as inputs.

Quantification and statistical analyses
All data are presented as mean ± SEM unless otherwise specified. Computational quantification
and statistical analyses were performed using custom scripts developed in Python. Due to the
non-normal distribution of behavioral and neural datasets, non-parametric statistical tests were
employed throughout the study, unless otherwise noted. Specifically, for comparisons between
two paired groups, the Wilcoxon signed-rank test was applied. Comparisons involving multiple
groups or experimental conditions were evaluated using the Friedman test. Spearman's rank
correlation coefficient was used for correlation analysis. Results were considered statistically
significant at $p < 0.05$. Comprehensive details regarding the specific statistical tests, exact p-
values, and sample sizes are summarized in Supplementary Table 2.

Resource Availability

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Henry H. Yin (hy43@duke.edu).

Materials Availability

This study did not generate new unique reagents.

Data and Code Availability

All data reported in this paper will be shared by the lead contact upon request.

All code used in this paper will be shared by the lead contact upon request.

Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Acknowledgments

This research was supported by NIH grant R01NS094754. We thank Rebecca Chung-Hui Yang and Lindsey Glickfeld for reading the manuscript and providing constructive feedback.

Author Contributions

SR and HHY conceived the study and designed experiments. SR built setups, collected data, established data-analysis pipelines, analyzed data, and wrote the manuscript. HHY supervised the project and revised the manuscript.

Declaration of Interests

The authors declare no competing interests.

Reference

1. Wang, X., Liu, Y., Li, X., Zhang, Z., Yang, H., Zhang, Y., Williams, P.R., Alwahab, N.S.A., Kapur, K., Yu, B., et al. (2017). Deconstruction of Corticospinal Circuits for Goal-Directed Motor Skills. Cell *171*, 440-455.e14. https://doi.org/10.1016/j.cell.2017.08.014.

2. Galiñanes, G.L., Bonardi, C., and Huber, D. (2018). Directional Reaching for Water as a Cortex-Dependent Behavioral Framework for Mice. Cell Reports *22*, 2767–2783. https://doi.org/10.1016/j.celrep.2018.02.042.

3. Lemke, S.M., Ramanathan, D.S., Guo, L., Won, S.J., and Ganguly, K. (2019). Emergent modular neural control drives coordinated motor actions. Nat Neurosci *22*, 1122–1131. https://doi.org/10.1038/s41593-019-0407-2.

4. Becker, M.I., and Person, A.L. (2019). Cerebellar Control of Reach Kinematics for Endpoint Precision. Neuron *103*, 335-348.e5. https://doi.org/10.1016/j.neuron.2019.05.007.

5. Sauerbrei, B.A., Guo, J.-Z., Cohen, J.D., Mischiati, M., Guo, W., Kabra, M., Verma, N., Mensh, B., Branson, K., and Hantman, A.W. (2020). Cortical pattern generation during dexterous movement is input-driven. Nature *577*, 386–391. https://doi.org/10.1038/s41586-019-1869-9.

6. Guo, J.-Z., Sauerbrei, B.A., Cohen, J.D., Mischiati, M., Graves, A.R., Pisanello, F., Branson, K.M., and Hantman, A.W. (2021). Disrupting cortico-cerebellar communication impairs dexterity. eLife *10*, e65906. https://doi.org/10.7554/eLife.65906.

7. Ruder, L., Schina, R., Kanodia, H., Valencia-Garcia, S., Pivetta, C., and Arber, S. (2021). A functional map for diverse forelimb actions within brainstem circuitry. Nature *590*, 445–450. https://doi.org/10.1038/s41586-020-03080-z.

8. Yang, W., Kanodia, H., and Arber, S. (2023). Structural and functional map for forelimb movement phases between cortex and medulla. Cell *186*, 162-177.e18. https://doi.org/10.1016/j.cell.2022.12.009.

9. Fricker, R.A., Annett, L.E., Torres, E.M., and Dunnett, S.B. (1996). The Placement of a Striatal Ibotenic Acid Lesion Affects Skilled Forelimb Use and the Direction of Drug-Induced Rotation. Brain Research Bulletin *41*, 409–416. https://doi.org/10.1016/S0361-9230(96)00083-4.

10. Kendall, A.L., David, F., Rayment, G., Torres, E.M., Annett, L.E., and Dunnett, S.B. (2000). The influence of excitotoxic basal ganglia lesions on motor performance in the common marmoset. Brain *123*, 1442–1458. https://doi.org/10.1093/brain/123.7.1442.

11. Dhawale, A.K., Wolff, S.B.E., Ko, R., and Ölveczky, B.P. (2021). The basal ganglia control the detailed kinematics of learned motor skills. Nat Neurosci *24*, 1256–1269. https://doi.org/10.1038/s41593-021-00889-3.

12. Lopez-Huerta, V.G., Denton, J.A., Nakano, Y., Jaidar, O., Garcia-Munoz, M., and Arbuthnott, G.W. (2021). Striatal bilateral control of skilled forelimb movement. Cell Reports *34*, 108651. https://doi.org/10.1016/j.celrep.2020.108651.

13. Inase, M., Buford, J.A., and Anderson, M.E. (1996). Changes in the control of arm position, movement, and thalamic discharge during local inactivation in the globus pallidus of the monkey. Journal of Neurophysiology. https://doi.org/10.1152/jn.1996.75.3.1087.

14. Wenger, K.K., Musch, K.L., and Mink, J.W. (1999). Impaired Reaching and Grasping After Focal Inactivation of Globus Pallidus Pars Interna in the Monkey. Journal of Neurophysiology. https://doi.org/10.1152/jn.1999.82.5.2049.

15. Desmurget, M., and Turner, R.S. (2008). Testing Basal Ganglia Motor Functions Through Reversible Inactivations in the Posterior Internal Globus Pallidus. Journal of Neurophysiology *99*, 1057–1076. https://doi.org/10.1152/jn.01010.2007.

16. Falasconi, A., Kanodia, H., and Arber, S. (2025). Dynamic basal ganglia output signals license and suppress forelimb movements. Nature *644*, 749–758. https://doi.org/10.1038/s41586-025-09066-z.

17. Albin, R.L., Young, A.B., and Penney, J.B. (1989). The functional anatomy of basal ganglia disorders.
Trends in Neurosciences *12*, 366–375. https://doi.org/10.1016/0166-2236(89)90074-X.
18. Alexander, G.E., and Crutcher, M.D. (1990). Functional architecture of basal ganglia circuits: neural
substrates of parallel processing. Trends in Neurosciences *13*, 266–271.
https://doi.org/10.1016/0166-2236(90)90107-L.
19. Rossi, M.A., Li, H.E., Lu, D., Kim, I.H., Bartholomew, R.A., Gaidis, E., Barter, J.W., Kim, N., Cai, M.T.,
Soderling, S.H., et al. (2016). A GABAergic nigrotectal pathway for coordination of drinking behavior.
Nat Neurosci *19*, 742–748. https://doi.org/10.1038/nn.4285.
20. Lee, J., Wang, W., and Sabatini, B.L. (2020). Anatomically segregated basal ganglia pathways allow
parallel behavioral modulation. Nat Neurosci *23*, 1388–1398. https://doi.org/10.1038/s41593-020-
00712-5.
21. McElvain, L.E., Chen, Y., Moore, J.D., Brigidi, G.S., Bloodgood, B.L., Lim, B.K., Costa, R.M., and
Kleinfeld, D. (2021). Specific populations of basal ganglia output neurons target distinct brain stem
areas while collateralizing throughout the diencephalon. Neuron *109*, 1721-1738.e4.
https://doi.org/10.1016/j.neuron.2021.03.017.
22. Foster, N.N., Barry, J., Korobkova, L., Garcia, L., Gao, L., Becerra, M., Sherafat, Y., Peng, B., Li, X.,
Choi, J.-H., et al. (2021). The mouse cortico–basal ganglia–thalamic network. Nature *598*, 188–194.
https://doi.org/10.1038/s41586-021-03993-3.
23. Arber, S., and Costa, R.M. (2022). Networking brainstem and basal ganglia circuits for movement.
Nat Rev Neurosci *23*, 342–360. https://doi.org/10.1038/s41583-022-00581-w.
24. Frost-Nylén, J., Thompson, W.S., Robertson, B., and Grillner, S. (2024). The Basal Ganglia
Downstream Control of Action - An Evolutionarily Conserved Strategy. Curr Neuropharmacol *22*,
1419–1430. https://doi.org/10.2174/1570159X21666230810141746.
25. Hikosaka, O., and Wurtz, R.H. (1983). Visual and oculomotor functions of monkey substantia nigra
pars reticulata. III. Memory-contingent visual and saccade responses. J Neurophysiol *49*, 1268–1284.
https://doi.org/10.1152/jn.1983.49.5.1268.
26. Hikosaka, O., and Wurtz, R.H. (1983). Visual and oculomotor functions of monkey substantia nigra
pars reticulata. IV. Relation of substantia nigra to superior colliculus. J Neurophysiol *49*, 1285–1301.
https://doi.org/10.1152/jn.1983.49.5.1285.
27. Mink, J.W. (1996). THE BASAL GANGLIA: FOCUSED SELECTION AND INHIBITION OF COMPETING
MOTOR PROGRAMS. Progress in Neurobiology *50*, 381–425. https://doi.org/10.1016/S0301-
0082(96)00042-1.
28. Redgrave, P., Prescott, T.J., and Gurney, K. (1999). The basal ganglia: a vertebrate solution to the
selection problem? Neuroscience *89*, 1009–1023. https://doi.org/10.1016/S0306-4522(98)00319-4.
29. Basso, M.A., and Wurtz, R.H. (2002). Neuronal Activity in Substantia Nigra Pars Reticulata during
Target Selection. J Neurosci *22*, 1883–1894. https://doi.org/10.1523/JNEUROSCI.22-05-01883.2002.
30. Fan, D., Rossi, M.A., and Yin, H.H. (2012). Mechanisms of Action Selection and Timing in Substantia
Nigra Neurons. J. Neurosci. *32*, 5534–5548. https://doi.org/10.1523/JNEUROSCI.5924-11.2012.
31. Markowitz, J.E., Gillis, W.F., Beron, C.C., Neufeld, S.Q., Robertson, K., Bhagat, N.D., Peterson, R.E.,
Peterson, E., Hyun, M., Linderman, S.W., et al. (2018). The Striatum Organizes 3D Behavior via
Moment-to-Moment Action Selection. Cell *174*, 44-58.e17.
https://doi.org/10.1016/j.cell.2018.04.019.
32. Albin, R.L., Young, A.B., and Penney, J.B. (1989). The functional anatomy of basal ganglia disorders.
Trends in Neurosciences *12*, 366–375. https://doi.org/10.1016/0166-2236(89)90074-X.
33. Kim, N., Barter, J.W., Sukharnikova, T., and Yin, H.H. (2014). Striatal firing rate reflects head
movement velocity. European Journal of Neuroscience *40*, 3481–3490.
https://doi.org/10.1111/ejn.12722.

34. Yttri, E.A., and Dudman, J.T. (2016). Opponent and bidirectional control of movement velocity in the basal ganglia. Nature *533*, 402–406. https://doi.org/10.1038/nature17639.

35. Dudman, J.T., and Krakauer, J.W. (2016). The basal ganglia: from motor commands to the control of vigor. Current Opinion in Neurobiology *37*, 158–166. https://doi.org/10.1016/j.conb.2016.02.005.

36. Barter, J.W., Castro, S., Sukharnikova, T., Rossi, M.A., and Yin, H.H. (2014). The role of the substantia nigra in posture control. European Journal of Neuroscience *39*, 1465–1473. https://doi.org/10.1111/ejn.12540.

37. Barter, J.W., Li, S., Sukharnikova, T., Rossi, M.A., Bartholomew, R.A., and Yin, H.H. (2015). Basal Ganglia Outputs Map Instantaneous Position Coordinates during Behavior. J. Neurosci. *35*, 2703–2716. https://doi.org/10.1523/JNEUROSCI.3245-14.2015.

38. Joseph, J.P., and Boussaoud, D. (1985). Role of the cat substantia nigra pars reticulata in eye and head movements. I. Neural activity. Exp Brain Res *57*, 286–296. https://doi.org/10.1007/BF00236534.

39. Handel, A., and Glimcher, P.W. (1999). Quantitative Analysis of Substantia Nigra Pars Reticulata Activity During a Visually Guided Saccade Task. Journal of Neurophysiology *82*, 3458–3475. https://doi.org/10.1152/jn.1999.82.6.3458.

40. Gulley, J.M., Kosobud, A.E.K., and Rebec, G.V. (2002). Behavior-related modulation of substantia nigra pars reticulata neurons in rats performing a conditioned reinforcement task. Neuroscience *111*, 337–349. https://doi.org/10.1016/S0306-4522(02)00018-0.

41. Lintz, M.J., and Felsen, G. (2016). Basal ganglia output reflects internally-specified movements. eLife *5*, e13833. https://doi.org/10.7554/eLife.13833.

42. Bryden, D.W., Johnson, E.E., Diao, X., and Roesch, M.R. (2011). Impact of expected value on neural activity in rat substantia nigra pars reticulata. European Journal of Neuroscience *33*, 2308–2317. https://doi.org/10.1111/j.1460-9568.2011.07705.x.

43. Cho, K., Merrienboer, B. van, Gulcehre, C., Bahdanau, D., Bougares, F., Schwenk, H., and Bengio, Y. (2014). Learning Phrase Representations using RNN Encoder-Decoder for Statistical Machine Translation. Preprint at arXiv, https://doi.org/10.48550/arXiv.1406.1078 https://doi.org/10.48550/arXiv.1406.1078.

44. Brown, J., Pan, W.-X., and Dudman, J.T. (2014). The inhibitory microcircuit of the substantia nigra provides feedback gain control of the basal ganglia output. eLife *3*, e02397. https://doi.org/10.7554/eLife.02397.

45. Mosberger, A.C., Sibener, L.J., Chen, T.X., Rodrigues, H.F.M., Hormigo, R., Ingram, J.N., Athalye, V.R., Tabachnik, T., Wolpert, D.M., Murray, J.M., et al. (2024). Exploration biases forelimb reaching strategies. Cell Reports *43*. https://doi.org/10.1016/j.celrep.2024.113958.

46. Lalive, A.L., Lien, A.D., Roseberry, T.K., Donahue, C.H., and Kreitzer, A.C. (2018). Motor thalamus supports striatum-driven reinforcement. eLife *7*, e34032. https://doi.org/10.7554/eLife.34032.

47. Isett, B.R., Nguyen, K.P., Schwenk, J.C., Yurek, J.R., Snyder, C.N., Vounatsos, M.V., Adegbesan, K.A., Ziausyte, U., and Gittis, A.H. (2023). The indirect pathway of the basal ganglia promotes transient punishment but not motor suppression. Neuron *111*, 2218-2231.e4. https://doi.org/10.1016/j.neuron.2023.04.017.

48. Cui, G., Jun, S.B., Jin, X., Pham, M.D., Vogel, S.S., Lovinger, D.M., and Costa, R.M. (2013). Concurrent activation of striatal direct and indirect pathways during action initiation. Nature *494*, 238–242. https://doi.org/10.1038/nature11846.

49. Mink, J.W., and Thach, W.T. (1991). Basal ganglia motor control. II. Late pallidal timing relative to movement onset and inconsistent pallidal coding of movement parameters. J Neurophysiol *65*, 301–329. https://doi.org/10.1152/jn.1991.65.2.301.

50. Mink, J.W., and Thach, W.T. (1993). Basal ganglia intrinsic circuits and their role in behavior. Curr Opin Neurobiol *3*, 950–957. https://doi.org/10.1016/0959-4388(93)90167-w.

51. Hou, L., Grijalva, S., Svedberg, D., and Assad, J. (2025). Congruent neuronal modulation across
competing actions challenges the role of the substantia nigra in action selection. Preprint at bioRxiv,
https://doi.org/10.1101/2025.02.18.638935 https://doi.org/10.1101/2025.02.18.638935.
52. Villalobos, C.A., and Basso, M.A. (2022). Optogenetic activation of the inhibitory nigro-collicular
circuit evokes contralateral orienting movements in mice. Cell Reports *39*, 110699.
https://doi.org/10.1016/j.celrep.2022.110699.
53. Kaneda, K., Isa, K., Yanagawa, Y., and Isa, T. (2008). Nigral Inhibition of GABAergic Neurons in Mouse
Superior Colliculus. J. Neurosci. *28*, 11071–11078. https://doi.org/10.1523/JNEUROSCI.3263-
08.2008.
54. Doykos, T.K., Yamauchi, T., Buteau, A., Hanson, S., Dudman, J.T., Felsen, G., and Stubblefield, E.A.
(2025). Correlates of head-fixed orienting movements in mouse superior colliculus and substantia
nigra pars reticulata. Preprint at bioRxiv, https://doi.org/10.1101/2025.05.02.651955
https://doi.org/10.1101/2025.05.02.651955.
55. Nath, T., Mathis, A., Chen, A.C., Patel, A., Bethge, M., and Mathis, M.W. (2019). Using DeepLabCut
for 3D markerless pose estimation across species and behaviors. Nat Protoc *14*, 2152–2176.
https://doi.org/10.1038/s41596-019-0176-0.
56. Nagai, Y., Miyakawa, N., Takuwa, H., Hori, Y., Oyama, K., Ji, B., Takahashi, M., Huang, X.-P., Slocum,
S.T., DiBerto, J.F., et al. (2020). Deschloroclozapine, a potent and selective chemogenetic actuator
enables rapid neuronal and behavioral modulations in mice and monkeys. Nat Neurosci *23*, 1157–
1167. https://doi.org/10.1038/s41593-020-0661-3.
57. Dong, Z., Mau, W., Feng, Y., Pennington, Z.T., Chen, L., Zaki, Y., Rajan, K., Shuman, T., Aharoni, D.,
and Cai, D.J. (2022). Minian, an open-source miniscope analysis pipeline. eLife *11*, e70661.
https://doi.org/10.7554/eLife.70661.

A

B

C Motif Ethogram

D Motif Transition Matrix

E 3D Trajectory

F AAV9-Syn-FLEX-jGCaMP7f
473nm

G Ipsilateral | Open   Contralateral | Open

H Ipsilateral

Contralateral

Early Reach
Late Reach

I Zscore   Ipsilateral Contralateral   Ipsilateral   Contralateral
Kinematics Mean Peak Mean Peak   Mean   Peak   Mean   Peak
            Motif Trajectory Length Vm Vp Vm Vp Vm Vp Vm Vp

J Example Correaltion

Mean Zscore

Open Trajectory Length
R 0.66
P 0.0012

Mean Zscore

Grasp Mean Velocity
R 0.57
P 0.0068

Figure 1. Increase of SNr population activity underlying stereotyped forelimb reaching action sequence.

(A) Schematic of the free-moving behavioral setup utilizing two perpendicular high-speed cameras for simultaneous recording and 3D reconstruction of forelimb reaching kinematics.

(B) Decomposition of the reaching sequence into six distinct, stereotyped action motifs (**Lift**, **Reach**, **Open**, **Grasp**, **Retract**, **Drink**). Top and bottom rows show orthogonal views of the posture for each motif. Colors correspond to specific motifs throughout the figure (color legend inset). Trial-based task structure illustrated above.

(C) Representative motif ethogram (top) and corresponding motif probability (bottom) from an example session (150 trials), aligned to the first **Open** motif.

(D) Motif transition matrix from the representative session in (C). Each color block represents the transition probability from the motif on the y-axis to the motif on the x-axis.

(E) Reconstructed 3D trajectories of the forelimb for all trials in the representative session in (C), color-coded by motif.

(F) Schematic of the fiber photometry strategy. AAV9-Syn-FLEX-jGCaMP7f was injected into the SNr to record populational calcium dynamics in SNr GABAergic neurons during behavior.

(G) Representative heatmaps of Z-scored SNr calcium signals from ipsilateral (left) and contralateral (right) hemispheres, aligned to water delivery and sorted by the onset of the first **Open** motif (pink line). Error bars represent mean ± SEM.

(H) Average Z-scored SNr activity aligned to water delivery, stratified as early reaching trials and late reaching trials by reaction time (top 30% vs. bottom 30%). n = 21 sessions (2100 trials), 7 mice.

(I) Correlation matrix summarizing the relationship between motif-specific kinematics (Trajectory Travelled Length, Mean/Peak Velocity) and motif specific SNr Z-scores. Circle size denotes statistical significance (p-value). Color intensity indicates the correlation coefficient. Spearman's rank correlation coefficient was used (Supplementary Table 1). n = 21 sessions (2100 trials), 7 mice. Refer to Table 1 for summary statistics.

(J) Representative scatter plots showing significant positive correlations between SNr activity and specific kinematic parameters: **Open** motif trajectory length (top) and **Grasp** motif mean velocity (bottom). Colored outlines in (I) indicate the specific correlations displayed here. n = 21 sessions (2100 trials), 7 mice. Pearson correlation coefficient was used.

GtACR2 Opto Inhibition

hM4D Chemo Inhibition

Figure 2. SNr inhibition suppresses forelimb skilled reaching behavior.

(A) Schematics of the experimental strategies. Optogenetic inhibition: AAV1-hSyn-DIO-GtACR2-FusRed was injected into the SNr, and photoinhibition (473nm, 4s) was delivered starting at water delivery. The motif color map (inset) applies to all panels.

(B and C) Comparison of reach success parameters: **(B)** Probability of success and **(C)** Latency to success relative to reach onset.

(D) Superimposed motif ethograms aligned to the first **Lift** motif, comparing inhibition (saturated colors) and control (lighter colors) trials.

(E and F) Comparison of reach initiation dynamics: **(E)** Motif occurrence probability and **(F)** motif latency.

(G) Quantification of motif count per trial.

(H) Quantification of total motif duration.

(I) Forelimb velocity profiles (mean ± SEM) aligned to the onset of **Lift**, **Reach**, **Open**, **Grasp**, and **Retract** motifs (left to right). Orange lines: control; Purple lines: stimulation.

(J and K) Comparison of motif Mean Velocity **(J)** and Peak Velocity **(K)**. Top panels show the Cumulative Distribution Function (CDF) plots of all trials; bottom panels show the violin plots of all sessions.

(L) Schematics of the experimental strategies. Chemogenetic inhibition: AAV2-hSyn-DIO-hM4D(Gi)-mCherry was injected into the SNr; Deschloroclozapine (DCZ) served as the agonist, with saline as the control.

(M and N) as in **(B and C)** respectively.

(O) as in **(D)**.

(P and Q) as in **(E and F)** respectively.

(R) as in **(G)**.

(S) Quantification of motif distance travelled.

(T) as in **(I)**.

(U and V) as in **(J and K)** respectively.

Left panel (GtACR2 Optogenetic inhibition), n = 2,687 control trials and 1,094 stimulation trials from 21 sessions, 7 mice. Right panel (hM4D Chemogenetic inhibition), n = 1,125 control trials and 1,125 stimulation trials from n= 15 sessions, 5 mice. Data in bar plot represent mean ± SEM. Dots in the bar and violin plots represent individual session means. Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. Wilcoxon Signed-Rank Test was used. ns no significance, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

ChR2 Opto Activation

A AAV5-EF1a-DIO-ChR2-eYFP
473nm
Drink / Retract / Grasp / Open / Reach / Lift
ITI
4s

B Success Probability — ns — Ctr Stim

C Time to Success(s) — **** — Ctr Stim

D Motif Ethogram — Ctr / Stim — Probability — Time(s)

E Motif Probability — ns ns ns ns ns ns — Probability — Ctr Stim

F Motif Latency — ns ns * ns ns ns — Time(s) — Ctr Stim

G Motif Duration — **** **** ns ns ns ns — ms — Ctr Stim

H Motif Distance Travelled — ** ns ** * ns — mm — Ctr Stim

I Motif Velocity — Ctr / Stim — mm/s — Time(s)

J Motif Mean Velocity — * ** ns ** ** — CDF — mm/s — Ctr Stim

K Motif Peak Velocity — ns *** ** ** *** — CDF — mm/s — Ctr Stim

hM3D Chemo Activation

L AAV2-hSyn-DIO-hM3D(Gq)-mCherry
Saline(Ctr)
DCZ(Stim)
Drink / Retract / Grasp / Open / Reach / Lift
ITI

M Success Probability — ns — Ctr Stim

N Time to Success(s) — ** — Ctr Stim

O Motif Ethogram — Ctr / Stim — Probability — Time(s)

P Motif Probability — ns ns ns ns ns ns — Probability — Ctr Stim

Q Motif Latency — ns ns ns ns ns ns — Time(s) — Ctr Stim

R Motif Duration — * ** ns ** ** ns — ms — Ctr Stim

S Motif Distance Travelled — ns ns ** ns ns ns — mm — Ctr Stim

T Motif Velocity — Ctr / Stim — mm/s — Time(s)

U Motif Mean Velocity — ** *** ** * ** — CDF — mm/s — Ctr Stim

V Motif Peak Velocity — ns *** ** * * — CDF — mm/s — Ctr Stim

Figure 3. SNr activation promotes forelimb skilled reaching behavior.

(A) Schematics of the experimental strategies. Optogenetic activation: AAV5-EF1a-DIO-ChR2-eYFP was injected into the SNr, and photoactivation (473nm, 4s) was delivered starting at water delivery. The motif color map (inset) applies to all panels.

(B and C) Comparison of reach success parameters: **(B)** Probability of success and **(C)** Latency to success relative to reach onset.

(D) Superimposed motif ethograms aligned to the first **Lift** motif, comparing activation (saturated colors) and control (lighter colors) trials.

(E and F) Comparison of reach initiation dynamics: **(E)** Motif occurrence probability and **(F)** motif latency.

(G) Quantification of motif duration.

(H) Quantification of total motif duration.

(I) Forelimb velocity profiles (mean ± SEM) aligned to the onset of **Lift**, **Reach**, **Open**, **Grasp**, and **Retract** motifs (left to right). Orange lines: control; Purple lines: stimulation.

(J and K) Comparison of motif Mean Velocity **(J)** and Peak Velocity **(K)**. Top panels show the Cumulative Distribution Function (CDF) plots of all trials; bottom panels show the violin plots of all sessions.

(L) Schematics of the experimental strategies. Chemogenetic activation: AAV2-hSyn-DIO-hM3D(Gq)-mCherry was injected into the SNr; Deschloroclozapine (DCZ) served as the agonist, with saline as the control.

(M and N) as in **(B and C)** respectively.

(O) as in **(D)**.

(P and Q) as in **(E and F)** respectively.

(R) as in **(G)**.

(S) as in **(H)**.

(T) as in **(I)**.

(U and V) as in **(J and K)** respectively.

Left panel (ChR2 Optogenetic inhibition), n = 1,824 control trials and 691 stimulation trials from 21 sessions, 4 mice. Right panel (hM3D Chemogenetic inhibition), n = 900 control trials and 900 stimulation trials from n= 12 sessions, 4 mice. Data in bar plot represent mean ± SEM. Dots in the bar and violin plots represent individual session means. Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. Wilcoxon Signed-Rank Test was used. ns no significance, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

GtACR2 100 ms Brief Pulse Inhibition During Reaching

A. AAV1-hSyn-SIO-GtACR-FusRed

B. Motif Transition Matrix Difference

C. Motif Duration

D. Motif Distance Travelled

E. 3D Trajectory

F. Coordinate

G. Distance to Target

H. Velocity

GtACR2 12.5 ms Momentary Pulse Inhibition During Reaching

I. AAV1-hSyn-SIO-GtACR-FusRed

J. Motif Transition Matrix Difference

K. Motif Duration

L. Motif Distance Travelled

M. 3D Trajectory

N. Coordinate

O. Distance to Target

P. Velocity

Figure 4. Coincident, momentary SNr inhibitory perturbation modulates online reaching kinematics.

(**A**) Schematic of the experimental strategy. An inhibitory opsin (AAV1-hSyn-DIO-GtACR2-FusRed) was injected into the SNr, and an optical fiber was implanted above the region. A single 100ms photoinhibition pulse was delivered at a randomized time after water delivery onset to target reaching action sequence. The motif color map (inset) applies to all panels.

(B) Motif transition matrix difference maps (stimulation - control). Each color block represents the difference between two conditions in transition probability from the motif on the y-axis to the motif on the x-axis. Motif sequence from a 300ms window (-40ms to 260ms aligned to laser onset) was used for quantification. The color map is shown on the right.

(C and D) Comparison of motif duration **(C)** and motif distance travelled **(D)**. Top panels show the Cumulative Distribution Function plots of all trials; bottom panels show the violin plots of all sessions.

(E) Average 3D forelimb trajectories (from -200ms to 800ms aligned to laser onset). The blue dot indicates laser pulse onset. Orange lines: control trials; Purple lines: stimulation trials.

(F) Forelimb coordinates (X, Y, Z) aligned to laser pulse onset. Left: temporal traces (mean ± SEM). The blue-shaded region represents laser duration. Right: paired comparison of mean and maximum/minimum coordinates over a 200ms window starting at laser onset (Seg).

(G and H) Forelimb distance to target **(G)** and velocity **(H)** aligned to laser pulse onset. Left: temporal traces (mean ± SEM). The blue-shaded region represents laser duration. Right: paired comparison of mean and maximum/minimum values over the designated analysis window (Seg).

(I) as in **(A)**. But a single 12.5ms photoinhibition pulse was used instead.

(J) as in **(B)**.

(K and L) as in **(C and D)** respectively.

(M) as in **(E)**.

(N) as in **(F)**.

(O and P) as in **(G and H)** respectively.

Left panel (300 ms inhibition), n = 902 control trials and 1,319 stimulation trials from 33 sessions, 7 mice. Right panel (12.5 ms inhibition), n = 761 control trials and 911 stimulation trials from n= 26 sessions, 7 mice. Dots in the violin plots represent individual session means. Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. Wilcoxon Signed-Rank Test was used. ns no significance, *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

ChR2 300 ms Brief Pulse Activation During Reaching

A AAV5-EF1a-DIO-ChR2-eYFP

B Motif Transition Matrix Difference

C 3D Trajectory

D Coordinate

E Distance to Target

F Velocity

G Motif Mean Velocity

H Motif Peak Velocity

ChR2 12.5 ms Momentary Pulse Activation During Reaching

I AAV5-EF1a-DIO-ChR2-eYFP

J Motif Transition Matrix Difference

K 3D Trajectory

L Coordinate

M Distance to Target

N Velocity

O Motif Mean Velocity

P Motif Peak Velocity

Figure 5. Coincident, momentary SNr excitatory perturbation modulates online reaching kinematics.

(**A**) Schematic of the experimental strategy. An excitatory opsin (AAV5-EF1a-DIO-ChR2-eYFP) was injected into the SNr, and an optical fiber was implanted above the region. A single 300ms photoactivation pulse was delivered at a randomized time after water delivery onset to target reaching action sequence. The motif color map (inset) applies to all panels.

(**B**) Motif transition matrix difference maps (stimulation - control). Each color block represents the difference between two conditions in transition probability from the motif on the y-axis to the motif on the x-axis. Motif sequence from a 300ms window (-40ms to 260ms aligned to laser onset) was used for quantification. The color map is shown on the right.

(**C**) Average 3D forelimb trajectories (from -200ms to 800ms aligned to laser onset). The blue dot indicates laser pulse onset. Orange lines: control trials; Purple lines: stimulation trials.

(**D**) Forelimb coordinates (X, Y, Z) aligned to laser pulse onset. Left: temporal traces (mean ± SEM). The blue-shaded region represents laser duration. Right: paired comparison of mean coordinates averaged over designated analysis windows (Seg 1: 300ms to 350ms; Seg 2:400ms to 800ms).

(**E and F**) Forelimb distance to target (**E**) and velocity (**F**) aligned to laser pulse onset. Left: temporal traces (mean ± SEM). The blue-shaded region represents laser duration. Right: paired comparison of mean values over the designated window (distance to target: Seg 1 (300ms to 350ms), Seg 2 (400ms to 800ms); velocity: Seg 1(100ms to 300ms), Seg 2(300ms to 400ms).

(**G and H**) Comparison of motif Mean Velocity (**G**) and Peak Velocity (**H**). Top panels: Cumulative Distribution Function plots of all trials; bottom panels: violin plots of all sessions.

(**I**) as in (**A**). But a single 12.5ms photoactivation pulse was used instead.

(**J**) as in (**B**).

(**K**) as in (**C**).

(**L**) as in (**D**).

(**M and N**) as in (**E and F**) respectively.

(**O and P**) as in (**G and H**) respectively.

Left panel (300 ms excitation), n = 493 control trials and 483 stimulation trials from 13 sessions, 4 mice. Right panel (12.5 ms excitation), n = 1,213 control trials and 1,193 stimulation trials from n= 30 sessions, 4 mice. Dots in the violin plots represent individual session means. Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. Wilcoxon Signed-Rank Test was used. ns no significance, *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

A AAV9-Syn-FLEX-jGCaMP7f

B Population Average
   Reach Aligned

C Reach

D Behavior Modulated Neurons
   Increased Decreased Unchanged

Lift   Reach   Open
30.8% 34.9% 35.2% 32.1% 37.1% 30.2%
       34.3%   32.7%   32.7%

Grasp   Retract   Drink
37.8% 29.8% 38.1% 30.5% 40.6% 33.7%
       32.4%   31.4%   25.7%

E

F Kinematics Prediction using Population Activity
   Real Predicted

   X   Y
   Z   Velocity

G Regression R² Score

H Weights In Regression Model

Figure 6. The SNr population acts as a continuous controller for moment-by-moment 3D kinematics during forelimb reaching.

(A) Schematic of the miniscope recording strategy. AAV9-Syn-FLEX-jGCaMP7f was injected into the SNr to monitor calcium dynamics in GABAergic projection neurons at single-cell resolution during behavior.

(B) Average population activity aligned to the onset of the Reach motif. The solid line represents the mean, and the shaded area represents SEM.

(C) Heatmap of trial-averaged, Reach-aligned calcium activity for all recorded neurons.

(D) Proportion of neurons exhibiting significantly increased (orange), decreased (purple), or unchanged (gray) activity during specific action motifs compared to baseline periods (Wilcoxon Signed-Rank Test).

(E) Heatmap of the change in activity (ΔZscore; Behavior - Baseline) for all neurons across the six action motifs, sorted in descending order by modulation magnitude during the Retract motif.

(F) Representative temporal traces comparing actual (orange) and predicted (light purple) forelimb spatial coordinates (X, Y, Z) and velocity. Predictions were generated using a linear regression model trained on concurrent SNr population activity.

(G) Quantification of the linear regression model's performance (R^2 score) in predicting moment-by-moment kinematics across sessions. Violin plots display the full data distribution density, with internal box plots representing the median (center marker) and the interquartile range.

(H) Heatmap of normalized regression weights for each neuron across the four predicted kinematic features. Neurons (rows) are sorted in descending order by their assigned weight in the velocity model.

n = 315 neurons from 600 trials, 10 sessions, 5 mice.

Figure S1. Establishment of a GRU (Gated recurrent unit)-based classifier for automated

reaching sequence segmentation and motif-specific analysis, related to Figure 1.

(A) Schematic of the behavioral analysis pipeline. DeepLabCut (DLC) was used to track six key

body parts (digits 2–4, paw, mouth, and nose). Extracted coordinates were used to calculate

pairwise inter-bodyparts coordinate and then were augmented with derived kinematic features,

including velocity, acceleration. The augmented features were used as input to a GRU classifier

for automated motif segmentation and subsequent analysis.

(B–M) Quantification of temporal and kinematic parameters across action motifs from a large

dataset (n=24498 control trials, 251 sessions from 28 mice). Violin plots display the full data

distribution density; internal box plots represent the median (center line) and the interquartile

range. Parameters are grouped into: **(B–E)** temporal statistics (motif count, duration, latency,

and total duration); **(F–I)** coordinate related metrics (distance to target, inter-digit distance, and

trajectory travelled length); and **(J–M)** velocity related metrics (mean and peak velocities for the

forelimb and inter-digit spread). All quantified parameters exhibited significant variation across

motifs (Friedman test, $****p < 0.0001$)

A
AAV1-hSyn-SIO-GtACR2-FusRed
473nm

Drink
Retract
Grasp
Open
Reach
Lift

ITI
10 mw, 5 mw, 2.5 mw, 1.25 mw
4s

B
Within Laser
Reach Probability
****
Probability
Ctr 1.25 2.5 5 10 mw

C
After Laser
Reach Probability
****
Probability
Ctr 1.25 2.5 5 10 mw

D
Latency to Reach
****
Time(s)
Ctr 1.25 2.5 5 10 mw

E
Motif Ethogram
Ctr
1.25 mw
2.5 mw
5 mw
10 mw
Probability
Time (s)

F
Motif Transition Matrix Difference
1.25 mW   2.5 mW   5 mW   10 mW
Preceding Motif
None, Lift, Reach, Open, Grasp, Retract, Drink
Current Motif
Probability Difference(Stim-Ctr)

G
Motif Velocity
mm/s
Ctr
1.25 mw
2.5 mw
Time(s)

H
Motif Mean Velocity
CDF
Ctr
20 Hz
40 Hz
****   ****   ****   ***   ns
mm/s
Ctr 1.25 mw 2.5 mw   Ctr 1.25 mw 2.5 mw   Ctr 1.25 mw 2.5 mw   Ctr 1.25 mw 2.5 mw   Ctr 1.25 mw 2.5 mw

I
Motif Peak Velocity
CDF
Ctr
20 Hz
40 Hz
*   ***   ***   ****   *
mm/s
Ctr 1.25 mw 2.5 mw   Ctr 1.25 mw 2.5 mw   Ctr 1.25 mw 2.5 mw   Ctr 1.25 mw 2.5 mw   Ctr 1.25 mw 2.5 mw

Figure S2. Intensity-dependent modulation of forelimb reaching behavior by SNr

photoinhibition, related to Figure 2.

(A) Schematic of the experimental strategy. An inhibitory opsin (AAV1-hSyn-DIO-GtACR2-

FusRed) was injected into the SNr and an optical fiber was implanted above the region.

Photoinhibition was delivered using a 473 nm blue laser at four different intensities (1.25 mW,

2.5 mW, 5 mW, 10 mW), applied randomly across trials starting at water delivery and lasting for

4 seconds. The motif color map is provided (inset).

(B–D) Quantification of reach initiation dynamics: **(B)** Probability of reaching during laser

delivery (Within Laser Reach Probability); **(C)** Probability of reaching following laser cessation

(After Laser Reach Probability); and **(D)** Latency to Reach. (n = 21 sessions, 7 mice; Friedman

test).

(E) Motif ethogram aligned to the first Lift motif, stratified by inhibition intensity (n = 21 sessions,

7 mice).

(F) Motif transition matrix difference maps, stratified by laser intensity. Each matrix displays the

difference in transition probability (Inhibition - Control). n = 21 sessions, 7 mice.

(G) Forelimb velocity profiles (mean ± SEM) aligned to the onset of Lift, Reach, Open, Grasp,

and Retract motifs (left to right). Only low-intensity trials (1.25 mW and 2.5 mW) are shown, as

higher intensities blocked the majority of reaching attempts (n = 19 sessions, 7 mice).

(H and I) Quantification of motif Mean Velocity and Peak Velocity across different laser

intensities. Top panels show the Cumulative Distribution Function (CDF) plots; bottom panels

show the violin plots. (n = 19 sessions, 7 mice; Friedman test).

1017

1018 Data in bar plot are shown as mean ± SEM. Dots in the bar and violin plots represent individual

1019 session means. Violin plots display the full data distribution density; internal box plots represent

1020 the median (center line) and the interquartile range. ns no significance, $*p < 0.05$, $**p < 0.01$,

1021 $***p < 0.001$, $****p < 0.0001$.

Figure S3. Rebound reaching following SNr inhibition exhibits enhanced kinematic signatures, related to Figure 2.

(A) Schematic of variable-duration optogenetics inhibition. A 473 nm laser was delivered at water delivery onset for randomly interleaved durations (1 s, 2 s, 3 s, 4 s). The motif color map (inset) applies to all panels.

(B) Motif ethograms aligned to laser onset (left) and the laser offset (right), stratified by inhibition duration in descending order. Shaded regions indicate laser-on periods (n = 808 trials, 14 sessions,5 mice).

(C) Quantification of reach probability during (Within Laser Reach Probability) and immediately following (After Laser Reach Probability) inhibition (n = 14 sessions, 5 mice).

(D) Cumulative Distribution Function (CDF) plots of reach latency of all trials, stratified by inhibition duration (dark to light purple) compared to controls (orange) (n = 1,652 control and 808 stimulation trials; 14 sessions, 5 mice).

(E) Motif transition matrices. Left: reaching in control trials; right: rebound reaching in stimulation trials; middle: difference between rebound and control transition matrices. (n = 1,652 control trials and 808 stimulation trials from 14 sessions, 5 mice).

(F and G) Comparison of reach success probability **(F)** and latency to success **(G)** between control and rebound reaches (n = 14 sessions, 4 mice; Wilcoxon Signed-Rank Test).

(H) Superimposed motif ethograms aligned to the first Lift motif, comparing rebound (saturated colors) and control (lighter colors) reaches (n = 1,652 control and 808 stimulation trials; 14 sessions, 5 mice).

(I) Session average forelimb velocity profiles (mean ± SEM) aligned to the onset of Lift, Reach, Open, Grasp, and Retract motifs (left to right). Orange lines: control reaches; Purple lines: rebound reaches. (n = 14 sessions, 4 mice).

(J and K) Quantification of motif Mean Velocity **(J)** and Peak Velocity **(K)** for control vs rebound reaches. Top panels: Cumulative Distribution Function (CDF) (n = 1,652 control and 808 stimulation trials; 14 sessions, 5 mice); bottom panels: violin plots (n = 12 sessions, 4 mice; Wilcoxon Signed-Rank Test). Dots in the bar and violin plots represent individual session means.

Data in bar plot are shown as mean ± SEM. Dots in the bar and violin plots represent individual session means. Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. ns no significance, $p < 0.05$, $p < 0.01$, $p < 0.001$, $p < 0.0001$.

Figure S4. Frequency-dependent modulation of forelimb reaching behavior by SNr photoactivation, related to Figure 3.

(A) Schematic of the optogenetic activation strategy. An excitatory opsin (AAV5-EF1a-DIO-ChR2-eYFP) was injected into the SNr and an optical fiber was implanted above the region. Photoactivation was delivered using a 473 nm blue laser at two different frequencies (20Hz, 40Hz), applied randomly across trials starting at water delivery and lasting for 4 seconds. The motif color map (inset) is used throughout the figure.

(B–D) Quantification of reach initiation dynamics: **(B)** Probability of reaching during laser delivery (Within Laser Reach Probability); **(C)** Probability of reaching following laser cessation (After Laser Reach Probability); and **(D)** Latency to Reach. (n = 12 sessions, 4 mice; Wilcoxon Signed-Rank Test).

(E and F) Quantification of reach success parameters: **(E)** Probability of success during laser delivery (Within Laser Success Probability); **(F)** Latency to success from reaching during laser delivery (Latency to Success). (n = 12 sessions, 4 mice; Wilcoxon Signed-Rank Test).

(G) Motif transition matrix difference maps, stratified by laser frequency. Each matrix displays the difference in transition probability (Activation - Control). n = 12 sessions, 4 mice.

(H) Motif ethogram aligned to the first Lift motif, stratified by inhibition intensity (n = 12 sessions, 4 mice).

(I) Session average forelimb velocity profiles (mean ± SEM) aligned to the onset of Lift, Reach, Open, Grasp, and Retract motifs (left to right). Orange lines represent control trials, and purple lines represent activation trials.

(J and K) Quantification of motif Mean Velocity and Peak Velocity across different laser frequencies. Top panels show the Cumulative Distribution Function (CDF) plots; bottom panels show the violin plots. (n = 12 sessions, 4 mice; Wilcoxon Signed-Rank Test).

Data in bar plot are shown as mean ± SEM. Dots in the bar and violin plots represent individual session means. Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. ns no significance, *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

A AAV1-hSyn-SIO-GtACR-FusRed

B Reach Probability — ns — Post Ctr, Post Stim

C Reach Latency(s) — ns — Post Ctr, Post Stim

D Success Probability — ns — Post Ctr, Post Stim

E Success Latency(s) — ns — Post Ctr, Post Stim

H 3D Trajectory — Post Ctr, Post Stim

F Motif Transition Matrix — Post Ctr, Post Stim - Post Ctr, Post Stim

G Motif Ethogram — Post Ctr, Post Stim

I X, Y, Z — Post Ctr, Post Stim

J Motif Onset Aligned Velocity — Post Ctr, Post Stim

Figure S5. **SNr photoinhibition does not produce carryover effects on subsequent control trials, related to Figure 2.**

(A) Schematic of the experimental strategy to assess carryover effects. Top: Viral expression and fiber implantation in the SNr. Middle: Diagram of the trial-based reaching task and optogenetics protocol. Bottom: Definition of trial categories; Post Ctr denotes control trials immediately following a control trial, and Post Stim denotes control trials immediately following a stimulation trial. The motif color map (inset) applies to all panels.

(B–E) Quantification of reach dynamics comparing Post Ctr and Post Stim conditions: **(B)** Probability of reaching; **(C)** Latency to Reach; **(D)** Probability of success; and **(E)** Latency to Success. Data in bar plot are shown as mean ± SEM. Dots in the bar and violin plots represent individual session means. Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. No significant differences were observed (n = 21 sessions, 7 mice; Wilcoxon Signed-Rank Test).

(F) Motif transition matrices for Post Ctr Trials (left) and Post Stim Trials (right). The difference matrix (middle) indicates no substantial deviation in motif sequence progression (n = 1,917 Post Ctr and 770 Post Stim Trials; 21 sessions, 7 mice).

(G) Superimposed motif ethograms aligned to the first Lift motif, comparing Post Ctr (lighter colors) and Post Stim (saturated colors) Trials (n = 1,917 Post Ctr Trials and 770 Post Stim Trials from 21 sessions, 7 mice.

(H and I) Forelimb reaching kinematics aligned to the first Lift motif: **(H)** Average 3D forelimb trajectories and **(I)** corresponding X, Y, Z coordinate traces. Orange lines: Post Ctr; Pink lines: Post Stim (n = 21 sessions, 7 mice).

(J) Session average forelimb velocity profiles (mean ± SEM) aligned to the onset of Lift, Reach, Open, Grasp, and Retract motifs (left to right). Orange lines represent control trials, and purple lines represent activation trials. (n = 21 sessions, 7 mice).

A AAV5-EF1a-DIO-ChR2-eYFP

Drink
Retract
Grasp
Open
Reach
Lift

B Reach Probability — ns — Post Ctr, Post Stim

C Reach Latency(s) — *** — Post Ctr, Post Stim

D Success Probability — ns — Post Ctr, Post Stim

E Success Latency(s) — ns — Post Ctr, Post Stim

H 3D Trajectory — Post Ctr, Post Stim

F Motif Transition Matrix Post Stim - Post Ctr — Post Ctr, Post Stim — Preceding Motif / Next Motif — Probability

G Motit Ethogram — Post Ctr, Post Stim

I Coordinate (mm) — X, Y, Z — Post Ctr, Post Stim

J Motif Velocity — Post Ctr, Post Stim

Figure S6. SNr photoactivation does not produce carryover effects on subsequent control trials, related to Figure 3.

(A) Schematic of the experimental strategy to assess carryover effects. Top: Viral expression and fiber implantation in the SNr. Middle: Diagram of the trial-based reaching task and optogenetics protocol. Bottom: Definition of trial categories; Post Ctr denotes control trials immediately following a control trial, and Post Stim denotes control trials immediately following a stimulation trial. The motif color map (inset) applies to all panels.

(B–E) Quantification of reach dynamics comparing Post Ctr and Post Stim conditions: **(B)** Probability of reaching; **(C)** Latency to Reach; **(D)** Probability of success; and **(E)** Latency to Success. Data in bar plot are shown as mean ± SEM. Dots in the bar and violin plots represent individual session means. Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. (n = 12 sessions, 4 mice; Wilcoxon Signed-Rank Test). ns no significance, $p < 0.05$, $p < 0.01$, $p < 0.001$, $p < 0.0001$.

(F) Motif transition matrices for Post Ctr Trials (left) and Post Stim Trials (right). The difference matrix (middle) indicates no substantial deviation in motif sequence progression (n = 1,320 Post Ctr and 504 Post Stim Trials; 14 sessions, 4 mice).

(G) Superimposed motif ethograms aligned to the first Lift motif, comparing Post Ctr (lighter colors) and Post Stim (saturated colors) Trials (n = 1, 320 Post Ctr Trials and 504 Post Stim Trials from 14 sessions, 4 mice.

(H and I) Forelimb reaching kinematics aligned to the first Lift motif: **(H)** Average 3D forelimb trajectories and **(I)** corresponding X, Y, Z coordinate traces. Orange: Post Ctr; Purple: Post Stim (n = 14 sessions, 4 mice).

(J) Session average forelimb velocity profiles (mean ± SEM) aligned to the onset of Lift, Reach, Open, Grasp, and Retract motifs (left to right). Orange lines represent control trials, and purple lines represent activation trials. (n = 14 sessions, 4 mice).

Figure S7. Brief SNr pulse photoinhibition produces real-time modulation of reaching kinematics, related to Figure 4.

(A) Schematic of the experimental strategy. A single 12.5 ms photoinhibition pulse was delivered at a randomized time after water delivery onset. Trials were grouped and analyzed post-hoc based on the action motif coincident with pulse onset (Lift, Reach, Open, Grasp, or Retract). The motif color map (inset) applies to all panels.

(B) Motif ethograms (top) and corresponding probability (bottom) aligned to laser pulse onset, stratified by the coincident motif (left to right: Lift, Reach, Open, Grasp, or Retract). The blue-shaded region indicates the laser duration. (Total n = 898 trials from 26 sessions, 7 mice; Lift: 240, Reach: 280, Open: 134, Grasp: 70, Retract: 186 trials).

(C) Average 3D forelimb trajectories aligned to laser pulse onset (blue dots), stratified by the coincident motif. Orange lines: control trials; Purple lines: stimulation trials. (n = 26 sessions, 7 mice).

(D) Forelimb coordinates (X, Y, Z) aligned to laser pulse onset. Left: temporal traces (mean ± SEM). Right: paired comparison of mean coordinates averaged over a 100 ms window starting at laser onset (Seg). Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. Orange: control trials; Purple: stimulation trials. (n = 26 sessions, 7 mice; Wilcoxon Signed-Rank Test). ns no significance, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

(E) Forelimb distance to target aligned to laser pulse onset. Left: temporal traces (mean ± SEM). Right: paired comparison of mean distance averaged over a 100 ms window starting at laser onset (Seg). Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. Orange: control trials; Purple: stimulation trials. (n = 26 sessions, 7 mice; Wilcoxon Signed-Rank Test). ns no significance, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Figure S8. Brief SNr pulse photoactivation produces real-time modulation of reaching kinematics, related to Figure 5.

(A) Schematic of the experimental strategy. A single 12.5 ms photoinhibition pulse was delivered at a randomized time after water delivery onset. Trials were grouped and analyzed post-hoc based on the action motif coincident with pulse onset (Lift, Reach, Open, Grasp, or Retract). The motif color map (inset) applies to all panels.

(B) Motif ethograms (top) and corresponding probability (bottom) aligned to laser pulse onset, stratified by the coincident motif (left to right: Lift, Reach, Open, Grasp, or Retract). The blue-shaded region indicates the laser duration. (Total n = 1193 trials from 30 sessions, 4 mice; Lift: 115, Reach: 311, Open: 191, Grasp: 170, Retract: 406 trials).

(C) Average 3D forelimb trajectories aligned to laser pulse onset (blue dots), stratified by the coincident motif. Orange lines: control trials; Purple lines: stimulation trials. (n = 30 sessions, 4 mice).

(D) Forelimb coordinates (X, Y, Z) aligned to laser pulse onset. Left: temporal traces (mean ± SEM). Right: paired comparison of mean coordinates averaged over a defined window (Seg). Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. Orange: control trials; Purple: stimulation trials. (n = 30 sessions, 4 mice; Wilcoxon Signed-Rank Test). ns no significance, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

(E) Forelimb velocity aligned to laser pulse onset. Left: temporal traces (mean ± SEM). Right: paired comparison of mean velocity averaged over a defined window (Seg). Violin plots display the full data distribution density; internal box plots represent the median (center line) and the interquartile range. Orange: control trials; Purple: stimulation trials. (n = 30 sessions, 4 mice; Wilcoxon Signed-Rank Test). ns no significance, $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

Supplementary Video 1. Reaching task structure and skilled forelimb motifs. Example trial

demonstrating the behavioral setup, task progression, and motor motifs in the skilled forelimb

reaching task. Related to Figure 1.

Supplementary Video 2. 3D trajectory reconstructions of forelimb reaching motifs.

Reconstructed 3D trajectories of all trials from an example session, color-coded by action motif.

The video shows 500 total frames per trial, beginning 100 frames prior to the first Grasp motif

following water delivery. Trajectory tails are set to a 50-frame fade. Related to Figure 1.

Supplementary Video 3. Behavioral effects of SNr manipulation. Representative trials

showing the effects of 4-s optogenetic inhibition, chemoinhibition, 4-s optogenetic activation,

and chemoactivation on forelimb skilled behavior. Related to Figures 2 and 3.

Supplementary Video 4. Session-averaged 3D trajectories under SNr manipulation.

Reconstructed 3D trajectories (session averages) comparing experimental (purple) and control

(orange) conditions. Data includes 4-s optogenetic inhibition, chemoinhibition, 4-s optogenetic

activation, and chemoactivation. Each curve represents a single session. Video shows 250

frames per session, starting 50 frames prior to laser onset. Related to Figures 2 and 3.

Supplementary Video 5. Behavioral effects of short optogenetic manipulation. Example

trials demonstrating the effects of short optogenetics manipulation coinciding with specific motor

motifs: 100-ms and 12.5-ms inhibition coinciding with Lift; 100-ms and 12.5-ms inhibition

coinciding with Reach; and 300-ms and 12.5-ms activation coinciding with Reach. Related to

Figures 4 and 5.

Supplementary Video 6. Reconstructed 3D trajectories under short optogenetic

manipulation. 3D trajectories of all trials from representative mice during short optogenetic

manipulations. Includes 100-ms and 12.5-ms inhibition (Lift and Reach) and 300-ms and 12.5-

ms activation (Lift-to-Retract). Purple: Inhibition; Orange: Control. Each curve represents one

trial color-coded by action motif. Video shows 250 frames per trial, starting 50 frames prior to

laser onset. Related to Figures 4 and 5.

1213

	Ipsilateral		Contralateral		Ipsilateral				Contralateral			
Zscore	Mean	Peak	Mean	Peak	Mean	Peak	Mean	Peak	Mean	Peak	Mean	Peak
	Motif Trajectory Length				Vm	Vp	Vm	Vp	Vm	Vp	Vm	Vp
Lift	R: 0.140 p: 0.544	R: 0.110 p: 0.634	R: 0.061 p: 0.793	R: 0.090 p: 0.695	R: -0.349 p: 0.121	R: -0.238 p: 0.300	R: -0.347 p: 0.124	R: -0.258 p: 0.258	R: -0.266 p: 0.243	R: -0.230 p: 0.316	R: -0.260 p: 0.256	R: -0.206 p: 0.369
Reach	R: 0.105 p: 0.650	R: 0.049 p: 0.832	R: 0.404 p: 0.069	R: 0.518 p: 0.016	R: 0.208 p: 0.366	R: 0.119 p: 0.606	R: 0.114 p: 0.622	R: 0.057 p: 0.806	R: 0.300 p: 0.186	R: 0.401 p: 0.071	R: 0.390 p: 0.081	R: 0.505 p: 0.019
Open	R: 0.526 p: 0.014	R: 0.526 p: 0.014	R: 0.560 p: 0.008	R: 0.588 p: 0.005	R: 0.496 p: 0.022	R: 0.553 p: 0.009	R: 0.479 p: 0.028	R: 0.547 p: 0.010	R: 0.409 p: 0.066	R: 0.617 p: 0.003	R: 0.436 p: 0.048	R: 0.638 p: 0.002
Grasp	R: 0.009 p: 0.969	R: -0.032 p: 0.889	R: 0.591 p: 0.005	R: 0.604 p: 0.004	R: 0.406 p: 0.067	R: 0.390 p: 0.081	R: 0.365 p: 0.104	R: 0.348 p: 0.122	R: 0.573 p: 0.007	R: 0.562 p: 0.008	R: 0.577 p: 0.006	R: 0.562 p: 0.008
Retract	R: 0.094 p: 0.687	R: 0.103 p: 0.658	R: 0.625 p: 0.002	R: 0.630 p: 0.002	R: 0.503 p: 0.020	R: 0.291 p: 0.201	R: 0.509 p: 0.018	R: 0.294 p: 0.197	R: 0.622 p: 0.003	R: 0.504 p: 0.020	R: 0.622 p: 0.003	R: 0.512 p: 0.018

1216

Fig	Statistic Method	Sample Size	P value
1I	Spearman's rank correlation coefficient	21 sessions from 7 mice	**See Table 1**
1J	Pearson correlation coefficient	21 sessions from 7 mice	**Top:** R = 0.571, p = 0.007 **Bottom:** R = 0.658, p = 0.001
2B	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	< 0.0001
2C	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	0.272
2E	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	**From left to right:** < 0.0001 | < 0.0001 | < 0.0001 < 0.0001 | < 0.0001 | < 0.0001
2F	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	**From left to right:** < 0.0001 | < 0.0001 | < 0.0001 < 0.0001 | < 0.0001 | 0.0002
2G	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	**From left to right:** 0.014 | < 0.0001 | < 0.0001 < 0.0001 | < 0.0001 | < 0.0001
2H	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	**From left to right:** 0.046 | < 0.0001 | 0.0002 < 0.0001 | < 0.0001 | < 0.0001
2J	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	**From left to right:** 0.0002 | 0.0003 | < 0.0001 | < 0.0001 | 0.070
2K	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	**From left to right:** 0.002 | 0.0002 | < 0.0001 | < 0.0001 | 0.035

2M	Wilcoxon Signed-Rank Test	15 sessions from 5 mice	0.389
2N	Wilcoxon Signed-Rank Test	15 sessions from 5 mice	0.0001
2P	Wilcoxon Signed-Rank Test	15 sessions from 5 mice	**From left to right:** 0.598 | 0.086 | 0.124 | 0.125 | 0.098 | 0.035
2Q	Wilcoxon Signed-Rank Test	15 sessions from 5 mice	**From left to right:** 0.720 | 0.359 | 0.229 | 0.303 | 0.252 | 0.022
2R	Wilcoxon Signed-Rank Test	15 sessions from 5 mice	**From left to right:** 0.0004 | 0.002 | 0.004 | 0.005 | 0.002 | 0.0001
2S	Wilcoxon Signed-Rank Test	15 sessions from 5 mice	**From left to right:** 0.0001 | 0.010 | 0.048 | 0.005 | 0.012 | 0.0004
2U	Wilcoxon Signed-Rank Test	15 sessions from 5 mice	**From left to right:** 0.002 | < 0.0001 | 0.389 | 0.083 | 0.026
2V	Wilcoxon Signed-Rank Test	15 sessions from 5 mice	**From left to right:** < 0.0001 | < 0.0001 | 0.003 | 0.018 | 0.008
3B	Wilcoxon Signed-Rank Test	21 sessions from 4 mice	0.733
3C	Wilcoxon Signed-Rank Test	21 sessions from 4 mice	< 0.0001
3E	Wilcoxon Signed-Rank Test	21 sessions from 4 mice	**From left to right:** 0.683 | 0.683 | 0.683 | 0.733 | 0.865 | 0.562
3F	Wilcoxon Signed-Rank Test	21 sessions from 4 mice	**From left to right:** 0.393 | 0.203 | 0.035 | 0.179 | 0.320 | 0.065
3G	Wilcoxon Signed-Rank Test	21 sessions from 4 mice	**From left to right:** < 0.0001 | < 0.0001 | 0.229 0.946 | 0.060 | 0.168
3H	Wilcoxon Signed-Rank Test	21 sessions from 4 mice	**From left to right:** 0.006 | 0.243 | 0.002 | 0.016 | 0.355 | 0.096
3J	Wilcoxon Signed-Rank Test	21 sessions from 4 mice	**From left to right:** 0.018 | 0.002 | 0.119 | 0.007 | 0.004
3K	Wilcoxon Signed-Rank Test	21 sessions from 4 mice	**From left to right:** 0.272 | 0.0006 | 0.006 | 0.002 | 0.0004
3M	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	0.110
3N	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	0.007
3P	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**From left to right:** 0.212 | 0.306 | 0.350 | 0.266 | 0.306 | 0.358
3Q	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**From left to right:** 0.970 | 0.622 | 0.301 | 0.266 | 0.380 | 0.092
3R	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**From left to right:** 0.016 | 0.007 | 0.569 | 0.622 | 0.005 | 0.970
3S	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**From left to right:** 0.470 | 0.850 | 0.003 | 0.424 | 0.151 | 0.470
3U	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**From left to right:** 0.005 | 0.001 | 0.005 | 0.042 | 0.002
3V	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**From left to right:** 0.380 | 0.001 | 0.0015 | 0.027 | 0.027
4C	Wilcoxon Signed-Rank Test	33 sessions from 7 mice	**Left to Right (Lift to Retract):** 0.0002 | 0.0001 | 0.016 | 0.012 | 0.009
4D	Wilcoxon Signed-Rank Test	33 sessions from 7 mice	**Left to Right (Lift to Retract):** 0.012 | 0.0001 | 0.005 | < 0.0001 | < 0.0001
4F	Wilcoxon Signed-Rank Test	33 sessions from 7 mice	**X Seg Mean:** 0.0002 **X Seg |Peak|:** < 0.0001

			Y Seg Mean: 0.0004 **Y Seg |Peak|:** < 0.0001 **Z Seg Mean:** < 0.0001 **Z Seg |Peak|:** < 0.0001
4G	Wilcoxon Signed-Rank Test	33 sessions from 7 mice	**Mean:** < 0.0001 **Peak:** < 0.0001
4H	Wilcoxon Signed-Rank Test	33 sessions from 7 mice	**Mean:** < 0.0001 **Min:** < 0.0001
4K	Wilcoxon Signed-Rank Test	26 sessions from 7 mice	**Left to Right (Lift to Retract):** < 0.0001 | < 0.0001 | 0.002 | 0.007 | 0.012
4L	Wilcoxon Signed-Rank Test	26 sessions from 7 mice	**Left to Right (Lift to Retract):** 0.005 | < 0.0001 | 0.277 | 0.0001 | 0.006
4N	Wilcoxon Signed-Rank Test	26 sessions from 7 mice	**X Seg Mean:** 0.006 **X Seg |Peak|:** 0.006 **Y Seg Mean:** 0.025 **Y Seg |Peak|:** 0.089 **Z Seg Mean:** < 0.0001 **Z Seg |Peak|:** < 0.0001
4O	Wilcoxon Signed-Rank Test	26 sessions from 7 mice	**Mean:** 0.0005 **Peak:** 0.012
4P	Wilcoxon Signed-Rank Test	26 sessions from 7 mice	**Mean:** 0.014 **Min:** 0.0001
5D	Wilcoxon Signed-Rank Test	13 sessions from 4 mice	**X Seg 1:** 0.048 **X Seg 2:** 0.017 **Y Seg 1:** 0.017 **Y Seg 2:** 0.027 **Z Seg 1:** 0.376 **Z Seg 2:** 0.001
5E	Wilcoxon Signed-Rank Test	13 sessions from 4 mice	**Seg 1:** 0.027 **Seg 2:** 0.0007
5F	Wilcoxon Signed-Rank Test	13 sessions from 4 mice	**Seg 1:** 0.0005 **Seg 2:** 0.001
5G	Wilcoxon Signed-Rank Test	13 sessions from 4 mice	**Left to Right (Lift to Retract):** 0.970 | 0.0002 | 0.305 | 0.002 | 0.001
5H	Wilcoxon Signed-Rank Test	13 sessions from 4 mice	**Left to Right (Lift to Retract):** 0.380 | 0.0002 | 0.216 | 0.005 | 0.001
5L	Wilcoxon Signed-Rank Test	30 sessions from 4 mice	**X Seg 1:** 0.019 **X Seg 2:** 0.503 **Y Seg 1:** 0.015 **Y Seg 2:** 0.360 **Z Seg 1:** 0.903 **Z Seg 2:** 0.198
5M	Wilcoxon Signed-Rank Test	30 sessions from 4 mice	**Seg 1:** 0.040 **Seg 2:** 0.221
5N	Wilcoxon Signed-Rank Test	30 sessions from 4 mice	**Seg 1:** 0.004 **Seg 2:** 0.919
5O	Wilcoxon Signed-Rank Test	30 sessions from 4 mice	**Left to Right (Lift to Retract):** 0.784 | 0.382 | 0.077 | 0.005 | 0.114
5P	Wilcoxon Signed-Rank Test	30 sessions from 4 mice	**Left to Right (Lift to Retract):** 0.571 | 0.584 | 0.003 | 0.008 | 0.008
S1B	Friedman test	251 sessions from 28 mice	< 0.0001
S1C	Friedman test	251 sessions from 28 mice	< 0.0001

S1D	Friedman test	251 sessions from 28 mice	< 0.0001
S1E	Friedman test	251 sessions from 28 mice	< 0.0001
S1F	Friedman test	251 sessions from 28 mice	< 0.0001
S1G	Friedman test	251 sessions from 28 mice	< 0.0001
S1H	Friedman test	251 sessions from 28 mice	< 0.0001
S1I	Friedman test	251 sessions from 28 mice	< 0.0001
S1J	Friedman test	251 sessions from 28 mice	< 0.0001
S1K	Friedman test	251 sessions from 28 mice	< 0.0001
S1L	Friedman test	251 sessions from 28 mice	< 0.0001
S1M	Friedman test	251 sessions from 28 mice	< 0.0001
S2B	Friedman test	21 sessions from 7 mice	< 0.0001
S2C	Friedman test	21 sessions from 7 mice	< 0.0001
S2D	Friedman test	21 sessions from 7 mice	< 0.0001
S2H	Friedman test	19 sessions from 7 mice	< 0.0001 | < 0.0001 | < 0.0001 | 0.0005 | 0.065
S2I	Friedman test	19 sessions from 7 mice	0.018 | 0.0003 | 0.0004 | < 0.0001 | 0.029
S3C	Wilcoxon Signed-Rank Test	14 sessions from 5 mice	**Within Laser:** 0.0001 **After Laser:** 0.0001
S3F	Wilcoxon Signed-Rank Test	14 sessions from 5 mice	0.583
S3G	Wilcoxon Signed-Rank Test	14 sessions from 5 mice	0.0004
S3J	Wilcoxon Signed-Rank Test	14 sessions from 5 mice	**Left to right:** 0.009 | 0.007 | 0.042 | 1.000 | 0.626
S3K	Wilcoxon Signed-Rank Test	14 sessions from 5 mice	**Left to right:** 0.013 | 0.011 | 0.013 | 0.855 | 0.626
S4B	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**20Hz - Control vs 40Hz - Control:** 0.301 **20 Hz vs Control:** 0.424 **40Hz vs Control:** 0.339
S4C	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**20Hz- Control vs 40Hz - Control:** 0.722 **20 Hz vs Control:**0.970 **40Hz vs Control:** 0.569
S4D	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**20Hz - Control vs 40Hz - Control** 0.233 **20 Hz vs Control:**0.910 **40Hz vs Control:** 0.110
S4E	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**20Hz - Control vs 40Hz - Control:** 0.077 **20 Hz vs Control:** 0.027 **40Hz vs Control:** 0.151

S4F	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**20Hz - Control vs 40Hz - Control:** 0.970 **20 Hz vs Control:** 0.001 **40Hz vs Control:** 0.007
S4J	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**20Hz- Control vs 40Hz-Control:** 0.233 | 0.002 | 0.677 | 0.077 | 0.204 **20 Hz vs Control:** 0.129 | 0.034 | 0.176 | 0.077 | 0.034 **40 Hz vs Control:** 0.064 | 0.002 | 0.110 | 0.009 | 0.009
S4K	Wilcoxon Signed-Rank Test	12 sessions from 4 mice	**20Hz- Control vs 40Hz-Control:** 1.000 | 0.012 | 0.110 | 0.034 | 0.034 **20 Hz vs Control:** 0.129 | 0.012 | 0.266 | 0.064 | 0.027 **40 Hz vs Control:** 0.424 | 0.007 | 0.002 | 0.005 | 0.002
S5B	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	0.881
S5C	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	0.838
S5D	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	0.683
S5E	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	0.157
S6B	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	0.708
S6C	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	0.0002
S6D	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	0.609
S6E	Wilcoxon Signed-Rank Test	21 sessions from 7 mice	0.838
S7D	Wilcoxon Signed-Rank Test	26 sessions from 7 mice	**X Seg Mean:** 0.007 | 0.143 | 0.899 | 0.216 | 0.191 **Y Seg Mean:** 0.080 | < 0.0001 | 0.248 | 0.027 | 0.312 **Z Seg Mean:** < 0.0001 | < 0.0001 | 0.002 | 0.002 | 0.120
S7E	Wilcoxon Signed-Rank Test	26 sessions from 7 mice	**Left to right:** < 0.0001 | < 0.0001 | 0.085 | 0.546 | 0.634
S8D	Wilcoxon Signed-Rank Test	30 sessions from 4 mice	**X Seg Mean** 0.635 | 0.038 | 0.003 | 0.324 | 0.730 **Y Seg Mean** 0.898 | 0.0005 | 0.051 | 0.324 | 0.700 **Z Seg Mean** 0.689 | 0.029 | 0.701 | 0.013 | 0.038
S8E	Wilcoxon Signed-Rank Test	30 sessions from 4 mice	**Left to right:** 0.303 | 0.019 | 0.001 | < 0.0001 | 0.016

1218