Research Article

Iota-carrageenan protects the ocular surface from desiccation-

induced cell death and tissue damage **in vitro** *and* **ex vivo**

Sabine Roch-Nakowitsch[a], Antonella Russo[a], Christiane Koller[a], Andrea Dolischka[a], Marielle König-

Schuster[a], Hanna Dellago[a], Eva Prieschl-Grassauer[a]

[a] Marinomed Biotech AG, Hovengasse 25, 2100 Korneuburg, Austria

Short Title: Iota-carrageenan eye drops

Corresponding Author:

Eva Prieschl-Grassauer, Marinomed Biotech AG, Hovengasse 25, 2100 Korneuburg, Austria.

Tel +43 226290300

E-mail: office@marinomed.com

Keywords: iota-carrageenan, Carragelose®, eye drops, dry eye disease, dry eye syndrome,

moistening, moisturizing.

1. Abstract

Iota-carrageenan is a natural polymer with moisturizing, mucoadhesive and shear-thinning properties. In this study, we aimed to evaluate the protective effects of iota-carrageenan on ocular surface against dehydration, to demonstrate its suitability for the use in lubricant eye drops. We utilized a human epithelial corneal cell culture model to test if pre-incubation with iota-carrageenan solution could protect cells from desiccation-induced cell death, to compare its effect with other natural polymers commonly used in artificial tear products, and to determine the optimal iota-carrageenan concentration. An *ex vivo* porcine eye model was established to confirm the protective effect of iota-carrageenan against dehydration on ocular tissue. Pre-incubation with 1.2 mg/ml iota-carrageenan increased the survival half-life of human corneal epithelial cells upon dehydration by three-fold; the effect was in the same range as observed for large molecular weight hyaluronic acid, and superior to all other tested natural polymers. The highest tested concentration of iota-carrageenan, 1.6 mg/ml, extended the cellular survival half-life by eight-fold while maintaining healthy cellular morphology. Repeated *ex vivo* instillation of an iota-carrageenan-based ophthalmic formulation into porcine eyes significantly protected the ocular surface from desiccation-induced corneal damage, as shown by corneal fluorescein staining These data suggest that Iota-carrageenan effectively moisturizes and protects the ocular surface, supporting its potential as a promising novel ingredient for eye drops in the management of dry eye disease.

2. Introduction

Dry eye disease (DED), also termed dry eye syndrome (DES) or keratoconjunctivitis sicca (KCS), is a multifactorial disease of the ocular surface characterized by a loss of homeostasis of the tear film and tear film instability. Symptoms include ocular dryness, irritation, redness, discharge, and blurred vision, can range from mild and occasional to severe and continuous, and have been shown to negatively impact work productivity and daily activities (Kanellopoulos and Asimellis, 2016; Liu et al., 2022; Sheppard et al., 2023).

The tear film of the eye, consisting of the lipid layer, the aqueous layer, and the mucous layer, is responsible for the maintenance of the corneal epithelium. The aqueous layer is responsible for the osmolality of the ocular surface and changes thereto lead to a disturbed homeostasis of the tear film. The mucous layer has viscoelastic and lubricating properties and detains the moisture on the corneal epithelium and is responsible to form a barrier against drugs, toxins, virus particles or allergens (Račić and Krajišnik, 2023; Willcox et al., 2017).

Mild to moderate cases of DED are usually treated with lubricant eye drops, gels or ointments that can be easily purchased over the counter. The majority of the available OTC lubricating products are gel-like ophthalmic formulations composed of ingredients that contain multifunctional large biopolymers such as hyaluronic acid, carboxymethyl cellulose or hydroxypropylmethylcellulose in order to increase the viscosity of the tear film, to moisturize and prolong the residence time of an aqueous layer on the cornea (Cassano et al., 2021; Račić and Krajišnik, 2023; Zhang et al., 2017). An ideal compound for the treatment of DED that aims to enhance tear film stability should possess properties similar to the mucous layer and in addition should be mucoadhesive to remain on the ocular surface. Here, we propose iota-carrageenan as a promising, safe and effective new alternative for the treatment of DED. Carrageenans are natural polysaccharides built of D-galactose and 3, 6-anhydro-D-galactose units which are extracted from certain red seaweeds of the Rhodophyceae class. In the food, cosmetic and pharmaceutical industry, carrageenans are widely utilized due to

their excellent physical functional properties, such as gelling, thickening, emulsifying and stabilizing

abilities (Li et al., 2014). There are three main types of carrageenan based on the number of sulfate

groups attached to the galactose molecule, which differ with regard to their solubility and their

gelling and thickening capacity: kappa- (one sulfate group per two galactose molecules), iota- (two

sulfate groups per two galactose molecules) and lambda- (three sulfate groups per two galactose

molecules) carrageenan. The toxicological profile of carrageenans has been thoroughly evaluated,

and they have been established to have minimal or no adverse physiological effects (Cohen and Ito,

2002). No health hazards have been documented for iota-carrageenan so far, and it has been

designated as "Generally Recognized As Safe" (GRAS status) by the FDA for food and topical

applications (U.S. Food & Drug Administration, SCOGS (Select Committee on GRAS Substances),

2020).

Iota-carrageenan is certified for marketing in the EU, parts of Asia and Australia under the brand

name Carragelose® as a component of nasal sprays, throat sprays and lozenges. Previous pre-clinical

and clinical studies led by us and others have demonstrated the broad, non-specific antiviral and

barrier-forming properties of iota-carrageenan (Bichiri et al., 2021; Eccles et al., 2010; Fazekas et al.,

2012; Fröba et al., 2021; Graf et al., 2018; Grassauer et al., 2008; Koenighofer et al., 2014; Ludwig et

al., 2013; Morokutti-Kurz et al., 2021; Unger-Manhart et al., 2024).

In addition, iota-carrageenan has great potential to be used in lubricating and moisturizing eye drops

based on the following properties: Firstly, it forms an *in situ* gelling system upon exposure to

physiological conditions (Akash Muktiram Shendge*, Vikas B Wamane, Jayshree R. Shejul, 2024;

Rupenthal et al., 2011) and hence retains moisture on the ocular surface due to its excellent water-

binding capacity (Thommes et al., 2009), counteracting evaporation. Secondly, it is a viscosity

enhancer that exhibits viscoelastic behavior (Hilliou, 2021; HOSSAIN et al., 1997). Thirdly, its

mucoadhesive properties are similar like that of the mucous layer (Račić and Krajišnik, 2023; Vigani

et al., 2019; Volod'ko et al., 2024). The second and third property additionally improves the

retention time on the ocular surface and thereby may provide a prolonged stay on the ocular surface

compared to aqueous solutions. Fourthly, it is a polymer with non-Newtonian properties and as such

it is per definition shear-thinning, hence reducing resistance against movements of the eyelid when

blinking (Ludwig, 2005). Finally, iota-carrageenan acts locally and is not absorbed by the body,

therefore it may be used also during pregnancy and lactation.

The suitability of iota-carrageenan for ophthalmic applications has already been demonstrated in

various contexts: Carrageenans have been tested and proposed as promising carriers for ocular drug

delivery (Bonferoni et al., 2004; Omran et al., 2023; Volod'ko et al., 2024). A commercial iota-

carrageenan-containing viscous gel has been proven to be safe and effective when used to protect

and hydrate the corneal surface during ophthalmic surgery (Giardini et al., 2018; Mencucci et al.,

2019). Hence, the potential of iota-carrageenan in multiple ophthalmic therapeutic areas may

deserve further attention. But to date, no study has evaluated the suitability of iota-carrageenan as

lubricant and moisturizing compound for treatment of DED.

The present research evaluated the moisturizing properties and the protective effects of iota-

carrageenan against air-induced desiccation. Using both *in vitro* cell culture and *ex vivo* porcine eye

model systems, we assessed the protective effect of iota-carrageenan in comparison to established

substances commonly used for the treatment of DED. Additionally, we demonstrated a

concentration-dependent protective effect of iota-carrageenan against dehydration and determined

the concentration of iota-carrageenan for use in an ophthalmic formulation.

3. Methods

3.1 Cell culture

We have adapted a cell culture model that has previously been used in the development of DED treatments to assess prevention of cell death from desiccation (Hill-Bator et al., 2014). Immortalized Human Corneal Epithelial Cell Line (IM-HCEpiC; Innoprot, Derio (Bizkaia), Spain; # P10871-IM) were cultured according to the supplier's instructions. Briefly, cells were maintained in IM-Ocular Epithelial Cell Medium (Innoprot; P60189) supplemented with growth factors and antibiotics in standard tissue culture flasks. Cells were incubated at 37°C in a humidified atmosphere containing 5% CO_2. The culture medium was exchanged with fresh medium every 2-3 days, and cells were passaged upon reaching 80-90% confluency using trypsin-EDTA.

3.2 *In vitro* dehydration assay

Human immortalized cornea epithelial cells (IM-HCEpiC) were seeded into 96-well plates (TPP; 92696) at a density of $1x10^4$ cells/well on the day before the experiment. The next day, medium was replaced with the sample solutions described below or with fresh cell culture medium only as control, and incubation under cell culture conditions was continued for 2 h. Subsequently, supernatants were removed, and culture plates were placed without lid in a dry incubator at 33°C for defined timespans of 2-40 min to allow dehydration. The following controls were included in the experiment: i) internal positive control (iPC) – cells were incubated with medium but not submitted to dehydration; ii) sample positive control (sPC) – cells were incubated with respective sample solution but not submitted to dehydration; iii) negative control (NC) – cells were incubated with cell culture medium and submitted to dehydration. Cell viability following dehydration was assessed fluorometrically using a resazurin based assay (Alamar Blue, Acros; 189900010). Cells were incubated with a 1x Alamar blue solution for 30 min at 37°C in a humidified atmosphere containing 5% CO_2. After incubation, the fluorescence of reduced resazurin was measured using a fluorescence

plate reader (BMG Fluostar Omega) with excitation at 544 nm and emission at 590 nm. Cell viability

was calculated for each sample relative to the corresponding sPC viability. For each tested sample

solution, a dose-response analysis was performed and the respective cellular survival half-life, i.e.,

the time point when 50 % of the metabolic signal was still detectable, was calculated based on the

dose-response curves. The goodness-of-fit of the dose-response analysis is expressed as confidence

interval of the calculated half-life.

3.3 Sample solutions used in the in vitro dehydration assay

For comparison of the dehydration-protective effect of natural polymers, the following substances

were dissolved in distilled water (aqua destillata, AD; Aqua Braun; B.Braun; 0082479E) and then

mixed 1:1 with cell culture medium to achieve a final assay concentration of 1.2 mg/ml: fucoidan

(two types: derived from Fucus vesiculosus (Sigma; F8190) and from Undaria pinnatifida (Sigma;

F8315)), dextran sulfate sodium salt (Sigma; D6001), carboxymethyl cellulose (Hercules; 7MF

Pharm), hydroxypropyl methylcellulose (Fagron; 102517), hyaluronic acid (three different molecular

weight species: 300-500 kDa (Sigma; 00941), 750-1000 kDa (Sigma; 53163) and 1.2-1.5MDa (Sigma;

53747), kappa-carrageenan (Du PontNutrition), and iota-carrageenan (Du PontNutrition).

For investigation of an iota-carrageenan dose effect, iota-carrageenan was dissolved in AD at a

concentration of 3.2 mg/ml and this stock solution was further diluted with AD to prepare solutions

with concentrations down to 0.15 mg/ml. These iota-carrageenan solutions were mixed 1:1 with cell

culture medium to obtain the following final assay concentrations of iota-carrageenan: 1.6, 1.4, 1.2,

1.0, 0.8, 0.6, 0.3, 0.15, and 0.075 mg/ml.

3.4 Cytoskeletal and nuclear staining after treatment and dehydration

Cell treatment and dehydration: IM-HCEpiC were seeded at a density of 8×10^4 cells/well on

coverslips (VWR; ECN631-1578) in a 24-well plate (TPP; 92024) the day before the experiment. Cells

were incubated for 2 h with 3.2 mg/ml iota-carrageenan in an isotonic buffer, mixed 1:1 with cell

culture medium to give a final assay concentration of 1.6 mg/ml. After the 2 h incubation, specific

wells were designated for dehydration condition. Cell culture medium and treatment solution was

aspirated from designated wells. Subsequently, the plate was placed without lid in a dry incubator at

33°C for 10 or 30 min. Following dehydration, cells were washed with PBS (Sigma; D8537), fixed with

Roti Histofix (Roth; P087.6) for 10 min, washed with PBS (3x), permeabilized with 0.1% Triton X-100

(Serva Feinbiochemica; 37240) for 5 min, and washed with PBS (3x). Coverslips were then stained for

1 h with 1x Phalloidin-iFluor 488 (Abcam; AB176753) plus 1 µg/ml of DAPI (Sigma; D9542) in PBS plus

1% BSA (Roth; 8076.4) on parafilm in a humidified petri dish. Coverslips were washed with PBS (3x)

and mounted on glass slides (Henry Schein; 900-2573) with mounting medium (Roti-Mount Aqua;

Roth; 2848.1) or epifluorescence microscopy (Axio ExaminerD1 with axiocam 202 camera; Carl

Zeiss).

3.5 *Ex vivo* dehydration assay

The induction of ocular surface damage by desiccation is based on a published model (Choy et al.,

2004). Porcine eyes including the surrounding tissue of the eye socket were extracted from skulls of

pigs (German-Large-White) slaughtered at a local slaughterhouse. Eyes were examined for corneal

damage by distributing one drop (25 µL) of sodium fluorescein (Fluka, 46960; 0.1% w/v in PBS) on

the ocular surface and scoring the surface damage under a black light torch l λ_{Em}: 395 nm according

to the Oxford grading scale. Only undamaged eyes (Oxford scale grading 0 or 1) were used for the

experiment. Eyes judged as undamaged were pinned open and incubated in a laminar flow hood for

6 h at room temperature and approx. 50% humidity. Every 15 min during this 6 h period, a single

drop of sample solution (described below) or 0.9% NaCl (B.Braun, 3570130) (negative control), was

dropped onto the cornea and spread across the surface by blinking with the nictitating membrane.

At the end of the 6 h dehydration period, eyes were again submitted to Oxford grading after

instillation of 1 drop of sodium fluorescein (0.1% w/v in PBS). Next, eyeballs were excised from the

surrounding tissue and immersed into sodium fluorescein (0.1% w/v in PBS) for 3 min, followed by

ten rinsing steps in PBS. Corneas were dissected, and 0.8 cm diameter punches were taken from the

center. Fluorescein was extracted from cornea punches in Triton X-100 (Sigma, T8787-50mL) (1 %

v/v in PBS) at 4°C for 48 h. Relative fluorescence units of extracts were measured at Ex/Em 485/520

nm using a BMG Fluostar Omega plate reader, and fluorescein concentration [µg/g] of each sample

was calculated based on a 20-point standard curve with two-fold serial dilutions. P values were

derived from one-way ANOVA with Tukey's post-hoc test.

3.6 Sample solutions used in the *ex vivo* dehydration assay

Iota-carrageenan dissolved in isotonic buffer composed of disodium-EDTA (Titriplex III) 1.00 mg/ml,

citric acid monohydrate 0.46 mg/ml, disodium hydrogen phosphate dihydrate 3.67 mg/ml, sorbitol

40.00 mg/ml, at concentrations 0.6 mg/ml, 2.0 mg/ml, and 3.2 mg/ml.

4. Results

Based on the inherent properties of iota-carrageenan and its demonstrated suitability for

ophthalmic applications [Bonferoni et al., 2004; Giardini et al., 2018; Mencucci et al., 2019; Omran et

al., 2023; Volod'ko et al., 2024], we set out to examine the potential of iota-carrageenan for use in

the treatment of DED.

To quantify and contextualize the moisturizing capacity of iota-carrageenan, we employed a cell

culture dehydration assay using human immortalized cornea epithelial cells (IM-HCEpiC). Cells were-

preincubated for 2 h with iota-carrageenan or other polymers that are currently used in products for

the treatment of DED, all at a final assay concentration of 1.2 mg/ml. Subsequently, cells were

submitted to dehydration for various time periods, and viability after dehydration was determined

fluorometrically using a resazurin based assay. Cellular survival half-lives upon dehydration, i.e. the

duration of dehydration when cell viability reached 50% of the respective non-dehydrated control,

were calculated for each sample. As shown in **figure 1**, pre-incubation of IM-HCEpiC with iota-

carrageenan protected cells from desiccation-induced cell death and led to a threefold increase of

the cellular survival half-life upon dehydration. This effect was specific to iota-carrageenan and was

not observed with kappa-carrageenan. The magnitude of effect was comparable with that of large

molecular weight hyaluronic acid (1.2-1.5 MDa) and was superior to the effect of all other tested

polymers, including fucoidan, dextran sulfate, carboxymethyl cellulose, hydroxypropyl

methylcellulose, as well as low and medium molecular weight hyaluronic acid (0.3-1 MDa).

Next, we aimed at determining if there is a concentration dependence of iota-carrageenan for

protection against dehydration, again using the IM-HCEpiC *in vitro* dehydration model. In this case,

cells were-preincubated for 2 h with various concentrations of iota-carrageenan spanning final assay

concentrations from 1.6 mg /ml down to 0.075 mg/ml, followed by dehydration for various

durations, and viability assessment based on resazurin (Alamar Blue). Our results demonstrate that

pre-incubation with iota-carrageenan enhanced cell viability after dehydration in a dose-dependent

manner. As shown for selected concentrations in **Figure 2a**, cells pre-incubated with medium alone

could compensate for only a 2 min dehydration period without loss of viability and had already lost

50% viability within less than 5 min of dehydration. In contrast, cells pre-incubated with the highest

tested concentration of 1.6 mg /ml iota-carrageenan were able to withstand dehydration and

maintain high viability levels of > 90% for at least 20 min and showed an increase in half-life survival

by eight-fold in comparison to the negative control (**Figure 2b**). Lower iota-carrageenan

concentrations had similar, though proportionally weaker effects. Iota-carrageenan concentrations

below 1 mg/ml led to markedly reduced viability already after 5-10 min of dehydration but still

extended the dehydration survival half-life, albeit to a lower extent. Concentrations below 0.15

mg/ml yielded similar results as the negative control (data not shown). In sum, the protective effect

of iota-carrageenan against dehydration shows a clear dose-dependence.

As shown in **figure 3,** dehydration caused cells to shrink and subsequently die, leading to reduced

cell density, when they had been pre-treated with medium only, while pre-incubation with 1.6

mg/ml iota-carrageenan prevents these changes and maintains cell density and healthy cell

morphology.

Next, we applied a porcine *ex vivo* model to validate the results obtained *in vitro*. **Figure 4a-c**

illustrates the experimental set-up. Eyes were extracted from the skulls of freshly slaughtered swine.

After transport to the laboratory, eyes were visually inspected for ocular surface damage, and only

undamaged eyes were used. Eye lids were pinned open and eyes were dehydrated by exposure to

laminar air flow for 6 h. During this period, eyes were treated in 15 min intervals with 1 drop of

either 0.9% NaCl or of 0.6, 2.0 or 3.2 mg/ml iota-carrageenan dissolved in isotonic buffer. In contrast

to the *in vitro* cell culture experiment where iota-carrageenan was dissolved in AD and 1:1 mixed

with cell culture medium to maintain cell viability, for *ex vivo* application iota-carrageenan was

dissolved in an isotonic buffered solution compatible for ophthalmic application, i.e., with neutral pH

containing sorbitol, di-sodium hydrogen phosphate dihydrate, citric acid monohydrate, and sodium-

ethylenediaminetetraacetic acid (EDTA) in AD. After 6 hours of dehydration, eyes were stained with

fluorescein to assess and quantify corneal damage caused by the desiccation. Fluorescein penetrates

poorly into the lipid layer of the corneal epithelium, and therefore, does not stain intact cornea, but

can only penetrate and stain damaged cornea where cell-to-cell junctions are disrupted (Srinivas and

Rao, 2023). As shown in **figure 5a**, no obvious cornea damage was seen before dehydration. After 6

h dehydration, a very small extent of damage was observed in eyes repeatedly treated with iota-

carrageenan eye drops during dehydration. In contrast, the control eyes treated with 0.9% NaCl had

acquired widespread corneal damage, as indicated by white arrows. The extent of damage can be

quantified by dye extraction from cornea punches after fluorescein staining. **Figure 5b** shows the

amount of fluorescein extracted from cornea punches, expressed as percentage of NaCl treated

control eyes. The amount of dye that penetrates the corneal epithelium and hence can be extracted

from cornea samples is proportional to the corneal damage acquired during dehydration. Repeated

treatment with iota-carrageenan protected porcine eyes from dehydration-induced cornea damage.

With all three tested concentrations of iota-carrageenan, the amount of extracted dye, i.e., of

corneal damage, was significantly reduced compared to control eyes treated with physiologic saline

solution. As shown before *in vitro*, there was a clear trend towards dose dependence of the effect,

even though the difference in effect size between concentrations was just short of being statistically

significant (p value for comparison between 0.6 and 3.2 mg/ml = 0.059) based on the available

number of biologic replicates. The baseline value has been derived from eyes not subjected to

dehydration and demonstrates the effectiveness of the dehydration treatment for inflicting corneal

damage.

In sum, our data demonstrate the capacity of iota-carrageenan to protect against dehydration in an

in vitro cell culture system of human cornea cells as well as in an *ex vivo* animal model. Based on

these *in vitro* and *ex vivo* results, 3.2 mg/ml was determined to be the most effective concentration

of iota-carrageenan to protect ocular cells and tissue from dehydration-induced damage.

5. Discussion

To restore and maintain a healthy ocular surface, it is essential that both the cornea and conjunctiva remain continuously hydrated - a function normally provided by the tear film. Recent consensus identifies tear film instability as the core mechanism underlying DED. This understanding has led to the development of tear film-oriented diagnosis, which seeks to clarify the specific causes of DED in individual patients by focusing on the distinct layers of the tear film. Consequently, tear-film-oriented DED therapy focuses on restoring tear film homeostasis and ensuring adequate lubrication of the ocular surface (Dash and Choudhury, 2024; Safir et al., 2024; Shimazaki, 2018).

The research presented in this article investigated the protective effect of iota-carrageenan against air-induced desiccation and evaluates its potential as an ocular lubricant for the relief of mild to moderate DED.

Our data show that in a cell culture model of corneal epithelial cells, pre-treatment of cultured cells with an iota-carrageenan solution dramatically enhanced cell survival after dehydration. The magnitude of the protective effect was comparable to high molecular weight hyaluronic acid, and was superior to all other tested natural polymers, including other widely used ingredients of eye drops used for DED treatments. Moreover, our results highlighted significant differences in the effectiveness of hyaluronic acid based on its molecular weight. The protective effect of iota-carrageenan was dose-dependent, as higher concentrations correlated with increased survival rates. Moreover, iota-carrageenan significantly inhibited the desiccation-stress induced accumulation of corneal damage in an *ex vivo* porcine eye model also in a dose dependent manner. In sum, the data presented here suggest that iota-carrageenan has a moisturizing and protective effect that prevents cornea cells from dehydration and, in consequence, from dehydration-induced cell death and tissue damage. Of note, the highest concentration used in the *ex vivo* experiment (3.2 mg/ml; maximum possible concentration for technical reasons) exceeds the highest concentration in the cell culture experiment described above (1.6 mg/ml). This was intended to compensate for the dilution with the

lacrimal fluid and the rapid restoration of the tear film that washes away a considerable proportion

of the administered solution (Gaudana et al., 2010).

The texture properties of carrageenans have long been known and exploited in the food and the

cosmetics industry. Carrageenans are known to retain moisture, enhance viscosity combined with

shear thinning behavior, and have mucoadhesive properties that are also exploited in various forms

of drug delivery (Bonferoni et al., 2004; Volod'ko et al., 2024). We hypothesize that these properties

together contribute to the effect observed in our experiments: Carrageenan forms a mucoadhesive

in situ film on the cellular monolayer and on the *ex vivo* porcine eye, it enhances viscosity of the

solution, improves surface retention, and counteracts dehydration due to water-retention. We claim

that these mechanisms should work equally well in the human eye, however this shall be subject to

further studies.

Hyaluronic acid containing eye drops have become the first choice for tear substitution in many

countries. According to literature, the average molecular weight of hyaluronic acid determines the

therapeutic efficacy of such eye drops (Salzillo et al., 2016), and high molecular weight hyaluronic

acid has enhanced mucoadhesion and moisturizing capacity compared to the low molecular weight

form (Guarise et al., 2023). Hyaluronic acid is a viscosity enhancer used to form gel-like ophthalmic

solutions which may cause, especially with high molecular weight, blurred vision and a foreign body

sensation, especially immediately after instillation (Müller-Lierheim, 2020). In contrast, in situ gelling

systems are generally considered more comfortable because they are installed as low-viscosity

liquids causing minimized immediate vision disruption (Cassano et al., 2021). In our experiments, the

viscosity of the high molecular weight hyaluronic acid solution was 9 mPas and that of the iota-

carrageenan solution 2 mPas, measured at 33°C and a shear rate of 10 1/s with an Rotavisc lo-vi

(IKA) with ELVAS-1 extension (data not shown). We have shown *in vitro* that iota-carrageenan is

equal to high molecular weight (1.2-1.5 MDa) hyaluronic acid in terms of cytoprotection under

dehydration at a concentration of 1.2 mg/ml and therefore expect similar therapeutic efficacy of

iota-carrageenan-based eye drops as reported for hyaluronic acid-based eye drops. It is important to

note that our *ex vivo* model represents only a simplified approximation of the eye *in vivo*, where

continuous secretion from lacrimal and Meibomian glands as well as regular blinking occur. To

partially compensate for the lack of natural blinking, our experimental set-up included mechanic

blinking with the lid to facilitate the distribution of test solutions. No tears replenishment was

performed in the *ex vivo* eyes which on one hand prevented wash out of the treatment but also led

to rapid cornea dehydration. We tried to compensate for this with repeated installations of the

respective test solutions. Nevertheless, this set up does not sufficiently consider the natural tear

clearance and the washout processes that occur *in vivo* .

Further studies are required to shed light on the safety, tolerability and therapeutic effectiveness of

iota-carrageenan eye drops in the treatment of DED in humans.

6. Conclusions

Iota-carrageenan exhibits significant dose-dependent protection against desiccation-induced cell

death and damage in both in vitro cultured human corneal cells and an ex vivo porcine eye model.

Iota-carrageenan closely mimics natural tear fluid due to its mucoadhesive properties, enabling it to

substitute lost tear fluid. By lubricating and hydrating the cornea and conjunctiva, it supports a

healthy ocular surface and additionally protects the eye surface from dehydration-induced damage.

Our findings position iota-carrageenan as a promising novel ingredient for DED-targeted eye drops.

Its dual capacity to replace tear fluid and shield the ocular surface from dehydration underscores its

potential as an advanced therapeutic agent.

7. Declarations

7.1 Acknowledgement

The authors thank Dorian Winter, Nicole Reiterer-Felber and Fabian Wallisch for excellent technical and scientific support.

7.2 Statement of Ethics

Regulations on the protection of animals used for scientific purposes do not apply to the present study, since tissues of animal origin used in this study were derived from the local slaughterhouse as waste product from food production under adherence to national laws on the protection of animals in connection with slaughtering.

7.3 Funding Sources

7.4 Disclosure

All authors are employees of Marinomed Biotech AG. EPG, SRN and AR are holders of a relevant patent: EP24167732. After completion of this investigation, Marinomed's iota-carrageenan product portfolio, including iota-carrageenan eyedrops, has transitioned to Unither Pharmaceuticals (Paris, France).

7.5 Data Availability Statement

Data related to this manuscript can be made available from the corresponding author upon reasonable request.

7.6 Author Contributions

SRN: conceptualization, project administration, supervision, data analysis, writing – review and editing; AR: methodology, supervision, data analysis, writing – review and editing; CK: methodology,

8. References

Akash Muktiram Shendge*, Vikas B Wamane, Jayshree R. Shejul, 2024. A Review on In Situ Gelling System.

Bichiri, D., Rente, A.R., Jesus, Â., 2021. Safety and efficacy of iota-carrageenan nasal spray in treatment and prevention of the common cold. Medicine and pharmacy reports 94 (1), 28–34.

Bonferoni, M.C., Chetoni, P., Giunchedi, P., Rossi, S., Ferrari, F., Burgalassi, S., Caramella, C., 2004. Carrageenan-gelatin mucoadhesive systems for ion-exchange based ophthalmic delivery: in vitro and preliminary in vivo studies. Eur. J. Pharm. Biopharm. 57 (3), 465–472.

Cassano, R., Di Gioia, M.L., Trombino, S., 2021. Gel-Based Materials for Ophthalmic Drug Delivery. Gels 7 (3).

Choy, E.P.Y., To, T.S.S., Cho, P., Benzie, I.F.F., Choy, C.K.M., 2004. Viability of porcine corneal epithelium ex vivo and effect of exposure to air: a pilot study for a dry eye model. Cornea 23 (7), 715–719.

Cohen, S.M., Ito, N., 2002. A critical review of the toxicological effects of carrageenan and processed eucheuma seaweed on the gastrointestinal tract. Crit. Rev. Toxicol. 32 (5), 413–444.

Dash, N., Choudhury, D., 2024. Dry Eye Disease: An Update on Changing Perspectives on Causes, Diagnosis, and Management. Cureus 16 (5), e59985.

Eccles, R., Meier, C., Jawad, M., Weinmüllner, R., Grassauer, A., Prieschl-Grassauer, E., 2010. Efficacy and safety of an antiviral Iota-Carrageenan nasal spray: a randomized, double-blind, placebo-controlled exploratory study in volunteers with early symptoms of the common cold. Respir. Res. 11, 108.

Fazekas, T., Eickhoff, P., Pruckner, N., Vollnhofer, G., Fischmeister, G., Diakos, C., Rauch, M., Verdianz, M., Zoubek, A., Gadner, H., Lion, T., 2012. Lessons learned from a double-blind randomised placebo-controlled study with a iota-carrageenan nasal spray as medical device in children with acute symptoms of common cold. BMC Complement. Altern. Med. 12, 147.

Fröba, M., Große, M., Setz, C., Rauch, P., Auth, J., Spanaus, L., Münch, J., Ruetalo, N., Schindler, M., Morokutti-Kurz, M., Graf, P., Prieschl-Grassauer, E., Grassauer, A., Schubert, U., 2021. Iota-Carrageenan Inhibits Replication of SARS-CoV-2 and the Respective Variants of Concern Alpha, Beta, Gamma and Delta. Int. J. Mol. Sci. 22 (24).

Gaudana, R., Ananthula, H.K., Parenky, A., Mitra, A.K., 2010. Ocular drug delivery. The AAPS journal 12 (3), 348–360.

Giardini, P., Hauranieh, N., Gatto, C., D'Amato Tóthová, J., 2018. Tripolymeric Corneal Coating Gel Versus Balanced Salt Solution Irrigation During Cataract Surgery: A Retrospective Analysis. Cornea 37 (4), 431–435.

Graf, C., Bernkop-Schnürch, A., Egyed, A., Koller, C., Prieschl-Grassauer, E., Morokutti-Kurz, M., 2018. Development of a nasal spray containing xylometazoline hydrochloride and iota-carrageenan for the symptomatic relief of nasal congestion caused by rhinitis and sinusitis. International journal of general medicine 11, 275–283.

Grassauer, A., Weinmuellner, R., Meier, C., Pretsch, A., Prieschl-Grassauer, E., Unger, H., 2008. Iota-Carrageenan is a potent inhibitor of rhinovirus infection. Virol. J. 5, 107.

Guarise, C., Acquasaliente, L., Pasut, G., Pavan, M., Soato, M., Garofolin, G., Beninatto, R., Giacomel, E., Sartori, E., Galesso, D., 2023. The role of high molecular weight hyaluronic acid in

mucoadhesion on an ocular surface model. Journal of the mechanical behavior of biomedical
materials 143, 105908.

Hill-Bator, A., Misiuk-Hojło, M., Marycz, K., Grzesiak, J., 2014. Trehalose-based eye drops preserve
viability and functionality of cultured human corneal epithelial cells during desiccation. BioMed
research international 2014, 292139.

Hilliou, L., 2021. Structure-Elastic Properties Relationships in Gelling Carrageenans. Polymers 13 (23).

HOSSAIN, K.S., NEMOTO, N., NISHINARI, K., 1997. Dynamic Viscoelasticity of Iota Carrageenan
Gelling System near Sol-Gel Transition. Nihon Reoroji Gakkaishi(Journal of the Society of
Rheology, Japan) 25 (3), 135–142.

Kanellopoulos, A.J., Asimellis, G., 2016. In pursuit of objective dry eye screening clinical techniques.
Eye and vision (London, England) 3.

Koenighofer, M., Lion, T., Bodenteich, A., Prieschl-Grassauer, E., Grassauer, A., Unger, H., Mueller,
C.A., Fazekas, T., 2014. Carrageenan nasal spray in virus confirmed common cold: individual
patient data analysis of two randomized controlled trials. Multidiscip. Respir. Med. 9 (1), 57.

Li, L., Ni, R., Shao, Y., Mao, S., 2014. Carrageenan and its applications in drug delivery. Carbohydr.
Polym. 103 (1), 1–11.

Liu, S.-H., Saldanha, I.J., Abraham, A.G., Rittiphairoj, T., Hauswirth, S., Gregory, D., Ifantides, C., Li, T.,
2022. Topical corticosteroids for dry eye. Cochrane Database Syst. Rev. 2022 (10).

Ludwig, A., 2005. The use of mucoadhesive polymers in ocular drug delivery. Adv. Drug Deliv. Rev. 57
(11), 1595–1639.

Ludwig, M., Enzenhofer, E., Schneider, S., Rauch, M., Bodenteich, A., Neumann, K., Prieschl-
Grassauer, E., Grassauer, A., Lion, T., Mueller, C.A., 2013. Efficacy of a carrageenan nasal spray in
patients with common cold: a randomized controlled trial. Respir. Res. 14, 124.

Mencucci, R., Cennamo, M., Favuzza, E., Rechichi, M., Rizzo, S., 2019. Triphasic polymeric corneal
coating gel versus a balanced salt solution irrigation during cataract surgery: A postoperative
anterior segment optical coherence tomography analysis and confocal microscopy evaluation. J.
Cataract Refract. Surg. 45 (8), 1148–1155.

Morokutti-Kurz, M., Fröba, M., Graf, P., Große, M., Grassauer, A., Auth, J., Schubert, U., Prieschl-
Grassauer, E., 2021. Iota-carrageenan neutralizes SARS-CoV-2 and inhibits viral replication in
vitro. PLoS ONE 16 (2).

Müller-Lierheim, W.G.K., 2020. Why Chain Length of Hyaluronan in Eye Drops Matters. Diagnostics
(Basel, Switzerland) 10 (8).

Omran, S., Elnaggar, Y.S.R., Abdallah, O.Y., 2023. Carrageenan tethered ion sensitive smart nanogel
containing oleophytocubosomes for improved ocular luteolin delivery. Int. J. Pharm. 646,
123482.

Račić, A., Krajišnik, D., 2023. Biopolymers in Mucoadhesive Eye Drops for Treatment of Dry Eye and
Allergic Conditions: Application and Perspectives. Pharmaceutics 15 (2).

Rupenthal, I.D., Green, C.R., Alany, R.G., 2011. Comparison of ion-activated in situ gelling systems for
ocular drug delivery. Part 1: physicochemical characterisation and in vitro release. Int. J. Pharm.
411 (1-2), 69–77.

Safir, M., Twig, G., Mimouni, M., 2024. Dry eye disease management. BMJ 384, e077344.

Salzillo, R., Schiraldi, C., Corsuto, L., D'Agostino, A., Filosa, R., Rosa, M. de, La Gatta, A., 2016.
Optimization of hyaluronan-based eye drop formulations. Carbohydr. Polym. 153, 275–283.

Sheppard, J., Shen Lee, B., Periman, L.M., 2023. Dry eye disease: identification and therapeutic
strategies for primary care clinicians and clinical specialists. Ann. Med. 55 (1), 241–252.
Shimazaki, J., 2018. Definition and Diagnostic Criteria of Dry Eye Disease: Historical Overview and
Future Directions. Invest. Ophthalmol. Vis. Sci. 59 (14), DES7-DES12.
Srinivas, S.P., Rao, S.K., 2023. Ocular surface staining: Current concepts and techniques. Indian
journal of ophthalmology 71 (4), 1080–1089.
Thommes, M., Ely, D.R., Kleinebudde, P., 2009. The water binding behavior of kappa-carrageenan
determined by three different methods. Pharm. Dev. Technol. 14 (3), 249–258.
U.S. Food & Drug Administration, SCOGS (Select Committee on GRAS Substances), 2020. SCOGS
database of Generally Recognized As Safe (GRAS) food substances: Entry for Carrageenan.
https://www.cfsanappsexternal.fda.gov/scripts/fdcc/?set=SCOGS&sort=Sortsubstance&order=A
SC&startrow=1&type=basic&search=carrageenan.
Unger-Manhart, N., Morokutti-Kurz, M., Zieglmayer, P.U., Lemell, P., Savli, M., Zieglmayer, R.,
Prieschl-Grassauer, E., 2024. Carrageenan-Containing Nasal Spray Alleviates Allergic Symptoms in
Participants with Grass Pollen Allergy: A Randomized, Controlled, Crossover Clinical Trial.
International journal of general medicine 17, 419–428.
Vigani, B., Faccendini, A., Rossi, S., Sandri, G., Bonferoni, M.C., Gentile, M., Ferrari, F., 2019.
Development of a Mucoadhesive and In Situ Gelling Formulation Based on κ-Carrageenan for
Application on Oral Mucosa and Esophagus Walls. I. A Functional In Vitro Characterization. Mar.
Drugs 17 (2).
Volod'ko, A.V., Son, E.Y., Glazunov, V.P., Davydova, V.N., Alexander-Sinkler, E.I., Aleksandrova, S.A.,
Blinova, M.I., Yermak, I.M., 2024. Carrageenan films as promising mucoadhesive ocular drug
delivery systems. Colloids and surfaces. B, Biointerfaces 237, 113854.
Willcox, M.D.P., Argüeso, P., Georgiev, G.A., Holopainen, J.M., Laurie, G.W., Millar, T.J., Papas, E.B.,
Rolland, J.P., Schmidt, T.A., Stahl, U., Suarez, T., Subbaraman, L.N., Uçakhan, O.Ö., Jones, L.,
2017. TFOS DEWS II Tear Film Report. The ocular surface 15 (3), 366–403.
Zhang, X., Jeyalatha M, V., Qu, Y., He, X., Ou, S., Bu, J., Jia, C., Wang, J., Wu, H., Liu, Z., Li, W., 2017.
Dry Eye Management: Targeting the Ocular Surface Microenvironment. Int. J. Mol. Sci. 18 (7).

Iota-carrageenan protects the ocular surface from desiccation-induced cell death and tissue damage in vitro *and* ex vivo

Figure Captions

Figure 1: Iota-carrageenan (white column) protect corneal cells from dehydration. IM-HCEpiC were pre-incubated with natural polymers in cell culture medium at a concentration of 1.2 mg/ml for 2h and then dehydrated for 2-40 min by removing the medium and placing the culture plates without lid in a dry incubator. Viability was assessed after dehydration photometrically using a resazurin based assay. Survival half-life was defined as time after which 50% of cells were still metabolic active. Error bars represent 95% CI (n= min. 3). NC = negative control, cells pre-treated with cell culture medium w/o polymers, followed by dehydration.

Figure 2: Iota-carrageenan protects corneal cells from dehydration in a concentration-dependant manner. IM-HCEpiC were pre-incubated with indicated iota-carrageenan concentrations in cell culture medium for 2h and then dehydrated for 2-40 min by removing the medium and placing the culture plates without lid in a dry incubator.

a: Viability was assessed photometrically using a resazurin based assay after dehydrating cells for indicated time periods. Viability is given relative to the viability of the respective sample positive control (i.e., cells pre-treated with same iota-carrageenan concentration as respective sample but did not undergo dehydration). Error bars represent the standard deviation across all replicates (n = 6-12). The horizontal red line represent 50% cell viability.

b: Survival half-life defined as time after which 50% of cells were still metabolic active after pretreatment with the tested concentrations of iota-carrageenan. Error bars represent 95% CI.

NC = negative control, cells pre-treated with cell culture medium w/o iota-carrageenan,

followed by dehydration.

Figure 3: Pre-treatment with iota-carrageenan eye drops protects corneal cells from dehydration-

induced morphological changes. IM-HCEpiC were pre-incubated with 3.2 mg/ml iota-

carrageenan mixed 1:1 with cell culture medium (giving a final iota-carrageenan concentration

of 1.6 mg/ml) for 2h and then dehydrated for 10 or 30 min. Actin fibers were stained with

phalloidin (green) and nuclei with DAPI (blue) to visualize morphology. Scale bare is 20 μm.

Figure 4: Experimental set-up of induction and detection of ocular surface damage in a porcine *ex*

***vivo* model.** Eyes extracted from freshly slaughtered swine were dehydrated by exposure to

laminar air flow for 6 h under repeated instillation of 0.9% NaCl or iota-carrageenan in isotonic

buffer. Eyes were visually inspected for ocular surface damage before and after dehydration.

Corneal damage was quantified by dye extraction from cornea punches after fluorescein

staining of eyeballs.

a-c Handling of porcine eyes. **a:** Porcine eye were pinned open and exposed to air for 6 h. Every 15

min, 1 drop of 0.9% NaCl or iota-carrageenan solution was instilled into the eye and evenly

spread by closing and opening the nictitating membrane. **b:** Immersion of eyeballs in 0.1%

fluorescein solution. **c**: Washing of fluorescein-stained eyeballs. After fluorescein staining,

corneas were dissected, circular punches were taken, and fluorescein was extracted from

punches and quantified.

Figure 5: Iota-carrageenan provides dose-dependent protection of the cornea from dehydration-

induced ocular surface damage in a porcine ex vivo ocular dehydration model.

a: Freshly extracted porcine eyes were examined for cornea damage under a black light torch after

instilling 25 µl fluorescein solution (left column). Only undamaged eyes (Oxford scale grading 0

or 1) were used in the experiment. After 6 hours of dehydration under repeated instillation of

0.9% NaCl or 3.2 mg/ml iota-carrageenan eye drops, eyes were again visually inspected under

black light after instilling of 25 µl fluorescein solution (right column). Widespread corneal

damage is seen after dehydration in NaCl-treated eyes, but not in eyes treated with 3.2 mg/ml

iota-carrageenan eye drops (white arrows indicate corneal damage).

b: The amount of fluorescein extracted from cornea punch samples after dehydration and

immersion into a 0.1% fluorescein solution, expressed as percentage of NaCl treated controls, is

proportional to the corneal damage acquired during dehydration. Error bars represent the SEM.

N=number of eyes used per treatment group. * $p \leq 0.05$; ** $p \leq 0.01$;*** $p \leq 0.001$. Baseline

value (=before dehydration) is significantly different from all other samples.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

a   b   c

Fig. 5

a

	Before dehydration	After 6 hours of dehydration and repeated treatment
Iota-carrageenan eye drops		
0.9% NaCl		

b